



Holomorphic bundles on complex manifolds with boundary

Andrei Teleman

► To cite this version:

| Andrei Teleman. Holomorphic bundles on complex manifolds with boundary. 2022. hal-03614586

HAL Id: hal-03614586

<https://hal.science/hal-03614586>

Preprint submitted on 20 Mar 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

HOLOMORPHIC BUNDLES ON COMPLEX MANIFOLDS WITH BOUNDARY

ANDREI TELEMAN

ABSTRACT. Let Ω be a complex manifold, and let $X \subset \Omega$ be an open submanifold whose closure \bar{X} is a (not necessarily compact) submanifold with smooth boundary.

Let G be a complex Lie group, Π be a differentiable principal G -bundle on Ω and J a formally integrable bundle almost complex structure on the restriction $\bar{P} := \Pi|_{\bar{X}}$. We prove that, if the boundary of \bar{X} is strictly pseudoconvex, J extends to a holomorphic structure on the restriction of Π to a neighborhood of \bar{X} in Ω . This answers positively and generalizes a problem stated in [Do]. We obtain a gauge theoretical interpretation of the quotient $\mathcal{C}^\infty(\partial\bar{X}, G)/\mathcal{O}^\infty(\bar{X}, G)$ associated with any compact Stein manifold with boundary \bar{X} endowed with a Hermitian metric.

For a fixed differentiable G -bundle \bar{P} on a complex manifold \bar{X} with non-pseudoconvex boundary, we study the set of formally integrable almost complex structures on \bar{P} which admit formally holomorphic local trivializations at boundary points. We give an example where a “generic” formally integrable almost complex on \bar{P} admits formally holomorphic local trivializations at *no* boundary point, whereas the set of formally integrable almost complex structures which admit formally holomorphic local trivializations at *all* boundary points is dense.

CONTENTS

1. Introduction	1
2. Strong integrability at pseudoconvex boundary points	5
3. Extension of holomorphic bundles to collar neighborhoods	6
4. Strong integrability at non-pseudoconvex boundary points	10
5. Appendix. Extending a section defined on a manifold with boundary	14
References	17

1. INTRODUCTION

A complex manifold with boundary is a manifold with boundary $\bar{X} = X \cup \partial\bar{X}$ endowed with a smooth (up to boundary) almost complex structure (ACS) j which is *formally integrable*, i.e. whose Neihenhuis tensor vanishes. Note that the “up to the boundary” version of the Newlander-Nirenberg theorem does not hold in general. More precisely the formal integrability of j does not guarantee the existence of j -compatible charts around the boundary points [Hi]. For this reason Hill uses the terminology “complex manifold with abstract boundary” for a pair (\bar{X}, j) as above [Hi].

2020 *Mathematics Subject Classification.* 32L05, 32G13, 32T15.

I am indebted to Mauro Nacinovich for a very useful exchange of mails on the collar neighborhood theorem. I thank Matei Toma for interesting discussions which led me to the problems considered in this article.

A smooth map $f : (\bar{X}, \partial\bar{X}) \rightarrow (\bar{Y}, \partial\bar{Y})$ between complex manifolds with boundary, or a smooth map $f : \bar{X} \rightarrow Y$ from a complex manifold with boundary to a complex manifold, will be called *formally holomorphic* if it is pseudo-holomorphic, i.e. if its differential at any point commutes with the two almost complex structures. Such a map is holomorphic in the classical sense (hence analytic) at any $x \in X$, but, in general, not at a boundary point; this justifies the chosen terminology *formally* holomorphic. If $Y = \mathbb{C}$, the formal holomorphy condition becomes $\bar{\partial}f = 0$.

Let G be a complex Lie group and $p : \bar{P} \rightarrow \bar{X}$ be a differentiable principal G -bundle on \bar{X} . The boundary $\partial\bar{P}$ of \bar{P} is the pre-image $p^{-1}(\partial\bar{X})$. By definition, a bundle ACS on \bar{P} is an ACS which makes the projection $\bar{P} \rightarrow \bar{X}$ and the G -action $\bar{P} \times G \rightarrow \bar{P}$ pseudo-holomorphic. In the particular case $G = \mathrm{GL}(r, \mathbb{C})$, the data of a bundle ACS J on \bar{P} is equivalent to the data of a Dolbeault operator $\delta : A^0(\bar{X}, E) \rightarrow A^{0,1}(\bar{X}, E)$ on the associated vector bundle E , and the formal integrability condition for J is equivalent to the formal integrability condition $\delta^2 = 0$ for δ .

The main goal of this article is the following extension theorem:

Theorem 3.1. *Let Ω be a complex manifold, and let $X \subset \Omega$ be an open submanifold whose closure \bar{X} is a submanifold with smooth boundary. Let G be a complex Lie group, Π be a differentiable principal G -bundle on Ω and J be a formally integrable bundle ACS on the restriction $\bar{P} := \Pi|_{\bar{X}}$*

If the boundary of \bar{X} is strictly pseudoconvex, there exists an open neighborhood Ω' of \bar{X} in Ω and an integrable bundle ACS J' on $\Pi|_{\Omega'}$ which extends J .

I came to this statement thinking of the following problem stated in [Do, p. 102]:

Problem. (Donaldson's problem) Let $X \subset \mathbb{C}^n$ be a relatively compact domain with smooth, strictly pseudoconvex boundary, and let E be a topologically trivial holomorphic vector bundle on \bar{X} . Prove that E has a global smooth trivialization on \bar{X} which is holomorphic on X .

In the special case $n = 2$ the claim is proved in [Do, Appendix A]. As noted at the beginning of Donaldson's proof, the claim follows easily from Grauert's classification theorem for holomorphic bundles on Stein manifolds if one can prove that E extends to an open neighborhood of \bar{X} in \mathbb{C}^n .

Therefore a special case of Theorem 3.1 gives the needed extension property, so it solves Donaldson's problem.

The proof of Theorem 3.1 is inspired by the beautiful proof of the collar neighborhood theorem of [HiNa] for complex manifolds with strictly pseudoconvex boundary. I am deeply indebted to Mauro Nacinovich for answering in detail my questions and clearing up a technical difficulty in the proof of [HiNa, Theorem 1]; the same difficulty occurs in general when one uses Zorn's lemma to prove extension theorems for objects defined on manifolds with boundary. The Appendix explains in detail this difficulty in a general framework which includes both extensions problems; the main result of this section (Lemma 5.5) is essentially due to Nacinovich [Nac].

Note that the conclusion of Theorem 3.1 holds for any, not necessarily reductive, complex Lie group G and for any open, not necessarily relatively compact, submanifold X with smooth, strictly pseudoconvex boundary of a complex manifold Ω . Moreover, our formal integrability condition for a bundle ACS does not require the existence of formally holomorphic local trivializations of \bar{P} at boundary points $x \in \partial\bar{X}$. This *strong integrability condition* is required in Donaldson's definition of a holomorphic bundle on a complex manifold with boundary [Do, p. 91] so, in fact, it is part of the hypothesis of the original Donaldson's problem. On the other hand, under the assumptions of Theorem 3.1, this condition is superfluous, because (see section 2):

Proposition 2.3. *Let \bar{X} be a complex manifold with boundary, G be a complex Lie group and $p : \bar{P} \rightarrow \bar{X}$ be a principal G -bundle on \bar{X} endowed with a formally integrable bundle ACS J . Let $x \in \partial\bar{X}$ and $y \in \bar{P}_x$. If \bar{X} is (weakly) pseudoconvex around x , there exists an open neighborhood U of x in \bar{X} and a formally holomorphic section $s : U \rightarrow \bar{P}$ such that $s(x) = y$.*

In other words *any* formally integrable bundle ACS on a bundle over a complex manifold with pseudoconvex boundary satisfies Donaldson's strong integrability condition.

We now recall the motivation behind Donaldson's problem: to give a gauge-theoretical interpretation of the quotient of the group of smooth maps $\partial\bar{X} \rightarrow \mathrm{GL}(r, \mathbb{C})$ by the subgroup of those maps which admit a formally holomorphic extension $\bar{X} \rightarrow \mathrm{GL}(r, \mathbb{C})$ (see [Do, p. 102]). Using Theorem 3.1 and the generalization of Donaldson's [Do, Theorem 1] to the Hermitian framework [Xi], one obtains the following generalization of this isomorphism theorem:

Theorem 3.3. *Let K be a compact Lie group and G be its complexification. Let $\bar{X} = X \cup \partial\bar{X}$ be a compact Stein manifold with boundary. Endow \bar{X} with any (not necessarily Kählerian) Hermitian metric g .*

Let $\mathcal{O}^\infty(\bar{X}, G)$ be the group of formally holomorphic G -valued maps on \bar{X} , identified with a subgroup of $\mathcal{C}^\infty(\partial\bar{X}, G)$ via the restriction map.

There exists a natural bijection between the moduli space of boundary framed Hermitian-Yang-Mills connections on the trivial K -bundle on (\bar{X}, g) and the quotient $\mathcal{C}^\infty(\partial\bar{X}, G)/\mathcal{O}^\infty(\bar{X}, G)$.

Note that in this statement \bar{X} is just an abstract complex manifold with boundary. However, by the collar neighborhood theorem mentioned above (see [HiNa] for the general case and [NaOh, Theorem p. 706], [Oh], [Ca2, Theorem 6] for the compact case), the proof of Theorem 3.3 can make use of Theorem 3.1.

The moduli spaces intervening in Theorem 3.3 will play an important role in a joint project developed in collaboration with Matei Toma dedicated to the generalization of the concept "bounded family of coherent sheaves" in non-algebraic complex geometry. We will be especially interested in the case when \bar{X} is a compact neighborhood of a 2-codimensional analytic set in a complex n -manifold. For $n \geq 3$, such a neighborhood will not be pseudoconvex in general. Therefore we come to the following two natural questions:

- (Q1) What is the role of Donaldson's strong integrability condition in the theory?
- (Q2) To what extent is the pseudoconvexity assumption in Proposition 2.3 necessary? If \bar{X} is not assumed to be pseudoconvex, what can be said about the abundance of the ACS which satisfy this strong integrability condition within the whole space of formally integrable bundle ACS on a bundle \bar{P} ?

Concerning (Q1): A complex manifold with boundary \bar{X} has the structure of a locally ringed space: it can be endowed with the sheaf $\mathcal{O}_{\bar{X}}^\infty$ of formally holomorphic \mathcal{C}^∞ -functions. The restriction of this sheaf to X coincides with \mathcal{O}_X and its restriction to $\partial\bar{X}$ always contains the constant sheaf $\mathbb{C}_{\partial\bar{X}}$.

The sheaf \mathcal{E}^∞ of formally holomorphic sections of a rank r vector bundle E on \bar{X} endowed with a formally integrable ACS (Dolbeault operator) is naturally a sheaf of $\mathcal{O}_{\bar{X}}^\infty$ -modules.

Remark 1.1. (1) The following conditions are equivalent:

- (a) E satisfies Donaldson's strong integrability condition.
- (b) The natural map $\mathcal{E}_x^\infty \rightarrow E_x$ is surjective for any $x \in \partial\bar{X}$.

- (2) If one of these two equivalent conditions is satisfied, \mathcal{E}^∞ is locally free of rank r .
- (3) The assignment $E \mapsto \mathcal{E}^\infty$ defines an equivalence between the groupoid of formally holomorphic vector bundles satisfying the strong integrability condition and the groupoid of locally free sheaves of \mathcal{O}_X^∞ -modules.

This remark shows that Donaldson's strong integrability condition for formally integrable bundle ACS is very natural: it defines the class of vector bundles which (as do holomorphic bundles in classical complex geometry) correspond to locally free sheaves on the base manifold, regarded as a ringed space.

On the other hand *neither* the main result of [Do] *nor* its generalization to the Hermitian framework really needs this strong integrability condition. One can see this in [Xi, p. 317-318]: a local holomorphic frame is only used to give explicit formulae for the connection matrix of a Chern connection (formula (2.2)) or for the Hermitian-Einstein flow (formula (2.6')). These explicit formulae show that the Hermitian-Einstein flow is a semi-linear strictly parabolic system. But similar (slightly more complicated) formulae are obtained using an arbitrary smooth local frame. More precisely, let E be a rank r vector bundle on \bar{X} endowed with a formally integrable Dolbeault operator δ , (f_1, \dots, f_r) be a smooth frame defined on an open set $U \subset \bar{X}$ and $\alpha \in A^{0,1}(U, \text{gl}(r\mathbb{C}))$ the matrix valued $(0,1)$ -form defined by

$$\delta(f_1, \dots, f_r) = (f_1, \dots, f_r)\alpha.$$

Putting $H_{ij} = h(f_i, f_j)$ we see that, in the local frame (f_1, \dots, f_r) , the connection form of Chern connection is $H^{-1}\partial H - H^{-1}{}^t\bar{\alpha}H + \alpha$, and the Hermitian-Einstein flow reads:

$$\begin{aligned} \dot{H} = & -2i\Lambda\bar{\partial}\partial H + 2i\Lambda(\bar{\partial}H \wedge (H^{-1}\partial H) - \bar{\partial}H \wedge (H^{-1}{}^t\bar{\alpha}H) + \bar{\partial}({}^t\bar{\alpha})H + {}^t\bar{\alpha} \wedge \bar{\partial}H \\ & - H\partial\alpha - \partial H \wedge \alpha + {}^t\bar{\alpha}H \wedge \alpha - (H\alpha H^{-1}) \wedge \partial H + (H\alpha H^{-1}) \wedge {}^t\bar{\alpha}H). \end{aligned}$$

This is also a semi-linear, strictly parabolic system.

Concerning (Q2): For a \mathcal{C}^∞ vector bundle E on a complex manifold with boundary \bar{X} , let Δ_E be the space of formally integrable Dolbeault operators on E ; for $x \in \partial\bar{X}$ denote by Δ_E^x the subspace of those $\delta \in \Delta_E$ which admit a formally δ -holomorphic frame (satisfies the strong integrability condition) around x . The space we are interested in is the intersection $\Delta_E^{\text{si}} := \bigcap_{x \in \partial\bar{X}} \Delta_E^x$ of formally integrable Dolbeault operators which are strongly integrable (admit formally holomorphic frames) around all boundary points.

The example below shows that, in general, on non-pseudoconvex manifolds, the strong integrability condition defines a set which is infinite codimensional, meagre (of first Baire category), but dense. Moreover, in our example, a “generic” formally integrable Dolbeault operator does *not* admit formally holomorphic frames at *any* boundary point.

Proposition 4.5. *Let \bar{X} be the complement of the standard ball $B \subset \mathbb{C}^2$ in $\mathbb{P}_{\mathbb{C}}^2$, L be a trivial \mathcal{C}^∞ complex line bundle on \bar{X} .*

- (1) *The union $\bigcup_{x \in \partial\bar{X}} \Delta_L^x$ is a first Baire category subset of Δ_L , in particular its subsets Δ_L^x (for $x \in \partial\bar{X}$), Δ_L^{si} have the same property.*
- (2) *For any $x \in \partial\bar{X}$, Δ_L^x is an infinite codimensional affine subspace of Δ_L .*
- (3) *Δ_L^{si} is a first Baire category, infinite codimensional, but dense affine subspace of Δ_L .*

This shows that, in general, on non-pseudoconvex manifolds, Donaldson's strong integrability condition might not define a locally closed subset of Δ_E . This complication should be taken into account in the construction of the moduli space of boundary framed holomorphic bundles on a non-pseudoconvex \bar{X} .

The proof of Proposition 4.5 is related to Lewy's celebrated example of a first order differential operator L with analytic coefficients on \mathbb{R}^3 such that, for "generic" smooth second term f , the "inhomogeneous equation" $Lu = f$ is non-solvable on any non-empty open set [Le], [Hö].

2. STRONG INTEGRABILITY AT PSEUDOCONVEX BOUNDARY POINTS

Let $\bar{X} = X \cup \partial\bar{X}$ be a complex n -manifold with boundary. As explained in the introduction, the formal integrability of j does not guarantee the existence of formally j -holomorphic charts around the boundary points [Hi]. By the main result of [Ca1], this difficulty vanishes if we assume (weak) pseudoconvexity of the boundary:

Let $x \in \partial\bar{X}$. We'll say that \bar{X} has pseudoconvex boundary around x if there exists an open neighborhood U of x in X and a real smooth non-positive function r on U such that $r^{-1}(0) = \partial\bar{X} \cap U$, $d_ur \neq 0$ for any $u \in \partial\bar{X} \cap U$, and $\partial\bar{\partial}r(a, \bar{a}) \geq 0$ for any $a \in T_{u,X}^{1,0} \cap T_{u,\partial\bar{X}}^C$, $u \in \partial\bar{X} \cap U$. [Ca1, Theorem, p. 234] can be reformulated as follows

Theorem 2.1. *Suppose that \bar{X} has pseudoconvex boundary around x . There exists an open neighborhood U of x and a formally j -holomorphic embedding $f : U \rightarrow \mathbb{C}^n$.*

By "embedding" we mean here an immersion which defines a homeomorphism on its image. It follows that $f(U)$ is an n -dimensional submanifold with boundary of \mathbb{C}^n whose boundary (interior) is $f(\partial\bar{X} \cap U)$ (respectively $f(U \cap X)$) and the induced map $U \rightarrow f(U)$ is a diffeomorphism which restricts to a biholomorphism between the interiors. As mentioned above,

Remark 2.2. If we assume that \bar{X} has *strictly* pseudoconvex boundary one can prove a stronger result: \bar{X} admits a holomorphic collar neighborhood, i.e. it can be embedded holomorphically in a complex manifold Ω [HiNa].

Proposition 2.3. *Let $p : \bar{P} \rightarrow \bar{X}$ be principal G bundle on \bar{X} endowed with a formally integrable bundle ACS J . Let $x \in \partial\bar{X}$, $y \in \bar{P}_x$ and suppose that \bar{X} has pseudoconvex boundary around x . There exists an open neighborhood U of x in \bar{X} and a formally holomorphic section $s : U \rightarrow \bar{P}$ such that $s(x) = y$.*

Proof. Regard \bar{P} as complex manifold with boundary $\partial\bar{P} = p^{-1}(\partial\bar{X})$, and note that this boundary is pseudoconvex around y . To see this, use the pull-back of a boundary defining function r for \bar{X} around x .

By Catlin's Theorem 2.1 there exists an open neighborhood Q of y in \bar{P} and a formally J -holomorphic embedding $F : Q \rightarrow \mathbb{C}^N$, where $N = n + \dim(G)$. The formal holomorphy condition implies that the restriction of F to any complex submanifold contained in the boundary is holomorphic. Therefore we obtain a holomorphic embedding $F|_{\bar{P}_x \cap Q} : \bar{P}_x \cap Q \rightarrow \mathbb{C}^N$.

Let $A \subset \mathbb{C}^N$ be an n -dimensional affine subspace which contains $F(y)$ and is transversal to $F(\bar{P}_x \cap Q)$ at this point. The pre-image via F of the germ $(A, F(y))$ is the germ at y of n -dimensional complex submanifold S of Q which is transversal to the fiber \bar{P}_x at y . Therefore S defines a holomorphic local section around x . ■

Corollary 2.4. *Let E be complex vector bundle on \bar{X} endowed with a Dolbeault operator $\delta : \Gamma(E) \rightarrow \Gamma(\Lambda^{0,1} \otimes E)$ satisfying the formal integrability condition $\delta^2 = 0$. Then E admits a formally δ -holomorphic trivialization around any boundary point around which \bar{X} has pseudoconvex boundary.*

Proof. The claim follows from Proposition 2.3 taking into account that, for a Dolbeault operator δ on E , the formal integrability condition for the associated bundle ACS on the frame bundle \bar{P}_E , reads $\delta^2 = 0$. ■

Remark 2.5. As the example studied in section 4 shows, the existence of a formal holomorphic trivialization around a boundary point does not hold without the pseudoconvexity condition (even if \bar{X} admits a holomorphic collar neighborhood).

3. EXTENSION OF HOLOMORPHIC BUNDLES TO COLLAR NEIGHBORHOODS

The goal of this section is the proof of the following extension theorem:

Theorem 3.1. *Let (Ω, j) be a complex manifold, and let $X \subset \Omega$ be an open submanifold whose closure \bar{X} is a submanifold with smooth boundary. Let G be a complex Lie group, $\Pi \xrightarrow{\pi} \Omega$ be a differentiable principal G -bundle on Ω and J be a formally integrable bundle ACS on the restriction $\bar{P} := \Pi|_{\bar{X}}$*

If the boundary of \bar{X} is strictly pseudoconvex, there exists an open neighborhood Ω' of \bar{X} in Ω and an integrable bundle ACS J' on $\Pi|_{\Omega'}$ which extends J .

Our proof is inspired by the proof of the collar neighborhood theorem for complex manifolds with strictly pseudo-convex boundary [HiNa], but uses a new ingredient: the étale space associated with a sheaf of sets and Godement's theorem [Go, Théorème 3.3.1 p. 150], which yields a general extension principle for a section defined on a subset of the base of an étale space. This allows us to give a simple construction of the inductively ordered set (to which Zorn's lemma is applied), which does not need factorization by an equivalence relation. Note also that in the proof we will point out the role of Lemma 5.5 to overcome the difficulty explained in the appendix.

We start with a remark which will allow us to control the derivatives of a bundle ACS J in a global way, without making use of charts and local trivializations.

Denoting by $V \subset T_\Pi$ the vertical tangent subbundle, we obtain the short exact sequence

$$0 \rightarrow V \xrightarrow{j} T_\Pi \xrightarrow{\pi_*} \pi^*(T_\Omega) \rightarrow 0$$

of G -vector bundles on Π . Factorizing by G , we obtain the short exact sequence

$$0 \rightarrow \text{ad}(\Pi) := \Pi \times_{\text{Ad}} \mathfrak{g} \xrightarrow{\tilde{j}} Q := T_\Pi/G \xrightarrow{\tilde{\pi}_*} T_\Omega \rightarrow 0 \quad (1)$$

of vector bundles on Ω (compare with [At, Theorem 1, p. 187]).

Remark 3.2. The data of a bundle ACS J on Π is equivalent to the data of a section $s_J \in \Gamma(\Omega, \text{End}(Q))$ with $s_J^2 = -\text{id}_Q$ which makes the bundle morphisms \tilde{j} and $\tilde{\pi}_*$ fiberwise \mathbb{C} -linear. Therefore one can use the formalism explained in Section 5 to define globally the derivatives of a bundle ACS J : one uses linear connections on T_Ω^* and $\text{End}(Q)$ and the associated differential operators D^k applied to s_J .

Proof. (of Theorem 3.1) Let \mathcal{J} be the space of germs of integrable locally defined bundle ACS on Π . Therefore, as a set, $\mathcal{J} = \coprod_{x \in \Omega} \mathcal{J}_x$ where

$$\mathcal{J}_x = \varinjlim_{\substack{U \ni x \\ U \text{ open}}} \mathcal{J}(U)$$

and $\mathcal{J}(U)$ denotes the set of integrable bundle ACSs on $\Pi|_U$. The set \mathcal{J} comes with an obvious surjection $q : \mathcal{J} \rightarrow \Omega$. For an element $\mathcal{J} \in \mathcal{J}(U)$ we denote by $\hat{\mathcal{J}} : U \rightarrow \mathcal{J}$ the associated section of q given by $\hat{\mathcal{J}}(u) = \mathcal{J}_u$. The topology of \mathcal{J} is generated by the images of the sections of this form, and this topology makes q local homeomorphisms. In the terminology of [Go, Section II.1.2] \mathcal{J} is the étale space associated with the sheaf of sets on Ω defined by $U \mapsto \mathcal{J}(U)$. Note also that, for any open set $U \subset \Omega$, the map $\mathcal{J} \mapsto \hat{\mathcal{J}}$ identifies $\mathcal{J}(U)$ with the set $\Gamma(U, \mathcal{J})$ of continuous sections of \mathcal{J} defined on U [Go, Théorème 1.2.1, p 111].

The restriction J_X of J to $\bar{P}|_X = \Pi|_X$ defines a continuous section $\hat{J}_X : X \rightarrow \mathcal{J}$, but we have *no* obvious extension of this section to \bar{X} , because J , although is defined and smooth up to the boundary, does not give integrable bundle ACSs locally (with respect to Ω) around boundary points.

Consider the set

$$\mathcal{R} := \{(V, \alpha) \mid V \text{ open in } \bar{X}, X \subset V \subset \bar{X}, \alpha \in \Gamma(V, \mathcal{J}), \alpha|_X = \hat{J}_X\}$$

of *continuous* extensions of \hat{J}_X to open (with respect to the relative topology) subsets of \bar{X} . \mathcal{R} is obviously non-empty and has an obvious partial order which satisfies the hypothesis of Zorn's lemma: every chain in \mathcal{R} has an upper bound. Therefore, by this lemma, there exists a maximal element $(V^{\max}, \alpha^{\max})$ of \mathcal{R} .

Claim. $V^{\max} = \bar{X}$.

The first step for proving the claim uses [Go, Théorème 3.3.1 p. 150]. According to this theorem¹, there exists an open set O of Ω containing V^{\max} and $\mathcal{J} \in \mathcal{J}(O)$ such that $\hat{\mathcal{J}}|_{V^{\max}} = \alpha^{\max}$. The maximality property of $(V^{\max}, \alpha^{\max})$ implies

$$\bar{X} \cap O = V^{\max}. \quad (2)$$

Note now that $J|_X = \mathcal{J}|_X$, so the corresponding sections (see Remark 3.2)

$$s_J \in \Gamma(\bar{X}, \text{End}(Q)), \quad s_{\mathcal{J}} \in \Gamma(O, \text{End}(Q))$$

agree on X , and the results of Section 5 apply. According to Lemma 5.5 we may suppose (by replacing O by a smaller open set still satisfying (2), if necessary) that the triple $(s_J, O, s_{\mathcal{J}})$ satisfies property \mathbf{P}_k for any $k \geq 0$. We will need this later in the proof.

Assume now by *reductio ad absurdum* that the claim is false and let $x_0 \in \bar{X} \setminus V^{\max}$. Since $V^{\max} \supset X$, we have $x_0 \in \partial \bar{X}$, so, by (2), it follows

$$x_0 \in \partial \bar{X} \setminus O. \quad (3)$$

By Whitney's theorem there exists a smooth non-negative real function φ on Ω such that $\varphi^{-1}(0) = \Omega \setminus O$.

Let B be a relatively compact open neighborhood of x_0 in Ω and $\rho : B \rightarrow \mathbb{R}$ be a smooth, strictly plurisubharmonic defining function for $X \cap B$. In other words $\rho^{-1}((-\infty, 0)) = X \cap B$, $\rho^{-1}(0) = \partial \bar{X} \cap B$ and $d\rho$ is nowhere vanishing on this intersection. We may suppose that ρ has a smooth extension around \bar{B} which is a strictly plurisubharmonic submersion.

For sufficiently small $\varepsilon > 0$ the function $\rho_\varepsilon := \rho - \varepsilon\varphi \in \mathcal{C}^\infty(B, \mathbb{R})$ will still be strictly plurisubharmonic submersion, so $X_\varepsilon^B := \rho_\varepsilon^{-1}((-\infty, 0))$ will be the interior of a strictly pseudoconvex manifold $\bar{X}_\varepsilon^B := \rho_\varepsilon^{-1}((-\infty, 0])$ with boundary $\partial \bar{X}_\varepsilon^B = \rho_\varepsilon^{-1}(0)$. The superscript B on the left emphasizes that the closure is taken with respect to the relative topology of B .

Note that the submanifolds X_ε^B , \bar{X}_ε^B of B have the properties

$$\bar{X} \cap O \cap B \subset X_\varepsilon^B \subset O \cap B. \quad (4)$$

$$\bar{X}_\varepsilon^B \subset (\bar{X} \cup O) \cap B. \quad (5)$$

Indeed, for the first inclusion in (4) note that, for a point $w \in \bar{X} \cap O$, we have $\varphi(w) > 0$ because, by construction, φ is positive on O , and $\rho(w) \leq 0$ because

¹The hypothesis of Godement's theorem requires the existence of fundamental system of paracompact neighborhoods of V^{\max} in Ω . Note that, in general, an open subspace of a paracompact space is not necessarily paracompact. In this article all manifolds are supposed second-countable by definition. This condition implies paracompactness and is "hereditary" with respect to open embeddings.

$x \in \bar{X}$. For the second inclusion in (4) note that, if a point $x \in X_\varepsilon^B$ does not belong to O , then $\varphi(x) = 0$, so the condition $\rho_\varepsilon(x) < 0$ becomes $\rho(x) < 0$ which implies $x \in X \subset O$. A similar argument proves (5).

Formulae (4), (5) show that $X_\varepsilon^B \subset O \cap B$ and

$$X \cap B \subset (\bar{X} \cap B) \cap (O \cap B) \subset X_\varepsilon^B \subset \bar{X}_\varepsilon^B \subset (\bar{X} \cap B) \cup (O \cap B),$$

so \bar{X}_ε^B interpolates between $\bar{X} \cap B$ and $O \cap B$ in the sense of Definition 5.3 given in the Appendix (see Fig. 1). We know that the triple $(O \cap B, s_{\mathcal{J}}|_{O \cap B})$ satisfies property \mathbf{P}_k for any $k \geq 0$. By Remark 5.4 explained in the appendix it follows that there exists a *smooth* almost complex structure J_ε on $\Pi|_{\bar{X}_\varepsilon^B}$ which agrees with J on $\bar{X} \cap B$ and with \mathcal{J} on X_ε^B . Note that, as explained in the appendix, the smoothness of J_ε does not follow using only the smoothnesses of \mathcal{J} and J .

Note that J_ε is formally integrable, because J and \mathcal{J} have this property. We have $x_0 \in \partial \bar{X}_\varepsilon^B$, because, by (3), $\rho(x) = \varphi(x) = 0$. By Proposition 2.3 applied to $(\Pi|_{\bar{X}_\varepsilon^B}, J_\varepsilon)$ there exists an open neighborhood U of x_0 in \bar{X}_ε^B and a smooth section $s_\varepsilon \in \Gamma(U, \Pi)$ which is formally holomorphic with respect to J_ε . Since J_ε agrees with \mathcal{J} on X_ε^B , it follows that s_ε is \mathcal{J} holomorphic on $X_\varepsilon^B \cap U$.




FIGURE 1. The picture in the case $V^{\max} = \bar{X} \setminus \{x_0\}$.

By Remark 5.1, the section s_ε can be smoothly extended across the boundary, i.e. there exists a smooth extension $\tilde{s}_\varepsilon \in \Gamma(\tilde{U}, \Pi)$ of s_ε to an open set \tilde{U} of B such that $\bar{X}_\varepsilon^B \cap \tilde{U} = U$.

Endow the restriction $\Pi|_{\tilde{U}}$ with the integrable bundle ACS $J_{\tilde{s}_\varepsilon}$ which makes \tilde{s}_ε holomorphic. The section \tilde{s}_ε is \mathcal{J} -holomorphic on $X_\varepsilon^B \cap U$, because on this set it coincides with s_ε ; it follows that $J_{\tilde{s}_\varepsilon}$ agrees with \mathcal{J} above this *open* set, so

$$\hat{J}_{\tilde{s}_\varepsilon}|_{X_\varepsilon^B \cap U} = \hat{\mathcal{J}}|_{X_\varepsilon^B \cap U}. \quad (6)$$

On the other hand we have $V^{\max} \cap \tilde{U} \subset V^{\max} \cap B \subset X_\varepsilon^B$ (we used (2) and (4)), so $V^{\max} \cap \tilde{U} \subset X_\varepsilon^B \cap \tilde{U} = X_\varepsilon^B \cap \bar{X}_\varepsilon^B \cap \tilde{U} = X_\varepsilon^B \cap U$, so (6) gives

$$\hat{J}_{\tilde{s}_\varepsilon}|_{V^{\max} \cap \tilde{U}} = \hat{\mathcal{J}}|_{V^{\max} \cap \tilde{U}} = \alpha^{\max}|_{V^{\max} \cap \tilde{U}}.$$

Therefore $\hat{J}_{\tilde{s}_\varepsilon}$ defines an extension of α^{\max} on $V^{\max} \cup \tilde{U}$, so on $V^{\max} \cup (\tilde{U} \cap \bar{X})$; this latter extension is continuous, because α^{\max} and $\hat{J}_{\tilde{s}_\varepsilon}$ are continuous and V^{\max} ,

$\tilde{U} \cap \bar{X}$ are open in \bar{X} . Since $x_0 \in V^{\max} \cup (\tilde{U} \cap \bar{X})$, this contradicts the maximality of α^{\max} , and the claim is proved.

We now know that $V^{\max} = \bar{X}$. But then $\Omega' := O$ is an open neighborhood of \bar{X} in Ω , and $J' := \mathcal{J}$ is an integrable bundle ACS on $\Pi_{\Omega'}$ which extends J . ■

As explained in the introduction, the motivation behind the extension Theorem 3.1 is the following generalization of Donaldson's gauge theoretical interpretation of the quotient $\mathcal{C}^\infty(\partial\bar{X}, \text{GL}(r, \mathbb{C}))/\mathcal{O}^\infty(\bar{X}, \text{GL}(r, \mathbb{C}))$ associated to a compact, strictly pseudoconvex submanifold with boundary in \mathbb{C}^n (see [Do, p. 102]):

Theorem 3.3. *Let K be a compact Lie group and G be its complexification. Let $\bar{X} = X \cup \partial\bar{X}$ be a compact Stein manifold with boundary. Endow \bar{X} with any (not necessarily Kählerian) Hermitian metric g .*

Let $\mathcal{O}^\infty(\bar{X}, G)$ be the group of formally holomorphic G -valued maps on \bar{X} , identified with a subgroup of $\mathcal{C}^\infty(\partial\bar{X}, G)$ via the restriction map.

There exists a natural bijection between the moduli space of boundary framed Hermitian-Yang-Mills connections on the trivial K -bundle on (\bar{X}, g) and the quotient $\mathcal{C}^\infty(\partial\bar{X}, G)/\mathcal{O}^\infty(\bar{X}, G)$.

We recall from [Do] that a (differentiably trivial) boundary framed holomorphic bundle on \bar{X} is a pair (J, θ) consisting of a formally integrable bundle ACS on the trivial \mathcal{C}^∞ -bundle $\bar{X} \times G$ and a smooth section (or, equivalently, a trivialization) of the restriction of this bundle to $\partial\bar{X}$. Let \mathcal{M} be the moduli space of pairs (J, θ) as above, modulo the natural action of the gauge group $\text{Aut}(\bar{X} \times G) = \mathcal{C}^\infty(\bar{X}, G)$ on the space of such pairs.

Taking into account Donaldson's isomorphism theorem [Do, Theorem 1'] which gives a gauge theoretical interpretation of the moduli space \mathcal{M} in terms of boundary framed Hermitian-Yang-Mills connections (and the generalization of this theorem to the Hermitian framework), the claim of Theorem 3.3 will follow as in Donaldson's article from the following

Lemma 3.4. *The natural map $R : \mathcal{C}^\infty(\partial\bar{X}, G)/\mathcal{O}^\infty(\bar{X}, G) \rightarrow \mathcal{M}$ given by*

$$R([f]) := [(J_0, f)],$$

where J_0 is the trivial holomorphic structure on $\bar{X} \times G$, is bijective.

Note that $[(J_0, f)] = [(J_0, f')]$ in \mathcal{M} if and only if $[f] = [f'] \bmod \mathcal{O}^\infty(\bar{X}, G)$, so R is well defined and injective. The image of R is the space of gauge classes $[(J, \theta)]$ such that J admits a global formally holomorphic trivialization on \bar{X} .

Therefore, Theorem 3.3 will follow from:

Lemma 3.5. *Let J be a formally integrable bundle ACS on $\bar{X} \times G$. Then $(\bar{X} \times G, J)$ admits a global formally holomorphic trivialization on \bar{X} .*

Proof. By the collar neighborhood theorem [HiNa], \bar{X} can be embedded in a complex manifold Ω . \bar{X} is a compact Stein manifold with boundary, so, by definition, there exists a strictly pseudoconvex smooth nonpositive real function ρ on \bar{X} such that $\partial\bar{X} = \rho^{-1}(0)$ and 0 is a regular value of ρ . Let $\tilde{\rho}$ be any smooth extension of ρ to a sufficiently small open neighborhood Ω' of \bar{X} in Ω such that $\tilde{\rho}$ is positive on $\Omega' \setminus \bar{X}$. Therefore the fibre $\tilde{\rho}^{-1}(0) = \partial\bar{X}$ is compact and 0 is a regular value of $\tilde{\rho}$. By the bicollar neighborhood theorem for compact real hypersurfaces (see for instance [Ma, Theorem 2.32, p. 68]) there exists $\eta > 0$, an open neighborhood U of $\partial\bar{X}$ in Ω' and a diffeomorphism $h : \partial\bar{X} \times (-\eta, \eta) \rightarrow U$ such that $h(x, 0) = x$ and $\tilde{\rho}(h(x, t)) = t$ for any $(x, t) \in \partial\bar{X} \times (-\eta, \eta)$. Choose $\varepsilon \in (0, \eta)$ sufficiently small such that $\tilde{\rho}$ remains strictly pseudoconvex on the pre-image

$$\bar{X}_\varepsilon := (\tilde{\rho}|_U)^{-1}((-\infty, \varepsilon]).$$

It follows that \bar{X}_ε is still a compact Stein manifold with boundary.

Let now $\Pi = \Omega \times G$ be the trivial \mathcal{C}^∞ -bundle on Ω and J a formally integrable bundle ACS on the restriction $\Pi|_{\bar{X}} = \bar{X} \times G$.

By Theorem 3.1, there exists an open neighborhood Ω' of \bar{X} and an integrable bundle ACS J' on $\Pi|_{\Omega'}$ extending J . Choosing ε sufficiently small, we may suppose $\Omega' = X_\varepsilon$. The pair $(\Pi|_{\Omega'}, J')$ is a topologically trivial holomorphic G -bundle on the Stein manifold X_ε , so it is holomorphically trivial by Grauert's classification theorem of holomorphic bundles on Stein manifolds (see [Gr], [For, Theorem 8.2.1, p. 356]). The restriction to \bar{X} of a global holomorphic trivialization $(\Pi|_{\Omega'}, J')$ will be a global formally holomorphic trivialization of $(\bar{X} \times G, J)$. ■

4. STRONG INTEGRABILITY AT NON-PSEUDOCONVEX BOUNDARY POINTS

Let $L = \bar{X} \times \mathbb{C}$ be the trivial line bundle on \bar{X} and let σ_0 the section $x \mapsto (x, 1)$. For a Dolbeault operator δ on L let α_δ be $(0,1)$ -form defined by $\delta\sigma_0 = \alpha_\delta\sigma_0$. The map $\delta \mapsto \alpha_\delta$ identifies the space Δ_L of formally integrable Dolbeault operators on L with the closed subspace $Z^{01}(\bar{X}) \subset A^{01}(\bar{X})$ of $\bar{\partial}$ -closed $(0,1)$ -forms on \bar{X} .

A local frame $\sigma = \varphi\sigma_0 \in \Gamma(U, L)$ is formally holomorphic with respect to δ if and only if $\bar{\partial}\varphi + \alpha_\delta\varphi = 0$. If U is simply connected we may write $\varphi = e^\psi$, and the above formula becomes $\bar{\partial}\psi = -\alpha_\delta$. This shows that

Remark 4.1. Let $\delta \in \Delta_L$ and $U \subset \bar{X}$ be a simply connected open set. L admits a formally δ -holomorphic frame on U if and only if $\alpha_\delta|_U$ is $\bar{\partial}$ -exact.

Unfortunately, in general, the $\bar{\partial}$ -Poincaré lemma does not hold at boundary points [AH].

For the rest of this section \bar{X} will be the complement of the standard ball $B \subset \mathbb{C}^2$ in $\mathbb{P}_{\mathbb{C}}^2$. The boundary complex [FK, Chapter V], [Fo] on the sphere $S := \partial\bar{X}$ reduces to

$$0 \rightarrow B^{00} = \mathcal{C}^\infty(S^3, \mathbb{C}) \xrightarrow{\bar{\partial}_S} B^{01} = \Gamma(S, \Lambda_H^{01}) \rightarrow 0,$$

where $H := T_S \cap jT_S$ and $\Lambda_H^{01} \subset H^* \otimes \mathbb{C}$ is the bundle of $(0,1)$ -forms on H . Regarding S^3 as an S^1 -bundle on $\mathbb{P}_{\mathbb{C}}^1$, H is just the horizontal distribution of the connection which corresponds to the Chern connection of the tautological line bundle on $\mathbb{P}_{\mathbb{C}}^1$.

Remark 4.2. Let $r : A^{01}(\bar{X}) \rightarrow B^{01}$ be the morphism induced by “restriction to the boundary”.

- (1) The morphism $H^{01}(\bar{X}) \rightarrow H^{01}(S)$ induced by r is an isomorphism.
- (2) The morphism $Z^{01}(\bar{X}) \rightarrow B^{01}$ induced by r is surjective.

Proof. (1) The injectivity of $H^{01}(\bar{X}) \rightarrow H^{01}(S)$ follows by [AH, Theorem 5, p. 355] taking into account that $H^{01}(\mathbb{P}_{\mathbb{C}}^2) = 0$. The surjectivity follows from [AH, Theorem 6(c), p. 355] taking into account that $H^{02}(\mathbb{P}_{\mathbb{C}}^2) = 0$ and $H^{01}(\bar{B}) = 0$. The latter is obtained using the vanishing of the sheaf cohomology space $H^1(B, \mathcal{O}_B)$ (which is obvious because B is Stein) and the comparison theorem [FK, Theorem 4.3.1, p. 77] taking into account the discussion on p. 57 in the same book.

(2) Let $\beta \in B^{01}$. The surjectivity of $H^{01}(\bar{X}) \rightarrow H^{01}(S)$ shows that there exists $\alpha \in Z^{01}(\bar{X})$ and $\varphi \in B^{00}$ such that $\beta = r(\alpha) + \bar{\partial}_S\varphi$. Choosing a smooth extension $\psi \in \mathcal{C}^\infty(\bar{X}, \mathbb{C})$ of φ we see that

$$\beta = r(\alpha) + \bar{\partial}_S(\psi|_S) = r(\alpha + \bar{\partial}\psi),$$

which proves the claim. ■

The generalized Cayley transform

$$C : \mathbb{C}^2 \setminus (\{-1\} \times \mathbb{C}) \rightarrow \mathbb{C}^2 \setminus (\{-i\} \times \mathbb{C}), \quad C(z_1, z_2) := \left(i \frac{1 - z_1}{1 + z_1}, \frac{z_2}{1 + z_1} \right)$$

identifies biholomorphically the unit ball $B \subset \mathbb{C}^2$ with the Siegel upper half-space

$$\mathcal{U} := \{(w_1, w_2) \in \mathbb{C}^2 \mid \Im(w_1) > |w_2|^2\},$$

and the punctured sphere $S \setminus \{(-1, 0)\}$ with the real hypersurface

$$\Sigma := \{(w_1, w_2) \in \mathbb{C}^2 \mid \Im(w_1) = |w_2|^2\} \subset \mathbb{C}^2$$

(the “unbounded realization” of the 3-sphere, see [Kr, p. 112]).

Since C is biholomorphic it follows that the boundary operator $\bar{\partial}_S$ on $S \setminus \{(-1, 0)\}$ corresponds via C to the boundary operator $\bar{\partial}_\Sigma$. Using the diffeomorphism

$$k : \mathbb{R} \times \mathbb{C} \rightarrow \Sigma, \quad k(t, w) = (t + i|w|^2, w)$$

we see [AH, p. 359-361] that $\bar{\partial}_\Sigma$ is given explicitly by

$$\bar{\partial}_\Sigma(\psi) = (L(\psi \circ k) \circ k^{-1}) d\bar{w}_2, \quad (7)$$

where $L : \mathcal{C}^\infty(\mathbb{R} \times \mathbb{C}, \mathbb{C}) \rightarrow \mathcal{C}^\infty(\mathbb{R} \times \mathbb{C}, \mathbb{C})$ is the Lewy operator [Le]

$$L\eta = \frac{\partial \eta}{\partial \bar{w}} - iw \frac{\partial \eta}{\partial t}.$$

The composition

$$h := k^{-1} \circ C : S \setminus \{(-1, 0)\} \rightarrow \mathbb{R} \times \mathbb{C}$$

is a chart of the sphere, and, in this chart, the operator $\bar{\partial}_S$ is given by

$$\bar{\partial}_S(\varphi) = (L(\varphi \circ h^{-1}) \circ h) C^*(d\bar{w}_2). \quad (8)$$

Lemma 4.3. *With the notations above we have:*

(1) *Let $V \subset S$ be a non-empty open subset. The set*

$$M^V := \{\beta \in B^{01} \mid \text{the equation } \bar{\partial}_S u = \beta|_V \text{ has a distribution solution } u \in \mathcal{D}'(V)\}$$

is a first Baire category subset of the Fréchet space B^{01} .

(2) *The union*

$$M := \bigcup_{\substack{\emptyset \neq V \subset S \\ V \text{ open}}} M^V$$

is a first Baire category subset of the Fréchet space B^{01} .

Proof. (1) We adapt the elegant proof of [Hö, Theorem 3.2, p. 135] to our situation explaining the fundamental ideas and the necessary changes. Note that our M is a subset of the Fréchet space B^{01} of sections of the bundle Λ_S^{01} on the compact manifold S , whereas the set M considered in Hörmander’s proof is a subset of the Fréchet space

$$\dot{B}(\Omega) := \{\varphi \in \mathcal{C}^\infty(\Omega, \mathbb{C}) \mid \forall \gamma \in \mathbb{N}^n \forall \varepsilon > 0 \exists K \subset \Omega \text{ compact s.t. } \sup_{\Omega \setminus K} |D^\gamma \varphi| < \varepsilon\}$$

associated with an open set $\Omega \subset \mathbb{R}^n$.

For an open set $\omega \subset S \setminus \{(-1, 0)\}$ and $N \in \mathbb{N}^*$ put

$$\begin{aligned} \mathcal{D}_N(\omega) &:= \left\{ u \in \mathcal{D}(\omega) \mid \sum_{|\gamma| \leq N} \sup |D^\gamma(\varphi \circ h^{-1})| \leq \frac{1}{N} \right\}, \\ \mathcal{D}'_N(\omega) &:= \{ u \in \mathcal{D}'(\omega) \mid \forall \varphi \in \mathcal{D}_N(\omega), |u(\varphi)| \leq 1 \} = \\ &= \{ u \in \mathcal{D}'(\omega) \mid \forall \varphi \in \mathcal{D}(\omega), |u(\varphi)| \leq N \sum_{|\gamma| \leq N} \sup |D^\gamma(\varphi \circ h^{-1})| \}. \end{aligned}$$

A theorem of Schwartz (see for instance [Vo, Theorem 10.12 p. 87]) shows that $\mathcal{D}_N(\omega)$ is a neighborhood of 0 in $\mathcal{D}(\omega)$, so, by Alaoglu–Bourbaki’s Theorem (see [Vo, Theorem 4.7 p. 32]), $\mathcal{D}'_N(\omega)$ is weakly compact in $\mathcal{D}'(\omega)$.

It follows that

$$M_N^\omega := \{\beta \in B^{01} \mid \exists u \in \mathcal{D}'_N(\omega), \bar{\partial}_S u = \beta|_\omega\}$$

is a closed subset of B^{01} . We claim that $M_N^\omega = \emptyset$. Indeed, note first that, taking into account (8), the properties of Lewy’s operator explained in [Hö, p. 136] and [Hö, Theorem 3.1], there exists $g \in B^{01}$ with $\text{supp}(g) \subset \omega$ such that the equation $\bar{\partial}_S u = g$ has no solution in $\mathcal{D}'(\omega)$. If an interior point $\beta \in M_N^\omega$ existed, we could find $\varepsilon > 0$ such that $\beta + \varepsilon g \in M_N^\omega$. Since M_N^ω is symmetric and convex, we obtain $\frac{\varepsilon}{2}g \in M_N^\omega$, which contradicts the way in which g has been chosen.

Now fix a non-empty open subset $\omega \subset S$ with $\bar{\omega} \subset V \setminus \{(-1, 0)\}$. Therefore $\bar{\omega}$ is a compact subset of V which is contained in the domain of the chart h . Taking into account the definition of $\mathcal{D}'(V)$, it follows that for any $u \in \mathcal{D}'(V)$ there exists $N \in \mathbb{N}$ such that $u|_\omega \in \mathcal{D}'_N(\omega)$. This implies

$$M^V \subset \bigcup_{N \in \mathbb{N}} M_N^\omega,$$

so M^V , being a subset of a first category subset, is itself of first category.

(2) Use a countable basis for the topology of S . ■

For an open set $U \subset \bar{X}$ put $Z^U := \{\alpha \in Z^{01}(\bar{X}) \mid \alpha|_U \text{ is } \bar{\partial}\text{-exact}\}$.

Proposition 4.4. *The union*

$$Z := \bigcup_{\substack{U \subset \bar{X} \text{ open} \\ U \cap \partial \bar{X} \neq \emptyset}} Z^U$$

is a first Baire category subset of $Z^{01}(\bar{X})$.

Proof. Let $r : Z^{01}(\bar{X}) \rightarrow B^{01}$ be the restriction morphism. The restriction to $U \cap S$ of a $\bar{\partial}$ -exact form on U is $\bar{\partial}_S$ -exact. This shows that, with the notation introduced in Lemma 4.3, we have

$$Z^U \subset r^{-1}(M^{U \cap S}), \quad Z \subset r^{-1}(M).$$

The claim follows now from Lemma 4.3 taking into account that

- (a) by Remark 4.2 and the Open Mapping Theorem, r is an open map.
- (b) Since r is open and continuous, the correspondence r^{-1} (pre-image via r) commutes with interior and closure. ■

Let now Δ_L be the space of formally integrable Dolbeault operators on L . For $x \in \partial \bar{X}$ let Δ_L^x be the subspace of those $\delta \in \Delta_L$ which admit a formally δ -holomorphic frame around x . The intersection $\Delta_L^{\text{si}} := \bigcap_{x \in \partial \bar{X}} \Delta_L^x$ is precisely the space of formally integrable Dolbeault operators on L which are strongly integrable (admit formally holomorphic frames) around all boundary points. We can now prove:

Proposition 4.5. *Let \bar{X} be the complement of the standard ball $B \subset \mathbb{C}^2$ in $\mathbb{P}_{\mathbb{C}}^2$, L be a trivial \mathcal{C}^∞ complex line bundle on \bar{X} .*

- (1) *The union $\bigcup_{x \in \partial \bar{X}} \Delta_L^x$ is a first Baire category subset of Δ_L , in particular its subsets Δ_L^x (for $x \in \partial \bar{X}$), Δ_L^{si} have the same property.*
- (2) *For any $x \in \partial \bar{X}$, Δ_L^x is an infinite codimensional affine subspace of Δ_L .*

- (3) Δ_E^{si} is a first Baire category, infinite codimensional but dense affine subspace of Δ_L .

Proof. (1) The first claim follows from Remark 4.1 and Proposition 4.4 taking into account that $\bigcup_{x \in \partial \bar{X}} \Delta_E^x$ corresponds to Z via the isomorphism

$$\Delta_L \ni \delta \mapsto \alpha_\delta \in Z^{01}(\bar{X}).$$

- (2) Taking into account Remarks 4.1, 4.2, it suffices to prove that the linear subspace

$$B^x := \{\beta \in B^{01} \mid \text{the germ of } \beta \text{ at } x \text{ is } \bar{\partial}_S\text{-exact}\}$$

is infinite codimensional in B^{01} . We may suppose $x = (1, 0)$ whose image via the chart $h = k^{-1} \circ C$ is the origin of $\mathbb{R} \times \mathbb{C}$. Let $\Lambda \subset B^{01}$ be the linear subspace defined by

$$\Lambda := \{\beta \in B^{01} \mid \text{the germ of } (\beta/C^*(d\bar{w}_2)) \circ h^{-1} \text{ at } 0 \text{ depends only on } t\}.$$

The main theorem of [Le] shows that, for a function $f \in \mathcal{C}^\infty(\mathbb{R} \times \mathbb{C})$ whose germ at 0 depends only on t , if the equation

$$Lu = f$$

is smoothly solvable around 0, then this germ is real analytic. Taking into account formula (8) and Lewy's result it follows that

$$B^x \cap \Lambda \subset \{\beta \in \Lambda \mid \text{the germ of } (\beta/C^*(d\bar{w}_2)) \circ h^{-1} \text{ at } 0 \text{ is analytic}\},$$

whose codimension in Λ is infinite. Therefore B^x is infinite codimensional in B^{01} .

- (3) The first idea would be to use the subspace $Z_{\text{an}}^{01}(\bar{X})$ of real analytic $\bar{\partial}$ -closed forms on \bar{X} . Unfortunately, it is not clear if this space is dense in $Z^{01}(\bar{X})$.

Instead we will use the pre-image of the space B_{an}^{01} of real analytic sections of the bundle Λ_H^{01} , which is dense in B^{01} . Since $r : Z^{01}(\bar{X}) \rightarrow B^{01}$ is continuous and open, it follows that $r^{-1}(B_{\text{an}}^{01})$ is a dense subset of $Z^{01}(\bar{X})$.

Taking into account again Remark 4.1 it suffices to prove that for any $\alpha \in r^{-1}(B_{\text{an}}^{01})$ and any $x \in \partial \bar{X}$ there exists $U \subset \bar{X}$ open such that $x \in U$ and $\alpha|_U$ is $\bar{\partial}$ -exact.

Since $\alpha|_S$ is real analytic, by the first order Cauchy–Kovalevskaya theorem, there exists an open neighborhood V of x in S and an analytic solution of the equation $\bar{\partial}_S u = \alpha|_S$ on V . Let W be Stein neighborhood of x in \mathbb{C}^2 such that $W \cap S \subset V$. Put

$$W^\pm := \{x \in W \mid \pm(1 - \|x\|^2) \geq 0\}.$$

Therefore

$$W^+ = W \cap \bar{B}, \quad W^- = W \cap \bar{X}.$$

Applying [AH, Theorem 5, p. 355] to the triple (W, W^-, W^+) we see that the restriction morphism

$$H^{01}(W^-) \rightarrow H^{01}(W \cap S)$$

is injective. Since the restriction of $\alpha|_S$ to $W \cap S$ is $\bar{\partial}_S$ -exact, it follows that the restriction of α to W^- is $\bar{\partial}$ -exact, which proves the claim. ■

5. APPENDIX. EXTENDING A SECTION DEFINED ON A MANIFOLD WITH BOUNDARY

We begin with a natural formalism for characterizing the smoothness sections in vector bundles in an invariant way, without making use of charts and local trivializations. Instead we will use connections.

Let Ω be a an n -dimensional differentiable manifold, E be differentiable real vector bundle on Ω and ∇, ∇_E linear connections on T_Ω^* and E respectively.

We define inductively differential operators (acting on local sections of E) by $D^1 = \nabla_E$, $D^{k+1} = (\nabla^{\otimes k} \otimes \nabla_E) \circ D^k$. A continuous section $s : U \rightarrow E$ (on an open set $U \subset \Omega$) is of class \mathcal{C}^1 on if and only $D^1 s$ exists and is continuous. Proceeding inductively we see that s is of class \mathcal{C}^{k+1} if and only if it is of class \mathcal{C}^k and $D^k s$ if of class \mathcal{C}^1 . This formalism is useful for characterizing smoothness of sections on a submanifold with boundary $\bar{X} \subset \Omega$ obtained as the closure of an open submanifold $X \subset \Omega$. Using [Se] and a partition of unity subordinate to an open cover with domains of charts we obtain:

Remark 5.1. A smooth section $s \in \Gamma(X, E)$ extends to a smooth section on \bar{X} if and only if $D^k s \in \Gamma(X, T_\Omega^{*\otimes k} \otimes E)$ extends continuously to \bar{X} for any $k \geq 0$. This condition is equivalent to the existence of a smooth extension of s on a collar neighborhood of \bar{X} in Ω .

Let $X \subset \Omega$ be an open submanifold whose closure \bar{X} is a smooth submanifold with boundary $\partial\bar{X} = \bar{X} \setminus X$ and let $s \in \Gamma(\bar{X}, E)$ be a smooth section defined on \bar{X} . Our problem is to extend s across an open piece $\partial\bar{X} \cap O$ of the boundary using a *given* smooth section σ defined on an open set $O \supset X$ which agrees with s on X (see Fig. 2).




FIGURE 2. A sequence $(u_n)_n$ of O converging to a point $x \in \bar{X} \setminus O$.

Therefore, let $O \subset \Omega$ be a open set containing X and $\sigma \in \Gamma(O, E)$ be a smooth section such that

$$\sigma|_X = s|_X. \quad (9)$$

The closure of X in O is $\bar{X} \cap O$, so the assumption (9) implies

$$\sigma|_{\bar{X} \cap O} = s|_{\bar{X} \cap O}. \quad (10)$$

Formula (10) suggests that we can glue together s and σ to obtain a section on the union $\bar{X} \cup O$ which agrees with s on \bar{X} and with σ on O . The problem is that

Difficulty. The section $s \vee \sigma : \bar{X} \cup O \rightarrow E$ obtained in this way might not even be continuous. Continuity on O (which is open, and on which $s \vee \sigma$ coincides with σ) is obvious, but not at the points of $\bar{X} \setminus O$.

Remark 5.2. The section $s \vee \sigma$ is continuous if and only if the triple (s, O, σ) satisfies the property:

P₀. For every sequence $(u_n)_n$ of O which converges to a point $x \in \bar{X} \setminus O$, the limit $\lim_{n \rightarrow \infty} \sigma(u_n)$ exists and coincides with $s(x)$ (see Fig. 2).

Indeed, if **P₀** holds, then for any sequence $(u_n)_n$ of $\bar{X} \cup O$ converging to a point $x \in \bar{X} \setminus O$, the sequence $((s \vee \sigma)(u_n))_n$ admits a subsequence converging to $(s \vee \sigma)(x)$, which implies continuity at x .

Our goal is to construct a *smooth*, not only continuous, extension of s across $\partial X \cap O$ using σ . Note that, by Remark 5.1, the assumption (9) also implies

$$D^k \sigma|_{\bar{X} \cap O} = D^k s|_{\bar{X} \cap O} \quad \forall k \geq 0, \quad (11)$$

which suggests that a smoothness criterion for $s \vee \sigma$ (similar to Remark 5.2) should hold. The problem is that, in general, the union $\bar{X} \cup O$ has no natural manifold structure, so smoothness is not defined for sections on this set. We are interested in extensions of s to a larger submanifold with boundary $\bar{Y} \supset \bar{X}$ which interpolates between \bar{X} and O in the following sense:

Definition 5.3. Let $Y \subset O$ be an open submanifold whose closure \bar{Y} is a smooth submanifold with boundary. We say that \bar{Y} interpolates between \bar{X} and O if

$$X \subset \bar{X} \cap O \subset Y \subset \bar{Y} \subset \bar{X} \cup O. \quad (12)$$




FIGURE 3. A submanifold with boundary interpolating between \bar{X} and O .

Note that if Y has this property, then

$$\bar{X} \cap O = \bar{X} \cap (\bar{X} \cap O) \subset \bar{X} \cap Y \subset \bar{X} \cap O,$$

so $\bar{X} \cap Y = \bar{X} \cap O$, i.e. Y and O define the same relatively open subset of \bar{X} , so the same open piece $\partial \bar{X} \cap Y = \partial \bar{X} \cap O$ of the boundary.

Taking into account Remark 5.1, we obtain:

Remark 5.4. The restriction of $s \vee \sigma$ to any submanifold with boundary \bar{Y} interpolating between \bar{X} and O is smooth provided for any $k \in \mathbb{N}$ the triple (s, O, σ) satisfies the property

P_k. For every sequence $(u_n)_n$ of O which converges to a point $x \in \bar{X} \setminus O$, the limit $\lim_{n \rightarrow \infty} (D^k \sigma)(u_n)$ exists and coincides with $(D^k s)(x)$.

The goal of this section is the following lemma, which is essentially due to Mauro Nacinovich [Nac]. Nacinovich's result concerns the extension problem for almost complex structures considered in [HiNa]; this problem can be handled taking $E = \text{End}(T_\Omega)$ in our formalism. The method of proof in the general case is the same.

The lemma states that, replacing O by a smaller open set O' with $\bar{X} \cap O' = \bar{X} \cap O$, the triple $(s, O', \sigma|_{O'})$ will satisfy **P_k** for any k , so Remark 5.4 applies, giving a smooth extension of s to \bar{Y} for any submanifold with boundary \bar{Y} interpolating between \bar{X} and O' . Note that, for any such \bar{Y} we'll have $\bar{X} \cap Y = \bar{X} \cap O' = \bar{X} \cap O$, so we obtain extensions across the original open piece $\partial \bar{X} \cap O$ of the boundary.

Lemma 5.5. Let $X \subset \Omega$ be an open submanifold whose closure \bar{X} is a smooth submanifold with boundary and let $s \in \Gamma(\bar{X}, E)$ be a smooth section defined on \bar{X} .

Let $O \subset \Omega$ be an open submanifold containing X and $\sigma \in \Gamma(O, E)$ be a smooth section such that $\sigma|_X = s|_X$.

There exists an open subset $O' \subset O$ such that $\bar{X} \cap O' = \bar{X} \cap O$ and the triple $(s, O', \sigma|_{O'})$ satisfies **P_k** for any $k \geq 0$.

Proof. Let N be the normal line bundle of $\partial \bar{X}$, let

$$\begin{array}{ccc} N & & \\ \nu \downarrow & \searrow f & \\ \partial \bar{X} & \hookrightarrow & \Omega \end{array}$$

be a tubular neighborhood of $\partial \bar{X}$ in Ω [Hir, Section 4.5] and $N_+ \subset N$ be the outer side of the normal bundle. Replacing Ω by $\bar{X} \cup f(N_+)$ and O by its intersection with $\bar{X} \cup f(N_+)$, we may suppose that there exists a continuous retraction $r : \Omega \rightarrow \bar{X}$.

Fix a Riemannian metric g on Ω and define $\delta : \Omega \rightarrow \mathbb{R}$ by $\delta(\omega) := d_g(\omega, \bar{X} \setminus O)$, where d_g is the distance associated with g . It is a well known general fact (valid in the general framework of metric spaces) that such a function is 1-Lipschitz, so continuous. Moreover, since $\bar{X} \setminus O$ is closed, $\delta^{-1}(0) = \bar{X} \setminus O$.

For any $r > 0$ the set $V_r := \delta^{-1}([0, r])$ is an open neighborhood of $\bar{X} \setminus O$ and $\bigcap_{r>0} V_r = \bar{X} \setminus O$. Let $(\rho_n)_n, (r_n)_n$ be decreasing sequences of positive numbers converging to 0 such that $\forall n \in \mathbb{N}, \rho_n < r_n$. For any $n \in \mathbb{N}$ let χ_n be a smooth $[0, 1]$ -valued function on Ω such that

$$V_{\rho_n} \subset \chi_n^{-1}(1) \subset \text{supp}(\chi_n) \subset V_{r_n}.$$

Let d_k be any metric on the total space of the bundle $T_\Omega^{*\otimes k} \otimes E$ which is compatible with its topology. A natural way to obtain such a metric is to choose the connections ∇, ∇_E compatible with g and an Euclidian structure h on E respectively, and to note that these data endow the total space of $T_\Omega^{*\otimes k} \otimes E$ with a Riemannian metric \mathbf{g}_k which makes the projection $T_\Omega^{*\otimes k} \otimes E \rightarrow \Omega$ a Riemannian submersion and the inclusions $T_u^{*\otimes k} \otimes E_u \hookrightarrow T_\Omega^{*\otimes k} \otimes E$ isometric embeddings.

Note first that the sum

$$\sum_{k \geq 0} \chi_k(u) d_k(D^k \sigma(u), D^k s(r(u))) \quad (13)$$

is locally finite on O ; indeed, for a point $v \in O$, we have $\delta(v) > 0$ and the function χ_k vanishes on $\delta^{-1}(\frac{1}{2}\delta(v), +\infty)$ (which is an open neighborhood of v) for any $k \in \mathbb{N}$ for which $r_k < \frac{1}{2}\delta(v)$. Since r is continuous, it follows that (13) defines a continuous

function on O , so the subset

$$O' := \{u \in O \mid \sum_{k \geq 0} \chi_k(u) d_k(D^k \sigma(u), D^k s(r(u))) < \delta(u)\} \quad (14)$$

of O is open. For a point $x \in \bar{X} \cap O$ we have $r(x) = x$ and $D^k \sigma(x) = D^k s(x)$ for any $k \geq 0$ by (11), so $x \in O'$. Therefore $\bar{X} \cap O = \bar{X} \cap O'$ as claimed.

We now prove that the triple $(s, O', \sigma|_{O'})$ satisfies \mathbf{P}_κ for any $\kappa \geq 0$. Fix $\kappa \geq 0$ and let $(u_n)_n$ be a sequence of O' such that $\lim_{n \rightarrow \infty} u_n = x \in \bar{X} \setminus O = \bar{X} \setminus O'$. Since $\lim_{n \rightarrow \infty} u_n \in \bar{X} \setminus O$, it follows $\lim_{n \rightarrow \infty} \delta(u_n) = 0$, so there exists $n_\kappa \in \mathbb{N}$ such that $\delta(u_n) < \rho_\kappa$, hence $\chi_\kappa(u_n) = 1$, for any $n \geq n_\kappa$. Since $u_n \in O'$, formula (14) shows that

$$\forall n \geq n_\kappa, d_\kappa(D^\kappa \sigma(u_n), D^\kappa s(r(u_n))) < \delta(u_n),$$

so

$$\lim_{n \rightarrow \infty} d_\kappa(D^\kappa \sigma(u_n), D^\kappa s(r(u_n))) = 0. \quad (15)$$

But $\lim_{n \rightarrow \infty} u_n = x$ implies $\lim_{n \rightarrow \infty} r(u_n) = r(x) = x$, so

$$\lim_{n \rightarrow \infty} D^\kappa s(r(u_n)) = D^\kappa s(x), \quad (16)$$

because $D^\kappa s$ is continuous on \bar{X} . By (15), (16) we have $\lim_{n \rightarrow \infty} D^\kappa \sigma(u_n) = D^\kappa s(x)$, as claimed. ■

REFERENCES

- [AH] A. Andreotti, C. D. Hill, E. E. Levi convexity and the Hans Lewy problem. Part I : reduction to vanishing theorems, *Annali della Scuola Normale Superiore di Pisa, Classe di Scienze* 3e série, tome 26, no 2 (1972), 325-363.
- [At] M. Atiyah, Complex analytic connections in fibre bundles, *Trans. Amer. Math. Soc.* 85 (1957), 181-207
- [Ca1] D. Catlin, A Newlander-Nirenberg theorem for manifolds with boundary, *Mich. Math. J.*, 35 (1988), 233-240.
- [Ca2] D. Catlin, Extension of CR structures, *Proceedings of Symposia in Pure Mathematics*, Vol. 52 (1991), part 3, 27-34.
- [Do] S. Donaldson, Boundary value problems for Yang-Mills fields, *Journal of Geometry and Physics* 8 (1992) 89-122.
- [Fo] G. B. Folland, The tangential Cauchy-Riemann complex on spheres, *Transactions of the AMS*, Vol. 171 (1972), 83-133.
- [FK] G. B. Folland, J. J. Kohn, The Neumann Problem for the Cauchy-Riemann Complex, *Annals of Mathematics Studies* 75, Princeton University Press, Year (1972).
- [For] F. Forstnerič, *Stein Manifolds and Holomorphic Mappings – The Homotopy Principle in Complex Analysis*, *Ergebnisse der Mathematik und ihrer Grenzgebiete Vol.56*, Springer, Second edition (2017).
- [Gr] H. Grauert, Analytische Faserungen über holomorph-vollständigen Räumen. *Math. Ann.* 135 (1958), 263-273.
- [Go] R. Godement, *Topologie algébrique et théorie des faisceaux*, Hermann, Paris (1973).
- [Hi] C. D. Hill, What is the notion of a complex manifold with a smooth boundary?, *Prospect in Algebraic Analysis*, Vol. 1 (Kashiwara and Kawai, eds.) Academic Press, New York (1988), 185-201.
- [Hir] M. W. Hirsch, *Differential Topology*, Graduate Texts in Mathematics 33, Springer-Verlag (1991).
- [HiNa] C. Hill, M. Nacinovich, A collar neighborhood theorem for a complex manifold, *Rendiconti del Seminario Matematico della Università di Padova*, tome 91 (1994), 23-30.
- [Hö] L. Hörmander, Differential Operators of Principal Type, *Math. Annalen* 140, (1960) 124-146.
- [NaOh] S. Nakano, T. Ohsawa, Strongly Pseudoconvex Manifolds and Strongly Pseudoconvex Domains, *Publ. RIMS, Kyoto Univ.* 20 (1984), 705-715.
- [Le] H. Lewy, "An example of a smooth linear partial differential equation without solution", *Annals of Mathematics*, Vol. 66, No. 1 (1957), 155-158.

- [Kr] S. G. Krantz, Function theory of several complex variables, second edition, AMS Chelsea Publishing, AMS, Providence, Rhode Island (1992).
- [Ma] Y. Matsumoto, An Introduction to Morse Theory, Translations of Mathematical Monographs, Vol. 208, AMS (2002).
- [Nac] M. Nacinovich, private email with detailed proof, March 13, 2022.
- [Oh] T. Ohsawa, Holomorphic embedding of compact s.p.c. manifolds into complex manifolds as real hypersurfaces, in Differential Geometry of Submanifolds, Lecture Notes in Math., Vol. 1090. Springer-Verlag, Berlin and New York (1984), 64-76.
- [Se] R. T. Seeley, Extension of C^∞ functions defined on the half space, Proc. Amer. Math. Soc. 15 (1964), 625-626.
- [Vo] J. Voigt, A Course on Topological Vector Spaces, Compact Textbooks in Mathematics, Birkhäuser (2020).
- [Xi] Z. Xi, Hermitian–Einstein metrics on holomorphic vector bundles over Hermitian manifolds, Journal of Geometry and Physics 53 (2005) 315-335.

AIX MARSEILLE UNIV, CNRS, I2M, MARSEILLE, FRANCE.
Email address: `andrei.teleman@univ-amu.fr`