

Nodal sets of eigenfunctions of sub-Laplacians

Suresh Eswarathasan, Cyril Letrouit

▶ To cite this version:

Suresh Eswarathasan, Cyril Letrouit. Nodal sets of eigenfunctions of sub-Laplacians. 2023. hal-03619388v3

HAL Id: hal-03619388 https://hal.science/hal-03619388v3

Preprint submitted on 15 Sep 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Nodal sets of eigenfunctions of sub-Laplacians

Suresh ESWARATHASAN^{*} and Cyril LETROUIT[†]

September 9, 2023

Abstract

Nodal sets of eigenfunctions of elliptic operators on compact manifolds have been studied extensively over the past decades. In this note, we initiate the study of nodal sets of eigenfunctions of hypoelliptic operators on compact manifolds, focusing on sub-Laplacians. A standard example is the sum of squares of bracket-generating vector fields on compact quotients of the Heisenberg group. Our results show that nodal sets behave in an anisotropic way which can be analyzed with standard tools from sub-Riemannian geometry such as sub-Riemannian dilations, nilpotent approximation and desingularization at singular points. Furthermore, we provide a simple example demonstrating that for sub-Laplacians, the Hausdorff measure of nodal sets of eigenfunctions cannot be bounded above by $\sqrt{\lambda}$, which is the bound conjectured by Yau for Laplace-Beltrami operators on smooth manifolds.

1 Introduction and main results

1.1 Eigenfunctions and nodal sets

Let Ω be a bounded open subset of \mathbb{R}^N and let Δ be the (non-positive) Laplacian on Ω with Dirichlet boundary conditions on $\partial \Omega$. It has a compact resolvent, and the spectrum of $-\Delta$, denoted by

$$0 \leqslant \lambda_1 \leqslant \lambda_2 \leqslant \ldots \leqslant \lambda_n \leqslant \ldots \to +\infty,$$

is discrete and tends to $+\infty$. The nodal set $Z_{\varphi_{\lambda}} = \varphi_{\lambda}^{-1}(0)$ of a mode φ_{λ} satisfying $-\Delta \varphi_{\lambda} = \lambda \varphi_{\lambda}$ is physically interpreted as the set of nodes of the vibration profile φ_{λ} of a drum of shape Ω , and it can be observed, when N = 2, by pouring sand on a vibrating drum of shape Ω (the famous Chladni experiment).

On the mathematical side, nodal sets of eigenfunctions on Euclidean domains and Riemannian manifolds have been studied extensively over the past decades. Let us only mention Courant's nodal domain theorem and Pleijel's asymptotic bound on the number of nodal domains, Donnelly and Fefferman's proof of Yau's conjecture in the analytic setting, and the recent advances by Logunov and Malinnikova concerning the smooth case of the Yau conjecture. For all these results and a global overview of the subject we refer the reader to [19] and the references therein. A stream of results has also emerged in the past twenty years regarding nodal sets of random waves (i.e., random linear combinations of eigenfunctions); for this we refer the reader to the recent survey [27] and the references therein.

In the present paper, we initiate the study of nodal sets of eigenfunctions for operators more general than Laplacians, specifically a certain class of *non-elliptic* operators called sub-Laplacians. Sub-Riemannian geometry is a natural extension of Riemannian geometry whose tools are suited to the analysis of sub-Laplacians.

^{*}Dalhousie University, Halifax NS B3H 4R2 (sr766936@dal.ca)

[†]Massachussets Institute of Technology, Cambridge MA 02139 (letrouit@mit.edu)

1.2 Sub-Laplacians

We start with the general definition of sub-Laplacians, and give immediately after concrete examples of sub-Laplacians.

Let $N \in \mathbb{N}^* = \mathbb{N} \setminus \{0\}$ and let M be either \mathbb{R}^N or a smooth, connected, compact manifold of dimension N without boundary, endowed with a smooth volume μ . Let X_1, \ldots, X_m be smooth vector fields on M satisfying Hörmander's bracket-generating condition:

The vector fields X_1, \ldots, X_m and their iterated Lie brackets $[X_i, X_j], [X_i, [X_j, X_k]]$, etc. span the tangent space $T_x M$ at every point $x \in M$. (1.1)

In particular, X_1, \ldots, X_m are not assumed to span TM. The sub-Laplacian Δ is then defined as

$$\Delta = -\sum_{i=1}^{m} X_i^* X_i \tag{1.2}$$

where $X_i^* = -X_i - \operatorname{div}_{\mu}(X_i)$ is the adjoint of X_i in $L^2(M, \mu)$. Sub-Laplacians are in general not elliptic since X_1, \ldots, X_m are not assumed to span TM.

Instead of the usual elliptic estimates, sub-Laplacians satisfy subelliptic estimates of the form¹ (see [13, Inequality (3.4)], [17, Theorem 1.5])

$$\|u\|_{H^{2/r}(M)} \leqslant C(\|u\|_{L^{2}(M)} + \|\Delta u\|_{L^{2}(M)}), \tag{1.3}$$

where $r \in \mathbb{N}^*$ is called the step of the sub-Laplacian Δ and will be defined later. We only mention that when X_1, \ldots, X_m span TM, then r = 1 and we recover usual elliptic estimates. In turn the estimate (1.3) implies the following form of regularity, called hypoellipticity (see [13]): if $\Delta u \in C^{\infty}(U)$ for some open set $U \subset M$, then $u \in C^{\infty}(U)$.

Sub-Laplacians are a natural generalization of Euclidean Laplacians and of the Laplace-Beltrami operator in Riemannian geometry.² They have been studied extensively since the 1960's and Hörmander's seminal work [13]. Let us illustrate the definition with several examples.

Example 1.1. On $M = (-1, 1)_x \times \mathbb{T}_y$ (where $\mathbb{T} = \mathbb{R}/2\pi\mathbb{Z}$), we set

$$\Delta_{BG} = \partial_x^2 + x^2 \partial_y^2.$$

This sub-Laplacian is the so-called Baouendi-Grushin operator. In this case, $X_1 = \partial_x$, $X_2 = x\partial_y$ and μ is the Lebesgue measure dxdy. Then Δ_{BG} is elliptic except along the singular line x = 0, where $X_2 = 0$. Along this line, $X_1 = \partial_x$ and $[X_1, X_2] = \partial_y$ span the tangent space, so that (1.1) is satisfied.

More generally, for any $\alpha \in \mathbb{N}^*$ we can consider the sub-Laplacian $\partial_x^2 + x^{2\alpha}\partial_y^2$, which is also elliptic except along $\{x = 0\}$. In this case $X_1 = \partial_x$, $X_2 = x^{\alpha}\partial_y$, and the bracket $[X_1, [X_1, \ldots, [X_1, X_2] \ldots]$ where X_1 appears α times generates the missing direction ∂_y .

¹ if $M = \mathbb{R}^N$, for any compact set $K \subset \mathbb{R}^N$ there exists a constant C such that (1.3) holds for any smooth function supported in K.

²To see that the Laplace-Beltrami operator on a Riemannian manifold (M, g) is a sub-Laplacian, take μ to be the Riemannian volume and take a partition of unity $1 = \sum_{k=0}^{K-1} \chi_k(x)^2$ where χ_k is smooth and supported in a chart where g admits an orthonormal frame (Y_1^k, \ldots, Y_N^k) . Then set m = KN and $X_{Nk+j} = \chi_k Y_j^k$ for $j \in \{1, \ldots, N\}$ and $0 \leq k \leq K - 1$. One can check that with this construction $\sum_{i=1}^m X_i^* X_i = -\Delta_g$.

Example 1.2. Sub-Laplacians can be defined on real Heisenberg groups in a natural way. Given $d \in \mathbb{N}^*$, we denote by \mathbf{H}_d the Heisenberg group of dimension 2d + 1, i.e., \mathbb{R}^{2d+1} endowed with the group law

$$(x, y, z) \cdot (x', y', z') := (x + x', y + y', z + z' - \sum_{j=1}^{a} x_j y'_j),$$
(1.4)

where $x, y, x', y' \in \mathbb{R}^d$ and $z, z' \in \mathbb{R}$. When d = 1, this group law comes from the representation of \mathbf{H}_1 as the group of 3×3 matrices of the form

$$\begin{pmatrix}
1 & x & -z \\
0 & 1 & y \\
0 & 0 & 1
\end{pmatrix}$$

for $x, y, z \in \mathbb{R}$, endowed with the usual product of matrices (the higher-dimensional case has a similar representation).

The set of left-invariant vector fields on \mathbf{H}_d is generated by ∂_z together with the 2d vector fields

$$X_j = \partial_{x_j}, \quad Y_j = \partial_{y_j} - x_j \partial_z, \qquad for \ j = 1, \dots, d.$$

The Heisenberg group \mathbf{H}_d admits lattices, such as $\Gamma = (\sqrt{2\pi}\mathbb{Z})^{2d} \times 2\pi\mathbb{Z}$. Left-invariant vector fields can be considered as vector fields on the compact left-quotient $\Gamma \setminus \mathbf{H}_d$. We define the sub-Laplacian on $\Gamma \setminus \mathbf{H}_d$ by

$$\Delta_{\Gamma \setminus \boldsymbol{H}_d} = \sum_{j=1}^d X_j^2 + Y_j^2$$

We note that $[X_j, Y_j] = -\partial_z$ for any j, hence (1.1) is verified.

Example 1.3. More generally, sub-Laplacians arise naturally in the setting of Carnot groups, which are a family of nilpotent Lie groups whose simplest non-Euclidean examples are real Heisenberg groups (see Example 1.2). A Carnot group G of step r is a connected, simply connected, finite-dimensional Lie group whose Lie algebra \mathfrak{g} admits a step-r stratification, meaning that there exist nontrivial linear subspaces V_1, \ldots, V_r such that

$$\mathfrak{g} = V_1 \oplus \ldots \oplus V_r$$
, $[V_1, V_i] = V_{i+1}$ for $i = 1, \ldots, r-1$ and $[V_1, V_r] = 0$.

The stratum V_i contains iterated brackets of length *i* between elements of V_1 . Taking a basis of V_1 composed of left-invariant vector fields X_1, \ldots, X_m (where *m* is the dimension of V_1), and taking μ to be the Haar measure on *G*, we obtain a sub-Laplacian thanks to the formula (1.2). Quotienting by an appropriate lattice and using the left-invariance of Δ , we can define sub-Laplacians on compact quotients of *G* (see Example 1.2).

To make this example more concrete, we now explain how G can be identified to (\mathbb{R}^N, \star) for $N = \sum_{i=1}^r \dim(V_i)$ and \star a specific group law. For each $i \in \{1, \ldots, r\}$ we pick a basis of vector fields of V_i , which all together form a basis X_1, \ldots, X_N of \mathfrak{g} . Then the exponential map \exp is a globally defined diffeomorphism from $\mathfrak{g} \cong \mathbb{R}^N$ to G (since G is connected and nilpotent):

$$\exp:(x_1,\ldots,x_N)\mapsto \exp\left(\sum_{i=1}^N x_i X_i\right)\in G.$$

This allows to identify G with \mathbb{R}^N , and the group law \star derives from the Baker-Campbell-Hausdorff formula. Following this procedure with $X_1 = \partial_x$, $X_2 = \partial_y - x\partial_z$ and $X_3 = \partial_z$, the group which is obtained is isomorphic to the Heisenberg group \mathbf{H}_1 of Example 1.2. Besides Heisenberg groups, other examples of Carnot groups include the Engel group, for which r = 3. Our results concern the nodal sets of eigenfunctions of Δ (i.e., the set where an eigenfunction φ vanishes) and the nodal components (i.e., the connected components of $\{\varphi \neq 0\}$). They extend results which are well-known for eigenfunctions of Laplace-Beltrami operators.

1.3 Main results: Courant's theorem and density of the nodal set

Our results address the validity of Courant's nodal domain theorem [9] for eigenfunctions of sub-Laplacians and the $1/\sqrt{\lambda}$ -density of the nodal set with respect to an adapted distance, called the sub-Riemannian distance.

Let $\Omega \subset M$ be a connected open subset, assumed to be bounded and with Lipschitz boundary if $M = \mathbb{R}^N$. We denote by $C_c^{\infty}(\Omega)$ the set of smooth functions whose support is contained in Ω . The operator $\Delta : C_c^{\infty}(\Omega) \to C_c^{\infty}(\Omega)$ is non-positive, symmetric and densely defined in $L^2(\Omega, \mu)$. In the sequel, we denote by $(\Delta_\Omega, \mathcal{D}(\Delta_\Omega))$ its Friedrichs extension (see [23] and Section 2 for reminders). When $\partial\Omega \neq \emptyset$, this naturally enforces Dirichlet boundary conditions.

Proposition 1.4. The selfadjoint operator $(-\Delta_{\Omega}, \mathcal{D}(\Delta_{\Omega}))$ has discrete point spectrum

 $0 \leqslant \lambda_1 \leqslant \lambda_2 \leqslant \ldots \leqslant \lambda_n \leqslant \ldots \to +\infty$

(with repetitions according to multiplicities). There exists an orthonormal basis $\{\varphi_n\}_{n\in\mathbb{N}}$ of $L^2(\Omega,\mu)$ such that for every $n\in\mathbb{N}$, $\varphi_n\in\mathcal{D}(\Delta_\Omega)$ and $-\Delta_\Omega\varphi_n=\lambda_n\varphi_n$.

In our next results, we are going to rely on two assumptions.

Assumption 1.5. We assume that either $\Omega = M$, or the boundary of Ω is smooth and noncharacteristic, meaning that for any $x \in \partial \Omega$, there exists $i \in \{1, ..., m\}$ such that $X_i(x) \notin T_x \partial \Omega$.

Under Assumption 1.5, the eigenfunctions of $-\Delta_{\Omega}$ are smooth up to the boundary $\partial\Omega$ (see [16, Theorem III, point (4)]), which will be important in the proofs.

Assumption 1.6. We assume that one of the following holds:

- 1. The (topological) dimension of M is N = 2;
- 2. or the manifold M, the volume μ and the vector fields X_1, \ldots, X_m are real-analytic.

Under this assumption, it follows from [26, Theorem 1] (in case 1) and [7] (in case 2) that any u satisfying $(\Delta_{\Omega} + \lambda)u = 0$ in Ω for some $\lambda \in \mathbb{R}$ and vanishing in a non-empty open subset $U \subset \Omega$ vanishes in fact everywhere in Ω . Actually, to prove this unique continuation property in case 2, instead of working directly with the operator $\Delta_{\Omega} + \lambda$ which does not satisfy the assumptions of [7], we work with $\partial_t^2 + \Delta_{\Omega}$ instead. We write that $v = ue^{i\lambda t}$ is a solution of the equation $(\partial_t^2 + \Delta_{\Omega})v = 0$ in $\Omega \times \mathbb{R}$, which vanishes in the non-empty open subset $U \times \mathbb{R}$. Hence v vanishes everywhere in $\Omega \times \mathbb{R}$ according to [26, Theorem 1] (in case 1) and [7, Corollary 4.1] (in case 2), and consequently u vanishes everywhere in Ω . This unique continuation property will be used in the proof of the second part of Theorem 1.

We denote by

$$Z_f = \{f = 0\} \subset \overline{\Omega}$$

the zero set, or commonly the *nodal set* of a function f. A *nodal domain* of f is a connected component of $\Omega \setminus Z_f$. We are in position to state an analogue of the Courant nodal domain theorem for sub-Laplacians:

Theorem 1. Under Assumption 1.5, for any $n \in \mathbb{N}$, any eigenfunction of $-\Delta_{\Omega}$ with eigenvalue λ_n has at most $n + mult(\lambda_n) - 1$ nodal domains, where $mult(\lambda_n)$ denotes the multiplicity of λ_n . If moreover Assumption 1.6 is satisfied, a stronger bound holds: the number of nodal domains of any eigenfunction with eigenvalue λ_n is bounded above by n.

For our second result we introduce the sub-Riemannian metric: it is defined for $q \in M$ and $v \in T_q M$ as

$$g_q(v) = \inf\left\{\sum_{i=1}^m u_i^2 \mid v = \sum_{i=1}^m u_i X_i\right\}.$$
 (1.5)

This metric is finite if and only if $v \in \mathcal{D} = \text{Span}(X_1, \ldots, X_m)$. It induces a notion of distance $d: M \times M \to \mathbb{R}_+$, and the distance between two points is always finite thanks to the bracketgenerating condition (1.1) (due to the Chow-Rashevsky theorem [5, Theorem 2.4]). The sub-Riemannian balls are then defined as

$$B_{\varepsilon}(q) = \left\{ q' \in M, \ d(q,q') < \varepsilon \right\}$$
(1.6)

for $q \in M$ and $\varepsilon > 0$.

Theorem 2. Under Assumption 1.5, there exists C > 0 depending only on Ω such that for any $\lambda \in \mathbb{R}$ and any eigenfunction φ_{λ} with eigenvalue λ , the nodal set $Z_{\varphi_{\lambda}}$ intersects any sub-Riemannian ball of radius greater than $C\lambda^{-1/2}$.

Theorem 2 is illustrated by a simple example in Section 5.3 below.

An important distinguishing factor of sub-Riemannian geometry from Riemannian geometry is that sub-Riemannian balls are *anisotropic* whilst Riemannian balls are isotropic. The ball-box theorem captures this difference in more precise language (see [5, Corollary 7.35]): it states that at any point $q \in \Omega$, there exist a system of privileged coordinates $\psi_q = (x_1, \ldots, x_N) : U \to \mathbb{R}^N$ defined in a neighborhood U of q, positive integers w_1, \ldots, w_N , and constants $C_q, \varepsilon_q > 0$ such that for any $\varepsilon < \varepsilon_q$, the sub-Riemannian ball $B_{\varepsilon}(q)$ verifies

$$\operatorname{Box}(C_q^{-1}\varepsilon) \subset (\psi_q)_*(B_\varepsilon(q)) \subset \operatorname{Box}(C_q\varepsilon)$$
(1.7)

where

$$Box(\varepsilon) = [-\varepsilon^{w_1}, \varepsilon^{w_1}]_{x_1} \times \ldots \times [-\varepsilon^{w_N}, \varepsilon^{w_N}]_{x_N} \subset \mathbb{R}^N$$
(1.8)

(note that C_q does not depend on ε). Sub-Riemannian balls can thus be approximated by Euclidean rectangles in an appropriate coordinate system, with side-lengths scaling differently in each direction. Section 5.3's example serves as a useful illustrator of the ball-box theorem.

As a consequence, roughly speaking, if Ω is some subset of \mathbb{R}^N , starting from a fixed point $q \in \Omega$ and following a given direction $\vec{\ell}$, one expects to cross the nodal set many more times in some directions $\vec{\ell}$ than some others, when the eigenvalue is large. This phenomenon is also explicitly expressed by Section 5.3's example.

1.4 Yau-type bounds for sub-Laplacians

In this section we explain that Yau's conjecture, unless properly modified, is not true for sub-Laplacians. Yau conjectured that for any smooth closed Riemannian manifold (M,g) of dimension N, there exist constants c, C > 0 such that for any eigenfunction φ_{λ} of the associated Laplace-Beltrami operator $-\Delta_g$ with eigenvalue λ ,

$$c\sqrt{\lambda} \leqslant \mathscr{H}^{N-1}(Z_{\varphi_{\lambda}}) \leqslant C\sqrt{\lambda}.$$
(1.9)

where \mathscr{H}^{N-1} is the (N-1)-dimensional Hausdorff measure. This conjecture has been proved in the real-analytic setting by Donnelly and Fefferman [10], but it is open in the case of general smooth manifolds.

In sub-Riemannian geometry, it is known, under the condition that the w_i in (1.8) are independent of q^{-3} , that the Hausdorff dimension with respect to the sub-Riemannian metric (1.5) of M is equal to

$$Q = \sum_{i=1}^{N} w_i \tag{1.10}$$

(see [20]), which is strictly greater than N as soon as the vector fields X_1, \ldots, X_m do not span TM. For instance, in the Heisenberg group \mathbf{H}_1 of Example 1.2, there holds N = 3, $w_1 = 1$, $w_2 = 1$, $w_3 = 2$, and hence $\mathcal{Q} = 4$. It is also known that hypersurfaces have Hausdorff dimension $\mathcal{Q} - 1$ with respect to the sub-Riemannian metric (1.5) (see [12, Section 0.6.C]).

The following result shows that Yau-type bounds like (1.9) (with N replaced by Q) do not hold for sub-Laplacians: in general we do not expect better bounds than

$$c\sqrt{\lambda} \leqslant \mathscr{H}^{\mathcal{Q}-1}(Z_{\varphi_{\lambda}}) \leqslant C\lambda^{r/2}$$

where the step r is defined in Section 4.1 below and is equal to 2 in the Heisenberg case. Here and in the sequel, \mathscr{H}^{K} denotes the Hausdorff measure of dimension K.

Theorem 3. Let $M = \Gamma \setminus \mathbf{H}_1$ as in Example 1.2, endowed with the Lebesgue measure and the vector fields $X_1 = \partial_{x_1}$ and $Y_1 = \partial_{y_1} - x_1 \partial_{z_1}$. Let \mathscr{H}^3 denote the Hausdorff measure of dimension 3 with respect to the associated sub-Riemannian metric on M. There exist an open subset $\Omega \subset M$, constants $c_1, c_2 > 0$ and sequences of eigenfunctions $(\varphi_{1,m})_{m \in \mathbb{N}}$ and $(\varphi_{2,m})_{m \in \mathbb{N}}$ of $-\Delta_{\Omega}$ with respective eigenvalues $(\lambda_{1,m})_{m \in \mathbb{N}}$ and $(\lambda_{2,m})_{m \in \mathbb{N}}$ tending to $+\infty$, such that

$$\mathscr{H}^{3}(Z_{\varphi_{1,m}}) \leqslant c_{1}\sqrt{\lambda_{1,m}} \tag{1.11}$$

and

$$\mathscr{H}^3(Z_{\varphi_{2,m}}) \geqslant c_2 \lambda_{2,m}.$$
(1.12)

The fact that the two bounds in our result are not of the same order is due to the fact that some sequences of eigenfunctions oscillate much more in the directions needing brackets to be generated (like ∂_z in the Heisenberg group) than what is possible for Laplacians. This demonstrates the need for a reformulation of Yau's Conjecture.

1.5 Open questions

1.5.1 Pleijel bound

The Courant bound for Laplace-Beltrami operators is known to be non-optimal as the eigenvalue tends to $+\infty$: the Pleijel bound asserts that when *n* becomes large, the number of nodal domains of an eigenfunction with eigenvalue λ_n is at most *cn* for some explicit constant c < 1 (see [21] for the case of 2-dimensional Euclidean domains, and [6]). An analogous result for sub-Laplacians would require new ideas since the appropriate Faber-Krahn inequality that one uses in the usual proof of the Pleijel bound is not known.

 $^{^{3}}$ this is equivalent to the condition that the sub-Riemannian flag is equiregular, see Section 4.1 for definitions.

1.5.2 Relation between elliptic and subelliptic bounds

In the analytic case, Yau's conjectured bound (1.9) for the Laplace-Beltrami operator is known to hold (see [10]). If we consider a family of Laplace-Beltrami operators of the form

$$\Delta_{g_{\varepsilon}} = \Delta_{\mathrm{sR}} + \varepsilon^2 \Delta_h \tag{1.13}$$

where Δ_{sR} is a fixed analytic sub-Laplacian and Δ_h is a fixed analytic Laplace-Beltrami operator (defined on the same domain of \mathbb{R}^N), how do the constants c and C in (1.9) behave as $\varepsilon \to 0$? This question is motivated by the fact that the "sub-Riemannian limit" of operators (1.13) has already attracted attention in other related contexts, see for instance [25].

1.6 Organization of the paper and new contributions

The proofs of our results follow the same broad strokes as those for elliptic Laplacians. However, to the best of our knowledge, it is the novel application of tools and techniques unique to sub-Riemannian geometry and hypoelliptic equations within the subject of nodal geometry that is our main contribution. We elaborate on this in our outline.

In Section 2, we prove Proposition 1.4: our proof mostly relies on the compactness of the resolvent of sub-Laplacians, which follows from well-known subelliptic estimates due to Hörmander. In Section 3 we prove Theorem 1 using the usual strategy for proving Couranttype bounds, and a combination of a robust argument by Colette Anné [3] with subelliptic estimates. In Section 4, we gather tools coming from sub-Riemannian geometry, namely the nilpotent approximation and the desingularization procedure. In Section 5, we prove Theorem 2 using the aforementioned tools as basic building blocks for estimating the first eigenvalue of a sub-Laplacian in a small ball. Finally, in Section 6 we prove Theorem 3 by constructing a sub-Laplacian and explicit examples of eigenfunctions.

Acknowledgments. We thank Hajer Bahouri, Yves Colin de Verdière, Valentina Franceschi, Bernard Helffer, Thomas Letendre, Eugenia Malinnikova, Iosif Polterovich and Luca Rizzi for interesting discussions. The first author is supported by an NSERC Discovery Grant and the second author is supported by the Simons Foundation Grant 601948, DJ.

2 Proof of Proposition 1.4

We first recall briefly the classical Friedrichs extension construction ([23]). We denote by q_{Ω} the quadratic form on $C_c^{\infty}(\Omega)$ given by $q_{\Omega}(v,w) = (\Delta v,w)$ where (\cdot,\cdot) denotes the $L^2(\Omega,\mu)$ scalar product. It is closable and we denote by \hat{q}_{Ω} its closure. Explicitly, denoting by H the Hilbert space completion of $C_c^{\infty}(\Omega)$ with respect to the scalar product $(v,w)_H = (v,w) + q_{\Omega}(v,w)$, the inclusion map $\iota : C_c^{\infty}(\Omega) \hookrightarrow L^2(\Omega,\mu)$ extends by continuity to a linear map $\hat{\iota} : H \to L^2(\Omega,\mu)$. The quadratic form q_{Ω} also extends by continuity to a quadratic form \hat{q}_{Ω} over H, so that if \bar{v} and \bar{w} denote the equivalence classes of $\{v_n\}$ and $\{w_n\}$ in H, then $\hat{q}(\bar{v},\bar{w}) = \lim_{n\to\infty} q_{\Omega}(v_n,w_n)$. One can check that $\hat{\iota}$ is injective, hence \hat{q}_{Ω} can be seen as a quadratic form on $L^2(\Omega,\mu)$, with domain $\mathcal{D}(\hat{q}_{\Omega}) = \hat{\iota}(H)$. More concretely, the domain $\mathcal{D}(\hat{q}_{\Omega})$ consists of those $v \in L^2(\Omega,\mu)$ such that there exists $\{v_n\} \subset C_c^{\infty}(\Omega)$ such that $v_n \to v$ in $L^2(\Omega,\mu)$ and $q_{\Omega}(v_n - v_\ell) \to 0$ as $\ell, n \to \infty$.

Then, the Friedrichs extension of $(\Delta, C_c^{\infty}(\Omega))$ is the operator $(\Delta_{\Omega}, \mathcal{D}(\Delta_{\Omega}))$ where

$$\mathcal{D}(\Delta_{\Omega}) = \{ v \in \mathcal{D}(\widehat{q}_{\Omega}) : \widehat{q}_{\Omega}(v, \cdot) \text{ is } L^{2}(\Omega, \mu) - \text{continuous} \}$$
(2.1)

and $\Delta_{\Omega} v \in L^2(\Omega, \mu)$ is defined through the Riesz representation theorem by the relation $(\Delta_{\Omega} v, w) = \widehat{q}_{\Omega}(v, w)$ for any $w \in \mathcal{D}(\widehat{q}_{\Omega})$ (note that $\mathcal{D}(\widehat{q}_{\Omega})$ is dense in $L^2(\Omega, \mu)$). By the

same procedure, for any bounded open set $U \subset M$ (including the cases of U = M and $U = \Omega$) we obtain the Friedrichs extension $(\Delta_U, \mathcal{D}(\Delta_U))$ of the non-positive symmetric operator $\Delta : C_c^{\infty}(U) \to C_c^{\infty}(U)$ (densely defined on $L^2(U, \mu)$).

In the next lemma, for any $v \in \mathcal{D}(\widehat{q}_{\Omega})$, we denote by Ev its extension by 0 in $M \setminus \Omega$.

Lemma 2.1. Let $v \in \mathcal{D}(\widehat{q}_{\Omega})$. Then $Ev \in \mathcal{D}(\widehat{q}_M)$ and $\widehat{q}_M(Ev) = \widehat{q}_{\Omega}(v)$.

Proof. Let $v_n \in C_c^{\infty}(\Omega)$ such that $v_n \to v$ in $L^2(\Omega, \mu)$ and $q_{\Omega}(v_n - v_\ell) \to 0$ as $n, \ell \to \infty$. We have $Ev_n \to Ev$ in $L^2(M, \mu)$ and

$$q_M(Ev_n - Ev_\ell) = (\Delta(Ev_n - Ev_\ell), Ev_n - Ev_\ell) = (\Delta(v_n - v_\ell), v_n - v_\ell) = q_\Omega(v_n - v_\ell) \xrightarrow[n,\ell \to \infty]{} 0.$$

Thus $Ev \in \mathcal{D}(\widehat{q}_M)$ and $\widehat{q}_M(Ev) = \widehat{q}_{\Omega}(v)$.

We start the proof of Proposition 1.4 with the following particular case of the subelliptic estimate [24, Estimate (17.20)]: whenever a and b are in $C_c^{\infty}(M)$ with sufficiently small support and b = 1 on the support of a, there exist s > 0, C > 0 such that

$$||au||_{H^s(M)} \leq C \sum_{j=1}^m (||bX_ju||_{L^2(M)} + ||bu||_{L^2(M)})$$

Using a partition of unity and the fact that commutators of X_j with smooth cutoff functions are multiplication operators, we can globalize this inequality: there exist s > 0, C > 0 such that

$$\|u\|_{H^{s}(M)} \leq C \sum_{j=1}^{m} (\|X_{j}u\|_{L^{2}(M)} + \|u\|_{L^{2}(M)}).$$

$$(2.2)$$

Squaring this inequality, we obtain $||u||_{H^s(M)}^2 \leq C(-\Delta_M u + u, u)_{L^2(M)}$, which together with the Cauchy-Schwarz inequality implies that

$$||u||_{H^s(M)} \leq C ||(\mathrm{Id} - \Delta_M)u||_{L^2(M)}.$$
 (2.3)

Case 1: *M* is compact. In this case, the Rellich–Kondrachov theorem gives that the resolvent $(\operatorname{Id} - \overline{\Delta}_M)^{-1}$ is compact from $L^2(M)$ to $L^2(M)$. Using [22, Theorem XIII.64 p.245], this implies that $\mu_n(\Delta_M) \to +\infty$ where

$$\mu_n(\Delta_M) = \inf_{\substack{W \subset \mathcal{D}(\widehat{q}_M) \\ \dim(W) = n}} \max_{\substack{v \in W \\ \|v\| = 1}} \widehat{q}_M(v).$$
(2.4)

Note that in [22, Theorem XIII.1 p. 76] the quantity $\mu_n(\Delta_M)$ is defined differently, but it is well-known that $\mu_n(\Delta_M)$ is in fact also equal to (2.4) (see for instance [18, Theorem 5.37]). This implies thanks to Lemma 2.1 that

$$\mu_n(\Delta_{\Omega}) = \inf_{\substack{W' \subset \mathcal{D}(\widehat{q}_{\Omega}) \\ \dim(W') = n}} \max_{\substack{v \in W' \\ \|v\| = 1}} \widehat{q}_{\Omega}(v)$$

tends to $+\infty$ as $n \to +\infty$: indeed, by extending all elements of $W' \subset \mathcal{D}(\hat{q}_{\Omega})$ by 0 outside Ω , we obtain an *n* dimensional subspace $W \subset \mathcal{D}(\hat{q}_M)$, hence $\mu_n(\Delta_M) \leq \mu_n(\Delta_\Omega)$ for any $n \in \mathbb{N}$. Applying again [22, Theorem XIII.64 p.245], this time to $(\Delta_\Omega, \mathcal{D}(\Delta_\Omega))$, we obtain the existence of a complete orthonormal basis $\{\varphi_n\}_{n=1}^{\infty}$ in $\mathcal{D}(\Delta_\Omega)$ so that $-\Delta_\Omega \varphi_n = \lambda_n \varphi_n$ with $0 \leq \lambda_1 \leq \ldots \leq \lambda_n \leq \ldots \to +\infty$. <u>Case 2:</u> $M = \mathbb{R}^N$. In this case, recall that we assumed in the introduction that Ω is bounded and has Lipschitz boundary. The inequality (2.3) applied to $u \in C_c^{\infty}(\Omega)$, together with the density of $C_c^{\infty}(\Omega)$ in $L^2(\Omega)$, proves that $(\mathrm{Id} - \Delta_{\Omega})^{-1} : L^2(\Omega) \to H^s(\Omega) \cap \mathcal{D}(\Delta_{\Omega})$ is continuous. Since the injection from $H^s(\Omega)$ to $L^2(\Omega)$ is compact (we use here the fact that the boundary of Ω is Lipschitz), we obtain directly that the resolvent is compact, and we conclude by [22, Theorem XIII.64 p.245].

3 Proof of Theorem 1

We start the proof of Theorem 1 with a lemma containing an integration-by-parts formula. We do not give its proof, which follows from the definition of the quadratic form \hat{q}_{Ω} recalled above.

Lemma 3.1. If $u \in \mathcal{D}(\widehat{q}_{\Omega})$, then $X_i u \in L^2(\Omega, \mu)$. Moreover if $u, v \in \mathcal{D}(\widehat{q}_{\Omega})$, then

$$\widehat{q}_{\Omega}(u,v) = -\sum_{i=1}^{m} \int_{\Omega} (X_{i}u)(X_{i}v)d\mu$$

We denote by E_{λ_k} the eigenspace associated to the eigenvalue λ_k of $-\Delta_{\Omega}$. We have the following min-max principle:

Lemma 3.2. 1. $\varphi \in \mathcal{D}(\widehat{q}_{\Omega}) \setminus \{0\}$ belongs to E_{λ_1} if and only if it minimizes over $\mathcal{D}(\widehat{q}_{\Omega}) \setminus \{0\}$ the Rayleigh quotient

$$R(\varphi) = \frac{\sum_{i=1}^{m} \|X_i\varphi\|_{L^2(\Omega,\mu)}^2}{\|\varphi\|_{L^2(\Omega,\mu)}^2}.$$
(3.1)

In this case $R(\varphi) = \lambda_1$.

2. If $\varphi \in \mathcal{D}(\widehat{q}_{\Omega}) \setminus \{0\}$ is orthogonal to $E_{\lambda_1}, \ldots, E_{\lambda_{k-1}}$ and $R(\varphi) = \lambda_k$, then $\varphi \in E_{\lambda_k}$.

The proof is standard and follows for the first point from the computation of $R(\varphi + \varepsilon \psi)$ for $\psi \in \mathcal{D}(\hat{q}_{\Omega})$ and $\varepsilon \to 0$, and for the second point from the decomposition of φ in the orthonormal basis given by Proposition 1.4.

The next two lemmas are classical in the Riemannian setting but their proofs require some care in the present sub-Riemannian (sR) context.

Lemma 3.3. Let $D \subset M$ be a connected open set with $\partial D \neq \emptyset$. Then $\lambda_1(D) > 0$, and there exists an eigenfunction of $-\Delta_D$ with eigenvalue $\lambda_1(D)$ which is non-negative.

Proof. Assume for the sake of a contradiction that $\lambda_1(D) = 0$ and let $u \neq 0$ be an eigenfunction $\Delta u = 0$. Then $(\Delta u, u) = 0$ hence by definition $\hat{q}_D(u, u) = 0$, which implies $||X_i u||_{L^2(D,\mu)} = 0$ for any *i* thanks to Lemma 3.1. But thanks to hypoelliptic regularity [13] we know that $u \in C^{\infty}(D)$ (a priori not up to the boundary if *D* is arbitrary) hence $X_i u \equiv 0$ in *D*. Then $[X_{i_1}, [X_{i_2}, \ldots] \ldots] u \equiv 0$ for any bracket of the vector fields, hence by the Hörmander bracket-generating condition *u* is constant in *D*.

Let us prove that the only constant u which belongs to $\mathcal{D}(\hat{q}_D)$ is 0. We choose $(u_\ell)_{\ell \in \mathbb{N}}$ such that $u_\ell \in C_c^{\infty}(D)$ and $u_\ell \to u$, $X_i u_\ell \to 0$ in $L^2(D)$ as $\ell \to +\infty$. We denote by \underline{u} (resp. \underline{u}_ℓ) the extension of u (resp. u_ℓ) to M by 0 in $M \setminus D$. First, $\underline{u} \in \mathcal{D}(\hat{q}_M)$ according to Lemma 2.1. Let $v \in \mathcal{D}(\hat{q}_M)$, and $v_\ell \in C_c^{\infty}(M)$ such that $v_\ell \to v$ in $L^2(M, \mu)$ and $q_M(v_n - v_\ell) \to 0$ as $\ell, n \to +\infty$. Then

$$\widehat{q}_M(\underline{u}, v) = \lim_{\ell \to +\infty} \sum_{i=1}^m (X_i \underline{u}_\ell, X_i v_\ell)_{L^2(M,\mu)} = \lim_{\ell \to +\infty} \sum_{i=1}^m (X_i u_\ell, X_i v_\ell)_{L^2(\operatorname{supp}(u_\ell),\mu)} = 0$$

since $(X_i v_\ell)_{\ell \in \mathbb{N}}$ is bounded in $L^2(M, \mu)$, and thus in $L^2(\operatorname{supp}(u_\ell), \mu)$. Hence $\underline{u} \in \mathcal{D}(\Delta_M)$ and $\Delta_M \underline{u} = 0$. By hypoellipticity, $\underline{u} \in C^{\infty}(M)$, which is impossible since \underline{u} is not smooth across $\partial D \neq \emptyset$.

Let $u_0 \neq 0$ be in the first eigenspace of Δ_D . Then $|u_0| \in \mathcal{D}(\widehat{q}_D)$ and for any $i \in \{1, \ldots, m\}$,

$$X_{i}|u_{0}| = \begin{cases} X_{i}u_{0} & \text{a.e. in } \{u_{0} > 0\} \\ 0 & \text{a.e. in } \{u_{0} = 0\} \\ -X_{i}u_{0} & \text{a.e. in } \{u_{0} < 0\} \end{cases}$$
(3.2)

Both statements follow from [11, Chapter 5, Exercise 17]: the main steps are to apply the chain rule to $F_{\varepsilon} \circ u_0$ where $F_{\varepsilon}(z) = \sqrt{z^2 + \varepsilon^2} - \varepsilon$ and then use the dominated convergence theorem.

From (3.2) we deduce that $R(|u_0|) = R(u_0)$ where the Rayleigh quotient R is defined in (3.1). According to Lemma 3.2, this implies that $|u_0|$ is also in the first eigenspace of Δ_D . \Box

Lemma 3.4. Let $u \in \mathcal{D}(\Delta_{\Omega})$ satisfying $-\Delta_{\Omega}u = \lambda u$ in Ω . Let D be a nodal domain of u. Then the restriction \underline{u} of u to D belongs to $\mathcal{D}(\Delta_D)$, and it is an eigenfunction of the Dirichlet problem in D, associated to the smallest eigenvalue $\lambda = \lambda_1(D)$.

Proof. Thanks to Assumption 1.5, u is smooth up to the boundary of Ω (see [16, Theorem III, point (4)]), hence \underline{u} is smooth up to the boundary of D. Without loss of generality, we assume that \underline{u} is non-negative. We follow the proof of Lemma 2.0.1 in [3], which does not use any regularity on the boundary of the nodal domain.

We fix a Riemannian metric g_R on M, which induces a distance $\operatorname{dist}_{g_R}$ and a gradient ∇_{g_R} , in order to conveniently conduct our local analysis. Let $\chi_n \in C_c^{\infty}(\overline{D})$ be a cut-off function such that there exists a constant C > 0 independent of n such that

- $\chi_n(x) = 1$ for dist_{g_R} $(x, \partial D) \ge 1/n$
- $\chi_n(x) = 0$ for dist_{*q_R*} $(x, \partial D) \leq 1/(2n)$
- $\|\nabla_{g_R}\chi_n\|_{L^{\infty}} \leq Cn$
- $\|\nabla^2_{q_P}\chi_n\|_{L^{\infty}} \leq Cn^2$

(the existence of χ_n is shown in [3]). Since X_1, \ldots, X_m are smooth, this implies that $||X_i\chi_n||_{L^{\infty}} \leq Cn$ and $||X_i^*\chi_n||_{L^{\infty}} \leq Cn$ for any *i*, and $||\Delta\chi_n||_{L^{\infty}} \leq Cn^2$ (the constant *C* may have changed).

It suffices to show that \underline{u} arises from a Cauchy sequence in $\mathcal{D}(\hat{q}_{\Omega})$ and that \underline{u} generates a continuous linear functional on $L^2(D,\mu)$ via the quadratic form \hat{q}_D . We set $u_n = \chi_n \underline{u}$. We have $u_n \to \underline{u}$ in $L^2(D,\mu)$. Let us prove that $q_D(u_n - u_\ell) \to 0$ as $\ell, n \to \infty$. To simplify notations, we set $\alpha_{n,\ell} = \chi_n - \chi_\ell$. We have

$$\begin{aligned} |q_D(u_n - u_\ell)| &= \int_D \sum_{i=1}^m (X_i(u_n - u_\ell))^2 \, d\mu = \int_D \sum_{i=1}^m \alpha_{n,\ell} u X_i^* ((X_i \alpha_{n,\ell}) u + \alpha_{n,\ell} (X_i u)) d\mu \\ &= -\int_D \alpha_{n,\ell} u^2 \Delta \alpha_{n,\ell} \, d\mu - \int_D \alpha_{n,\ell}^2 u \Delta u \, d\mu \\ &- \int_D \left(\sum_{i=1}^m (X_i \alpha_{n,\ell}) (X_i^* u) \alpha_{n,\ell} u + (X_i^* \alpha_{n,\ell}) (X_i u) \alpha_{n,\ell} u \right) d\mu \end{aligned}$$
(3.3)
$$= I_1 + I_2 + I_3.$$

We show that $I_j \to 0$ for j = 1, 2, 3, a $n, \ell \to +\infty$. We denote by A_n the support of $\nabla \chi_n$, in particular $\operatorname{vol}(A_n) \to 0$. We have

$$\left| \int_{D} \alpha_{n,\ell} u^2 \Delta \alpha_{n,\ell} \, d\mu \right| \leqslant C\ell^2 \int_{A_\ell} u^2 \, d\mu + Cn^2 \int_{A_n} u^2 \, d\mu. \tag{3.4}$$

We use the fact that u is smooth up to the boundary to get that $u^2 \leq C\ell^{-2}$ in A_ℓ and similarly $u^2 \leq Cn^{-2}$ in A_n . Hence

$$|I_1| = \left| \int_D \alpha_{n,\ell} u^2 \Delta \alpha_{n,\ell} \, d\mu \right| \leqslant C(\operatorname{vol}(A_\ell) + \operatorname{vol}(A_n)) \underset{n,\ell \to +\infty}{\longrightarrow} 0.$$

Then, we have

$$\left| \int_{D} \alpha_{n,\ell}^2 u \Delta u \, d\mu \right| = \lambda \int_{D} \alpha_{n,\ell}^2 u^2 \, d\mu \leqslant C(\operatorname{vol}(A_\ell) + \operatorname{vol}(A_n)) \int_{D} u^2 \, d\mu \underset{n,\ell \to +\infty}{\longrightarrow} 0$$

One can also check that

$$I_{3} = -\frac{1}{2} \int_{D} u^{2} \left(\sum_{i=1}^{m} \alpha_{n,\ell} X_{i}^{2} \alpha_{n,\ell} + (X_{i} \alpha_{n,\ell})^{2} + \alpha_{n,\ell} (X_{i}^{*})^{2} \alpha_{n,\ell} + (X_{i}^{*} \alpha_{n,\ell})^{2} + \alpha_{n,\ell}^{2} (\operatorname{div}_{\mu}(X_{i}))^{2} \right) d\mu$$

and once again

$$|I_3| \leqslant C\ell^2 \int_{A_\ell} u^2 \, d\mu + Cn^2 \int_{A_n} u^2 \, d\mu \leqslant C(\operatorname{vol}(A_\ell) + \operatorname{vol}(A_n)) \underset{n,\ell \to +\infty}{\longrightarrow} 0.$$

All in all, $|q_D(u_n - u_\ell)| \to 0$ as $n, \ell \to +\infty$. Hence $\underline{u} \in \mathcal{D}(\widehat{q}_D)$.

Next, we have to check that $|\widehat{q}_D(\underline{u}, v)| \leq C(\underline{u}, v)_{L^2(D,\mu)}$ for any $v \in \mathcal{D}(\widehat{q}_D)$. It is sufficient to check it for $v \in C_c^{\infty}(D)$ and then extend it by density to $\mathcal{D}(\widehat{q}_D)$. Let $v \in C_c^{\infty}(D)$. We have

$$\widehat{q}_D(\underline{u}, v) = -\lim_{n \to +\infty} \int_D \sum_{i=1}^m X_i(\chi_n u) X_i v \ d\mu = \lim_{n \to +\infty} (\Delta u_n, v)_{L^2(D,\mu)}$$
$$= (\Delta \underline{u}, v)_{L^2(D,\mu)} = -\lambda(\underline{u}, v)_{L^2(D,\mu)}$$

since $u_n = \underline{u}$ on Supp(v) for n sufficiently large. Hence $\underline{u} \in \mathcal{D}(\Delta_D)$ and \underline{u} is an eigenfunction with eigenvalue λ .

By Lemma 3.3 we know that $\lambda_1(D) > 0$. Assume for the sake of a contradiction that $\lambda > \lambda_1(D)$. Let us denote by u_0 a non-negative function in the first eigenspace of $-\Delta_D$, which exists thanks to Lemma 3.3. Then according to Proposition 1.4, u_0 and \underline{u} are orthogonal for the $L^2(D,\mu)$ scalar product. At the beginning of the proof, we assumed without loss of generality that \underline{u} is non-negative, but since $D \subset \Omega \setminus Z_{\overline{u}}$, we even know that \underline{u} is strictly positive in D. Hence $u_0 = 0$ a.e. in D, which is a contradiction with the fact that $\lambda_1(D) > 0$. We conclude that $\lambda_1(D) = \lambda$.

We start the proof of Theorem 1 by proving its second part. For this, we follow the arguments of [9, Chapter VI (p. 453-454)] (see also [6, Appendix D]). Suppose that $u \in E_{\lambda_n}$ has at least (n + 1) nodal domains D_1, \ldots, D_{n+1} . We also assume $\lambda_{n-1} < \lambda_n$. For $1 \leq i \leq n$, we denote by u_i the restriction of u to D_i , which lies in the first eigenspace of the Dirichlet problem in D_i according to Lemma 3.4. In particular its Rayleigh quotient $R(u_i)$ is equal to λ_n due to Point 1. of Lemma 3.2. We extend u_i by 0 in $\Omega \setminus D_i$, and we still denote by u_i this extension, which belongs to $\mathcal{D}(\hat{q}_\Omega)$ according to Lemma 2.1. We can determine $(a_1, \ldots, a_n) \in \mathbb{R}^n \setminus \{0\}$ such that $f = \sum_{i=1}^{n} a_i u_i$ is orthogonal in $L^2(\Omega, \mu)$ to the (n-1) first eigenfunctions $\varphi_1, \ldots, \varphi_{n-1}$ of $-\Delta_{\Omega}$ on Ω . We have $R(f) = \lambda_n$, hence f is an eigenfunction for λ_n according to the min-max principle (Point 2 of Lemma 3.2). But f vanishes in the open set D_{n+1} in contradiction with the unique continuation property of eigenfunctions which is satisfied thanks to Assumption 1.6.

The first part of Theorem 1 follows from similar arguments, except that we avoid using the unique continuation property in the end.⁴ Assume for the sake of a contradiction that $u \in E_{\lambda_n}$ has at least $m_n = n + \text{mult}(\lambda_n)$ nodal domains D_1, \ldots, D_{m_n} . By standard linear algebra, there exist $\text{mult}(\lambda_n) + 1$ linearly independent functions $f_j, j = 1, \ldots, \text{mult}(\lambda_n) + 1$ of the form

$$f_j = \sum_{i=1}^{m_n} a_{j,i} u_i,$$

with $a_{j,i} \in \mathbb{R}$, u_i is the extension by 0 of the restriction of u to D_i , and f_j is orthogonal in $L^2(\Omega, \mu)$ to the (n-1) first eigenfunctions $\varphi_1, \ldots, \varphi_{n-1}$ of $-\Delta_{\Omega}$ on Ω . For any j we have $R(f_j) = \lambda_n$, hence f_j is an eigenfunction for λ_n according to the min-max principle (Point 2 of Lemma 3.2). The f_j are mult $(\lambda_n) + 1$ linearly independent eigenfunctions with eigenvalue λ_n , which is impossible.

Remark 3.5. Counterexamples to the unique continuation property are known when Assumption 1.6 is not satisfied, for operators of the form $\Delta + V$ where V is a smooth function on Ω (see [4]).

4 Sub-Riemannian tools

This section introduces the notations, the terminology and the tools of sub-Riemannian (sR) geometry which will be needed in the proof of Theorem 2. For a more comprehensive introduction to sR geometry, we refer to [2] and [5].

4.1 Sub-Riemannian flag

First, we define the sR distribution

$$\mathcal{D} = \operatorname{Span}(X_1, \dots, X_m)$$

and then the sR flag as follows: $\mathcal{D}^0 = \{0\}, \mathcal{D}^1 = \mathcal{D}$, and, for any $j \ge 1$,

$$\mathcal{D}^{j+1} = \mathcal{D}^j + [\mathcal{D}, \mathcal{D}^j].$$

For any $q \in M$, this gives a flag

$$\{0\} = \mathcal{D}_q^0 \subset \mathcal{D}_q^1 \subset \ldots \subset \mathcal{D}_q^{r-1} \subsetneq \mathcal{D}_q^{r(q)} = T_q M.$$

$$(4.1)$$

where \mathcal{D}_q^i denotes \mathcal{D}^i taken at point q. The integer r(q) is called the *step*, or *non-holonomic* order, of \mathcal{D} at q. In the case of Example 1.1 it is equal to 1 except on the singular line $\{x = 0\}$, where it is equal to 2 (or α). In the case of Example 1.3 it coincides with the step r defined there; in particular it is equal to 2 in Example 1.2 at any point.

For $i \in \{0, \ldots, r(q)\}$, we set

$$n_i(q) = \dim \mathcal{D}_q^i.$$

The sequence $(n_i(q))_{0 \leq i \leq r(q)}$ is called the growth vector at point q. We say that $q \in M$ is regular if the growth vector $(n_i(q'))_{0 \leq i \leq r(q')}$ at q' is constant for q' in a neighborhood of q. Otherwise, q is said to be singular. If any point $q \in M$ is regular, we say that the structure is equiregular.

 $^{{}^{4}}$ We would like to thank Iosif Polterovich for sharing with us this argument, itself communicated to him by Dan Mangoubi.

Remark 4.1. The Heisenberg sub-Laplacian of Example 1.2 is equiregular (with $n_1 = 2d$, $n_2 = 2d + 1$), but the Baouendi-Grushin sub-Laplacian of Example 1.1 is not equiregular.

The number

$$Q(q) = \sum_{i=1}^{r(q)} i(n_i(q) - n_{i-1}(q)), \qquad (4.2)$$

coincides at any regular point q with the Hausdorff dimension of the metric space induced by the sR distance on M near q (see [20]).

We define a non-decreasing sequence of weights $w_i(q)$. Roughly speaking, $w_i(q)$ is the minimal length of the brackets of X_1, \ldots, X_m needed to generate *i* independent directions at *q*. Formally, given any $i \in \{1, \ldots, N\}$ and $q \in M$, there exists a unique $j \in \{1, \ldots, r(q)\}$ such that $n_{j-1}(q) +$ $1 \leq i \leq n_j(q)$. We set $w_i(q) = j$. It is not difficult to check that (4.2) coincides with $\sum_{i=1}^N w_i(q)$ (see (1.10)).

4.2 Nilpotentization

The aim of the following paragraphs is to introduce a system of local coordinates, called privileged coordinates, in which it is natural to write Taylor expansions of vector fields defined on the sR manifold (see [5, Section 4], [14, Chapter 2]). The first order term in the Taylor expansion of a vector field in privileged coordinates is called the nilpotent approximation of the vector field. A typical example of privileged coordinates system is given by some exponential coordinates with respect to a frame of $T_q M$ which is "adapted" to the sR flag (4.1) (see (4.5) below).

4.2.1 Non-holonomic orders.

The non-holonomic order of a smooth germ of function is

$$\operatorname{ord}_q(f) = \min\{p \in \mathbb{N} : \exists i_1, \dots, i_p \in \{1, \dots, m\} \text{ such that } (X_{i_1} \dots X_{i_p} f)(q) \neq 0\}$$

where we adopt the convention that $\min \emptyset = +\infty$.

The non-holonomic order of a smooth germ of vector field X at q, denoted by $\operatorname{ord}_q(X)$, is the real number

$$\operatorname{ord}_q(X) = \sup\{\sigma \in \mathbb{R} : \operatorname{ord}_q(Xf) \ge \sigma + \operatorname{ord}_q(f), \forall f \in C^{\infty}(q)\} \in \mathbb{Z}.$$

In other words, applying X to a function f "increases" the non-holonomic order by at least $\operatorname{ord}_q(X)$ (we put quotation marks to indicate that since $\operatorname{ord}_q(X)$ may be negative, applying X may in fact decrease the non-holonomic order).

There holds $\operatorname{ord}_q(fg) = \operatorname{ord}_q(f) + \operatorname{ord}_q(g)$, $\operatorname{ord}_q(fX) = \operatorname{ord}_q(f) + \operatorname{ord}_q(X)$ and $\operatorname{ord}_q([X, Y]) \ge \operatorname{ord}_q(X) + \operatorname{ord}_q(Y)$. As a consequence, every X which has the property that $X(q') \in \mathcal{D}^i_{q'}$ for any q' in a neighborhood of q is of non-holonomic order $\ge -i$.

Example 4.2. Let us illustrate these definitions on Example 1.1 where $X_1 = \partial_x$ and $X_2 = x\partial_y$. At (x, y) = 0, the non-holonomic order of the function f(x, y) = x is 1 and the non-holonomic order of f(x, y) = y is 2 since $X_1X_2y = 1$. The non-holonomic order of ∂_x is -1, the non-holonomic order of ∂_y is -2, and the non-holonomic order of $x\partial_y$ is -1.

4.2.2 Privileged coordinates.

A system of *privileged coordinates* at q is a system of local coordinates (x_1, \ldots, x_N) near q verifying

$$\operatorname{ord}_q(x_i) = w_i, \qquad \text{for } 1 \leqslant i \leqslant N.$$
 (4.3)

In particular, privileged coordinates satisfy $\partial_{x_i} \in \mathcal{D}_q^{w_i(q)} \setminus \mathcal{D}_q^{w_i(q)-1}$ at q, meaning that privileged coordinates are adapted to the flag (see definition below). One can also check that for any $(\alpha_1, \ldots, \alpha_N) \in \mathbb{N}^N, (\beta_1, \ldots, \beta_N) \in \mathbb{N}^N$,

$$\operatorname{ord}_{q}(x_{1}^{\alpha_{1}}\dots x_{N}^{\alpha_{N}}\partial_{x_{1}}^{\beta_{1}}\dots \partial_{x_{N}}^{\beta_{N}}) = \sum_{i=1}^{N} (\alpha_{i} - \beta_{i})w_{i}.$$
(4.4)

We now describe a construction showing that privileged coordinates systems exist at any $q \in M$. A family (Z_1, \ldots, Z_N) of vector fields is said to be *adapted* to the sR flag at q if it is a frame of $T_q M$ at q and if $Z_i(q) \in \mathcal{D}_q^{w_i(q)}$ for any $i \in \{1, \ldots, N\}$. In other words, for any $i \in \{1, \ldots, r(q)\}$, the vectors $Z_1, \ldots, Z_{n_i(q)}$ at q span \mathcal{D}_q^i .

If (Z_1, \ldots, Z_N) is an adapted frame at q, it is proved in [14, Appendix B] that the inverse of the local diffeomorphism

$$(x_1, \dots, x_n) \mapsto \exp(x_1 Z_1) \dots \exp(x_N Z_N)(q) \tag{4.5}$$

defines privileged coordinates at q (called exponential coordinates of the second kind).

4.2.3 Dilations.

As we mentioned in the introduction, sR geometries are anisotropic. The natural sR dilations that we now define are thus also anisotropic.

Fix $q \in M$. For every $\varepsilon \in \mathbb{R} \setminus \{0\}$, the dilation $\delta_{\varepsilon} : \mathbb{R}^N \to \mathbb{R}^N$ is defined by

$$\delta_{\varepsilon}(x) = (\varepsilon^{w_1(q)} x_1, \dots, \varepsilon^{w_N(q)} x_N)$$

for every $x = (x_1, \ldots, x_N)$ - we omit the dependance in q in the notation.

A dilation δ_{ε} acts also on functions and vector fields on \mathbb{R}^N by pull-back: $\delta_{\varepsilon}^* f = f \circ \delta_{\varepsilon}$ and $\delta_{\varepsilon}^* X$ is the vector field such that $(\delta_{\varepsilon}^* X)(\delta_{\varepsilon}^* f) = \delta_{\varepsilon}^* (Xf)$ for any $f \in C^1(\mathbb{R}^N)$. In particular, given a system of privileged coordinates $\psi_q : U \to \mathbb{R}^N$, for any vector field X

In particular, given a system of privileged coordinates $\psi_q : U \to \mathbb{R}^N$, for any vector field X in U of non-holonomic order k there holds $\delta_{\varepsilon}^*(\psi_q)_* X = \varepsilon^{-k}(\psi_q)_* X$. We will use this property many times for vector fields of the form (4.4).

4.2.4 Nilpotent approximation.

We now turn to the definition of the nilpotent approximation, which is a first-order approximation of vector fields in privileged coordinates near a point $q \in M$. An explicit example of computation of nilpotent approximation is given in [14, Example 2.8], it may help to understand the definitions which follow.

Fix a system of privileged coordinates $\psi_q = (x_1, \ldots, x_N) : U \to \mathbb{R}^N$ defined in a neighborhood U of q. Coming back to the vector fields X_1, \ldots, X_m , we write the Taylor expansion

$$(\psi_q)_* X_i(x) \sim \sum_{\substack{\alpha \in \mathbb{N}^N \\ j \in \{1, \dots, N\}}} a_{\alpha, j} x^\alpha \partial_{x_j}.$$
(4.6)

Since $X_i \in \mathcal{D}$, its non-holonomic order is -1. Hence, when $a_{\alpha,j} \neq 0$ for some $\alpha = (\alpha_1, \ldots, \alpha_N) \in \mathbb{N}^N$ and $j \in \{1, \ldots, N\}$, the monomial vector field $x^{\alpha} \partial_{x_j}$ has non-holonomic order ≥ -1 , which implies that $\sum_{i=1}^N w_i(q) \alpha_i \geq w_j(q) - 1$ according to (4.4). Therefore, we may write X_i as a formal series

$$(\psi_q)_* X_i = X_i^{(-1)} + X_i^{(0)} + X_i^{(1)} + \dots$$
(4.7)

where $X_i^{(k)}$ is a homogeneous vector field of degree k, meaning that

$$\delta_{\varepsilon}^* X_i^{(k)} = \varepsilon^k X_i^{(k)}. \tag{4.8}$$

We set

$$\widehat{X}_i^q = X_i^{(-1)}, \qquad 1 \leqslant i \leqslant m \tag{4.9}$$

which is a vector field on \mathbb{R}^N . Then \widehat{X}_i^q is homogeneous of degree -1 with respect to dilations, meaning that $\delta_{\varepsilon}^* \widehat{X}_i^q = \varepsilon^{-1} \widehat{X}_i^q$ for $\varepsilon \neq 0$. For $\varepsilon > 0$ small enough we have

$$X_i^{\varepsilon} := \varepsilon \delta_{\varepsilon}^* (\psi_q)_* X_i = \widehat{X}_i^q + \varepsilon R_{i,\varepsilon}^q$$
(4.10)

where $R_{i,\varepsilon}^q$ depends smoothly on ε for the C^{∞} topology (see also [2, Lemma 10.58]).

Finally, the nilpotent approximation of X_1, \ldots, X_m at q is defined as $\widehat{M}^q \simeq \mathbb{R}^N$ endowed with the vector fields $\widehat{X}_1^q, \ldots, \widehat{X}_m^q$. This definition does not depend on the choice of privileged coordinates at q because two sets of such coordinates produce two "sR-isometric" sR structures. An important property is that $(\widehat{X}_1^q, \ldots, \widehat{X}_m^q)$ generates a nilpotent Lie algebra of step r(q) (see [14, Proposition 2.3]).

The nilpotent approximation of a measure μ on M at $q \in M$ is the measure on \mathbb{R}^N

$$\widehat{\mu}^{q} = \lim_{\varepsilon \to 0} \varepsilon^{-\mathcal{Q}(q)} \delta^{*}_{\varepsilon}(\psi_{q})_{*} \mu$$
(4.11)

where the convergence is understood in the vague topology. It follows from this definition that $\hat{\mu}^q$ is proportional to the Lebesgue measure.

4.3 Desingularization

The estimates we will need at some point in the proof of Theorem 2 blow-up at singular points. However, when $q \in M$ is a singular point, it is possible to lift locally in a neighborhood U of q the vector fields X_1, \ldots, X_m to vector fields $\widetilde{X}_1, \ldots, \widetilde{X}_m$ on $\widetilde{U} = U \times \mathbb{R}^K$, so that the lift $\widetilde{q} = (q, 0)$ of q is a *regular point* in \widetilde{U} and many properties of the vector fields are preserved. This lifting procedure will allow us to recover uniform estimates in Section 5.2.

Lemma 4.3. [14, Lemma 2.5 and Theorem 2.9] Let q be a point in M. Then there exist $K \in \mathbb{N}$, a neighborhood $U \subset M$ of q, coordinates (x, y) on $\widetilde{U} = U \times \mathbb{R}^K$ and smooth vector fields

$$\widetilde{X}_i(x,y) = X_i(x) + \sum_{j=1}^K b_{ij}(x,y)\partial_{y_j}, \qquad i = 1,\dots,m,$$

on \widetilde{U} such that

- $\widetilde{X}_1, \ldots, \widetilde{X}_m$ satisfy Hörmander's bracket-generating condition in \widetilde{U} ;
- every \widetilde{p} in \widetilde{U} is regular;

• denoting by $\pi : \widetilde{U} \to U$ the canonical projection, and by \widetilde{d} the sR distance defined by $\widetilde{X}_1, \ldots, \widetilde{X}_m$ on \widetilde{U} , we have $\pi_* \widetilde{X}_i = X_i$, and for $p \in U$ and ε small enough,

$$B_{\varepsilon}(p) = \pi \left(B_{\varepsilon}^{\widetilde{d}}((p,0)) \right).$$
(4.12)

Example 4.4. A possible (global) desingularization of the vector fields $X_1 = \partial_x$ and $X_2 = x \partial_y$ on \mathbb{R}^2 is given by the vector fields $\widetilde{X}_1 = \partial_x$ and $\widetilde{X}_2 = \partial_z + x \partial_y$ on \mathbb{R}^3 .

5 Proof of Theorem 2

The proof of Theorem 2 splits into two steps. The first one consists in proving an asymptotic upper bound for the first eigenvalue of the Dirichlet sub-Laplacian in a sR ball centered at $q \in M$ whose radius tends to 0. This upper bound is uniform in q when q is a regular point. The second step is to use Lemma 4.3 (i.e., a desingularization) to conclude.

5.1 The first eigenvalue of the sub-Laplacian in a small sR ball

We fix $q \in M$ and we take a chart $\psi_q : U \to \mathbb{R}^N$ of privileged coordinates at q, with $\psi_q(q) = 0$. We denote by R(q) the maximal radius such that $B_{R(q)}(q) \subset U$.

As seen in Section 4.2.4, the nilpotent approximations of X_1, \ldots, X_m (resp. of μ) at q are vector fields \widehat{X}_i^q (resp. a measure $\widehat{\mu}^q$) in \mathbb{R}^N .

For $\varepsilon \leq R(q)$, we set

$$L_{\varepsilon}^{2} = L^{2}((\psi_{q})_{*}B_{\varepsilon}(q), (\psi_{q})_{*}\mu)$$

We also fix $u_1 \in C_c^{\infty}(\text{Box}(1/2))$ such that $u_1(0) \neq 0$. Finally, we set

$$u_{\varepsilon}(x) = \varepsilon^{-\mathcal{Q}(q)/2} u_1(\delta_{1/\varepsilon} x) \tag{5.1}$$

and we have $u_{\varepsilon} \in C_c^{\infty}(\text{Box}(\varepsilon/2)).$

Lemma 5.1. If X is a vector field on $\psi_q(U) \subset \mathbb{R}^N$ which is homogeneous of degree $k \in \mathbb{R}$ (in the sense of (4.8)), then there exist $c(q) > 0, \varepsilon(q) > 0$ such that for any $\varepsilon \leq \varepsilon(q)$ and any $1 \leq i \leq m$,

$$\|Xu_{\varepsilon}\|_{L^{2}_{\varepsilon}} \leqslant c(q)\varepsilon^{k}\|u_{\varepsilon}\|_{L^{2}_{\varepsilon}}$$

$$(5.2)$$

Proof. First, the ball-box theorem (1.7) (see also [5], [14, Corollary 2.1]) yields the existence of $0 < \alpha \leq 1$ and $\varepsilon(q) > 0$ (both depending on q) such that for any $0 < \varepsilon < \varepsilon(q)$,

$$\operatorname{Box}(\alpha\varepsilon) \subset (\psi_q)_* B_{\varepsilon}(q) \subset \operatorname{Box}(\alpha^{-1}\varepsilon).$$
(5.3)

We use (5.1) and the homogeneity in ε of X, Box(ε) and $\hat{\mu}^q$. Due to (4.11) this implies the following two convergences

$$\varepsilon^{-\mathcal{Q}(q)/2}\varepsilon^{-k}\|Xu_{\varepsilon}\|_{L^{2}(\operatorname{Box}(\varepsilon),(\psi_{q})_{*}\mu)} = \varepsilon^{-\mathcal{Q}(q)/2}\|Xu_{1}\|_{L^{2}(\operatorname{Box}(1),\delta_{\varepsilon}^{*}(\psi_{q})_{*}\mu)} \xrightarrow{}_{\varepsilon \to 0} \|Xu_{1}\|_{L^{2}(\operatorname{Box}(1),\widehat{\mu}^{q})}$$
$$\varepsilon^{-\mathcal{Q}(q)/2}\|u_{\varepsilon}\|_{L^{2}(\operatorname{Box}(\alpha^{2}\varepsilon),(\psi_{q})_{*}\mu)} = \varepsilon^{-\mathcal{Q}(q)/2}\|u_{1}\|_{L^{2}(\operatorname{Box}(\alpha^{2}),\delta_{\varepsilon}^{*}(\psi_{q})_{*}\mu)} \xrightarrow{}_{\varepsilon \to 0} \|u_{1}\|_{L^{2}(\operatorname{Box}(\alpha^{2}),\widehat{\mu}^{q})}.$$

Taking the ratio of the two convergences (justified by the fact that $u_1(0) \neq 0$ hence the last limit is $\neq 0$), we obtain

$$\|Xu_{\varepsilon}\|_{L^{2}(\mathrm{Box}(\varepsilon),(\psi_{q})*\mu)} \leq c(q)\varepsilon^{k}\|u_{\varepsilon}\|_{L^{2}(\mathrm{Box}(\alpha^{2}\varepsilon),(\psi_{q})*\mu)}.$$
(5.4)

Using twice (5.3), we obtain

$$\|Xu_{\varepsilon}\|_{L^{2}_{\varepsilon}} \leq \|Xu_{\varepsilon}\|_{L^{2}(\operatorname{Box}(\alpha^{-1}\varepsilon),(\psi_{q})*\mu)} \leq c(q)\varepsilon^{k}\|u_{\varepsilon}\|_{L^{2}(\operatorname{Box}(\alpha\varepsilon),(\psi_{q})*\mu)} \leq c(q)\varepsilon^{k}\|u_{\varepsilon}\|_{L^{2}_{\varepsilon}}$$

which implies the lemma.
$$\Box$$

Corollary 5.2. For any $q \in M$, there exist c(q) > 0 and $\varepsilon(q) > 0$ such that for any $\varepsilon \leq \varepsilon(q)$, there holds $\lambda_1(B_{\varepsilon}(q)) \leq c(q)\varepsilon^{-2}$.

Proof. We fix $q \in M$ and $1 \leq i \leq m$. According to (5.3), we know that $u_{\alpha\varepsilon}$ is supported in $(\psi_q)_*B_{\varepsilon}(q)$. We set $R_i^q = (\psi_q)_*X_i - \hat{X}_i^q$, which is a vector field on $(\psi_q)_*(U) \subset \mathbb{R}^N$. Let us assume for the moment that we have proved the existence of C(q) such that for ε sufficiently small,

$$\|R_i^q u_{\alpha\varepsilon}\|_{L^2_{\varepsilon}}^2 \leqslant C(q) \|u_{\alpha\varepsilon}\|_{L^2_{\varepsilon}}^2.$$

$$(5.5)$$

Then we can write

$$\begin{aligned} \|(\psi_q)_* X_i u_{\alpha\varepsilon}\|_{L^2_{\varepsilon}}^2 &\leq 2\left(\|\widehat{X}_i^q u_{\alpha\varepsilon}\|_{L^2_{\varepsilon}}^2 + \|R_i^q u_{\alpha\varepsilon}\|_{L^2_{\varepsilon}}^2\right) \\ &\leq (c(q)\varepsilon^{-2} + C(q))\|u_{\alpha\varepsilon}\|_{L^2_{\varepsilon}}^2 \\ &\leq c'(q)\varepsilon^{-2}\|u_{\alpha\varepsilon}\|_{L^2_{\varepsilon}}^2 \end{aligned}$$
(5.6)

where in the second inequality we used Lemma 5.1 with $X = \hat{X}_i^q$, and (5.5). By the min-max principle (Lemma 3.2), we get the result.

There remains to prove (5.5). We can write $R_i^q = \sum_{k=1}^N a_{i,k}^q \partial_{x_k}$ where the $a_{i,k}^q$ are smooth functions of $x \in \mathbb{R}^N$. We then apply Taylor's theorem for multivariate functions with exact remainder to each $a_{i,k}^q$. Recalling (4.7), (4.9) and the fact that $R_i^q = (\psi_q)_* X_i - \hat{X}_i^q$ has homogeneous components of order ≥ 0 only, this yields a decomposition ("factorizing out from R_i^q monomial vector fields of order 0")

$$R_i^q = \sum_{j \in \mathcal{J}} b_{i,j}^q Y_j \tag{5.7}$$

where $b_{i,j}^q$ is a continuous function of x. Here \mathcal{J} is a finite set such that $(Y_j)_{j \in \mathcal{J}}$ consists of all monomial vector fields Y which are homogeneous of degree 0, i.e. Y is of the form $Y = x_i \partial_{x_k}$ for some indices $i, k \in \{1, \ldots, n\}$ satisfying $w_i = w_k$. Then we can write

$$\|R_i^q u_{\alpha\varepsilon}\|_{L^2_{\varepsilon}} \leqslant \sum_{j \in \mathcal{J}} \|b_{i,j}^q Y_j u_{\alpha\varepsilon}\|_{L^2_{\varepsilon}} \leqslant \sum_{j \in \mathcal{J}} \|b_{i,j}^q\|_{L^{\infty}((\psi_q)_* B_{\varepsilon}(q))} \|Y_j u_{\alpha\varepsilon}\|_{L^2_{\varepsilon}} \leqslant C(q) \|u_{\alpha\varepsilon}\|_{L^2_{\varepsilon}}$$

where the last inequality comes from Lemma 5.1 applied with k = 0 (and the fact that $b_{i,j}^q$ is smooth, hence bounded). This concludes the proof of (5.5).

Remark 5.3. One can in fact prove that $\varepsilon^2 \lambda_1(B_{\varepsilon}(q))$ converges to the first eigenvalue of the Dirichlet sub-Laplacian $\widehat{\Delta}^q$ on $L^2(\widehat{B}_1(q), \widehat{\mu}^q)$, which is a stronger statement than Corollary 5.2. Here, $\widehat{\Delta}^q = \sum_{i=1}^m (\widehat{X}_i^q)^2$ and $\widehat{B}_1(q) \subset \mathbb{R}^N$ denotes the sR ball computed with the metric obtained by replacing in (1.5) the vector fields X_i by the nilpotentized ones \widehat{X}_i^q .

However, what we will need for our purpose is the uniformity of the convergence with respect to q, and it is easier to prove the uniformity of c(q) with respect to q (see Lemma 5.4) than the uniformity of the convergence of $\varepsilon^2 \lambda_1(B_{\varepsilon}(q))$ with respect to q. This is why we prefered to keep our weaker statements.

Lemma 5.4. When q is regular, the constants c(q) and $\varepsilon(q)$ in Corollary 5.2 can be taken uniform in a small neighborhood of q.

Proof. We use the fact that taking a nilpotent approximation is a "uniform" procedure near a regular point (but it is not uniform near a singular point). This fact is described in Section 2.2.2 in [14], and it mainly relies on the property that q being regular, there exists a smooth frame $q' \mapsto (Z_1(q'), \ldots, Z_N(q')) \in (T_{q'}M)^N$ which is an adapted frame at every point q' in some

neighborhood V of q. Using (4.5), this yields a smoothly varying system of privileged coordinates in V, and a smooth nilpotent approximation in V (see Definition 2.9 in [14]). As remarked in [8, Section 4.4] where a similar uniformity argument as ours is carried out, the continuity in Definition 2.9 of [14] can be replaced by smoothness.

It follows from Theorem 2.3 in [14] that the constants α and $\varepsilon(q')$ in (5.3) can be taken uniform over $q' \in V$. These uniform constants are respectively denoted by $\alpha \in (0, 1]$ and $\varepsilon(V) > 0$. We will deduce that the inequality (5.4) remains true in V with a uniform constant c(V). To state this property rigorously (see (5.8)), we notice that given any family of vector fields $V \ni q' \mapsto X^{q'}$ which is smooth in $q' \in V$, since $\{\widehat{\mu}^{q'}\}_{q' \in V}$ is a smooth family of measures (see [1, Section 4.1]), the constant

$$c(q') = \frac{\|X^{q'}u_1\|_{L^2(\text{Box}(1),\hat{\mu}^{q'})}}{\|u_1\|_{L^2(\text{Box}(\alpha^2),\hat{\mu}^{q'})}}$$

in (5.4) is continuous over V, and its supremum over V is denoted by c(V). Moreover, the convergence (4.11) is also uniform over V, due to the smoothness of μ . We assume that each vector field $X^{q'}$ is homogeneous of degree $k \in \mathbb{R}$. Following the proof of Lemma 5.1, we obtain that for any $q' \in V$ and any $0 < \varepsilon < \varepsilon(V)$ there holds

$$\|X^{q'}u_{\varepsilon}\|_{L^{2}(\mathrm{Box}(\varepsilon),(\psi_{q'})*\mu)} \leq c(V)\varepsilon^{k}\|u_{\varepsilon}\|_{L^{2}(\mathrm{Box}(\alpha^{2}\varepsilon),(\psi_{q'})*\mu)}.$$
(5.8)

Applying (5.8) to $X^{q'} = \widehat{X}_i^{q'}$ we obtain that $\|\widehat{X}_i^{q'} u_{\alpha\varepsilon}\|_{L^2_{\varepsilon}} \leq c(V)\varepsilon^{-2}$ for any $q' \in V$. Since $q' \mapsto (\psi_{q'})_* X_i(q')$ and $q' \mapsto \widehat{X}_i^{q'}$ are smooth, the map

$$V \ni q' \mapsto (\psi_{q'})_* X_i(q') - \widehat{X}_i^{q'}(q') = R_i^{q'}$$

is also smooth. Therefore the functions $b_{i,j}^{q'}$ defined in (5.7) depend smoothly on q'. Besides, the vector fields $x_i \partial_{x_k}$ for $i, k \in \{1, \ldots, N\}$, which appear in (5.7), do not depend on q'. This implies that the constant C(q') in (5.5) can be taken uniform over V. More precisely, this means that there exists C(V) > 0 such that for any $\varepsilon < \varepsilon(V)$ and any $q' \in V$,

$$\|R_{i}^{q'}u_{\alpha\varepsilon}\|_{L^{2}((\psi_{q'})_{*}B_{\varepsilon}(q'),(\psi_{q'})_{*}\mu)}^{2} \leqslant C(V)\|u_{\alpha\varepsilon}\|_{L^{2}((\psi_{q'})_{*}B_{\varepsilon}(q'),(\psi_{q'})_{*}\mu)}^{2}.$$
(5.9)

This implies that the constants $c(q'), \varepsilon(q')$ in Corollary 5.2 can be taken uniform over $q' \in V$, which proves Lemma 5.4.

Remark 5.5. Corollary 5.2 establishes an upper bound for the first Dirichlet eigenvalue as $r \rightarrow 0$. Note that lower bounds on the first Neumann eigenvalue were established in [15], this is equivalent to Poincaré's inequality.

5.2 End of the proof of Theorem 2

We assume that φ_{λ} is an eigenfunction of $-\Delta_{\Omega}$ not belonging to the first eigenspace E_{λ_1} . We denote by D_j its nodal domains. According to Lemma 3.4, the restriction of φ_{λ} to each domain D_j is an eigenfunction of the Dirichlet sub-Laplacian Δ_{D_j} , it belongs to its first eigenspace, and $\lambda_1(D_j) = \lambda$ for each D_j . Now, if $x \in \Omega$ and $d(q, Z_{\varphi_{\lambda}}) > \varepsilon$, where d is the sR distance, then $B_{\varepsilon}(q) \subset D_j$ for some j. By the min-max principle, it implies that $\lambda = \lambda_1(D_j) \leq \lambda_1(B_{\varepsilon}(q))$. But $\lambda_1(B_{\varepsilon}(q)) \leq c(q)\varepsilon^{-2}$ thanks to Corollary 5.2, hence $\varepsilon \leq c(q)\lambda^{-1/2}$.

If q is regular, using Lemma 5.4, we obtain that the constant c(q') above is in fact uniform for q' in a neighborhood of q. Hence any sR ball centered in a neighborhood V of q and of radius $\geq c'(V)\lambda^{-1/2}$ will intersect $Z_{\varphi_{\lambda}}$, which concludes the proof of the theorem "locally near q" in this case.

If $q \in \Omega$ is a singular point, the idea is to desingularize the vector fields at q thanks to Lemma 4.3 in order to recover a regular neighborhood but in a higher-dimensional space, and be able to apply the result we just obtained in the regular case. Following the notations and defined quantities of Lemma 4.3, we consider

$$\widetilde{X}_i(x,y) = X_i(x) + \sum_{j=1}^K b_{ij}(x,y)\partial_{y_j}, \qquad i = 1,\dots,m,$$

for $(x, y) \in U \times \mathbb{R}^K$.

We will build a sub-Laplacian $\widetilde{\Delta}$ satisfying the following key properties:

- It is defined on the bounded set $\Omega \times \mathbb{T}^K$ where $\mathbb{T} = \mathbb{R}/2\pi\mathbb{Z}$.
- In a neighborhood of (q, 0) we have $\widetilde{\Delta} = -\sum_{i=1}^{m} \widetilde{X}_{i}^{*} \widetilde{X}_{i}$. Its domain $D(\widetilde{\Delta})$ is constructed as in Section 2.
- The vector fields defining $\widetilde{\Delta}$ satisfy Hörmander's bracket-generating condition everywhere in $\Omega \times \mathbb{T}^{K}$.

The construction of $\widetilde{\Delta}$ is achieved through cut-offs and extensions of the vector fields \widetilde{X}_i to vector fields which are periodic in the y_j variables and thus defined on $\Omega \times \mathbb{T}^K$. In the sequel, \mathbb{T} is identified with $[-\pi, \pi)$ (with periodic boundary).

Without loss of generality we assume that U is contained in the fundamental domain $\Omega \times [-\pi,\pi)^K$. We fix a compact set $V \subset U$ which is a neighborhood of (q,0). We consider cut-off functions satisfying:

- $\chi_0: \Omega \times \mathbb{T}^K \to \mathbb{R}^+$ is a smooth function which is equal to 1 in V and 0 in $\Omega \times \mathbb{T}^K \setminus U$.
- $\chi_1: \Omega \times \mathbb{T}^K \to \mathbb{R}^+$ is a smooth function which is equal to 0 in V and > 0 outside V.

We consider the vector fields

$$\underline{\widetilde{X}}_{i}(x,y) = X_{i} + \sum_{j=1}^{K} \chi_{0}(x,y) b_{ij}(x,y) \partial_{y_{j}}$$
(5.10)

for $i \in \{1, \ldots, m\}$ and $Y_j = \chi_1(x, y)\partial_{y_j}$ for $j \in \{1, \ldots, K\}$ on $\Omega \times \mathbb{T}^K$, and the sub-Laplacian on $\Omega \times \mathbb{T}^K$ defined by

$$\widetilde{\Delta} = -\sum_{i=1}^{m} (\underline{\widetilde{X}}_{i})^{*} \underline{\widetilde{X}}_{i} - \sum_{j=1}^{K} Y_{j}^{*} Y_{j}.$$

Lemma 5.6. The family of vector fields $\underline{X}_1, \ldots, \underline{X}_m, Y_1, \ldots, Y_K$ satisfies Hörmander's bracketgenerating condition (1.1), it is regular at (q, 0), and the non-characteristic boundary condition (Assumption 1.5) is verified on $\partial(\Omega \times \mathbb{T}^K)$.

Proof. In V there holds $\underline{\widetilde{X}}_i = \widetilde{X}_i$. By Lemma 4.3, this implies that the Hörmander bracketgenerating condition is satisfied in V. We notice that for any $i_1, \ldots, i_\ell \in \{1, \ldots, m\}$,

$$[\underline{\widetilde{X}}_{i_1}, [\underline{\widetilde{X}}_{i_2}, [\dots, \underline{\widetilde{X}}_{i_\ell}]] = [X_{i_1}, [X_{i_2}, [\dots, X_{i_\ell}]] \mod Y_1, \dots, Y_K.$$

Since the Y_j do not vanish outside V, using that the vector fields X_1, \ldots, X_m satisfy Hörmander's bracket-generating condition in Ω , we obtain that the family of vector fields $\tilde{X}_1, \ldots, \tilde{X}_m, Y_1, \ldots, Y_K$ satisfies Hörmander's bracket-generating condition outside V.

The regularity of the family at (q, 0) follows from the fact that in V there holds $\underline{\widetilde{X}}_i = \widetilde{X}_i$ and $Y_j = 0$ for any $i \in \{1, \ldots, m\}$ and any $j \in \{1, \ldots, K\}$. By definition of the desingularized vector fields \widetilde{X}_i , they form an equiregular family, so in particular a regular family at (q, 0). The non-characteristic boundary condition on $\partial(\Omega \times \mathbb{T}^K)$ follows from the non-characteristic boundary condition satisfied by the vector fields X_i on $\partial\Omega$ (Assumption 1.5) and the fact that $\underline{\widetilde{X}}_i = X_i$ on $\partial(\Omega \times \mathbb{T}^K)$.

Let $\varphi_{\lambda}(x)$ be an eigenfunction of $-\Delta_{\Omega}$, with eigenvalue λ . We consider $\psi_{\lambda} : \Omega \times \mathbb{T}^K \to \mathbb{R}$ defined by $\psi_{\lambda}(x, y) = \varphi_{\lambda}(x)$. This is an eigenfunction of $-\widetilde{\Delta}$ with eigenvalue λ .

We apply the arguments of the beginning of Section 5.2 to Δ : they imply that there exist a neighborhood $\widetilde{V} \subset \Omega \times \mathbb{T}^K$ and a constant $c(\widetilde{V}) > 0$ independent of λ such that

$$\forall \widetilde{q} \in \widetilde{V}, \ \forall \varepsilon \ge c(\widetilde{V})\lambda^{-1/2}, \quad Z_{\psi_{\lambda}} \cap \widetilde{B}_{\varepsilon}(\widetilde{q}) \neq \emptyset.$$
(5.11)

These sR balls are computed with the vector fields defining $\widetilde{\Delta}$, and these vector fields coincide near (q, 0) with \widetilde{X}_i thanks to (5.10); hence it is equivalent to compute the sR balls with the vector fields \widetilde{X}_i since we are considering small balls near (q, 0), with radius much smaller than δ . By the projection property (4.12), since $Z_{\psi_{\lambda}} = Z_{\varphi_{\lambda}} \times \mathbb{T}^K$, we finally obtain that

$$\forall \widetilde{q} \in \widetilde{V}, \ \forall \varepsilon \ge c(\widetilde{V})\lambda^{-1/2}, \quad Z_{\varphi_{\lambda}} \cap B_{\varepsilon}(\pi(\widetilde{q})) \neq \emptyset$$
(5.12)

where $\pi : \Omega \times \mathbb{T}^K$ is the canonical projection. The constant involved in (5.12) is thus uniform in a neighborhood of q. Since this uniformity is true in a neighborhood of any point $q \in M$ (either regular or singular), using the compactness of $\overline{\Omega}$ we obtain the result.

Remark 5.7. One could wonder why we do not simply consider in the proof the sub-Laplacian $-\sum_{i=1}^{m} \widetilde{X}_{i}^{*} \widetilde{X}_{i}$ on $U \times \mathbb{R}^{K}$ instead of $\widetilde{\Delta}$ (the adjoint being computed with respect to $\widetilde{\mu} = \mu \otimes \mathscr{L}_{\mathbb{R}^{K}}$, where $\mathscr{L}_{\mathbb{R}^{K}}$ is the Lebesgue measure on \mathbb{R}^{K}). In fact this does not work for our purposes, since the formula $\psi_{\lambda}(x, y) = \varphi_{\lambda}(x)$ does not define an $L^{2}(U \times \mathbb{R}^{K})$ eigenfunction.

5.3 An example

In this section we illustrate Theorem 2 with an example.

Fix $\alpha \in \mathbb{N}^*$ and consider the generalized Baouendi-Grushin sub-Laplacian $\Delta_{BG} = \partial_x^2 + x^{2\alpha} \partial_y^2$ on $(-1,1)_x \times \mathbb{T}_y$. For $k \in \mathbb{Z}$ we denote by ψ_k a non-trivial element of the lowest energy eigenspace of the 1D operator $H_k = -\partial_x^2 + k^2 x^{2\alpha}$ on $(-1,1)_x$. The associated eigenvalue satisfies

$$c_{\alpha}|k|^{2/(\alpha+1)} \leqslant \mu_k \leqslant C_{\alpha}|k|^{2/(\alpha+1)}$$

as $k \to +\infty$ for some constants $c_{\alpha}, C_{\alpha} > 0$. Then $\Psi_k : (x, y) \mapsto \psi_k(x) \cos(ky)$ is an eigenfunction of $-\Delta_{\text{BG}}$ with eigenvalue μ_k . Its nodal set is

$$Z_{\Psi_k} = (-1,1)_x \times \left(\bigcup_{n \in \mathbb{Z}} \pi \frac{n}{k} \mathbb{Z}\right)_y$$

i.e. it is a union of "horizontal" lines separated by π/k (in Euclidean distance).

Using the ball-box theorem (1.7), the sR ball centered at a point $(0, y_0)$ on the singular line, and of radius ε , can be compared with ball-boxes of the form $[-C\varepsilon, C\varepsilon]_x \times [y_0 - (C\varepsilon)^{\alpha+1}, y_0 + (C\varepsilon)^{\alpha+1}]_y$:

the weights w_i in this case are $w_1 = 1$ and $w_2 = \alpha + 1$. The sR ball is more squeezed in the y direction due to the fact that brackets are needed to span this direction; we refer the reader to [5, Section 3.1] and the picture in [5, Section 3.3] for the case $\alpha = 1$. Hence, the statement that any sR ball of radius $c/\sqrt{\mu_k}$ intersects the nodal set of Ψ_k is true for c large enough but false for c small enough. This proves the sharpness of Theorem 2.

6 Proof of Theorem 3

We use the setting defined in Example 1.2 for d = 1 (and we drop the indices: x, y, z replace x_1, y_1, z_1): we consider $M = \Gamma \setminus \mathbf{H}_1$, with coordinates x, y, z, and endowed with the Lebesgue measure $\mu = dx dy dz$. We denote by Ω the open subset of M containing all points $(x, y, z) \in M$ such that $x \notin \sqrt{2\pi}(\mathbb{Z} + \frac{1}{2})$ (the factor $\frac{1}{2}$ becomes clear in the proof). The boundary condition is then at $x = \pm \sqrt{\frac{\pi}{2}} \pmod{2\pi}$.

We consider the vector fields $X = \partial_x$ and $Y = \partial_y - x\partial_z$ on M (see Example 1.2). The sub-Laplacian is

$$\Delta = -X^*X - Y^*Y = X^2 + Y^2.$$

Proposition 1.4 applies; we denote by $(\Delta_{\Omega}, \mathcal{D}(\Delta_{\Omega}))$ the domain of Δ acting on functions on Ω (with Dirichlet boundary conditions). We now proceed to explicit computations.

For $c \in \mathbb{R}$ we introduce the sets

$$A_c = \{ (x, y, z) \in \Omega \mid y = c \}, \qquad B_c = \{ (x, y, z) \in \Omega \mid z = c \}.$$

We make the following observations:

- All these hypersurfaces have Hausdorff dimension 3 (see [12, Section 0.6.C]).
- All hypersurfaces A_c have the same 3D Hausdorff measure, which we denote by a. This follows from the fact that for any $t \in \mathbb{R}$, the multiplication on the left by (0, t, 0) is an isometry, which sends (x, y, z) to (x, y + t, z), and thus A_c to A_{c+t} for any $c \in \mathbb{R}$. Since it is an isometry, it preserves balls and Hausdorff measures, and thus A_c and A_{c+t} have the same 3D Hausdorff measure, for any $c, t \in \mathbb{R}$.
- All hypersurfaces B_c have the same 3D Hausdorff measure, which we denote by b. This follows from the fact that for any $t \in \mathbb{R}$, the multiplication on the left by (0,0,t) is an isometry, which sends (x, y, z) to (x, y, z + t), and thus B_c to B_{c+t} for any $c \in \mathbb{R}$. Since it is an isometry, it preserves balls and Hausdorff measures, and thus B_c and B_{c+t} have the same 3D Hausdorff measure, for any $c, t \in \mathbb{R}$.

We consider two sequences of eigenfunctions of Δ_{Ω} ; it is not difficult to check that they are indeed well-defined on Ω and that they satisfy Dirichlet boundary conditions.

We first consider eigenfunctions of the form $\varphi_{1,m}(x, y, z) = \sin(\sqrt{2\pi}x)\sin(\sqrt{2\pi}my)$ for $m \in \mathbb{Z}_{\geq 0}$, for which the corresponding eigenvalue is $\lambda_{1,m} = 2\pi(1+m^2)$. The nodal set in Ω^5 is the disjoint union of $\{x = 0\}$ (whose Hausdorff measure is denoted by a_0) with the sets A_c for $c = \frac{k}{m}\sqrt{\frac{\pi}{2}}, k \in \{-m+1, \ldots, m-1\}$. The Hausdorff measure of this nodal set is $a_0 + (2m-1)a$ which is bounded above by $Ca\sqrt{\lambda_{1,m}}$ for some C > 0 independent of m.

Secondly, we consider for $m \in \mathbb{Z}_{\geq 0}$ the eigenfunction defined by extending to Ω by periodization with the group law (1.4) the function on the fundamental cell $\left(-\sqrt{\frac{\pi}{2}}, \sqrt{\frac{\pi}{2}}\right) \times \left[-\sqrt{\frac{\pi}{2}}, \sqrt{\frac{\pi}{2}}\right) \times \left[-\pi, \pi\right)$ given by $\varphi_{2,m}(x, y, z) = \psi_m(x) \sin(mz)$, where ψ_m denotes a non-null element of the

⁵since $x \in (-\sqrt{\pi/2}, \sqrt{\pi/2})$, x = 0 is the only nodal set coming from $\sin(\sqrt{2\pi}x)$.

first eigenspace of the 1D operator $-d^2/dx^2 + m^2x^2$ on $(-\sqrt{\frac{\pi}{2}}, \sqrt{\frac{\pi}{2}})$ with Dirichlet boundary conditions. The first eigenvalue $\lambda_{2,m}$ of this harmonic oscillator, which is also the eigenvalue associated to $\varphi_{2,m}$, is m + o(1) as $m \to +\infty$. Since ψ_m does not vanish, the nodal set of $\varphi_{2,m}$ is the (disjoint) union of the sets B_c for $c = k\pi/m$, $k \in \{-m + 1, \ldots, m - 1\}$. Its Hausdorff measure is (2m-1)b which is bounded below by $C\lambda_{2,m}b$ for $m \neq 0$ and some C > 0 independent of m.

This completes the proof of Theorem 3.

References

- Andrei Agrachev, Davide Barilari and Ugo Boscain. On the Hausdorff volume in sub-Riemannian geometry. *Calculus of Variations and Partial Differential Equations*, 2012, vol. 43, no 3-4, p. 355-388.
- [2] Andrei Agrachev, Davide Barilari and Ugo Boscain. A comprehensive introduction to sub-Riemannian geometry. Cambridge University Press, 2019.
- [3] Colette Anné. Bornes sur la multiplicité. Prépublications EPFL, mars 1992.
- [4] Hajer Bahouri. Non prolongement unique des solutions d'opérateurs "somme de carrés". Annales de l'institut Fourier, 1986, tome 36, no 4, p. 137-155.
- [5] André Bellaïche. The tangent space in sub-Riemannian geometry. In Sub-Riemannian geometry, Birkhäuser Basel, 1996. p. 1-78.
- [6] Pierre Bérard and Daniel Meyer. Inégalités isopérimétriques et applications. Annales scientifiques de l'École Normale Supérieure, 1982, vol. 15, no 3, p. 513-541.
- [7] Jean-Michel Bony. Principe du maximum, inégalité de Harnack et unicité du problème de Cauchy pour les opérateurs elliptiques dégénérés. Annales de l'institut Fourier, 1969, vol. 19, no 1, p. 277-304.
- [8] Yves Colin de Verdière, Luc Hillairet, and Emmanuel Trélat. Small-time asymptotics of hypoelliptic heat kernels near the diagonal, nilpotentization and related results. Annales Henri Lebesgue, 2021, vol. 4, p. 897-971.
- [9] Richard Courant and David Hilbert. Methods of mathematical physics: Partial Differential Equations, Volume 2. John Wiley & Sons, 2008.
- [10] Harold Donnelly and Charles Fefferman. Nodal sets of eigenfunctions on Riemannian manifolds. *Inventiones Mathematicae*, 1988, vol. 93, p. 161-183.
- [11] Lawrence C. Evans. Partial differential equations. American Mathematical Society, 2010.
- [12] Mikhael Gromov. Carnot-Carathéodory spaces seen from within. In Sub-Riemannian geometry, Birkhäuser Basel, 1996. p. 79-323.
- [13] Lars Hörmander. Hypoelliptic second order differential equations. Acta Mathematica, 1967, vol. 119, p. 147-171.
- [14] Frédéric Jean. Control of nonholonomic systems: from sub-Riemannian geometry to motion planning. Springer Briefs in Mathematics, Springer, 2014.

- [15] David Jerison. The Poincaré inequality for vector fields satisfying Hörmander's condition. Duke Mathematical Journal, 1986, vol. 53, no 2, p. 503-523.
- [16] Joseph J. Kohn and Louis Nirenberg. Non-coercive boundary value problems. *Communications on Pure and Applied Mathematics*, 1965, vol. 18, no 3, p. 443-492.
- [17] Camille Laurent and Matthieu Léautaud. Tunneling estimates and approximate controllability for hypoelliptic equations. *Memoirs of the American Mathematical Society*, 2022, vol. 1357, vi+95p.
- [18] Mathieu Lewin. Théorie spectrale et mécanique quantique. Mathématiques et Applications (SMAI). Springer International Publishing, 2022.
- [19] Alexander Logunov and Eugenia Malinnikova. Review of Yau's conjecture on zero sets of Laplace eigenfunctions. In: *Current Developments in Mathematics*, 2018, vol. 2018, no 1, p. 179-212.
- [20] John Mitchell. On Carnot-Caratheodory metrics. Journal of Differential Geometry, 1985, vol. 21, no 1, p. 35-45.
- [21] Arke Pleijel. Remarks on Courant's nodal theorem. Communications in Pure and Applied Mathematics, 1956, vol. 9, p. 543–550.
- [22] Michael Reed and Barry Simon. Methods of modern mathematical physics. IV. Analysis of operators. Academic Press, New York-London, 1978.
- [23] Frigyes Riesz and Béla Nagy. Functional Analysis. Blackie & Son LTD, London, England 1956.
- [24] Linda P. Rothschild and Elias M. Stein. Hypoelliptic differential operators and nilpotent groups. Acta Mathematica, 1976, vol. 137, p. 247-320.
- [25] Michel Rumin. Sub-Riemannian limit of the differential form spectrum of contact manifolds. Geometric & Functional Analysis GAFA, 2000, vol. 10, no 2, p. 407-452.
- [26] Kinji Watanabe. Sur l'unicité du prolongement des solutions des équations elliptiques dégénérées. Tôhoku Math. J., 1982, vol. 34, no 2, p. 239-249.
- [27] Igor Wigman. On the nodal structures of random fields-a decade of results. ArXiv preprint arXiv:2206.10020, 2022.