



HAL
open science

Biocatalytic oxidation of fatty alcohols into aldehydes for the flavors and fragrances industry

David Ribeaucourt, Bastien Bissaro, Fanny Lambert, Mickael Lafond,
Jean-Guy Berrin

► **To cite this version:**

David Ribeaucourt, Bastien Bissaro, Fanny Lambert, Mickael Lafond, Jean-Guy Berrin. Biocatalytic oxidation of fatty alcohols into aldehydes for the flavors and fragrances industry. *Biotechnology Advances*, 2022, 56, pp.107787. 10.1016/j.biotechadv.2021.107787 . hal-03619407

HAL Id: hal-03619407

<https://hal.science/hal-03619407>

Submitted on 4 May 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Biocatalytic oxidation of fatty alcohols into aldehydes for the** 2 **flavors and fragrances industry**

3 David Ribeaucourt^{1,2,3*}, Bastien Bissaro¹, Fanny Lambert², Mickael Lafond³ and Jean-Guy
4 Berrin^{1*}

¹ INRAE, Aix Marseille Univ, UMR1163 Biodiversité et Biotechnologie Fongiques, 13009, Marseille, France

² V. Mane Fils, 620 route de Grasse, 06620 Le Bar sur Loup, France

³ Aix Marseille Univ, CNRS, Centrale Marseille, iSm2, Marseille, France

5

6 *corresponding authors:

David Ribeaucourt (ribeaucourt.david@gmail.com)

Jean-Guy Berrin (jean-guy.berrin@inrae.fr)

7

8 Keywords: Flavors and Fragrances, Fragrant aldehydes, Biocatalysis, Enzymes, Copper-
9 Radical Oxidase, Fatty Alcohol, Fatty Aldehyde, AA5_2

10

11 **Abstract**

12 From Egyptian mummies to the Chanel n°5 perfume, fatty aldehydes have long been used and
13 keep impacting our senses in a wide range of foods, beverages and perfumes. Natural sources
14 of fatty aldehydes are threatened by qualitative and quantitative variability while traditional
15 chemical routes are insufficient to answer the society shift toward more sustainable and natural
16 products. The production of fatty aldehydes using biotechnologies is therefore the most
17 promising alternative for the flavors and fragrances industry. In this review, after drawing the
18 portrait of the origin and characteristics of fragrant fatty aldehydes, we present the three main
19 classes of enzymes that catalyze the reaction of fatty alcohols oxidation into aldehydes, namely
20 alcohol dehydrogenases, flavin-dependent alcohol oxidases and copper radical alcohol
21 oxidases. The constraints, challenges and opportunities to implement these oxidative enzymes
22 in the flavors and fragrances industry are then discussed. By setting the scene on the biocatalytic
23 production of fatty aldehydes, and providing a critical assessment of its potential, we expect
24 this review to contribute to the development of biotechnology-based solutions in the flavors
25 and fragrances industry.

26 Contents

27		
28	Introduction	3
29	1. Aldehydes in the flavors and fragrances industry	4
30	1.1. A brief history of fragrant aldehydes.....	4
31	1.2. Characteristics and origin of commonly used fatty aldehydes	5
32	2. Overview of bio-oxidative catalysts	7
33	2.1. Alcohol dehydrogenases (ADHs).....	9
34	2.1.1. Source of ADHs	9
35	2.1.2. Structure and catalytic mechanism of ADHs	9
36	2.1.3. Substrate specificity of ADHs	10
37	2.1.4. ADHs from extremophiles	11
38	2.1.5. Heterologous production of ADHs.....	12
39	2.2. Flavin-dependent alcohol oxidases (FAD-AOXs)	12
40	2.2.1. Source of FAD-AOXs	13
41	2.2.2. Structure and catalytic mechanism of FAD-AOXs	13
42	2.2.3. Substrate specificity of FAD-AOXs.....	15
43	2.2.4. Heterologous expression of FAD-FAOXs	16
44	2.3. Copper Radical Alcohol Oxidases (CRO-AlcOx).....	16
45	2.3.1. Source of CRO-AlcOx	16
46	2.3.2. Structure and catalytic mechanism of CRO-AlcOx	17
47	2.3.3. Substrate specificity of CRO-AlcOx	18
48	2.3.4. Heterologous expression of CRO-AlcOx	19
49	3. Toward efficient natural biocatalysts	21
50	3.1. Exploration of biodiversity.....	21
51	3.2. Accessory enzymes	22
52	3.2.1. Co-factor recycling systems	22
53	3.2.2. Catalase	23
54	3.2.3. Peroxidase	23
55	3.3. Overoxidation.....	24
56	4. Biocatalytic system engineering.....	26
57	4.1. Protein engineering.....	26
58	4.2. Whole-cell catalysis	27
59	4.3. Biphasic systems and enzyme immobilization.....	28
60	5. Outlook.....	30
61	References	33
62		

63 **Introduction**

64 Flavors and fragrances (F&F) are found in a wide variety of consumer goods: foods and
65 beverages, fine fragrances, pharmaceuticals, cosmetics, soap, household products, etc. Today,
66 more than 3,000 molecules are used to manufacture products that affect our senses of smell and
67 taste (Zviely, 2012). While F&F usually make up only a minor quantitative fraction of the final
68 product, they play a major role in its qualitative perception by consumers. In 2019, the value of
69 the global F&F market reached 31.5 billion U.S. dollars and is predicted to grow by 3.5 % per
70 year up to 2024 (IAL Consultants, 2020). Current industrial production of F&F relies to a large
71 extent on chemical synthesis providing large quantities of material at cost-effective prices.
72 However, chemical synthesis has many drawbacks as (i) it is based on high energy consuming
73 processes, (ii) it requires the use of metallic or toxic catalysts, (iii) it often relies on fossil
74 material (Johannsen et al., 2020), and (iv) it generates large amounts of wastes and thus raises
75 environmental concerns (Braga et al., 2018; Silveira et al., 2019). Chemical processes are
76 usually poorly regio- and enantio-selective resulting in racemic mixtures that require further
77 downstream processing (e.g. purification) to avoid unwanted off-flavors (Berger, 2015; Longo
78 and Sanromán, 2006; Serra et al., 2005). The growing societal environmental concerns go along
79 with a steadily increasing demand for “natural” products (Román et al., 2017), a feature usually
80 not satisfied by established chemical routes. Many F&F are directly extracted from natural raw
81 materials, using empirical processes such as distillation. This approach provides natural
82 products with complex olfactive profiles but suffers from extremely low production yields. For
83 instance, vanilla requires the meticulous manual pollination of each flower (Frey, 2005; Gallage
84 and Møller, 2015). Raw material supply is also vulnerable to environmental (*i.e.* climate
85 variability, pathogens expansion, etc.), economic, social and political fluctuations (Braga et al.,
86 2018). Consequently, there is a growing demand to provide alternatives for the production of
87 natural F&F. Paradoxically, growing demand for food ingredients of natural origin has
88 occasionally led to shortage of some plant resources such as vanilla or peppermint (Dunkel et
89 al., 2014; Malik and Rawat, 2021). The tremendous increase of consumer demand over natural,
90 and sustainable products, combined with the limitations of chemical and extractive routes call
91 for a middle path, which likely lies in the rise of biotechnologies (Berger, 2009; Check Hayden,
92 2014; Pessôa et al., 2019; Schwarz and Wydra, 2020; Silveira et al., 2019; Waltz, 2015).

93 Among F&F compounds, aldehydes constitute a prominent class of molecules broadly
94 used as food products additives (Tsuzuki, 2019). Aldehydes are also key components of
95 perfumes (Surburg and Panten, 2016). Most notably, straight chain (C6 to C13) saturated

96 aldehydes, henceforth denoted as “fatty aldehydes”, occupy a central role in the F&F field (see
97 section 1), and will thus be the focus of the present review.

98 Biotechnological production of aldehydes can be tackled by two approaches: (i) *de novo*
99 synthesis, which relies on the metabolization of simple substrates (e.g. glucose) by natural or
100 engineered organisms to yield the final desired product (this approach will not be covered here
101 but has been reviewed elsewhere (Gounaris, 2010; Kunjapur and Prather, 2015)), or by (ii)
102 biotransformation/bioconversion, which consist in biocatalytic chemical modification of a
103 precursor structurally related to the product through one or several step(s) catalyzed by
104 enzyme(s) (Braga et al., 2018; Pessôa et al., 2019). In the first section of this review, we briefly
105 draw the portrait of aldehydes as key aroma and fragrance molecules. In the second section, we
106 review the main enzymes of interest for oxidation of fatty alcohols, namely: alcohol
107 dehydrogenases (ADHs - EC 1.1.1.1), flavin-dependent alcohol oxidases (FAD-AOXs - EC
108 1.1.3.13) and copper radical alcohol oxidases (CRO-AlcOx - EC 1.1.3.13). We then discuss
109 the available resources to find new biocatalysts and the critical enzymatic and process criteria
110 to consider for implementation of efficient bioprocesses. In the last section, we discuss some
111 problematics regarding transfer from lab-scale to industrial F&F exploitation.

112

113 **1. Aldehydes in the flavors and fragrances industry**

114 **1.1. A brief history of fragrant aldehydes.**

115 First recorded uses of fragrant substances date back to the Antiquity. In that time, odorants
116 materials (*e.g.* tree barks, aromatic plants, resins...) were usually burnt to release and spread
117 their scent. Ancient Egyptians (3150-30 BC) used to prepare cedarwood oil for the embalming
118 process to protect mummies from insects and microbial attacks (Balasubramanian, 2015;
119 Sawamura, 2011). During most of the Human history, essential oils have been the main source
120 of fragrant materials and were often reserved to an elite and/or dedicated to specific usages (*e.g.*
121 religious practices). Essential oils are still today of great importance for perfumers and
122 flavorists. From late 19th century and over the 20th century, a modern era for F&F, marked by
123 the democratization of their use opened up, notably thanks to the advent of both analytical and
124 organic chemistry (Brenna and Parmeggiani, 2017; Sell, 2015; Taylor, 2002).

125 Aldehydes are intimately linked to the early days of fragrance chemistry.
126 Cinnamaldehyde (Chiozza, 1856; Richmond, 1950) and benzaldehyde were some of the first

127 isolated and characterized chemical compounds. The latter being the first perfumery molecule
128 isolated from nature and synthetically produced in 1832 (Wöhler and Liebig, 1832). Natural
129 benzaldehyde from bitter almond, pits of apricots, peaches or plums is recovered in low yields
130 by enzymatic hydrolysis of amygdalin (a cyanogenic glycoside), a process that must be
131 conducted carefully to get rid of the poisonous cyanide (Clark, 1995). (Bio)catalytic production
132 processes yielding simpler and safer product profiles are therefore of obvious interest. In 1874,
133 Tiemann and Haarmann reported the first isolation and chemical synthesis of vanillin (Tiemann
134 and Haarmann, 1874). In 1921, the introduction of a combination at high dosage of the C10,
135 C11 and C12 saturated fatty aldehydes and of 2-methylundecanal, allowed the design of the
136 landmark famous perfume “Chanel N°5”. This perfume truly popularized the use of synthetic
137 ingredients in modern perfumery.

138 **1.2. Characteristics and origin of commonly used fatty aldehydes**

139 Despite their importance as reactive intermediates in various chemical routes (Zhou et al.,
140 2020), the valorization of aldehydes *per se* is rare. Exceptions to this are some low molecular
141 weight aldehydes (*e.g.* acetaldehyde, formaldehyde) and aldehydes used as perfumes,
142 fragrances ingredients and flavoring agents in the F&F industry. In this domain, aldehydes are
143 a staple, ubiquitous in food (Parker, 2015; Tsuzuki, 2019), and necessary for the elaboration of
144 most perfumes (Kohlpaintner et al., 2013), illustrated by a dedicated odor descriptive term –
145 *i.e.* “aldehydic” (Surburg and Panten, 2016). Aldehydes used in F&F are very odorant
146 compounds with a low odor threshold and are therefore highly diluted in final products (Pessôa
147 et al., 2019; Tsuzuki, 2019).

148 Among the short-chain aliphatic aldehydes, acetaldehyde is a key compound in fruity
149 flavoring (Rowe, 2011). Fatty aldehydes \geq C6 display a dual “citrus-like/fatty” character. Their
150 industrial relevance is mainly restricted to C6, & C8-C13 aldehydes (Kohlpaintner et al., 2013;
151 Surburg and Panten, 2016). Indeed, increase of the alkyl chain length entails the predominance
152 of the “fatty”, unpleasant, character, rendering the use of compounds $>$ C13 inappropriate for
153 most applications as F&F.

154 Hexanal (C6) releases a “green-note” odor, reminiscent of a “cut-grass” (Table 1). Fatty
155 aldehydes with even number (C8, C10, C12) occur naturally in the oil glands of citrus peel (*i.e.*
156 in the flavedo) yielding “green”, “fresh” and “citrus-like” notes. In these oils, C8 and C10
157 aldehydes usually provide the “fresh/citrus” character while C12 gives the floral note (Truong
158 et al., 2017). Octanal (C8) is also of importance in orange juice, being one of the main

159 contributors of its flavor (Averbeck and Schieberle, 2009; Porat et al., 2016), Overall, citrus
 160 aromas represent the most important tonality of the flavoring market (20 % of the demand)
 161 (IAL Consultants, 2020).

162 **Table 1: Overview of main fatty aldehydes used as flavors and fragrances.**

Names	Number of C atoms	Chemical structure	Scent	Main natural sources
Hexanal, Caproaldehyde	6		Green, grassy fruity	Occurs in many plants and fruits
Octanal, Caprylaldehyde	8		Green, citrus-like fruity, floral, lemon, green grassy	Sweet orange peel oil
Nonanal, Pelargonaldehyde	9		Soapy, citrus-like, rose-like, floral	Orange and rose oils
Decanal Caprinaldehyde	10		Green, citrus-like, fatty, peelly	Large occurrence in various essential oils: citrus peel oils, neroli oil...
Dodecanal Lauric aldehyde	12		Waxy, fatty, soapy, reminiscent of violets	Citrus and <i>Pinus</i> oils

163

164 Traditional extraction of citrus essential oil is performed by “cold press” extraction (also
 165 referred to as “expression”), by applying physical means to shatter the oil gland and recover the
 166 so-called “peel-oil”, which is the most widely produced essential oil worldwide. Production of
 167 citrus peel oil is usually conducted concomitantly with production of citrus juice. Most of the
 168 global production of these juices relies on Brazilian and Florida citrus crops (Berger, 2007;
 169 Diaz et al., 2005; Porat et al., 2016; Sell, 2015) and the emergence of the citrus greening disease
 170 – Huánglóngbìng disease (lit. “Yellow dragon disease”) – raises concern about our dependency
 171 on this major source of natural molecules. The bacteria *Candidatus Liberibacter* spp. are the
 172 causal agent of this severe disease that affects fruit flavor and causes premature death of the
 173 infected trees (Baldwin et al., 2010; Dala-Paula et al., 2019).

174 Overall, socio-environmental concerns together with the urge for a renewal of supply
 175 sources call for the development of alternative, greener routes for the production of fragrant
 176 aldehydes. Biotechnology and more specifically biocatalysis, offers promising perspectives to

177 tackle this challenge. In the following sections, we present some of the major available
178 oxidative enzymes for the production of fatty aldehydes.



179

180 **2. Overview of bio-oxidative catalysts**

181 The use of enzymes to produce F&F molecules has been for a long time restricted to a
182 small set of reactions. However, over the past decades, considerable improvements have been
183 made in the development of biotechnological solutions and the relevance of the enzymatic
184 approach is now widely recognized (Fasim et al., 2021; Musa and Phillips, 2011). Powerful
185 tools have been developed to improve biocatalysts and their use including computer-assisted
186 evolution of enzymes (Arnold, 2019), immobilization techniques, biphasic systems,
187 thermostability engineering, large-scale recombinant enzyme production and isolation of highly
188 enantio- and regio-selective enzymes (see section 3. and 4.) (Cao et al., 2021; Wu et al., 2020).

189 Using enzymes, aldehydes can be obtained through either an oxidative or a reductive
190 route, starting from an alcohol or a carboxylic acid as precursor substrate, respectively.
191 Carboxylic acid reductases perform the reduction of carboxylic acid to aldehydes at the expense
192 of ATP and NADPH as co-factors. Such expensive organic co-factors constrain their use in
193 whole cell systems. Yet, aldehydes are often toxic for the cells and thus, are further reduced to
194 alcohols or reoxidized to carboxylic acids by endogenous enzymes. The reductive route will
195 not be further discussed here, and readers are directed to some recent reviews on this topic
196 (Derrington et al., 2019; Maurer et al., 2019; Winkler, 2018).

197 Three main classes of enzymes are used for alcohol oxidation and will be discussed
198 hereinafter: ADHs, FAD-AOXs and CRO-AlcOx. On the one hand, ADHs are broadly
199 available, deeply studied and catalyze a reversible reaction using expensive organic
200 nicotinamide cofactor as electron acceptor (commonly NAD(P)⁺). On the other hand, despite
201 being less studied and available, FAD-AOXs and CRO-AlcOx use only molecular O₂ as
202 inorganic final electron acceptor to catalyze the irreversible oxidation of the alcohol into the
203 corresponding aldehyde. Such characteristics entail significant practical differences in the
204 implementation of the biocatalytic systems, as discussed hereinafter. An overview of the main
205 advantages and drawbacks of each biocatalytic system is presented in Figure 1. In Table S1, we
206 provide the reader with more information on the enzymes discussed hereafter, including notably
207 their origin, biochemical properties, 3D structures, kinetic parameters and conversion
208 efficiencies.



209

210 **Figure 1. Comparative overview of (A) the main advantages and drawbacks for each enzyme class**
 211 **discussed in this review and of (B) catalytic efficiencies of some CRO-AlcOx (blue bar), FAD-AOXs**
 212 **(yellow bars) and ADHs (pink bars) toward a range of fatty alcohols (C6-C12).** For each enzyme, the
 213 k_{cat}/K_M values are shown with a different symbol depending on the substrate chain length (C6 to C12,
 214 see legend key in the figure). For illustrative purposes, we show a colored bar in the background that
 215 corresponds to the highest k_{cat}/K_M value measured for each enzyme. An asterisk (*) indicates when an
 216 X-ray crystallographic structure of the enzyme is available. See Table S1 for more details on properties

217 and kinetic parameters of the displayed enzymes Note that on the Y-axis (k_{cat}/K_M) is displayed in
218 logarithmic scale.

219

220 **2.1. Alcohol dehydrogenases (ADHs)**


221 2.1.1. Source of ADHs

222 ADHs are ubiquitous in living organisms (Thompson et al., 2018). ADHs, which are
223 mostly intracellular, are involved in various metabolic pathways. In anaerobic bacteria and
224 yeast, ADHs reduce acetaldehyde to ethanol, whose toxicity inhibits the growth of other
225 microorganisms (Hernández-Tobías et al., 2011) while recycling NAD(P)⁺ (Reid and Fewson,
226 1994). In humans, ADHs promote the opposite reaction to detoxify deleterious ethanol
227 (Edenberg, 2007).

228

229 2.1.2. Structure and catalytic mechanism of ADHs

230 In terms of 3D structure, ADHs share a conserved Rossmann fold for cofactor binding,
231 but otherwise display a high structural diversity. The three main classes of ADHs are
232 distinguished according to sequence length and structural/mechanistic features: short-chain
233 dehydrogenases (SDRs) encompass small ADHs condensed in a single domain while medium-
234 chain dehydrogenases (MDRs), and long-chain dehydrogenases (LDRs), both harbor a co-
235 factor binding domain and a catalytic domain (An et al., 2019). SDRs are metal-free enzymes
236 while MDRs usually make use of two Zn co-factors (one involved in catalysis and the second
237 one playing a structural role – see Figure 2A). LDRs – a more heterogenous class – can require
238 either Zn and/or Fe as co-factor, or sometime no metal ion at all (Sellés Vidal et al., 2018). The
239 Zn-containing ADHs are generally found as dimers (mostly in higher plants and mammals) or
240 tetramers (in bacteria and yeasts) (An et al., 2019). ADHs operate through a hydride transfer
241 from the substrate to the NAD(P)⁺ co-factor with concomitant release of a proton (Figure 2B)
242 (Crichton, 2012). The reaction they catalyze is reversible. These enzymes can both oxidize
243 alcohol or reduce carbonyl compounds (note that some ADHs are also called ketone reductases
244 - KRED), depending on the enzyme and substrate, the oxidation or reduction route can be
245 preferred and the pH of the medium can influence the direction of the reaction (D'Auria, 2002).
246 Yet, from a thermodynamic standpoint, ADH-catalyzed reductive reaction is usually strongly
247 favored over alcohol oxidation (Hollmann et al., 2012).



248

249 **Figure 2. Example of an ADH structure and overview of ADHs catalytic cycle. (A) Cartoon**
 250 **representation of one HADH monomer in complex with NADH** (PDB 2OHX (Al-Karadaghi et al.,
 251 1994)). NADH is displayed as yellow sticks. Zinc (Zn) ions are displayed as blue spheres; the Zn ion
 252 close to NADH plays a role in catalysis, the second Zn ion is structural. Note that HADH structure is
 253 used for illustrative purposes and is not meant to represent the entire, diverse class of ADHs. **(B)**
 254 **Catalytic mechanism of ADH dehydrogenation of primary alcohols** (pink arrows) (and reverse
 255 reduction reaction of aldehydes, shown in blue arrows) coupled to *in situ* NAD(P)H regeneration. In the
 256 NAD(P)⁺/NAD(P)H chemical structure only the nicotinamide group is shown (R corresponds to the
 257 adenine (monophosphate) dinucleotide chain)

258

259 2.1.3. Substrate specificity of ADHs

260 ADHs are the most intensively studied group of oxidoreductases. Among well-studied
 261 ADHs, the prominent horse liver ADH (HADH) was shown to accept 1-octanol as substrate
 262 ($k_{cat}/K_M = 5.8 \times 10^4 \text{ M}^{-1} \cdot \text{s}^{-1}$; Cea et al., 2009; Tani et al., 2000; Whitaker et al., 2002). Yeast ADHs
 263 (YADHs) from *S. cerevisiae* YADH1 and YADH2 were also found active on C2 to C10
 264 alcohols (de Smidt et al., 2008; Dickinson and Dack, 2001; Leskovac et al., 2002; Ottone et al.,
 265 2018; Pal et al., 2009; Schöpp and Aurich, 1976). YADHs are known to be unstable enzymes
 266 (Trivedi et al., 2005). – Of note, both HADH and YADH exist through multiple isoforms. The
 267 isoform will be specified hereinafter when possible. No specification means that the enzyme is
 268 either a mixture of isoforms or that no clear identification was provided in the original reference.
 269 Comparison between HADH, human liver ADH and YADH on a wide range of substrates, was
 270 studied in the 1970s (Pietruszko et al., 1973): HADH exhibited ca. 10-fold higher catalytic
 271 efficiency than both human liver ADH and YADH. Overall, a multitude of studies reported
 272 expression and characterization of new ADHs with observed activity on fatty alcohols (from

273 C6 to C8) often among other substrates included in the initial specificity screening.
274 Unfortunately, most of these studies are restricted to spectrophotometric monitoring of
275 NAD(P)⁺ reduction/NAD(P)H oxidation, and thus lack direct analysis of product formation.
276 Among the characterized ADHs, potent catalysts for oxidation of fatty alcohols very likely
277 exist, but are still untapped as studies for applicability of these enzymes in biocatalysis are
278 predominantly focused on their use for asymmetric reduction processes and for their
279 enantioselective oxidation of secondary alcohols (Dong et al., 2018; Matsuda et al., 2009;
280 Turner, 2011). In particular, the production of ketones using ADHs is of interest for other F&F
281 applications such as the synthesis of carvone (de Carvalho and da Fonseca, 2006; Werf et al.,
282 1999).

283

284 2.1.4. ADHs from extremophiles

285 In this paragraph, we focus on some ADHs from extremophiles with interesting traits.
286 Indeed, while many ADHs from extremophiles have been described, some encompass activity
287 on fatty alcohols (Hirano et al., 2005; Holt et al., 2000; Inoue et al., 2005; Ma et al., 1994; Ma
288 and Adams, 1999; Yanai et al., 2009) and can exhibit attractive biochemical characteristics such
289 as high thermostability and tolerance to solvent, two assets for the oxidation of poorly soluble
290 substrates. A recombinant ADH from *Bacillus stearothermophilus* was used in an amination
291 cascade reaction to convert fatty alcohols to aldehydes and was able to convert 50 mM C6
292 alcohol with a yield > 99%. The enzyme was also able to convert partially C8 (50 % yield) and
293 C10 (2% yield) alcohols, and addition of 1,2-dimethoxyethane (10% (v/v)) as co-solvent
294 improved the conversion of C10 up to 25 % (Sattler et al., 2012). *Thermococcus hydrothermalis*
295 ADH can be driven preferentially toward reductive or oxidative reaction by adjusting the pH of
296 the reaction with pH 10.5 for an optimal alcohol oxidation vs pH 7.5 for an optimal reductive
297 reaction. Interestingly, this enzyme found as a dimer at pH 7.5 forms a tetramer at pH 10.5.
298 This ADH showed best activity on hexanol (>C6 not assayed) and was also able to oxidize
299 some aromatic alcohols (*i.e.* benzyl alcohol) and monoterpenes (*i.e.* geraniol and nerol)
300 (Antoine et al., 1999). The ADH from *Aeropyrum pernix* K1 showed kinetic constants that
301 increased with the alkyl chain length (tested up to C6 with a k_{cat}/K_M of 2.5 mM⁻¹.s⁻¹) and strong
302 thermostability with 24% residual activity after 30 min incubation at 98°C (Hirakawa et al.,
303 2004). A thermostable enzyme from the psychrotolerant *Flavobacterium frigidimaris* KUC-1
304 showed increasing activity from C6 to C10, with high temperature resilience, being active from

305 0 to 85°C (optimum at 70°C) (Kazuoka et al., 2007). However, no heterologous production of
306 the latter enzyme has been described. Alkanes/alkanols-degrading microorganisms can be a
307 source of ADHs active on fatty alcohols as demonstrated by several studies (Álvarez et al.,
308 2011; Barth and Künkel, 2007; Fox et al., 1992; Nagashima et al., 1996; Tassin and
309 Vandecasteele, 1972; Ueda and Tanaka, 1990; Weimer et al., 1993; Yamada et al., 1980).
310 Noteworthy, two ADHs from long-chain alkane-degrading strain *Geobacillus*
311 *thermodenitrificans* NG80-2 exhibited activity toward fatty alcohols including C6-C10 (Liu et
312 al., 2009).

313 314 2.1.5. Heterologous production of ADHs

315 Commercial ADHs (Hummel and Hummel, 1999) mainly originate from either animal
316 tissues (horse liver), *S. cerevisiae* or other recombinant hosts, and can encompass multiple
317 isoenzymes when purified from native host. Despite potential issues pertaining to the generation
318 of aggregated proteins (Utekal et al., 2014), *E. coli* is predominantly used for the heterologous
319 production of ADHs (Höllrigl et al., 2008; Holt et al., 2000; Kim et al., 2019; Liu et al., 2009)
320 in the cytosolic compartment. For instance, the well-studied HADH (EE isoenzyme) has been
321 recombinantly produced in *E. coli* (Huang et al., 2018; Park and Plapp, 1991; Quaglia et al.,
322 2012) but its expression at large scale is still challenging (Hummel et al., 2003; Ottone et al.,
323 2018). Yet, successful recombinant expression of other ADHs in *E. coli* (Wolberg et al., 2001)
324 and utilization at commercial scale have been reported for other applications (Davis et al., 2005;
325 Patel, 2009), after having conducted protein engineering (see section 4.1).

326 **2.2. Flavin-dependent alcohol oxidases (FAD-AOXs)**

327 FAD-AOXs are found in both the large superfamily of glucose–methanol–choline (GMC)
328 oxidoreductases and the superfamily of Vanillyl Alcohol Oxidase/p-cresol methylhydroxylase
329 (VAO/PCMH). Some FAD-AOXs are also classified into the Carbohydrate-Active enZymes
330 database (CAZy, www.cazy.org; Levasseur et al., 2013; Lombard et al., 2014) in the Auxiliary
331 Activity 3 (AA3) and AA4 families. The AA3 CAZy family, initially built for carbohydrate
332 oxidoreductases (e.g., cellobiose dehydrogenase, glucose 1-oxidase), contains
333 phylogenetically-related enzymes active on either aryl alcohols (AA3_2 sub-family) or on small
334 primary alcohols (e.g., methanol, ethanol; AA3_3 sub-family) (Sützl et al., 2018). The AA4
335 family contains VAO that belong to subgroup 4 of the VAO/PCMH superfamily (Ewing et al.,
336 2017). AA4 will not be further discussed in this review because they are specifically active on
337 *para*-substituted phenolic compounds and not active on aliphatic substrates. Yet, these enzymes

338 have been harnessed for other major F&F applications such as the production of vanillin
339 (Lambert et al., 2013, 2014).

340

341 2.2.1. Source of FAD-AOXs

342 FAD-AOXs are widespread (Ewing et al., 2015), and some of the best-known FAD-
343 AOXs originate from peroxisomes of methylotrophic yeasts, such as *Pichia pastoris*
344 (*Komagataella phaffii*), *Candida boidinii* or *Hansenula* sp. where they play a key role in
345 methanol metabolism (Ozimek et al., 2005; Zavec et al., 2021). Alternatively, a number of
346 “long-chain alcohol oxidases” (LCAOs) also called “fatty-alcohol oxidases” (FAO)
347 (EC:1.1.3.20) have been described (Ewing et al., 2015). Owing to heterogenous nomenclature,
348 these enzymes will hereinafter be referred to as “FAD-FAOX”. FAD-FAOXs have been mostly
349 isolated from “alkane-growing” yeasts (Fukuda and Ohta, 2017), fungi (Alvarado-Caudillo et
350 al., 2002; Kumar and Goswami, 2006; Pickl et al., 2015; Savitha and Ratledge, 1991; Silva-
351 Jiménez et al., 2009) and plants (Banthorpe et al., 1976; Cheng et al., 2004; Moreau and Huang,
352 1979; Zhao et al., 2008). Early studies described interesting activities on fatty alcohols, albeit
353 individual enzymes were not isolated/characterized. First evidences of FAD-FAOX enzymes
354 (from yeasts) were described in the 80-90’s (Blasig et al., 1988; Il’chenko, 1984; Il’chenko and
355 Tsfasman, 1988; Krauzova et al., 1986, 1985), followed by studies on the microsomal fractions
356 of *Aspergillus flavipes*, *Candida bombicola*, *C. maltose*, *C. apicola*, *C. tropicalis* and *Yarrowia*
357 *lipolytica* (Hommel and Ratledge, 1990; Hommel et al., 1994; Kemp et al., 1991, 1990, 1988;
358 Mauersberger et al., 1992).


359

360 2.2.2. Structure and catalytic mechanism of FAD-AOXs

361 GMC-type FAD-AOXs are multimeric enzymes usually found as homooctamers
362 (Ozimek et al., 2005; Silva-Jiménez et al., 2009) or sometimes as homodimers (Dickinson and
363 Wadforth, 1992; Heath et al., 2019) in yeasts while fungal enzymes, which are less investigated,
364 can be found as heterodimers (Kumar and Goswami, 2009). Overall, all FAD-AOXs share a
365 conserved FAD binding domain displaying both the sequence motif GxGxxG/A and a
366 Rossmann fold (Dijkman et al., 2013). The flavin co-factor is usually non-covalently bound to
367 the enzymes of the GMC superfamily in contrast to the members of the VAO/PCMH
368 superfamily (Ewing et al., 2020, 2017; Romero et al., 2018). The C-terminal substrate binding

369 domain is more variable, allowing accommodation of different types of substrates (Sützl et al.,
370 2019). Although several crystallographic structures of FAD-AOXs from methylotrophic yeasts
371 have been resolved (see Figure 3A), equivalent information for FAD-FAOXs is still lacking
372 (Cheng et al., 2005). Intriguingly FAD-FAOXs of *Candida* sp. and *Arabidopsis thaliana* were
373 reported to possess a heme binding motif “CXXC” and a possible covalently bound heme
374 (Cheng et al., 2005, 2004; Vanhanen et al., 2000), the function of which remains to be
375 elucidated. The general catalytic cycle of FAD-AOXs can be divided in two half reactions,
376 where FAD plays the role of an electron shuttle. In the reductive half reaction, two electrons
377 are transferred from the substrate to the oxidized flavin *via* a hydride transfer mechanism
378 (Wongnate and Chaiyen, 2013). Then, FADH₂ is regenerated to FAD during the oxidative half
379 reaction *via* the two electrons reduction of O₂ into H₂O₂ (Figure 3B). As stated before, the
380 oxidation of the alcohol is strictly irreversible.

381



382

383 **Figure 3. Example of a FAD-AOX structure and overview of FAD-AOXs catalytic cycle. (A)**
384 **Cartoon representation of a FAD-AOX monomer from *P. pastoris* in complex with FAD (shown**
385 **as yellow stick) (PDB 5HSA). This enzyme is shown as an example and is not meant to represent**
386 **the entire, diverse class of FAD-AOXs. (B) Generic catalytic mechanism of AOX-catalyzed**
387 **oxidation of primary alcohols and H₂O₂ removal by catalase.**

388

389 2.2.3. Substrate specificity of FAD-AOXs

390 The substrate scope of FAD-AOXs is broad with enzymes active on aliphatic alcohols,
391 aromatic alcohols, carbohydrates, steroids, etc. (Ewing et al., 2020). Only enzymes that fit in
392 the focus of this review will be described here. We nevertheless underscore that aryl alcohol
393 oxidases from the AA3_2 subfamily exhibit some activity on unsaturated fatty alcohols
394 (Urlacher and Koschorreck, 2021). Among the FAD-AOXs active on saturated aliphatic
395 alcohols, some of the most prominent members are from methylotrophic yeasts, which are
396 efficient at oxidizing methanol and other short-chain alcohols, while their activity gradually
397 decreases with extension of the alkyl-chain (Ozimek et al., 2005). In aqueous medium, FAD-
398 AOX from *P. pastoris* is found as an homooctamer where the subunits interact via their
399 substrate binding domain (Koch et al., 2016). As a result, the active site appears buried into the
400 octameric structure. These FAD-AOXs are thus not competent catalysts for fatty-alcohols
401 oxidation in aqueous conditions, although their substrate scope can be extended by protein
402 engineering (see section 4.1) or by carrying out reactions in biphasic system (see section 4.3).
403 In contrast, FAD-FAOXs are more efficient on fatty alcohols. Indeed, yeast FAD-FAOX from
404 *Candida tropicalis* exhibited activity on C12 and longer alcohols (but not on C8 and C10)
405 (Cheng et al., 2005). A FAD-FAOXs from another strain of *C. tropicalis* (ATCC 20336)
406 showed activity over C6-C16 primary alcohols, with highest specific activity on C8, followed
407 by C12 alcohols and low K_M on both substrates (Table S1) (Eirich et al., 2004; Eirich and Craft,
408 2007). A plant FAD-FAOX from jojoba (*Simmondsia chinensis*) displayed activity on a range
409 of fatty alcohols, notably C8, and exhibited best performances on C12, with one of the lowest
410 K_M hitherto reported for fatty-alcohols active oxidases (see table S1) (Rajangam et al., 2013).
411 Among fungal FAD-FAOXs, an enzyme purified from the microsomal fraction *Aspergillus*
412 *terreus* (MTCC 6324) was biochemically characterized, showing best activity on C7 and a
413 broad activity on aromatic and secondary alcohols (Kumar and Goswami, 2009, 2008, 2006).
414 We also underscore that a FAD-FAOX from *A. fumigatus* was successfully applied in a cascade
415 reaction where C6-C8 fatty alcohols (10 mM) were fully converted to aldehydes (Pickl et al.,
416 2015).

417

418 2.2.4. Heterologous expression of FAD-FAOXs

419 Heterologous expression of FAD-FAOXs became possible only two decades ago when
420 genes encoding an enzyme from *C. tropicalis* (NCYC 470) and two isoenzymes from *C.*
421 *cloacae* were cloned (Slabas et al., 1999; Vanhanen et al., 2000). Since then, several fungal,
422 yeast and plant FAD-FAOXs have subsequently been successfully produced in *E. coli* (Cheng
423 et al., 2004; Eirich et al., 2004; Rajangam et al., 2013; Takahashi et al., 2016). FAD-FAOXs
424 are multimeric, membrane-bound enzymes showing biophysical complexity (Kumar and
425 Goswami, 2008) which can complicate their heterologous expression and hamper their
426 implementation at large scale. Solubilization of membrane-bound FAD-FAOXs can lead to
427 significant decrease in activity (Kakoti et al., 2012) although it was reported that preservation
428 of activity is possible depending on the enzyme (Cheng et al., 2004). Furthermore, a
429 recombinant FAD-FAOX from *A. fumigatus* (LCAO_af), was successfully employed in a
430 cascade with a ω -transaminase for conversion of fatty alcohols to amines (Pickl et al., 2015).
431 In this study, lyophilized recombinant *E. coli* cells were used to avoid the solubilization and
432 purification of the enzyme and to provide an expression system compatible with the scaled-up
433 production of such FAD-FAOXs. So far, only *E. coli* has been used as recombinant production
434 host and there is no data as to whether the utilization of eukaryotic hosts (e.g. *P. pastoris*) may
435 improve protein expression.

436

437 **2.3. Copper Radical Alcohol Oxidases (CRO-AlcOx)**

438 Copper radical oxidases (CROs) are classified into the CAZy database as members of the
439 Auxiliary Activity 5 family (AA5). They have been further segregated into two subfamilies,
440 namely, AA5_1 containing glyoxal oxidases (GLOX - EC 1.2.3.15) that catalyze the oxidation
441 of aldehydes (most certainly through their hydrated *gem*-diol form) to carboxylic acids, and
442 AA5_2 that gathers enzymes acting on the alcohol moieties of diverse substrate scaffolds to
443 generate aldehydes (see below the “substrate specificity” paragraph for more details). GLOXs
444 were well covered elsewhere (Daou and Faulds, 2017; Kersten and Cullen, 2014), and will
445 therefore not be further discussed here.

446 2.3.1. Source of CRO-AlcOx


447 To date, all characterized CROs originate from filamentous fungi. The only exceptions
448 are a CRO (distantly related to AA5_1) from the filamentous bacterium *Streptomyces lividans*,
449 the substrate preference of which remains to be clarified (Chaplin et al., 2017, 2015; Petrus et al.,

450 2016) and galactose oxidase-like enzymes from *A. thaliana* (Šola et al., 2019; Šola et al., 2021).
451 Within the AA5_2 subfamily, CRO-AlcOx originate from plant pathogens ascomycetes (e.g.
452 *Colletotrichum* sp. or *Magnaporthe* sp.). Recent investigations on the biological context of
453 these enzymes revealed an essential role of *Colletotrichum orbiculare* and *Magnaporthe oryzae*
454 CRO-AlcOx in the penetration of the plant through oxidation of plant cuticular long chain
455 alcohols (Bissaro et al., 2021).

456

457 2.3.2. Structure and catalytic mechanism of CRO-AlcOx

458 Currently, all the structurally characterized CROs belong to the AA5_2 subfamily as no
459 structure of GLOX (AA5_1) is available. CROs are monomeric enzymes with a conserved
460 architecture, displaying a characteristic conserved seven bladed β -propeller fold (“Kelch motif”
461 – Pfam 01344) (Figure 4A). The predominance of β -sheets in their structure has been suggested
462 to contribute to their strong stability (Ito et al., 1991). The catalytic module exhibits a
463 remarkably solvent-exposed active site (Firbank et al., 2001). The rather open and solvent-
464 exposed topology of the CRO active site, which is in striking contrast with FAD-AOX and
465 ADH (Figure 5), may explain why CRO can act on bulky substrates such as fatty alcohols
466 (Ribeaucourt et al., 2021a) or plant waxes (Bissaro et al., 2021). CRO-AlcOx are generally
467 composed of a single catalytic module unlike most other CROs and aside from the CRO-AlcOx
468 from *Magnaporthe oryzae*, that harbors a cell wall integrity and stress response (WSC) module
469 (Oide et al., 2019). The role of this WSC module is still unclear and may not be directly involved
470 in the catalytic mechanism, as its deletion did not affect the activity, nor the specificity of the
471 enzyme (Oide et al., 2019). CROs are monocopper metalloenzymes that catalyze the two-
472 electron aerobic oxidation of alcohols into aldehydes (Figure 4B). No organic co-factor is
473 involved in the catalytic mechanism, which distinguish them from previously discussed ADHs
474 and FAD-AOXs. To catalyze the two electrons transfer, a crosslinked 3’-(S-cysteinyl)-tyrosine
475 free radical (Cys-Tyr[•]) acts in combination with the Cu^{2+} ion, hence the term “copper radical”.
476 CROs rely on a characteristic “ping-pong” mechanism composed of a reductive half-reaction –
477 i.e. substrate oxidation and reduction of the active site – and the complementary reverse
478 oxidative half-reaction – i.e. reoxidation of the active site and concomitant reduction of O_2 to
479 H_2O_2 (Figure 4B). CROs can also be found in an inactive “semi-reduced” form which could
480 increase their stability or be involved in activity regulation *in vivo* (Bissaro et al., 2021).



481
 482 **Figure 4. Example of a CRO-AlcOx structure and overview of CRO-AlcOx catalytic cycle. (A)**
 483 **Cartoon representation of *CgrAlcOx* (PDB 5C92; (Yin et al., 2015)). Copper is displayed as orange**
 484 **sphere. Active sites residues (including copper coordinating residues) are shown as yellow sticks. (B)**
 485 **Generic catalytic mechanism of CRO-catalyzed oxidation of primary alcohols, H₂O₂ consumption**
 486 **by catalase and CRO activation by a peroxidase.**

487
 488 2.3.3. Substrate specificity of CRO-AlcOx


489 CROs have been for long restricted to very few archetypal members and their scope of
 490 application accordingly limited. Indeed, since their discovery in 1959, the founding member of
 491 CROs, namely the galactose oxidase (EC 1.1.3.9) from *Fusarium graminearum* (*FgrGalOx*)
 492 has been the main object of most studies. Numerous applications have been described for the
 493 *FgrGalOx* and the boundaries of its substrate specificity have been pushed far beyond its natural
 494 carbohydrate preference (Birmingham and Turner, 2018; Deacon et al., 2004; Escalettes and
 495 Turner, 2008; Herter et al., 2015; Matthey et al., 2020; McKenna et al., 2015; Sun et al., 2002;
 496 Vilím et al., 2018; Wilkinson et al., 2004). It is only recently, with the advent of natural diversity
 497 exploration, notably rendered possible by the availability of abundant genomic data, that the
 498 true potential of CROs scope of action was unveiled (Andberg et al., 2017; Mathieu et al., 2020;
 499 Mollerup et al., 2019; Paukner et al., 2015, 2014; Yin et al., 2015). Remarkably, a subgroup of
 500 AA5_2 CROs, so-called CRO-AlcOx, appeared to be of particular relevance for fragrant
 501 aldehydes production (Forget et al., 2020; Oide et al., 2019; Ribeaucourt et al., 2021b, 2021a;
 502 Yin et al., 2015). These enzymes, mainly isolated from *Colletotrichum* species, all exhibit a
 503 similar substrate specificity profile and comparable efficiencies. They can oxidize a large panel

504 of primary alcohols, aryl alcohols and activated aliphatic alcohols being excellent substrates.
505 Importantly, CRO-AlcOx can also convert non-activated alcohols with increasing affinity
506 towards longer alkyl chains (Ribeaucourt et al., 2021a; Yin et al., 2015). For instance, 1-hexanol
507 and 1-octanol (3 mM) could be fully converted by the AlcOx from *Colletotrichum graminicola*
508 (*CgrAlcOx*) and *C. destructivum* (*CdeAlcOx*) within 15 min and under mild conditions (23°C,
509 in aqueous media) (Ribeaucourt et al., 2021b, 2021a). Despite stable/increasing catalytic
510 efficiencies up to C10, conversion of C10 alcohol was only partial compared to C6-C8
511 (unpublished results; see Table S1), which might be the result of increasing insolubility of the
512 substrate or stronger inhibition of *CgrAlcOx* by the hydrated form of the aldehyde (Scheme 1)
513 (Forget et al., 2020; Ribeaucourt et al., 2021a). Preparative-scale conversion was also
514 demonstrated using a crude preparation of *CgrAlcOx* (supernatant of a recombinant *P. pastoris*
515 strain) yielding 0.72 g of octanal from 2 g of 1-octanol (Ribeaucourt et al., 2021a).

516

517 2.3.4. Heterologous expression of CRO-AlcOx

518 AA5_2 are secreted fungal enzymes and necessitate post-translational modification such
519 as the formation of a thioether bond for the Cys-Tyr^{*} co-factor. *E. coli* and *P. pastoris* have been
520 the main expression systems used, although *Aspergillus nidulans* (Baron et al., 1994;
521 McPherson et al., 1993), *A. oryzae* and *Fusarium venenatum* (Xu et al., 2000) have also been
522 reported as possible hosts for the archetypal *FgrGalOx*. Detailed comparison of *E. coli* and *P.*
523 *pastoris* expression systems have been described elsewhere (Spadiut et al., 2010) for the
524 *FgrGalOx*. Poor expression yields were obtained in *E. coli* unless a custom-variant strain was
525 used. To date, satisfactory yields of CRO-AlcOx have exclusively been obtained using *P.*
526 *pastoris* as host. Scaled-up production in *P. pastoris* has been developed in bioreactor for both
527 *CgrAlcOx* and *FgrGalOx* (Anasontzis et al., 2014; Ribeaucourt et al., 2021a). Post translational
528 activation of the enzyme is usually done by incubation with copper under aerobic conditions.
529 This process can be done directly *in situ* by adding copper salt into the expression medium
530 (Spadiut et al., 2010; Yin et al., 2015). CRO-AlcOx, like other CROs, are usually recovered
531 after purification in their “semi-reduced” inactive state and must therefore be oxidized to
532 generate the fully active form prior to utilization (Forget et al., 2020; Whittaker, 2005;
533 Whittaker and Whittaker, 1988). This activation is usually achieved by incubation with a
534 peroxidase (EC 1.11.1.7) (see section 3.2.3).



535

536 **Figure 5. Surface views of crystal structures of (A) *A. pernix* ADH active site (pdb 1H2B)**
 537 **(chain A) in complex with octanoic acid (inhibitor; shown as green stick); (B) *S. cerevisiae***
 538 **YADH1 active site (pdb 4W6Z) (chain A) in complex with trifluoroethanol (shown as green**
 539 **stick); (C) *C. graminicola* CRO-AlcOx (pdb 5C92 – copper ion shown as orange sphere) in**
 540 **complex with 1-octanol (shown as green stick; docking model adapted from Ribeaucourt et al.,**
 541 **2021a); (D) *P. pastoris* FAD-AOX (pdb 5HSA) homooctamer. The red arrow indicates the**
 542 **position of the buried substrate binding cavity. For all enzymes, surface was colored according**
 543 **to the ratio of non-polar/polar (NPP) residues using Protein-Sol web tool ([https://protein-](https://protein-sol.manchester.ac.uk/)**
 544 **[sol.manchester.ac.uk/](https://protein-sol.manchester.ac.uk/); Hebdtich et al., 2017). The NPP ratio scale, indicative of increasing**
 545 **hydrophobicity level from 0.6 to 2.3, displayed at the bottom left hand corner of panel C applies**
 546 **to all panels. Note that the structures displayed are not shown at the same scale.**

547 **3. Toward efficient natural biocatalysts**

548 Nature remains the main source of enzymatic diversity providing a wide range of
549 ingenious “ready-to-use” catalysts. In this paragraph, we tackle the question of the discovery
550 and exploitation of efficient natural fatty alcohols-active biocatalysts.

551 **3.1. Exploration of biodiversity**

552 In the quest for suitable catalysts, the exploration of biodiversity can prove to be a
553 powerful approach. A great potential, still largely untapped, exists in the microbial kingdom,
554 as exemplified by the discovery and application of FAD-FAOXs from alkane-growing yeasts
555 and fungi (Kemp et al., 1991, 1988; Pickl et al., 2015) (see section 2.2). Diversity among CROs
556 was also identified while investigating phylogenetical differences in fungal sequences of the
557 GalOx family (Andberg et al., 2017; Mathieu et al., 2020; Yin et al., 2015). The tremendous
558 amount of genomic information available today (Sharon and Banfield, 2013) combined with,
559 powerful bioinformatics/modeling tools and platforms for high throughput (HTP) recombinant
560 expression (Haon et al., 2015; Vincentelli, 2019) allow to generate wide libraries of natural
561 enzyme candidates. Once access to enzyme diversity is achieved, the next most challenging
562 task is most likely the design of a suitable, HTP-compatible screening method to detect desired
563 substrate specificities and/or properties. Most generic HTP methods are often indirect and fail
564 to provide fine details as to the sought-after reaction products. Getting a higher level of
565 information is often at the expense of the throughput frequency. For instance, for substrate
566 screening targeting fatty alcohol oxidases, the well-established ABTS (or equivalent Amplex
567 Red)/horseradish peroxidase (HRP) coupled assay can be used. This assay has the advantage of
568 enabling real-time kinetic monitoring of the enzymatic reaction. However, extensive controls
569 are required to avoid false positive/negative reactions owing to the indirect nature of the assay
570 and to the reactivity of the radicals generated. Other indirect assays based on the detection of
571 H₂O₂ exist such as the xylenol orange-based assay (Ewing et al., 2018), this method avoids the
572 use of a secondary enzyme, but does not allow direct real-time monitoring of the reaction. A
573 direct assay detecting the production of aldehydes would be more suitable, and applicable to all
574 classes of enzymes. Commercial colorimetric assays are available and an HTP method using
575 amino benzamidoxime has been developed recently and validated on CARs enzyme (Horvat et
576 al., 2020; Ressmann et al., 2019). A HTP-derivatization strategy coupled to liquid- and
577 supercritical fluid-chromatography was also implemented for screening of GalOx variants,
578 which enabled product analysis. However, this method requires a more complex procedure

579 (derivatization) and equipment (Huffman et al., 2019). Redirecting “nature’s technology” for
580 our needs require thoughtful reaction set-up. Parameters that can significantly affect the
581 efficiency of the biocatalysts for fatty aldehydes production (co-factor regeneration, need for
582 accessory enzymes, overoxidation) are discussed hereafter.

583 3.2. Accessory enzymes

584 Alcohol dehydrogenases and oxidases often require the use of accessory enzymes to
585 harness the best of their ability. An overview of the most important accessory enzymes is given
586 hereinafter.

587 3.2.1. Co-factor recycling systems

588 One of the main hurdles for the industrial application of ADH is the need for expensive
589 organic co-factors (*e.g.* NAD(P)⁺) where stoichiometric addition is not an affordable option.
590 Furthermore, the oxidation of alcohols is thermodynamically disfavored compared to the
591 reduction of aldehydes by ADH (Hummel and Hummel, 1999). Both problems can be solved
592 by *in situ* co-factor regeneration. This can be achieved by chemical approaches, although
593 coupled enzyme systems are most commonly applied. Yet, the enzymatic regeneration of the
594 oxidized form of the cofactors (NAD(P)⁺) – needed for ADH-catalyzed oxidation reaction (Fig.
595 2) – is less investigated than the regeneration of the reduced cofactors (NAD(P)H) (useful when
596 running ADH in “reductase mode”). This lack is often a critical obstacle in the implementation
597 of ADH-catalyzed oxidation in industry. Among the enzymes described for this purpose,
598 glutamate dehydrogenase, lactate dehydrogenase, and NAD(P)H oxidases (NOX) are the most
599 prominent ones. In particular, NOX have attracted a lot of attention in the past years. They can
600 directly oxidize the reduced cofactor using oxygen as co-substrate and releasing H₂O or H₂O₂.
601 Recently, the use of an ADH/NOX coupled-system was demonstrated at pilot-scale for the
602 oxidation of lactols (Bartsch et al., 2020), with the use of an engineered NOX enzyme
603 (Petschacher et al., 2014). Alternatively, regeneration can be achieved by a substrate-coupled
604 approach, where the ADH itself reoxidizes the co-factor by reducing a secondary substrate (see
605 Figure 2). To this end, a cheap secondary substrate must be chosen and added in excess (*e.g.*
606 acetone, which also acts as substrate-solubilizing agent) (Höllrigl et al., 2008). Care must be
607 taken to avoid enzyme inhibition or any other drawbacks for downstream processes by these
608 additional compounds. Case to case study is essential to select the most suitable approach. For
609 instance, Vrsalović and colleagues studied the conversion of 1-hexanol by either isolated
610 YADH1 or with permeabilized baker’s yeast cells (Vrsalović Presečki et al., 2012; Vrsalović

611 Presečki and Vasić-Rački, 2009). Substrate-coupled regeneration was probed, but the
612 “sacrificial” oxidant (i.e. acetaldehyde) strongly inhibited the isolated YADH and utilization of
613 a coupled enzyme system was more suitable. More exhaustive coverage of recycling systems
614 can be found in recent reviews (Liu et al., 2018; Puetz et al., 2020; Rehn et al., 2016; Wu et al.,
615 2013).

616 3.2.2. Catalase

617 Both, FAD-AOXs and CRO-AlcOx, as part of their catalytic cycle, release H₂O₂ as
618 byproduct, which, when accumulated, is generally deleterious for the enzyme and must be
619 therefore eliminated. Catalases (EC 1.11.1.6), which belong to a catalytically very efficient
620 class of enzymes, are already used at industrial scale in textile bleaching processes (Lončar and
621 Fraaije, 2015), and are widely applied to circumvent AOXs inactivation. Catalases conveniently
622 recycle H₂O₂ into H₂O and ½ O₂, supplying back molecular oxygen which can be rate-limiting
623 in the reaction due to its limited solubility in aqueous media (0.25 mM at room temperature).
624 Yet, some enzymes seem to possess an intrinsic resistance to H₂O₂ as described for a
625 microsome-bound preparation of a FAD-FAOX from *A. terreus*, able to convert ≥85% of C7-
626 C12 alcohols (10 mM) without any addition of catalase (Kakoti et al., 2012).

627 3.2.3. Peroxidase

628 CROs are usually recovered in their inactive “semi-reduced” form and require activation
629 prior to utilization. Peroxidases or chemical oxidants (*e.g.* ferricyanide, ammonium
630 hexachloroiridate, Mn(III) compounds etc...) can be used to this end (Forget et al., 2020;
631 Hamilton et al., 1978; Johnson et al., 2021; T. Pedersen et al., 2015). However, these chemical
632 compounds are often less efficient than peroxidases and are required in large amounts, Thus, to
633 date no equivalent alternative to peroxidase has been found and more investigations are still
634 needed. The HRP is the most common enzyme used as activating agent, mainly for its
635 commercial availability and stability in reaction mixtures. Of note, recent work demonstrated
636 that natural fungal peroxidases are better activators of CRO-AlcOx than HRP (Bissaro et al.,
637 2021). Strikingly, although the activation of CRO by peroxidases has been known for long, the
638 underlying mechanism remains obscure. Current state of the art points toward an oxidative
639 activation mechanism independent of any peroxidase catalytic activity but most likely mediated
640 by a protein-protein interaction (Bissaro et al., 2021; Forget et al., 2020). Indeed, no peroxidase
641 substrate seems involved into the activation, which suggest that chemical oxidants may act in a
642 similar way. We underscore that the required amount of peroxidase greatly varies depending

643 on the substrates: only minimal amounts are needed for activated alcohols while stoichiometric
644 amounts (relative to the AlcOx) or more are required for oxidation of non-activated alcohols
645 (Ribeaucourt et al., 2021a). The necessity to add large amount of peroxidase to foster the
646 reaction is unambiguously the most important barrier for the implementation of CROs-
647 catalyzed oxidation of fatty alcohols at industrial scale. Peroxidases are expensive enzymes as
648 their recombinant production remains challenging. Deeper understanding of this mechanism is
649 needed to provide access to alternative activation methods. Very recently, an electrochemical
650 activation method was investigated to substitute the peroxidase (Zhang et al., 2021). This
651 approach proved to be efficient for the conversion of various aromatic alcohols and aliphatic
652 polyols catalyzed by several *FgrGalOx* mutants.

653

654 **3.3. Overoxidation**


655 Overoxidation of aldehydes into carboxylic acids is frequently observed and affect – in a
656 substrate-, pH and enzyme-dependent manner – every class of enzymes aforementioned,
657 namely ADHs (Hinson and Neal, 1975; Hollmann et al., 2011; Velonia and Smonou, 2000),
658 FAD-AOXs (Ferreira et al., 2010; Martin et al., 2020; Romero and Gadda, 2014) and CROs
659 (Birmingham and Turner, 2018; Kelleher and Bhavanandan, 1986; Parikka and Tenkanen,
660 2009). The aldehydes cannot be oxidized directly by these enzymes, as their H atom is not
661 abstractable *per se* and need an adduct brought either by the enzyme (Olson et al., 1996), or by
662 the environment. In most cases, the observed oxidation of aldehydes is in fact directly
663 dependent on the propension of the aldehyde to undergo hydration to form the corresponding
664 *geminal*-diol (*gem*-diol) (Scheme 1) (Dong et al., 2018) and the subsequent ability of the
665 enzyme to accept the latter as a substrate. Destabilization of the dipole at the carbonyl group
666 favor nucleophilic addition of H₂O, consequently shifting the equilibrium toward the hydrated
667 species. Electron-withdrawing groups will thence favor *gem*-diol, while electron-donating
668 groups or resonance effect should stabilize the carbonyl group (Bell, 1966; Hutchings and
669 Gasteiger, 1986). Aldehydes bearing an alkyl group, like fatty aldehydes are known to be prone
670 to hydration (Buschmann et al., 1982, 1980; McClelland and Coe, 1983).

671 Aldehyde overoxidation and disproportionation (i.e., when two molecules of aldehyde
672 yields one molecule of acid and one molecule of alcohol, also known as “Cannizzaro reaction”)
673 by ADHs (Velonia and Smonou, 2000) has been investigated for some applications in
674 pharmaceuticals or food additives (Contente et al., 2020; Könst et al., 2012; Tassano et al.,

675 2018). Production of carboxylic acid has been reported early for the prominent HADH,
 676 exhibiting a different behavior depending on the substrate. HADH oxidation of ethanol resulted
 677 in a mixture of aldehyde and carboxylic acid, while for 1-octanol, octanoic acid was the only
 678 detectable end-product (Hinson and Neal, 1975, 1972; Oppenheimer and Henehan, 1995).

679 Overoxidation in FAD-AOXs-catalyzed reactions must also be carefully considered.
 680 Indeed, in a recent study, a mutant of the *Phanerochaete chrysosporium* FAD-AOX (*PcAOX*)
 681 yielded >99 % of carboxylic acid when oxidizing C6 alcohol (Martin et al., 2020). In contrast,
 682 a choline oxidase variant (*AcCO6*) engineered for fatty alcohol oxidation (see section 4.1)
 683 (Heath et al., 2019) revealed only traces of acid for C6 and between 1.5 - 8 % for C7-C12 (Table
 684 S1) despite expectable strong activity on *gem*-diols (Gadda, 2008). Very few studies on LCAO
 685 included products analyses and, therefore, overoxidation has rarely been evaluated. The
 686 aforementioned study on LCAO_Af revealed no overoxidation for C4 to C8 but led to 5-26%
 687 carboxylic acid for C9 to C11 (Pickl et al., 2015).

688 Overoxidation was also monitored when using CRO-AlcOx and in this case, different
 689 mechanisms were observed. While a *gem*-diol dependent, CRO-catalyzed overoxidation was
 690 monitored for benzyl substrates, the overoxidation of fatty aldehydes was rather triggered by
 691 exposure to HRP.



692

693 **Scheme 1: Aldehyde hydration and overoxidation routes.** Formation of *gem*-diol upon
 694 aldehyde hydration in acidic or alkaline conditions and subsequent enzymatic oxidation of the
 695 *gem*-diol to carboxylic acid.

696 Overoxidation can be circumvented or limited by different level of control on reaction
 697 conditions. The substrate structure will firstly give indication on the propension of the aldehyde
 698 to undergo hydration and subsequent probability of overoxidation. In terms of hydration rate,
 699 *gem*-diols seem readily generated in a matter of seconds or less upon exposure to H₂O at room

700 temperature (Hilal et al., 2005). Setting the pH of the reaction around neutrality, when possible,
701 could slow down the rate of *gem*-diol formation (Doussin and Monod, 2013). *Gem*-diols are
702 often disfavored substrates compared to the initial alcohols, especially for enzymes with
703 preferences upon primary alcohols, as *gem*-diols are structurally similar to secondary alcohols.
704 Thus, tight control of alcohol concentration and reaction time may be one of the main levers to
705 avoid the formation of unwanted carboxylic acid (Ribeaucourt et al., 2021a). The
706 implementation of a biphasic system can also be an alternative to remove the aldehyde from the
707 aqueous phase and prevent its hydration (see section 4.2).

708 Aside from the enzymatic-catalyzed overoxidation, intrinsic reactivity of aldehydes can
709 result in autooxidation (Marteau et al., 2013; Sankar et al., 2014) or side reactions (e.g.
710 formation of hemi-acetals) which require careful controls of reaction conditions and product
711 validation.

712

713 **4. Biocatalytic system engineering**

714 **4.1. Protein engineering**

715 Protein engineering is a complimentary approach to the exploration of biodiversity
716 (section 3.1), enabling to enhance or reprogram a natural enzymatic starting scaffold for a
717 targeted process (Woodley, 2019). Native enzymes are designed for *in vivo* finely tuned
718 reactions. Early engineering approaches mostly relied on rational engineering based on
719 structural and mechanistic information (Wilkinson et al., 1984). Directed evolution of enzymes
720 though the introduction of random diversity was then widely adopted using error-prone PCR
721 and DNA shuffling (Chen and Arnold, 1993; Cobb et al., 2013; Stemmer, 1994). The large
722 libraries of generated mutants require HTP screening methods raising similar problematics than
723 for biodiversity screening. The field is now rapidly evolving toward semi-rational approaches,
724 harnessing structure/mechanistic information and computational modeling to focus the directed
725 evolution effort on the most relevant residues (Vogel, 2019). The resulting “smart” libraries are
726 smaller allowing to apply more specific detection methods, closer to the final sought-after
727 properties (Lutz, 2010; Sheldon and Brady, 2019).

728 In the context of this review, enzyme engineering is an asset to (i) improve substrate and
729 organic solvent tolerance, (ii) improve thermal stability, (iii) switch enzyme specificity to non-
730 natural substrates, (iv) improve expression, (v) adapt enzyme to specific conditions (pH,

731 temperature, presence of inhibitors) etc. For instance, a choline oxidase from *Arthrobacter*
732 *cholorphenolicus* was recently engineered using structure-guided evolution (Heath et al., 2019).
733 With six mutated residues, the AcCO6 variant gained activity toward 1-hexanol and a broad
734 array of saturated, unsaturated, aliphatic, cyclic and benzylic alcohols (Heath et al., 2019).
735 Notably, C6–C10 unsaturated alcohols were identified as good substrates and were converted
736 (with little overoxidation) in 24 h at 30°C in air-saturated buffer, without any added catalase.
737 This enzyme variant also displayed improved thermostability and solvent tolerance as
738 demonstrated by the implementation of biphasic biotransformation with heptane.

739 The enlargement of the substrate binding pocket has been successfully applied on several
740 enzymes of interest to increase their activities towards bulky primary alcohols. With only two
741 single-point mutations on YADH1, kinetic constants were improved by one order of magnitude
742 on octanol, ($k_{cat}/K_M = 5.5 \times 10^4 \text{ M}^{-1} \text{ s}^{-1}$) for the T48S-W39A variant), reaching a catalytic
743 efficiency comparable to HADH (EE isoenzyme) (Green et al., 1993). Similarly, enlargement
744 of the substrate pocket of *PcAOX* enabled oxidation of 1-hexanol and diols, when the wild-
745 type enzyme showed only poor activity on these substrates (Linke et al., 2014; Martin et al.,
746 2020; Nguyen et al., 2018).

747 The CRO *FgrGalOx* enzyme has been the subject of many engineering studies, bending
748 its properties far from its initial characteristics, yielding for instance, enantioselective oxidation
749 of secondary alcohols or glucose-6-oxidase activity (Escalalettes and Turner, 2008; Sun et al.,
750 2002) (see section 2.3.2). The *FgrGalOx* was also recently engineered to catalyze the first step
751 of a cascade to produce a drug for HIV treatment (Huffman et al., 2019). Twelve rounds of
752 mutations starting from a previously generated variant (Rannes et al., 2011) were run leading
753 to 34 mutated residues in the evolved enzyme enabling an 11-fold improvement for 2-
754 ethynylglycerol oxidation and reversing the enzyme enantio-specificity. This study, among the
755 numerous studies performed on the *FgrGalOx*, let envision a great potential for CRO-AlcOx
756 enzymes, which naturally possess attractive features for fatty alcohol conversion and can
757 benefit from the knowledge gathered on structurally/phylogenetically related CROs.

758

759 **4.2. Whole-cell catalysis**

760 When considering industrial applications, whole-cell and isolated enzymes approaches
761 must be put into balance (Hoyos et al., 2017). The whole-cell approach is attractive for some
762 enzymes like ADHs that require *in situ* supply and regeneration of a cofactor (Romano et al.,

763 2012; Sheldon and Brady, 2019). Implementation of multi-enzymatic systems into a chassis
764 through synthetic biology is a powerful approach to reprogram metabolic pathways, as
765 exemplified by recent work on the production of sclareol in pathway-engineered *E. coli* (Schalk,
766 2014; Schalk et al., 2012). Yet, this approach has inherent drawbacks such as low enzyme
767 activity levels, cell membrane permeability-related issues, growth inhibition by substrate or
768 product toxicity etc. (Garzón-Posse et al., 2018). Indeed, aldehydes are particularly hard to
769 produce through this approach as they are undesirable intermediates for most cells and are often
770 readily converted to carboxylic acids or alcohols by endogenous enzymes (Liu et al., 2018).
771 Two studies using permeabilized *S. cerevisiae* cells producing YADH outlined such
772 difficulties: the conversion of C6 alcohol yielded only hexanoic acid due to overoxidation
773 attributed to the endogenous aldehyde dehydrogenase (Vrsalović Presečki et al., 2012;
774 Vrsalović Presečki and Vasić-Rački, 2009). Whole-cell aldehyde production for F&F has been
775 recently reviewed elsewhere (Ben Akacha and Gargouri, 2015; Kunjapur and Prather, 2015)
776 and will not be further discussed here.

777


778 **4.3. Biphasic systems and enzyme immobilization**

779 Biphasic systems are attractive to trap the produced aldehydes in the organic phase and
780 avoid their hydration and subsequent overoxidation, especially for hydrophobic fatty aldehydes
781 (Gandolfi et al., 2001; Heath et al., 2019). Biphasic systems are also a convenient way to (i)
782 overcome low water solubility of the substrates, (ii) limit product inhibition and (iii) facilitate
783 product recovery through simple distillation (Domínguez de María and Hollmann, 2015). The
784 use of biphasic systems to avoid overoxidation can yet be limited, when hydration occurs
785 directly within the active site as reported for a FAD-AOX (Gadda, 2003; Salvi et al., 2014).
786 Another limitation of these systems lies on the intrinsic enzyme tolerance to the water-
787 immiscible solvent, their related resistance to interfacial tension, the rate of phase transfer, and
788 the level of O₂ available in the case of AOXs. However, in some cases, utilization of an organic
789 layer can improve or modify the activity of a given enzyme. The microsomal fraction of *C.*
790 *tropicalis* was for instance much more efficient in oxidizing C10 and C12 alcohols in an
791 hydrophobic organic solvent containing 1-10% water as compared to aqueous media (Kemp et
792 al., 1991). The FAD-AOX from *P. pastoris* – active on methanol and other short-chain alcohols
793 – has been used for conversion of 1-hexanol and bulkier substrates thanks to the implementation
794 of biphasic systems (Karra-Chaabouni et al., 2003; Murray and Duff, 1990). Noteworthy, the

795 substrate itself can sometimes be employed as the organic layer (de Almeida et al., 2019). Using
796 naturally resilient or engineered enzymes, reactions can be run in water-saturated organic
797 medium.

798 Another alternative to improve enzyme industrial applicability is immobilization.
799 Immobilization techniques are intended to facilitate downstream processes, improve turnover
800 number by enhancing catalyst stability and reusability. They also can be applied to improve
801 resistance of enzymes to organic solvents. Some examples are given elsewhere (Basso and
802 Serban, 2019; Pessôa et al., 2019). Immobilization of ADH enzymes has been extensively
803 studied (Bolivar et al., 2006; Vasić et al., 2020) but with the use of fatty alcohols as substrates
804 is less common (Cea et al., 2009). YADH immobilized on glyoxyl agarose crosslinked with a
805 polymer improved its stability and its activity on fatty alcohols (Ottone et al., 2018). A study
806 conducted in 2010 combined immobilization of HADH and YADH and the use of organic
807 solvents, including the use of the substrate itself as solvent (Snijder-Lambers et al., 2010). In
808 organic medium, YADH showed better activity on bulkier substrates compared to aqueous
809 environment. Immobilization of YADH on amberlite XAD-8, resulted in 93-99 % loss of
810 activity despite optimization attempts, while immobilized HADH remained fully active
811 (Snijder-Lambers et al., 2010). The aforementioned FAD-AOX AcCO6 has been immobilized
812 on an affinity resin *via* the His-tag used for purification (Thompson et al., 2019) and on an
813 epoxy resin (Mattey et al., 2020), improving its reusability. Only one study reported the
814 immobilization of a FAD-FAOX (isolated from microsomal fraction of *A. terreus*) into a
815 polyurethane foam matrix but resulted in only 20% conversion of C7 alcohol (Kakoti et al.,
816 2012). Immobilization studies were performed on the M₁ and M₃₋₅ variants of the CRO
817 *FgrGalOx* and improved their thermal stability, reusability and organic solvent tolerance
818 (Mattey et al., 2020). To our knowledge, no attempts have yet been made for CRO-AlcOx. It is
819 likely that immobilization of CRO-AlcOx and FAD-FAOX could greatly enhance their use in
820 bi-phasic system or hydrophobic medium, avoiding solubility and overoxidation issues.

821 The main problematics and possible solutions (discussed in sections 3. and 4.) to consider
822 for the applications of ADHs; FAD-AOXs or CRO-AlcOx for the oxidation of fatty alcohols
823 are summarized in Figure 6.



824

825 **Figure 6. Overview of some challenges and possible solutions for implementation of**
 826 **enzymatic oxidation of fatty alcohols.** While bringing answers to certain challenges, some
 827 solutions might cause other unwanted drawbacks that are represented by red dashed lines.

828

829 5. Outlook

830 The current state of the art reveals that a great number of enzymes candidates already
 831 exists for the challenging oxidation of non-activated fatty alcohols. Some clear trends stand out
 832 (see Figure 1):

- 833 (i) ADHs constitute a tremendous reservoir of enzymes but scarce information exists
 834 on their biocatalytic performances for fatty alcohols turnover. The need for a
 835 nicotinamide co-factor adds complexity to this biocatalytic system.
- 836 (ii) CRO-AlcOx are promising catalysts with well-established large-scale recombinant
 837 production but require a peroxidase activation step for oxidation of unactivated
 838 alcohols. Research on CRO-AlcOx is still in its infancy since their discovery in 2015
 839 by Yin et al. and better understanding of these microbial enzyme systems could

840 bring alternative activation methods. The recent success in activating some
841 *FgrGalOx* mutants by electrochemical means (Zhang et al., 2021), open a new
842 avenue to overcome the need of peroxidase and could be assessed on CRO-AlcOx
843 likewise.

844 (iii) FAD-FAOX might be today the best suited enzymes for oxidation of fatty alcohols.
845 They display high catalytic efficiencies toward fatty alcohols (Figure 1B) and only
846 require the “industrial-compatible” catalase as accessory enzyme. However, only
847 scarce information is available regarding their large-scale recombinant production
848 and their performances in biocatalytic systems.

849 To foster the translation of fatty aldehydes production at industrial scale, we draw the
850 attention towards some crucial criteria that must be taken into account early in the research and
851 development process: (i) Culture media used for growth of microorganisms should be simple
852 and composed of low-cost ingredients (e.g. malt). Complex bioreactor media supplemented
853 with vitamins or metals are costly when shifting to large scale processes for F&F and any
854 additional compounds (chemicals, metals, accessory enzymes, etc.) will increase reaction costs
855 and must be balanced with substantial gain in yields. (ii) Upscaled reactions in large tank
856 reactors (e.g. 30-80 m³) require important investments and solutions that are easily set up at lab
857 scale (e.g., work in pressurized reactor, addition of large volume or organic solvent, etc.) can
858 be difficult to transfer to industry or must be useful for multiple processes to justify new
859 investments. (iii) The duration of the process has also a direct effect in the cost. Long-time
860 immobilization of the material will impair profitability and could increase the probability of a
861 contamination. Consequently, every step must be optimized: culture of the microorganism,
862 production and recovery of the biocatalyst, conversion reaction and product recovery. (iv) The
863 extraction procedure is also crucial and full conversion of the substrate is thus highly desirable
864 to simplify downstream processes.

865 In the future, the biotechnological production of fatty aldehydes will increase to secure
866 the production using greener systems. To set up a sustainable approach, natural sources of fatty
867 alcohols must be considered. Although cheap chemical sources are available (Surburg and
868 Panten, 2016), natural sources of fatty alcohols are very limited today. Some new reservoirs
869 could arise from forestry by-products and from the pulp and paper industry (Corberán et al.,
870 2014) and alternative sources could be established by implementing cascade reactions from
871 natural esters (using for instance well-known lipases; Johannsen et al., 2020) or carboxylic acid

872 as starting raw material. In any case, the extraction of the substrate from raw materials must be
873 done in compliance with local regulation to preserve their natural origin.

874 The hunt for *ad-hoc* biocatalysts to produce fatty aldehydes calls for highly integrated
875 approaches harnessing fundamental knowledge on enzymes and their biological context, taking
876 into account industrial constraints. We hope that the background information provided in this
877 review will encourage new investigations to foster the sustainable development of
878 biotechnological solutions for the F&F industry.

879

880 **Funding**

881 This study was supported by the French National Agency for Research (“Agence
882 Nationale de la Recherche”) through the “Projet de Recherche Collaboratif International” ANR-
883 NSERC (FUNTASTIC project, ANR-17-CE07-0047). PhD fellowship of D.R. was funded by
884 V. MANE Fils and the “Association Nationale Recherche Technologie” (ANRT) through the
885 *Convention Industrielle de Formation par la Recherche* (CIFRE) grant no. 2017/1169.

886

887

- 889
890 Al-Karadaghi, S., Cedergren-Zeppezauer, E.S., Hovmöller, S., Petratos, K., Terry, H., Wilson,
891 K.S., 1994. Refined crystal structure of liver alcohol dehydrogenase–NADH complex
892 at 1.8 Å resolution. *Acta Cryst.* D50, 793–807.
893 <https://doi.org/10.1107/S0907444994005263>
- 894 Alvarado-Caudillo, Y., Torres, J.C.B., Novoa, V.Z., Jiménez, H.S., Torres-Guzmán, J.C.,
895 Gutiérrez-Corona, J.F., Zazueta-Sandoval, R., 2002. Presence and physiologic
896 regulation of alcohol oxidase activity in an indigenous fungus isolated from petroleum-
897 contaminated soils. *Appl Biochem Biotechnol* 98, 243–255.
898 <https://doi.org/10.1385/ABAB:98-100:1-9:243>
- 899 Álvarez, L., Acevedo, F., Illanes, A., 2011. Induction of NAD⁺ dependent alcohol
900 dehydrogenases with activity towards long chain aliphatic alcohols in mesophilic,
901 thermophilic and extreme thermophilic microorganisms. *Process Biochem.* 46, 1342–
902 1349. <https://doi.org/10.1016/j.procbio.2011.03.002>
- 903 An, J., Nie, Y., Xu, Y., 2019. Structural insights into alcohol dehydrogenases catalyzing
904 asymmetric reductions. *Crit. Rev. Biotechnol.* 39, 366–379.
905 <https://doi.org/10.1080/07388551.2019.1566205>
- 906 Anasontzis, G.E., Salazar Penã, M., Spadiut, O., Brumer, H., Olsson, L., 2014. Effects of
907 temperature and glycerol and methanol-feeding profiles on the production of
908 recombinant galactose oxidase in *Pichia pastoris*. *Biotechnol Progress* 30, 728–735.
909 <https://doi.org/10.1002/btpr.1878>
- 910 Andberg, M., Mollerup, F., Parikka, K., Koutaniemi, S., Boer, H., Juvonen, M., Master, E.,
911 Tenkanen, M., Kruus, K., 2017. A Novel *Colletotrichum graminicola* Raffinose
912 Oxidase in the AA5 Family. *Appl. Environ. Microbiol.* 83.
913 <https://doi.org/10.1128/AEM.01383-17>
- 914 Antoine, E., Rolland, J.-L., Raffin, J.-P., Dietrich, J., 1999. Cloning and over-expression in
915 *Escherichia coli* of the gene encoding NADPH group III alcohol dehydrogenase from
916 *Thermococcus hydrothermalis*. *Eur. J. Biochem.* 264, 880–889.
917 <https://doi.org/10.1046/j.1432-1327.1999.00685.x>
- 918 Arnold, F.H., 2019. Innovation by Evolution: Bringing New Chemistry to Life (Nobel Lecture).
919 *Angew. Chem., Int. Ed.* 58, 14420–14426. <https://doi.org/10.1002/anie.201907729>
- 920 Balasubramanian, N., 2015. Scented Oils and Perfumes, in: *Chemical Technology in Antiquity*,
921 ACS Symposium Series. American Chemical Society, pp. 219–244.
922 <https://doi.org/10.1021/bk-2015-1211.ch008>
- 923 Baldwin, E., Plotto, A., Manthey, J., McCollum, G., Bai, J., Irey, M., Cameron, R., Luzio, G.,
924 2010. Effect of Liberibacter Infection (Huanglongbing Disease) of Citrus on Orange
925 Fruit Physiology and Fruit/Fruit Juice Quality: Chemical and Physical Analyses. *J.*
926 *Agric. Food Chem.* 58, 1247–1262. <https://doi.org/10.1021/jf9031958>
- 927 Banthorpe, D.V., Cardemil, E., Del Carmen Contreras, M., 1976. Purification and properties of
928 alcohol oxidase from *Tanacetum vulgare*. *Phytochemistry* 15, 391–394.
929 [https://doi.org/10.1016/S0031-9422\(00\)86829-6](https://doi.org/10.1016/S0031-9422(00)86829-6)
- 930 Baron, A.J., Stevens, C., Wilmot, C., Seneviratne, K.D., Blakeley, V., Dooley, D.M., Phillips,
931 S.E., Knowles, P.F., McPherson, M.J., 1994. Structure and mechanism of galactose
932 oxidase. The free radical site. *J. Biol. Chem.* 269, 25095–25105.
- 933 Barth, G., Künkkel, W., 2007. Alcohol dehydrogenase (ADH) in yeasts II. NAD⁺- and NADP⁺-
934 dependent alcohol dehydrogenases in *Saccharomycopsis lipolytica*. *Z Allg Mikrobiol*
935 19, 381–390. <https://doi.org/10.1002/jobm.19790190603>

- 936 Bartsch, S., Brummund, J., Köpke, S., Straatman, H., Vogel, A., Schürmann, M., 2020.
937 Optimization of Alcohol Dehydrogenase for Industrial Scale Oxidation of Lactols.
938 Biotechnol. J. 15, 2000171. <https://doi.org/10.1002/biot.202000171>
- 939 Basso, A., Serban, S., 2019. Industrial applications of immobilized enzymes—A review. Mol.
940 Catal. 479, 110607. <https://doi.org/10.1016/j.mcat.2019.110607>
- 941 Bell, R.P., 1966. The Reversible Hydration of Carbonyl Compounds, in: Gold, V. (Ed.),
942 Advances in Physical Organic Chemistry. Academic Press, pp. 1–29.
943 [https://doi.org/10.1016/S0065-3160\(08\)60351-2](https://doi.org/10.1016/S0065-3160(08)60351-2)
- 944 Ben Akacha, N., Gargouri, M., 2015. Microbial and enzymatic technologies used for the
945 production of natural aroma compounds: Synthesis, recovery modeling, and
946 bioprocesses. Food Bioprod. Process. 94, 675–706.
947 <https://doi.org/10.1016/j.fbp.2014.09.011>
- 948 Berger, R.G., 2015. Biotechnology as a source of natural volatile flavours. Curr. Opin. Food
949 Sci., 1, 38–43. <https://doi.org/10.1016/j.cofs.2014.09.003>
- 950 Berger, R.G., 2009. Biotechnology of flavours—the next generation. Biotechnol Lett 31, 1651.
951 <https://doi.org/10.1007/s10529-009-0083-5>
- 952 Berger, R.G. (Ed.), 2007. Flavours and Fragrances: Chemistry, Bioprocessing and
953 Sustainability. Springer-Verlag, Berlin Heidelberg.
- 954 Birmingham, W.R., Turner, N.J., 2018. A Single Enzyme Oxidative “Cascade” via a Dual-
955 Functional Galactose Oxidase. ACS Catal. 8, 4025–4032.
956 <https://doi.org/10.1021/acscatal.8b00043>
- 957 Bissaro, B., Kodama, S., Hage, H., Ribeaucourt, D., Haon, M., Grisel, S., Simaan, J., Beisson,
958 F., Forget, S., Brumer, H., Rosso, M.-N., O’Connell, R., Lafond, M., Kubo, Y., Berrin,
959 J.-G., 2021. Unravelling the role of alcohol copper radical oxidases in fungal plant
960 pathogens. <https://doi.org/10.21203/rs.3.rs-493001/v1>
- 961 Blasig, R., Mauersberger, S., Riege, P., Schunck, W.-H., Jockisch, W., Franke, P., Müller, H.-
962 G., 1988. Degradation of long-chain n-alkanes by the yeast *Candida maltosa*. Appl
963 Microbiol Biotechnol 28, 589–597. <https://doi.org/10.1007/BF00250418>
- 964 Bolivar, J.M., Wilson, L., Ferrarotti, S.A., Guisán, J.M., Fernández-Lafuente, R., Mateo, C.,
965 2006. Improvement of the stability of alcohol dehydrogenase by covalent
966 immobilization on glyoxyl-agarose. J. Biotechnol. 125, 85–94.
967 <https://doi.org/10.1016/j.jbiotec.2006.01.028>
- 968 Braga, A., Guerreiro, C., Belo, I., 2018. Generation of Flavors and Fragrances Through
969 Biotransformation and De Novo Synthesis. Food Bioprocess Technol. 11.
970 <https://doi.org/10.1007/s11947-018-2180-8>
- 971 Brenna, E., Parmeggiani, F., 2017. Biotechnological Production of Flavors, in: Industrial
972 Biotechnology. John Wiley & Sons, Ltd, pp. 271–308.
973 <https://doi.org/10.1002/9783527807833.ch9>
- 974 Buschmann, H.-J., Dutkiewicz, E., Knoche, W., 1982. The Reversible Hydration of Carbonyl
975 Compounds in Aqueous Solution Part II: The Kinetics of the Keto/*Gem*-diol Transition.
976 Ber Bunsen Phys Chem 86, 129–134. <https://doi.org/10.1002/bbpc.19820860208>
- 977 Buschmann, H.-J., Földner, H.-H., Knoche, W., 1980. The Reversible Hydration of Carbonyl
978 Compounds in Aqueous Solution. Part I, The Keto/*Gem*-diol Equilibrium. Ber Bunsen
979 Phys Chem 84, 41–44. <https://doi.org/10.1002/bbpc.19800840109>
- 980 Cao, Y., Li, X., Ge, J., 2021. Enzyme Catalyst Engineering toward the Integration of
981 Biocatalysis and Chemocatalysis. Trends Biotechnol.
982 <https://doi.org/10.1016/j.tibtech.2021.01.002>
- 983 Cea, G., Wilson, L., Bolívar, J.M., Markovits, A., Illanes, A., 2009. Effect of chain length on
984 the activity of free and immobilized alcohol dehydrogenase towards aliphatic alcohols.

985 Enzyme Microb. Technol. 44, 135–138.
986 <https://doi.org/10.1016/j.enzmictec.2008.10.021>
987 Chaplin, A.K., Petrus, M.L.C., Mangiameli, G., Hough, M.A., Svistunenko, D.A., Nicholls, P.,
988 Claessen, D., Vijgenboom, E., Worrall, J.A.R., 2015. GlxA is a new structural member
989 of the radical copper oxidase family and is required for glycan deposition at hyphal tips
990 and morphogenesis of *Streptomyces lividans*. *Biochem. J.* 469, 433–444.
991 <https://doi.org/10.1042/BJ20150190>
992 Chaplin, A.K., Svistunenko, D.A., Hough, M.A., Wilson, M.T., Vijgenboom, E., Worrall,
993 J.A.R., 2017. Active-site maturation and activity of the copper-radical oxidase GlxA are
994 governed by a tryptophan residue. *Biochem. J.* 474, 809–825.
995 <https://doi.org/10.1042/BCJ20160968>
996 Check Hayden, E., 2014. Synthetic-biology firms shift focus. *Nature News* 505, 598.
997 <https://doi.org/10.1038/505598a>
998 Chen, K., Arnold, F.H., 1993. Tuning the activity of an enzyme for unusual environments:
999 sequential random mutagenesis of subtilisin E for catalysis in dimethylformamide.
1000 *PNAS* 90, 5618–5622
1001 Cheng, Q., Liu, H.-T., Bombelli, P., Smith, A., Slabas, A.R., 2004. Functional identification of
1002 AtFao3, a membrane bound long chain alcohol oxidase in *Arabidopsis thaliana*. *FEBS*
1003 *Lett.* 574, 62–68. <https://doi.org/10.1016/j.febslet.2004.07.086>
1004 Cheng, Q., Sanglard, D., Vanhanen, S., Liu, H.T., Bombelli, P., Smith, A., Slabas, A.R., 2005.
1005 *Candida* yeast long chain fatty alcohol oxidase is a c-type haemoprotein and plays an
1006 important role in long chain fatty acid metabolism. *Biochim. Biophys. Acta - Molecular*
1007 *and Cell Biology of Lipids* 1735, 192–203. <https://doi.org/10.1016/j.bbalip.2005.06.006>
1008 Chiozza, L., 1856. Sur la production artificielle de l'essence de cannelle. *Comptes rendus*
1009 *hebdomadaires des séances de l'Académie des science* 1, 222.
1010 Clark, G.S., 1995. Benzaldehyde. *Perfumer & Flavorist*.
1011 Cobb, R.E., Chao, R., Zhao, H., 2013. Directed Evolution: Past, Present and Future. *AIChE J*
1012 59, 1432–1440. <https://doi.org/10.1002/aic.13995>
1013 Contente, M., Fiore, N., Cannazza, P., Roura Padrosa, D., Molinari, F., Gourlay, L., Paradisi,
1014 F., 2020. Uncommon overoxidative catalytic activity in a new halo-tolerant alcohol
1015 dehydrogenase. *ChemCatChem* 12, 5679–5685.
1016 <https://doi.org/10.1002/cctc.202001112>
1017 Corberán, V.C., González-Pérez, M.E., Martínez-González, S., Gómez-Avilés, A., 2014. Green
1018 oxidation of fatty alcohols: Challenges and opportunities. *Appl. Catal., A, B. Delmon*
1019 *Honor Special Issue* 474, 211–223. <https://doi.org/10.1016/j.apcata.2013.09.040>
1020 Crichton, R.R., 2012. Chapter 12 - Zinc – Lewis Acid and Gene Regulator, in: Crichton, R.R.
1021 (Ed.), *Biological Inorganic Chemistry (Second Edition)*. Elsevier, Oxford, pp. 229–246.
1022 <https://doi.org/10.1016/B978-0-444-53782-9.00012-7>
1023 Dala-Paula, B.M., Plotto, A., Bai, J., Manthey, J.A., Baldwin, E.A., Ferrarezi, R.S., Gloria,
1024 M.B.A., 2019. Effect of Huanglongbing or Greening Disease on Orange Juice Quality,
1025 a Review. *Front. Plant Sci.* 9. <https://doi.org/10.3389/fpls.2018.01976>
1026 Daou, M., Faulds, C.B., 2017. Glyoxal oxidases: their nature and properties. *World J Microbiol*
1027 *Biotechnol* 33, 87. <https://doi.org/10.1007/s11274-017-2254-1>
1028 D'Auria, S., 2002. Alcohol Dehydrogenase, in: *Handbook of Food Enzymology*. CRC Press,
1029 pp. 458–467. <https://doi.org/10.1201/9780203910450-34>
1030 Davis, S.C., Fox, R.J., Gavrilovic, V., Huisman, G.W., Newman, L.M., 2005. Improved
1031 halohydrin dehalogenases and related polynucleotides. *WO2005017141A1*.
1032 de Carvalho, C.C.C.R., da Fonseca, M.M.R., 2006. Carvone: Why and how should one bother
1033 to produce this terpene. *Food Chem.* 95, 413–422.
1034 <https://doi.org/10.1016/j.foodchem.2005.01.003>

- 1035 de Smidt, O., du Preez, J.C., Albertyn, J., 2008. The alcohol dehydrogenases of *Saccharomyces*
1036 *cerevisiae*: a comprehensive review. FEMS Yeast Res. 8, 967–978.
1037 <https://doi.org/10.1111/j.1567-1364.2008.00387.x>
- 1038 Deacon, S.E., Mahmoud, K., Spooner, R.K., Firbank, S.J., Knowles, P.F., Phillips, S.E.V.,
1039 McPherson, M.J., 2004. Enhanced Fructose Oxidase Activity in a Galactose Oxidase
1040 Variant. ChemBioChem 5, 972–979. <https://doi.org/10.1002/cbic.200300810>
- 1041 de Almeida, T.P., van Schie, M.M.C.H., Ma, A., Tieves, F., Younes, S.H.H., Fernández-Fueyo,
1042 E., Arends, I.W.C.E., Riul, A., Hollmann, F., 2019. Efficient Aerobic Oxidation of *trans*
1043 *-2-Hexen-1-ol* using the Aryl Alcohol Oxidase from *Pleurotus eryngii*. Adv. Synth.
1044 Catal. 361, 2668–2672. <https://doi.org/10.1002/adsc.201801312>
- 1045 Derrington, S.R., Turner, N.J., France, S.P., 2019. Carboxylic acid reductases (CARs): An
1046 industrial perspective. J. Biotechnol. 304, 78–88.
1047 <https://doi.org/10.1016/j.jbiotec.2019.08.010>
- 1048 Diaz, S., Espinosa, S., Brignole, E.A., 2005. Citrus peel oil deterpenation with supercritical
1049 fluids: Optimal process and solvent cycle design. J. Supercrit. Fluids 35, 49–61.
1050 <https://doi.org/10.1016/j.supflu.2004.12.002>
- 1051 Dickinson, F.M., Dack, S., 2001. The activity of yeast ADH I and ADH II with long-chain
1052 alcohols and diols. Chem.-Biol. Interact. 130–132, 417–423.
1053 [https://doi.org/10.1016/S0009-2797\(00\)00266-0](https://doi.org/10.1016/S0009-2797(00)00266-0)
- 1054 Dickinson, F.M., Wadforth, C., 1992. Purification and some properties of alcohol oxidase from
1055 alkane-grown *Candida tropicalis*. Biochem J 282, 325–331.
- 1056 Dijkman, W.P., de Gonzalo, G., Mattevi, A., Fraaije, M.W., 2013. Flavoprotein oxidases:
1057 classification and applications. Appl Microbiol Biotechnol 97, 5177–5188.
1058 <https://doi.org/10.1007/s00253-013-4925-7>
- 1059 Domínguez de María, P., Hollmann, F., 2015. On the (Un)greenness of Biocatalysis: Some
1060 Challenging Figures and Some Promising Options. Front. Microbiol. 6.
1061 <https://doi.org/10.3389/fmicb.2015.01257>
- 1062 Dong, J., Fernández-Fueyo, E., Hollmann, F., Paul, C.E., Pesic, M., Schmidt, S., Wang, Y.,
1063 Younes, S., Zhang, W., 2018. Biocatalytic Oxidation Reactions: A Chemist’s
1064 Perspective. Angew Chem Int Ed Engl 57, 9238–9261.
1065 <https://doi.org/10.1002/anie.201800343>
- 1066 Doussin, J.-F., Monod, A., 2013. Structure–activity relationship for the estimation of OH-
1067 oxidation rate constants of carbonyl compounds in the aqueous phase. Atmos. Chem.
1068 Phys. 13, 11625–11641. <https://doi.org/10.5194/acp-13-11625-2013>
- 1069 Dunkel, A., Steinhaus, M., Kotthoff, M., Nowak, B., Krautwurst, D., Schieberle, P., Hofmann,
1070 T., 2014. Nature’s Chemical Signatures in Human Olfaction: A Foodborne Perspective
1071 for Future Biotechnology. Angew Chem Int Ed Engl 53, 7124–7143.
1072 <https://doi.org/10.1002/anie.201309508>
- 1073 Edenberg, H.J., 2007. The Genetics of Alcohol Metabolism: Role of Alcohol Dehydrogenase
1074 and Aldehyde Dehydrogenase Variants. Alcohol Res Health 30, 5–13.
- 1075 Eirich, L.D., Craft, D.L., 2007. Fatty alcohol oxidase genes and proteins from *Candida*
1076 *tropicalis* and methods relating thereto. US7160708B2.
- 1077 Eirich, L.D., Craft, D.L., Steinberg, L., Asif, A., Eschenfeldt, W.H., Stols, L., Donnelly, M.I.,
1078 Wilson, C.R., 2004. Cloning and characterization of three fatty alcohol oxidase genes
1079 from *Candida tropicalis* strain ATCC 20336. Appl. Environ. Microbiol. 70, 4872–4879.
1080 <https://doi.org/10.1128/AEM.70.8.4872-4879.2004>
- 1081 Escalettes, F., Turner, N.J., 2008. Directed Evolution of Galactose Oxidase: Generation of
1082 Enantioselective Secondary Alcohol Oxidases. ChemBioChem 9, 857–860.
1083 <https://doi.org/10.1002/cbic.200700689>

- 1084 Ewing, T.A., Fraaije, M., van Berkel, W.J.H., 2015. Oxidation using alcohol oxidases, in:
1085 Faber, K., Fessner, W.-D., Turner, N.J. (Eds.), Science of Synthesis: Biocatalysis in
1086 Organic Synthesis. Thieme, pp. 157–185.
- 1087 Ewing, T.A., Fraaije, M.W., Mattevi, A., van Berkel, W.J.H., 2017. The VAO/PCMH
1088 flavoprotein family. Arch. Biochem. Biophys., Flavoproteins: Beyond the Classical
1089 Paradigms 632, 104–117. <https://doi.org/10.1016/j.abb.2017.06.022>
- 1090 Ewing, T.A., Gygli, G., Fraaije, M.W., van Berkel, W.J.H., 2020. Chapter Four - Vanillyl
1091 alcohol oxidase, in: Chaiyen, P., Tamanoi, F. (Eds.), The Enzymes, Flavin-Dependent
1092 Enzymes: Mechanisms, Structures and Applications. Academic Press, pp. 87–116.
1093 <https://doi.org/10.1016/bs.enz.2020.05.003>
- 1094 Ewing, T.A., Van Noord, A., Paul, C.E., Van Berkel, W.J.H., 2018. A Xylenol Orange-Based
1095 Screening Assay for the Substrate Specificity of Flavin-Dependent *para*-Phenol
1096 Oxidases. Molecules 23, 164. <https://doi.org/10.3390/molecules23010164>
- 1097 Fasim, A., More, V.S., More, S.S., 2021. Large-scale production of enzymes for biotechnology
1098 uses. Curr. Opin. Biotechnol. 69, 68–76. <https://doi.org/10.1016/j.copbio.2020.12.002>
- 1099 Ferreira, P., Hernández-Ortega, A., Herguedas, B., Rencoret, J., Gutiérrez, A., Martínez, M.J.,
1100 Jiménez-Barbero, J., Medina, M., Martínez, Á.T., 2010. Kinetic and chemical
1101 characterization of aldehyde oxidation by fungal aryl-alcohol oxidase. Biochem J 425,
1102 585–593. <https://doi.org/10.1042/BJ20091499>
- 1103 Firbank, S.J., Rogers, M.S., Wilmot, C.M., Dooley, D.M., Halcrow, M.A., Knowles, P.F.,
1104 McPherson, M.J., Phillips, S.E., 2001. Crystal structure of the precursor of galactose
1105 oxidase: an unusual self-processing enzyme. PNAS. 98, 12932–12937.
1106 <https://doi.org/10.1073/pnas.231463798>
- 1107 Forget, S., Xia, R. (Fan), Hein, J.E., Brumer, H., 2020. Determination of biocatalytic parameters
1108 of a copper radical oxidase using real-time reaction progress monitoring. Org. Biomol.
1109 Chem. <https://doi.org/10.1039/C9OB02757B>
- 1110 Fox, M.G.A., Dickinson, F.M., Ratledge, C., 1992. Long-chain alcohol and aldehyde
1111 dehydrogenase activities in *Acinetobacter calcoaceticus* strain HO1-N. Microbiology,
1112 138, 1963–1972. <https://doi.org/10.1099/00221287-138-9-1963>
- 1113 Frey, C., 2005. Natural Flavors and Fragrances: Chemistry, Analysis, and Production, in:
1114 Natural Flavors and Fragrances, ACS Symposium Series. American Chemical Society,
1115 pp. 3–19. <https://doi.org/10.1021/bk-2005-0908.ch001>
- 1116 Fukuda, R., Ohta, A., 2017. Enzymes for Aerobic Degradation of Alkanes in Yeasts, in: Rojo,
1117 F. (Ed.), Aerobic Utilization of Hydrocarbons, Oils and Lipids, Handbook of
1118 Hydrocarbon and Lipid Microbiology. Springer International Publishing, Cham, pp. 1–
1119 14. https://doi.org/10.1007/978-3-319-39782-5_7-1
- 1120 Gadda, G., 2008. Hydride Transfer Made Easy in the Reaction of Alcohol Oxidation Catalyzed
1121 by Flavin-dependent Oxidases. Biochemistry 47, 13745–13753.
1122 <https://doi.org/10.1021/bi801994c>
- 1123 Gadda, G., 2003. Kinetic mechanism of choline oxidase from *Arthrobacter globiformis*.
1124 Biochim. Biophys. Acta, Proteins Proteomics 1646, 112–118.
1125 [https://doi.org/10.1016/S1570-9639\(03\)00003-7](https://doi.org/10.1016/S1570-9639(03)00003-7)
- 1126 Gallage, N.J., Møller, B.L., 2015. Vanillin–Bioconversion and Bioengineering of the Most
1127 Popular Plant Flavor and Its *De Novo* Biosynthesis in the Vanilla Orchid. Mol. Plant, 8,
1128 40–57. <https://doi.org/10.1016/j.molp.2014.11.008>
- 1129 Gandolfi, R., Ferrara, N., Molinari, F., 2001. An easy and efficient method for the production
1130 of carboxylic acids and aldehydes by microbial oxidation of primary alcohols.
1131 Tetrahedron Lett. 42, 513–514. [https://doi.org/10.1016/S0040-4039\(00\)02008-6](https://doi.org/10.1016/S0040-4039(00)02008-6)

- 1132 Garzón-Posse, F., Becerra-Figueroa, L., Hernández-Arias, J., Gamba-Sánchez, D., 2018.
1133 Whole Cells as Biocatalysts in Organic Transformations. *Molecules* 23.
1134 <https://doi.org/10.3390/molecules23061265>
- 1135 Gounaris, Y., 2010. Biotechnology for the production of essential oils, flavours and volatile
1136 isolates. A review. *Flavour Fragr J.* 25, 367–386. <https://doi.org/10.1002/ffj.1996>
- 1137 Green, D.W., Sun, H.W., Plapp, B.V., 1993. Inversion of the substrate specificity of yeast
1138 alcohol dehydrogenase. *J. Biol. Chem.* 268, 7792–7798.
- 1139 Hamilton, G.A., Adolf, P.K., De Jersey, J., DuBois, G.C., Dyrkacz, G.R., Libby, R.D., 1978.
1140 Trivalent copper, superoxide, and galactose oxidase. *J. Am. Chem. Soc.* 100, 1899–
1141 1912. <https://doi.org/10.1021/ja00474a042>
- 1142 Haon, M., Grisel, S., Navarro, D., Gruet, A., Berrin, J.-G., Bignon, C., 2015. Recombinant
1143 protein production facility for fungal biomass-degrading enzymes using the yeast *Pichia*
1144 *pastoris*. *Front. Microbiol.* 6. <https://doi.org/10.3389/fmicb.2015.01002>
- 1145 Heath, R.S., Birmingham, W.R., Thompson, M.P., Taglieber, A., Daviet, L., Turner, N.J., 2019.
1146 An Engineered Alcohol Oxidase for the Oxidation of Primary Alcohols. *ChemBioChem*
1147 20, 276–281. <https://doi.org/10.1002/cbic.201800556>
- 1148 Hebditch, M., Carballo-Amador, M.A., Charonis, S., Curtis, R., Warwicker, J., 2017. Protein–
1149 Sol: a web tool for predicting protein solubility from sequence. *Bioinformatics* 33,
1150 3098–3100. <https://doi.org/10.1093/bioinformatics/btx345>
- 1151 Hernández-Tobías, A., Julián-Sánchez, A., Piña, E., Riveros-Rosas, H., 2011. Natural alcohol
1152 exposure: Is ethanol the main substrate for alcohol dehydrogenases in animals? *Chem.-*
1153 *Biol. Interact., Enzymology and Molecular Biology of Carbonyl Metabolism* 191, 14–
1154 25. <https://doi.org/10.1016/j.cbi.2011.02.008>
- 1155 Herter, S., McKenna, S.M., Frazer, A.R., Leimkühler, S., Carnell, A.J., Turner, N.J., 2015.
1156 Galactose Oxidase Variants for the Oxidation of Amino Alcohols in Enzyme Cascade
1157 Synthesis. *ChemCatChem* 7, 2313–2317. <https://doi.org/10.1002/cctc.201500218>
- 1158 Hilal, S.H., Bornander, L.L., Carreira, L.A., 2005. Hydration Equilibrium Constants of
1159 Aldehydes, Ketones and Quinazolines. *QSAR Comb. Sci.* 24, 631–638.
1160 <https://doi.org/10.1002/qsar.200430913>
- 1161 Hinson, J.A., Neal, R.A., 1975. An examination of octanol and octanal metabolism to octanoic
1162 acid by horse liver alcohol dehydrogenase. *Biochim. Biophys. Acta, Enzymol.* 384, 1–
1163 11. [https://doi.org/10.1016/0005-2744\(75\)90090-X](https://doi.org/10.1016/0005-2744(75)90090-X)
- 1164 Hinson, J.A., Neal, R.A., 1972. An Examination of the Oxidation of Aldehydes by Horse Liver
1165 Alcohol Dehydrogenase. *J. Biol. Chem.* 247, 7106–7107.
- 1166 Hirakawa, H., Kamiya, N., Kawarabayashi, Y., Nagamune, T., 2004. Properties of an alcohol
1167 dehydrogenase from the hyperthermophilic archaeon *Aeropyrum pernix* K1. *J. Biosci.*
1168 *Bioeng.* 97, 202–206. [https://doi.org/10.1016/S1389-1723\(04\)70191-7](https://doi.org/10.1016/S1389-1723(04)70191-7)
- 1169 Hirano, J., Miyamoto, K., Ohta, H., 2005. Purification and characterization of the alcohol
1170 dehydrogenase with a broad substrate specificity originated from 2-phenylethanol-
1171 assimilating *Brevibacterium* sp. KU 1309. *J. Biosci. Bioeng.* 100, 318–322.
1172 <https://doi.org/10.1263/jbb.100.318>
- 1173 Hollmann, F., Arends, I.W.C.E., Buehler, K., Schallmeyer, A., Bühler, B., 2011. Enzyme-
1174 mediated oxidations for the chemist. *Green Chem.* 13, 226–265.
1175 <https://doi.org/10.1039/C0GC00595A>
- 1176 Hollmann, F., Bühler, K., Bühler, B., 2012. Oxidation of Alcohols, Aldehydes, and Acids, in:
1177 *Enzyme Catalysis in Organic Synthesis*. John Wiley & Sons, Ltd, pp. 1325–1437.
1178 <https://doi.org/10.1002/9783527639861.ch32>
- 1179 Höllrigl, V., Hollmann, F., Kleeb, A.C., Buehler, K., Schmid, A., 2008. TADH, the
1180 thermostable alcohol dehydrogenase from *Thermus* sp. ATN1: a versatile new

1181 biocatalyst for organic synthesis. *Appl Microbiol Biotechnol* 81, 263–273.
1182 <https://doi.org/10.1007/s00253-008-1606-z>

1183 Holt, P.J., Williams, R.E., Jordan, K.N., Lowe, C.R., Bruce, N.C., 2000. Cloning, sequencing
1184 and expression in *Escherichia coli* of the primary alcohol dehydrogenase gene from
1185 *Thermoanaerobacter ethanolicus* JW200. *FEMS Microbiol Lett* 190, 57–62.
1186 <https://doi.org/10.1111/j.1574-6968.2000.tb09262.x>

1187 Hommel, R., Ratledge, C., 1990. Evidence for two fatty alcohol oxidases in the biosurfactant-
1188 producing yeast *Candida (Torulopsis) bombicola*. *FEMS Microbiol Lett* 70, 183–186.
1189 <https://doi.org/10.1111/j.1574-6968.1990.tb13975.x>

1190 Hommel, R.K., Lassner, D., Weiss, J., Kleber, H.-P., 1994. The inducible microsomal fatty
1191 alcohol oxidase of *Candida (Torulopsis) apicola*. *Appl Microbiol Biotechnol* 40, 729–
1192 734. <https://doi.org/10.1007/BF00173336>

1193 Horvat, M., Larch, T.-S., Rudroff, F., Winkler, M., 2020. Amino Benzamidoxime (ABAO)-
1194 Based Assay to Identify Efficient Aldehyde-Producing *Pichia pastoris* Clones. *Adv.*
1195 *Synth. Catal.* 362, 4673–4679. <https://doi.org/10.1002/adsc.202000558>

1196 Hoyos, P., Hernáiz, M.J., Alcántara, A.R., 2017. 3.28 - Biocatalyzed Production of Fine
1197 Chemicals, in: Moo-Young, M. (Ed.), *Comprehensive Biotechnology (Third Edition)*.
1198 Pergamon, Oxford, pp. 334–373. [https://doi.org/10.1016/B978-0-12-809633-8.09153-](https://doi.org/10.1016/B978-0-12-809633-8.09153-6)
1199 6

1200 Huang, L., Sayoga, G.V., Hollmann, F., Kara, S., 2018. Horse Liver Alcohol Dehydrogenase-
1201 Catalyzed Oxidative Lactamization of Amino Alcohols. *ACS Catal.* 8, 8680–8684.
1202 <https://doi.org/10.1021/acscatal.8b02355>

1203 Huffman, M.A., Fryszkowska, A., Alvizo, O., Borra-Garske, M., Campos, K.R., Canada, K.A.,
1204 Devine, P.N., Duan, D., Forstater, J.H., Grosser, S.T., Halsey, H.M., Hughes, G.J., Jo,
1205 J., Joyce, L.A., Kolev, J.N., Liang, J., Maloney, K.M., Mann, B.F., Marshall, N.M.,
1206 McLaughlin, M., Moore, J.C., Murphy, G.S., Nawrat, C.C., Nazor, J., Novick, S., Patel,
1207 N.R., Rodriguez-Granillo, A., Robaire, S.A., Sherer, E.C., Truppo, M.D., Whittaker,
1208 A.M., Verma, D., Xiao, L., Xu, Y., Yang, H., 2019. Design of an *in vitro* biocatalytic
1209 cascade for the manufacture of islatravir. *Science* 366, 1255–1259.
1210 <https://doi.org/10.1126/science.aay8484>

1211 Hummel, W., Abokitse, K., Drauz, K., Rollmann, C., Gröger, H., 2003. Towards a Large-Scale
1212 Asymmetric Reduction Process with Isolated Enzymes: Expression of an (*S*)-Alcohol
1213 Dehydrogenase in *E. coli* and Studies on the Synthetic Potential of this Biocatalyst. *Adv.*
1214 *Synth. Catal.* 345, 153–159. <https://doi.org/10.1002/adsc.200390001>

1215 Hummel, W., Hummel, W., 1999. Large-scale applications of NAD(P)-dependent
1216 oxidoreductases: recent developments. *Trends Biotechnol.* 17, 487–492.
1217 [https://doi.org/10.1016/S0167-7799\(98\)01207-4](https://doi.org/10.1016/S0167-7799(98)01207-4)

1218 Hutchings, M.G., Gasteiger, J., 1986. Correlation analyses of the aqueous-phase acidities of
1219 alcohols and *gem*-diols, and of carbonyl hydration equilibria using electronic and
1220 structural parameters. *J. Chem. Soc., Perkin Trans.* 2 455–462.
1221 <https://doi.org/10.1039/P29860000455>

1222 IAL Consultants, 2020. *An Overview of the Global Flavours & Fragrances Market*, 12th
1223 Edition.

1224 Il'chenko, A., 1984. Oxidase of higher alcohols in the yeast *Torulopsis candida* grown on
1225 hexadecane. *Mikrobiologiya* 53, 903–906.

1226 Il'chenko, A., Tsfasman, I., 1988. Isolation and characterization of alcohol oxidase from higher
1227 alcohols of the yeast *Torulopsis candida* grown on hexadecane. *Biokhimiya* 53, 263–
1228 271.

1229 Inoue, K., Makino, Y., Itoh, N., 2005. Purification and Characterization of a Novel Alcohol
1230 Dehydrogenase from *Leifsonia* sp. Strain S749: a Promising Biocatalyst for an

1231 Asymmetric Hydrogen Transfer Bioreduction. *Appl Environ Microbiol* 71, 3633–3641.
1232 <https://doi.org/10.1128/AEM.71.7.3633-3641.2005>

1233 Ito, N., Phillips, S.E., Stevens, C., Ogel, Z.B., McPherson, M.J., Keen, J.N., Yadav, K.D.,
1234 Knowles, P.F., 1991. Novel thioether bond revealed by a 1.7 Å crystal structure of
1235 galactose oxidase. *Nature* 350, 87–90. <https://doi.org/10.1038/350087a0>

1236 Johannsen, J., Meyer, F., Fieg, G., Waluga, T., 2020. Process for the Preparation of Fatty
1237 Aldehydes from Renewable Resources. *Chem Ing Tech* 92, 777–781.
1238 <https://doi.org/10.1002/cite.201900111>

1239 Johnson, H.C., Zhang, S., Fryszkowska, A., Ruccolo, S., Robaire, S.A., Klapars, A., Patel, N.R.,
1240 Whittaker, A.M., Huffman, M.A., Strotman, N.A., 2021. Biocatalytic oxidation of
1241 alcohols using galactose oxidase and a manganese(III) activator for the synthesis of
1242 islatravir. *Org. Biomol. Chem.* <https://doi.org/10.1039/D0OB02395G>

1243 Kakoti, A., Kumar, A.K., Goswami, P., 2012. Microsome-bound alcohol oxidase catalyzed
1244 production of carbonyl compounds from alcohol substrates. *J. Mol. Catal. B: Enzym.*
1245 78, 98–104. <https://doi.org/10.1016/j.molcatb.2012.03.013>

1246 Karra-Chaabouni, M., Pulvin, S., Meziani, A., Thomas, D., Touraud, D., Kunz, W., 2003.
1247 Biooxidation of *n*-hexanol by alcohol oxidase and catalase in biphasic and micellar
1248 systems without solvent. *Biotechnol. Bioeng.* 81, 27–32.
1249 <https://doi.org/10.1002/bit.10452>

1250 Kazuoka, T., Oikawa, T., Muraoka, I., Kuroda, S., Soda, K., 2007. A cold-active and
1251 thermostable alcohol dehydrogenase of a psychrotolerant from Antarctic seawater,
1252 *Flavobacterium frigidimaris* KUC-1. *Extremophiles* 11, 257–267.
1253 <https://doi.org/10.1007/s00792-006-0034-1>

1254 Kelleher, F.M., Bhavanandan, V.P., 1986. Re-examination of the products of the action of
1255 galactose oxidase. Evidence for the conversion of raffinose to 6''-carboxyraffinose. *J.*
1256 *Biol. Chem.* 261, 11045–11048.

1257 Kemp, G.D., Dickinson, F.M., Ratledge, C., 1991. Activity and substrate specificity of the fatty
1258 alcohol oxidase of *Candida tropicalis* in organic solvents. *Appl Microbiol Biotechnol*
1259 34, 441–445. <https://doi.org/10.1007/BF00180567>

1260 Kemp, G.D., Dickinson, F.M., Ratledge, C., 1990. Light sensitivity of the *n*-alkane-induced
1261 fatty alcohol oxidase from *Candida tropicalis* and *Yarrowia lipolytica*. *Appl Microbiol*
1262 *Biotechnol* 32, 461–464. <https://doi.org/10.1007/BF00903783>

1263 Kemp, G.D., Dickinson, F.M., Ratledge, C., 1988. Inducible long chain alcohol oxidase from
1264 alkane-grown *Candida tropicalis*. *Appl Microbiol Biotechnol* 29, 370–374.
1265 <https://doi.org/10.1007/BF00265821>

1266 Kersten, P., Cullen, D., 2014. Copper radical oxidases and related extracellular oxidoreductases
1267 of wood-decay Agaricomycetes. *Fungal Genetics and Biology* 72, 124–130.
1268 <https://doi.org/10.1016/j.fgb.2014.05.011>

1269 Kim, M., Jang, B., Ahn, Y.-J., 2019. Enhanced Production of Recombinant Alcohol
1270 Dehydrogenase Using the Genetically Engineered *Escherichia coli* Strain that
1271 Heterologously Expresses Carrot Heat Shock Protein 70. *Curr Microbiol* 76, 1338–
1272 1344. <https://doi.org/10.1007/s00284-019-01730-8>

1273 Koch, C., Neumann, P., Valerius, O., Feussner, I., Ficner, R., 2016. Crystal Structure of Alcohol
1274 Oxidase from *Pichia pastoris*. *PLOS ONE* 11, e0149846.
1275 <https://doi.org/10.1371/journal.pone.0149846>

1276 Kohlpaintner, C., Schulte, M., Falbe, J., Lappe, P., Weber, J., Frey, G.D., 2013. Aldehydes,
1277 Aliphatic, in: *Ullmann's Encyclopedia of Industrial Chemistry*. American Cancer
1278 Society. https://doi.org/10.1002/14356007.a01_321.pub3

1279 Könst, P., Merckens, H., Kara, S., Kochius, S., Vogel, A., Zuhse, R., Holtmann, D., Arends,
1280 I.W.C.E., Hollmann, F., 2012. Enantioselective Oxidation of Aldehydes Catalyzed by

1281 Alcohol Dehydrogenase. *Angew. Chem., Int. Ed.* 51, 9914–9917.
1282 <https://doi.org/10.1002/anie.201203219>

1283 Krauzova, V.I., Il'chenko, A.P., Sharyshev, A.A., Lozinov, A.B., 1985. Possible pathways of
1284 the oxidation of higher alcohols by membrane fractions of yeasts cultured on
1285 hexadecane and hexadecanol. *Biokhimiya* 50, 726–732.

1286 Krauzova, V.I., Kuvichkina, T.N., Sharyshev, A.A., Romanova, I.B., Lozinov, A.B., 1986.
1287 Formation of lauric acid and NADH in the oxidation of dodecanol by membrane
1288 fractions of the yeast *Candida maltosa* cultured on hexadecane. *Biokhimiya* 51, 23–27.

1289 Kumar, A.K., Goswami, P., 2009. Dissociation and Reconstitution Studies of a Broad Substrate
1290 Specific Multimeric Alcohol Oxidase Protein Produced by *Aspergillus terreus*. *J.*
1291 *Biochem.* 145, 259–265. <https://doi.org/10.1093/jb/mvn163>

1292 Kumar, A.K., Goswami, P., 2008. Purification and properties of a novel broad substrate specific
1293 alcohol oxidase from *Aspergillus terreus* MTCC 6324. *Biochim. Biophys. Acta,*
1294 *Proteins Proteomics* 1784, 1552–1559. <https://doi.org/10.1016/j.bbapap.2008.06.009>

1295 Kumar, A.K., Goswami, P., 2006. Functional characterization of alcohol oxidases from
1296 *Aspergillus terreus* MTCC 6324. *Appl Microbiol Biotechnol* 72, 906–911.
1297 <https://doi.org/10.1007/s00253-006-0381-y>

1298 Kunjapur, A.M., Prather, K.L.J., 2015. Microbial Engineering for Aldehyde Synthesis. *Appl*
1299 *Environ Microbiol* 81, 1892–1901. <https://doi.org/10.1128/AEM.03319-14>

1300 Lambert, F., Zucca, J., Mane, J., 2013. System for the production of aromatic molecules in
1301 *Streptomyces*. US8344119B2.

1302 Lambert, F., Zucca, J., Ness, F., Aigle, M., 2014. Production of ferulic acid and coniferyl
1303 alcohol by conversion of eugenol using a recombinant strain of *Saccharomyces*
1304 *cerevisiae*. *Flavour and Fragrance Journal* 29, 14–21. <https://doi.org/10.1002/ffj.3173>

1305 Leskovac, V., Trivić, S., Peričin, D., 2002. The three zinc-containing alcohol dehydrogenases
1306 from baker's yeast, *Saccharomyces cerevisiae*. *FEMS Yeast Res* 2, 481–494.
1307 <https://doi.org/10.1111/j.1567-1364.2002.tb00116.x>

1308 Levasseur, A., Drula, E., Lombard, V., Coutinho, P.M., Henrissat, B., 2013. Expansion of the
1309 enzymatic repertoire of the CAZy database to integrate auxiliary redox enzymes.
1310 *Biotechnol Biofuels* 6, 41. <https://doi.org/10.1186/1754-6834-6-41>

1311 Linke, D., Lehnert, N., Nimtz, M., Berger, R.G., 2014. An alcohol oxidase of *Phanerochaete*
1312 *chrysosporium* with a distinct glycerol oxidase activity. *Enzyme Microb. Technol.* 61–
1313 62, 7–12. <https://doi.org/10.1016/j.enzmictec.2014.04.001>

1314 Liu, J., Wu, S., Li, Z., 2018. Recent advances in enzymatic oxidation of alcohols. *Curr Opin*
1315 *Chem Biol* 43, 77–86. <https://doi.org/10.1016/j.cbpa.2017.12.001>

1316 Liu, X., Dong, Y., Zhang, J., Zhang, A., Wang, L., Feng, L., 2009. Two novel metal-
1317 independent long-chain alkyl alcohol dehydrogenases from *Geobacillus*
1318 *thermodenitrificans* NG80-2. *Microbiology*, 155, 2078–2085.
1319 <https://doi.org/10.1099/mic.0.027201-0>

1320 Lombard, V., Golaconda Ramulu, H., Drula, E., Coutinho, P.M., Henrissat, B., 2014. The
1321 carbohydrate-active enzymes database (CAZy) in 2013. *Nucleic Acids Res* 42, D490–
1322 D495. <https://doi.org/10.1093/nar/gkt1178>

1323 Lončar, N., Fraaije, M.W., 2015. Catalases as biocatalysts in technical applications: current
1324 state and perspectives. *Appl Microbiol Biotechnol* 99, 3351–3357.
1325 <https://doi.org/10.1007/s00253-015-6512-6>

1326 Longo, M., Sanromán, M., 2006. Production of Food Aroma Compounds: Microbial and
1327 Enzymatic Methodologies. *Food Technol. Biotechnol.* 44.

1328 Lutz, S., 2010. Beyond directed evolution—semi-rational protein engineering and design. *Curr.*
1329 *Opin. Biotechnol.* 21, 734–743. <https://doi.org/10.1016/j.copbio.2010.08.011>

- 1330 Ma, K., Adams, M.W.W., 1999. An Unusual Oxygen-Sensitive, Iron- and Zinc-Containing
1331 Alcohol Dehydrogenase from the Hyperthermophilic Archaeon *Pyrococcus furiosus*. *J*
1332 *Bacteriol* 181, 1163–1170.
- 1333 Ma, K., Robb, F.T., Adams, M.W., 1994. Purification and characterization of NADP-specific
1334 alcohol dehydrogenase and glutamate dehydrogenase from the hyperthermophilic
1335 archaeon *Thermococcus litoralis*. *Appl. Environ. Microbiol.* 60, 562–568.
- 1336 Malik, T., Rawat, S., 2021. Biotechnological Interventions for Production of Flavour and
1337 Fragrance Compounds, in: Venkatramanan, V., Shah, S., Prasad, R. (Eds.), *Sustainable*
1338 *Bioeconomy: Pathways to Sustainable Development Goals*. Springer, Singapore, pp.
1339 131–170. https://doi.org/10.1007/978-981-15-7321-7_7
- 1340 Marteau, C., Ruyffelaere, F., Aubry, J.-M., Penverne, C., Favier, D., Nardello-Rataj, V., 2013.
1341 Oxidative degradation of fragrant aldehydes. Autoxidation by molecular oxygen.
1342 *Tetrahedron* 69, 2268–2275. <https://doi.org/10.1016/j.tet.2013.01.034>
- 1343 Martin, C., Trajkovic, M., Fraaije, M., 2020. Production of Hydroxy Acids through Selective
1344 Double Oxidation of Diols by a Flavoprotein Alcohol Oxidase. *Angew. Chem., Int. Ed.*
1345 59, 4869–4872. <https://doi.org/10.1002/anie.201914877>
- 1346 Mathieu, Y., Offen, W.A., Forget, S.M., Ciano, L., Viborg, A.H., Blagova, E., Henrissat, B.,
1347 Walton, P.H., Davies, G.J., Brumer, H., 2020. Discovery of a fungal copper radical
1348 oxidase with high catalytic efficiency towards 5-hydroxymethylfurfural and benzyl
1349 alcohols for bioprocessing. *ACS Catal.* 10, 3042–3058.
1350 <https://doi.org/10.1021/acscatal.9b04727>
- 1351 Matsuda, T., Yamanaka, R., Nakamura, K., 2009. Recent progress in biocatalysis for
1352 asymmetric oxidation and reduction. *Tetrahedron: Asymmetry* 20, 513–557.
1353 <https://doi.org/10.1016/j.tetasy.2008.12.035>
- 1354 Matthey, A.P., Sangster, J.J., Ramsden, J.I., Baldwin, C., Birmingham, W.R., Heath, R.S.,
1355 Angelastro, A., Turner, N.J., Cosgrove, S.C., Flitsch, S.L., 2020. Natural heterogeneous
1356 catalysis with immobilised oxidase biocatalysts. *RSC Adv.* 10, 19501–19505.
1357 <https://doi.org/10.1039/D0RA03618H>
- 1358 Mauersberger, S., Drechsler, H., Oehme, G., Müller, H.-G., 1992. Substrate specificity and
1359 stereoselectivity of fatty alcohol oxidase from the yeast *Candida maltosa*. *Appl*
1360 *Microbiol Biotechnol* 37, 66–73. <https://doi.org/10.1007/BF00174205>
- 1361 Maurer, S., Schewe, H., Schrader, J., Buchhaupt, M., 2019. Investigation of fatty aldehyde and
1362 alcohol synthesis from fatty acids by α Dox- or CAR-expressing *Escherichia coli*. *J.*
1363 *Biotechnol.* 305, 11–17. <https://doi.org/10.1016/j.jbiotec.2019.08.011>
- 1364 McClelland, R.A., Coe, M., 1983. Structure-reactivity effects in the hydration of
1365 benzaldehydes. *J. Am. Chem. Soc.* 105, 2718–2725.
1366 <https://doi.org/10.1021/ja00347a033>
- 1367 McKenna, S.M., Leimkühler, S., Herter, S., Turner, N.J., Carnell, A.J., 2015. Enzyme cascade
1368 reactions: synthesis of furandicarboxylic acid (FDCA) and carboxylic acids using
1369 oxidases in tandem. *Green Chem.* 17, 3271–3275.
1370 <https://doi.org/10.1039/C5GC00707K>
- 1371 McPherson, M.J., Stevens, C., Baron, A.J., Ogel, Z.B., Seneviratne, K., Wilmot, C., Ito, N.,
1372 Brocklebank, I., Phillips, S.E., Knowles, P.F., 1993. Galactose oxidase: molecular
1373 analysis and mutagenesis studies. *Biochem. Soc. Trans.* 21 (Pt 3), 752–756.
- 1374 Mollerup, F., Aumala, V., Parikka, K., Mathieu, Y., Brumer, H., Tenkanen, M., Master, E.,
1375 2019. A family AA5_2 carbohydrate oxidase from *Penicillium rubens* displays
1376 functional overlap across the AA5 family. *PLOS ONE* 14, e0216546.
1377 <https://doi.org/10.1371/journal.pone.0216546>

- 1378 Moreau, R.A., Huang, A.H.C., 1979. Oxidation of fatty alcohol in the cotyledons of jojoba
1379 seedlings. Arch. Biochem. Biophys. 194, 422–430. <https://doi.org/10.1016/0003->
1380 9861(79)90636-2
- 1381 Murray, W.D., Duff, S.J.B., 1990. Bio-oxidation of aliphatic and aromatic high molecular
1382 weight alcohols by *Pichia pastoris* alcohol oxidase. Appl Microbiol Biotechnol 33,
1383 202–205. <https://doi.org/10.1007/BF00176525>
- 1384 Musa, M.M., Phillips, R.S., 2011. Recent advances in alcohol dehydrogenase-catalyzed
1385 asymmetric production of hydrophobic alcohols. Catal. Sci. Technol. 1, 1311–1323.
1386 <https://doi.org/10.1039/C1CY00160D>
- 1387 Nagashima, H., Inoue, J., Sasaki, E., Yamamoto, S., Sasaki, Y., Yamauchi-Inomata, Y.,
1388 Harayama, S., 1996. Long-chain *n*-alkanol dehydrogenase from *Pseudomonas putida*.
1389 J. Ferment. Bioeng. 82, 328–333. [https://doi.org/10.1016/0922-338X\(96\)89145-8](https://doi.org/10.1016/0922-338X(96)89145-8)
- 1390 Nguyen, Q.-T., Romero, E., Dijkman, W.P., de Vasconcellos, S.P., Binda, C., Mattevi, A.,
1391 Fraaije, M.W., 2018. Structure-Based Engineering of *Phanerochaete chrysosporium*
1392 Alcohol Oxidase for Enhanced Oxidative Power toward Glycerol. Biochemistry 57,
1393 6209–6218. <https://doi.org/10.1021/acs.biochem.8b00918>
- 1394 Oide, S., Tanaka, Y., Watanabe, A., Inui, M., 2019. Carbohydrate-binding property of a cell
1395 wall integrity and stress response component (WSC) domain of an alcohol oxidase from
1396 the rice blast pathogen *Pyricularia oryzae*. Enzyme Microb. Technol. 125, 13–20.
1397 <https://doi.org/10.1016/j.enzmictec.2019.02.009>
- 1398 Olson, L.P., Luo, J., Almarsson, Ö., Bruice, T.C., 1996. Mechanism of Aldehyde Oxidation
1399 Catalyzed by Horse Liver Alcohol Dehydrogenase. Biochemistry 35, 9782–9791.
1400 <https://doi.org/10.1021/bi952020x>
- 1401 Oppenheimer, N.J., Henahan, G.T.M., 1995. Horse Liver Alcohol Dehydrogenase-Catalyzed
1402 Aldehyde Oxidation, in: Weiner, H., Holmes, R.S., Wermuth, B. (Eds.), Enzymology
1403 and Molecular Biology of Carbonyl Metabolism 5, Advances in Experimental Medicine
1404 and Biology. Springer US, Boston, MA, pp. 407–415. <https://doi.org/10.1007/978-1->
1405 4615-1965-2_48
- 1406 Ottone, C., Bernal, C., Serna, N., Illanes, A., Wilson, L., 2018. Enhanced long-chain fatty
1407 alcohol oxidation by immobilization of alcohol dehydrogenase from *S. cerevisiae*. Appl
1408 Microbiol Biotechnol 102, 237–247. <https://doi.org/10.1007/s00253-017-8598-5>
- 1409 Ozimek, P., Veenhuis, M., Klei, I.J. van der, 2005. Alcohol oxidase: A complex peroxisomal,
1410 oligomeric flavoprotein. FEMS Yeast Research 5, 975–983.
1411 <https://doi.org/10.1016/j.femsyr.2005.06.005>
- 1412 Pal, S., Park, D.-H., Plapp, B.V., 2009. Activity of Yeast Alcohol Dehydrogenases on Benzyl
1413 Alcohols and Benzaldehydes. Characterization of ADH1 from *Saccharomyces*
1414 *carlsbergensis* and Transition State Analysis. Chem Biol Interact 178, 16–23.
1415 <https://doi.org/10.1016/j.cbi.2008.10.037>
- 1416 Parikka, K., Tenkanen, M., 2009. Oxidation of methyl α -D-galactopyranoside by galactose
1417 oxidase: products formed and optimization of reaction conditions for production of
1418 aldehyde. Carbohydrate Research 344, 14–20.
1419 <https://doi.org/10.1016/j.carres.2008.08.020>
- 1420 Park, D.H., Plapp, B.V., 1991. Isoenzymes of horse liver alcohol dehydrogenase active on
1421 ethanol and steroids. cDNA cloning, expression, and comparison of active sites. J. Biol.
1422 Chem. 266, 13296–13302. [https://doi.org/10.1016/S0021-9258\(18\)98838-1](https://doi.org/10.1016/S0021-9258(18)98838-1)
- 1423 Parker, J.K., 2015. 1 - Introduction to aroma compounds in foods, in: Parker, J.K., Elmore, J.S.,
1424 Methven, L. (Eds.), Flavour Development, Analysis and Perception in Food and
1425 Beverages, Woodhead Publishing Series in Food Science, Technology and Nutrition.
1426 Woodhead Publishing, pp. 3–30. <https://doi.org/10.1016/B978-1-78242-103-0.00001-1>

1427 Patel, J.M., 2009. Biocatalytic synthesis of atorvastatin intermediates. *J. Mol. Catal. B: Enzym*
1428 61, 123–128. <https://doi.org/10.1016/j.molcatb.2009.07.004>

1429 Paukner, R., Staudigl, P., Choosri, W., Haltrich, D., Leitner, C., 2015. Expression, purification,
1430 and characterization of galactose oxidase of *Fusarium sambucinum* in *E. coli*. *Protein*
1431 *Expr Purif* 108, 73–79. <https://doi.org/10.1016/j.pep.2014.12.010>

1432 Paukner, R., Staudigl, P., Choosri, W., Sygmund, C., Halada, P., Haltrich, D., Leitner, C., 2014.
1433 Galactose Oxidase from *Fusarium oxysporum* - Expression in *E. coli* and *P. pastoris*
1434 and Biochemical Characterization. *PLOS ONE* 9, e100116.
1435 <https://doi.org/10.1371/journal.pone.0100116>

1436 Pessôa, M.G., Paulino, B.N., Molina, G., Pastore, G.M., 2019. Prospective Research and
1437 Current Technologies for Bioflavor Production, in: *Bioprocessing for Biomolecules*
1438 *Production*. John Wiley & Sons, Ltd, pp. 93–123.
1439 <https://doi.org/10.1002/9781119434436.ch5>

1440 Petrus, M.L.C., Vijgenboom, E., Chaplin, A.K., Worrall, J.A.R., van Wezel, G.P., Claessen,
1441 D., 2016. The DyP-type peroxidase DtpA is a Tat-substrate required for GlxA
1442 maturation and morphogenesis in *Streptomyces*. *Open Biology* 6, 150149.
1443 <https://doi.org/10.1098/rsob.150149>

1444 Petschacher, B., Staunig, N., Müller, M., Schürmann, M., Mink, D., De Wildeman, S., Gruber,
1445 K., Glieder, A., 2014. Cofactor Specificity Engineering of *Streptococcus mutans* NADH
1446 Oxidase 2 For NAD(P)⁺ Regeneration in Biocatalytic Oxidations. *Comput. Struct.*
1447 *Biotechnol. J.* 9, e201402005. <https://doi.org/10.5936/csbj.201402005>

1448 Pickl, M., Fuchs, M., Glueck, S.M., Faber, K., 2015. Amination of ω-Functionalized Aliphatic
1449 Primary Alcohols by a Biocatalytic Oxidation–Transamination Cascade.
1450 *ChemCatChem* 7, 3121–3124. <https://doi.org/10.1002/cctc.201500589>

1451 Pietruszko, R., Crawford, K., Lester, D., 1973. Comparison of substrate specificity of alcohol
1452 dehydrogenases from human liver, horse liver, and yeast towards saturated and 2-enoic
1453 alcohols and aldehydes. *Arch. Biochem. Biophys.* 159, 50–60.
1454 [https://doi.org/10.1016/0003-9861\(73\)90428-1](https://doi.org/10.1016/0003-9861(73)90428-1)

1455 Porat, R., Deterre, S., Giampaoli, P., Plotto, A., 2016. The flavor of citrus fruit, in:
1456 *Biotechnology in Flavor Production*. John Wiley & Sons, Ltd, pp. 1–31.
1457 <https://doi.org/10.1002/9781118354056.ch1>

1458 Puetz, H., Puchl'ová, E., Vranková, K., Hollmann, F., 2020. Biocatalytic Oxidation of Alcohols.
1459 *Catalysts* 10, 952. <https://doi.org/10.3390/catal10090952>

1460 Quaglia, D., Irwin, J.A., Paradisi, F., 2012. Horse Liver Alcohol Dehydrogenase: New
1461 Perspectives for an Old Enzyme. *Mol Biotechnol* 52, 244–250.
1462 <https://doi.org/10.1007/s12033-012-9542-7>

1463 Rajangam, A.S., Gidda, S.K., Craddock, C., Mullen, R.T., Dyer, J.M., Eastmond, P.J., 2013.
1464 Molecular Characterization of the Fatty Alcohol Oxidation Pathway for Wax-Ester
1465 Mobilization in Germinated Jojoba Seeds. *Plant Physiol.* 161, 72–80.
1466 <https://doi.org/10.1104/pp.112.208264>

1467 Rannes, J.B., Ioannou, A., Willies, S.C., Grogan, G., Behrens, C., Flitsch, S.L., Turner, N.J.,
1468 2011. Glycoprotein labeling using engineered variants of galactose oxidase obtained by
1469 directed evolution. *J. Am. Chem. Soc.* 133, 8436–8439.
1470 <https://doi.org/10.1021/ja2018477>

1471 Rehn, G., Pedersen, A.T., Woodley, J.M., 2016. Application of NAD(P)H oxidase for cofactor
1472 regeneration in dehydrogenase catalyzed oxidations. *J. Mol. Catal. B: Enzym.* 134, 331–
1473 339. <https://doi.org/10.1016/j.molcatb.2016.09.016>

1474 Reid, M.F., Fewson, C.A., 1994. Molecular Characterization of Microbial Alcohol
1475 Dehydrogenases. *Crit. Rev. Microbiol.* 20, 13–56.
1476 <https://doi.org/10.3109/10408419409113545>

1477 Ressmann, A.K., Schwendenwein, D., Leonhartsberger, S., Mihovilovic, M.D., Bornscheuer,
1478 U.T., Winkler, M., Rudroff, F., 2019. Substrate-Independent High-Throughput Assay
1479 for the Quantification of Aldehydes. *Adv. Synth. Catal.* 361, 2538–2543.
1480 <https://doi.org/10.1002/adsc.201900154>

1481 Ribeaucourt, D., Bissaro, B., Guallar, V., Yemloul, M., Haon, M., Grisel, S., Alphand, V.,
1482 Brumer, H., Lambert, F., Berrin, J.-G., Lafond, M., 2021a. Comprehensive Insights into
1483 the Production of Long Chain Aliphatic Aldehydes Using a Copper-Radical Alcohol
1484 Oxidase as Biocatalyst. *ACS Sustainable Chem. Eng.* 9, 4411–4421.
1485 <https://doi.org/10.1021/acssuschemeng.0c07406>

1486 Ribeaucourt, D., Saker, S., Navarro, D., Bissaro, B., Drula, E., Correia, L.O., Haon, M., Grisel,
1487 S., Lapalu, N., Henrissat, B., O'Connell, R.J., Lambert, F., Lafond, M., Berrin, J.-G.,
1488 2021b. Time-resolved secretome analysis of three *Colletotrichum* species identifies
1489 copper radical alcohol oxidases for the production of fatty aldehydes. *bioRxiv*
1490 2021.06.09.447826. <https://doi.org/10.1101/2021.06.09.447826>

1491 Richmond, H.H., 1950. Preparation of cinnamaldehyde. US2529186A.

1492 Román, S., Sánchez-Siles, L.M., Siegrist, M., 2017. The importance of food naturalness for
1493 consumers: Results of a systematic review. *Trends Food Sci. Technol.* 67, 44–57.
1494 <https://doi.org/10.1016/j.tifs.2017.06.010>

1495 Romano, D., Villa, R., Molinari, F., 2012. Preparative Biotransformations: Oxidation of
1496 Alcohols. *ChemCatChem* 4, 739–749. <https://doi.org/10.1002/cctc.201200042>

1497 Romero, E., Gadda, G., 2014. Alcohol oxidation by flavoenzymes. *Biomol. Concepts* 5, 299–
1498 318. <https://doi.org/10.1515/bmc-2014-0016>

1499 Romero, E., Gómez Castellanos, J.R., Gadda, G., Fraaije, M.W., Mattevi, A., 2018. Same
1500 Substrate, Many Reactions: Oxygen Activation in Flavoenzymes. *Chem. Rev.* 118,
1501 1742–1769. <https://doi.org/10.1021/acs.chemrev.7b00650>

1502 Rowe, D., 2011. Overview of Flavor and Fragrance Materials, in: *Practical Analysis of Flavor*
1503 *and Fragrance Materials*. John Wiley & Sons, Ltd, pp. 1–22.
1504 <https://doi.org/10.1002/9781444343137.ch1>

1505 Salvi, F., Wang, Y.-F., Weber, I.T., Gadda, G., 2014. Structure of choline oxidase in complex
1506 with the reaction product glycine betaine. *Acta Crystallogr., Sect. D: Biol. Crystallogr.*
1507 70, 405–413. <https://doi.org/10.1107/S1399004713029283>

1508 Sankar, M., Nowicka, E., Carter, E., Murphy, D.M., Knight, D.W., Bethell, D., Hutchings, G.J.,
1509 2014. The benzaldehyde oxidation paradox explained by the interception of peroxy
1510 radical by benzyl alcohol. *Nat. Commun.* 5, 3332. <https://doi.org/10.1038/ncomms4332>

1511 Sattler, J.H., Fuchs, M., Tauber, K., Mutti, F.G., Faber, K., Pfeffer, J., Haas, T., Kroutil, W.,
1512 2012. Redox Self-Sufficient Biocatalyst Network for the Amination of Primary
1513 Alcohols. *Angew. Chem., Int. Ed.* 51, 9156–9159.
1514 <https://doi.org/10.1002/anie.201204683>

1515 Savitha, J., Ratledge, C., 1991. Alcohol oxidase of *Aspergillus flavipes* grown on hexadecanol.
1516 *FEMS Microbiol Lett* 80, 221–224. <https://doi.org/10.1111/j.1574-6968.1991.tb04665.x>

1517

1518 Sawamura, M., 2011. *Citrus Essential Oils: Flavor and Fragrance*. John Wiley & Sons.

1519 Schalk, M., 2014. Method for producing sclareol. US20140162332A1.

1520 Schalk, M., Pastore, L., Mirata, M.A., Khim, S., Schouwey, M., Deguerry, F., Pineda, V.,
1521 Rocci, L., Daviet, L., 2012. Toward a Biosynthetic Route to Sclareol and Amber
1522 Odorants. *J. Am. Chem. Soc.* 134, 18900–18903. <https://doi.org/10.1021/ja307404u>

1523 Schöpp, W., Aurich, H., 1976. Kinetics and reaction mechanism of yeast alcohol
1524 dehydrogenase with long-chain primary alcohols. *Biochem J* 157, 15–22.

1525 Schwarz, A., Wydra, S., 2020. Bio-based flavours and fragrances | Advanced Technologies for
1526 Industry. EUROPEAN COMMISSION.

- 1527 Sell, C.S., 2015. The Chemistry of Fragrances: From Perfumer to Consumer. Royal Society of
1528 Chemistry.
- 1529 Sellés Vidal, L., Kelly, C.L., Mordaka, P.M., Heap, J.T., 2018. Review of NAD(P)H-dependent
1530 oxidoreductases: Properties, engineering and application. *Biochim. Biophys. Acta,*
1531 *Proteins Proteomics* 1866, 327–347. <https://doi.org/10.1016/j.bbapap.2017.11.005>
- 1532 Serra, S., Fuganti, C., Brenna, E., 2005. Biocatalytic preparation of natural flavours and
1533 fragrances. *Trends Biotechnol.* 23, 193–198.
1534 <https://doi.org/10.1016/j.tibtech.2005.02.003>
- 1535 Sharon, I., Banfield, J.F., 2013. Genomes from Metagenomics. *Science* 342, 1057–1058.
1536 <https://doi.org/10.1126/science.1247023>
- 1537 Sheldon, R.A., Brady, D., 2019. Broadening the Scope of Biocatalysis in Sustainable Organic
1538 Synthesis. *ChemSusChem* 12, 2859–2881. <https://doi.org/10.1002/cssc.201900351>
- 1539 Silva-Jiménez, H., Zazueta-Novoa, V., Durón-Castellanos, A., Rodríguez-Robelo, C., Leal-
1540 Morales, C.A., Zazueta-Sandoval, R., 2009. Intracellular distribution of fatty alcohol
1541 oxidase activity in *Mucor circinelloides* YR-1 isolated from petroleum contaminated
1542 soils. *Antonie van Leeuwenhoek* 96, 527. <https://doi.org/10.1007/s10482-009-9368-x>
- 1543 Silveira, B.M.P., Barcelos, M.C.S., Vespermann, K.A.C., Pelissari, F.M., Molina, G., 2019. An
1544 Overview of Biotechnological Processes in the Food Industry, in: *Bioprocessing for*
1545 *Biomolecules Production.* John Wiley & Sons, Ltd, pp. 1–19.
1546 <https://doi.org/10.1002/9781119434436.ch1>
- 1547 Slabas, A.R., Elborough, K., Vanhanen, S., West, M., Cheng, Q., Lindner, N., Casey, J.,
1548 Sanglard, D., 1999. Improvements in or relating to fatty acid metabolism.
1549 WO1999047685A1.
- 1550 Snijder-Lambers, A.M., Vulfson, E.N., Doddema, H.J., 2010. Optimization of alcohol
1551 dehydrogenase activity and NAD(H) regeneration in organic solvents. *Recl. Trav.*
1552 *Chim. Pays-Bas* 110, 226–230. <https://doi.org/10.1002/recl.19911100517>
- 1553 Šola, K., Dean, G.H., Li, Y., Lohmann, J., Movahedan, M., Gilchrist, E.J., Adams, K.L.,
1554 Haughn, G.W., 2021. Expression Patterns and Functional Characterisation of
1555 *Arabidopsis* GALACTOSE OXIDASE-LIKE (GOXL) Genes Suggest Specialised
1556 Roles for Galactose Oxidases in Plants. *Plant Cell Physiol.*
1557 <https://doi.org/10.1093/pcp/pcab073>
- 1558 Šola, K., Gilchrist, E.J., Ropartz, D., Wang, L., Feussner, I., Mansfield, S.D., Ralet, M.-C.,
1559 Haughn, G.W., 2019. RUBY, a Putative Galactose Oxidase, Influences Pectin
1560 Properties and Promotes Cell-To-Cell Adhesion in the Seed Coat Epidermis of
1561 *Arabidopsis*. *Plant Cell* 31, 809–831. <https://doi.org/10.1105/tpc.18.00954>
- 1562 Spadiut, O., Olsson, L., Brumer, H., 2010. A comparative summary of expression systems for
1563 the recombinant production of galactose oxidase. *Microb Cell Fact* 9, 68.
1564 <https://doi.org/10.1186/1475-2859-9-68>
- 1565 Stemmer, W.P.C., 1994. Rapid evolution of a protein in vitro by DNA shuffling. *Nature* 370,
1566 389–391. <https://doi.org/10.1038/370389a0>
- 1567 Sun, L., Bulter, T., Alcalde, M., Petrounia, I.P., Arnold, F.H., 2002. Modification of Galactose
1568 Oxidase to Introduce Glucose 6-Oxidase Activity. *ChemBioChem* 3, 781–783.
1569 [https://doi.org/10.1002/1439-7633\(20020802\)3:8<781::AID-CBIC781>3.0.CO;2-8](https://doi.org/10.1002/1439-7633(20020802)3:8<781::AID-CBIC781>3.0.CO;2-8)
- 1570 Surburg, H., Panten, J., 2016. *Common Fragrance and Flavor Materials: Preparation, Properties*
1571 *and Uses.* John Wiley & Sons.
- 1572 Sützl, L., Foley, G., Gillam, E.M.J., Bodén, M., Haltrich, D., 2019. The GMC superfamily of
1573 oxidoreductases revisited: analysis and evolution of fungal GMC oxidoreductases.
1574 *Biotechnol. Biofuels* 12, 118. <https://doi.org/10.1186/s13068-019-1457-0>

1575 Sützl, L., Laurent, C.V.F.P., Abrera, A.T., Schütz, G., Ludwig, R., Haltrich, D., 2018.
1576 Multiplicity of enzymatic functions in the CAZy AA3 family. *Appl Microbiol*
1577 *Biotechnol* 102, 2477–2492. <https://doi.org/10.1007/s00253-018-8784-0>
1578 T. Pedersen, A., Birmingham, W.R., Rehn, G., Charnock, S.J., Turner, N.J., Woodley, J.M.,
1579 2015. Process Requirements of Galactose Oxidase Catalyzed Oxidation of Alcohols.
1580 *Org. Process Res. Dev.* 19, 1580–1589. <https://doi.org/10.1021/acs.oprd.5b00278>
1581 Takahashi, F., Igarashi, K., Hagihara, H., 2016. Identification of the fatty alcohol oxidase FAO1
1582 from *Starmerella bombicola* and improved novel glycolipids production in an FAO1
1583 knockout mutant. *Appl Microbiol Biotechnol* 100, 9519–9528.
1584 <https://doi.org/10.1007/s00253-016-7702-6>
1585 Tani, A., Sakai, Y., Ishige, T., Kato, N., 2000. Thermostable NADP⁺-Dependent Medium-
1586 Chain Alcohol Dehydrogenase from *Acinetobacter* sp. Strain M-1: Purification and
1587 Characterization and Gene Expression in *Escherichia coli*. *Appl. Environ. Microbiol.*
1588 66, 5231–5235. <https://doi.org/10.1128/AEM.66.12.5231-5235.2000>
1589 Tassano, E., Faber, K., Hall, M., 2018. Biocatalytic Parallel Interconnected Dynamic
1590 Asymmetric Disproportionation of α -Substituted Aldehydes: Atom-Efficient Access to
1591 Enantiopure (S)-Profens and Profenols. *Adv. Synth. Catal.* 360, 2742–2751.
1592 <https://doi.org/10.1002/adsc.201800541>
1593 Tassin, J.-P., Vandecasteele, J.-P., 1972. Separation and characterization of long-chain alcohol
1594 dehydrogenase isoenzymes from *Pseudomonas aeruginosa*. *Biochim. Biophys.*
1595 *Acta, Enzymol.* 276, 31–42. [https://doi.org/10.1016/0005-2744\(72\)90005-8](https://doi.org/10.1016/0005-2744(72)90005-8)
1596 Taylor, A.J., 2002. *Food Flavour Technology*. John Wiley & Sons.
1597 Thompson, C.E., Freitas, L.B., Salzano, F.M., 2018. Molecular evolution and functional
1598 divergence of alcohol dehydrogenases in animals, fungi and plants. *Genet Mol Biol* 41,
1599 341–354. <https://doi.org/10.1590/1678-4685-GMB-2017-0047>
1600 Thompson, M.P., Derrington, S.R., Heath, R.S., Porter, J.L., Mangas-Sanchez, J., Devine, P.N.,
1601 Truppo, M.D., Turner, N.J., 2019. A generic platform for the immobilisation of
1602 engineered biocatalysts. *Tetrahedron* 75, 327–334.
1603 <https://doi.org/10.1016/j.tet.2018.12.004>
1604 Tiemann, F., Haarmann, W., 1874. Ueber das Coniferin und seine Umwandlung in das
1605 aromatische Princip der Vanille. *Ber. Dtsch. Chem. Ges.* 7, 608–623.
1606 <https://doi.org/10.1002/cber.187400701193>
1607 Trivedi, A., Heinemann, M., Spiess, A.C., Daussmann, T., Büchs, J., 2005. Optimization of
1608 adsorptive immobilization of alcohol dehydrogenases. *J. Biosci. Bioeng.* 99, 340–347.
1609 <https://doi.org/10.1263/jbb.99.340>
1610 Truong, K.-N., Weger, L.B., Stahl, W., Mouhib, H., 2017. Favored Conformations of Carbonyl
1611 Compounds: A Structural Study of *n*-Octanal. *ChemPhysChem* 18, 2631–2636.
1612 <https://doi.org/10.1002/cphc.201700341>
1613 Tsuzuki, S., 2019. Higher Straight-Chain Aliphatic Aldehydes: Importance as Odor-Active
1614 Volatiles in Human Foods and Issues for Future Research. *J. Agric. Food Chem.* 67,
1615 4720–4725. <https://doi.org/10.1021/acs.jafc.9b01131>
1616 Turner, N.J., 2011. Enantioselective Oxidation of C–O and C–N Bonds Using Oxidases. *Chem.*
1617 *Rev.* 111, 4073–4087. <https://doi.org/10.1021/cr200111v>
1618 Ueda, M., Tanaka, A., 1990. [28] Long-chain alcohol dehydrogenase of *Candida* yeast, in:
1619 *Methods in Enzymology, Hydrocarbons and Methylotrophy*. Academic Press, pp. 171–
1620 175. [https://doi.org/10.1016/0076-6879\(90\)88030-E](https://doi.org/10.1016/0076-6879(90)88030-E)
1621 Urlacher, V.B., Koschorreck, K., 2021. Peculiarities and applications of aryl-alcohol oxidases
1622 from fungi. *Appl Microbiol Biotechnol.* <https://doi.org/10.1007/s00253-021-11337-4>
1623 Utekal, P., Tóth, C., Illésová, A., Koiš, P., Bocánová, L., Turňa, J., Drahovská, H., Stuchlík, S.,
1624 2014. Expression of soluble *Saccharomyces cerevisiae* alcohol dehydrogenase in

- 1625 *Escherichia coli* applicable to oxido-reduction bioconversions. *Biologia* 69, 722–726.
 1626 <https://doi.org/10.2478/s11756-014-0376-6>
- 1627 Vanhanen, S., West, M., Kroon, J.T.M., Lindner, N., Casey, J., Cheng, Q., Elborough, K.M.,
 1628 Slabas, A.R., 2000. A Consensus Sequence for Long-chain Fatty-acid Alcohol Oxidases
 1629 from *Candida* Identifies a Family of Genes Involved in Lipid ω -Oxidation in Yeast with
 1630 Homologues in Plants and Bacteria. *J. Biol. Chem.* 275, 4445–4452.
 1631 <https://doi.org/10.1074/jbc.275.6.4445>
- 1632 Vasić, K., Knez, Ž., Leitgeb, M., 2020. Immobilization of alcohol dehydrogenase from
 1633 *Saccharomyces cerevisiae* onto carboxymethyl dextran-coated magnetic nanoparticles:
 1634 a novel route for biocatalyst improvement via epoxy activation. *Sci. Rep.* 10, 19478.
 1635 <https://doi.org/10.1038/s41598-020-76463-x>
- 1636 Velonia, K., Smonou, I., 2000. Dismutation of aldehydes catalyzed by alcohol dehydrogenases.
 1637 *J. Chem. Soc., Perkin Trans.* 1 2283–2287. <https://doi.org/10.1039/B001039L>
- 1638 Vilím, J., Knaus, T., Mutti, F.G., 2018. Catalytic Promiscuity of Galactose Oxidase: A Mild
 1639 Synthesis of Nitriles from Alcohols, Air, and Ammonia. *Angew. Chem.* 130, 14436–
 1640 14440. <https://doi.org/10.1002/ange.201809411>
- 1641 Vogel, A., 2019. Enzyme Development Technologies, in: *Industrial Enzyme Applications*. John
 1642 Wiley & Sons, Ltd, pp. 25–45. https://doi.org/10.1002/9783527813780.ch1_2
- 1643 Vincentelli, R. (Ed.), 2019. *High-Throughput Protein Production and Purification: Methods and*
 1644 *Protocols, Methods in Molecular Biology*. Humana Press. [https://doi.org/10.1007/978-](https://doi.org/10.1007/978-1-4939-9624-7)
 1645 [1-4939-9624-7](https://doi.org/10.1007/978-1-4939-9624-7)
- 1646 Vrsalović Presečki, A., Makovšek, K., Vasić-Rački, Đ., 2012. Coenzyme Regeneration in
 1647 Hexanol Oxidation Catalyzed by Alcohol Dehydrogenase. *Appl Biochem Biotechnol*
 1648 167, 595–611. <https://doi.org/10.1007/s12010-012-9712-x>
- 1649 Vrsalović Presečki, A., Vasić-Rački, Đ., 2009. Mathematical modelling of the dehydrogenase
 1650 catalyzed hexanol oxidation with coenzyme regeneration by NADH oxidase. *Process*
 1651 *Biochem.* 44, 54–61. <https://doi.org/10.1016/j.procbio.2008.09.007>
- 1652 Waltz, E., 2015. Engineers of scent. *Nat. Biotechnol.* 33, 329–332.
 1653 <https://doi.org/10.1038/nbt.3191>
- 1654 Weimer, E.P., Rao, E., Brendel, M., 1993. Molecular structure and genetic regulation of SFA,
 1655 a gene responsible for resistance to formaldehyde in *Saccharomyces cerevisiae*, and
 1656 characterization of its protein product. *Molec. Gen. Genet.* 237, 351–358.
 1657 <https://doi.org/10.1007/BF00279438>
- 1658 Werf, M.J. van der, Ven, C. van der, Barbirato, F., Eppink, M.H.M., Bont, J.A.M. de, Berkel,
 1659 W.J.H. van, 1999. Stereoselective Carveol Dehydrogenase from *Rhodococcus*
 1660 *erythropolis* DCL14: A Novel Nicotinoprotein Belonging to the Short Chain
 1661 Dehydrogenase/Reductase Superfamily. *J. Biol. Chem.* 274, 26296–26304.
 1662 <https://doi.org/10.1074/jbc.274.37.26296>
- 1663 Whitaker, J.R., Voragen, A.G.J., Wong, D.W.S., 2002. *Handbook of Food Enzymology*. CRC
 1664 Press.
- 1665 Whittaker, J.W., 2005. The radical chemistry of galactose oxidase. *Arch. Biochem. Biophys.*
 1666 433, 227–239. <https://doi.org/10.1016/j.abb.2004.08.034>
- 1667 Whittaker, M.M., Whittaker, J.W., 1988. The active site of galactose oxidase. *J. Biol. Chem.*
 1668 263, 6074–6080.
- 1669 Wilkinson, A.J., Fersht, A.R., Blow, D.M., Carter, P., Winter, G., 1984. A large increase in
 1670 enzyme–substrate affinity by protein engineering. *Nature* 307, 187–188.
 1671 <https://doi.org/10.1038/307187a0>
- 1672 Wilkinson, D., Akumanyi, N., Hurtado-Guerrero, R., Dawkes, H., Knowles, P.F., Phillips,
 1673 S.E.V., McPherson, M.J., 2004. Structural and kinetic studies of a series of mutants of

1674 galactose oxidase identified by directed evolution. *Protein Eng. Des. Sel.* 17, 141–148.
1675 <https://doi.org/10.1093/protein/gzh018>

1676 Winkler, M., 2018. Carboxylic acid reductase enzymes (CARs). *Curr Opin Chem Biol*, 43, 23–
1677 29. <https://doi.org/10.1016/j.cbpa.2017.10.006>

1678 Wöhler, Liebig, 1832. Untersuchungen über das Radikal der Benzoesäure. *Ann. Pharm.* 3, 249–
1679 282. <https://doi.org/10.1002/jlac.18320030302>

1680 Wolberg, M., Hummel, W., Müller, M., 2001. Biocatalytic Reduction of β,δ -Diketo Esters: A
1681 Highly Stereoselective Approach to All Four Stereoisomers of a Chlorinated β,δ -
1682 Dihydroxy Hexanoate. *Chem. - Eur. J.* 7, 4562–4571. [https://doi.org/10.1002/1521-3765\(20011105\)7:21<4562::AID-CHEM4562>3.0.CO;2-4](https://doi.org/10.1002/1521-3765(20011105)7:21<4562::AID-CHEM4562>3.0.CO;2-4)

1683

1684 Wongnate, T., Chaiyen, P., 2013. The substrate oxidation mechanism of pyranose 2-oxidase
1685 and other related enzymes in the glucose–methanol–choline superfamily. *FEBS J.* 280,
1686 3009–3027. <https://doi.org/10.1111/febs.12280>

1687 Woodley, J.M., 2019. Accelerating the implementation of biocatalysis in industry. *Appl*
1688 *Microbiol Biotechnol* 103, 4733–4739. <https://doi.org/10.1007/s00253-019-09796-x>

1689 Wu, H., Tian, C., Song, X., Liu, C., Yang, D., Jiang, Z., 2013. Methods for the regeneration of
1690 nicotinamide coenzymes. *Green Chem.* 15, 1773–1789.
1691 <https://doi.org/10.1039/C3GC37129H>

1692 Wu, S., Snajdrova, R., Moore, J.C., Baldenius, K., Bornscheuer, U., 2020. Biocatalysis:
1693 Enzymatic Synthesis for Industrial Applications. *Angew. Chem., Int. Ed.*
1694 <https://doi.org/10.1002/anie.202006648>

1695 Xu, F., Golightly, E.J., Schneider, P., Berka, R.M., Brown, K.M., Johnstone, J.A., Baker, D.H.,
1696 Fuglsang, C.C., Brown, S.H., Svendsen, A., Klotz, A.V., 2000. Expression and
1697 characterization of a recombinant *Fusarium* spp. galactose oxidase. *Appl Biochem*
1698 *Biotechnol* 88, 23. <https://doi.org/10.1385/ABAB:88:1-3:023>

1699 Yamada, T., Nawa, H., Kawamoto, S., Tanaka, A., Fukui, S., 1980. Subcellular localization of
1700 long-chain alcohol dehydrogenase and aldehyde dehydrogenase in *n*-alkane-grown
1701 *Candida tropicalis*. *Arch. Microbiol.* 128, 145–151.
1702 <https://doi.org/10.1007/BF00406151>

1703 Yanai, H., Doi, K., Ohshima, T., 2009. *Sulfolobus tokodaii* ST0053 Produces a Novel
1704 Thermostable, NAD-Dependent Medium-Chain Alcohol Dehydrogenase. *Appl Environ*
1705 *Microbiol* 75, 1758–1763. <https://doi.org/10.1128/AEM.01392-08>

1706 Yin, D., Urresti, S., Lafond, M., Johnston, E.M., Derikvand, F., Ciano, L., Berrin, J.-G.,
1707 Henrissat, B., Walton, P.H., Davies, G.J., Brumer, H., 2015. Structure–function
1708 characterization reveals new catalytic diversity in the galactose oxidase and glyoxal
1709 oxidase family. *Nat. Commun.* 6. <https://doi.org/10.1038/ncomms10197>

1710 Zavec, D., Troyer, C., Maresch, D., Altmann, F., Hann, S., Gasser, B., Mattanovich, D., 2021.
1711 Beyond alcohol oxidase: The methylotrophic yeast *Komagataella phaffii* utilizes
1712 methanol also with its native alcohol dehydrogenase Adh2. *FEMS Yeast Research.*
1713 <https://doi.org/10.1093/femsyr/foab009>

1714 Zhang, S., Ruccolo, S., Fryszkowska, A., Klapars, A., Marshall, N., Strotman, N.A., 2021.
1715 Electrochemical Activation of Galactose Oxidase: Mechanistic Studies and Synthetic
1716 Applications. *ACS Catal.* 7270–7280. <https://doi.org/10.1021/acscatal.1c01037>

1717 Zhao, S., Lin, Z., Ma, W., Luo, D., Cheng, Q., 2008. Cloning and Characterization of Long-
1718 Chain Fatty Alcohol Oxidase LjFAO1 in *Lotus japonicus*. *Biotechnol. Prog.* 24, 773–
1719 779. <https://doi.org/10.1021/bp0703533>

1720 Zhou, J., Chen, Z., Wang, Y., 2020. Bioaldehydes and beyond: Expanding the realm of
1721 bioderived chemicals using biogenic aldehydes as platforms. *Curr Opin Chem Biol* 59,
1722 37–46. <https://doi.org/10.1016/j.cbpa.2020.04.007>

1723 Zviely, M., 2012. Aroma Chemicals, in: Kirk-Othmer Encyclopedia of Chemical Technology.
1724 John Wiley & Sons, pp. 1–33.
1725 <https://doi.org/10.1002/0471238961.0118151326220905.a01.pub2>
1726