

# **Microplastic contamination in tropical fishes: an assessment of different feeding habits**

Anne K.S. Justino, Véronique Lenoble, Latifa Pelage, Guilherme V.B. Ferreira, Rafaela Passarone, Thierry Frédou, Flávia Lucena Frédou

## **To cite this version:**

Anne K.S. Justino, Véronique Lenoble, Latifa Pelage, Guilherme V.B. Ferreira, Rafaela Passarone, et al.. Microplastic contamination in tropical fishes: an assessment of different feeding habits. Regional Studies in Marine Science, 2021, 45, pp.101857. 10.1016/j.rsma.2021.101857. hal-03619681

# **HAL Id: hal-03619681 <https://hal.science/hal-03619681>**

Submitted on 25 Mar 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

- Microplastic contamination in tropical fishes: an assessment of different feeding habits
- 2 Anne K. S. Justino <sup>1,2</sup>, Véronique Lenoble <sup>2</sup>, Latifa Pelage <sup>1</sup>, Guilherme V. B Ferreira <sup>1</sup>, Rafaela Passarone <sup>1</sup>,
- 3 Thierry Frédou<sup>1</sup>, Flávia Lucena Frédou<sup>1</sup>
- <sup>1</sup>Departamento de Pesca e Aquicultura da Universidade Federal Rural de Pernambuco DEPAQ/UFRPE, Rua Dom Manuel de Medeiros, s/n, 52171-900, Recife, Brasil.
- <sup>2</sup> Université de Toulon, Aix Marseille Univ., CNRS/INSU, IRD, MIO UM 110, Mediterranean Institute of Oceanography, CS 60584, 83041 – TOULON, France
- 8 Corresponding author: anne.karen@hotmail.com
- Abstract

Marine ecosystems are reported to be contaminated by microplastics (MPs) (< 5 mm), however the ecological mechanisms involved in the ingestion of debris by marine organisms are relatively unknown. By developing and optimising an appropriate protocol of gut digestion for fish species, this study explores a tropical estuarine environment to unriddle the processes responsible for the different ingestion rates of plastic debris. A total of 82 fishes with different 15 feeding habits were analysed, *Centropomus undecimalis* ( $n = 30$ ; Piscivore), *Bairdiella ronchus* 16 (n = 21; Zoobenthivore) and *Gobionellus stomatus* (n = 31; Detritivore). The microplastic 17 ingestion varied with the feeding strategy; C. undecimalis, the predator, was the most contaminated species. Overall, most MPs were fibres (47%), followed by pellets (40%) and fragments (13%), although these proportions varied among species. A high level of contamination was found in the Estuarine Complex of Santa Cruz Channel, Northeast of Brazil, with many sources of MPs contaminants, as for many other tropical estuaries, which probably accumulate in the sediment and water column, with unknown consequences for human health.

Keywords: Trophic transfer, Plastic extraction, Trophic guild, Plastic pollution, Tropical Atlantic

#### Introduction

Estuaries are well known for providing ecosystem services by supplying essential goods (raw materials and food) and offering an attractive environment for population growth and cultural activities (Atkins et al., 2011). In addition, they carry out an important ecological function since fishes use these environments for protection, feeding, reproduction, settlement, and nursery (Krumme et al., 2008; Potter et al., 2013; Lima et al., 2016b; Ramos et al., 2016;

Ferreira et al., 2019). Worldwide, estuaries are usually surrounded by large metropolises, and the expansion of the urban population is directly associated with impacts on the coastal ecosystems (Freeman et al., 2019). Plastic pollution is one of the most significant environmental problems of the 21st century since large quantities of these materials are being mismanaged and/or illegally disposed of in marine ecosystems (Dauvergne, 2018; Ostle et al., 2019). Once 36 in the environment, these plastic materials are weakened by natural processes  $(e.g.,)$ hydrodynamic forces, solar radiation, and biological actions) (Thompson et al., 2004; Jambeck et al., 2015) and fragmented into smaller parts known as microplastics (< 5 mm).

Plastic debris poses several risks to marine biota (Galloway et al., 2017). Ingestion can be hazardous, causing digestive injuries, decreasing predatory efficiency, or inducing toxic effects (de Sá et al., 2015; Moore, 2008; Teuten et al., 2007; Barboza et al., 2018). Microplastics can adsorb pollutants available in the water column, such as persistent organic pollutants (POPs) (Oehlmann et al., 2009; Frias et al., 2010; Rochman et al., 2013) or heavy metals (Ashton et al., 2010; Holmes et al., 2012), which may be further bioaccumulated and biomagnified in the food web (Batel et al., 2016; Teuten et al., 2009). Furthermore, microplastics may be transferred in the trophic chain by predating contaminated prey (Ferreira et al., 2016; 2019; Chagnon et al., 2018). The trophic transfer has already been pointed out as a relevant contamination mechanism for estuarine species (Athey et al., 2020).

Although the long-term effects of microplastic contamination are still unknown, the scientific community continually emphasises the importance of using reliable and replicable methods of investigations (Hermsen et al., 2018; Markic et al., 2019). A useful approach to obtain plastic in marine wildlife is based on chemical digestion protocols, which are efficient and low cost to work with (Karami et al., 2017; Kühn et al., 2017). Digestion protocols are a practical and secure way to extract and isolate microplastics in organisms, being widely used in the investigation of fish contamination by plastics (Foekema et al., 2013; Tanaka & Takada 2016; Bellas et al., 2016; Hermsen et al., 2017; Bessa et al., 2018; Pellini et al., 2018; Herrera et al., 2019; Su et al., 2019). However, the lack of security procedures (e.g., cleaned workroom and blanks procedures) during the implementation of digestion protocols may lead to the overestimation of contaminants since samples are more prone to airborne and cross-contamination (Torre et al., 2016; Hermsen et al., 2017; Hermsen et al., 2018). Moreover, the procedure must ensure data reliability through an effective and careful extraction of the microplastics and the implementation of a robust sample size with a minimum of 10 samples (Markic et al., 2019).

Several studies conducted in South Atlantic estuaries evaluated the ingestion of plastics by marine organisms, mammals (Attademo et al., 2015), mussels (Santana et al., 2016; Birnstiel et al., 2019), turtles (Guebert-Bartholo et al., 2011), and microplastics interactions with ichthyoplankton (Lima et al., 2016a), and fish assemblages (Vendel et al., 2017). However, few studies investigated the ecological and biological dynamics associated with microplastic intakes on wild fishes (Ferreira et al., 2016; 2018; Amorim et al., 2020; Dantas et al., 2020). Furthermore, the bioaccumulation of microplastics in marine species is highly influenced by feeding strategies (Miller et al., 2020). However, the possible correlation between feeding habits and MPs ingestion is not yet well known.

Three estuarine species were chosen to test our hypothesis that microplastic ingestion varies according to the feeding strategies: (1) Centropomus undecimalis (Bloch, 1792); (2) Bairdiella ronchus (Cuvier, 1830); and (3) Gobionellus stomatus (Starks, 1913). C. undecimalis is an essential economic living resource for the commercial and subsistence fisheries in South America (Carpenter, 2002). While adults of C. undecimalis inhabit coastal areas and migrate towards the estuary, their juvenile stage uses the estuarine areas as a nursery ground (Ferreira et al., 2019). C. undecimalis is classified as opportunistic predator, feeding on a large variety of available preys in the environment, with a piscivorous tendency (Lira et al., 2017; Ferreira et al., 2019). B. ronchus is classified as zoobenthivore (Ferreira et al., 2019), which preys on 82 invertebrates associated with the sediment (Elliott et al., 2007). G. *stomatus* is a detritivore fish (Ferreira et al., 2019) consuming detritus and microphytobenthos (Elliott et al., 2007). Although 84 B. ronchus and G. stomatus were not of economic importance, they play a significant ecological 85 role within the estuarine ecosystem and are the main energy source for the C. *undecimalis* (Lira et al., 2018; Gonzalez et al., 2019).

Understanding the role of microplastics as a component of anthropic pollution in this ecosystem is crucial to assess adverse impacts on regional biodiversity and the quality of fisheries resources that are being traded and consumed. In this context, the present study aims 90 to  $(i)$  apply an adapted extraction protocol to assess MPs in the digestive tract of fishes, assuring the integrated quality control and appropriate sampling size, (ii) describe microplastics

92 contamination in fishes in estuarine waters, and *(iii)* identify the main types of microplastics considering the different feeding strategies.

#### Materials and Methods

#### Study area

The Estuarine Complex of the Santa Cruz Channel (ECSC) (Fig.1) is located along the northeast Brazilian coast. The climate is classified as tropical, hot and humid, with an average 98 of 26 $^{\circ}$ C ( $\pm$ 2.8 $^{\circ}$ C) annual air temperature and two seasons defined according to the level of precipitation (rainy and dry seasons) (Medeiros et al., 2001).

The ECSC is a tidal channel that surrounds the Itamaracá Island, separating it from the mainland, with a total area of 22 km, a maximum width of 1.5 km and a depth between 4 and 5 m (Lira et al., 2017). The main tributaries are formed by the Arataca, Botafogo and Igarassu Rivers, and the predominant vegetation is the mangrove forest (Medeiros et al., 2001). The ECSC is surrounded by two cities (Itapissuma and Itamaracá), which have their economies mainly focused on the industrial and agricultural sectors (IBGE, 2011), whereas the local economy is primarily supported by artisanal fisheries, aquaculture, and tourism (de Moura et al., 2009).



#### 

Figure 1. Map of the Estuarine Complex of the Santa Cruz Channel (ECSC) adapted from Pelage et al., 2019.

#### Sampling and Laboratory procedures

Three demersal species were selected for this study, given their commercial and or subsistence importance for riverine populations and their ecological interactions in the trophic chain (Lira et al., 2017; 2018): Gobionellus stomatus, Bairdiella ronchus, and Centropomus undecimalis, classified as Detritivores, Zoobenthivores and Piscivores respectively, according 115 to Ferreira et al. (2019) for the area. For the common Snook C. *undecimalis*, individuals were collected in the juvenile stage (Total Length < 26.3 cm) (Ferreira et al., 2019), ensuring that the obtained specimens had not left the estuarine ecosystem. This procedure was carried out to ensure that contaminants' intake occurred within the estuarine boundary since adults of C. *undecimalis* perform migrations toward coastal areas. As *B. ronchus* and *G. stomatus* are not caught outside the estuary of ECSC (Ferreira et al., 2019), we did not fix these species' ontogeny, and juveniles and adults were analysed. A total of 82 individuals were obtained by local fishermen. After each sampling, the specimens were labelled, and the individuals were

frozen. In the laboratory, individuals were identified (Menezes & Figueiredo, 2000), measured and weighted. Each individual had their organs (stomach and intestine) carefully removed, weighed and stored again for the digestion analysis. The minimum sample size for each species was 20 individuals to avoid the bias of a low sample size, which is twice the number suggested by Markic et al. (2019).

## Quality control and Extraction protocol

Firstly, to guarantee quality control and avoid potential airborne contamination, several steps were implemented. The entire process of digestion, filtration, identification and storage of the microplastic samples was carried out in a cleaned and reserved room within the main laboratory, reserved only for microplastic analysis. The flow of people was limited; cotton lab coats and disposable latex gloves were worn during the entire process. Also, all used tools were previously cleaned with alcohol 70% and rinsed with filtered distilled water. The solutions utilised in the various procedures were filtered using a vacuum pump system (equipped with 136 laboratory glassware) through a 47 mm GF/F 0.7 µm glass fibre filter (Whatman).

The digestive tracts were rinsed with distilled water before being placed in a beaker, submerged in NaOH (1 mol/L; PA 97%) solution, and covered by a glass lid (Fig. 2). The entire digestive tracts were submitted to NaOH without further dissection of those organs to avoid airborne contamination during handling. The proportion used was 1:100 w/v for 1 g of digestive 141 tract weight, 100 ml of NaOH (1 mol/L) solution. The mixture was oven-dried at  $60^{\circ}$  C for 24 h, mixed from time to time with a glass stick. The samples digested in the previous step, and the procedural blanks were filtered using a vacuum pump system through a 47 mm glass fibre filter (GF/F 0.7 µm Whatman). After filtration, filters were carefully set in a Petri dish (47 mm 145 diameter) and covered. These filters were oven-dried at 60° C for 24 h. The microplastics were identified using a stereomicroscope (Zeiss Stemi 508) with 6.3 – 50 times magnification with a 147 detection limit of 20 µm, photographed (Axiocam 105 Color), measured (Zeiss Zen 3.2) from 148 the filter and stored in covered Petri dishes (Fig. 2). They were then categorised by type:  $(i)$ fibres (filamentous shape), (ii) fragments (irregular shape) or (iii) pellets (spherical shape). The digestion protocol is a useful tool to separate the organic materials and facilitates visual identification, although it is not sufficient for identifying the polymers. Thereby, we also applied the method described by Ferreira et al. (2019) to confirm plastic debris by drying the samples in an oven to verify whether their physical characteristics changed or not.

Procedural blanks were made for each day of analysis before beginning the sample digestion. For blanks, a beaker was filled with 50 ml of NaOH (1 mol/L) solution and covered with a glass lid, and these blanks were exposed to the same protocol applied to the samples. A total of 10 blanks were made; among them, four blanks were observed with eight tiny particles  $(\leq 100 \mu m)$  considered as paint fragments. Thereby, all particles further identified with any resemblance to those observed in the blanks were excluded from posterior analyses.



Figure 2. Flowchart with the stepwise of the extraction protocol.

## Data Analysis

Kruskal-Wallis tests were used to verify if ingested microplastics (total MPs and different types) presented significant differences in number and length in relation to species (C. undecimalis, B. ronchus and G. stomatus). When the Kruskal-Wallis showed significant 164 differences, the *post hoc* pairwise comparisons Dunn's test was performed to investigate the sources of variance (Dunn, 1964). All statistical analyses were carried out with the software R version 3.6.3 (R Core Team, 2020) and were conducted considering a level of significance of 5%.

#### 168 Results

A total of 30 individuals of Centropomus undecimalis, 31 Gobionellus stomatus, 21 Bairdiella ronchus were analysed. Microplastics were present in 77% of C. undecimalis, 74% 171 of G. stomatus, and 67% of B. ronchus individuals. A total of 176 particles of MPs were recovered from 82 fishes (Table I).

173 According to the number of MPs, ingestion significantly differed between species (p-174 value  $\leq$  0.05), C. undecimalis being the most contaminated (3.3  $\pm$  2.9 MPs fish<sup>-1</sup>), followed by 175 G. stomatus  $(1.7 \pm 1.5 \text{ MPs fish}^{-1})$  and B. ronchus  $(1.2 \pm 1.3 \text{ MPs fish}^{-1})$  (Fig. 3). Significant 176 differences were recorded between C. *undecimalis* and B. *ronchus* (chi-squared = 7.873, df = 177 2, p-value  $\leq$  0.05). Concerning the length of ingested MPs, no significant differences among 178 the same type of MPs were observed when comparing MPs' size between species.

Regarding the types of MPs ingested by fishes, most were fibres (47%), followed by pellets (40%) and fragments (13%), and proportions varied between the species. C. undecimalis registered 68% of pellets, 28% of fibres, and 4% of fragments, B. ronchus and G. stomatus registered 23% and 4% of pellets, 62% and 71% of fibres and 15% and 25% of fragments, respectively (Fig. 4 and 5).



185 Figure 3. Mean number of microplastics ingested by fishes collected in the Estuarine Complex of the Santa Cruz Channel. The asterisks represent the statistical differences with a significance of 0.05.



Figure 4. Different types of microplastics (fibres, fragments, and pellets) ingested by fish species, expressed as a percentage.

# Types of microplastics



Figure 5. Types of microplastics ingested by fishes of the estuarine complex of Santa Cruz Channel. a) fragment (irregular 191 shape); b) pellet (spherical shape); and c) fibre (filamentous shape).



#### Discussion

### **Quality Assurance and Quality Control**

In our study, we applied an analytical method to extract microplastics (MPs) in the digestive tract of fishes, involving a careful procedure of quality control, using sodium hydroxide (NaOH 1 M), and a sampling size with a minimum of 20 individuals for each species, following the recent recommendations of Markic et al. (2019) and Hermsen et al. (2018).

The application of digestion protocols to extract microplastics in marine biota has been growing worldwide (Lusher et al., 2017). Digestion protocol is a reliable method for isolating microplastics in the biota, facilitating the observation of non-organic materials. For fish, the most common method used is alkaline digestion. Indeed, for many authors, this is considered the most suitable method to remove organic material and isolate plastic debris (Karami et al., 2017; Kühn et al., 2017; Schirinzi et al., 2020). Although the efficiency of potassium hydroxide (KOH 10%) has been more often tested, the use of sodium hydroxide (NaOH 1M) has also been widely tested and demonstrated to be very useful (Bellas et al., 2016; Baalkhuyur et al., 2018; Budimir et al., 2018; Morgana et al., 2018; Wieczorek et al., 2018; Su et al., 2019), and ensures 221 polymer integrity after chemical digestion (Budimir et al., 2018).

Although adaptations of digestion protocol have been carried out around the world (Kühn et al., 2017; Karami et al., 2017), so far, only three studies in estuaries of South America have used a chemical digestion protocol for microplastics extraction in the digestive tract of fishes, with the implementation of adequate quality assurance and quality control (QA/QC) (Arias et al., 2019; Ribeiro-Brasil et al., 2020; Garcés-Ordóñez, et al., 2020). Implementing 227 QA/QC procedures is essential to avoid cross-contamination of microplastic samples (Lusher et al., 2017; Hermsen et al., 2017; O'Connor et al., 2019). This treatment is necessary to minimise over/underestimation of microplastics due to airborne contamination and or loss of particles during sample handling. Moreover, the sample size is also an essential factor in the analysis. Among the studies in South America, only Arias et al. (2019) have chosen a sample 232 size  $n > 10$ . Such bias is unlikely to occur in our study, where the sample size was the highest in South America. Therefore, our results are expected to adequately reflect the contamination in the Estuarine Complex of Santa Cruz Channel (ECSC), and the protocol here used can safely be replicated in other studies.

#### Microplastic ingestion by fishes

In our study, microplastic (MP) contamination rates were high (frequency of occurrence of 73%) for the three analysed demersal estuarine species. Overall, demersal estuarine species are frequently reported to have a high ingestion rate of MPs (Ferreira et al., 2016; 2019; Arias et al., 2019). Demersal species inhabit and feed on the fauna associated with the substrate (Elliot et al., 2017), which might be in direct contact with contaminated sediment. Indeed, the estuarine sediment is a significant accumulation zone for MPs (Zhang, 2017). As estuaries are a transitional ecosystem, often surrounded by urban areas and exposed to domestic sewage discharge, they receive contaminants of both riverine and tidal inputs (Lebreton et al., 2017). Thereby, these ecosystems are more prone to be contaminated by MPs, especially fibres (Browne et al., 2011; Bessa et al., 2018). In addition to the sewage discharge, fibres can also be originated by the use, maintenance, discarding and loss of fisheries gear (Lima et al., 2014). MPs fibres occurred the most in the estuarine bottom, possibly due to the rapid sinking of these types of MPs (Lima et al., 2014).

Overall, the most ingested MP type were fibres, as observed in most of the studies worldwide (Wright et al., 2013; Foekema et al., 2013; Bellas et al., 2016; Herrera et al., 2019; Bessa et al., 2018). Although all types of plastics were ingested by the species analysed in this 253 study, ingestion rates varied. Fibres were the most frequent type in G. stomatus (representing 254 71% of the ingested MPs) and B. ronchus ( $68\%$ ). G. stomatus and B. ronchus are detritivore and zoobenthivore species, respectively, depending on the organisms associated with the substrate or the organic matter available. Consequently, they are more vulnerable to the MPs fibres contaminating estuarine sediments. However, pellets were the most frequent type in C. *undecimalis* (68%), differently from the results observed by Ferreira et al. (2019) for the same species, which registered mostly ingestion of filaments. Pellets also dominated the diet of fish along the coast of Salvador (Brazil) (Miranda & Carvalho-Souza,2016) and from the Amazon estuary (Pegado et al., 2018). Different from the other types of MP ingested, pellets are primary microplastics (Fendall et al., 2009), which are manufactured as MPs mainly for the cosmetics industry (e.g., microbeads). This type of MP can be accidentally discharged into the environment during the transport of this raw material (Ogata et al., 2009) or by the release of domestic sewage (Tanaka & Takada 2016). The increase in urbanisation in the Santa Cruz Channel (Pelage et al., 2019) surely amplified the sewage discharge in this area. Pellets can be

found floating in the water column, and they can even have fish eggs attached to them (Ivar do 268 Sul et al., 2013). Predatory fishes such as C. *undecimalis* can ingest the pellets directly by confusing them with their natural prey. Moreover, when feeding, opportunistic predators ingest a large amount of prey, which might increase the momentary build-up of MPs particles prior to egestion.

Predator species are more vulnerable to microplastic contamination due to the trophic transference, which occurs when they ingest contaminated prey (Eriksson and Burton, 2003; Ferreira et al., 2016; Chagnon et al., 2018; Nelms et al., 2018; Ferreira et al., 2019). Consequently, as we observed among the three analysed demersal fish species, the predator, C. *undecimalis*, had the highest contamination rate  $(3.3 \pm 2.9 \text{ MPs fish}^{-1})$  despite being in its early 277 life stage (juvenile), followed by G. stomatus (1.7  $\pm$  1.5 MPs fish<sup>-1</sup>) and B. ronchus (1.2  $\pm$  1.3 MPs fish<sup>-1</sup>). Differences in plastic ingestion might be associated with the feeding strategy of species (Wright et al., 2013; Miller et al., 2020). Our study tested and confirmed the hypothesis that microplastic ingestion varies with the different feeding strategies. However, regardless of the diet preferences of the species, in our study area, it is reported many sources of MPs contaminants, which probably accumulate in the sediment and water column, negatively affecting the life-strategies of fish species and mostly the juvenile stages which utilise estuaries as a nursery ground. Thus, the predators are more prone to be contaminated by microplastics through two main exposure routes: (1) the highly contaminated estuarine habitats and (2) the ingestion of contaminated prey.

Anthropogenic activities in the ECSC (urban areas, manufacturers, aquaculture plants, and sugarcane fields) are found surrounding the whole floodplain, and these activities might be an important source of microplastics and other contaminants (e.g., pesticides and heavy metals). Indeed, the ECSC has registered many impacts such as habitat loss and mercury releases (Pelage et al., 2019; Araújo et al., 2019; Albuquerque et al., 2019), which likely affect the estuarine community. Besides, our study has identified high contamination by microplastics in fish species that are a relevant source of protein locally and regionally. Despite being a Marine Protected Area, which provides essential ecosystems services, there is a lack of awareness and public policies, highlighting the importance of monitoring and management policies to control and mitigate social and health problems.

Further studies are needed regarding the microplastic impacts on marine fauna and whether they could transfer adsorbed pollutants such as persistent organic pollutants (POPs), heavy metals, and plastic additives to the food web are necessary, as microplastics particles can be transferred along the trophic chain, the chances to accumulate other pollutants in the food web increases. In fishery resources, this question is a public health matter because it is linked to human uptake of these pollutants. Our study also emphasises the importance of implementing protocols to extract microplastics in biological samples, which guarantee the quality of samples, avoid under or overestimation and airborne contamination, and which can be easily replicable.

#### Acknowledgement

This study was supported by the LMI TAPIOCA program CAPES/COFECUB (88881.142689/2017-01); FUNBIO and Humanize under the grant "Programa Bolsas Funbio - Conservando o Futuro 2019 (02/2019)"; FACEPE (Fundação de Amparo à Ciência e Tecnologia do Estado de Pernambuco); and MicroplastiX Project JPI-Oceans/CONFAP/FACEPE (n°APQ-0035-1.08/19). We thank the CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico), which provided research grants to Flávia Lucena Frédou (CNPq n° 308554/2019-1) and Thierry Frédou (CNPq 307422/2020-8); FACEPE for granting a scholarship (BFP-0049-5.06/20) and CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior) for granting a scholarship at doctoral level for the first author. All authors thanks to the members of the laboratory BIOIMPACT for the support in the sampling procedures and the fishermen of Itapissuma for the sampling collect.

#### REFERENCES

Amorim, A. L. A., Ramos, J. A., & Júnior, M. N., 2020. Ingestion of microplastic by ontogenetic phases of Stellifer brasiliensis (Perciformes, Sciaenidae) from the surf zone of tropical beaches. Marine pollution bulletin, 158, 111214. doi.org/10.1016/j.marpolbul.2020.111214

Albuquerque, P. T., Frédou, T., Arruda, G. N., Silva Filho, C. A., Nascimento, A. F., da Silva, M. J., & De França, E. J., 2019. Tracking Hg historical inputs by Pb-210 geochronology for the Itapessoca Estuarine Complex, Pernambuco, Brazil. Journal of Radioanalytical and Nuclear Chemistry, 321(3), 875-883. doi.org/10.1007/s10967-019-06665-9

- Araújo, P. R. M., Biondi, C. M., do Nascimento, C. W. A., da Silva, F. B. V., & Alvarez, A. M., 2019. Bioavailability and sequential extraction of mercury in soils and organisms of a mangrove contaminated by a chlor-alkali plant. Ecotoxicology and environmental safety, 183, 109469. doi.org/10.1016/j.ecoenv.2019.109469
- Arias, A.H., Ronda, A.C., Oliva, A.L., 2019. Evidence of Microplastic Ingestion by Fish from the Bahía Blanca Estuary in Argentina, South America. Bull Environ Contam Toxicol 102, 750–756. doi:10.1007/s00128-019-02604-2
- Ashton, K., Holmes, L., & Turner, A., 2010. Association of metals with plastic production pellets in the marine environment. Marine Pollution Bulletin, 60(11), 2050-2055. doi.org/10.1016/j.marpolbul.2010.07.014
- Athey, S. N., Albotra, S. D., Gordon, C. A., Monteleone, B., Seaton, P., Andrady, A. L., Taylor, A. R. & Brander, S. M., 2020. Trophic transfer of microplastics in an estuarine food chain and the effects of a sorbed legacy pollutant. Limnology and Oceanography Letters, 5(1), 154-162. doi: 10.1002/lol2.10130
- Atkins, J. P., Burdon, D., Elliott, M., & Gregory, A. J., 2011. Management of the marine environment: Integrating ecosystem services and societal benefits with the DPSIR framework in a systems approach. Marine Pollution Bulletin, 62(2), 215–226. doi.org/10.1016/j.marpolbul.2010.12.012
- Attademo, F. L. N., Balensiefer, D. C., da Bôaviagem Freire, A. C., de Sousa, G. P., da Cunha, F. A. G. C., & de Oliveira Luna, F., 2015. Debris ingestion by the Antillean Manatee (Trichechus manatus manatus). Marine pollution bulletin, 101(1), 284-287. doi.org/10.1016/j.marpolbul.2015.09.040
- Baalkhuyur, F. M., Dohaish, E. J. A. B., Elhalwagy, M. E., Alikunhi, N. M., AlSuwailem, A. M., Røstad, A., ... & Duarte, C. M., 2018. Microplastic in the gastrointestinal tract of fishes along the Saudi Arabian Red Sea coast. Marine Pollution Bulletin, 131, 407-415. doi.org/10.1016/j.marpolbul.2018.04.040
- Barboza, L. G. A., Vieira, L. R., & Guilhermino, L., 2018. Single and combined effects of microplastics and mercury on juveniles of the European seabass (Dicentrarchus labrax): Changes in behavioural responses and reduction of swimming velocity and resistance time. Environmental Pollution, 236, 1014-1019. doi.org/10.1016/j.envpol.2017.12.082
- Batel, A., Linti, F., Scherer, M., Erdinger, L., & Braunbeck, T., 2016. Transfer of benzo [a] pyrene from microplastics to Artemia nauplii and further to zebrafish via a trophic food
- web experiment: CYP1A induction and visual tracking of persistent organic pollutants. Environmental Toxicology and Chemistry, 35(7), 1656-1666. doi.org/10.1002/etc.3361
- Bellas, J., Martínez-Armental, J., Martínez-Cámara, A., Besada, V., & Martínez-Gómez, C., 2016. Ingestion of microplastics by demersal fish from the Spanish Atlantic and Mediterranean coasts. Marine pollution bulletin, 109(1), 55-60. doi.org/10.1016/j.marpolbul.2016.06.026
- Bessa, F., Barría, P., Neto, J. M., Frias, J. P., Otero, V., Sobral, P., & Marques, J. C., 2018. Occurrence of microplastics in commercial fish from a natural estuarine environment. Marine pollution bulletin, 128, 575-584. doi.org/10.1016/j.marpolbul.2018.01.044
- Birnstiel, S., Soares-Gomes, A., & da Gama, B. A., 2019. Depuration reduces microplastic content in wild and farmed mussels. Marine pollution bulletin, 140, 241-247. doi:10.1016/j.marpolbul.2019.01.044
- Browne, M. A., Crump, P., Niven, S. J., Teuten, E., Tonkin, A., Galloway, T., & Thompson, R., 2011. Accumulation of Microplastic on Shorelines Woldwide: Sources and Sinks. Environmental Science & Technology, 45(21), 9175–9179. doi:10.1021/es201811s
- Budimir, S., Setälä, O., & Lehtiniemi, M., 2018. Effective and easy to use extraction method shows low numbers of microplastics in offshore planktivorous fish from the northern Baltic Sea. Marine pollution bulletin, 127, 586-592. doi.org/10.1016/j.marpolbul.2017.12.054
- Carpenter, K. E., & De Angelis, N., 2002. The living marine resources of the Western Central Atlantic (Vol. 2, pp. 602-1373). Rome: Food and agriculture organisation of the United Nations.
- Chagnon, C., Thiel, M., Antunes, J., Ferreira, J. L., Sobral, P., & Ory, N. C., 2018. Plastic ingestion and trophic transfer between Easter Island flying fish (Cheilopogon rapanouiensis) and yellowfin tuna (Thunnus albacares) from Rapa Nui (Easter Island). Environmental Pollution, 243, 127-133. doi.org/10.1016/j.envpol.2018.08.042
- Dantas, N. C., Duarte, O. S., Ferreira, W. C., Ayala, A. P., Rezende, C. F., & Feitosa, C. V.,
- 2020. Plastic intake does not depend on fish eating habits: Identification of microplastics in the stomach contents of fish on an urban beach in Brazil. Marine Pollution Bulletin,
- 153, 110959. doi.org/10.1016/j.marpolbul.2020.110959
- de Moura, A. R. L. U., Candeias, A. L. B., Limongi, C. M., 2009. A multi-temporal remote sensing and gis based inventory of the mangroves at itamaracá estuarine system, Northeastern Brazil. Tropical Oceanography, 37(1-2).
- de Sá, L. C., Luís, L. G., & Guilhermino, L., 2015. Effects of microplastics on juveniles of the common goby (Pomatoschistus microps): confusion with prey, reduction of the predatory performance and efficiency, and possible influence of developmental conditions. Environmental Pollution, 196, 359-362. doi.org/10.1016/j.envpol.2014.10.026
- Dauvergne, P., 2018. Why is the global governance of plastic failing the oceans?. Global Environmental Change, 51, 22-31. doi.org/10.1016/j.gloenvcha.2018.05.002
- Dunn, O. J., 1964. Multiple comparisons using rank sums. Technometrics, 6(3), 241-252.
- Elliott, M., Whitfield, A. K., Potter, I. C., Blaber, S. J., Cyrus, D. P., Nordlie, F. G., & Harrison,
- T. D., 2007. The guild approach to categorising estuarine fish assemblages: a global review. Fish and fisheries, 8(3), 241-268. doi:10.1111/j.1467-2679.2007.00253.x
- Eriksson, C., & Burton, H., 2003. Origins and biological accumulation of small plastic particles in fur seals from Macquarie Island. AMBIO: A Journal of the Human Environment, 32(6), 380-384. doi.org/10.1579/0044-7447-32.6.380
- Fendall, L.S., Sewell, M.A., 2009. Contributing to marine pollution by washing your face: Microplastics in facial cleansers. Mar. Pollut. Bull. 58, 1225–1228. doi.org/10.1016/j.marpolbul.2009.04.025
- Ferreira, G. V., Barletta, M., Lima, A. R., Dantas, D. V., Justino, A. K., & Costa, M. F., 2016. Plastic debris contamination in the life cycle of Acoupa weakfish (Cynoscion acoupa) in a tropical estuary. ICES Journal of Marine Science: Journal du Conseil, fsw108.
- doi.org/10.1093/icesjms/fsw108
- Ferreira, G. V., Barletta, M., Lima, A. R., Morley, S. A., Justino, A. K., & Costa, M. F., 2018. High intake rates of microplastics in a Western Atlantic predatory fish, and insights of a direct fishery effect. Environmental Pollution, 236, 706-717. doi.org/10.1016/j.envpol.2018.01.095
- Ferreira, G. V., Barletta, M., Lima, A. R., Morley, S. A., & Costa, M. F., 2019. Dynamics of marine debris ingestion by profitable fishes along the estuarine ecocline. Scientific reports, 9(1), 1-12. doi.org/10.1038/s41598-019-49992-3
- Ferreira, V., Loc'h, L., Ménard, F., Frédou, T., & Frédou, F. L., 2019. Composition of the fish fauna in a tropical estuary: the ecological guild approach. Scientia Marina, 83(2), 133- 142. doi: 10.3989/scimar.04855.25A
- Foekema, E. M., De Gruijter, C., Mergia, M. T., van Franeker, J. A., Murk, A. J., & Koelmans, A. A., 2013. Plastic in North Sea fish. Environmental science & technology, 47(15), 8818-8824. doi.org/10.1021/es400931b
- Freeman, L. A., Corbett, D. R., Fitzgerald, A. M., Lemley, D. A., Quigg, A., & Steppe, C. N., 2019. Impacts of urbanisation and development on estuarine ecosystems and water quality. Estuaries and Coasts, 42(7), 1821-1838. doi.org/10.1007/s12237-019-00597-z
- Frias, J. P. G. L., Sobral, P., & Ferreira, A. M., 2010. Organic pollutants in microplastics from two beaches of the Portuguese coast. Marine Pollution Bulletin, 60(11), 1988-1992. doi.org/10.1016/j.marpolbul.2010.07.030
- Galloway, T. S., Cole, M., & Lewis, C., 2017. Interactions of microplastic debris throughout the marine ecosystem. Nature Ecology & Evolution, 1(5). doi:10.1038/s41559-017-0116
- Garcés-Ordóñez, O., Mejía-Esquivia, K. A., Sierra-Labastidas, T., Patiño, A., Blandón, L. M., & Espinosa Díaz, L. F., 2020. Prevalence of microplastic contamination in the digestive tract of fishes from mangrove ecosystem in Cispata, Colombian Caribbean. Marine Pollution Bulletin, 154, 111085. doi:10.1016/j.marpolbul.2020.111085
- Guebert-Bartholo, F. M., Barletta, M., Costa, M. F., & Monteiro-Filho, E. L. A., 2011. Using gut contents to assess foraging patterns of juvenile green turtles Chelonia mydas in the Paranaguá Estuary, Brazil. Endangered Species Research, 13(2), 131-143. doi: 10.3354/esr00320
- Gonzalez, J. G., Ménard, F., Le Loc'h, F., de Andrade, H. A., Viana, A. P., Ferreira, V., ... & Frédou, T., 2019. Trophic resource partitioning of two snook fish species (Centropomidae) in tropical estuaries in Brazil as evidenced by stable isotope analysis. Estuarine, Coastal and Shelf Science, 226, 106287. doi.org/10.1016/j.ecss.2019.106287
- Hermsen, E., Mintenig, S. M., Besseling, E., & Koelmans, A. A., 2018. Quality criteria for the analysis of microplastic in biota samples: a critical review. Environmental science & technology, 52(18), 10230-10240. doi.org/10.1021/acs.est.8b01611
- Hermsen, E., Pompe, R., Besseling, E., & Koelmans, A. A., 2017. Detection of low numbers of microplastics in North Sea fish using strict quality assurance criteria. Marine pollution bulletin, 122(1-2), 253-258. doi.org/10.1016/j.marpolbul.2017.06.051
- Herrera, A., Ŝtindlová, A., Martínez, I., Rapp, J., Romero-Kutzner, V., Samper, M. D., Montoto, T., Aguiar-González, B., Packard, T., & Gómez, M., 2019. Microplastic ingestion by Atlantic chub mackerel (Scomber colias) in the Canary Islands coast. Marine pollution bulletin, 139, 127-135. doi.org/10.1016/j.marpolbul.2018.12.022
- Holmes, L. A., Turner, A., & Thompson, R. C., 2012. Adsorption of trace metals to plastic resin pellets in the marine environment. Environmental Pollution, 160, 42-48. doi.org/10.1016/j.envpol.2011.08.052
- Ivar do Sul, J. A., & Costa, M. F., 2013. The present and future of microplastic pollution in the marine environment. Environmental pollution, 185, 352-364. doi.org/10.1016/j.envpol.2013.10.036
- IBGE (Instituto Brasileiro de Geografia e Estatística) 2011. Estimativas da população nos municípios brasileiros em 1º de julho de 2011. Available in: http://www.ibge.gov.br/home/estatistica/populacao/estimativa2011/POP2011\_DOU.pdf
- Jambeck, J.R., Geyer, R., Wilcox, C., Siegler, T.R., Perryman, M., Andrady, A., Narayan, R., 2015. Plastic waste inputs from land into the ocean. Science 347 (6223), 768–771. doi:10.1126/science.1260352.
- Karami, A., Golieskardi, A., Choo, C. K., Romano, N., Ho, Y. B., & Salamatinia, B., 2017. A high-performance protocol for extraction of microplastics in fish. Science of the Total Environment, 578, 485-494. doi.org/10.1016/j.scitotenv.2016.10.213
- Krumme, U., Brenner, M. & Saint-Paul, U., 2008. Spring-neap cycle as a major driver of temporal variations in the feeding of intertidal fishes: evidence from the sea catfish Sciades herzbergii (Ariidae) of equatorial West Atlantic mangrove creeks. Journal of Experimental Marine Biology and Ecology 367, 91–99. doi.org/10.1016/j.jembe.2008.08.020
- Kühn, S., Van Werven, B., Van Oyen, A., Meijboom, A., Rebolledo, E. L. B., & Van Franeker, J. A., 2017. The use of potassium hydroxide (KOH) solution as a suitable approach to isolate plastics ingested by marine organisms. Marine pollution bulletin, 115(1-2), 86-90. doi.org/10.1016/j.marpolbul.2016.11.034
- Lima, A. R. A., Costa, M. F., & Barletta, M., 2014. Distribution patterns of microplastics within the plankton of a tropical estuary. Environmental Research, 132, 146-155. doi.org/10.1016/j.envres.2014.03.031
- Lima, A.R.A., Barletta, M., Costa, M.F., Ramos, J.A.A., Dantas, D.V., Justino, A.K.S., & Ferreira, G.V.B., 2016a. Changes in the composition of ichthyoplankton assemblage and plastic debris in mangrove creeks relative to moon phases. Journal of Fish Biology, 89, 619-640. doi.org/10.1111/jfb.12838
- Lima, A.R.A., Barletta, M., 2016b. Lunar influence on prey availability, diet shifts and niche overlap between Engraulidae larvae in tropical mangrove creeks. Journal of Fish Biology 89, 2133–2152. doi.org/10.1111/jfb.13121
- Lira, A. S., Frédou, F. L., Viana, A. P., Eduardo, L. N., Frédou, T., 2017. Feeding ecology of Centropomus undecimalis (Bloch, 1792) and Centropomus parallelus (Poey, 1860) in two tropical estuaries in Northeastern Brazil. Pan-American Journal of Aquatic Sciences, 12(2), 123-135.
- Lira, A., Angelini, R., Le Loc'h, F., Ménard, F., Lacerda, C., Frédou, T., & Frédou, F. L., 2018. Trophic flow structure of a neotropical estuary in northeastern Brazil and the comparison of ecosystem model indicators of estuaries. Journal of Marine Systems, 182, 31-45. doi.org/10.1016/j.jmarsys.2018.02.007
- Lebreton, L.C.M., van der Zwet, J., Damsteeg, J.-W., Slat, B., Andrady, A.L., Reisser, J., 2017. River plastic emissions to the world's oceans. Nat. Commun. 8, 15611. doi.org/10.1038/ncomms15611
- Lusher, A. L., Welden, N. A., Sobral, P., & Cole, M., 2017. Sampling, isolating, and identifying microplastics ingested by fish and invertebrates. Analytical Methods, 9(9), 1346-1360. doi:10.1039/C6AY02415G
- Markic, A., Gaertner, J. C., Gaertner-Mazouni, N., & Koelmans, A. A., 2019. Plastic ingestion by marine fish in the wild. Critical Reviews in Environmental Science and Technology, 50(7), 657-697. doi:10.1080/10643389.2019.1631990
- Menezes, N. A & J. L. Figueiredo., 2000. Manual de peixes marinhos do sudeste do Brasil. Teleostei (5). Museu de Zoologia, Universidade de São Paulo, p. 116.
- Medeiros, C., Kjerfve, B., Araujo, M., Neumann-Leitão, S., 2001. The itamaracá estuarine ecosystem, Brazil. In: Seeliger, U., Kjerfve, B. (Eds.), Coastal Marine Ecosystem of Latin America (Ecological Studies). Springer, New York, pp. 71–82.
- https://doi.org/10.1007/978-3-662-04482-7.
- Miller, M.E., Hamann, M., Kroon, F.J., 2020. Bioaccumulation and biomagnification of microplastics in marine organisms: A review and meta-analysis of current data. PLoS One. https://doi.org/10.1371/journal.pone.0240792
- Miranda, D. de A., & de Carvalho-Souza, G. F., 2016. Are we eating plastic-ingesting fish? Marine Pollution Bulletin, 103(1-2), 109–114. doi:10.1016/j.marpolbul.2015.12.035
- Moore, C. J., 2008. Synthetic polymers in the marine environment: a rapidly increasing, long-518 term threat. Environmental Research, 108(2), 131-139. doi.org/10.1016/j.envres.2008.07.025
- Morgana, S., Ghigliotti, L., Estévez-Calvar, N., Stifanese, R., Wieckzorek, A., Doyle, T., Garaventa, F., 2018. Microplastics in the Arctic: A case study with sub-surface water and fish samples off Northeast Greenland. Environmental Pollution, 242, 1078–1086. doi:10.1016/j.envpol.2018.08.001
- Nelms, S. E., Galloway, T. S., Godley, B. J., Jarvis, D. S., & Lindeque, P. K., 2018. Investigating microplastic trophic transfer in marine top predators. Environmental Pollution, 238, 999-1007. doi.org/10.1016/j.envpol.2018.02.016
- O'Connor, J. D., Mahon, A. M., Ramsperger, A. F. R. M., Trotter, B., Redondo-Hasselerharm, P. E., Koelmans, A. A., Murphy, S., 2019. Microplastics in Freshwater Biota: A Critical Review of Isolation, Characterisation, and Assessment Methods. Global Challenges,
- 1800118. doi:10.1002/gch2.201800118 Oehlmann, J., Schulte-Oehlmann, U., Kloas, W., Jagnytsch, O., Lutz, I., Kusk, K. O & Tyler, C. R., 2009. A critical analysis of the biological impacts of plasticisers on wildlife. Philosophical Transactions of the Royal Society of London B: Biological Sciences,
- 364(1526), 2047-2062. doi.org/10.1098/rstb.2008.0242
- Ostle, C., Thompson, R. C., Broughton, D., Gregory, L., Wootton, M., & Johns, D. G., 2019. The rise in ocean plastics evidenced from a 60-year time series. Nature communications, 10(1), 1-6. doi.org/10.1038/s41467-019-09506-1
- Ogata, Y., Takada, H., Mizukawa, K., Hiraia, H., Iwasaa, S., Endo, S., Mato, Y., Saha, M., Okuda, K., Nakashima, A., Murakami, M., Zurcher, N., Booyatumanondo, R., Zakaria, M.P., Dung, L.Q., Gordon, M., Miguez, C., Suzuki, S., Moore, C., Karapanagiotik, H.K., Weerts, S., McClurg, T., Burres, E., Smith, W., Van Velkenburg, M., Lang, J.S., Lang, R.C., Laursen, D., Danner, B., Stewardson, N., Thompson, R.C., 2009. International
- Pellet Watch: global monitoring of persistent organic pollutants (POPs) in coastal waters.
- 1. Initial phase data on PCBs, DDTs, and HCHs. Mar. Pollut. Bull. 58, 1437–1446. doi.org/10.1016/j.marpolbul.2009.06.014.
- Pegado, T. S. S., Schmid, K., Winemiller, K. O., Chelazzi, D., Cincinelli, A., Dei, L., & Giarrizzo, T., 2018. First evidence of microplastic ingestion by fishes from the Amazon River estuary. Marine pollution bulletin, 133, 814-821. doi.org/10.1016/j.marpolbul.2018.06.035
- Pelage, L., Domalain, G., Lira, A. S., Travassos, P., & Frédou, T., 2019. Coastal land use in Northeast Brazil: mangrove coverage evolution over three decades. Tropical Conservation Science, 12, 1940082918822411. doi.org/10.1177/1940082918822411
- Pellini, G., Gomiero, A., Fortibuoni, T., Ferrà, C., Grati, F., Tassetti, A. N., ... & Scarcella, G., 2018. Characterisation of microplastic litter in the gastrointestinal tract of Solea solea from the Adriatic Sea. Environmental pollution, 234, 943-952. doi.org/10.1016/j.envpol.2017.12.038
- Potter, I.C., Tweedley, J.R., Elliott, M., Whitfield, A.K., 2013. The ways in which fish use estuaries: a refinement and expansion of the guild approach. Fish and Fisheries 16, 230– 239. doi.org/10.1111/faf.12050
- R Core Team., 2020. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.
- Ramos, J.A.A., Barletta, M., Dantas, D.V., Costa, M.F., 2016. Seasonal and spatial ontogenetic movement of Gerreidae in a Brazilian tropical estuarine ecocline and its application for nursery habitat conservation. Journal of Fish Biology 89, 696–712. doi.org/10.1111/jfb.12872
- Ribeiro-Brasil, D. R. G., Torres, N. R., Picanço, A. B., Sousa, D. S., Ribeiro, V. S., Brasil, L. S., & de Assis Montag, L. F., 2020. Contamination of stream fish by plastic waste in the Brazilian Amazon. Environmental Pollution, 266, 115241. doi.org/10.1016/j.envpol.2020.115241
- Rochman, C. M., Hoh, E., Kurobe, T., & Teh, S. J., 2013. Ingested plastic transfers hazardous chemicals to fish and induces hepatic stress. Scientific Reports, 3. doi.org/10.1038/srep03263
- Santana, M. F. M., Ascer, L. G., Custódio, M. R., Moreira, F. T., & Turra, A., 2016. Microplastic contamination in natural mussel beds from a Brazilian urbanised coastal
- region: Rapid evaluation through bioassessment. Marine Pollution Bulletin, 106(1-2), 183–189. doi:10.1016/j.marpolbul.2016.02.074
- 577 Schirinzi, G. F., Pedà, C., Battaglia, P., Laface, F., Galli, M., Baini, M., ... Romeo, T., 2020. A new digestion approach for the extraction of microplastics from gastrointestinal tracts (GITs) of the common dolphinfish (Coryphaena hippurus) from the western Mediterranean Sea. Journal of Hazardous Materials, 122794. doi:10.1016/j.jhazmat.2020.122794
- Su, L., Nan, B., Hassell, K. L., Craig, N. J., & Pettigrove, V., 2019. Microplastics biomonitoring in Australian urban wetlands using a common noxious fish (Gambusia holbrooki). Chemosphere. doi:10.1016/j.chemosphere.2019.04.114
- Tanaka, K., & Takada, H., 2016. Microplastic fragments and microbeads in digestive tracts of planktivorous fish from urban coastal waters. Scientific reports, 6, 34351. doi: 10.1038/srep34351
- Teuten, E. L., Rowland, S. J., Galloway, T. S., & Thompson, R. C., 2007. Potential for plastics to transport hydrophobic contaminants. Environmental Science & Technology, 41(22), 7759-7764. doi.org/10.1021/es071737s
- Teuten, E. L., Saquing, J. M., Knappe, D. R., Barlaz, M. A., Jonsson, S., Björn, A., Rowland, S. J., Thompson, R. C., Galloway, T.S., Yamashita, R., Ochi, D., Watanuki, Y., Moore, C., Viet, P. H., Tana, S. T., Prudente, M., Boonyatumanond, R., Zakaria, M. P., Akkhavong, K., Ogata, Y., Hirai, H., Iwasa, S., Mizukawa, .K., Hagino, Y., Imamura, A., Saha, M., Takada, H., 2009. Transport and release of chemicals from plastics to the environment and to wildlife. Philosophical Transactions of the Royal Society B: Biological Sciences, 364(1526), 2027-2045. https://doi.org/10.1098/rstb.2008.0284
- Thompson, R. C., Olsen, Y., Mitchell, R. P., Davis, A., Rowland, S. J., John, A. W., McGonigle, D., & Russell, A. E., 2004. Lost at sea: where is all the plastic?. Science (Washington), 304(5672), 838.
- Torre, M., Digka, N., Anastasopoulou, A., Tsangaris, C., & Mytilineou, C., 2016. Anthropogenic microfibres pollution in marine biota. A new and simple methodology to minimise airborne contamination. Marine Pollution Bulletin, 113(1-2), 55–61. doi:10.1016/j.marpolbul.2016.07.050
- Vendel, A. L., Bessa, F., Alves, V. E. N., Amorim, A. L. A., Patrício, J., & Palma, A. R. T., 2017. Widespread microplastic ingestion by fish assemblages in tropical estuaries
- subjected to anthropogenic pressures. Marine Pollution Bulletin, 117(1-2), 448-455. doi.org/10.1016/j.marpolbul.2017.01.081.
- Wieczorek, A. M., Morrison, L., Croot, P. L., Allcock, A. L., MacLoughlin, E., Savard, O., ...
- & Doyle, T. K., 2018. Frequency of microplastics in mesopelagic fishes from the
- Northwest Atlantic. Frontiers in Marine Science, 5, 39.
- doi.org/10.3389/fmars.2018.00039
- Wright, S. L., Rowe, D., Thompson, R. C., & Galloway, T. S., 2013. Microplastic ingestion decreases energy reserves in marine worms. Current Biology, 23(23), R1031-R1033. doi.org/10.1016/j.cub.2013.10.068
- Zhang, H., 2017. Transport of microplastics in coastal seas. Estuarine, Coastal and Shelf
- Science, 199, 74-86. doi.org/10.1016/j.ecss.2017.09.032