



HAL
open science

Probing quantum entanglement from magnetic-sublevels populations: beyond spin squeezing inequalities

Guillem Müller-Rigat, Maciej Lewenstein, Irénée Frérot

► To cite this version:

Guillem Müller-Rigat, Maciej Lewenstein, Irénée Frérot. Probing quantum entanglement from magnetic-sublevels populations: beyond spin squeezing inequalities. 2022. hal-03622059v1

HAL Id: hal-03622059

<https://hal.science/hal-03622059v1>

Preprint submitted on 28 Mar 2022 (v1), last revised 7 Feb 2023 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Probing quantum entanglement from magnetic-sublevels populations: beyond spin squeezing inequalities

Guillem Müller-Rigat¹, Maciej Lewenstein^{1,2}, and Irénée Frérot³

¹ICFO - Institut de Ciències Fotoniques, The Barcelona Institute of Science and Technology, 08860 Castelldefels (Barcelona), Spain

²ICREA, Pg. Lluís Companys 23, 08010 Barcelona, Spain

³Univ Grenoble Alpes, CNRS, Grenoble INP, Institut Néel, 38000 Grenoble, France

Spin squeezing inequalities (SSI) represent a major tool to probe quantum entanglement among a collection of few-level atoms, and are based on collective spin measurements and their fluctuations. Yet, for atomic ensembles of spin- j atoms and ultracold spinor gases, many experiments can image the populations in all Zeeman sublevels $s = -j, -j + 1, \dots, j$, potentially revealing finer features of quantum entanglement not captured by SSI. Here we present a systematic approach which exploits Zeeman-sublevel population measurements in order to construct novel entanglement criteria, and illustrate our approach on ground states of spin-1 and spin-2 Bose-Einstein condensates. Beyond these specific examples, our approach allows one to infer, in a systematic manner, the optimal permutationally-invariant entanglement witness for any given set of collective measurements in an ensemble of d -level quantum systems.

1 Introduction

To prepare and detect quantum-entangled states has become a central goal for experimental many-body physics. On the one hand, it demonstrates the ability to probe regimes unthinkable within a classical framework, either at equilibrium or during the dynamics. On the other hand, it represents a crucial step towards using such systems as resources for quantum-enhanced applications, such as sensing [14, 26], or the simulation of quantum many-body problems [13].

Guillem Müller-Rigat: guillem.muller@icfo.eu

Irénée Frérot: irenee.ferot@neel.cnrs.fr

Here, our main focus are those many-body systems which are most easily probed by collective measurements, such as atomic ensembles and ultracold spinor gases. In such systems, entanglement can be famously revealed by so-called spin-squeezing inequalities (SSI), which involve measurements of first- and second-moments of collective spin operators [26, 32, 35]. SSI and their generalizations [32, 35] have found an impressively broad range of successful applications, from detecting entanglement in cold and hot atomic ensembles [19, 26] to unveiling many-body Bell non-locality [6, 25, 27, 30], from probing quantum phase transitions [9] and quantum quenches [4] to applications in quantum-enhanced metrology [26]. In particular, a handful of generalized SSI is sufficient to capture the complete palette of entanglement that can be revealed via collective-spin measurement in $j = 1/2$ spin ensembles [32]. In the context of $j = 1$ spinor Bose-Einstein condensates (BEC), by restricting the spin states to effective two-dimensional subspaces, entanglement has also been probed via those SSI [5, 15, 20, 21, 28]. In parallel, the last decade has witnessed the development of several experiments investigating much-larger- j spinor gases [1, 3, 7, 8, 11, 12, 15, 16, 20–24, 28, 33, 36, 37], mixtures [2, 17], or effective qudits ensembles [29]. In these systems, the populations of all Zeeman sublevels $s = -j, -j + 1, \dots, j$ can be measured, typically by fluorescence imaging after collective spin rotations and Stern-Gerlach splitting [7, 8, 12, 15, 20–22, 24, 28, 37]. The entanglement patterns incorporated by $j > 1/2$ spin ensembles is potentially much richer than what can be captured by SSI, with new classes of entangled states and new entanglement mechanisms. Yet, the optimal use of such collective imaging data to probe

entanglement beyond SSI has remained an open problem, for which our paper offers a novel approach.

In order to do so, we actually formulate and solve a more general problem: considering a set of single-atom observables, we assume that their average value and pair correlations, averaged over all permutations of the atoms, can be estimated. We then derive a single inequality based on these data to reveal entanglement. This inequality not only summarizes all known SSI [32, 34, 35], but also detects new forms of entanglement. Specializing then to Zeeman-sublevels population measurements in spinor gases, we apply this general approach to discover novel families of entanglement criteria, akin to but not captured by SSI. We illustrate our findings on representative states of a spin-1 BEC, as produced and characterized in several experiments [7, 8, 15, 28, 37]. Finally, we extend these findings to the largely unexplored case of a spin-2 BEC.

Framework. Our goal is to exclude the possibility to decompose $\hat{\rho}$, the many-body density operator of the system, as a statistical mixture of product states over individual atoms: $\hat{\rho} \neq \hat{\rho}_{\text{sep}} := \sum_{\lambda} p_{\lambda} \otimes_{i=1}^N \hat{\rho}_{\lambda}(i)$. Here, $\hat{\rho}_{\lambda}(i)$ is an arbitrary internal state (pure or mixed) for atom $i = 1, \dots, N$, and $p_{\lambda} > 0$ is an arbitrary probability distribution, defining a separable state $\hat{\rho}_{\text{sep}}$. Our starting point is to consider for each atom a set of K different observables $\{\hat{o}_k(i)\}_{k=1}^K$, some of which must be non-commuting in order to potentially detect entanglement. We then assume that data of the following form are available:

$$m_k = \frac{1}{N} \sum_{i=1}^N \langle \hat{o}_k(i) \rangle, \quad (1)$$

$$C_{kl} = \frac{1}{N(N-1)} \sum_{i \neq j} \langle \hat{o}_k(i) \hat{o}_l(j) \rangle, \quad (2)$$

where $\langle \dots \rangle = \text{Tr}(\hat{\rho} \dots)$ is an expectation value over many identically-prepared systems. In practice, as we illustrate below, the correlations C_{kl} might be available for only a subset of pairs (k, l) , something which can be accommodated within our approach. It is also central to our method to have a bound β s.t.

$$\frac{1}{N} \sum_{i=1}^N \sum_{k=1}^K \langle \hat{o}_k(i) \rangle^2 \leq \beta, \quad (3)$$

where $\langle \hat{o}_k(i) \rangle$ denotes the average value in an arbitrary internal state of atom i . Depending on the choice of observables $\hat{o}_k(i)$, the bound β might be computable analytically, but in general it may be found numerically, optimizing over the states of atom i . Our central result is the following inequality, valid for any separable state of the many-body system:

$$N \text{Tr}[P(C - \mathbf{m} \otimes \mathbf{m})] \geq \text{Tr}(C) - \beta, \quad (4)$$

where we introduced the notation $[\mathbf{m} \otimes \mathbf{m}]_{kl} = m_k m_l$, and P is any projector (namely a $K \times K$ symmetric matrix obeying $P^2 = P$). The detailed proof of Eq. (4) is given in Appendix A. The optimal choice to most strongly violate Eq. (4) is to project onto the subspace corresponding to all negative eigenvalues of the matrix $C - \mathbf{m} \otimes \mathbf{m}$. As a special case, Eq. (4) contains all generalized SSI of ref. [35] when choosing the spin in orthogonal directions x, y, z as local observables. As we shall see, when applied to suitable states of a spinor gas, it allows us to find novel entanglement criteria beyond SSI.

Zeeman-sublevels population measurements. In order to illustrate the relevance of our new approach to probe entanglement in a spin- j atomic ensemble, we assume that the populations $\hat{N}_s^{(a)}$ of all Zeeman sublevels $s = -j, \dots, j$ can be measured, for different quantization axes a (at least, for the purpose of entanglement detection, along two different orientations). This is typically achieved by first applying collective spin rotations to map a given orientation a onto the z axis, followed by a spatial splitting of the atoms in a magnetic-field gradient. The intensity of fluorescence imaging finally allows one to estimate the number of atoms in each Zeeman sublevel. For each orientation a , average populations $\langle \hat{N}_s^{(a)} \rangle$ and their correlations $\langle \hat{N}_s^{(a)} \hat{N}_{s'}^{(a)} \rangle$ can be inferred in this manner. In the general framework presented above, the single-atom observables correspond to the projectors onto the spin states $\hat{n}_s^{(a)}(i) = |s_a\rangle_i \langle s_a|_i$. Correspondingly, one reconstructs the average values $m_s^{(a)} = \langle \hat{N}_s^{(a)} \rangle / N$, and the diagonal blocks of the correlation matrix corresponding to a fixed orientation $C_{ss'}^{(a)} = \langle \hat{N}_s^{(a)} [\hat{N}_{s'}^{(a)} - \delta_{s,s'}] \rangle / [N(N-1)]$. The projector P in Eq. (4) is then chosen in a block-diagonal form, projecting onto the neg-

ative eigenspace of each block $C^{(a)} - \mathbf{m}^{(a)} \otimes \mathbf{m}^{(a)}$.

2 Applications to a spin-1 BEC

We first illustrate our approach on a BEC of spin-1 atoms, as investigated recently in several experiments [7, 8, 15, 28, 37]. We consider N spin-1 atoms with contact isotropic interactions in the single-mode approximation, adiabatically prepared from the so-called polar state $|0, N, 0\rangle = \otimes_{i=1}^N |0_z\rangle_i$ (with $|N_-, N_0, N_+\rangle$ the state written in the collective population basis along the z quantization axis). In an external magnetic field, the Hamiltonian reads (see Appendix B for details):

$$\hat{H}_{j=1} = \frac{c}{N} \hat{\mathbf{J}}^2 + q \hat{Q}^{(z)}, \quad (5)$$

where $\hat{J}^{(a)} = \sum_{s=-j}^j s \hat{N}_s^{(a)}$ is the collective spin in direction $a \in \{x, y, z\}$ and $\hat{\mathbf{J}}^2 = [\hat{J}^{(x)}]^2 + [\hat{J}^{(y)}]^2 + [\hat{J}^{(z)}]^2$; $\hat{Q}^{(z)} = \sum_{s=-j}^j s^2 \hat{N}_s^{(z)}$ is the collective quadrupole operator; and $\{x, y, z\}$ is an orthonormal basis of \mathbb{R}^3 . Notice that the linear Zeeman term, proportional to $\hat{J}^{(z)}$, is omitted as it commutes with $\hat{\mathbf{J}}^2$ and acts trivially on the initial state: $\hat{J}^{(z)}|0, N, 0\rangle = 0$. The initial state is also the ground state of $\hat{Q}^{(z)}$; and by varying the magnetic field intensity (q) one can adiabatically prepare the ground state of $\hat{H}_{j=1}$ for either ferromagnetic interactions ($c \leq 0$) or antiferromagnetic interactions ($c \geq 0$). Notice that in the case of ferromagnetic interactions, two quantum phase transitions occur at $q/|c| = \pm 4$, so that adiabaticity is possible only for finite N . As the Zeeman sublevel populations are related by $\sum_{s=-j}^j \hat{N}_s^{(a)} = N$ for all directions, $m_s^{(a)}$ and $C_{ss'}^{(a)}$ for all s, s' are not all independent. As local observables in Eqs. (1) and (2) we used the spin $\hat{S}^{(a)}(i) = \sum_{s=-j}^j s \hat{n}_s^{(a)}(i)$ and the projector onto the Zeeman sublevel $s = 0$, $\hat{n}_0^{(a)}(i)$. We use these data as input to Eq. (4), with bound $\beta_{j=1} = 3/2$ in Eq. (3) (see Appendix E). Notice that for $j = 1$, we have $\hat{Q}^{(a)} = \hat{N}_+^{(a)} + \hat{N}_-^{(a)} = N - \hat{N}_0^{(a)}$.

Antiferromagnetic interactions. For antiferromagnetic interactions ($c > 0$) and $q/c \lesssim 4.4$ ($N = 100$), we find violation of Eq. (4), with an




Figure 1: (a) Comparison of the population-based witness [Eq. (6)] (solid line) with the SSI $\text{Var}(\hat{\mathbf{J}}) \geq Nj$ (dashed line) for spin-1 states ($N = 100$ atoms). The light grey area are data points which cannot be achieved (see text and Appendix D). The dotted line indicates the ground state of Eq. (5) for both antiferromagnetic interactions ($c = 1$), varying the quadratic Zeeman term ($-10 \leq q \leq 10$). Three representative states are also indicated: the polar state $|0, N, 0\rangle$ ($q \rightarrow \infty$) which is separable, the spin singlet s.t. $\hat{\mathbf{J}}^2|\Psi\rangle = 0$ ($q = 0$), and the twin-Fock state $|N/2, 0, N/2\rangle$ ($q \rightarrow -\infty$). (b) Comparison of the population-based witness [Eq. (7)] with a related SSI (see text) for spin-1 states. Graphical conventions as in panel (a). Here, the ground state is computed for ferromagnetic interactions ($c = -1$). At $q = 0$, we obtain the Dicke state $\hat{\mathbf{J}}^2|\text{Dicke}\rangle = N(N+1)|\text{Dicke}\rangle$, $\hat{J}^{(z)}|\Psi\rangle = 0$.

entanglement witness (EW) which reads (see Appendix E):

$$\text{Var}(\hat{\mathbf{J}}) - \frac{1}{N-1} \langle \hat{\mathbf{N}}_0^2 \rangle \geq \frac{N(N-3)}{2(N-1)}, \quad (6)$$

with the notations $\text{Var}(\hat{\mathbf{J}}) = \text{Var}(\hat{J}^{(x)}) + \text{Var}(\hat{J}^{(y)}) + \text{Var}(\hat{J}^{(z)})$, $\text{Var}(\hat{O}) = \langle \hat{O}^2 \rangle - \langle \hat{O} \rangle^2$ and $\hat{\mathbf{N}}_0 = (\hat{N}_0^{(x)}, \hat{N}_0^{(y)}, \hat{N}_0^{(z)})$. The EW of Eq. (6) is maximally violated at $q = 0$, where the ground state of Eq. (5) is a spin singlet ($\text{Var}(\hat{\mathbf{J}}) = 0$). In particular, we also applied our method using as input the experimental data of ref. [8], where an almost perfect spin singlet is prepared, and all populations are measured; we also recovered Eq. (6) as the optimal violated EW. This EW is reminiscent of, yet generally more powerful than, the generalized SSI $\text{Var}(\hat{\mathbf{J}}) \geq Nj$ (with $j = 1$ here) derived in ref. [35]. For a quantitative comparison of both criteria, in Fig. 1(a) we illustrate their detection power for $N = 100$ and states with $\langle \hat{\mathbf{J}} \rangle = \mathbf{0}$ (vanishing mean spin, as is the case in the ground state of $\hat{H}_{j=1}$, Eq. (5)) in the plane ($x = \langle \hat{\mathbf{J}}^2 \rangle / N, y = \langle \hat{\mathbf{N}}_0^2 \rangle / [N(N-1)]$), where violation occurs respectively for $x - y < (N-3)/[2(N-1)]$ [Eq. (6)] and $x < 1$ (SSI). We plot the data points corresponding to the ground state of $\hat{H}_{j=1}$ for $c > 0$ and both $q > 0$ and $q < 0$. More generally, not all combinations of $\langle \hat{\mathbf{J}}^2 \rangle$ and $\langle \hat{\mathbf{N}}_0^2 \rangle$ are physically allowed. As detailed in Appendix D (see also [31]), the physically-allowed region can be obtained by considering all quantum ground states of Hamiltonians of the form $\lambda_1 \hat{\mathbf{J}}^2 + \lambda_2 \hat{\mathbf{N}}_0^2$ for all values of λ_1, λ_2 , and forming the convex envelope of the corresponding data points. The excluded (non-feasible) region is indicated in light grey in Fig. 1(a). Regarding the ground state of $\hat{H}_{j=1}$, three limiting cases are especially illustrative. For $q \rightarrow +\infty$, the ground state is the polar state $|0, N, 0\rangle = \otimes_{i=1}^N |0_z\rangle_i$, which is separable and therefore does not violate any EW. For $q = 0$, the ground state is a spin singlet, and violates both our witness Eq. (6) and the SSI $\text{Var}(\hat{\mathbf{J}}) \geq Nj$. In-between, we notice that our EW detects entanglement in the ground state of $\hat{H}_{j=1}$ for a larger range of values of q than SSIs. Finally, for $q \rightarrow -\infty$, the ground state is a twin Fock state $|N/2, 0, N/2\rangle$. This state does not violate any EW if only collective spin observables are measured (notice, though, that it does violate a generalized SSI

for $j = 1/2$ particles, when considering only the two-dimensional subspace spanned by Zeeman sublevels $s = \pm 1$). Yet, the twin-Fock states robustly violates the EW of Eq. (6) (with $\text{Var}(\hat{\mathbf{J}}) = N$ and $\langle \hat{\mathbf{N}}_0^2 \rangle = N(3N+2)/4$, so that the l.h.s is $\sim N/4$, while the r.h.s is $\sim N/2$). The polar, singlet and twin-Fock states are indicated on Fig. 1(a).

Ferromagnetic interactions. For ferromagnetic interactions ($c < 0$) and $q/|c| \lesssim 1.2$ ($N = 100$), we find violation of (see Appendix E):

$$(N-1)\text{Var}(\hat{J}^{(z)}) - N\langle \hat{Q}^{(z)} \rangle - \langle [\hat{J}^{(x)}]^2 + [\hat{J}^{(y)}]^2 \rangle + \frac{3N(N+1)}{2} \geq \langle \hat{\mathbf{N}}_0^2 \rangle. \quad (7)$$

Maximal violation is found close to $q = 0$ where the state is a Dicke state, such that $\hat{\mathbf{J}}^2|\text{Dicke}\rangle = Nj(Nj+1)|\text{Dicke}\rangle$ (maximal total spin), and $\hat{J}^{(z)}|\text{Dicke}\rangle = 0$. Eq. (7) can be compared to another SSI of ref. [35] violated by the Dicke state, namely $(N-1)\text{Var}(\hat{J}^{(z)}) - N\langle \hat{Q}^{(z)} \rangle - \langle [\hat{J}^{(x)}]^2 + [\hat{J}^{(y)}]^2 \rangle + Nj(Nj+1) \geq 0$. In Fig. 1(b), we compare their power in detecting entanglement in the ground state of $\hat{H}_{j=1}$. In particular, for $q \lesssim |c|$, the ground state of $\hat{H}_{j=1}$ is very close to the twin-Fock state $|N/2, 0, N/2\rangle$, which robustly violates Eq. (7): the l.h.s is equal to $N(N+1)/2 \sim N^2/2$, while on the r.h.s $\langle \hat{\mathbf{N}}_0^2 \rangle = N(3N+2)/4 \sim 4N^2/4$. To conclude on the spin-1 BEC, we find that collective population measurements in orthogonal Zeeman sublevels can reveal entanglement beyond collective spin observables, and our method can be used to infer novel EWs in an optimal and data-agnostic way from these measurements. These new EWs are generically more powerful than SSIs (which can be recovered as a special case in our approach). In particular, we exhibit two EWs [Eqs. (6) and (7)] robustly violated by the twin-Fock state $|N/2, 0, N/2\rangle$, stabilized as the ground state of $\hat{H}_{j=1}$ in the limit $q \ll |c|$; yet no generalized SSI for spin-1 ensembles is violated by the twin-Fock state.

3 Applications to a spin-2 BEC

The Hamiltonian of a BEC of $j = 2$ atoms interacting via pairwise isotropic interactions reads




Figure 2: Comparison of the population-based entanglement witness of Eq. (9) (solid line) with the related SSI $\text{Var}(\hat{\mathbf{J}}) \geq Nj$ (dashed line) for spin-2 ensembles ($N = 24$). The light grey area are data points which cannot be achieved (see text and Appendix D). Data in the ground state of Eq. (8) are computed for $c = 1$, and $p = 1$ (dotted line) or $p = -1$ (dashed-dotted line). Four representative states are indicated: the polar state $|0, 0, N, 0, 0\rangle$ (with $\langle \hat{\mathbf{J}}^2 / (2N) \rangle = 3$), the twin-Fock state $|N/2, 0, 0, 0, N/2\rangle$ and two forms of spin singlets ($\text{Var}(\hat{\mathbf{J}}) = 0$), obtained either as a pair condensate or a triplet condensate (see text).

(see Appendix B):

$$\hat{H}_{j=2} = \frac{c}{N} \hat{\mathbf{J}}^2 + \frac{p}{N} \hat{\Theta}_2^\dagger \hat{\Theta}_2 + q \hat{Q}^{(z)}. \quad (8)$$

In contrast to the spin-1 case [Eq. (5)], the Hamiltonian contains the additional pairing term $\hat{\Theta}_2^\dagger \hat{\Theta}_2$ where $\hat{\Theta}_2 = \sum_{s=-j}^j (-1)^s \hat{a}_s \hat{a}_{-s}$ annihilates time-reversal pairs $\pm s$, with \hat{a}_s annihilating an atom in the spin state $|s_z\rangle$. Notice that $[\hat{\mathbf{J}}^2, \hat{\Theta}_2^\dagger \hat{\Theta}_2] = 0$. We explored the ground state of $\hat{H}_{j=2}$ to find novel violated EW. Using as local operators $\hat{S}^{(a)}(i)$ (the spin) and $\hat{n}_0^{(a)}(i)$ (the projector onto the $s = 0$ spin state) for $a \in \{x, y, z\}$, the bound in Eq. (3) is numerically found as $\beta_2 \approx 4.3$. For this choice of local operators, we found that generalizations of Eqs. (6) and (7) are violated. These EW can be further generalized to arbitrary integer spins, and read (see Appendix E):

$$\text{Var}(\hat{\mathbf{J}}) - \frac{1}{N-1} \langle \hat{\mathbf{N}}_0(\hat{\mathbf{N}}_0 - \mathbf{1}) \rangle \geq N[j(j+1) - \beta_j], \quad (9)$$

and:

$$(N-1)\text{Var}(\hat{J}^{(z)}) - N\langle \hat{Q}^{(z)} \rangle - \langle [\hat{J}^{(x)}]^2 + [\hat{J}^{(y)}]^2 \rangle + N[(N-1)\beta_j + j(j+1)] \geq \langle \hat{\mathbf{N}}_0(\hat{\mathbf{N}}_0 - \mathbf{1}) \rangle, \quad (10)$$

where $\mathbf{1} = (1, 1, 1)$ and β_j is the corresponding bound in Eq. (3). As for the spin-1 case, one can compare the entanglement detection power of these witnesses with the related SSIs; this comparison is presented in Fig. 2 for Eq. (9), while for Eq. (10) we notice that the improvement with respect to the SSI is mild, and not systematic. Interestingly, as in the case of spin-1, the twin-Fock state $|N/2, 0, 0, 0, N/2\rangle$ (obtained in the limit $q \rightarrow -\infty$) violates Eq. (9), while it does not violate any generalized SSI for spin-2 collective spin observables. As a final exploration of the entanglement patterns in the ground state of a spin-2 BEC, we have focused on many-body singlets ($\langle \hat{\mathbf{J}}^2 \rangle = 0$), stabilized at $q = 0$ for antiferromagnetic interactions ($c > 0$). In the spin-1 case, there is a unique such state for a BEC, obtained as a “pair condensate” ($\hat{\Theta}_2^\dagger$) $^{N/2} |\text{vac.}\rangle$. This pair condensate is also stabilized in the spin-2 case for $p < 0$. Yet, for $p > 0$, the many-body singlet is instead a “triplet condensate” ($\hat{\Theta}_3^\dagger$) $^{N/3} |\text{vac.}\rangle$, where $\hat{\Theta}_3^\dagger = -9[(a_{-1}^\dagger)^2 a_2^\dagger + (a_1^\dagger)^2 a_{-2}^\dagger] + \sqrt{6} a_0^\dagger [-(a_0^\dagger)^2 + 3a_1^\dagger a_{-1}^\dagger + 6a_2^\dagger a_{-2}^\dagger]$ is the singlet-trio creation operator. Both states are entangled and violate Eq. (9). However, considering as local measurements $\{\hat{n}_{+2}^{(a)}(i), \hat{n}_{+1}^{(a)}(i), \hat{n}_{-1}^{(a)}(i), \hat{n}_{-2}^{(a)}(i)\}_{a \in \{x, y, z\}}$, we found the following EW violated only by the pair condensate and not by the triplet condensate (see Appendix F):

$$\langle \hat{\sigma}_2^2 + \hat{\sigma}_1^2 \rangle - \frac{N-1}{N} (\langle \hat{\sigma}_2 \rangle^2 + \langle \hat{\sigma}_1 \rangle^2) - \frac{2}{N} \langle \hat{\mathbf{N}}_{+2}^2 + \hat{\mathbf{N}}_{+1}^2 + \hat{\mathbf{N}}_{-1}^2 + \hat{\mathbf{N}}_{-2}^2 \rangle + (1 - \frac{2}{N}) \langle \hat{\mathbf{N}}_0 \cdot \mathbf{1} \rangle + 3 \geq 0, \quad (11)$$

where we denoted by $\hat{\sigma}_s^{(a)} = \hat{N}_s^{(a)} - \hat{N}_{-s}^{(a)}$ the population imbalance between opposed levels along direction a . Notice that $\{\hat{\sigma}_s^{(x)}, \hat{\sigma}_s^{(y)}, \hat{\sigma}_s^{(z)}\}$ generate an $su(2)$ subalgebra of $su(5)$. This result illustrates the difference in entanglement properties of two different spin singlets, which by definition cannot be distinguished if only collective spin observables are measured.

4 Conclusions

We have presented a new method to infer violated entanglement witnesses from the collective measurement of Zeeman-sublevel populations in spinor gases of arbitrary spin- j atoms. Our method recovers all known generalized spin squeezing inequalities [35] as a special case, in a data-agnostic way. But when considering appropriate correlations between Zeeman-sublevel populations, one can also construct novel entanglement criteria which have no analog if only the collective spin is measured. We have illustrated our method on the ground states of spin-1 and spin-2 BECs, interacting with contact two-body isotropic interactions in an external magnetic field. In particular, for both $j = 1$ and $j = 2$ we showed that the twin Fock state robustly violates an entanglement witness involving all populations in three orthogonal directions, while entanglement cannot be detected if only the collective spin is measured. For $j = 2$, we presented an entanglement witness distinguishing between two forms of many-body singlets, namely a pair condensate and a triplet condensate. Overall, our new approach appears especially suited to probe multipartite entanglement in high-spin spinor gases, for which entanglement detection has remained a considerable challenge. We would like to emphasize that, although we have focused on collective measurements which are invariant under the permutations of the atoms, our approach is not limited to this situation. It could also be used to probe entanglement in spatially-structured systems by using different local observables $\hat{o}_k(i)$ for each atom; a straightforward example is to introduce local phases $e^{i\phi_k(i)}$, leaving the bound β invariant in Eq. (3). As pointed out recently [10], this offers considerable flexibility to probe entanglement through components of the structure factors. Finally, a limitation of our current implementation is that there is no optimal and systematic way of incorporating all available first- and second-order moments of population measurements into our main inequality (4). Indeed, while adding more measurement operators *a priori* improves the entanglement detection power, this leads to an increase of the β bound in Eq. (3), and can make Eq. (4) harder to violate. The optimization of the choice of measurement operators is left open to future studies.

Code availability. The code used to diagonalize the spin-1 and spin-2 Hamiltonians, to compute the collective populations in the ground state, and to infer the optimal entanglement witnesses presented in this paper, is available at https://github.com/GuillemMRR/EW_Zeeman.

Acknowledgments

GM and ML acknowledges support from: ERC AdG NOQIA; Agencia Estatal de Investigación (R & D project CEX2019-000910-S, funded by MCIN/ AEI/10.13039/501100011033, Plan National FIDEUA PID2019-106901GB-I00, FPI, QUANTERA MAQS PCI2019-111828-2, Proyectos de I+D+I “Retos Colaboración” QUSPIN RTC2019-007196-7); Fundació Cellex; Fundació Mir-Puig; Generalitat de Catalunya through the European Social Fund FEDER and CERCA program (AGAUR Grant No. 2017 SGR 134, QuantumCAT U16-011424, co-funded by ERDF Operational Program of Catalonia 2014-2020); EU Horizon 2020 FET-OPEN OPTologic (Grant No 899794); National Science Centre, Poland (Symfonia Grant No. 2016/20/W/ST4/00314); European Union’s Horizon 2020 research and innovation programme under the Marie-Skłodowska-Curie grant agreement No 101029393 (STREDCH) and No 847648 (“La Caixa” Junior Leaders fellowships ID100010434: LCF/BQ/PI19/11690013, LCF/BQ/PI20/11760031, LCF/BQ/PR20/11770012, LCF/BQ/PR21/11840013). IF acknowledges support from the Agence Nationale de la Recherche (Qu-DICE project ANR-PRC-CES47) and the John Templeton Foundation (Grant No. 61835).

References

- [1] P. Bataille, A. Litvinov, I. Manai, J. Huckans, F. Wiotte, A. Kaladjian, O. Gorceix, E. Maréchal, B. Laburthe-Tolra, and M. Robert-de Saint-Vincent. Adiabatic spin-dependent momentum transfer in an $su(n)$ degenerate fermi gas. *Phys. Rev. A*, 102:013317, Jul 2020. DOI: [10.1103/PhysRevA.102.013317](https://doi.org/10.1103/PhysRevA.102.013317). URL <https://link.aps.org/doi/10.1103/PhysRevA.102.013317>.

- [2] Rohit Prasad Bhatt, Jan Kilinc, Lilo Höcker, and Fred Jendrzejewski. Stochastic dynamics of a few sodium atoms in presence of a cold potassium cloud. *Scientific Reports*, 12(1), Feb 2022. ISSN 2045-2322. DOI: [10.1038/s41598-022-05778-8](https://doi.org/10.1038/s41598-022-05778-8). URL <http://dx.doi.org/10.1038/s41598-022-05778-8>.
- [3] L. Chomaz, S. Baier, D. Petter, M. J. Mark, F. Wächtler, L. Santos, and F. Ferlaino. Quantum-fluctuation-driven crossover from a dilute bose-einstein condensate to a macrodroplet in a dipolar quantum fluid. *Phys. Rev. X*, 6:041039, Nov 2016. DOI: [10.1103/PhysRevX.6.041039](https://doi.org/10.1103/PhysRevX.6.041039). URL <https://link.aps.org/doi/10.1103/PhysRevX.6.041039>.
- [4] Tommaso Comparin, Fabio Mezzacapo, and Tommaso Roscilde. Robust spin squeezing from the tower of states of $u(1)$ -symmetric spin hamiltonians. *Phys. Rev. A*, 105:022625, Feb 2022. DOI: [10.1103/PhysRevA.105.022625](https://doi.org/10.1103/PhysRevA.105.022625). URL <https://link.aps.org/doi/10.1103/PhysRevA.105.022625>.
- [5] L.-M. Duan, J. I. Cirac, and P. Zoller. Quantum entanglement in spinor bose-einstein condensates. *Phys. Rev. A*, 65:033619, Feb 2002. DOI: [10.1103/PhysRevA.65.033619](https://doi.org/10.1103/PhysRevA.65.033619). URL <https://link.aps.org/doi/10.1103/PhysRevA.65.033619>.
- [6] Nils J. Engelsens, Rajiv Krishnakumar, Onur Hosten, and Mark A. Kasevich. Bell Correlations in Spin-Squeezed States of 500 000 Atoms. *Phys. Rev. Lett.*, 118(14):140401, April 2017. DOI: [10.1103/PhysRevLett.118.140401](https://doi.org/10.1103/PhysRevLett.118.140401). URL <https://link.aps.org/doi/10.1103/PhysRevLett.118.140401>.
- [7] Bertrand Evrard, An Qu, Jean Dalibard, and Fabrice Gerbier. From many-body oscillations to thermalization in an isolated spinor gas. *Phys. Rev. Lett.*, 126:063401, Feb 2021. DOI: [10.1103/PhysRevLett.126.063401](https://doi.org/10.1103/PhysRevLett.126.063401). URL <https://link.aps.org/doi/10.1103/PhysRevLett.126.063401>.
- [8] Bertrand Evrard, An Qu, Jean Dalibard, and Fabrice Gerbier. Observation of fragmentation of a spinor bose-einstein condensate. *Science*, 373(6561):1340–1343, 2021. DOI: [10.1126/science.abd8206](https://doi.org/10.1126/science.abd8206). URL <https://www.science.org/doi/abs/10.1126/science.abd8206>.
- [9] Irénée Frérot and Tommaso Roscilde. Quantum critical metrology. *Phys. Rev. Lett.*, 121:020402, Jul 2018. DOI: [10.1103/PhysRevLett.121.020402](https://doi.org/10.1103/PhysRevLett.121.020402). URL <https://link.aps.org/doi/10.1103/PhysRevLett.121.020402>.
- [10] Irénée Frérot, Flavio Baccari, and Antonio Acín. Unveiling quantum entanglement in many-body systems from partial information. *arXiv preprint arXiv:2107.03944*, 2021. DOI: <https://doi.org/10.48550/arXiv.2107.03944>. URL <https://arxiv.org/abs/2107.03944>.
- [11] Albert Frisch, Michael Mark, Kiyotaka Aikawa, Francesca Ferlaino, John L. Bohn, Constantinos Makrides, Alexander Petrov, and Svetlana Kotochigova. Quantum chaos in ultracold collisions of gas-phase erbium atoms. *Nature*, 507(7493):475–479, Mar 2014. ISSN 1476-4687. DOI: [10.1038/nature13137](https://doi.org/10.1038/nature13137). URL <http://dx.doi.org/10.1038/nature13137>.
- [12] Lucas Gabardos, Bihui Zhu, Steven Lepoutre, Ana Maria Rey, Bruno Laburthe-Tolra, and Laurent Vernac. Relaxation of the collective magnetization of a dense 3d array of interacting dipolar $s = 3$ atoms. *Phys. Rev. Lett.*, 125:143401, Sep 2020. DOI: [10.1103/PhysRevLett.125.143401](https://doi.org/10.1103/PhysRevLett.125.143401). URL <https://link.aps.org/doi/10.1103/PhysRevLett.125.143401>.
- [13] I. M. Georgescu, S. Ashhab, and Franco Nori. Quantum simulation. *Rev. Mod. Phys.*, 86:153–185, Mar 2014. DOI: [10.1103/RevModPhys.86.153](https://doi.org/10.1103/RevModPhys.86.153). URL <https://link.aps.org/doi/10.1103/RevModPhys.86.153>.
- [14] Kevin A. Gilmore, Matthew Affolter, Robert J. Lewis-Swan, Diego Barberena, Elena Jordan, Ana Maria Rey, and John J. Bollinger. Quantum-enhanced sensing of displacements and electric fields with two-dimensional trapped-ion crystals. *Science*, 373(6555):673–678, Aug 2021. ISSN 1095-9203. DOI: [10.1126/science.abi5226](https://doi.org/10.1126/science.abi5226). URL <http://dx.doi.org/10.1126/science.abi5226>.

- [15] C. D. Hamley, C. S. Gerving, T. M. Hoang, E. M. Bookjans, and M. S. Chapman. Spin-nematic squeezed vacuum in a quantum gas. *Nature Physics*, 8(4):305–308, Feb 2012. ISSN 1745-2481. DOI: [10.1038/nphys2245](https://doi.org/10.1038/nphys2245). URL <http://dx.doi.org/10.1038/nphys2245>.
- [16] Holger Kadau, Matthias Schmitt, Matthias Wenzel, Clarissa Wink, Thomas Maier, Igor Ferrier-Barbut, and Tilman Pfau. Observing the rosenweig instability of a quantum ferrofluid. *Nature*, 530(7589):194–197, Feb 2016. ISSN 1476-4687. DOI: [10.1038/nature16485](https://doi.org/10.1038/nature16485). URL <http://dx.doi.org/10.1038/nature16485>.
- [17] Valentin Kasper, Daniel González-Cuadra, Apoorva Hegde, Andy Xia, Alexandre Dauphin, Felix Huber, Eberhard Tiemann, Maciej Lewenstein, Fred Jendrzejewski, and Philipp Hauke. Universal quantum computation and quantum error correction with ultracold atomic mixtures. *Quantum Science and Technology*, 7(1):015008, 2021.
- [18] Yuki Kawaguchi and Masahito Ueda. Spinor bose–einstein condensates. *Physics Reports*, 520(5):253–381, 2012. ISSN 0370-1573. DOI: <https://doi.org/10.1016/j.physrep.2012.07.005>. URL <https://www.sciencedirect.com/science/article/pii/S0370157312002098>. Spinor Bose–Einstein condensates.
- [19] Jia Kong, Ricardo Jiménez-Martínez, Charikleia Troullinou, Vito Giovanni Lucivero, Géza Tóth, and Morgan W. Mitchell. Measurement-induced, spatially-extended entanglement in a hot, strongly-interacting atomic system. *Nature Communications*, 11(1), May 2020. ISSN 2041-1723. DOI: [10.1038/s41467-020-15899-1](https://doi.org/10.1038/s41467-020-15899-1). URL <http://dx.doi.org/10.1038/s41467-020-15899-1>.
- [20] Philipp Kunkel, Maximilian Prüfer, Helmut Strobel, Daniel Linnemann, Anika Frölian, Thomas Gasenzer, Martin Gärttner, and Markus K. Oberthaler. Spatially distributed multipartite entanglement enables epr steering of atomic clouds. *Science*, 360(6387):413–416, 2018. DOI: [10.1126/science.aao2254](https://doi.org/10.1126/science.aao2254). URL <https://www.science.org/doi/abs/10.1126/science.aao2254>.
- [21] Philipp Kunkel, Maximilian Prüfer, Stefan Lannig, Rodrigo Rosa-Medina, Alexis Bonnin, Martin Gärttner, Helmut Strobel, and Markus K. Oberthaler. Simultaneous read-out of noncommuting collective spin observables beyond the standard quantum limit. *Phys. Rev. Lett.*, 123:063603, Aug 2019. DOI: [10.1103/PhysRevLett.123.063603](https://doi.org/10.1103/PhysRevLett.123.063603). URL <https://link.aps.org/doi/10.1103/PhysRevLett.123.063603>.
- [22] Karsten Lange, Jan Peise, Bernd Lücke, Ilka Kruse, Giuseppe Vitagliano, Iagoba Apellaniz, Matthias Kleinmann, Géza Tóth, and Carsten Klempt. Entanglement between two spatially separated atomic modes. *Science*, 360(6387):416–418, 2018. DOI: [10.1126/science.aao2035](https://doi.org/10.1126/science.aao2035). URL <https://www.science.org/doi/abs/10.1126/science.aao2035>.
- [23] E. Lucioni, L. Tanzi, A. Fregosi, J. Catani, S. Gozzini, M. Inguscio, A. Fioretti, C. Gabbanini, and G. Modugno. Dysprosium dipolar bose-einstein condensate with broad feshbach resonances. *Phys. Rev. A*, 97:060701, Jun 2018. DOI: [10.1103/PhysRevA.97.060701](https://doi.org/10.1103/PhysRevA.97.060701). URL <https://link.aps.org/doi/10.1103/PhysRevA.97.060701>.
- [24] Xin-Yu Luo, Yi-Quan Zou, Ling-Na Wu, Qi Liu, Ming-Fei Han, Meng Khoon Tey, and Li You. Deterministic entanglement generation from driving through quantum phase transitions. *Science*, 355(6325):620–623, 2017. DOI: [10.1126/science.aag1106](https://doi.org/10.1126/science.aag1106). URL <https://www.science.org/doi/abs/10.1126/science.aag1106>.
- [25] Guillem Müller-Rigat, Albert Aloy, Maciej Lewenstein, and Irénée Frérot. Inferring nonlinear many-body bell inequalities from average two-body correlations: Systematic approach for arbitrary spin- j ensembles. *PRX Quantum*, 2:030329, Aug 2021. DOI: [10.1103/PRXQuantum.2.030329](https://doi.org/10.1103/PRXQuantum.2.030329). URL <https://link.aps.org/doi/10.1103/PRXQuantum.2.030329>.
- [26] Luca Pezzè, Augusto Smerzi, Markus K. Oberthaler, Roman Schmied, and Philipp Treutlein. Quantum metrology with nonclassical states of atomic ensembles. *Rev. Mod. Phys.*, 90:035005, Sep 2018. DOI: [10.1103/RevModPhys.90.035005](https://doi.org/10.1103/RevModPhys.90.035005). URL <https://link.aps.org/doi/10.1103/RevModPhys.90.035005>.
- [27] Angelo Piga, Albert Aloy, Maciej Lewen-

- stein, and Irénée Frérot. Bell correlations at ising quantum critical points. *Phys. Rev. Lett.*, 123:170604, Oct 2019. DOI: [10.1103/PhysRevLett.123.170604](https://doi.org/10.1103/PhysRevLett.123.170604). URL <https://link.aps.org/doi/10.1103/PhysRevLett.123.170604>.
- [28] An Qu, Bertrand Evrard, Jean Dalibard, and Fabrice Gerbier. Probing spin correlations in a bose-einstein condensate near the single-atom level. *Phys. Rev. Lett.*, 125:033401, Jul 2020. DOI: [10.1103/PhysRevLett.125.033401](https://doi.org/10.1103/PhysRevLett.125.033401). URL <https://link.aps.org/doi/10.1103/PhysRevLett.125.033401>.
- [29] Martin Ringbauer, Michael Meth, Lukas Postler, Roman Stricker, Rainer Blatt, Philipp Schindler, and Thomas Monz. A universal qudit quantum processor with trapped ions, 2021.
- [30] Roman Schmied, Jean-Daniel Bancal, Baptiste Allard, Matteo Fadel, Valerio Scarani, Philipp Treutlein, and Nicolas Sangouard. Bell correlations in a Bose-Einstein condensate. *Science*, 352(6284): 441–444, April 2016. ISSN 0036-8075, 1095-9203. DOI: [10.1126/science.aad8665](https://doi.org/10.1126/science.aad8665). URL <http://science.sciencemag.org/content/352/6284/441>.
- [31] Anders S. Sørensen and Klaus Mølmer. Entanglement and extreme spin squeezing. *Phys. Rev. Lett.*, 86:4431–4434, May 2001. DOI: [10.1103/PhysRevLett.86.4431](https://doi.org/10.1103/PhysRevLett.86.4431). URL <https://link.aps.org/doi/10.1103/PhysRevLett.86.4431>.
- [32] Géza Tóth, Christian Knapp, Otfried Gühne, and Hans J. Briegel. Spin squeezing and entanglement. *Phys. Rev. A*, 79: 042334, Apr 2009. DOI: [10.1103/PhysRevA.79.042334](https://doi.org/10.1103/PhysRevA.79.042334). URL <https://link.aps.org/doi/10.1103/PhysRevA.79.042334>.
- [33] A. Trautmann, P. Ilzhöfer, G. Durastante, C. Politi, M. Sohmen, M. J. Mark, and F. Ferlaino. Dipolar quantum mixtures of erbium and dysprosium atoms. *Phys. Rev. Lett.*, 121:213601, Nov 2018. DOI: [10.1103/PhysRevLett.121.213601](https://doi.org/10.1103/PhysRevLett.121.213601). URL <https://link.aps.org/doi/10.1103/PhysRevLett.121.213601>.
- [34] Giuseppe Vitagliano, Philipp Hyllus, Iñigo L. Egusquiza, and Géza Tóth. Spin squeezing inequalities for arbitrary spin. *Phys. Rev. Lett.*, 107:240502, Dec 2011. DOI: [10.1103/PhysRevLett.107.240502](https://doi.org/10.1103/PhysRevLett.107.240502). URL <https://link.aps.org/doi/10.1103/PhysRevLett.107.240502>.
- [35] Giuseppe Vitagliano, Iagoba Apellaniz, Iñigo L. Egusquiza, and Géza Tóth. Spin squeezing and entanglement for an arbitrary spin. *Phys. Rev. A*, 89: 032307, Mar 2014. DOI: [10.1103/PhysRevA.89.032307](https://doi.org/10.1103/PhysRevA.89.032307). URL <https://link.aps.org/doi/10.1103/PhysRevA.89.032307>.
- [36] Matthias Wenzel, Fabian Böttcher, Jan-Niklas Schmidt, Michael Eisenmann, Tim Langen, Tilman Pfau, and Igor Ferrier-Barbut. Anisotropic superfluid behavior of a dipolar bose-einstein condensate. *Physical Review Letters*, 121(3), Jul 2018. ISSN 1079-7114. DOI: [10.1103/PhysRevLett.121.030401](https://doi.org/10.1103/PhysRevLett.121.030401). URL <http://dx.doi.org/10.1103/PhysRevLett.121.030401>.
- [37] Yi-Quan Zou, Ling-Na Wu, Qi Liu, Xin-Yu Luo, Shuai-Feng Guo, Jia-Hao Cao, Meng Khoon Tey, and Li You. Beating the classical precision limit with spin-1 dicke states of more than 10,000 atoms. *Proceedings of the National Academy of Sciences*, 115(25):6381–6385, Jun 2018. ISSN 1091-6490. DOI: [10.1073/pnas.1715105115](https://doi.org/10.1073/pnas.1715105115). URL <http://dx.doi.org/10.1073/pnas.1715105115>.

A Derivation of the main entanglement witness inequality

In this section, we derive our central result, Eq. (4). In order to detect entanglement from the available data [Eqs. (1) and (2)], our starting point is to assume that the latter are explained by a separable state, namely a quantum state of the form:

$$\hat{\rho}_{\text{sep}} = \sum_{\lambda} p_{\lambda} \otimes_{i=1}^N \hat{\rho}_{\lambda}(i) , \quad (12)$$

and to arrive at a contradiction. We introduce the notations:

$$m_k(\lambda, i) := \text{Tr}[\hat{\rho}_{\lambda}(i) \hat{o}_k(i)] \quad (13)$$

$$m_k(\lambda) := \frac{1}{N} \sum_{i=1}^N m_k(\lambda, i) . \quad (14)$$

The key insight of our approach is that, if the state is separable, we may interpret the quantities $m_k(\lambda, i)$ as local classical variables, whose collective fluctuations must reproduce the observed fluctuations of collective observables. The constraints that such local classical variables must obey give rise to our central inequality [Eq. (4)]. Most importantly, the constraints of Eq. (3) can be expressed as:

$$\frac{1}{N} \sum_{i=1}^N \sum_k [m_k(\lambda, i)]^2 \leq \beta \quad (15)$$

for all λ . With the above notations, the one-body quantum data [Eq. (1)] read:

$$m_k = \frac{1}{N} \sum_{i=1}^N \langle \hat{o}_k(i) \rangle = \sum_{\lambda} p_{\lambda} m_k(\lambda) \quad (16)$$

We then introduce the vector notation $\mathbf{m} := (m_k)$, and the outer product notation $[\mathbf{u} \otimes \mathbf{v}]_{ab} = u_a v_b$. With these notations, the two-body quantum data [Eq. (2)] read:

$$C = \frac{1}{N(N-1)} \sum_{i \neq j} \sum_{\lambda} p_{\lambda} \mathbf{m}(\lambda, i)^* \otimes \mathbf{m}(\lambda, j) . \quad (17)$$

We now introduce the fluctuations of the $m_k(\lambda, i)$ variables:

$$A := \sum_{\lambda} p_{\lambda} [\mathbf{m}(\lambda) - \mathbf{m}]^* \otimes [\mathbf{m}(\lambda) - \mathbf{m}] \succeq 0 \quad (18)$$

$$a := \sum_{\lambda} p_{\lambda} \frac{1}{N} \sum_i [\mathbf{m}(\lambda, i) - \mathbf{m}(\lambda)]^* \otimes [\mathbf{m}(\lambda, i) - \mathbf{m}(\lambda)] \succeq 0 \quad (19)$$

The notation $M \succeq 0$ means the the matrix M is positive semidefinite (PSD), that is, for any vector \mathbf{u} , $\mathbf{u}^{\dagger} M \mathbf{u} \geq 0$. This PSD property is straightforward to verify, since A and a are both of the form $\sum_x w_x \mathbf{v} \mathbf{v}^{\dagger}$ with $w_x \geq 0$ and \mathbf{v} a one-dimensional vector.

The matrix C is then:

$$C = \frac{N}{N-1} [\mathbf{m}^* \otimes \mathbf{m} + A] - \frac{1}{N(N-1)} \sum_{\lambda} p_{\lambda} \sum_i \mathbf{m}(\lambda, i)^* \otimes \mathbf{m}(\lambda, i) \quad (20)$$

$$= \mathbf{m}^* \otimes \mathbf{m} + A - \frac{1}{N-1} a . \quad (21)$$

From Eq. (20) and Eq. (15)], we have:

$$\text{Tr}(C) \geq \frac{N}{N-1} \text{Tr}[\mathbf{m}^* \otimes \mathbf{m} + A] - \frac{\beta}{N-1} . \quad (22)$$

We then introduce a projector P , and write:

$$\text{Tr}[\mathbf{m}^* \otimes \mathbf{m} + A] = \text{Tr}[P(\mathbf{m}^* \otimes \mathbf{m} + A)] + \text{Tr}[(1 - P)(\mathbf{m}^* \otimes \mathbf{m} + A)] \quad (23)$$

$$= \text{Tr}[P(\mathbf{m}^* \otimes \mathbf{m} + A)] + \text{Tr}[(1 - P)(C + a/(N - 1))] \quad (24)$$

where we used Eq. (21). From the positivity of A and P , we have $\text{Tr}(PA) \geq 0$. Similarly, from the positivity of a and $1 - P$ (which is also a projector), we have $\text{Tr}[(1 - P)a] \geq 0$. Hence:

$$\text{Tr}[\mathbf{m}^* \otimes \mathbf{m} + A] \geq \text{Tr}[P(\mathbf{m}^* \otimes \mathbf{m})] + \text{Tr}[(1 - P)C] \quad (25)$$

Introducing this inequality into Eq. (22), and reorganizing the terms, this leads to our central result:

$$L(P) := N \text{Tr}[P(C - \mathbf{m}^* \otimes \mathbf{m})] \geq \text{Tr}(C) - \beta, \quad (26)$$

which is valid for any projector P whenever the state is separable [Eq. (12)]. This central inequality is a generalization of the results contained of Vitagliano *et al.* [35]. In order to detect entanglement at best, one has to find the optimal projector P such that inequality (26) is violated – which would invalidate our starting assumption, namely that the state is separable. This optimization is straightforward: in order to minimize L over the projector P , one has to choose P as the projector onto the subspace of negative eigenvalues of $C - \mathbf{m}^* \otimes \mathbf{m}$.

B The spinor Bose gas model

In this section, we present some background information, based on Ref. [18], on the BEC Hamiltonians used in the main text [Eqs. (5) and (8)]. We consider an ensemble of spin- j bosonic atoms with contact, two-body, spin-conserving interactions in \mathbb{R}^3 . In the single-mode approximation, its Hamiltonian reads:

$$\hat{H}_0(\{g_f\}_{f=0}^j) = \sum_{s_1 s_2, s_3 s_4 = -j}^j g_f C_{s_1 s_2, s_3 s_4}^{j \otimes j = 2f} \hat{a}_{s_1}^\dagger \hat{a}_{s_2}^\dagger \hat{a}_{s_3} \hat{a}_{s_4}, \quad (27)$$

where g_f is the scattering amplitude of the spin- $2f$ channel. The matrix element of the projection of a composite spin- $2f$ onto two spin- j atoms is $C_{s_1 s_2, s_3 s_4}^{j \otimes j = 2f} = \langle j, s_1 |_j \otimes \langle j, s_2 |_j \mathcal{P}^{j \otimes j = 2f} |j, s_3 \rangle_j \otimes |j, s_4 \rangle_j$, with $\mathcal{P}^{j \otimes j = 2f} = \sum_{S=-2f}^{2f} |2f, S \rangle_{j \otimes j} \langle 2f, S |_{j \otimes j}$, where S is the spin projection of the composite spin- $2f$. The operators $\hat{a}_s (\hat{a}_s^\dagger)$ annihilate (create) a bosonic mode with spin projection $s \in \{-j, -j + 1, \dots, j\}$. The total number of atoms $\hat{N} = \sum_{s=-j}^j \hat{N}_s$, where $\hat{N}_s = \hat{a}_s^\dagger \hat{a}_s$, is fixed to N .

In the following we will express the Hamiltonian (27) in terms of collective operators. This is done via the identification:

$$\hat{O} = \sum_{s_L, s_R = -j}^j \langle s_L | \hat{o} | s_R \rangle \hat{a}_{s_L}^\dagger \hat{a}_{s_R}, \quad (28)$$

where $\hat{O} = \sum_{i=1}^N \hat{o}^i$ is the collective operator corresponding to the local observable \hat{o} acting on a single-atom spin- j degree of freedom. Note that in particular the population operator \hat{N}_s is a collective operator with $\langle s_L | \hat{n}_s | s_R \rangle := \langle s_L | s \rangle \langle s | s_R \rangle = \delta_{s_L, s} \delta_{s, s_R}$.

Spin 1. For spin-1, the spin projection onto the cartesian basis $\{x, y, z\}$ reads:

$$\hat{J}^{(x)} = \frac{1}{\sqrt{2}} \begin{pmatrix} \hat{a}_{+1} & \hat{a}_0 & \hat{a}_{-1} \end{pmatrix} \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix} \begin{pmatrix} \hat{a}_{+1} \\ \hat{a}_0 \\ \hat{a}_{-1} \end{pmatrix} \quad (29)$$

$$\hat{J}^{(y)} = \frac{1}{\sqrt{2}} \begin{pmatrix} \hat{a}_{+1} & \hat{a}_0 & \hat{a}_{-1} \end{pmatrix} \begin{pmatrix} 0 & -\mathfrak{i} & 0 \\ \mathfrak{i} & 0 & -\mathfrak{i} \\ 0 & \mathfrak{i} & 0 \end{pmatrix} \begin{pmatrix} \hat{a}_{+1} \\ \hat{a}_0 \\ \hat{a}_{-1} \end{pmatrix} \quad (30)$$

$$\hat{J}^{(z)} = \begin{pmatrix} \hat{a}_{+1} & \hat{a}_0 & \hat{a}_{-1} \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix} \begin{pmatrix} \hat{a}_{+1} \\ \hat{a}_0 \\ \hat{a}_{-1} \end{pmatrix}, \quad (31)$$

where \mathfrak{i} is the imaginary unit, $\mathfrak{i}^2 = -1$. With this expressions, we can compute the total spin, using the bosonic commutation relations and $N_{+1}^{(z)} + N_0^{(z)} + N_{-1}^{(z)} = N$ we have:

$$\begin{aligned} \hat{\mathbf{J}}^2 &:= [\hat{J}^{(x)}]^2 + [\hat{J}^{(y)}]^2 + [\hat{J}^{(z)}]^2 = \\ &= [\hat{J}^{(z)}]^2 + \hat{N} + \hat{N}_0 + 2\hat{N}_0(N - \hat{N}_0) + 2((\hat{a}_0^\dagger)^2 \hat{a}_{+1} \hat{a}_{-1} + \text{h.c.}) \end{aligned} \quad (32)$$

Alternatively, by identifying the pair annihilation operator $\hat{\Theta}_2 = \hat{a}_0^2 - 2\hat{a}_{+1}\hat{a}_{-1}$, we can write $\hat{\mathbf{J}}^2 = N^2 - \hat{\Theta}^\dagger \hat{\Theta}$. As first observation, we notice that unlike the spin-1/2 case, $\hat{\mathbf{J}}^2$ is not diagonal in the occupation basis. Direct computation shows that the Hamiltonian (27) is equivalent to:

$$\hat{H}_0(c_1) = c_1 \hat{\mathbf{J}}^2, \quad (33)$$

where $c_1 = (g_1 - g_0)/3$.

Spin 2. For spin-2, there is a second relevant interaction channel. In this case, the Hamiltonian reads:

$$\hat{H}_0(c_2, p) = c_2 \hat{\mathbf{J}}^2 + p [\hat{N}_0(\hat{N}_0 - 1) + 4(\hat{N}_1 \hat{N}_{-1} + \hat{N}_2 \hat{N}_2) + 2(\hat{T}_{0,-2}^+ \hat{T}_{0,2}^+ - \hat{T}_{0,-1}^+ \hat{T}_{0,1}^+ - 2\hat{T}_{1,2}^+ \hat{T}_{-1,-2}^+ + (+ \leftrightarrow -))], \quad (34)$$

where $c_2 = (g_4 - g_2)/7$, $p = (7g_0 - 10g_2 + 3g_4)/35$ and we introduced the collective operators $\hat{T}_{s,s'}^+ = \hat{a}_s^\dagger \hat{a}_{s'} + \text{h.c.}$, $\hat{T}_{s,s'}^- = \mathfrak{i}(\hat{a}_s^\dagger \hat{a}_{s'} - \text{h.c.})$. Using the pair annihilation operator for spin-2, $\hat{\Theta}_2 = \hat{a}_0^2 - 2\hat{a}_1 \hat{a}_{-1} + 2\hat{a}_{+2} \hat{a}_{-2}$, the term coupling p can be interpreted as a pair amplitude $\hat{\Theta}_2^\dagger \hat{\Theta}_2$. This term commutes with $\hat{\mathbf{J}}^2$, although we do not have written it explicitly in a rotationally invariant way using cartesian coordinates.

Quadratic Zeeman shift. In this work we consider the gas is immersed in a uniform magnetic field of intensity B along the z direction. In this case, the total Hamiltonian becomes $\hat{H} = \hat{H}_0/N + q\hat{Q}^{(z)}$, where $q \propto B^2$ and $\hat{Q}^{(z)} = \sum_{i=1}^N [\hat{J}^{(z)}]^2$. We introduce a scaling to make the Hamiltonian extensive in N . We consider an experiment where all atoms are initially prepared in the state $\otimes_{i=1}^N |0_z\rangle$, so that we may remove the linear Zeeman term, proportional to $\hat{J}^{(z)}$ (as $\hat{J}^{(z)}$ is a symmetry of \hat{H} , and acts trivially on the state).

C Computation of the quantum data

In this appendix, we provide technical details on the computation of the quantum data used in the illustration of our method. Specifically, one has to compute one and two body Zeeman sublevel correlations of the form:

$$\langle \hat{N}_s^{(a)} \rangle, \langle \hat{N}_s^{(a)} \hat{N}_{s'}^{(a)} \rangle, \text{etc.}, \quad (35)$$

in the ground state of relevant Hamiltonians of spin- j ensembles.

Diagonalization of the Hamiltonians. The Hamiltonians considered in this paper are polynomials of collective operators $\hat{O}_k = \sum_{i=1}^N \hat{o}_k(i)$ with the spin projection $\hat{J}^{(z)}$ a symmetry. Since operators acting in different atoms commute, we notice that the collective operators $\{\hat{O}_k\}$ obey the same algebra as their their local counterparts, $\{\hat{o}_k(i)\}$. Accordingly, the problem reduces to compute the matrices of the irreducible representations (irreps) of the algebra, describing the collective operators. The local operators can be described by $d \times d$ Hermitian matrices, where $d = 2j + 1$. Let $\{\hat{\lambda}_\alpha\}_{\alpha=1}^{d^2-1}$ be a basis of

$d \times d$ Hermitian traceless matrices fulfilling the orthogonality condition $\text{Tr}[\hat{\lambda}_\alpha \hat{\lambda}_\beta] = \delta_{\alpha\beta}$, for instance the so-called generalized Gell-Mann matrices. The commutation relations:

$$\hat{\lambda}_\alpha \hat{\lambda}_\beta - \hat{\lambda}_\beta \hat{\lambda}_\alpha = \mathbb{i} \sum_{\gamma=1}^{d^2-1} f_{\alpha\beta\gamma} \hat{\lambda}_\gamma, \quad (36)$$

are characterized by the structure constants $f_{\alpha\beta\gamma}$. The structure constants are the same for the generators of the $SU(d)$ group with the Lie bracket $[\hat{\lambda}_\alpha, \hat{\lambda}_\beta] := \hat{\lambda}_\alpha \hat{\lambda}_\beta - \hat{\lambda}_\beta \hat{\lambda}_\alpha$. The representations of $SU(d)$ are labelled by partitions of N in d parts as the permutation group of N d -level atoms. In the totally symmetric sector with respect to the exchange of atoms, the matrix representation of the generators is computed $\{\hat{\Lambda}_\alpha = I[\hat{\lambda}_\alpha]\}_{\alpha=1}^{d^2-1}$. The operator $\hat{\Lambda}_\alpha$ is the collective counterpart of $\hat{\lambda}_\alpha$, that is $\hat{\Lambda}_\alpha = \sum_{i=1}^N \hat{\lambda}_\alpha(i)$. Consequently, for any other operator \hat{o} , the map I distributes linearly

$$I[\hat{o}] = N \text{Tr}[\hat{o}] \frac{\mathbb{1}}{D} + \sum_{\alpha=1}^{d^2-1} \text{Tr}[\hat{\lambda}_\alpha \hat{o}] \hat{\Lambda}_\alpha := \hat{O}, \quad (37)$$

where D is the dimension of the representation, and the identity element $\mathbb{1}$ is added in order to account for the trace of \hat{o} . The multiplicity N is introduced to consistently transform the identity, $I[\mathbb{1}] = \sum_{i=1}^N \mathbb{1} = N$. The expression (37) is an alternative of the second-quantization form (28) providing the advantage that higher moments are computed just by multiplying matrices, as specified in the following.

Gathering the correlations. The Hamiltonian is diagonalized using the previous matrix representation, allowing one to find the ground state $|\text{GS}\rangle$. Then, the computation of moments reduces to matrix multiplication matrices and computing expectation values in the ground state: $\langle \hat{O}_a \rangle = \langle \text{GS} | I[\hat{o}_a] | \text{GS} \rangle$, $\langle \hat{O}_a \hat{O}_b \rangle = \langle \text{GS} | I[\hat{o}_a] I[\hat{o}_b] | \text{GS} \rangle$, etc, for any collective operators \hat{O}_a . In this work in particular we probe the GS with magnetic-sublevel populations along a direction a , $\{\hat{N}_s^{(a)}\}_{s=-j}^j$. These are generated from the single-atom observable $\hat{n}_s^{(a)} = \prod_{s' \neq s}^j (s - \hat{j}^{(a)}) / (s - s')$, where $\hat{j}^{(a)}$ is the atomic spin projection along direction $a \in \{x, y, z\}$, i.e $\hat{j}^{(a)} := \mathbf{a} \cdot \hat{\mathbf{j}}$.

D Computation of the physically feasible region of a set of average values

In this section, we explain how to compute the feasible region showed in the figures of the main text. Formally, given two observables \hat{X} and \hat{Y} , acting on the Hilbert space for N qudits, our goal is find the possible combinations of “feasible” mean values ($x = \langle \hat{X} \rangle, y = \langle \hat{Y} \rangle$), that is, combinations of mean values which can be obtained in some (generally mixed) quantum state. Equivalently, one may look for the extremal values of $\langle \hat{Y} \rangle$, while keeping fixed the mean value $x = \langle \hat{X} \rangle$. By making statistical mixtures of the corresponding quantum states, the full convex hull of these extremal points is also feasible. This problem can be solved introducing a Lagrange multiplier λ_x , and finding the ground state of Hamiltonians of the form:

$$\hat{L}(\lambda_x) = \pm \hat{Y} - \lambda_x (\hat{X} - x \mathbb{1}). \quad (38)$$

Adjusting the multiplier λ_x ensures that the constraint $\langle \hat{X} \rangle = x$ is met in the ground state. More generally, the extremal points (x, y) are found as the expectation values of \hat{X}, \hat{Y} in the ground state of:

$$\hat{L}(\lambda_x, \lambda_y) = \lambda_x \hat{X} + \lambda_y \hat{Y}, \quad (39)$$

varying the parameters λ_x, λ_y between $-\infty$ and $+\infty$. By forming statistical mixtures of these ground states, the full convex hull of the extremal points are also feasible, which leads to the construction showed in the figures of the main text.

In order to find the ground state of \hat{L} , the same technique outlined above for the Bose gas Hamiltonian can be applied since in the cases of interest here, \hat{L} is a polynomial of collective operators.

E Derivation of the entanglement witness inequalities (6), (7) and their generalization to arbitrary integer spins Eqs. (9), (10)

These inequalities are found by choosing, as local measurements, the spin observables $\hat{S}^{(a)}(i) = \sum_{s=-j}^j s \hat{n}_s^{(a)}(i)$, as well as the projectors $\hat{n}_0^{(a)}(s)$ onto the $s = 0$ magnetic sublevels, along the three orthogonal directions $a \in \{x, y, z\}$. For completeness, we rewrite here the EW inequalities (9) and (10) of the main text:

$$\text{Var}(\hat{\mathbf{J}}) - \frac{1}{N-1} \langle \hat{\mathbf{N}}_0 (\hat{\mathbf{N}}_0 - \mathbf{1}) \rangle \geq N[j(j+1) - \beta_j], \quad (40)$$

and:

$$(N-1)\text{Var}(\hat{J}^{(z)}) - N\langle \hat{Q}^{(z)} \rangle - \langle [\hat{J}^{(x)}]^2 + [\hat{J}^{(y)}]^2 \rangle + N[(N-1)\beta_j + j(j+1)] \geq \langle \hat{\mathbf{N}}_0 (\hat{\mathbf{N}}_0 - \mathbf{1}) \rangle, \quad (41)$$

where $\mathbf{1} = (1, 1, 1)$ and β_j is the corresponding bound in Eq. (3). We evaluated this bound numerically, and the corresponding values are:

spin j	1	2	3	4	5
$\beta \{\mathbf{S}, \mathbf{n}_0\}$	1.5	4.3	9.2	16.15	25.22

Table 1: Numerical separable bound β for the basis of local measurements $\{\{\hat{S}^{(a)}(i), \hat{n}_0^{(a)}(i)\}_{a=\{x,y,z\}}\}_{i=\{1,2,\dots,N\}}$ for different spin- j .

In the specific case of $j = 1$, the EW inequalities can be simplified using the identities: $\sum_{a \in \{x,y,z\}} (\hat{N}_{+1}^{(a)} + \hat{N}_{-1}^{(a)}) = 3N - \sum_{a \in \{x,y,z\}} \hat{N}_0^{(a)} = \sum_{i=1}^N [\hat{\mathbf{S}}(i)]^2 = j(j+1)N = 2N$. Together with the bound $\beta_{j=1} = 3/2$, we obtain Eqs. (6), (7) from respectively Eqs. (40) and (41).

Derivation of $\text{Tr}[C]$. To derive these EWs from our central inequality, Eq. (4), one has to compute $\text{Tr}[C]$, which reads:

$$\text{Tr}[C] = \frac{1}{N(N-1)} \sum_{i \neq j} \sum_{a \in \{x,y,z\}} \langle \hat{S}^{(a)}(i) \hat{S}^{(a)}(j) + \hat{n}_0^{(a)}(i) \hat{n}_0^{(a)}(j) \rangle \quad (42)$$

$$= \frac{1}{N(N-1)} \sum_{a \in \{x,y,z\}} \left\langle [\hat{J}^{(a)}]^2 + [\hat{N}_0^{(a)}]^2 - \sum_{i=1}^N [[\hat{S}^{(a)}(i)]^2 + \hat{n}_0^{(a)}(i)] \right\rangle \quad (43)$$

$$= \frac{1}{N(N-1)} \left(\langle \hat{\mathbf{J}}^2 + \hat{\mathbf{N}}_0 (\hat{\mathbf{N}}_0 - \mathbf{1}) \rangle - j(j+1)N \right) \quad (44)$$

where $\hat{\mathbf{J}} = (\hat{J}^{(x)}, \hat{J}^{(y)}, \hat{J}^{(z)})$ is the collective spin, $\hat{\mathbf{N}}_0 = (\hat{N}_0^{(x)}, \hat{N}_0^{(y)}, \hat{N}_0^{(z)})$ and $\mathbf{1} = (1, 1, 1)$. We used that $\sum_a [\hat{S}^{(a)}(i)]^2 = j(j+1)$, and that $[\hat{n}_0^{(a)}(i)]^2 = \hat{n}_0^{(a)}(i)$. The two EW are obtained by choosing two different projectors in Eq. (4). Recall that, since correlations in different orientations are not observed, P has a block-diagonal form $(P^{(x)}, P^{(y)}, P^{(z)})$.

Derivation of Eq. (40). For Eq. (40) (tailored in particular to a spin- j many-body singlet), the optimal projection P is $P^{(x)} = P^{(y)} = P^{(z)} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} := P_j$ in the local basis described before, leading to

$$\text{Tr}[NP(C - \mathbf{m} \otimes \mathbf{m})] = \frac{1}{N-1} \left(\langle \hat{\mathbf{J}}^2 \rangle - j(j+1)N \right) - \frac{1}{N} \langle \hat{\mathbf{J}} \rangle^2. \quad (45)$$

Inserting this expression, together with Eq. (44), into our central inequality (4), we obtain the announced result Eq. (40).

Derivation of Eq. (41). For Eq. (41) (tailored in particular to the Dicke states $\mathbf{J}^2 = Nj(Nj + 1)$, $J^{(z)} = 0$), the optimal projector is $P^{(x)} = P^{(y)} = 0$, $P^{(z)} = P_j$, from which we find:

$$\text{Tr}[NP(C - \mathbf{m} \otimes \mathbf{m})] = \frac{1}{N-1} \langle (\hat{J}^{(z)})^2 - \hat{Q}_z \rangle - \frac{1}{N} \langle \hat{J}^{(z)} \rangle^2. \quad (46)$$

Inserting this expression, together with Eq. (44), into our central inequality (4), we obtain the announced result Eq. (41).

F Derivation of spin-2 EW, inequality (11) and generalization to arbitrary integer spins

In this appendix we show how to derive the entanglement witness presented in Eq. (11), and violated by the pair condensate, but not by the triplet condensate. In the present case, we use as local observables the projectors onto all magnetic sublevels, except the $s = 0$ sublevel: $\{\hat{n}_{+2}^{(a)}(i), \hat{n}_{+1}^{(a)}(i), \hat{n}_{-1}^{(a)}(i), \hat{n}_{-2}^{(a)}(i)\}_{a \in \{x,y,z\}}$. This choice leads to the $\text{Tr}(C)$ term:

$$\text{Tr}(C) = \frac{1}{N(N-1)} \sum_{a \in \{x,y,z\}} \sum_{s \neq 0} \langle [\hat{N}_s^{(a)}]^2 - \hat{N}_s^{(a)} \rangle \quad (47)$$

$$\frac{1}{N(N-1)} \sum_{a \in \{x,y,z\}} \left(\sum_{s \neq 0} \langle [\hat{N}_s^{(a)}]^2 \rangle - (N - \langle \hat{N}_0^{(a)} \rangle) \right) \quad (48)$$

$$= \frac{1}{N(N-1)} \sum_{a \in \{x,y,z\}} \left(\sum_{s \neq 0} \langle [\hat{N}_s^{(a)}]^2 \rangle + \langle \hat{N}_0^{(a)} \rangle \right) - \frac{3}{N-1} \quad (49)$$

The method reveals the optimal projector $P^{(x)} = P^{(y)} = P^{(z)} = \frac{1}{2} \begin{pmatrix} 1 & 0 & 0 & -1 \\ 0 & 1 & -1 & 0 \\ 0 & -1 & 1 & 0 \\ -1 & 0 & 0 & 1 \end{pmatrix}$. This

yields the terms:

$$\text{Tr}[NPC] = \frac{1}{2(N-1)} \left(\sum_{a \in \{x,y,z\}} \langle [\hat{N}_{+2}^{(a)} - \hat{N}_{-2}^{(a)}]^2 + [\hat{N}_{+1}^{(a)} - \hat{N}_{-1}^{(a)}]^2 + \hat{N}_0^{(a)} \rangle - 3N \right) \quad (50)$$

$$\text{Tr}[NP(\mathbf{m} \otimes \mathbf{m})] = \frac{1}{2N} \sum_{a \in \{x,y,z\}} \left(\langle \hat{N}_{+2}^{(a)} - \hat{N}_{-2}^{(a)} \rangle^2 + \langle \hat{N}_{+1}^{(a)} - \hat{N}_{-1}^{(a)} \rangle^2 \right). \quad (51)$$

We introduce the notation $\hat{\sigma}_s^{(a)} = \hat{N}_s^{(a)} - \hat{N}_{-s}^{(a)}$, the population imbalance between opposed levels along direction a . Together with the notation $\mathbf{1} = (1, 1, 1)$, we can rewrite these equations as:

$$2(N-1)\text{Tr}(C) = \frac{2}{N} \sum_{s \neq 0} \langle [\hat{\mathbf{N}}_s]^2 \rangle + \frac{2}{N} \langle \hat{\mathbf{N}}_0 \cdot \mathbf{1} \rangle - 6 \quad (52)$$

$$2N(N-1)\text{Tr}[PC] = \langle [\hat{\boldsymbol{\sigma}}_1]^2 + [\hat{\boldsymbol{\sigma}}_2]^2 \rangle + \langle \hat{\mathbf{N}}_0 \cdot \mathbf{1} \rangle - 3N \quad (53)$$

$$2N(N-1)\text{Tr}[P(\mathbf{m} \otimes \mathbf{m})] = \frac{N-1}{N} [\langle \hat{\boldsymbol{\sigma}}_1 \rangle^2 + \langle \hat{\boldsymbol{\sigma}}_2 \rangle^2]. \quad (54)$$

Together with the bound $\beta = 3/2$ found numerically, we find:

$$2(N-1)[N\text{Tr}[P(C - \mathbf{m} \otimes \mathbf{m})] - \text{Tr}(C) + \beta] = \langle [\hat{\boldsymbol{\sigma}}_1]^2 + [\hat{\boldsymbol{\sigma}}_2]^2 \rangle - \frac{N-1}{N} [\langle \hat{\boldsymbol{\sigma}}_1 \rangle^2 + \langle \hat{\boldsymbol{\sigma}}_2 \rangle^2] + \left(1 - \frac{2}{N}\right) \langle \hat{\mathbf{N}}_0 \cdot \mathbf{1} \rangle - \frac{2}{N} \sum_{s \neq 0} \langle [\hat{\mathbf{N}}_s]^2 \rangle + 3, \quad (55)$$

namely, Eq. (11).

Generalization to arbitrary even-level systems. A possible generalization to arbitrary integer spin is obtained by taking the projector to $\sum_{s=1}^j (\hat{n}_s^{(a)} - \hat{n}_{-s}^{(a)})$ for $a \in \{x, y, z\}$. The corresponding bound β for local measurements $\{\{\hat{n}_s^{(a)}\}_{s=-j \neq 0}^j\}_{a \in \{x, y, z\}}$ obtained numerically is summarized in the following table for several spin- j .

spin j	1	2	3	4	5
β for $\{\hat{n} \setminus \hat{\mathbf{n}}_0\}$	1.25	1.5	1.5	1.3369	1.2926

Table 2: Numerical separable bound β for the basis of local measurements $\{\{\hat{n}_s^{(a)}\}_{s=-j \neq 0}^j\}_{a \in \{x, y, z\}}$ for different spin- j .

Notice that for $j = 1$, we recover Eq. (6).