



HAL
open science

A dimensional mass transference principle for Borel probability measures and applications

Édouard Daviaud

► **To cite this version:**

Édouard Daviaud. A dimensional mass transference principle for Borel probability measures and applications. 2023. hal-03626178v2

HAL Id: hal-03626178

<https://hal.science/hal-03626178v2>

Preprint submitted on 14 Jul 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A DIMENSIONAL MASS TRANSFERENCE PRINCIPLE FOR BOREL PROBABILITY MEASURES AND APPLICATIONS

EDOUARD DAVIAUD

ABSTRACT. In this article, we establish a dimensional mass transference principle valid when the ambient measure is finite. We provide two applications of this result. First we study certain dynamical coverings associated with some self-similar IFS with overlaps and then we give an application in Diophantine approximation to rational approximation among points of $[0, 1]$ with few digit equal to 1 in their base-3 expansion.

1. INTRODUCTION

Estimating the Hausdorff dimension of limsup sets obtained from the contractions of the elements of a given family of sets is a natural question of metric approximation theory, which arises in many contexts. In this article, given a sequence of balls of \mathbb{R}^d , $\mathcal{B} = (B_n := B(x_n, r_n))_{n \in \mathbb{N}}$, we investigate in a very general frame the size properties of the limsup sets obtained from smaller sets, i.e sets of the form $\limsup_{n \rightarrow +\infty} U_n$, where $U_n \subset B_n$.

Let us recall that the historical example of Jarnik-Besicovitch's theorem deals with the case $U_n = B_n^\delta := B(x_n, r_n^\delta)$, where $\delta > 1$, x_n is a rational number $\frac{p}{q}$ and $r_n = \frac{1}{q^2}$.

Generalizations of Jarnik-Besicovitch's Theorem often consider a given sequence $(x_n)_{n \in \mathbb{N}}$ of points in \mathbb{R}^d , as well as a sequence of radii $(r_n)_{n \in \mathbb{N}}$ for which the associated limsup set $E_1 = \limsup_{n \rightarrow +\infty} B_n$ has a controlled size (in terms of Lebesgue measure or Hausdorff dimension for instance); then, given a sequence of sets $\mathcal{U} = (U_n)_{n \in \mathbb{N}}$ with, for every $n \in \mathbb{N}$, $U_n \subset B_n$, one estimates the Hausdorff dimension of the smaller limsup set $E(\mathcal{U}) = \limsup_{n \rightarrow +\infty} U_n$. The classical case above studies the case where $U_n = B_n^\delta$ but different shapes for U_n have also been considered (rectangles or ellipsoïds rather than balls for instance). Such problems are studied for instance in [26, 8, 7, 33, 28] among many references.

The same question arises in any topological dynamical system (X, T) endowed with some metric, when the sequence $(x_n)_{n \in \mathbb{N}}$ is the orbit $(T^n(x))_{n \in \mathbb{N}}$ of a well chosen point x . Some specific cases are for instance treated in [23, 29, 30]. In probability theory, the famous Dvoretzky covering problem consists in computing, when it is possible, the Hausdorff dimension of the limsup set associated with a sequence of random balls drawn independently and uniformly in a compact Baire space, see for instance [20, 15, 7]. In analysis, the value of the pointwise regularity exponents of measures and functions at a given point x often relies on the ability to understand how x is close to remarkable points x_n . The reader may refer to [26, 4, 6].

As mentioned above, in the largest part of the literature, a strong geometric measure theoretic condition is initially imposed on \mathcal{B} to obtain results, for instance that the Lebesgue measure of $\limsup_{n \rightarrow +\infty} B(x_n, r_n)$ is full (cf [8]). But there are

many situations in which the Lebesgue measure is not the relevant measure to work with (cf [5]).

Our purpose in this article is to obtain a general lower bound for the Hausdorff dimension $\limsup_{n \rightarrow +\infty} U_n$, where the sets (U_n) are open sets in some balls (B_n) satisfying the property called μ -asymptotically covering property, where μ is a probability measure on \mathbb{R}^d . This property, introduced in [14], is proved to be almost equivalent to verifying that $\mu(\limsup_{n \rightarrow +\infty} B_n) = 1$ (e.g [14]).

The results presented here extend, for instance, both the results of Koivusalo-Rams stated in [28] and the result of Barral-Seuret ([7]) which deals with balls and self-similar measures under the open set condition. It is worth noticing that the works of Koivusalo-Rams, Ghosh and Nandi and Bique-Eriksson in [28, 21, 16] highlighted the importance of the Hausdorff content to compute Hausdorff dimension of linsup sets and this article makes further use of this fact.

An important advantage of the lower bound obtained in the present paper is that its value is tractable in many cases. For instance, as a first application, a dimensional mass transference principle is given in the case where μ is a self-similar measure (we do not require any condition on the possible overlaps associated with such a μ).

Two other applications of our main result are treated in this article as well. The problem of self-similar dynamical coverings is studied when the corresponding iterated function system (in short IFS) is dimension-regular and has similarity dimension less than d (see Section 2.14, Definition 2.9).

Another application in Diophantine approximation is given. Let $K_{1/3}^{(0)}$ the set of points of $[0, 1]$ such that in their sequence of digits in basis 3, the asymptotic frequency of appearance of the digit 1 is infinitely many often close to 0 (note that this set contains the middle-third Cantor set $K_{1/3}$ and $\dim_H(K_{1/3}^{(0)}) = \dim_H(K_{1/3})$). We compute the Hausdorff dimension of points of $K_{1/3}^{(0)}$ well approximable by rationals (see Theorem 2.15 for a precise statement).

2. DEFINITIONS AND MAIN STATEMENTS

Let us start with some notations

Let $d \in \mathbb{N}$. For $x \in \mathbb{R}^d$, $r > 0$, $B(x, r)$ stands for the closed ball of $(\mathbb{R}^d, \|\cdot\|_\infty)$ of center x and radius r . Given a ball B , $|B|$ stands for the diameter of B . For $t \geq 0$, $\delta \in \mathbb{R}$ and $B = B(x, r)$, tB stand for $B(x, tr)$, i.e. the ball with same center as B and radius multiplied by t , and the δ -contracted ball B^δ is defined by $B^\delta = B(x, r^\delta)$.

Given a set $E \subset \mathbb{R}^d$, $\overset{\circ}{E}$ stands for the interior of the E , \overline{E} its closure and ∂E its boundary, i.e. $\partial E = \overline{E} \setminus \overset{\circ}{E}$. If E is a Borel subset of \mathbb{R}^d , its Borel σ -algebra is denoted by $\mathcal{B}(E)$.

Given a topological space X , the Borel σ -algebra of X is denoted $\mathcal{B}(X)$ and the space of probability measure on $\mathcal{B}(X)$ is denoted $\mathcal{M}(X)$.

The d -dimensional Lebesgue measure on $(\mathbb{R}^d, \mathcal{B}(\mathbb{R}^d))$ is denoted by \mathcal{L}^d .

For $\mu \in \mathcal{M}(\mathbb{R}^d)$, $\text{supp}(\mu) = \{x \in [0, 1] : \forall r > 0, \mu(B(x, r)) > 0\}$ is the topological support of μ .

Given $E \subset \mathbb{R}^d$, $\dim_H(E)$ and $\dim_P(E)$ denote respectively the Hausdorff and the packing dimension of E .

Now we recall some definitions.

Definition 2.1. Let $\zeta : \mathbb{R}^+ \mapsto \mathbb{R}^+$. Suppose that ζ is increasing in a neighborhood of 0 and $\zeta(0) = 0$. The Hausdorff outer measure at scale $t \in (0, +\infty]$ associated with ζ of a set E is defined by

$$(1) \quad \mathcal{H}_t^\zeta(E) = \inf \left\{ \sum_{n \in \mathbb{N}} \zeta(|B_n|) : |B_n| \leq t, B_n \text{ closed ball and } E \subset \bigcup_{n \in \mathbb{N}} B_n \right\}.$$

The Hausdorff measure associated with ζ of a set E is defined by

$$(2) \quad \mathcal{H}^\zeta(E) = \lim_{t \rightarrow 0^+} \mathcal{H}_t^\zeta(E).$$

For $t \in (0, +\infty]$, $s \geq 0$ and $\zeta : x \mapsto x^s$, one simply uses the usual notation $\mathcal{H}_t^\zeta(E) = \mathcal{H}_t^s(E)$ and $\mathcal{H}^\zeta(E) = \mathcal{H}^s(E)$, and these measures are called s -dimensional Hausdorff outer measure at scale $t \in (0, +\infty]$ and s -dimensional Hausdorff measure respectively. Thus,

$$(3) \quad \mathcal{H}_t^s(E) = \inf \left\{ \sum_{n \in \mathbb{N}} |B_n|^s : |B_n| \leq t, B_n \text{ closed ball and } E \subset \bigcup_{n \in \mathbb{N}} B_n \right\}.$$

The quantity $\mathcal{H}_\infty^s(E)$ (obtained for $t = +\infty$) is called the s -dimensional Hausdorff content of the set E .

Definition 2.2. Let $\mu \in \mathcal{M}(\mathbb{R}^d)$. For $x \in \text{supp}(\mu)$, the lower and upper local dimensions of μ at x are defined as

$$\underline{\dim}_{\text{loc}}(\mu, x) = \liminf_{r \rightarrow 0^+} \frac{\log(\mu(B(x, r)))}{\log(r)} \quad \text{and} \quad \overline{\dim}_{\text{loc}}(\mu, x) = \limsup_{r \rightarrow 0^+} \frac{\log(\mu(B(x, r)))}{\log(r)}.$$

Then, the lower and upper Hausdorff dimensions of μ are respectively defined by

$$(4) \quad \underline{\dim}_H(\mu) = \text{ess inf}_\mu(\underline{\dim}_{\text{loc}}(\mu, x)) \quad \text{and} \quad \overline{\dim}_P(\mu) = \text{ess sup}_\mu(\overline{\dim}_{\text{loc}}(\mu, x)).$$

It is known (for more details see [17]) that

$$\begin{aligned} \underline{\dim}_H(\mu) &= \inf \{ \dim_H(E) : E \in \mathcal{B}(\mathbb{R}^d), \mu(E) > 0 \} \\ \overline{\dim}_P(\mu) &= \inf \{ \dim_P(E) : E \in \mathcal{B}(\mathbb{R}^d), \mu(E) = 1 \}. \end{aligned}$$

When $\underline{\dim}_H(\mu) = \overline{\dim}_P(\mu)$, this common value is simply denoted by $\dim(\mu)$ and μ is said to be *exact dimensional*.

2.1. The μ -a.c property. We fix a sequence of closed balls $\mathcal{B} = (B_n)_{n \in \mathbb{N}}$ such that $\lim_{n \rightarrow +\infty} |B_n| = 0$ (otherwise the situation is trivial for the questions we consider).

The main property (introduced in [14]) used for the sequence of balls \mathcal{B} is meant to ensure that any set can be covered efficiently by the limsup of the B_n 's, with respect to a measure μ . This property is a general version of the key covering property used in the KGB Lemma of Beresnevitch and Velani, stated in [8], using a Borel probability measure μ . Observe that such properties (like the KGB Lemma) are usually key (cf [26, 8, 5] for instance) to prove ubiquity or mass transference results.

Definition 2.3. Let $\mu \in \mathcal{M}(\mathbb{R}^d)$. The sequence $\mathcal{B} = (B_n)_{n \in \mathbb{N}}$ of balls of \mathbb{R}^d is said to be μ -asymptotically covering (in short, μ -a.c) when there exists a constant $C > 0$ such that for every open set $\Omega \subset \mathbb{R}^d$ and $g \in \mathbb{N}$, there is an integer $N_\Omega \in \mathbb{N}$ as well as integers $\{n_i\}_{1 \leq i \leq N_\Omega}$ satisfying $g \leq n_1 \leq \dots \leq n_{N_\Omega}$ such that:

- (i) $\forall 1 \leq i \leq N_\Omega, B_{n_i} \subset \Omega$;
- (ii) $\forall 1 \leq i \neq j \leq N_\Omega, B_{n_i} \cap B_{n_j} = \emptyset$;
- (iii) also,

$$(5) \quad \mu\left(\bigcup_{i=1}^{N_\Omega} B_{n_i}\right) \geq C\mu(\Omega).$$

In other words, for any open set Ω and any integer $g \geq 1$, there exists a finite set of disjoint balls of $\{B_n\}_{n \geq g}$ supporting a fixed proportion of $\mu(\Omega)$.

This notion of μ -asymptotically covering is related to the way the balls of \mathcal{B} are distributed according to the measure μ . This property is a priori slightly stronger than having a limsup of full μ -measure when μ is not doubling, as suggested by the following proposition proved in [14].

Proposition 2.1. Let $\mu \in \mathcal{M}(\mathbb{R}^d)$ and $\mathcal{B} = (B_n := B(x_n, r_n))_{n \in \mathbb{N}}$ be a sequence of balls of \mathbb{R}^d with $\lim_{n \rightarrow +\infty} r_n = 0$.

- (1) If \mathcal{B} is μ -a.c, then $\mu(\limsup_{n \rightarrow +\infty} B_n) = 1$.
- (2) If there exists $v < 1$ such that $\mu(\limsup_{n \rightarrow +\infty} (vB_n)) = 1$, then \mathcal{B} is μ -a.c.

The second item will be used to apply our main theorem to self-similar measures. It follows from the proof of [8, Lemma 5] that these properties are equivalent when μ is doubling.

2.2. Essential content and statement of the main result. The key geometric notion for the mass transference principle developed in this paper is the following.

Definition 2.4. Let $\mu \in \mathcal{M}(\mathbb{R}^d)$, and $s \geq 0$. The s -dimensional μ -essential Hausdorff content at scale $t \in (0, +\infty]$ of a set $A \subset \mathcal{B}(\mathbb{R}^d)$ is defined as

$$(6) \quad \mathcal{H}_t^{\mu, s}(A) = \inf \{ \mathcal{H}_t^s(E) : E \subset A, \mu(E) = \mu(A) \}.$$

One will almost exclusively look at these contents at scale $t = +\infty$ and one refers to $\mathcal{H}_\infty^{\mu, s}(A)$ as the s -dimensional μ -essential Hausdorff content of A . Basic properties of these quantities are studied in Section 3.3, and precise estimates of $\mathcal{H}_\infty^{\mu, s}(A)$ are achieved for the Lebesgue measure and self-similar measures in Section 5.

Note that in [28, Theorem 3.1] the key underlying geometric notion used to handle the variety of shapes of the sets $(U_n)_{n \in \mathbb{N}}$ is the Hausdorff content. It is easily seen from (3) that the Hausdorff content also carries some “high scale” geometric information (because there is no restriction concerning the diameter of the balls (B_n) in (3)). This will also be the case in this article to handle not only the shape of the sets $(U_n)_{n \in \mathbb{N}}$ but also the geometric behavior related to the measure μ at high scale in the sets $(U_n)_{n \in \mathbb{N}}$.

The s -dimensional μ -essential Hausdorff content is now used to associate a critical exponent to any sequence of open sets $(U_n)_{n \in \mathbb{N}}$ such that $U_n \subset B_n$ for all $n \in \mathbb{N}$. This exponent is involved in our lower bound estimate for $\dim_H(\limsup_{n \rightarrow +\infty} U_n)$.

Definition 2.5. Let $\mu \in \mathcal{M}(\mathbb{R}^d)$. If B and U are Borel subsets of \mathbb{R}^d , the μ -critical exponent of (B, U) is defined as

$$(7) \quad s_\mu(B, U) = \sup \{s \geq 0 : \mathcal{H}_\infty^{\mu, s}(U) \geq \mu(B)\}.$$

Let $\mathcal{B} = (B_n)_{n \in \mathbb{N}}$ be a sequence of closed balls, $\mathcal{U} = (U_n)_{n \in \mathbb{N}}$ a sequence of Borel subsets of \mathbb{R}^d , and $s \geq 0$.

Let

$$(8) \quad \mathcal{N}_\mu(\mathcal{B}, \mathcal{U}, s) = \{n \in \mathbb{N} : s_\mu(B_n, U_n) \geq s\}.$$

The μ -critical exponent of $(\mathcal{B}, \mathcal{U})$ is given by

$$(9) \quad s(\mu, \mathcal{B}, \mathcal{U}) = \sup \{s \geq 0 : (B_n)_{n \in \mathcal{N}_\mu(\mathcal{B}, \mathcal{U}, s)} \text{ is } \mu\text{-a.c.}\}.$$

Note that, for $s' \leq s$, one has $\mathcal{N}_\mu(\mathcal{B}, \mathcal{U}, s) \subset \mathcal{N}_\mu(\mathcal{B}, \mathcal{U}, s')$ so that (9) makes sense.

The main result of this paper is the following.

Theorem 2.2. Let $\mathcal{B} = (B_n)_{n \in \mathbb{N}}$ be a sequence of closed balls of \mathbb{R}^d such that $|B_n| \rightarrow 0$ and let $\mathcal{U} = (U_n)_{n \in \mathbb{N}}$ be a sequence of open sets such that $U_n \subset B_n$ for all $n \in \mathbb{N}$.

Then, for every $\mu \in \mathcal{M}(\mathbb{R}^d)$ such that $\min \{s(\mu, \mathcal{B}, \mathcal{U}), \underline{\dim}_H(\mu)\} > 0$ there exists a gauge function $\zeta : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ such that $\lim_{r \rightarrow 0^+} \frac{\log \zeta(r)}{\log r} = \min \{s(\mu, \mathcal{B}, \mathcal{U}), \underline{\dim}_H(\mu)\}$ and

$$\mathcal{H}^\zeta(\limsup_{n \rightarrow +\infty} U_n) > 0.$$

In particular, for every $\mu \in \mathcal{M}(\mathbb{R}^d)$, one has

$$(10) \quad \dim_H \left(\limsup_{n \rightarrow +\infty} U_n \right) \geq \min \{s(\mu, \mathcal{B}, \mathcal{U}), \underline{\dim}_H(\mu)\}.$$

Remark 2.3. (1) It is easily verified that the lower-bound in Theorem 2.2 equals $-\infty$ if the sequence $(B_n)_{n \in \mathbb{N}}$ is not assumed to be μ -a.c. Consequently, for the previous result to give non trivial information one has to assume that $(B_n)_{n \in \mathbb{N}}$ is μ -a.c. The question is then to give explicit estimates of $s(\mu, \mathcal{B}, \mathcal{U})$ depending on the specificities of $(\mu, \mathcal{B}, \mathcal{U})$.

(2) It is proved in Section 3.3 that $s(\mu, \mathcal{B}, \mathcal{U}) \leq \overline{\dim}_H(\mu)$. This implies that for exact dimensional measures, $\min \{s(\mu, \mathcal{B}, \mathcal{U}), \underline{\dim}_H(\mu)\} = s(\mu, \mathcal{B}, \mathcal{U})$.

(3) The case where μ satisfies $\min \{s(\mu, \mathcal{B}, \mathcal{U}), \underline{\dim}_H(\mu)\} = 0$ could also be treated, but although (10) is still obviously true, some distinction should further be made when investigating the existence of the gauge function. If $\mathcal{H}_\infty^{\mu, s}(U_n) = 0$ for any $n \in \mathbb{N}$, the set $\limsup_{n \rightarrow +\infty} U_n$ could, for instance, be empty. On the other hand, if $(B_n)_{n \in \mathbb{N}}$ is μ -a.c and $s_\mu(B_n, U_n) > 0$ for any $n \in \mathbb{N}$, a gauge function can be constructed in a similar way than in the proof of Theorem 2.2. However the existence of such a gauge function in the case $\min \{s(\mu, \mathcal{B}, \mathcal{U}), \underline{\dim}_H(\mu)\} = 0$ is of little interest for practical applications, and is not treated in this article.

A quite direct, but useful, corollary of Theorem 2.2 is the following:

Corollary 2.4. Let $\mu \in \mathcal{M}(\mathbb{R}^d)$ and $\mathcal{B} = (B_n)_{n \in \mathbb{N}}$ be a μ -a.c. sequence of closed balls of \mathbb{R}^d . Let $\mathcal{U} = (U_n)_{n \in \mathbb{N}}$ be a sequence of open sets such that $U_n \subset B_n$ for all

$n \in \mathbb{N}$, and $0 \leq s \leq \underline{\dim}_H(\mu)$. If $\limsup_{n \rightarrow +\infty} \frac{\log \mathcal{H}_\infty^{\mu, s}(U_n)}{\log \mu(B_n)} \leq 1$, then $s(\mu, \mathcal{B}, \mathcal{U}) \geq s$, so that

$$\dim_H(\limsup_{n \rightarrow +\infty} U_n) \geq s.$$

In the classical case where the sets U_n are shrunk balls of the form B_n^δ (with $\delta \geq 1$), it is convenient to consider the following quantity:

Definition 2.6. Let $\mu \in \mathcal{M}(\mathbb{R}^d)$, $\varepsilon > 0$ and $\mathcal{B} = (B_n)_{n \in \mathbb{N}}$ be a sequence of balls of \mathbb{R}^d . For every $\delta \geq 1$, set

$$(11) \quad t(\mu, \delta, \varepsilon, \mathcal{B}) = \limsup_{n \rightarrow +\infty} \frac{\log(\mathcal{H}_\infty^{\mu, \underline{\dim}_H(\mu) - \varepsilon}(\mathring{B}_n^\delta))}{\log(|B_n^\delta|)}.$$

Then the (μ, δ) -exponent of the sequence \mathcal{B} is defined as

$$(12) \quad t(\mu, \delta, \mathcal{B}) = \lim_{\varepsilon \rightarrow 0} t(\mu, \delta, \varepsilon, \mathcal{B}).$$

It follows from the definitions that $t(\mu, \delta, \mathcal{B})$ exists as a limit, since $\varepsilon \mapsto t(\mu, \delta, \varepsilon, \mathcal{B})$ is monotonic. Moreover, one has $\underline{\dim}_H(\mu) \leq t(\mu, \delta, \mathcal{B})$ (see the proof of Corollary 2.5).

Next result provides an explicit lower bound for the Hausdorff dimension of the limsup of δ -contracted balls; it is a consequence of Corollary 2.4.

Corollary 2.5. Let $\mu \in \mathcal{M}(\mathbb{R}^d)$ and let $\mathcal{B} = (B_n)_{n \in \mathbb{N}}$ be a μ -a.c sequence of closed balls of \mathbb{R}^d . Suppose that $\underline{\dim}_H(\mu) > 0$. For every $\delta \geq 1$, setting

$$s_\delta = \frac{\underline{\dim}_H(\mu)}{\delta} \cdot \frac{\underline{\dim}_H(\mu)}{t(\mu, \delta, \mathcal{B})},$$

one has

$$s(\mu, (B_n)_{n \in \mathbb{N}}, (\mathring{B}_n^\delta)_{n \in \mathbb{N}}) \geq s_\delta,$$

hence

$$\dim_H(\limsup_{n \rightarrow +\infty} B_n^\delta) \geq \dim_H(\limsup_{n \rightarrow +\infty} \mathring{B}_n^\delta) \geq s_\delta.$$

2.3. Dimensional mass transference principle for self-similar measures.

Let us start by recalling the definition of a self-similar measure.

Definition 2.7. A self-similar IFS is a family $S = \{f_i\}_{i=1}^m$ of $m \geq 2$ contracting similarities of \mathbb{R}^d .

Let $(p_i)_{i=1, \dots, m} \in (0, 1)^m$ be a positive probability vector, i.e. $p_1 + \dots + p_m = 1$.

The self-similar measure μ associated with $\{f_i\}_{i=1}^m$ and $(p_i)_{i=1}^m$ is the unique probability measure such that

$$(13) \quad \mu = \sum_{i=1}^m p_i \mu \circ f_i^{-1}.$$

The topological support of μ is the attractor of S , that is the unique non-empty compact set $K \subset X$ such that $K = \bigcup_{i=1}^m f_i(K)$.

The existence and uniqueness of K and μ are standard results [25]. Recall that due to a result by Feng and Hu [19], any self-similar measure is exact dimensional.

The essential Hausdorff content of a self-similar measure μ can be estimated quite precisely.

Theorem 2.6. *Let S be a self-similar IFS of \mathbb{R}^d . Let K be the attractor of S . Let μ be a self-similar measure associated with S . For any $0 \leq s < \dim(\mu)$, there exists a constant $c = c(d, \mu, s) > 0$ depending on the dimension d , μ and s only, such that for any ball $B = B(x, r)$ centered on K and $r \leq 1$, any open set Ω , one has*

$$(14) \quad \begin{aligned} c(d, \mu, s)|B|^s &\leq \mathcal{H}_\infty^{\mu, s}(\overset{\circ}{B}) \leq \mathcal{H}_\infty^{\mu, s}(B) \leq |B|^s \text{ and} \\ c(d, \mu, s)\mathcal{H}_\infty^s(\Omega \cap K) &\leq \mathcal{H}_\infty^{\mu, s}(\Omega) \leq \mathcal{H}_\infty^s(\Omega \cap K). \end{aligned}$$

For any $s > \dim(\mu)$, $\mathcal{H}_\infty^{\mu, s}(\Omega) = 0$.

Note that The system S is not assumed to verify any separation condition.

Moreover, in the special case of the Lebesgue measure restricted to $[0, 1]^d$ (or some cube K), (14) implies that for any $0 \leq s \leq d$, the Lebesgue-essential s dimensional Hausdorff content is strongly equivalent to the usual s -dimensional Hausdorff content (it is even possible to take the constant $c(d, \mu, s)$ in (14), independent of s). In particular, Theorem 2.6 together with Theorem 2.2 in this special case implies the main theorem of Koivusalo and Rams, [28, Theorem 3.2], recalled below.

Theorem 2.7 ([28]). *Let $(B_n)_{n \rightarrow +\infty}$ be a sequence of balls of $[0, 1]^d$ verifying $|B_n| \rightarrow 0$ and $\mathcal{L}^d(\limsup_{n \rightarrow +\infty} B_n) = 1$.*

Let $(U_n)_{n \in \mathbb{N}}$ be a sequence of open sets satisfying $U_n \subset B_n$. For any $0 \leq s \leq d$ such that, for all $n \in \mathbb{N}$ large enough, $\mathcal{H}_\infty^s(U_n) \geq \mathcal{L}^d(B_n)$, it holds that

$$(15) \quad \dim_H(\limsup_{n \rightarrow +\infty} U_n) \geq s.$$

As a consequence of Theorem 2.6 and Corollary 2.4, one gets

Corollary 2.8. *Let $\mu \in \mathcal{M}(\mathbb{R}^d)$ be a self-similar measure and $\mathcal{B} = (B_n)_{n \in \mathbb{N}}$ be a μ -a.c. sequence of closed balls of \mathbb{R}^d centered in $\text{supp}(\mu)$. Let $\mathcal{U} = (U_n)_{n \in \mathbb{N}}$ be a sequence of open sets such that $U_n \subset B_n$ for all $n \in \mathbb{N}$, and $0 \leq s \leq \dim(\mu)$. If, for $n \in \mathbb{N}$ large enough, $\mathcal{H}_\infty^{\mu, s}(U_n) \geq \mu(B_n)$, then*

$$\dim_H(\limsup_{n \rightarrow +\infty} U_n) \geq s.$$

One also emphasizes that in the case of a self-similar measure, conversely, any $s \geq 0$ such that $\mathcal{H}_\infty^{\mu, s}(U_n) \leq \mu(B_n)$ for every n large enough is an upper-bound for $\dim_H(\limsup_{n \rightarrow +\infty} U_n)$ if \mathcal{B} verifies that, for any $p \in \mathbb{N}$, the balls B_n with $|B_n| \approx 2^{-p}$ does not overlap too much (which is the case for instance the case of the dynamic balls defined in Theorem 2.14 in the case of a real IFS satisfying the exponential separation and with similarity dimension smaller than 1). More precisely, in the companion paper of the present article, [14], the following result is proved.

Theorem 2.9 ([14]). *Let $\mu \in \mathcal{M}(\mathbb{R}^d)$ be a self-similar measure, K its support and $(B_n)_{n \rightarrow +\infty}$ be a weakly redundant sequence of balls of \mathbb{R}^d (see [7, Definition 1.5]) verifying $|B_n| \rightarrow 0$ and, for any $n \in \mathbb{N}$, $B_n \cap K \neq \emptyset$. Let $(U_n)_{n \in \mathbb{N}}$ be a sequence of open sets satisfying $U_n \subset B_n$. For any $0 \leq s < \dim(\mu)$ such that, for all large enough $n \in \mathbb{N}$, $\mathcal{H}_\infty^{\mu, s}(U_n) \leq \mu(B_n)$, it holds that*

$$(16) \quad \dim_H(\limsup_{n \rightarrow +\infty} U_n) \leq s.$$

Combining Theorem 2.2 and Corollary 2.5 with Theorem 2.6 and Proposition 2.1 yields the following consequence for self-similar measures.

Theorem 2.10. *Let S be a self-similar IFS of \mathbb{R}^d with attractor K and μ be a self-similar measure associated with S . Let $(B_n)_{n \in \mathbb{N}}$ be a sequence of closed balls centered on K , such that $\lim_{n \rightarrow +\infty} |B_n| = 0$.*

- (1) *Suppose that $(B_n)_{n \in \mathbb{N}}$ is μ -a.c. Then $t(\mu, \delta, (B_n)_{n \in \mathbb{N}}) \leq \dim(\mu)$; consequently $s(\mu, (B_n)_{n \in \mathbb{N}}, (\mathring{B}_n^\delta)_{n \in \mathbb{N}}) \geq \frac{\dim(\mu)}{\delta}$ and there exists a gauge function ζ such that $\lim_{r \rightarrow 0^+} \frac{\log(\zeta(r))}{\log(r)} \geq \frac{\dim(\mu)}{\delta}$ and $\mathcal{H}^\zeta(\limsup_{n \rightarrow \infty} \mathring{B}_n^\delta) > 0$. In particular*

$$(17) \quad \dim_H(\limsup_{n \rightarrow +\infty} \mathring{B}_n^\delta) \geq \frac{\dim(\mu)}{\delta}.$$

- (2) *Suppose that $\mu(\limsup_{n \rightarrow +\infty} B_n) = 1$. Then, (17) still holds but the existence of the gauge function is not ensured. Furthermore if μ is doubling, then $(B_n)_{n \in \mathbb{N}}$ is μ -a.c, so that the conclusion of item (1) holds.*

Remark 2.11. *Since no separation condition is assumed about the system S , Theorem 2.10 implies [7, Theorem 1.6] in the special case where the sequence of measures $(\mu_p)_{p \in \mathbb{N}}$ is constant and equal to some self-similar measure with open set condition μ and the sequence of contraction ratio $(\delta_p)_{p \in \mathbb{N}}$ is constant as well.*

Corollary 2.4 and Theorem 2.6 make it possible to deal with more general open sets $(U_n)_{n \geq 1}$ than contracted balls (\mathring{B}_n^δ) , if one is able to compare efficiently the s -dimensional Hausdorff contents of the sets $U_n \cap K$ with a power of $|B_n|$. It is then convenient to assume that K is the closure of its interior. Here is an example.

Let $1 \leq \tau_1 \leq \dots \leq \tau_d$ be d real numbers and $\tau = (\tau_1, \dots, \tau_d)$. One starts by defining a family of rectangles of \mathbb{R}^d associated with τ .

Definition 2.8. *Let $1 \leq \tau_1 \leq \dots \leq \tau_d$ and $\tau = (\tau_1, \dots, \tau_d)$. For any $x = (x_i)_{1 \leq i \leq d} \in \mathbb{R}^d$ and $r > 0$, the τ -rectangle centered in x and associated with r is defined by*

$$(18) \quad R_\tau(x, r) = \prod_{i=1}^d [x_i - \frac{r^{\tau_i}}{2}, x_i + \frac{r^{\tau_i}}{2}].$$

The theorem we get is the following:

Theorem 2.12. *Let S be a self-similar IFS of \mathbb{R}^d such that the attractor K of S satisfies $K = \bar{K}$. Let μ be a self-similar measure associated with S . Let $1 \leq \tau_1 \leq \dots \leq \tau_d$, $\tau = (\tau_1, \dots, \tau_d)$ and $(B_n := B(x_n, r_n))_{n \in \mathbb{N}}$ be a sequence of balls of \mathbb{R}^d satisfying $r_n \rightarrow 0$ and $\mu(\limsup_{n \rightarrow +\infty} B_n) = 1$. Define $R_n = \mathring{R}_\tau(x_n, r_n)$. Then*

$$(19) \quad \dim_H(\limsup_{n \rightarrow +\infty} R_n) \geq \min_{1 \leq i \leq d} \left\{ \frac{\dim(\mu) + \sum_{1 \leq j \leq i} \tau_i - \tau_j}{\tau_i} \right\}.$$

Remark 2.13. (1) *Since $(\tau_1, \dots, \tau_d) \mapsto \min_{1 \leq i \leq d} \left\{ \frac{\dim(\mu) + \sum_{1 \leq j \leq i} \tau_i - \tau_j}{\tau_i} \right\}$ is continuous, the result stands for the sequence of closed rectangles as well.*

(2) *One may also apply any rotation to the shrunk rectangles, this would not change the bound (since Hausdorff contents are invariant by rotation).*

(3) *Theorem 2.12 extends the results of [13], where the measure was quasi-Bernoulli or verifying the open set condition, and supported on $[0, 1]^d$.*

(4) When $K = \overline{\overset{\circ}{K}}$ it is easy to compute $\mathcal{H}_\infty^s(R_n \cap K)$. Without this assumption, the conclusion of Theorem 2.12 fails. Indeed, in general, no formula involving only the dimension of the measure and the contraction ratio can be accurate. For instance, consider a self-similar measure in \mathbb{R}^2 carried by a line D and a sequence $(B_n)_{n \in \mathbb{N}}$ of balls centered on the attractor K and verifying $\mu(\limsup_{n \rightarrow +\infty} B_n) = 1$. Then, consider the sequence of rectangles R_n with side-length $|B_n|^{\tau_1} \times |B_n|^{\tau_2}$, $1 \leq \tau_1 \leq \tau_2$ and where the largest side (of side-length $|B_n|^{\tau_1}$) is in the direction of D . In this case, Theorem 2.10 yields the lower-bound $\dim_H(\limsup_{n \rightarrow +\infty} R_n) \geq \frac{\dim_H(\mu)}{\tau_1}$. Then if \widehat{R}_n are the rectangles R_n rotated by $\frac{\pi}{2}$, Theorem 2.10 gives that $\dim_H(\limsup_{n \rightarrow +\infty} \widehat{R}_n) \geq \frac{\dim_H(\mu)}{\tau_2}$. Moreover, under additional conditions, these lower bounds are equalities.

2.4. Application to self-similar dynamical coverings. We deal with points well approximable by orbits under an IFS with no exact overlap and satisfying the condition introduced by Barral and Feng ([3]) of being *dimension regular* with similarity dimension less than d , after Hochman's work ([24]).

Definition 2.9. Let $S = \{f_1, \dots, f_m\}$ be a self-similar IFS of \mathbb{R}^d . Denote by $0 < c_1, \dots, c_m < 1$ the contraction ratio of f_1, \dots, f_m . The system S is said to be *dimension regular* if, for any probability vector (p_1, \dots, p_m) , the self-similar measure associated with S and the probability vector (p_1, \dots, p_m) verifies

$$\dim(\mu) = \min \left\{ d, \frac{\sum_{1 \leq i \leq m} p_i \log(p_i)}{\sum_{1 \leq i \leq m} p_i \log(c_i)} \right\}.$$

Denoting by $\dim(S)(K)$ the unique real number s satisfying $\sum_{i=1}^m c_i^s = 1$, one has $\dim_H(K) = \min \{d, \dim(S)(K)\}$.

Some notation useful when dealing with IFS are introduced now. These notations will be used repeatedly throughout this article.

Let $S = \{f_1, \dots, f_m\}$ be a self-similar IFS, $0 < c_1, \dots, c_m < 1$ the associated contraction ratios, and K the attractor of S . Let (p_1, \dots, p_m) be a probability vector with positive entries, μ the self-similar associated with S and (p_1, \dots, p_m) .

Let $\Lambda = \{1, \dots, m\}$ and $\Lambda^* = \bigcup_{k \geq 0} \Lambda^k$. For $k \geq 0$ and $\underline{i} := (i_1, \dots, i_k) \in \Lambda^k$, define

$$c_{\underline{i}} = c_{i_1} \cdots c_{i_k}, \quad f_{\underline{i}} = f_{i_1} \circ \cdots \circ f_{i_k},$$

$$\Lambda^{(k)} = \{\underline{i} = (i_1, \dots, i_s) \in \Lambda^* : c_{i_s} 2^{-k} < c_{\underline{i}} \leq 2^{-k}\}.$$

Theorem 2.14. Let $S = \{f_1, \dots, f_m\}$ be a dimension regular self-similar IFS with contraction ratio $0 < c_1, \dots, c_m < 1$ and such that the attractor K verifies $\dim(S)(K) = \dim_H(K)$. For any $x \in K$, for any $\delta \geq 1$,

$$(20) \quad \dim_H \left(\limsup_{\underline{i} \in \Lambda^*} B(f_{\underline{i}}(x), c_{\underline{i}}^\delta) \right) = \frac{\dim_H(K)}{\delta}.$$

This result extends some of the results obtained under the open set condition in [1] and [2]. Here are some examples of self-similar IFS satisfying the condition of Theorem 2.14 and some example of IFS for which the conclusion fails.

- Any self-similar IFS satisfying the open set condition. This case was already established by Hill and Velani in [23].

- Any self-similar IFS on \mathbb{R} satisfying the exponential separation condition (due to a result of Hochman [24]) or with algebraic contraction ratios, no exact overlaps $\dim(S) \leq 1$ (due to a result of Rapaport [31]).
- Let $0 < c_1, \dots, c_m < 1$ such that

$$\sum_{1 \leq i \leq m} c_i \leq 1.$$

For Lebesgue almost every $(a_1, \dots, a_d) \in \mathbb{R}^d$, the IFS

$$S = \{f_1(\cdot) = c_1 \cdot + a_1, \dots, f_m(\cdot) = c_m \cdot + a_m\}$$

is also dimension regular (due to a result of Hochman [24]).

- There exists some IFS satisfying $\dim(S) > d$ (so that $\dim_H(K) \neq \dim(S)$) and for which

$$\dim_H W(x_0, \delta) = d$$

for every $\delta \in [1, \delta_0]$ for some $\delta_0 > 1$ (an explicit example can be found in [2]).

Let us also mention that it is known that it is equivalent for a self-similar IFS S to satisfy the open set condition and to satisfy $\mathcal{H}^{\dim(S)}(K) > 0$, where K is the attractor of S (see [32]). In particular, each of these examples which satisfies $\dim_H(K) = \dim(S)$ but not the open set condition also satisfies that $\mathcal{H}^{\dim_H(K)}(K) = 0$. In consequence, in such cases, the classical mass transference principle from Beresnevich and Velani [8] cannot be applied.

2.5. A result motivated by a question of Mahler.

Let $\mathcal{Q} = \left\{ B\left(\frac{p}{q}, q^{-2}\right) \right\}_{q \in \mathbb{N}^*, 0 \leq p \leq q}$. Remember the following result in Diophantine approximation [27]:

- $\limsup_{B \in \mathcal{Q}} B = [0, 1]$.
- For any $\delta \geq 1$, $\dim_H(\limsup_{B \in \mathcal{Q}} B^\delta) = \frac{1}{\delta}$.

Unlike in the case of the points in $[0, 1]$, the approximation by rational numbers of elements of the middle third Cantor set $K_{1/3}$ set is not well understood yet. This question was raised by Mahler, and only some partial results are known (see [8], [7],[10]). Here we consider the set $K_{1/3}^{(0)}$ of points in $[0, 1]$ having an asymptotic lower frequency of appearance of the digit 1 in basis 3 equal to 0. This set contains $K_{1/3}$ and has the same Hausdorff dimension as $K_{1/3}$. We compute the Hausdorff dimension of sets of points in $K_{1/3}^{(0)}$ which are well approximable by rational numbers.

To describe more precisely the problem, let $S = \{f_1, f_2, f_3\}$ where f_1, f_2 and f_3 are the contracting affine maps of \mathbb{R} defined by $f_0(x) = \frac{1}{3}x$, $f_1(x) = \frac{1}{3}x + \frac{1}{3}$ and $f_2(x) = \frac{1}{3}x + \frac{2}{3}$. The attractor of S is $[0, 1]$. Let $\Lambda = \{0, 1, 2\}$.

The shift operation on the symbolic space $\Lambda^{\mathbb{N}}$ is defined by σ . The canonical projection from $\Lambda^{\mathbb{N}}$ to $[0, 1]$ is the mapping

$$(22) \quad \pi : x = (x_n)_{n \in \mathbb{N}} \mapsto \lim_{n \rightarrow +\infty} f_{(x_1, \dots, x_n)}(0).$$

The set $K_{1/3}$ is the attractor of $\{f_0, f_2\}$ and also the image by canonical projection of $\{0, 2\}^{\mathbb{N}}$.

Definition 2.10. Let $\phi : \Lambda^{\mathbb{N}} \rightarrow \{0, 1\}$ defined by $\begin{cases} \phi(x) = 1 & \text{if } x_1 = 1 \\ \phi(x) = 0 & \text{if } x_1 = 0 \text{ or } 2. \end{cases}$

and

$$K_{1/3}^{(0)} = \pi \left(\left\{ x \in \Lambda^{\mathbb{N}} : \liminf_{k \rightarrow +\infty} \frac{S_k \phi(x)}{k} = 0 \right\} \right),$$

where $(S_k \phi)_{k \in \mathbb{N}}$ stands for the sequence of Birkhoff sums of ϕ .

It is proved in [18] that $\dim_H K_{1/3}^{(0)} = \frac{\log 2}{\log 3}$ ($= \dim_H K_{1/3}$).

As we will study the approximation of numbers of $K_{1/3}^{(0)}$, one introduces the irrationality exponent of a number.

Definition 2.11. Let $x \in \mathbb{R}$, we define the irrationality of x $\xi(x)$ by

$$\xi(x) = \sup \left\{ \delta : x \in B\left(\frac{p}{q}, \frac{1}{q^\delta}\right) \text{ for infinitely many } (p, q) \in \mathbb{Z} \times \mathbb{N}^* \right\}.$$

Let us state the main results of this subsection.

Theorem 2.15. For every $\delta \geq 1$,

$$(23) \quad \begin{cases} \dim_H \left(\{x : \xi(x) \geq 2\delta\} \cap K_{1/3}^{(0)} \right) = \frac{\log 2}{\log 3} & \text{if } 1 \leq \delta \leq \frac{\log 3}{\log 2}, \\ \dim_H \left(\{x : \xi(x) = 2\delta\} \cap K_{1/3}^{(0)} \right) = \frac{1}{2\delta} & \text{if } \delta \geq \frac{\log 3}{\log 2}. \end{cases}$$

Observe that a saturation phenomenon occurs : for $1 \leq \delta \leq \frac{\log 3}{\log 2}$

$$\dim_H \left(\{x : \xi(x) \geq 2\delta\} \cap K_{1/3}^{(0)} \right) = \frac{\log 2}{\log 3}.$$

We also conjecture that for $\delta \in [1, \frac{\log 3}{\log 2}]$, (23) should hold for the level set $\{x : \xi(x) = 2\delta\}$.

In Section 3, the general mass transference principle, Theorem 2.2, is proved as well as Corollary 2.5. Section 4 gives estimations of essential contents in the self-similar case and Theorem 2.6 is proved.

Section 5 contains three applications to the main result, Theorem 2.2. More precisely, the mass transference principles for self-similar measures, Theorem 2.10 and Theorem 2.12, are proved in the first sub-section. The second sub-section treats the case of self-similar dynamical coverings for dimension regular IFS with similarity dimension less than d , e.g, Theorem 2.14 is proved. In the last sub-section, one gives an application in Diophantine approximation, Theorem 2.15 is proved.

3. PROOF OF THEOREM 2.2

3.1. Preliminary facts. We gather in this subsection a series of results on which we will base the proof of Theorem 2.2.

The following lemma, which is a version of Besicovitch covering Lemma, as well as the subsequent one, both established in [14], will be used several times.

Lemma 3.1. *For any $0 < v \leq 1$ there exists $Q_{d,v} \in \mathbb{N}^*$, a constant depending only on the dimension d and v , such that for every bounded subset $E \subset \mathbb{R}^d$, for every set $\mathcal{F} = \{B(x, r(x)) : x \in E, r(x) > 0\}$, there exists $\mathcal{F}_1, \dots, \mathcal{F}_{Q_{d,v}}$ finite or countable sub-families of \mathcal{F} such that:*

- $\forall 1 \leq i \leq Q_{d,v}$, $L \neq L' \in \mathcal{F}_i$, one has $\frac{1}{v}L \cap \frac{1}{v}L' = \emptyset$.
- E is covered by the families \mathcal{F}_i , i.e.

$$(24) \quad E \subset \bigcup_{1 \leq i \leq Q_{d,v}} \bigcup_{L \in \mathcal{F}_i} L.$$

Lemma 3.2 ([14]). *Let $0 < v \leq 1$, $\mathcal{B} = (B_n)_{n \in \mathbb{N}}$ a family of balls, and B a ball such that*

- (i) $\forall n \geq 1$, $|B_n| \geq \frac{1}{2}|B|$,
- (ii) $\forall n_1 \neq n_2 \geq 1$, $vB_{n_1} \cap vB_{n_2} = \emptyset$.

Then B intersects less than $Q_{d,v}$ elements of \mathcal{B} , where $Q_{d,v}$ can be taken equal to the constant considered in Lemma 3.1.

The following lemma will also be useful later on and is also proved in [14].

Lemma 3.3. *Let \mathcal{L} be a family of pairwise disjoint balls satisfying $\sup_{L \in \mathcal{L}} |L| < +\infty$. Then, for any $v \geq 1$, there exists sub-families $\mathcal{L}_1, \dots, \mathcal{L}_{Q_{d,v}}$ (where $Q_{d,v}$ is the constant of the same name in Lemma 3.1) of \mathcal{L} such that $\mathcal{L} = \bigcup_{1 \leq i \leq Q_{d,v}} \mathcal{F}_i$ and for any $L \neq L' \in \mathcal{L}_i$, $vL \cap vL' = \emptyset$.*

Recall the following version of Frostman Lemma, due to Carleson.

Proposition 3.4 ([11]). *Let $s \geq 0$. There is a constant $\kappa_d > 0$ depending only on the dimension d such that for any bounded set $E \subset \mathbb{R}^d$ with $\mathcal{H}_\infty^s(E) > 0$, there exists a probability measure supported by E , that we denote by m_E^s , such that*

$$(25) \quad \text{for every ball } B(x, r), \quad m_E^s(B(x, r)) \leq \kappa_d \frac{r^s}{\mathcal{H}_\infty^s(E)}.$$

For $s \geq 0$ and $E \subset \mathbb{R}^d$, a bounded subset such that $\mathcal{H}_\infty^s(E) > 0$, m_E^s will always denote such a measure associated with a (fixed) constant κ_d .

In the next two lemmas, the choice of the interval $[5, 6]$ is convenient to take enough space between the shrunk balls involved in the construction elaborated in Section 3.2.

Lemma 3.5. *Let $t \in (5, 6)$, $m \in \mathcal{M}(\mathbb{R}^d)$, and $\varepsilon > 0$. Let $x \in \mathbb{R}^d$ be such that $\overline{\dim}_{\text{loc}}(m, x) \leq \beta$. Let $C_{\beta, \varepsilon} = \frac{1}{2}6^{-\frac{\beta}{2\varepsilon}}$. There exists an integer n_x such that for every $n \geq n_x$,*

$$(26) \quad \frac{\#\{0 \leq k \leq n-1 : m(B(x, t^{-k-1})) \geq C_{\beta, \varepsilon} m(B(x, t^{-k}))\}}{n} \geq 1 - \varepsilon.$$

Previous lemma is a slight extension of result by Käenmäki [12, Lemma 2.2], which shows such a property at m -almost every point (where one has necessarily $\overline{\dim}_{\text{loc}}(m, x) \leq d$), and uses t integer (a choice that we could make).

Thus, points with a given local dimension with respect to a measure m are for most scales “locally doubling”.

Proof. Observe first that if for a constant $0 < C \leq 1$ and some integer $n \in \mathbb{N}$ one has

$$\frac{\#\{1 \leq k \leq n : m(B(x, t^{-k-1})) \geq Cm(B(x, t^{-k}))\}}{n} \leq 1 - \varepsilon,$$

then there necessarily exist $N = \lfloor (n-1)\varepsilon \rfloor$ integers $0 < k_1 < \dots < k_N < n$ such that for every $1 \leq i \leq N$, $m(B(x, t^{-k_i-1})) \leq Cm(B(x, t^{-k_i}))$.

In particular, writing $k_{N+1} = n$ and $k_0 = 0$, this implies that

$$\begin{aligned} m(B(x, t^{-n})) &= \prod_{i=0}^N \frac{m(B(x, t^{-k_{i+1}}))}{m(B(x, t^{-k_i}))} \leq \prod_{i=0}^N \frac{m(B(x, t^{-k_i-1}))}{m(B(x, t^{-k_i}))} \leq C^N \\ &\leq C^{(n-1)\varepsilon} \leq C^{n\varepsilon/2} = (t^{-n})^{\varepsilon \frac{-\log(C)}{2\log(t)}}. \end{aligned}$$

The inequality $C^{(n-1)\varepsilon} \leq C^{n\varepsilon/2}$ occurs when n is large enough. Recalling that $\overline{\dim}_{\text{loc}}(m, x) \leq \beta$, if this happens for infinitely many n , one should have

$$\beta \geq \limsup_{r \rightarrow 0^+} \frac{\log m(B(x, r))}{\log r} \geq \varepsilon \frac{-\log(C)}{2\log(t)},$$

which is equivalent to $C \geq t^{-\frac{\beta}{2\varepsilon}}$.

Setting $C_{\varepsilon, \beta} = \frac{1}{2}6^{-\frac{\beta}{2\varepsilon}}$, one concludes that there exists n_x such that for every $n \geq n_x$, one necessarily has

$$\frac{\#\{0 \leq k \leq n-1 : m(B(x, t^{-k-1})) \geq C_{\varepsilon, \beta} m(B(x, t^{-k}))\}}{n} \geq 1 - \varepsilon,$$

hence the result. \square

Lemma 3.6. *Let m and μ be two elements of $\mathcal{M}([0, 1])$, $\beta \geq 0$ and $\varepsilon > 0$. For every $x \in \mathbb{R}^d$ verifying $\overline{\dim}_{\text{loc}}(m, x) \leq \beta$, there exists $\rho_x > 0$ and $t_x \in (5, 6)$ so that for all $0 < r \leq \rho_x$ there exists $r \leq r' \leq r^{1-\varepsilon}$ such that*

$$(27) \quad m(B(x, r'/t_x)) \geq C_{\beta, \frac{\varepsilon}{2}} m(B(x, r')) \text{ and } \mu(\partial B(x, r'/t_x)) = 0.$$

Proof. Consider $x \in \mathbb{R}^d$ such that $\overline{\dim}_{\text{loc}}(m, x) \leq \beta$.

We apply Lemma 3.5 to x and the measure m , and for an arbitrary $t \in [5, 6]$ and $\varepsilon' = \frac{\varepsilon}{2}$: for $n \geq n_x$, there must be an integer n' such that $n(1 - \varepsilon) \leq n' \leq n$ and $m(B(x, t^{-n'+1})) \geq C_{\beta, \frac{\varepsilon}{2}} m(B(x, t^{-n'}))$.

Let $\rho_x = \min\{t^{-n_x-1}, t^{-\frac{1}{\varepsilon}}\}$. For $r \in (0, \rho_x]$, let n be the integer such that $t^{-n-1} < r \leq t^{-n}$. The previous claim yields an integer $n' \in [n(1 - \frac{\varepsilon}{2}), n]$ such that $m(B(x, t^{-n'})) \geq C_{\beta, \frac{\varepsilon}{2}} m(B(x, t^{-n'+1}))$. Also,

$$r \leq r' = t^{-n'+1} \leq t^{1-(1-\frac{\varepsilon}{2})n} = t^2 \cdot t^{-n-1} \cdot t^{\frac{\varepsilon}{2}n} \leq t^2 \cdot r \cdot r^{-\frac{\varepsilon}{2}} \leq r^{1-\varepsilon}.$$

Consequently,

$$m(B(x, r'/t)) \geq C_{\beta, \frac{\varepsilon}{2}} m(B(x, r')).$$

The desired conclusion holds if we choose $t_x \in (5, t)$ such that $\mu(\partial B(x, r'/t_x)) = 0$. \square

The previous lemma will be used in the case $\beta = d$ in our proof the main theorem (see step 2 of the construction in Section 3.2).

Next, we introduce some sets associated to a given element of $\mathcal{M}(\mathbb{R}^d)$, which will play a natural role in our construction.

Definition 3.1. Let $\beta \geq \alpha \geq 0$ be real numbers, $m \in \mathcal{M}(\mathbb{R}^d)$, and $\varepsilon, \rho > 0$ two positive real numbers. Then define

$$(28) \quad \widetilde{E}_m^{[\alpha, \beta], \rho, \varepsilon} = \left\{ x \in \mathbb{R}^d : \underline{\dim}_{\text{loc}}(m, x) \in [\alpha, \beta] \text{ and } \forall r \leq \rho, m(B(x, r)) \leq r^{\underline{\dim}_{\text{loc}}(m, x) - \varepsilon} \right\}$$

and

$$(29) \quad E_m^{[\alpha, \beta], \rho, \varepsilon} = \left\{ x \in \widetilde{E}_m^{[\alpha, \beta], \rho, \varepsilon} : \forall r \leq \rho, \frac{3}{4}m(B(x, r)) \leq m(B(x, r) \cap \widetilde{E}_m^{[\alpha, \beta], \rho, \varepsilon}) \right\}$$

Notice that, for every $0 < \rho < \rho'$, one has $E_m^{[\alpha, \beta], \rho', \varepsilon} \subset E_m^{[\alpha, \beta], \rho, \varepsilon}$.

Definition 3.2. Let $\beta \geq \alpha \geq 0$ be real numbers, $m \in \mathcal{M}(\mathbb{R}^d)$, and $\varepsilon > 0$. Define

$$(30) \quad E_m^{[\alpha, \beta], \varepsilon} = \bigcup_{n \geq 1} E_m^{[\alpha, \beta], \frac{1}{n}, \varepsilon}.$$

Proposition 3.7. For every $m \in \mathcal{M}(\mathbb{R}^d)$, every $\beta \geq \alpha \geq 0$ and $\varepsilon > 0$,

$$(31) \quad m(E_m^{[\alpha, \beta], \varepsilon}) = m(\{x : \underline{\dim}_{\text{loc}}(m, x) \in [\alpha, \beta]\}).$$

Notice that, for every $0 < \rho' < \rho$, one has $E_m^{[\alpha, \beta], \rho, \varepsilon} \subset E_m^{[\alpha, \beta], \rho', \varepsilon}$.

These sets play a key role in the proofs of Theorem 2.2 .

Proof. One first recalls the following result.

Lemma 3.8. [9] Let $m \in \mathcal{M}(\mathbb{R}^d)$ and A be a Borel set with $m(A) > 0$. For every $r > 0$, set

$$(32) \quad A(r) = \left\{ x \in A : \forall \tilde{r} \leq r, m(B(x, \tilde{r}) \cap A) \geq \frac{3}{4}m(B(x, \tilde{r})) \right\}$$

Then

$$(33) \quad m\left(\bigcup_{r > 0} A(r)\right) = m(A).$$

Note that it is clear from Definition 2.2 that

$$\{x : \underline{\dim}_{\text{loc}}(m, x) \in [\alpha, \beta]\} = \bigcup_{\rho > 0} \widetilde{E}_m^{[\alpha, \beta], \rho, \varepsilon}.$$

Let $\varepsilon' > 0$. By Definition 2.2, there exists $\rho_{\varepsilon'}$ small enough so that

$$(34) \quad m(\widetilde{E}_m^{[\alpha, \beta], \rho_{\varepsilon'}, \varepsilon}) \geq (1 - \varepsilon')m(\{x : \underline{\dim}_{\text{loc}}(m, x) \in [\alpha, \beta]\}).$$

By Lemma 3.8 (and the notations therein) applied to $\widetilde{E}_m^{[\alpha, \beta], \rho_{\varepsilon'}, \varepsilon}$, there exists $\tilde{\rho}_{\varepsilon'}$ such that

$$(35) \quad m(\widetilde{E}_m^{[\alpha, \beta], \rho_{\varepsilon'}, \varepsilon}(\tilde{\rho}_{\varepsilon'})) \geq (1 - \varepsilon')m(\widetilde{E}_m^{[\alpha, \beta], \rho_{\varepsilon'}, \varepsilon}).$$

Finally for $\rho = \min\{\rho_{\varepsilon'}, \tilde{\rho}_{\varepsilon'}\}$, by Definition 3.1 and (32), one has $(\widetilde{E}_m^{[\alpha, \beta], \rho_{\varepsilon'}, \varepsilon})_{\tilde{\rho}_{\varepsilon'}} \subset E_m^{[\alpha, \beta], \rho, \varepsilon}$, so that, by (34) and (35)

$$\begin{aligned} m(E_m^{[\alpha, \beta], \rho, \varepsilon}) &\geq m((\widetilde{E}_m^{[\alpha, \beta], \rho_{\varepsilon'}, \varepsilon}(\tilde{\rho}_{\varepsilon'}))) \geq (1 - \varepsilon')m(E_m^{[\alpha, \beta], \varepsilon}) \\ &\geq (1 - \varepsilon')^2 m(\{x : \underline{\dim}_{\text{loc}}(m, x) \in [\alpha, \beta]\}). \end{aligned}$$

In particular

$$m(\{x : \underline{\dim}_{\text{loc}}(m, x) \in [\alpha, \beta]\}) \geq m(E_m^{[\alpha, \beta], \varepsilon}) \geq (1 - \varepsilon')^2 m(\{x : \underline{\dim}_{\text{loc}}(m, x) \in [\alpha, \beta]\}).$$

Letting $\varepsilon' \rightarrow 0$ proves the result. \square

Corollary 3.9. *For every $m \in \mathcal{M}(\mathbb{R}^d)$, for $\alpha = \underline{\dim}_H(m)$ and $\beta = \overline{\dim}_H(m)$, for any $\varepsilon > 0$, one has*

$$(36) \quad m(E_m^{[\alpha, \beta], \varepsilon}) = 1.$$

3.2. Construction of the Cantor set and the measure. Recall that μ is a probability measure on \mathbb{R}^d , and that $\mathcal{B} = (B_n := B(x_n, r_n))_{n \in \mathbb{N}}$ is a μ -a.c sequence of balls of \mathbb{R}^d with $\lim_{n \rightarrow +\infty} r_n = 0$. Fix $\mathcal{U} = (U_n)_{n \in \mathbb{N}}$ a sequence of open sets satisfying $U_n \subset B_n$ for every $n \in \mathbb{N}$.

Set $\alpha = \underline{\dim}_H(\mu)$, and assume that $\min \{s(\mu, \mathcal{B}, \mathcal{U}), \alpha\} > 0$.

Our goal is to construct a gauge function $\zeta : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ such that $\lim_{r \rightarrow 0^+} \frac{\log \zeta(r)}{\log r} = \min \{s(\mu, \mathcal{B}, \mathcal{U}), \underline{\dim}_H(\mu)\}$ as well as $\eta \in \mathcal{M}(\mathbb{R}^d)$ supported on $\limsup_{n \rightarrow \infty} U_n$ such that for all $r \in (0, 1]$ and $x \in \mathbb{R}^d$ one has $\eta(B(x, r)) \leq \zeta(2r)$.

Let $(\varepsilon_k)_{k \in \mathbb{N}}$ be a sequence decreasing to 0 and such that $\varepsilon_1 < s(\mu, \mathcal{B}, \mathcal{U})$. For $k \geq 0$, set

$$(37) \quad s_k = \min \{s(\mu, \mathcal{B}, \mathcal{U}), \alpha\} - \varepsilon_k.$$

Along the construction of ζ , we only use that $s_k < s(\mu, \mathcal{B}, \mathcal{U})$ and the fact that $s_k < \alpha$ is used at the end of our analysis (see equation (73)).

Step 1. We need the following lemma.

Lemma 3.10 ([14]). *Let $\mu \in \mathcal{M}(\mathbb{R}^d)$ and $\mathcal{B} = (B_n := B(x_n, r_n))_{n \in \mathbb{N}}$ be a μ -a.c sequence of balls of \mathbb{R}^d with $\lim_{n \rightarrow +\infty} r_n = 0$.*

Then for every open set Ω and every integer $g \in \mathbb{N}$, there exists a subsequence $(B_{(n)}^{(\Omega)}) \subset \{B_n\}_{n \geq g}$ such that:

- (1) $\forall n \in \mathbb{N}, B_{(n)}^{(\Omega)} \subset \Omega$,
- (2) $\forall 1 \leq n_1 \neq n_2, B_{(n_1)}^{(\Omega)} \cap B_{(n_2)}^{(\Omega)} = \emptyset$,
- (3) $\mu \left(\bigcup_{n \geq 1} B_{(n)}^{(\Omega)} \right) = \mu(\Omega)$.

In addition, there exists an integer N_Ω such that for the balls $(B_{(n)}^{(\Omega)})_{n=1, \dots, N_\Omega}$, the conditions (1) and (2) are realized, and (3) is replaced by $\mu \left(\bigcup_{n=1}^{N_\Omega} B_{(n)}^{(\Omega)} \right) \geq \frac{3}{4} \mu(\Omega)$.

The last part of Lemma 3.10 simply follows from item (3) and the σ -additivity of μ .

Using Lemma 3.10 with, $(B_n)_{n \in \mathcal{N}_\mu(\mathcal{B}, \mathcal{U}, s_1)}$ (which is μ -a.c since $s_1 < s(\mu, \mathcal{B}, \mathcal{U})$), $g = 0$ and $\Omega = \mathbb{R}^d$, one finds integers N_1 and $n_1 < \dots < n_{N_1} \in \mathcal{N}_\mu(\mathcal{B}, \mathcal{U}, s_1)$ such that :

- (i) $\forall 1 \leq i \leq N_1, B_{n_i} \cap B_{n_j} = \emptyset$,
- (ii) $\mu \left(\bigcup_{1 \leq i \leq N_1} B_{n_i} \right) \geq \frac{1}{2}$.

By Lemma 3.3 applied to $\{B_{n_i}\}_{1 \leq i \leq N_1}$ and $v = 4$, the balls $\{B_{n_i}\}_{1 \leq i \leq N_1}$ can be sorted in $Q_{d,4}$ families of balls $\mathcal{L}_1, \dots, \mathcal{L}_{Q_{d,4}}$ such that

- for any $1 \leq i \leq Q_{d,4}$, any $L \neq L' \in \mathcal{L}_i, 4L \cap 4L' = \emptyset$,
- $\bigcup_{1 \leq i \leq Q_{d,4}} \mathcal{L}_i = \{B_{n_i}\}_{1 \leq i \leq N_1}$.

At least one of these families, \mathcal{L}_{i_0} , must satisfy

$$\mu \left(\bigcup_{L \in \mathcal{L}_{i_0}} L \right) \geq \frac{1}{2Q_{d,4}}.$$

In particular, if one must rename the balls of the family \mathcal{L}_{i_0} , we can assume that the family $\{B_{n_i}\}_{1 \leq i \leq N_1}$ satisfies

$$(i') : \text{ for any } 1 \leq i < j \leq N_1, 4B_{n_i} \cap 4B_{n_j} = \emptyset$$

(ii') : and

$$(38) \quad \mu\left(\bigcup_{1 \leq i \leq N_1} B_{n_i}\right) \geq \frac{1}{2Q_{d,4}}.$$

Set

$$\mathcal{W}_1 = \{U_{n_i}\}_{1 \leq i \leq N_1} \quad \text{and} \quad W_1 = \bigcup_{1 \leq i \leq N_1} U_{n_i}.$$

Along the construction of the Cantor set, for every $U \in \mathcal{U}$, the ball of \mathcal{B} naturally associated with U will be denoted $B^{[U]}$ (that is $B^{[U_n]} = B_n$).

The pre-measure η on the σ -algebra generated by the sets of \mathcal{W}_1 is defined by

$$(39) \quad \text{for every } U \in \mathcal{W}_1, \quad \eta(U) = \frac{\mu(B^{[U]})}{\sum_{\tilde{U} \in \mathcal{W}_1} \mu(B^{[\tilde{U}]})}.$$

It is obvious that $\eta(\mathbb{R}^d) = \eta(W_1) = 1$.

Recalling (8) and (9), since $s_1 < s(\mu, \mathcal{B}, \mathcal{U})$, the sub-sequence $(B_n)_{n \in \mathcal{N}_\mu(\mathcal{B}, \mathcal{U}, s_1)}$ is μ -a.-c. Recall also that $\lim_{n \rightarrow +\infty} r_n = 0$ and for every $n \in \mathbb{N}$, $|U_n| \leq r_n$.

So, for every $n \in \mathcal{N}_\mu(\mathcal{B}, \mathcal{U}, s_1)$,

$$(40) \quad \mathcal{H}_\infty^{\mu, s_1}(U_n) \geq \mu(B_n) \quad \text{and} \quad |U_n| \leq r_n.$$

In particular, by Definition 2.4, for every $n \in \mathcal{N}_\mu(\mathcal{B}, \mathcal{U}, s_1)$ for any set $E_n \subset U_n$ with $\mu(E_n) = \mu(U_n)$,

$$\mu(B_n) \leq \mathcal{H}_\infty^{\mu, s_1}(U_n) \leq \mathcal{H}_\infty^{s_1}(E_n).$$

By Lemma 3.4, and the notations therein, one has

$$m_{E_n}^{s_1}(U_n) = 1 \leq \frac{\kappa_d |U_n|^{s_1}}{\mathcal{H}_\infty^{s_1}(E_n)} \leq \frac{\kappa_d |U_n|^{s_1}}{\mu(B_n)}.$$

This implies that

$$(41) \quad \mu(B_n) \leq \kappa_d |U_n|^{s_1}.$$

By equation (41), recalling the fact that the sets $\mathcal{W}_1 \subset \{U_n\}_{n \in \mathbb{N}}$, one has for every $U \in \mathcal{W}_1$,

$$(42) \quad \eta(U) \leq \frac{\mu(B^{[U]})}{\frac{1}{2Q_{d,4}}} \leq 2Q_{d,4} \kappa_d |U|^{s_1}.$$

Step 2. This step (and all the following steps) is split into two sub-steps. First, into each open set U of \mathcal{W}_1 , smaller intermediary balls are selected according to the μ -essential content of U . Then in a second time, each intermediary ball will be covered by balls of the sequence $(B_n)_{n \in \mathbb{N}}$ according to the measure μ and, as in step 1, the sets U_n associated with this covering will form the generation \mathcal{W}_2 .

Let $g \in \mathbb{N}$ be such that for every $n \geq g$, $r_n \leq \frac{1}{3} \min(|U| : U \in \mathcal{W}_1)$.

As above, since $s_2 < s(\mu, \mathcal{B}, \mathcal{U})$, the sub-sequence $(B_n)_{n \in \mathcal{N}_\mu(\mathcal{B}, \mathcal{U}, s_2), n \geq g}$ is μ -a.-c. The same arguments as above yield for every $n \in \mathcal{N}_\mu(\mathcal{B}, \mathcal{U}, s_2)$,

$$(43) \quad \mathcal{H}_\infty^{\mu, s_2}(U_n) \geq \mu(B_n) \quad \text{and} \quad |U_n| \leq r_n$$

and

$$(44) \quad \mu(B_n) \leq \kappa_d |U_n|^{s_2}.$$

Covering with respect to the μ -essential content. Consider $U \in \mathcal{W}_1$. Set $\beta = \overline{\dim}_H(\mu)$. For $0 \leq k \leq \lfloor \frac{\beta-\alpha}{\varepsilon_2} \rfloor + 1$, define $\theta_k = \alpha + k\varepsilon_2$. Write

$$(45) \quad E_U = U \cap E_\mu^{[\alpha, \beta], \varepsilon_2} \cap \limsup_{n \rightarrow +\infty} B_n.$$

Notice that by Proposition 3.7 and by item (1) of Proposition 2.1, one has $\mu(E_U) = \mu(U)$.

In addition, using the definition (6) of $\mathcal{H}_\infty^{\mu, s_2}$, the fact that $E_U \subset U$ and $\mu(E_U) = \mu(U)$, and finally (7) applied with $B_n = B^{[U]}$, one gets

$$(46) \quad \mathcal{H}_\infty^{s_2}(E_U) \geq \mathcal{H}_\infty^{\mu, s_2}(U) \geq \mu(B^{[U]}) > 0.$$

This allows us to apply Proposition 3.4: there exists a Borel probability measure $m_{E_U}^{s_2}$ supported on E_U such that for every ball $B := B(x, r)$, one has

$$m_{E_U}^{s_2}(B) \leq \kappa_d \frac{r^{s_2}}{\mathcal{H}_\infty^{s_2}(E_U)}.$$

Also, since $m_{E_U}^{s_2}(E_U) = 1$ and $E_U \subset E_\mu^{[\alpha, \beta], \varepsilon_2}$, and recalling (30), for any $0 \leq k \leq \lfloor \frac{\beta-\alpha}{\varepsilon_2} \rfloor + 1$, there exists ρ_{k, ε_2} such that

$$m_{E_U}^{s_2}(E_\mu^{[\theta_k, \theta_{k+1}], \rho_{k, \varepsilon_2}, \varepsilon_2}) \geq \frac{1}{2} m_{E_U}^{s_2}(E_\mu^{[\theta_k, \theta_{k+1}], \varepsilon}).$$

Setting $\rho_U = \min_{0 \leq k \leq \lfloor \frac{\beta-\alpha}{\varepsilon_2} \rfloor + 1} \rho_{k, \varepsilon_2}$ one has, for any $0 \leq k \leq \lfloor \frac{\beta-\alpha}{\varepsilon_2} \rfloor + 1$,

$$(47) \quad m_{E_U}^{s_2}(E_\mu^{[\theta_k, \theta_{k+1}], \rho_U, \varepsilon_2}) \geq \frac{1}{2} m_{E_U}^{s_2}(E_\mu^{[\theta_k, \theta_{k+1}], \varepsilon}).$$

In particular,

$$(48) \quad m_{E_U}^{s_2}(E_\mu^{[\alpha, \beta], \rho_U, \varepsilon_2}) \geq \frac{1}{2}$$

Let

$$(49) \quad S_U := \bigcup_{0 \leq k \leq \lfloor \frac{\beta-\alpha}{\varepsilon_2} \rfloor + 1} E_\mu^{[\theta_k, \theta_{k+1}], \rho_U, \varepsilon_2} \cap E_U \cap \{x \in \mathbb{R}^d : \overline{\dim}_{\text{loc}}(m_{E_U}^{s_2}, x) \leq d\}.$$

Recalling that for every probability measure m , $m(\{x = \overline{\dim}_{\text{loc}}(m, x) \leq d\}) = 1$, one necessarily has $m_{E_U}^{s_2}(S_U) \geq 1/2$.

Let $x \in S_U$; consider $0 \leq k_x \leq \lfloor \frac{\beta-\alpha}{\varepsilon_2} \rfloor + 1$ such that $x \in E_\mu^{[\theta_{k_x}, \theta_{k_x+1}], \rho_U, \varepsilon_2}$. Applying Lemma 3.6, there exists $0 < r_x < \min(\rho_x, \frac{1}{3} \min\{|V| : V \in \mathcal{W}_1\})$ and $t_x \in (5, 6)$ such that:

$$(50) \quad 10r_x < \rho_U;$$

$$(51) \quad B(x, r_x) \subset U \quad \text{and} \quad \mu(\partial B(x, r_x/t_x)) = 0;$$

$$(52) \quad r_x^{-\varepsilon_2} \geq 5^d \frac{4Q_{d,1}}{C_{\varepsilon_3, d}} \frac{\eta(U)}{\mu(B^{[U]})} \geq 5^{s_2} \frac{4Q_{d,1}}{C_{\varepsilon_2, d}} \frac{\eta(U)}{\mu(B^{[U]})};$$

$$(53) \quad r_x^{\theta_{k_x} + 2\varepsilon_2} \leq \mu(B(x, r_x)) \leq r_x^{\theta_{k_x} - 2\varepsilon_2};$$

$$(54) \quad m_{E_U}^{s_2}(B(x, r_x/t_x)) \geq C_{\varepsilon_2, d} \cdot m_{E_U}^{s_2}(B(x, r_x)).$$

Note that in (52) the second inequality follows automatically from the first one since $s_2 \leq \alpha \leq d$ and the constant $C_{\varepsilon,d}$ is an increasing function of ε .

The family $\{B(x, r_x) : x \in S_U\}$ forms a covering of S_U . We apply Lemma 3.1 with $v = 1$ (i.e., the standard Besicovich covering Theorem) to this family to extract $Q_{d,1}$ subfamilies of balls, $\mathcal{G}_1^U, \dots, \mathcal{G}_{Q_{d,1}}^U$ such that:

- $\forall 1 \leq i \leq Q_{d,1}, \forall B \neq B' \in \mathcal{G}_i^U$, one has $B \cap B' = \emptyset$,
- $S_U \subset \bigcup_{i=1}^{Q_{d,1}} \bigcup_{B \in \mathcal{G}_i^U} B$.

In particular, $m_{E_U}^{s_2} \left(\bigcup_{i=1}^{Q_{d,1}} \bigcup_{B \in \mathcal{G}_i^U} B \right) \geq m_{E_U}^{s_2}(S_U) \geq 1/2$.

At least one of these families, say $\mathcal{G}_{i_0}^U$, verifies that

$$m_{E_U}^{s_2} \left(\bigcup_{B \in \mathcal{G}_{i_0}^U} B \right) \geq \frac{m_{E_U}^{s_2}(S_U)}{Q_{d,1}} \geq \frac{1}{2Q_{d,1}}.$$

Writing $\mathcal{G}_{i_0}^U = \{B_{i_0,k}^U\}_{k \in \mathbb{N}}$, one can find an integer N_U so large that

$$m_{E_U}^{s_2} \left(\bigcup_{1 \leq k \leq N_U} B_{i_0,k}^U \right) \geq \frac{1}{4Q_{d,1}}.$$

Remind that each $B_{i_0,k}^U$ is a ball $B(x, r_x)$ satisfying (51), (52) and (54).

Finally, setting $\mathcal{G}^U = \{B(x, r_x/t_x) : B(x, r_x) \in \mathcal{F}_{i_0}^U\}$, one has by construction

$$(55) \quad m_{E_U}^{s_2} \left(\bigcup_{B \in \mathcal{G}^U} B \right) = \sum_{B \in \mathcal{G}^U} m_{E_U}^{s_2}(B) \geq \frac{C_{\varepsilon_2,d}}{4Q_{d,1}}.$$

One then extends the pre-measure η to the Borel σ -algebra generated by the balls of \mathcal{G}^U , by the formula

$$(56) \quad \text{for every } B \in \mathcal{G}^U, \quad \eta(B) = \eta(U) \times \frac{m_{E_U}^{s_2}(B)}{\sum_{B' \in \mathcal{G}^U} m_{E_U}^{s_2}(B')}.$$

By construction, this formula is consistent since $\eta(U) = \sum_{B \in \mathcal{G}^U} \eta(B)$.

Observe that by (25), (55) and (46), one has for every $B \in \mathcal{G}^U$,

$$(57) \quad \eta(B) \leq \eta(U) \kappa_d \frac{|B|^{s_2}}{\mathcal{H}_{\infty}^{s_2}(E_U)} \frac{4Q_{d,1}}{C_{\varepsilon_2,d}} \leq \frac{4Q_{d,1} \kappa_d}{C_{\varepsilon_2,d}} \frac{\eta(U)}{\mu(B^{[U]})} |B|^{s_2} \leq |B|^{s_2 - \varepsilon_2},$$

where the second inequality of (52) was used.

This is achieved simultaneously for all $U \in \mathcal{W}_1$.

Covering with respect to μ . Now, in order to build the second generation of the Cantor set K , we select balls of \mathcal{B} that lie in the interior of these intermediate balls $B \in \mathcal{G}^U$.

Let $U \in \mathcal{W}_1$ and $B \in \mathcal{G}^U$ be one of these intermediary balls. Since \mathcal{B} is μ -a.c., the last part of Lemma 3.10 proves the existence of a finite family $\mathcal{F}^B = \{U_{n_i}\}_{1 \leq i \leq N_B}$ such that

(i_1) for every $1 \leq i \leq N_B$, one has $B_{n_i} \subset \mathring{B}$ and

$$(58) \quad \max \left\{ 2Q_{d,4} \frac{\eta(B)}{\mu(B)}, \frac{5^d 4Q_{d,1} \kappa_d}{C_{\varepsilon_3,d}} \right\} \leq r_{n_i}^{-\varepsilon_2},$$

(i_2) for every $1 \leq i \neq j \leq N_B$, one has $B_{n_i} \cap B_{n_j} = \emptyset$.

In addition, recalling that $\mu(\partial B) = 0$ by (51), one has

$$\mu(B_{n_i}) > 0 \quad \text{and} \quad \mu \left(\bigcup_{1 \leq i \leq N_B} B_{n_i} \right) \geq \frac{3\mu(\mathring{B})}{4} = \frac{3\mu(B)}{4}.$$

Recall the definitions (29) and (49) of the sets $E_\mu^{[a,b],\rho_U,\varepsilon_2}$ and S_U . By equations (50)-(54), there exists $\alpha \leq a \leq \beta$ such that the center of B belongs to $S_U \subset E_\mu^{[a,a+\varepsilon_2],\rho_U\varepsilon_2}$, and $|B| \leq \rho_U$, hence one has

$$\mu(B \cap \widetilde{E}_\mu^{[a,a+\varepsilon_2],\rho_U,\varepsilon_2}) \geq \frac{3}{4}\mu(B).$$

By (i_2), and recalling (29), one has

$$\begin{aligned} \mu \left(\bigcup_{B_{n_i} : B_{n_i} \cap \widetilde{E}_\mu^{[a,a+\varepsilon_2],\rho_U,\varepsilon_2} \neq \emptyset} B_{n_i} \right) &\geq \mu \left(\bigcup_{1 \leq i \leq N_B} B_{n_i} \cap \widetilde{E}_\mu^{[a,a+\varepsilon_2],\rho_U,\varepsilon_2} \right) \\ &= \mu \left(\bigcup_{1 \leq i \leq N_B} B_{n_i} \right) + \mu \left(\widetilde{E}_\mu^{[a,a+\varepsilon_2],\rho_U,\varepsilon_2} \right) - \mu \left(\widetilde{E}_\mu^{[a,a+\varepsilon_2],\rho_U,\varepsilon_2} \cup \bigcup_{1 \leq i \leq N_B} B_{n_i} \right) \\ &\geq \frac{3}{4}\mu(B) + \frac{3}{4}\mu(B) - \mu(B) = \frac{1}{2}\mu(B). \end{aligned}$$

By a slight abuse of notations, up to an extraction, we still denote by $\{B_{n_i}\}_{1 \leq i \leq N_B}$ the balls B_{n_i} such that $B_{n_i} \cap \widetilde{E}_\mu^{[a,a+\varepsilon_2],\varepsilon_2,\rho_U} \neq \emptyset$. The last inequality implies that the family of balls $\{B_{n_i}\}_{1 \leq i \leq N_B}$ can be chosen so that it verifies conditions (i_1) and (i_2), as well as the two following additional conditions:

$$(i_3) \quad \mu(B_{n_i}) > 0 \quad \text{and} \quad \mu \left(\bigcup_{1 \leq i \leq N_B} B_{n_i} \right) \geq \frac{\mu(\mathring{B})}{2} = \frac{\mu(B)}{2},$$

$$(i_4) \quad \text{for every } 1 \leq i \leq N_B, \quad B_{n_i} \cap \widetilde{E}_\mu^{[a,a+\varepsilon_2],\rho_U,\varepsilon_2} \neq \emptyset.$$

The obtained family is still denoted by \mathcal{F}^B .

Applying again Lemma 3.3 to \mathcal{F}^B with $v = 4$, as in step one (see (38), (i') and (ii')), if one must consider a subfamily, one can assume that the family \mathcal{F}^B satisfies (i_1) and (i_4) as well as the following condition (i'_2) and (i'_3):

$$(i'_2) : \text{for every } 1 \leq i \neq j \leq N_B, \text{ one has } 4B_{n_i} \cap 4B_{n_j} = \emptyset.$$

$$(i'_3) : \mu(B_{n_i}) > 0 \quad \text{and} \quad \mu \left(\bigcup_{1 \leq i \leq N_B} B_{n_i} \right) \geq \frac{\mu(\mathring{B})}{2Q_{d,4}} = \frac{\mu(B)}{2Q_{d,4}},$$

Finally one defines

$$\mathcal{W}_2 = \bigcup_{U \in \mathcal{W}_1} \bigcup_{B \in \mathcal{F}^{1,U}} \mathcal{F}^B \quad \text{and} \quad \mathcal{W}_2 = \bigcup_{L \in \mathcal{W}_2} L.$$

The pre-measure η is then extended to the σ -algebra generated by the elements of \mathcal{W}_2 by setting for every $U \in \mathcal{W}_1$, every $B \in \mathcal{G}^U$ and $V \in \mathcal{F}^B$,

$$(59) \quad \eta(V) = \eta(B) \times \frac{\mu(B^{[V]})}{\sum_{V' \in \mathcal{F}^B} \mu(B^{[V']})}.$$

By construction, one has $\sum_{V \in \mathcal{F}^B} \eta(V) = \eta(B)$. Also, (58), (59), and (i'_3) imply

$$(60) \quad \frac{\eta(V)}{\mu(B^{[V]})} \leq 2Q_{d,4} \frac{\eta(B)}{\mu(B)} \leq |V|^{-\varepsilon_2},$$

so that by (44) and (60) one has

$$(61) \quad \eta(V) \leq 2Q_{d,4} \frac{\eta(B)}{\mu(B)} \times \mu(B^{[V]}) \leq |B^{[V]}|^{-\varepsilon_2} |V|^{s_2} \leq |V|^{s_2 - \varepsilon_2}.$$

3.2.1. Recurrence scheme and end of the construction. Let $p \in \mathbb{N}^*$ be an integer, and set $\mathcal{W}_0 = \mathbb{R}^d$. Suppose that sets of balls $\mathcal{W}_1, \dots, \mathcal{W}_p$ as well as the measure η are constructed such that :

- (1) for every $1 \leq q \leq p$, $\mathcal{W}_q \subset \{U_n\}_{n \geq q}$, $\mathcal{W}_q \subset \mathcal{W}_{q-1}$, and η is defined on the σ -algebra generated by the elements of $\bigcup_{q=1}^p \mathcal{W}_q$.
- (2) For every $1 \leq q \leq p-1$, for every $U \in \mathcal{W}_q$, setting, as in step 2, $E_U = \limsup_{n \in \mathcal{N}_\mu(\mathcal{B}, \mathcal{M}, s_q)} B_n \cap U \cap E_\mu^{[\alpha, \beta], \varepsilon_q}$, then $\mathcal{H}_\infty^{s_q}(E_U) > 0$. If $m_{E_U}^{s_q}$ stands for the measure associated with E_U provided by Proposition 3.4, there exists $\rho_U > 0$ such that, for every $0 \leq k \leq \lfloor \frac{\beta - \alpha}{\varepsilon_q} \rfloor + 1$, setting $\theta_k = \theta_k^{(q)} = \alpha + k\varepsilon_q$, one has

$$m_{E_U}^{s_q}(E_U \cap E_\mu^{[\theta_k, \theta_{k+1}], \rho_U, \varepsilon_q}) \geq \frac{1}{2} m_{E_U}^{s_q}(E_U \cap E_\mu^{[\theta_k, \theta_{k+1}], \varepsilon_q}).$$

In particular,

$$m_{E_U}^{s_q}(E_U \cap \bigcup_{0 \leq k \leq \lfloor \frac{\beta - \alpha}{\varepsilon_q} \rfloor + 1} E_\mu^{[\theta_k, \theta_{k+1}], \rho_U, \varepsilon_q}) \geq \frac{1}{2}.$$

- (3) For every $1 \leq q \leq p-1$, for every $U \in \mathcal{W}_q$, there exists a finite family \mathcal{G}^U of balls $B(x, r_x/t_x)$, where $x, r_x < \frac{1}{3} \min \{|\tilde{U}| : \tilde{U} \in \mathcal{W}_q\}$ and t_x satisfy (50), (51), (52), (54) and (55). Also, if $B \neq B' \in \mathcal{G}^U$, $3B \cap 3B' = \emptyset$.

Also, for every $B \in \mathcal{G}^U$, (56) and (57) hold true. Moreover $\mathcal{W}_{q+1} \subset \bigcup_{U \in \mathcal{W}_q} \mathcal{G}^U$.

- (4) For every $1 \leq q \leq p-1$, for every $U \in \mathcal{W}_q$, for every $B \in \mathcal{G}^U$ there exists a family $\mathcal{F}^B \subset \{U_n\}_{n \geq q}$ of pairwise disjoint open sets such that :

- for every $\tilde{U} \neq \hat{U} \in \mathcal{F}^B$, one has

$$(62) \quad 4B^{[\tilde{U}]} \cap 4B^{[\hat{U}]} = \emptyset;$$

- for every $\tilde{U} \in \mathcal{F}^B$, $\tilde{U} \subset \mathring{B}$, (59) and (61) hold true, as well as

$$(63) \quad 2Q_{d,4} \frac{\eta(B)}{\mu(B)} \leq |B^{[\tilde{U}]}|^{-\varepsilon_{q+1}}$$

and

$$(64) \quad B^{[\tilde{U}]} \cap \tilde{E}_\mu^{[\theta_{k_B}, \theta_{k_B+1}], \rho_U, \varepsilon_{q+1}} \neq \emptyset;$$

- the following inequality also holds true:

$$(65) \quad \mu \left(\bigcup_{\tilde{U} \in \mathcal{F}^B} B^{[\tilde{U}]} \right) \geq \frac{\mu(B)}{2Q_{d,4}}.$$

In item (3), the fact that $3B \cap 3B' = \emptyset$ just follows from the choice of $B(x, r_x/t_x)$ instead of simply $B(x, r_x)$.

The proof follows then exactly and rigorously the same lines as in Step 2. We do not reproduce it here, the only differences are that $\mathcal{W}_1, \mathcal{W}_2$ and s_2 are replaced by $\mathcal{W}_p, \mathcal{W}_{p+1}$ and s_{p+1} .

Finally, define the Cantor set

$$K = \bigcap_{p \geq 1} W_p = \bigcap_{p \geq 1} \bigcup_{V \in \mathcal{W}_p} B^{[V]}.$$

Applying Caratheodory's extension Theorem to the pre-measure η yields a probability outer-measure on \mathbb{R}^d that we still denote by η , which is metric, so that Borel sets are η -measurable and its restriction to Borel sets belongs to $\mathcal{M}(\mathbb{R}^d)$. The so obtained measure η is fully supported on K . Also, for every $p \geq 2$, for any $U \in \mathcal{W}_p, B \in \mathcal{G}^U$, and $\tilde{U} \in \mathcal{F}^B$, the inequalities (56), (57), (59) and (61) holds with s_p and ε_p instead of s_2 and ε_2 .

3.2.2. Upper-bound for the mass of a ball. One first recall the following lemma (see, e.g., [8]).

Lemma 3.11. *Let $A = B(x, r)$ and $B = B(x', r')$ be two closed balls, $q \geq 3$ such that $A \cap B \neq \emptyset$ and $A \setminus (qB) \neq \emptyset$. Then $r' \leq r$ and $qB \subset 5A$.*

Define the gauge function $\zeta : \mathbb{R}^+ \mapsto \mathbb{R}^+$ as follows:

- if for some $p \geq 1$, $\frac{1}{3} \min \{|U| : U \in \mathcal{W}_{p+1}\} \leq r < \frac{1}{3} \min \{|U| : U \in \mathcal{W}_p\}$, then $\zeta(r) = 2Q_{d,4}10^d r^{s_p - 5\varepsilon_p}$,
- if $r \geq \frac{1}{3} \min \{|U| : U \in \mathcal{W}_1\}$, $\zeta(r) = 1$,
- $\zeta(0) = 0$.

Since $\varepsilon_p \rightarrow 0$, one checks that $\lim_{r \rightarrow 0^+} \frac{\log(\zeta(r))}{\log(r)} = \min \{s(\mu, \mathcal{B}, \mathcal{U}), \underline{\dim}_H(\mu)\}$.

Let A be a ball of radius r . If there exists $n \in \mathbb{N}$ such that A does not intersect K_n then $\eta(A) = \eta(A \cap K_n) = 0$. Suppose that for every $n \in \mathbb{N}$, A intersects K_n . The goal is to prove that $\eta(A) \leq \zeta(|A|)$ when $|A|$ is small.

Some cases must be distinguished.

First if for every $n \in \mathbb{N}$, A intersects only one contracted set V_n of K_n , then by (57)

$$\eta(A) \leq \eta(V_n) \leq |V_n|^{s_n - \varepsilon_n} \xrightarrow{n \rightarrow +\infty} 0.$$

In the other case, there exists $p \in \mathbb{N}$ such that A intersects only one element of \mathcal{W}_p , and at least two elements of \mathcal{W}_{p+1} . Denote by U the unique element of \mathcal{W}_p intersecting A .

(1) **Case 1:** If $|A| \geq |U|$, then by (61)

$$(66) \quad \eta(A) \leq \eta(U) \leq |U|^{s_p - \varepsilon_p} \leq \zeta(|A|).$$

- (2) **Case 2:** If $|A| < |U|$ and A intersects at least two balls of \mathcal{G}^U : Observe that when A intersects two balls B and B' of \mathcal{G}^U , since by item (3) of the recurrence scheme $3B \cap 3B' = \emptyset$, one necessarily has (by Lemma 3.11) $B \cup B' \subset 5A$. Hence, $\bigcup_{B \in \mathcal{G}^U: B \cap A \neq \emptyset} B \subset 5A$ and by (56) and (55),

$$\eta(A) = \eta(U) \times \frac{\sum_{B \in \mathcal{G}^U: B \cap A \neq \emptyset} m_{E_U}^{s_{p+1}}(B)}{\sum_{B' \in \mathcal{G}^U} m_{E_U}^{s_{p+1}}(B')} \leq \frac{4Q_{d,1}}{C_{\varepsilon_{p+1},d}} \eta(U) m_{E_U}^{s_{p+1}}(5A).$$

Then, by (25), (46), (58) and (60)

$$\begin{aligned} \eta(A) &\leq \frac{4Q_{d,1}}{C_{\varepsilon_{p+1},d}} \eta(U) \kappa_d \frac{(5|A|)^{s_{p+1}}}{\mathcal{H}_{\infty}^{\mu, s_{p+1}}(E_U)} \leq 5^{s_{p+1}} \frac{4Q_{d,1} \kappa_d}{C_{\varepsilon_{p+1},d}} \frac{\eta(U)}{\mu(B^{[U]})} |A|^{s_{p+1}} \\ (67) \quad &\leq |A|^{s_{p+1}} |U|^{-2\varepsilon_p} \leq |A|^{s_{p+1}-2\varepsilon_{p+1}} \leq \zeta(|A|), \end{aligned}$$

where we used that $|A| < |U|$, and the mappings $x \mapsto |U|^{-x}$ and $x \mapsto x^{-\varepsilon_{p+1}}$ are decreasing.

- (3) **Case 3:** If A intersects only one ball of \mathcal{G}^U : calling B this particular ball and r_B its radius (at this stage there should be no confusion with the radii of the terms of the sequence $(B_n)_{n \geq 1}$), two cases must again be distinguished:

(a) **Subcase 3.1:** $|B| \leq |A|$: by (57),

$$(68) \quad \eta(A) \leq \eta(B) \leq |B|^{s_{p+1}-\varepsilon_{p+1}} \leq |A|^{s_{p+1}-\varepsilon_{p+1}} \leq \zeta(|A|).$$

(b) **Subcase 3.2:** $|A| \leq |B|$: Denote by k_B the integer such that its center belongs to $E_{\mu}^{[\theta_{k_B}, \theta_{k_B+1}], \rho_U, \varepsilon_{p+1}}$.

The ball A must intersect at least two elements $V \neq V'$ of \mathcal{W}_{p+1} (by definition of p). Note that these sets must belong to \mathcal{F}^B (because A intersects only B). Applying Lemma 3.11 to the ball A with any of these balls $V \in \mathcal{F}_{p+1}$, since $A \cap V \neq \emptyset$ and $A \setminus B^{[V]} \neq \emptyset$ (because A intersects an other dilated ball, $B^{[V']}$ by hypothesis and two such balls verifies (62)), one has

$$(69) \quad \bigcup_{V \cap A \neq \emptyset} B^{[V]} \subset 5A.$$

Then, (59) and (65) imply that

$$(70) \quad \eta(A) = \eta(B) \cdot \frac{\sum_{V \in \mathcal{W}_{p+1}: V \cap A \neq \emptyset} \mu(B^{[V]})}{\sum_{V' \in \mathcal{F}^B} \mu(B^{[V']})} \leq 2Q_{d,4} \frac{\eta(B)}{\mu(B)} \mu(5A).$$

Recalling (69), the ball $5A$ contains some of the balls of \mathcal{F}^B : Hence, by (64), $\widetilde{E}_{\mu}^{[\theta_{k_B}, \theta_{k_B+1}], \rho_U, \varepsilon_{p+1}} \cap 5A \neq \emptyset$. Since $|A| \leq |B|$, by (50), since $r_B < \frac{1}{10} \rho_U$, for any $x \in \widetilde{E}_{\mu}^{[\theta_{k_B}, \theta_{k_B+1}], \rho_U, \varepsilon_{p+1}} \cap 5A$ one has

$$(71) \quad \mu(5A) \leq \mu(B(x, 10r)) \leq (10r)^{\theta_{k_B} - 2\varepsilon_{p+1}}.$$

Recalling (53) (applied to the ball B), one has

$$(72) \quad \mu(B) \geq (r_B)^{\theta_{k_B} + 2\varepsilon_{p+1}}.$$

Using (57) (applied to B) (70), (71) and (72), one obtains

$$\begin{aligned} \eta(A) &\leq 2Q_{d,4} r_B^{s_{p+1}-\varepsilon_{p+1}} \frac{(10r)^{\theta_{k_B}-2\varepsilon_{p+1}}}{r_B^{\theta_{k_B}+2\varepsilon_{p+1}}} \\ &= 2Q_{d,4} 10^{\theta_{k_B}-2\varepsilon_{p+1}} \frac{r_B^{s_{p+1}-\theta_{k_B}-\frac{\varepsilon_{p+1}}{\delta}-2\varepsilon_{p+1}}}{r_B^{s_{p+1}-\theta_{k_B}-\varepsilon_{p+1}-2\varepsilon_{p+1}}} r_B^{s_{p+1}-\theta_{k_B}-\varepsilon_{p+1}-4\varepsilon_{p+1}} \\ &\leq 2Q_{d,4} 10^{\theta_{k_B}-2\varepsilon_{p+1}} r_B^{s_{p+1}-5\varepsilon_{p+1}}. \end{aligned}$$

Finally, recalling (37), $s_{p+1} - 5\varepsilon_{p+1} \leq \alpha \leq \theta_k$, and since $r_B \geq r$ and $s_p \leq s_{p+1}$, one gets

$$\eta(A) \leq 2Q_{d,4} 10^{\theta_k-\varepsilon_{p+1}} (r)^{s_{p+1}-5\varepsilon_{p+1}} \leq 2Q_{d,4} 10^{d-\varepsilon_{p+1}} |A|^{s_p-5\varepsilon_p} \leq 2Q_{d,4} 10^{d-\varepsilon_{p+1}} |A|^{s_p-5\varepsilon_p},$$

hence

$$(73) \quad \eta(A) \leq \zeta(|A|).$$

Since for any $p \in \mathbb{N}$ and any ball A satisfying $|A| \leq \frac{1}{3} \min\{|U| : U \in \mathcal{W}_p\}$, if A intersects at most one element of \mathcal{W}_p , the inequalities (66), (67), (68), (73) proves that for any such ball, one has $\eta(A) \leq \zeta(|A|)$.

Hence recalling Definition 2.1, by the mass distribution principle, one deduces that $\mathcal{H}^\zeta(K) \geq 1$, which concludes the proof of Theorem 2.2.

3.3. Proof of Corollary 2.5.

3.3.1. *Some basic properties about the μ -essential Hausdorff content.* In this subsection, basic properties of the μ -essential content are established.

First, we work in this article with the $\|\cdot\|_\infty$ norm for convenience. Any other norm could have been chosen, the corresponding quantities would have been equivalent.

In (3), only closed balls are considered. Choosing open balls does not change the value of (6) in Definition 2.4.

The following propositions are directly derived from the properties of the standard Hausdorff measures.

Proposition 3.12. *Let $\mu \in \mathcal{M}(\mathbb{R}^d)$, $s \geq 0$ and $A \subset \mathbb{R}^d$ be a Borel set. The s -dimensional $\mathcal{H}_\infty^{\mu,s}(\cdot)$ outer measure satisfies the following properties:*

- (1) *If $|A| \leq 1$, the mapping $s \geq 0 \mapsto \mathcal{H}_\infty^{\mu,s}(A)$ is decreasing from $\mathcal{H}_\infty^{\mu,0}(A) = 1$ to $\lim_{t \rightarrow +\infty} \mathcal{H}_\infty^{\mu,t}(A) = 0$.*
- (2) $0 \leq \mathcal{H}_\infty^{\mu,s}(A) \leq \min\{|A|^s, \mathcal{H}_\infty^s(A)\}$.
- (3) *For every subset $B \subset A$ with $\mu(A) = \mu(B)$, $\mathcal{H}_\infty^{\mu,s}(A) = \mathcal{H}_\infty^{\mu,s}(B)$.*
- (4) *For every $\delta \geq 1$, $\mathcal{H}_\infty^{\mu,\frac{s}{\delta}}(A) \geq (\mathcal{H}_\infty^{\mu,s}(A))^{\frac{1}{\delta}}$.*
- (5) *For every $s > \overline{\dim}_H(\mu)$, $\mathcal{H}_\infty^{\mu,s}(A) = 0$.*

Proof. Items (1), (2), (3) directly follow from the definition. Item (4) is obtained by concavity of the mapping $x \mapsto |x|^{1/\delta}$.

(5) By Definition 2.2, for any $s > \overline{\dim}_H(\mu)$, there exists a set E with $\dim_H(E) < s$ and $\mu(E) = 1$. Using item (2), one has then $0 \leq \mathcal{H}_\infty^{\mu,s}(A) = \mathcal{H}_\infty^{\mu,s}(A \cap E) \leq \mathcal{H}_\infty^s(A \cap E) \leq \mathcal{H}^s(E) = 0$.

□

3.3.2. *Proof of Corollary 2.5.* One starts with a lemma, the proof of which can be found in [14].

Lemma 3.13. *Let $\mu \in \mathcal{M}(\mathbb{R}^d)$. Let $\mathcal{B} = (B_n := B(x_n, r_n))_{n \in \mathbb{N}}$ be a μ -a.c sequence of balls of \mathbb{R}^d . Then for every $\varepsilon > 0$, there exists a μ -a.c sub-sequence $(B_{\phi(n)})_{n \in \mathbb{N}}$ of \mathcal{B} such that for every $n \in \mathbb{N}$, $\mu(B_{\phi(n)}) \leq (r_{\phi(n)})^{\underline{\dim}_H(\mu) - \varepsilon}$.*

Proof of Corollary 2.5. (1) Observe that item (2) of Proposition 3.12 implies that $t(\mu, \delta, \varepsilon, \mathcal{B}) \geq \underline{\dim}_H(\mu) - \varepsilon$, and $t(\mu, \delta, \mathcal{B}) \geq \underline{\dim}_H(\mu)$.

Now choose $\varepsilon > 0$ so small that $\frac{(1-\varepsilon)\underline{\dim}_H(\mu)}{\delta \cdot t(\mu, \delta, \varepsilon, \mathcal{B})} \leq 1$. Recalling Lemma 3.13, up to an extraction, one can assume that for any $n \in \mathbb{N}$,

$$\mu(B_n) \leq |B_n|^{(1-\varepsilon^2) \cdot \underline{\dim}_H(\mu)}.$$

Due to (11), there exists $N_\varepsilon \in \mathbb{N}$ such that for any $n \geq N_\varepsilon$,

$$\mathcal{H}_\infty^{\mu, \underline{\dim}_H(\mu) - \varepsilon}(\mathring{B}_n^\delta) \geq |B_n^\delta|^{(1+\varepsilon) \cdot t(\mu, \delta, \varepsilon, \mathcal{B})}.$$

Then, Proposition 3.12 (4) implies that for every $n \geq N_\varepsilon$,

$$\begin{aligned} \mathcal{H}_\infty^{\mu, \frac{(1-\varepsilon)\underline{\dim}_H(\mu) \times (\underline{\dim}_H(\mu) - \varepsilon)}{\delta \cdot t(\mu, \delta, \varepsilon, \mathcal{B})}}(\mathring{B}_n^\delta) &\geq (\mathcal{H}_\infty^{\mu, \underline{\dim}_H(\mu) - \varepsilon}(B_n^\delta))^{\frac{(1-\varepsilon)\underline{\dim}_H(\mu)}{\delta \cdot t(\mu, \delta, \varepsilon, \mathcal{B})}} \\ &\geq |B_n^\delta|^{\frac{(1+\varepsilon)t(\mu, \delta, \varepsilon, \mathcal{B})}{\delta \cdot t(\mu, \delta, \varepsilon, \mathcal{B})} (1-\varepsilon) \cdot \underline{\dim}_H(\mu)} \\ &\geq |B_n|^{(1+\varepsilon)(1-\varepsilon)\underline{\dim}_H(\mu)} \geq \mu(B_n). \end{aligned}$$

Thus, setting $s_{\delta, \varepsilon} = \frac{(1-\varepsilon)\underline{\dim}_H(\mu) \times (\underline{\dim}_H(\mu) - \varepsilon)}{\delta \cdot t(\mu, \delta, \varepsilon, \mathcal{B})}$, Corollary 2.4 yields

$$\dim_H(\limsup_{n \rightarrow +\infty} \mathring{B}_n^\delta) \geq s_{\delta, \varepsilon}.$$

Since the result holds for any $\varepsilon > 0$, one gets the desired conclusion. □

4. ESTIMATION OF ESSENTIAL CONTENT FOR SELF-SIMILAR MEASURES

In this section one computes the Hausdorff content of balls in the case of the Lebesgue measure, and estimates it for any self-similar measure.

4.1. Computation of essential content for the Lebesgue measure. When the measure μ is the Lebesgue measure, the computations are quite easy.

Proposition 4.1. *Let $B = B(x, r)$ be a ball in \mathbb{R}^d , and \mathcal{L}^d be the d -dimensional Lebesgue measure. Then for any $0 \leq s \leq d$, $\mathcal{H}_\infty^{\mathcal{L}^d, s}(B) = \mathcal{H}_\infty^{\mathcal{L}^d, s}(\mathring{B}) = r^s$.*

Proof. One starts first by computing $\mathcal{H}_\infty^{\mathcal{L}^d, d}(B)$.

Let $\varepsilon > 0$, and let $E \subset B$ be a Borel set with $\mathcal{L}^d(E) = \mathcal{L}^d(B)$. Notice first that since B covers E , recalling that \mathbb{R}^d is endowed with $\|\cdot\|_\infty$ one has

$$\mathcal{H}_\infty^{\mathcal{L}^d, d}(E) \leq \mathcal{H}_\infty^d(B) \leq |B|^d.$$

Consider a sequence of balls $(L_n)_{n \in \mathbb{N}}$ such that

$$\mathcal{H}_\infty^d(E) \leq \sum_{n \geq 0} |L_n|^d \leq (1 + \varepsilon) \mathcal{H}_\infty^d(E).$$

This implies

$$\begin{aligned} (1 + \varepsilon)|B|^d &\geq (1 + \varepsilon)\mathcal{H}_\infty^d(B) \geq (1 + \varepsilon)\mathcal{H}_\infty^d(E) \geq \sum_{n \geq 0} |L_n|^d \\ &\geq \sum_{n \geq 0} \mathcal{L}^d(L_n) \geq \mathcal{L}^d(E) = \mathcal{L}^d(B) = |B|^d. \end{aligned}$$

Taking the infimum on the Borel sets $E \subset B$ such that $\mathcal{L}^d(E) = \mathcal{L}^d(A)$ gives

$$|B|^d \leq (1 + \varepsilon)\mathcal{H}_\infty^{\mathcal{L}^d, d}(B).$$

In particular,

$$\frac{1}{1 + \varepsilon}|B|^d \leq \mathcal{H}_\infty^{\mathcal{L}^d, d}(B) \leq |B|^d.$$

Letting $\varepsilon \rightarrow 0$ shows that $\mathcal{H}_\infty^{\mathcal{L}^d, d}(B) = |B|^d$. This implies, with item (4) of Proposition 3.12, that for any $\delta \geq 1$,

$$|B|^{\frac{d}{\delta}} \geq \mathcal{H}_\infty^{\mathcal{L}^d, \frac{d}{\delta}}(B) \geq (\mathcal{H}_\infty^{\mathcal{L}^d, d}(B))^{\frac{1}{\delta}} = |B|^{\frac{d}{\delta}},$$

hence the result. \square

4.2. Proof of Theorem 2.6.

Proposition 4.2. *Let μ be a self-similar measure. For any $0 < \varepsilon \leq \dim(\mu)$, there exists a constant $\kappa(d, \mu, \varepsilon) \in (0, 1)$ such that for any ball $B = B(x, r)$ with $x \in K$ (the attractor of the underlying IFS) and $r \leq 1$, one has*

$$\kappa(d, \mu, \varepsilon)|B|^{\dim(\mu) - \varepsilon} \leq \mathcal{H}_\infty^{\mu, \dim(\mu) - \varepsilon}(\dot{B}) \leq \mathcal{H}_\infty^{\mu, \dim(\mu) - \varepsilon}(B) \leq |B|^{\dim(\mu) - \varepsilon}.$$

In addition, for any $s > \dim(\mu)$ one has $\mathcal{H}_\infty^{\mu, s}(B) = 0$.

Proof. Let $\{f_1, \dots, f_m\}$ the underlying IFS. Denote by c_i the contraction ration of f_i , and (p_1, \dots, p_m) the probability vector with positive entries associated with μ so that (13) is satisfied. Set $\alpha = \dim(\mu)$ and $\Lambda = \{1, \dots, m\}$. For $k \geq 0$ and $\underline{i} := (i_1, \dots, i_k) \in \Lambda^k$, define

- $c_{\underline{i}} = c_{i_1} \dots c_{i_k}$, $f_{\underline{i}} = f_{i_1} \circ \dots \circ f_{i_k}$ and $K_{\underline{i}} = f_{\underline{i}}(K)$, so that $|K_{\underline{i}}| = c_{\underline{i}}|K|$.
- $\Lambda^{(k)} = \{\underline{i} := (i_1, \dots, i_s) : c_{i_s} 2^{-k} < c_{\underline{i}} \leq 2^{-k}\}$.

Note first that item (5) of Proposition 3.12 implies that for any $s > \dim(\mu)$, $\mathcal{H}_\infty^{\mu, s}(B) = 0$.

Let us consider $0 \leq s < \dim_H(\mu)$ and start by few remarks.

Recalling (29) and Proposition 3.7, let us fix ρ_ε so that $\mu(E_\mu^{[\alpha, \alpha], \rho_\varepsilon, \varepsilon}) \geq \frac{1}{2}$ and write $E = E_\mu^{[\alpha, \alpha], \rho_\varepsilon, \varepsilon}$.

Set $\Lambda^* := \bigcup_{k \geq 0} \Lambda^k$, and for $\underline{i} \in \Lambda^*$, define $E_{\underline{i}} = f_{\underline{i}}(E)$ and $\mu_{\underline{i}} = \mu(f_{\underline{i}}^{-1})$. One has

$$\begin{aligned} (74) \quad E_{\underline{i}} &= \{f_{\underline{i}}(x) \in \mathbb{R}^d : x \in K \forall r \leq \rho_\varepsilon, \mu(B(x, r)) \leq r^{\alpha - \varepsilon}\} \\ &= \left\{ f_{\underline{i}}(x) : x \in K, \forall, c_{\underline{i}} r \leq c_{\underline{i}} \rho_\varepsilon, \mu(f_{\underline{i}}^{-1}(B(f_{\underline{i}}(x), r c_{\underline{i}}))) \leq \left(\frac{r c_{\underline{i}}}{c_{\underline{i}}}\right)^{\alpha - \varepsilon} \right\} \\ &= \left\{ y \in K_{\underline{i}} : \forall r' \leq c_{\underline{i}} \rho_\varepsilon, \mu_{\underline{i}}(B(y, r')) \leq \left(\frac{r'}{c_{\underline{i}}}\right)^{\alpha - \varepsilon} \right\}, \end{aligned}$$

Also, $\mu_{\underline{i}}(E_{\underline{i}}) = \mu(E) \geq \frac{1}{2}$.

One emphasizes that iterating the self-similarity equation gives

$$\mu = \sum_{i' \in \Lambda^k} p_{i'} \mu_{i'},$$

which implies that $\mu_{\underline{i}}$ is absolutely continuous with respect to μ (since all $p_{\underline{i}}$'s are strictly positive).

We are now ready to estimate the μ -essential content of a ball B centered on K .

Let $B = B(x, r)$ with $x \in K$ and $r \leq \min_{1 \leq i \leq m} c_i$.

Since $x \in K$, there exists an \underline{i} such that $\min_{1 \leq j \leq m} c_j r \leq c_{\underline{i}} |K| \leq r$ and $K_{\underline{i}} \subset \overset{\circ}{B}$.

By construction, $E_{\underline{i}} \subset \overset{\circ}{B}$.

Consider a Borel set $A \subset B$ such that $\mu(A) = \mu(B)$. One aims at giving a lower-bound of the Hausdorff content of A which does not depends on A .

Consider a sequence of balls $(L_n = B(z_n, \ell_n))_{n \geq 1}$ covering $A \cap E_{\underline{i}}$, such that $\ell_n < \rho_\varepsilon c_{\underline{i}}$ and $z_n \in A \cap E_{\underline{i}}$. Since $\mu_{\underline{i}}$ is absolutely continuous with respect to μ , it holds that $\mu_{\underline{i}}(A) = 1$.

By (74) applied to every $n \in \mathbb{N}$, one has $\left(\frac{|L_n|}{c_{\underline{i}}}\right)^{\alpha-\varepsilon} \geq \mu_{\underline{i}}(L_n)$, so that

$$(75) \quad \sum_{n \in \mathbb{N}} |L_n|^{\alpha-\varepsilon} \geq \sum_{n \in \mathbb{N}} c_{\underline{i}}^{\alpha-\varepsilon} \mu_{\underline{i}}(L_n) \geq c_{\underline{i}}^{\alpha-\varepsilon} \mu_{\underline{i}}\left(\bigcup_{n \in \mathbb{N}} L_n\right) \geq c_{\underline{i}}^{\alpha-\varepsilon} \mu_{\underline{i}}(E_{\underline{i}}) \geq \frac{1}{2} c_{\underline{i}}^{\alpha-\varepsilon}.$$

This series of inequalities holds for any sequence of balls $(L_n)_{n \in \mathbb{N}}$ with radius less than $\rho_\varepsilon c_{\underline{i}}$ centered on $A \cap E_{\underline{i}}$ and covering $A \cap E_{\underline{i}}$.

Now, assume that $(L_n)_{n \in \mathbb{N}}$ is a sequence of balls covering $A \cap E_{\underline{i}}$, which still verifies $\ell_n < \rho_\varepsilon c_{\underline{i}}$ but z_n does not necessarily belongs to $A \cap E_{\underline{i}}$.

Let $n \in \mathbb{N}$. One constructs recursively a sequence of balls $(L_{n,j})_{1 \leq j \leq J_n}$ such that the following properties hold for any $1 \leq j \leq J_n$:

- $L_{n,j}$ is centered on $A \cap E_{\underline{i}} \cap L_n$;
- $A \cap E_{\underline{i}} \cap L_n \subset \bigcup_{1 \leq j \leq J_n} L_{n,j}$;
- for all $1 \leq j \leq J_n$, $|L_{n,j}| = |L_n|$;
- the center of $L_{n,j}$ does not belong to any $L_{n,j'}$ for $1 \leq j' \neq j \leq J_n$.

To achieve this, simply consider $y_1 \in A \cap E_{\underline{i}} \cap L_n$ and set $L_{1,n} = B(y_1, \ell_n)$. If $A \cap E_{\underline{i}} \cap L_n \not\subset L_{1,n}$, consider $y_2 \in A \cap E_{\underline{i}} \cap L_n \setminus L_{1,n}$ and set $L_{2,n} = B(y_2, \ell_n)$. If $A \cap E_{\underline{i}} \cap L_n \not\subset L_{1,n} \cup L_{2,n}$, consider $y_3 \in A \cap E_{\underline{i}} \cap L_n \setminus L_{1,n} \cup L_{2,n}$ and set $L_{3,n} = B(y_3, \ell_n)$, and so on...

Note that, for any $1 \leq j \leq J_n$, any ball $L_{j,n}$ has radius ℓ_n , intersects L_n (which also has radius ℓ_n) and, because $y_j \notin \bigcup_{1 \leq j' \neq j \leq J_n} L_{j',n}$, it holds that, for any $j \neq j'$, $\frac{1}{3}L_{n,j} \cap \frac{1}{3}L_{n,j'} = \emptyset$. By Lemma 3.2, this implies that $J_n \leq Q_{d, \frac{1}{3}}$.

Hence, denoting by $(\tilde{L}_n)_{n \in \mathbb{N}}$ the collection of the corresponding balls centered on $A \cap E_{\underline{i}}$ associated with all the balls L_n , one has by (75) applied to $(\tilde{L}_n)_{n \in \mathbb{N}}$:

$$\sum_{n \in \mathbb{N}} |L_n|^{\alpha-\varepsilon} \geq \frac{1}{Q_{d, \frac{1}{3}}} \sum_{n \in \mathbb{N}} |\tilde{L}_n|^{\alpha-\varepsilon} \geq \frac{1}{2Q_{d, \frac{1}{3}}} c_{\underline{i}}^{\alpha-\varepsilon}.$$

Remark also that any ball of radius smaller than r can be covered by at most $\left(\frac{2}{\rho_\varepsilon}\right)^d$ balls of radius $r\rho_\varepsilon$.

This proves that, for any sequence of balls \widehat{L}_n covering $A \cap E_i$, since $c_i \geq \frac{\min_{1 \leq j \leq m} c_j}{|K|} |B|$, it holds that

$$(76) \quad \sum_{n \in \mathbb{N}} |\widehat{L}_n|^{\alpha-\varepsilon} \geq \frac{\rho_\varepsilon^d}{2^{d+1} Q_{d, \frac{1}{3}}} c_i^{\alpha-\varepsilon} \geq \frac{\min_{1 \leq j \leq m} c_j^{\alpha-\varepsilon} \rho_\varepsilon^d}{|K|^{\alpha-\varepsilon} 2^{d+1} Q_{d, \frac{1}{3}}} |B|^{\alpha-\varepsilon}.$$

Recall (3). Since (76) is valid for any covering $(\widehat{L}_n)_{n \in \mathbb{N}}$ of $A \cap E_i$, one has

$$(77) \quad |B|^{\alpha-\varepsilon} \geq \mathcal{H}_\infty^{\alpha-\varepsilon}(A) \geq \mathcal{H}_\infty^{\alpha-\varepsilon}(A \cap E_i) \geq \frac{\min_{1 \leq j \leq m} c_j^{\alpha-\varepsilon} \rho_\varepsilon^d}{|K|^{\alpha-\varepsilon} 2^{d+1} Q_{d, \frac{1}{3}}} |B|^{\alpha-\varepsilon}.$$

Taking the infimum over all the Borel sets $A \subset B$ satisfying $\mu(A) = \mu(B)$, one gets

$$|B|^{\alpha-\varepsilon} \geq \mathcal{H}_\infty^{\mu, \alpha-\varepsilon}(B) \geq \frac{\min_{1 \leq j \leq m} c_j^{\alpha-\varepsilon} \rho_\varepsilon^d}{|K|^{\alpha-\varepsilon} 2^{d+1} Q_{d, \frac{1}{3}}} |B|^{\alpha-\varepsilon}.$$

The result stands for balls of diameter less than $\min_{1 \leq j \leq m} c_j$. Then for any ball B centered on K with $|B| \leq 1$, remarking that

$$|B|^{\alpha-\varepsilon} \geq \mathcal{H}_\infty^{\mu, \alpha-\varepsilon}(B) \geq \mathcal{H}_\infty^{\mu, \alpha-\varepsilon}\left(\min_{1 \leq j \leq m} c_j B\right)$$

and setting $\kappa(d, \mu, \varepsilon) = \frac{\min_{1 \leq j \leq m} c_j^{2(\alpha-\varepsilon)} \rho_\varepsilon^d}{|K|^{\alpha-\varepsilon} 2^{d+1} Q_{d, \frac{1}{3}}}$ yields the desired inequality. \square

Remark 4.3. Note that in the proof of Proposition 4.2, the estimate of $\mathcal{H}_\infty^{\mu, s}(B)$ for $s < \dim(\mu)$ only relies on the absolute continuity of $\mu(f_i^{-1}(\cdot))$, for any $i \in \Lambda^*$. In particular, the same estimate holds for any quasi-Bernoulli measures (which are proved to be exact-dimensional, see [22]).

This result in hand, one establishes the more general Theorem 2.6.

Proof of Theorem 2.6. Note first, that by item (5) of Proposition 3.12, for any $s > \dim(\mu)$ and any set E , one has $\mathcal{H}_\infty^{\mu, s}(E) = 0$.

Let us fix $s < \dim(\mu)$ and set $\varepsilon = \dim(\mu) - s > 0$. Since $K \cap \Omega \subset \Omega$ and $\mu(K \cap \Omega) = \mu(\Omega)$, it holds that

$$\mathcal{H}_\infty^{\mu, s}(\Omega) \leq \mathcal{H}_\infty^s(\Omega \cap K).$$

It remains to show that there exists a constant $c(d, \mu, s)$ such that for any open set Ω , the converse inequality

$$c(d, \mu, s) \mathcal{H}_\infty^s(\Omega \cap K) \leq \mathcal{H}_\infty^{\mu, s}(\Omega)$$

holds.

Let $E \subset \Omega$ be a Borel set such that $\mu(E) = \mu(\Omega)$ and

$$(78) \quad \mathcal{H}_\infty^s(E) \leq 2\mathcal{H}_\infty^{\mu, s}(\Omega).$$

Let $\{L_n\}_{n \in \mathbb{N}}$ be a covering of E by balls verifying

$$(79) \quad \mathcal{H}_\infty^s(E) \leq \sum_{n \geq 0} |L_n|^s \leq 2\mathcal{H}_\infty^s(E).$$

The covering $(L_n)_{n \in \mathbb{N}}$ will be modified to get a covering $(\widetilde{L}_n)_{n \in \mathbb{N}}$ which verifies the following properties:

- $K \cap \Omega \subset \bigcup_{n \in \mathbb{N}} \widetilde{L}_n$;

- $\bigcup_{n \in \mathbb{N}} L_n \subset \bigcup_{n \in \mathbb{N}} \tilde{L}_n$;
- one has

$$\sum_{n \geq 0} |\tilde{L}_n|^s \leq 8 \cdot 2^s \frac{Q_{d,1}^2}{\kappa(d, \mu, \varepsilon)} \sum_{n \geq 0} |L_n|^s,$$

where $\kappa(d, \mu, \varepsilon)$ is the constant introduced in Proposition 4.2 and $Q_{d,1}$ is the constant arising in Proposition 3.8 applied with $v = 1$. Last item together with (78) and (79) then immediately imply that

$$\frac{\kappa(d, \mu, \varepsilon)}{8 \cdot 2^s Q_{d,1}^2} \mathcal{H}_\infty^s(K \cap \Omega) \leq \mathcal{H}_\infty^{\mu, s}(\Omega),$$

and setting $c(d, \mu, s) = \frac{\kappa(d, \mu, \dim(\mu) - s)}{8 \cdot 2^s Q_{d,1}^2}$ concludes the proof.

Let us start the construction of the sequence of balls $(\tilde{L}_n)_{n \in \mathbb{N}}$. Let $X = (K \setminus \bigcup_{n \in \mathbb{N}} L_n) \cap \Omega$. For every $x \in X$, fix $0 < r_x \leq 1$ such that $B(x, r_x) \subset \Omega$. One of the following alternatives must occur:

- (1) for any ball L_n such that $L_n \cap B(x, r_x) \neq \emptyset$, it holds that $|L_n| \leq r_x$, or
- (2) there exists $n_x \in \mathbb{N}$ such that $L_{n_x} \cap B(x, r_x) \neq \emptyset$ and $|L_{n_x}| \geq r_x$.

Consider the set S_1 be the set of points in X for which the first alternative holds. By Lemma 3.1 applied with $v = 1$, it is possible to extract from the covering of S_1 , $\{B(x, r_x), x \in S_1\}$, $Q_{d,1}$ families of pairwise disjoint balls, $\mathcal{F}_1, \dots, \mathcal{F}_{Q_{d,1}}$, such that

$$S_1 \subset \bigcup_{1 \leq i \leq Q_{d,1}} \bigcup_{L \in \mathcal{F}_i} L.$$

Now, any ball L_n intersecting a ball $L \in \bigcup_{1 \leq i \leq Q_{d,1}} \mathcal{F}_i$ must satisfy $|L_n| \leq |L|$. In particular, since for any $1 \leq i \leq Q_{d,1}$ the balls of \mathcal{F}_i are pairwise disjoint, applying Lemma 3.2 to the balls of \mathcal{F}_i intersecting L_n we get that L_n intersects at most $Q_{d,1}$ balls of \mathcal{F}_i , hence at most $Q_{d,1}^2$ balls of $\bigcup_{1 \leq i \leq Q_{d,1}} \mathcal{F}_i$.

Let $L \in \bigcup_{1 \leq i \leq Q_{d,1}} \mathcal{F}_i$. One aims at replacing all the balls L_n intersecting L by the ball $2L$.

For any $1 \leq i \leq Q_{d,1}$ and any ball $L \in \mathcal{F}_i$, denote by \mathcal{G}_L the set of balls L_n intersecting L . Since $E \subset \bigcup_{n \in \mathbb{N}} L_n$ and $\mu(E) = \mu(\Omega)$, one has $E \cap L \subset \bigcup_{B \in \mathcal{G}_L} B$ and $\mu(E \cap L) = \mu(L)$. By Definition 2.4 and Proposition 4.2, this implies that

$$(80) \quad \kappa(d, \mu, \varepsilon) |L|^s \leq \mathcal{H}_\infty^{\mu, s}(L) \leq \sum_{B \in \mathcal{G}_L} \mathcal{H}_\infty^{\mu, s}(B) \leq \sum_{B \in \mathcal{G}_L} |B|^s.$$

Replace the balls of \mathcal{G}_L by the ball $\hat{L} = 2L$ (recall that $\bigcup_{B \in \mathcal{G}_L} B \subset 2L$). The new sequence of balls so obtained by the previous construction applied to all the balls $L \in \bigcup_{1 \leq i \leq Q_{d,1}} \mathcal{F}_i$ is denoted by $(\hat{L}_k)_{1 \leq k \leq K}$, where $0 \leq K \leq +\infty$.

It follows from the construction and (80) that $S_1 \subset \bigcup_{1 \leq k \leq K} \hat{L}_k$ and

$$(81) \quad \sum_{1 \leq k \leq K} \left(\frac{|\hat{L}_k|}{2} \right)^s \leq \frac{Q_{d,1}^2}{\kappa(d, \mu, \varepsilon)} \sum_{n \geq 0} |L_n|^s.$$

On the other hand, since for any $x \in S_2 = X \setminus S_1$, there exists $n_x \in \mathbb{N}$ such that $L_{n_x} \cap B(x, r_x) \neq \emptyset$ and $r_x \leq |L_{n_x}|$, one has $S_2 \subset \bigcup_{n \in \mathbb{N}} 2L_n$, so that

$$\left(\bigcup_{n \in \mathbb{N}} L_n \right) \cup \left(K \cap \Omega \setminus \bigcup_{n \in \mathbb{N}} L_n \right) \subset \left(\bigcup_{1 \leq k \leq K} \widehat{L}_k \right) \cup \left(\bigcup_{n \in \mathbb{N}} 2L_n \right).$$

Putting the elements of $(\widehat{L}_k)_{1 \leq k \leq K}$ and $(2L_n)_{n \geq 0}$ in a single sequence $(\widehat{L}_n)_{n \geq 0}$, writing $(\widetilde{L}_n := 2\widehat{L}_n)_{n \in \mathbb{N}}$, by construction, $K \cap \Omega \subset \bigcup_{n \in \mathbb{N}} \widetilde{L}_n$ and due to (81):

$$\mathcal{H}_\infty^s(K \cap \Omega) \leq \sum_{n \in \mathbb{N}} |\widetilde{L}_n|^s \leq 2^s \left(\frac{Q_{d,1}^2}{\kappa(d, \mu, \varepsilon)} + 1 \right) \sum_{n \in \mathbb{N}} |L_n|^s \leq 8 \cdot 2^s \frac{Q_{d,1}^2}{\kappa(d, \mu, \varepsilon)} \mathcal{H}_\infty^{\mu,s}(\Omega).$$

□

Remark 4.4. *The proof of Theorem 2.6 only uses Proposition 2.10. In particular, Theorem 2.6 holds for any measure $\mu \in \mathcal{M}(\mathbb{R}^d)$ supported on K and verifying, for any $\underline{i} \in \Lambda^*$, $\mu(f_{\underline{i}}^{-1}(\cdot))$ is absolutely continuous with respect to μ .*

5. APPLICATIONS OF THEOREM 2.2

5.1. Mass transference principle for self-similar measures.

5.1.1. *Proof of Theorem 2.10.* Let μ be a self-similar measure with support K , and set $\alpha = \dim(\mu)$. Let $(B_n := B(x_n, r_n))_{n \in \mathbb{N}}$ be a sequence of balls such that $x_n \in K$ for all $n \in \mathbb{N}$, $\lim_{n \rightarrow +\infty} r_n = 0$ and $\mu(\limsup_{n \rightarrow +\infty} B_n) = 1$.

Fix $\varepsilon > 0$, $v > 1$ and $\delta \geq 1$ and set $\mathcal{B}_v = \{vB_n\}_{n \in \mathbb{N}}$. Proposition 2.1 shows that \mathcal{B}_v is μ -a.c. Then, by Proposition 4.2, for n large enough, one has

$$\mathcal{H}_\infty^{\mu, \alpha - \varepsilon}((v\overset{\circ}{B}_n)^\delta) \geq \kappa(d, \mu, \varepsilon)(vr_n)^{\delta(\alpha - \varepsilon)} \geq (vr_n)^{\delta(\alpha - \frac{\varepsilon}{2})}.$$

Consequently,

$$t(\mu, \delta, \varepsilon, \mathcal{B}_v) = \limsup_{n \rightarrow +\infty} \frac{\log \mathcal{H}_\infty^{\mu, \alpha - \varepsilon}((v\overset{\circ}{B}_n)^\delta)}{\delta \log |vB_n|} \leq \alpha - \frac{\varepsilon}{2}$$

so $t(\mu, \delta, \varepsilon, \mathcal{B}_v) \leq \alpha$. Due to Corollary 2.5, one concludes that

$$\dim_H(\limsup_{n \rightarrow +\infty} (vB_n)^\delta) \geq \frac{\alpha}{\delta}.$$

But for any $\varepsilon' > 0$, $\limsup_{n \rightarrow +\infty} (vB_n)^\delta \subset \limsup_{n \rightarrow +\infty} B_n^{\delta - \varepsilon'}$, so that

$$\dim_H(\limsup_{n \rightarrow +\infty} B_n^{\delta - \varepsilon'}) \geq \frac{\alpha}{\delta}.$$

It follows that for any $\varepsilon' > 0$ and $\delta \geq 1$ one has

$$\dim_H(\limsup_{n \rightarrow +\infty} B_n^\delta) \geq \frac{\alpha}{\delta + \varepsilon'}.$$

Letting $\varepsilon' \rightarrow 0$ proves that $\dim_H(\limsup_{n \rightarrow +\infty} B_n^\delta) \geq \frac{\dim(\mu)}{\delta}$, hence the result.

Remark 5.1. *If the sequence of balls $(B_n)_{n \in \mathbb{N}}$ is not assumed to be μ -a.c, but only to verify $\mu(\limsup_{n \rightarrow +\infty} B_n) = 1$, then the same lower-bound estimate holds for $\dim_H(\limsup_{n \rightarrow +\infty} B_n^\delta)$, but the existence of a gauge function as in Theorem 2.2 does not hold in general.*

- Let us also notice that the computation in the proof of Theorem 2.10 actually shows that, under the assumption that $\lim_{n \rightarrow +\infty} \frac{\log \mu(B_n)}{\log |B_n|} = \dim(\mu)$, it holds that, for n large enough, $\mathcal{H}_\infty^{\mu, s}(B_n^\delta) \geq \mu(B_n) \Leftrightarrow s < \frac{\dim(\mu)}{\delta}$.

5.1.2. *Proof of Theorem 2.12.* Given $\tau_1 = 1 \leq \tau_2 \leq \dots \leq \tau_d$ and $s \geq 0$, set $\tau = (\tau_1, \dots, \tau_d)$ and

$$g_\tau(s) = \max_{1 \leq k \leq d} \left\{ s\tau_k - \sum_{1 \leq i \leq k} \tau_k - \tau_i \right\}.$$

We will need the following lemma (one refers to [28], Proposition 2.1 for the proof, although it is stated in terms of singular values functions).

Lemma 5.2. *Let $\tau_1 = 1 \leq \tau_2 \leq \dots \leq \tau_d$.*

The are two positive constants C_1 and C_2 depending on d only such that for all $s \geq 0$, $r > 0$ and $x \in \mathbb{R}^d$ one has

$$C_1 r^{g_\tau(s)} \leq \mathcal{H}_\infty^s(R_\tau(x, r)) = \mathcal{H}_\infty^s(\mathring{R}_\tau(x, r)) \leq C_2 r^{g_\tau(s)}.$$

Recall that K is the closure of its interior, and note that since the weights p_i are taken positive in Definition 2.7, one must have $\mu(\mathring{K}) > 0$.

Denote $\tilde{\mu} = \frac{\mu(\cdot)}{\mu(\mathring{K})}$ and $\alpha = \dim(\mu) = \dim(\tilde{\mu})$. It is easily verified that the computation made in the proof of Theorem 2.6 implies that, for any open set $\Omega \subset \mathring{K}$, there exists a constant $c(d, \mu, s)$ given by Theorem 2.6, so that

$$(82) \quad \begin{cases} c(d, \mu, s) \mathcal{H}_\infty^s(\Omega) \leq \mathcal{H}_\infty^{\tilde{\mu}, s}(\Omega) \leq \mathcal{H}_\infty^s(\Omega) & \text{if } s < \alpha \\ \mathcal{H}_\infty^{\tilde{\mu}, s}(\Omega) = 0 & \text{if } s > \alpha. \end{cases}$$

Also, $\tilde{\mu}$ being absolutely continuous with respect to μ , the sequence $(B_n)_{n \in \mathbb{N}}$ is $\tilde{\mu}$ -a.c. Furthermore, up to a $\tilde{\mu}$ -a.c extraction, we can assume that each ball $(B_n)_{n \in \mathbb{N}}$ is included in \mathring{K} (and we will do so).

Let $\varepsilon > 0$. Set $\mathcal{R} = \{R_n\}_{n \geq 0}$. By Lemma 3.13, up to a $\tilde{\mu}$ -a.c extraction, one can assume that for every $n \in \mathbb{N}$, the ball B_n satisfies

$$\tilde{\mu}(B_n) \leq r_n^{\alpha - \varepsilon}.$$

Setting $\tau' = (\frac{\tau_i}{\tau_1})_{1 \leq i \leq d}$, for all $0 \leq s < \alpha - \varepsilon$, one has

$$g_{\tau'}(s) = \max_{1 \leq k \leq d} \left\{ \frac{s\tau_k - \sum_{1 \leq i \leq k} \tau_k - \tau_i}{\tau_1} \right\}.$$

From equation (82) and Lemma 5.2, one deduces that

$$(83) \quad C_1 c(d, \mu, s) r_n^{\tau_1 g_{\tau'}(s)} \leq \mathcal{H}_\infty^{\tilde{\mu}, s}(R_n).$$

In particular, for any s verifying

$$(84) \quad \tau_1 g_{\tau'}(s) \leq \alpha - \frac{\varepsilon}{2},$$

if $r_n \leq 1$ one has

$$C_1 c(d, \mu, s) r_n^{\alpha - \frac{1}{2}\varepsilon} \leq C_1 c(d, \mu, s) r_n^{\tau_1 g_{\tau'}(s)} \leq \mathcal{H}_\infty^{\tilde{\mu}, s}(R_n).$$

Since $r_n \rightarrow 0$, for n large enough, this yields

$$(85) \quad \tilde{\mu}(B_n) \leq r_n^{\alpha - \varepsilon} \leq C_1 c(d, \mu, s) r_n^{\tau_1 g_{\tau'}(s)} \leq \mathcal{H}_\infty^{\tilde{\mu}, s}(R_n),$$

hence $(B_n)_{n \in \mathcal{N}_{\tilde{\mu}}(\mathcal{B}, \mathcal{R}, s)}$ is $\tilde{\mu}$ -a.c., and $s(\tilde{\mu}, \mathcal{R}, \mathcal{B}) \geq s$.

It remains to note that

$$\begin{aligned}
(84) &\Leftrightarrow \max_{1 \leq k \leq d} \left\{ \frac{s\tau_k - \sum_{1 \leq i \leq k} \tau_k - \tau_i}{\tau_1} \right\} \leq \frac{\alpha - \frac{\varepsilon}{2}}{\tau_1} \\
&\Leftrightarrow \forall 1 \leq k \leq d, \frac{s\tau_k - \sum_{1 \leq i \leq k} \tau_k - \tau_i}{\tau_1} \leq \frac{\alpha - \frac{\varepsilon}{2}}{\tau_1} \\
&\Leftrightarrow \forall 1 \leq k \leq d, s \leq \frac{\alpha - \frac{1}{2}\varepsilon + \sum_{1 \leq i \leq k} \tau_k - \tau_i}{\tau_k} \\
(86) &\Leftrightarrow s \leq \min_{1 \leq k \leq d} \left\{ \frac{\alpha - \frac{1}{2}\varepsilon + \sum_{1 \leq i \leq k} \tau_k - \tau_i}{\tau_k} \right\}.
\end{aligned}$$

Since $\varepsilon > 0$ was arbitrary, this implies that

$$s(\tilde{\mu}, \mathcal{R}, \mathcal{B}) \geq \min_{1 \leq k \leq d} \left\{ \frac{\alpha + \sum_{1 \leq i \leq k} \tau_k - \tau_i}{\tau_k} \right\},$$

and applying Theorem 2.2 gives the desired lower bound estimate.

Remark 5.3. *Note that the estimates made in the proof of Theorem 2.12, together with Lemma 5.2, can be used to show that, under the assumption that $\lim_{n \rightarrow \infty} \frac{\log \mu(B_n)}{\log |B_n|} = \dim(\mu)$, one has the following properties:*

If $s < \min_{1 \leq k \leq d} \left\{ \frac{\dim(\mu) + \sum_{1 \leq i \leq k} \tau_k - \tau_i}{\tau_k} \right\}$ then, for $n \in \mathbb{N}$ large enough, $\mathcal{H}_{\infty}^{\mu, s}(R_n) \geq \mu(B_n)$. If $s > \min_{1 \leq k \leq d} \left\{ \frac{\dim(\mu) + \sum_{1 \leq i \leq k} \tau_k - \tau_i}{\tau_k} \right\}$ then, for n large enough, $\mathcal{H}_{\infty}^{\mu, s}(R_n) \leq \mu(B_n)$.

5.2. Application to self-similar dynamical coverings.

Proof. In this section, Theorem 2.14 is proved and we adopt the notation of the proof of Proposition 4.2.

Set $s = \dim_H(K)$. Note that, for each $k \in \mathbb{N}$, the set $\{B(f_{\underline{i}}(x), 2|K|c_{\underline{i}})\}_{\underline{i} \in \Lambda^k}$ covers K . In particular, for any measure μ supported on K , the family $\{B(f_{\underline{i}}(x), 2|K|c_{\underline{i}})\}_{\underline{i} \in \Sigma^*}$ is μ -a.c.

Set $B_{\underline{i}} = B(f_{\underline{i}}(x), 2|K|c_{\underline{i}})$. One now focuses on proving that, for any $\delta \geq 1$, $\dim_H(\limsup_{\underline{i} \in \Sigma^*} B_{\underline{i}}^{\delta}) = \frac{s}{\delta}$. If this holds, since for any $\varepsilon > 0$,

$$(87) \quad \limsup_{\underline{i} \in \Sigma^*} B_{\underline{i}}^{\delta + \varepsilon} \subset \limsup_{\underline{i} \in \Sigma^*} B(f_{\underline{i}}(x), c_{\underline{i}}^{\delta}) \subset \limsup_{\underline{i} \in \Sigma^*} B_{\underline{i}}^{\delta},$$

it also holds that $\dim_H(\limsup_{\underline{i} \in \Sigma^*} B(f_{\underline{i}}(x), c_{\underline{i}}^{\delta})) = \frac{s}{\delta}$.

Note that s satisfies the equation $\sum_{1 \leq i \leq m} c_i^s = 1$. Let also be ν_s , the measure on $(\Sigma, \mathcal{B}(\Sigma))$ associated with the probability vector $(p_i = c_i^s)_{1 \leq i \leq m}$ and μ_s its projection on K by the canonical coding map.

Let $0 < t < \min_{1 \leq i \leq m} c_i$ and

$$\Lambda_t^{(k)} = \{\underline{i} = (i_1, \dots, i_\ell) \in \Lambda^* : c_{i_\ell} t^k < c_{\underline{i}} \leq t^k\}.$$

If $\underline{i} \in \Lambda_t^{(k)}$, then for any $\ell \in \Lambda$, the word $\underline{i}\ell \notin \Lambda_t^{(k)}$. This implies that for any $\underline{i} \neq \underline{j} \in \Lambda_t^{(k)}$, $[\underline{i}] \cap [\underline{j}] = \emptyset$.

Then for any $\delta \geq 1$, one has, for any $\varepsilon > 0$,

$$\begin{aligned} \sum_{k \geq 0} \sum_{i \in \Lambda_t^{(k)}} (|B_i|^\delta)^{\frac{s+\varepsilon}{\delta}} &= (4|K|)^{s+\varepsilon} \sum_{k \geq 0} \sum_{i \in \Lambda_t^{(k)}} c_i^{s+\varepsilon} \\ &\leq (4|K|)^{s+\varepsilon} \sum_{k \geq 0} \sum_{i \in \Lambda_t^{(k)}} t^{k\varepsilon} \nu_s([i]). \end{aligned}$$

Since $\sum_{i \in \Lambda_t^{(k)}} \nu_s([i]) \leq 1$, one obtains

$$\sum_{k \geq 0} \sum_{i \in \Lambda_t^{(k)}} (|B_i|^\delta)^{\frac{s+\varepsilon}{\delta}} \leq (4|K|)^{s+\varepsilon} \sum_{k \geq 0} t^{k\varepsilon} < +\infty.$$

This shows that

$$(88) \quad \dim_H \left(\limsup_{i \in \Lambda^*} B_i^\delta \right) \leq \frac{s}{\delta}.$$

One now establishes the lower-bound estimate. By the dimension regularity assumption (see Definition 2.9), $\dim_H(\mu_s) = s$. Since $(B_i)_{i \in \Sigma^*}$ is μ_s -a.c, Theorem 2.10 yields $\dim_H(\limsup_{i \in \Sigma^*} B_i^\delta) \geq \frac{s}{\delta}$. \square

5.3. Study of a problem related to a question of Mahler. Let us first notice that by Theorem 21, one has

$$\dim_H \left(\limsup_{B \in \mathcal{Q}} B^\delta \cap K_{1/3}^{(0)} \right) \leq \min \left\{ \frac{1}{\delta}, \frac{\log 2}{\log 3} \right\}.$$

In particular, this proves that the expected upper-bound in Theorem 2.15 stands.

Before showing that the lower-bound also holds, let us start with some facts and remarks.

Remark 5.4. • One has $\mathcal{H}_\infty^{\frac{\log 2}{\log 3}}(K_{1/3}) > 0$ (this is well known and easily follows from the fact that $K_{1/3}$ carries an Alfhors regular measure of dimension $\frac{\log 2}{\log 3}$).

Moreover, for every $k \in \mathbb{N}$, setting $\mathcal{K}_k = \{f_i([0, 1])\}_{i \in \{0, 2\}^k}$, one has

$$(89) \quad 1 = \sum_{I \in \mathcal{K}_k} |I|^{\frac{\log 2}{\log 3}}.$$

• For every $k \in \mathbb{N}$, let us define

$$(90) \quad \Omega_k = \bigcup_{I \in \mathcal{K}_k} \overset{\circ}{I}.$$

Since $\mathcal{H}_\infty^{\frac{\log 2}{\log 3}}(\bigcup_{I \in \mathcal{K}_k} I \setminus \Omega_k) = 0$ (it is a finite set of points), it follows from (89) that

$$(91) \quad C \mathcal{H}_\infty^{\frac{\log 2}{\log 3}}(\Omega_k) \leq \mathcal{H}_\infty^{\frac{\log 2}{\log 3}}(K_{1/3}) \leq \mathcal{H}_\infty^{\frac{\log 2}{\log 3}} \left(\bigcup_{I \in \mathcal{K}_k} I \right) = \mathcal{H}_\infty^{\frac{\log 2}{\log 3}}(\Omega_k) \leq 1,$$

with $C = \mathcal{H}_\infty^{\frac{\log 2}{\log 3}}(K_{1/3}) > 0$.

• If $n \in \mathbb{N}$ and $T \in \mathcal{T}_n = \{[\frac{k}{3^n}, \frac{k+1}{3^n}[, 0 \leq k \leq 3^n - 1\}$ is a triadic interval of generation n , denote by F_T the canonical homothetical mapping which sends $[0, 1]$

to \bar{T} . For every $I \in \bigcup_{J \in \mathcal{K}_k} F_T(J)$, for all $n \leq k' \leq n+k$ and all $x = (x_n)_{n \in \mathbb{N}} \in \Sigma$ such that $\pi(x) \in I$, one has

$$(92) \quad \frac{S_{n+k'}\phi(x)}{n+k'} = \frac{S_n\phi(x)}{n} \times \frac{n}{n+k'}.$$

We are now ready to finish the proof of Theorem 2.15.

Let $(\varepsilon_q)_{q \in \mathbb{N}}$ be a positive sequence such that $\lim_{q \rightarrow \infty} \varepsilon_q = 0$. One constructs a family $\{U_{p,q,\delta}\}_{\delta \geq 1, q \in \mathbb{N}, 0 \leq p \leq q}$ of open sets as follows: Let $\delta \geq 1$, $q \in \mathbb{N}^*$ and $0 \leq p \leq q$. Consider T a triadic interval of generation $n_q = \lfloor \log_3(q^{2\delta}) \rfloor + 1$ included in $B(\frac{p}{q}, q^{-2\delta})$. Let $N_{p,q,\delta}$ be large enough to ensure that for any $x \in \Sigma$ verifying $\pi(x) \in T$, one has

$$(93) \quad \frac{S_{n_q}\phi(x)}{n_q} \times \frac{n_q}{n_q + N_{p,q,\delta}} \leq \varepsilon_q.$$

Set

$$(94) \quad U_{p,q,\delta} = F_T(\Omega_{N_{p,q,\delta}}).$$

By (92) and (93), for all $x \in U_{p,q,\delta}$ one has

$$\frac{S_{n_q + N_{p,q,\delta}}\phi(x)}{n_q + N_{p,q,\delta}} \leq \varepsilon_q.$$

This implies that

$$\bigcap_{Q \geq 1} \bigcup_{q \geq Q} \bigcup_{0 \leq p \leq q} U_{p,q,\delta} \subset K_{1/3}^{(0)} \cap \bigcap_{Q \geq 1} \bigcup_{q \geq Q} \bigcup_{0 \leq p \leq q} B\left(\frac{p}{q}, q^{-2\delta}\right).$$

Since $U_{p,q,\delta}$ is an homothetic copy of $\Omega_{N_{p,q,\delta}}$ (see (94)), by (91), due to the choice of n_q there exists $\tilde{C} > 0$ independent of p, q and δ such that

$$(95) \quad \mathcal{H}_{\infty}^{\frac{\log 2}{\log 3}}(U_{p,q,\delta}) \geq \tilde{C} q^{-2\delta \frac{\log(2)}{\log(3)}}.$$

For $1 \leq \delta \leq \frac{\log 3}{\log 2}$, it follows that

$$(96) \quad \mathcal{H}_{\infty}^{\frac{\log 2}{\log 3}}(U_{p,q,\delta}) \geq \tilde{C} q^{-2} = \mathcal{L}\left(B\left(\frac{p}{q}, q^{-2}\right)\right).$$

For $\delta \geq \frac{\log 3}{\log 2}$, by concavity of $x \mapsto x^{\frac{\log 3}{\delta \log 2}}$,

$$(97) \quad \mathcal{H}_{\infty}^{\frac{1}{\delta}}(U_{p,q,\delta}) \geq \left(\mathcal{H}_{\infty}^{\frac{\log 2}{\log 3}}(U_{p,q,\delta})\right)^{\frac{\log 3}{\delta \log 2}} \geq \tilde{C} (q^{-2\delta \frac{\log 2}{\log 3}})^{\frac{\log 3}{\delta \log 2}} = \tilde{C} \mathcal{L}\left(B\left(\frac{p}{q}, q^{-2}\right)\right).$$

By Theorem 2.2 applied to $\mathcal{Q} = (B(\frac{p}{q}, \frac{1}{q^{2\delta}}))_{q \in \mathbb{N}^*, 0 \leq p \leq q}$, $\mathcal{U} = (U_{p,q,\delta})_{q \in \mathbb{N}^*, 0 \leq p \leq q}$ and the Lebesgue measure, there exists a gauge function $\zeta : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ satisfying $\lim_{r \rightarrow 0^+} \frac{\log \zeta(r)}{\log r} = s_0$ and

$$\mathcal{H}^{\zeta}\left(\limsup_{q \in \mathbb{N}^*, 0 \leq p \leq q} U_{p,q,\delta}\right) > 0.$$

Since $\limsup_{q \in \mathbb{N}^*, 0 \leq p \leq q} B(\frac{p}{q}, \frac{1}{q^{2\delta}}) \subset \{x : \xi(x) \geq 2\delta\}$, for $\delta \in [1, \frac{\log 3}{\log 2}[$ one gets

$$\dim_H \left(\{x : \xi(x) \geq 2\delta\} \cap K_{1/3}^{(0)} \right) = \frac{\log 2}{\log 3}.$$

For $\delta \geq \frac{\log 3}{\log 2}$, observe that

$$\{x : \xi(x) = 2\delta\} = \limsup_{q \in \mathbb{N}^*, 0 \leq p \leq q} B\left(\frac{p}{q}, \frac{1}{q^{2\delta}}\right) \setminus \bigcup_{\delta' > \delta} \limsup_{q \in \mathbb{N}^*, 0 \leq p \leq q} B\left(\frac{p}{q}, \frac{1}{q^{2\delta'}}\right).$$

Since for every $\delta' > \delta$, $\dim_H \limsup_{q \in \mathbb{N}^*, 0 \leq p \leq q} B\left(\frac{p}{q}, \frac{1}{q^{2\delta'}}\right) \cap K_{\frac{1}{3}}^{(0)} \leq \frac{1}{\delta'}$ and $\lim_{r \rightarrow 0^+} \frac{\log \zeta(r)}{\log r} = \frac{1}{\delta}$, one also has

$$\begin{aligned} & \mathcal{H}^\zeta(\{x : \xi(x) = 2\delta\} \cap K_{\frac{1}{3}}^{(0)}) \\ &= \mathcal{H}^\zeta\left(\left(\limsup_{q \in \mathbb{N}^*, 0 \leq p \leq q} B\left(\frac{p}{q}, \frac{1}{q^{2\delta}}\right) \setminus \bigcup_{\delta' > \delta} \limsup_{q \in \mathbb{N}^*, 0 \leq p \leq q} B\left(\frac{p}{q}, \frac{1}{q^{2\delta'}}\right)\right) \cap K_{\frac{1}{3}}^{(0)}\right) > 0 \end{aligned}$$

so that

$$\dim_H \left(\{x : \xi(x) = 2\delta\} \cap K_{\frac{1}{3}}^{(0)} \right) = \frac{1}{\delta}.$$

6. REMARKS, COMMENTS AND PERSPECTIVES

6.1. Some remarks about Hausdorff measures associated with MTP. The historical mass transference principle from Beresnevich-Velani (established in [8]) provides estimates on the Hausdorff measure rather than only on the dimension. In the case where the ambient measure is not Alfhors regular, one could also establish the following analog of the Beresnevich-Velani's mass transference principle.

Theorem 6.1. *Let $\mu \in \mathcal{M}(\mathbb{R}^d)$ and $\mathcal{B} = (B_n)_{n \in \mathbb{N}}$ be a μ -a.c. sequence of closed balls of \mathbb{R}^d . Let $\mathcal{U} = (U_n)_{n \in \mathbb{N}}$ be a sequence of open sets such that $U_n \subset B_n$ for all $n \in \mathbb{N}$, and $f : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ a doubling¹ gauge function such that for μ -almost every $x \in \text{supp}(\mu)$,*

$$(98) \quad \lim_{r \rightarrow 0^+} \frac{\mu(B(x, r))}{f(r)} = 0.$$

If $\lim_{n \rightarrow +\infty} \frac{\mu(U_n)}{\mu(B_n)} = 0$ and for every $n \in \mathbb{N}$,

$$\mathcal{H}_\infty^{\mu, f}(U_n) \geq \mu(B_n)$$

then

$$\mathcal{H}^f\left(\limsup_{n \rightarrow +\infty} U_n\right) = +\infty.$$

The only issue with such a formulation is that, in general finding a gauge as in (98) is not an easy task. Only one such non trivial example is known by the author. In the case of self-similar measure μ of dimension $\alpha \geq 0$ associated with

¹To prove a full Hausdorff measure statement, in order to build floors of small sets as in [8], one would need for each $n \in \mathbb{N}$ to find some compact sets $K_n \subset U_n$ with roughly the same essential content as U_n . To be able to do that, Lemma 3.4 is crucial and the doubling assumption is needed in its proof.

an IFS satisfying the open set condition, it was established in [6] that the mapping $f : \mathbb{R}^+ \rightarrow \mathbb{R}^+$

$$f(r) = r^{\alpha - \sqrt{\frac{\log \log |\log r|}{\log |\log r|}}}.$$

satisfies the desired properties. But attractors associated with IFS satisfying the open set condition are known to be Ahlfors-regular, so in this case this result is not useful for the dynamical covering problem we considered. We finally add that, when the measure μ is not doubling, even verifying for $\delta > 1$ $\lim_{n \rightarrow +\infty} \frac{\mu(B_n^\delta)}{\mu(B_n)} = 0$ is not free in general.

In addition, we recall again that it is established in [32] that, given a self-similar IFS S of attractor K , $\mathcal{H}^{\dim(S)}(K) > 0$ if and only if S satisfies the open set condition. This in particular implies that $\mathcal{H}^{\dim_H(H)}(K) = 0$ for every IFS S satisfying the hypothesis of Theorem 2.14 but not the open set condition. Is that true that, in the settings of Theorem 2.14,

$$\mathcal{H}^{\frac{\dim_H(K)}{\delta}}(W(x_0, \delta)) > 0 \Leftrightarrow S \text{ satisfies the open set condition?}$$

6.2. Some perspective in Diophantine approximation. The problem raised by Mahler of understanding the rational approximation in the middle-third Cantor set seems currently out of reach. In this sens, Theorem 2.15 raises the following natural interesting problem suggested by Bugeaud. Fix $\varepsilon > 0$ and denote

$$K_{1/3}^{0,\varepsilon} = \left\{ x \in [0, 1] : \liminf_{k \rightarrow +\infty} \phi_{1,k}(x) = 0 \text{ and } \limsup_{k \rightarrow +\infty} \phi_{1,k}(x) \leq \varepsilon \right\}.$$

Can we get an analog of Theorem 2.15 for $K_{1/3}^{0,\varepsilon}$?

6.3. Study of different dynamical coverings. Theorem 2.14 establishes the Hausdorff dimension of certain dynamical coverings associated with self-similar IFS. In such results, the estimates of the essential Hausdorff content associated with self-similar measures plays a key role. Can we estimate such contents for conformal measures or self-affine measures? If so, can we get an analog of Theorem 2.14 in such cases?

REFERENCES

- [1] D. Allen and S. Bárány. On the Hausdorff measure of shrinking target sets on self-conformal sets. *accepted Mathematika*, 2021.
- [2] S. Baker. Intrinsic Diophantine approximation for overlapping iterated function systems. *submitted*, 2022.
- [3] J. Barral and D. Feng. On multifractal formalism for self-similar measures with overlaps. *Math. Z.*, (298):359–383, 2021.
- [4] J. Barral and S. Seuret. Sums of dirac masses and conditioned ubiquity. *C. R. Acad. Sci. Paris, Sér. I* 339:787–792, 2004.
- [5] J. Barral and S. Seuret. Heterogeneous ubiquitous systems in \mathbb{R}^d and Hausdorff dimensions. *Bull. Brazilian Math. Soc.*, 38(3):467–515, 2007.
- [6] J. Barral and S. Seuret. The multifractal nature of heterogeneous sums of dirac masses. *Math. Proc. Camb. Phil. Soc.*, 144(3):707–727, 2008.
- [7] J. Barral and S. Seuret. Ubiquity and large intersections properties under digit frequencies constraints. *Math. Proc. Camb. Phil. Soc.*, 145(3):527–548, 2008.
- [8] V. Beresnevitch and S. Velani. A mass transference principle and the Duffin-Schaeffer conjecture for Hausdorff measures. *Ann. Math.*, 164(3):22 pages, 2006.

- [9] A.S. Besicovitch. A general form of the covering principle and relative differentiation of additive functions. *Math. Proc. Camb. Phil. Soc.*, 41:103–110, 1945.
- [10] Y. Bugeaud. Metric Diophantine approximation on the middle-third Cantor set. *J. Europ. Math. Soc.*, 2016.
- [11] L. Carleson. *Selected Problems on Exceptional Sets*. Van Nostrand, 1967.
- [12] M. Csörnyei, A. Käenmäki, T. Rajala, and V. Suomala. Upper conical density densities for general measure on \mathbb{R}^n . *Proc. Edinburgh Math. Soc.*, 53:971–992, 2010.
- [13] E. Daviaud. An anisotropic inhomogeneous ubiquity theorem. *Submitted, arXiv:2112.10413*, 2021.
- [14] E. Daviaud. Extraction of optimal sub-sequences of balls and application to optimality estimates of mass transference principles. *Submitted, arXiv:2204.01304*, 2022.
- [15] F. Ekström and T. Persson. Hausdorff dimension of random limsup sets. *Journal of the London Mathematical Society*, 98:661–686, 2018.
- [16] S. Eriksson-Bique. A new Hausdorff content bound for limsup sets. *preprint*, 2022.
- [17] K. Falconer. *Fractal geometry*. John Wiley & Sons, Inc., Hoboken, NJ, second edition, 2003. Mathematical foundations and applications.
- [18] A.-H. Fan, D.J. Feng, and J. Wu. Recurrence, dimension and entropy. *J. London Math. Soc.*, 64 (1):229–244, 2001.
- [19] D. Feng and H. Hu. Dimension theory of iterated function systems. *Comm. Pure Appl. Math.*, 62:1435–1500, 2009.
- [20] D. Feng, E. Jarvenpaa, M. Jarvenpaa, and M. Suomala. Dimensions of random covering sets in Riemann manifolds. *Ann. Probab.*, 46:1542–1596, 2018.
- [21] A. Ghosh and D. Nandi. Diophantine approximations, large intersections and geodesics in negative curvature. *arXiv:1911.01406*, 2019.
- [22] Y. Heurteaux. Estimations de la dimension inférieure et de la dimension supérieure des mesures. *Ann. Inst. H. Poincaré Probab. Statist.*, 34:309–338, 1998.
- [23] R. Hill and S. Velani. The ergodic theory of shrinking targets. *Inv. Math.*, 119:175–198, 1995.
- [24] M. Hochman. On self-similar sets with overlaps and inverse theorems for entropy in \mathbb{R}^d . *To appear in Memoires of the AMS*, 2022.
- [25] J.E. Hutchinson. Fractals and self similarity. *Indiana Univ. Math. J.*, 30:713–747, 1981.
- [26] S. Jaffard. Wavelet techniques in multifractal analysis. In *Fractal Geometry and Applications: A Jubilee of Benoît Mandelbrot, M. Lapidus and M. van Frankenhuysen, Eds., Proc. Symposia in Pure Mathematics*, volume 72(2), pages 91–152. AMS, 2004.
- [27] V. Jarnik. Diophantischen approximationen und Hausdorffsches mass. *Mat. Sbornik*, 36:371–381, 1929.
- [28] H. Koivusalo and M. Rams. Mass transference principle: From balls to arbitrary shapes. *To appear in I.R.M.N*, 2020.
- [29] L. Liao and S. Seuret. Diophantine approximation by orbits of expanding Markov maps. *Ergod. Th. Dyn. Syst.*, 33:585–608, 2013.
- [30] T. Persson and M. Rams. On shrinking targets for piecewise expanding interval maps. *Ergod. Th. Dyn. Syst.*, 37:646–663, 2017.
- [31] A. Rapaport. Proof of the exact overlaps conjecture for systems with algebraic contractions. *Annales Scientifique de l'ENS*, pages 55, 1357–1377, 2022.
- [32] A. Schief. Separation properties for self-similar sets. *Proc. Amer. Math. Soc.*, 122, 1, 1994.
- [33] B. Wang, J. Wu, and J. Xu. Mass transference principle for limsup sets generated by rectangles. *Math. Proc. Camb. Phil. Soc.*, 158:419–443, 2015.