



HAL
open science

Geodesics in first-passage percolation cross any pattern

Antonin Jacquet

► **To cite this version:**

| Antonin Jacquet. Geodesics in first-passage percolation cross any pattern. 2022. hal-03630139v2

HAL Id: hal-03630139

<https://hal.science/hal-03630139v2>

Preprint submitted on 18 Jul 2022 (v2), last revised 3 Mar 2023 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Geodesics in first-passage percolation cross any pattern

Antonin Jacquet*

Abstract

In first-passage percolation, one places nonnegative i.i.d. random variables $(T(e))$ on the edges of \mathbb{Z}^d . A geodesic is an optimal path for the passage times $T(e)$. Consider a local property of the time environment. We call it a pattern. We investigate the number of times a geodesic crosses a translate of this pattern. Under mild conditions, we show that, apart from an event with exponentially small probability, this number is linear in the distance between the extremities of the geodesic.

Contents

1	Introduction and main result	2
1.1	Settings	2
1.2	Patterns	2
1.3	Main result	3
1.4	Some applications	4
1.5	Sketch of the proof	5
1.6	Some tools and notations	7
2	Proofs of generalizations of modification arguments in [5]	8
2.1	Modification proof for the Euclidean length of geodesics	8
2.2	Modification proof for the strict concavity of the expected passage times as a function of the weight shifts	12
3	Unbounded case	14
3.1	Proof of Proposition 1.10 in the unbounded case	14
3.2	Typical boxes crossed by geodesics	21
3.3	Modification argument	24
4	Bounded case	27
4.1	Oriented pattern	27
4.2	Proof of Proposition 1.10 in the bounded case	28
4.3	Properties of a typical box	32
4.4	Modification argument	34
4.4.1	Mains ideas of the proof	34
4.4.2	First modification	35
4.4.3	Construction of π	37
4.4.4	Second modification	39
4.4.5	Proof of the second and third items of Lemma 4.5	42
4.4.6	Every geodesic takes the pattern.	44
A	Construction of the path π for the modification in the unbounded case	47
B	Overlapping pattern	49

*Institut Denis Poisson, UMR-CNRS 7013, Université de Tours, antonin.jacquet@univ-tours.fr

1 Introduction and main result

1.1 Settings

Fix an integer $d \geq 2$. In this article, we consider the model of first passage percolation on the hypercubic lattice \mathbb{Z}^d . We denote by 0 the origin of \mathbb{Z}^d and by \mathcal{E} the set of edges in this lattice. The edges in \mathcal{E} are those connecting two vertices x and y such that $\|x - y\|_1 = 1$. A finite path $\pi = (x_0, \dots, x_k)$ is a sequence of adjacent vertices of \mathbb{Z}^d , i.e. for all $i = 0, \dots, k - 1$, $\|x_{i+1} - x_i\|_1 = 1$. We say that π goes from x_0 to x_k . Sometimes we identify a path with the sequence of the edges that it visits, writing $\pi = (e_1, \dots, e_k)$ where for $i = 1, \dots, k$, $e_i = \{x_{i-1}, x_i\}$. We say that k is the length of π and we denote $|\pi| = k$.

The basic random object consists of a family $T = \{T(e) : e \in \mathcal{E}\}$ of i.i.d. non-negative random variables defined on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$, where $T(e)$ represents the passage time of the edge e . Their common distribution is denoted by F . The passage time $T(\pi)$ of a path $\pi = (e_1, \dots, e_k)$ is the sum of the variables $T(e_i)$ for $i = 1, \dots, k$.

For two vertices x and y , we define the geodesic time

$$t(x, y) = \inf\{T(\pi) : \pi \text{ is a path from } x \text{ to } y\}. \quad (1.1)$$

A self-avoiding path γ such that $T(\gamma) = t(x, y)$ is called a geodesic between x and y .

For the following and for the existence of geodesics, we need some assumptions on F . Let t_{\min} denote the minimum of the support of F . We recall a definition introduced in [8]. A distribution F with support in $[0, \infty)$ is called useful if the following holds:

$$\begin{aligned} F(t_{\min}) &< p_c \text{ when } t_{\min} = 0, \\ F(t_{\min}) &< \vec{p}_c \text{ when } t_{\min} > 0, \end{aligned}$$

where p_c denotes the critical probability for the Bernoulli bond percolation model on \mathbb{Z}^d and \vec{p}_c the critical probability for the oriented Bernoulli bond percolation.

In the whole article, we assume that F has support in $[0, \infty)$, is useful, and that

$$\mathbb{E} \min [T_1^d, \dots, T_{2d}^d] < \infty, \quad (1.2)$$

where T_1, \dots, T_{2d} are independent with distribution F .

As F is useful, $F(0) < p_c$. By Proposition 4.4 in [2], we thus know that geodesics between any points exist with probability one.

1.2 Patterns

Let $\ell^\Lambda \geq 1$ be an integer. We fix $\Lambda = B_\infty(0, \ell^\Lambda) = \{-\ell^\Lambda, \dots, \ell^\Lambda\}^d$ and u^Λ and v^Λ two points on the boundary of Λ , i.e. on the set $\{y \in \mathbb{Z}^d : \|y\|_\infty = \ell^\Lambda\}$. These points u^Λ and v^Λ are called endpoints. Then we fix an event \mathcal{A}^Λ , with positive probability, only depending on the passage time of the edges joining two vertices of Λ . We say that $\mathfrak{P} = (\Lambda, u^\Lambda, v^\Lambda, \mathcal{A}^\Lambda)$ is a pattern. Let $x \in \mathbb{Z}^d$. Define:

- for $y \in \mathbb{Z}^d$, $\theta_x y = y - x$,
- for $e = \{u, v\}$ an edge connecting two vertices u and v , $\theta_x e = \{\theta_x u, \theta_x v\}$.

Similarly, if $\pi = (x_0, \dots, x_k)$ is a path, we define $\theta_x \pi = (\theta_x x_0, \dots, \theta_x x_k)$. Then $\theta_x T$ denotes the environment T translated by $-x$, i.e. the family of random variables indexed by the edges of \mathbb{Z}^d defined for all $e \in \mathcal{E}$ by

$$(\theta_x T)(e) = T(\theta_{-x} e).$$

Let π be a self-avoiding path and $x \in \mathbb{Z}^d$. We say that x satisfies the condition $(\pi; \mathfrak{P})$ if these two conditions are satisfied:

1. $\theta_x \pi$ visits u^Λ and v^Λ , and the subpath of $\theta_x \pi$ between u^Λ and v^Λ is entirely contained in Λ ,
2. $\theta_x T \in \mathcal{A}^\Lambda$.

Note that, if x satisfies the condition $(\pi; \mathfrak{P})$ when π is a geodesic, then $(\theta_x \pi)_{u^\Lambda, v^\Lambda}$ is one of the optimal paths from u^Λ to v^Λ entirely contained in Λ in the environment $\theta_x T$. When the pattern is given, we also say " π takes the pattern in $\theta_{-x} \Lambda$ " for " x satisfies the condition $(\pi; \mathfrak{P})$ ". We denote:

$$N^{\mathfrak{P}}(\pi) = \sum_{x \in \mathbb{Z}^d} \mathbb{1}_{\{x \text{ satisfies the condition } (\pi; \mathfrak{P})\}}. \quad (1.3)$$

Note that the number of terms in this sum is actually bounded from above by the number of vertices in π . If $N^{\mathfrak{P}}(\pi) \geq 1$, we say that π takes the pattern. The aim of the article is to investigate, under reasonable conditions on \mathfrak{P} , the behavior of $N^{\mathfrak{P}}(\gamma)$ for all geodesics γ from 0 to x with $\|x\|_1$ large. The first step is to determine these reasonable conditions, that is why we define the notion of valid patterns.

Definition 1.1. *We say that a pattern is valid if the following two conditions hold:*

- \mathcal{A}^Λ has a positive probability,
- the support of F is unbounded or $u^\Lambda = -\ell^\Lambda \varepsilon_1$ and $v^\Lambda = \ell^\Lambda \varepsilon_1$, where ε_1 is the first coordinate vector.

Remark 1.2. *In the case where the support of F is bounded, the condition on the endpoints of the pattern can be relaxed. See Lemma 4.1 and subsequent remarks.*

1.3 Main result

The main result of this paper is the following.

Theorem 1.3. *Let $\mathfrak{P} = (\Lambda, u^\Lambda, v^\Lambda, \mathcal{A}^\Lambda)$ be a valid pattern and assume that F is useful and satisfies (1.2). Then there exist $\alpha > 0$, $\beta_1 > 0$ and $\beta_2 > 0$ such that for all $x \in \mathbb{Z}^d$,*

$$\mathbb{P}(\exists \text{ a geodesic } \gamma \text{ from } 0 \text{ to } x \text{ such that } N^{\mathfrak{P}}(\gamma) < \alpha \|x\|_1) \leq \beta_1 e^{-\beta_2 \|x\|_1}.$$

The first result of this kind appears in an article by van den Berg and Kesten [8]. Let us recall their setting. Let F be a finite mean distribution on $[0, +\infty)$. Denote by $\mu(F)$ the associated time constant, that is

$$\mu(F) = \lim_{n \rightarrow \infty} \frac{\mathbb{E}[t(0, n\varepsilon_1)]}{n}$$

where t is a first-passage percolation model associated with F and where ε_1 is the first vector of the canonical basis of \mathbb{R}^d . Let \tilde{F} be another finite mean distribution on $[0, +\infty)$. Assume F useful, $F \neq \tilde{F}$ and $d \geq 2$. If \tilde{F} is more variable¹ than F , then $\mu(\tilde{F}) < \mu(F)$. This is the main result of [8]. One of the key intermediate results is the existence of a constant $c > 0$ such that, for all large n ,

$$\mathbb{E}[N^{\mathfrak{P}}(\pi(n))] \geq cn \quad (1.4)$$

where \mathfrak{P} is a properly designed pattern and where $\pi(n)$ is the first geodesic from 0 to $n\varepsilon_1$ (geodesics are ordered in an arbitrary way). This is the content of Proposition 5.22 in [8]. The proof relies on a modification argument.

Estimates similar to (1.4) are established, for different patterns, in [7] (see Lemma 1 in [7] which is used to prove Theorem 1) and in [5] (see Sections 5 and 6). We give more details on the latter below as we wish to strengthen some of their results to illustrate the use of Theorem 1.3.

Establishing the above mentioned results is rather technical and sometimes quite involved. Theorem 1.3 thus answers a need for a result handling any valid pattern. It is moreover stronger on two aspects:

1. It provides an at least linear growth of the number of crossed patterns out of an event of exponentially small probability,
2. It gives the result for all geodesics and not only for a specific geodesic (see Theorem 1.8 for an example where this is useful).

¹One says that \tilde{F} is more variable than F if there exists two random variables T – with distribution F – and \tilde{T} – with distribution \tilde{F} – such that $\mathbb{E}[\tilde{T}|T] \leq T$. See Definition (2.1) and Theorem 2.6 in [8].

A result fulfilling both items above appears in an article by Andjel and Vares [1] for the number of edges with large time crossed by a geodesic.

Theorem 1.4 (Theorem 2.3 in [1]). *Let F be a useful distribution on $[0, +\infty)$ with unbounded support. Then, for each M positive there exists $\varepsilon = \varepsilon(M) > 0$ and $\alpha = \alpha(M) > 0$ so that for all x , we have*

$$\mathbb{P}\left(\exists \text{ geodesic } \pi \text{ from } 0 \text{ to } x \text{ such that } \sum_{e \in \pi} \mathbb{1}_{T(e) \geq M} \leq \alpha \|x\|_1\right) \leq e^{-\varepsilon \|x\|_1}. \quad (1.5)$$

Theorem 1.3 is a generalization of this theorem since, for example, we can take the pattern $\mathfrak{P} = (\{-1, 0, 1\}^d, -\varepsilon_1, \varepsilon_1, \mathcal{A}^\Lambda)$ with \mathcal{A}^Λ the event on which each edge of $\{-1, 0, 1\}^d$ has a passage time greater than M to get this result. The proof of Theorem 1.3 is partly inspired by the proof of this theorem and by the proof of (1.4) in [8].

Even if it is stated for distributions with unbounded support, one can check that Theorem 2.3 in [1] holds for F with bounded support with the same proof. Then, we get the following theorem which is useful in the proof of Theorem 1.3 when the support of F is bounded (Section 4.2).

Theorem 1.5. *Let F be a useful distribution on $[0, +\infty)$ with bounded support. Then, for each M positive such that $F([M, +\infty)) > 0$, there exists $\varepsilon = \varepsilon(M) > 0$ and $\alpha = \alpha(M) > 0$ so that for all x , we have (1.5).*

1.4 Some applications

Several of the main results recently obtained in [5] are based on modification arguments leading to results of the type (1.4). We take advantage of Theorem 1.3 to slightly improve some of these results. The purpose of this section is primarily to illustrate the use of Theorem 1.3, the details of the proofs are postponed to the next section.

Euclidean length of geodesics

Consider the following two assumptions² on the distribution F :

(H1) There exist strictly positive integers k and ℓ and atoms $r'_1, \dots, r'_{k+2\ell}, s'_1, \dots, s'_k$ (not necessarily distinct) such that

$$\sum_{i=1}^{k+2\ell} r'_i = \sum_{j=1}^k s'_j. \quad (1.6)$$

(H2) There exist strictly positive integers k and ℓ and atoms $r < s$ such that $(k + 2\ell)r = ks$.

Note that (H2) is stronger than (H1) and that (H1) holds as soon as 0 is an atom. For $x \in \mathbb{Z}^d$, we denote by $\underline{L}_{0,x}$ (resp. $\overline{L}_{0,x}$) the minimal (resp. maximal) Euclidean length of self-avoiding geodesics from 0 to x . In [5], Krishnan, Rassoul-Agha and Seppäläinen prove the following theorem.

Theorem 1.6 (Theorem 6.2 in [5]). *Assume that $\mathbb{P}(T(e) = t_{\min}) < p_c$ and $\mathbb{E} \min [T_1^p, \dots, T_{2d}^p] < \infty$ with $p > 1$. Furthermore, assume one of the following two assumptions:*

- *the support of F is unbounded and (H1) is satisfied,*
- *the support of F is bounded (H2) is satisfied.*

Then, there exist constants $0 < D, \delta, M < \infty$ such that

$$\mathbb{P}(\overline{L}_{0,x} - \underline{L}_{0,x} \geq D \|x\|_1) \geq \delta \text{ for } \|x\|_1 \geq M. \quad (1.7)$$

We use Theorem 1.3 to prove the following result. It generalizes in a way Theorem 1.6 since in the case of bounded support, we have a less restrictive assumption and since the lower bound in (1.8) is exponentially close to one in the distance instead of the uniform lower bound in (1.7). However, the assumption on the moment is less restrictive in Theorem 1.6.

²Assuming that $\mathbb{P}(T(e) = t_{\min}) < p_c$, the assumption (i) or (ii) or (iii) in Assumption 6.1 in [5] is equivalent to the assumption (H1) or (H2). We prove that by distinguishing the case where 0 is an atom or not.

Theorem 1.7. *Assume that F is useful and $\mathbb{E} \min [T_1^d, \dots, T_{2d}^d] < \infty$. Furthermore, assume (H1). Then there exist constants $0 < \beta_1, \beta_2, D < \infty$ such that*

$$\mathbb{P}(\overline{L}_{0,x} - \underline{L}_{0,x} \geq D\|x\|_1) \geq 1 - \beta_1 e^{-\beta_2 \|x\|_1}. \quad (1.8)$$

The proof of this theorem is the aim of Section 2.1.

Strict concavity of the expected passage times as a function of the weight shifts

For $b \in \mathbb{R}$, define the b -shifted weights by

$$T^{(b)} = \{T^{(b)}(e) : e \in \mathcal{E}\} \text{ with } T^{(b)}(e) = T(e) + b \text{ for all } e \in \mathcal{E}.$$

Following the notations of [5] (see Section 2.2 in [5]), all the quantities associated with the passage times $T^{(b)}$ acquire the superscript. For example, $t^{(b)}(x, y)$ is the geodesic time between x and y defined at (1.1), where the infimum is only on self-avoiding paths. Theorem A.1 in [5] gives the existence of a constant $\varepsilon_0 > 0$ with which we have an extension of the Cox-Durrett shape theorem for the shifted weights $T^{(-b)}$ for $b < t_{\min} + \varepsilon_0$ (note that here the weights can be negative). Note that (ii) in Theorem A.1 in [5] guarantees that $\mathbb{E}[t_{0,x}^{(-b)}]$ is finite if $b \in (0, t_{\min} + \varepsilon_0)$.

Theorem 1.8. *Assume F useful. Furthermore, assume that the support of F is bounded and that it contains at least two strictly positive reals. Then, there exists a finite positive constant M and a function $D(b) > 0$ of $b > 0$ such that the following bounds hold for all $b \in (0, t_{\min} + \varepsilon_0)$ and all $\|x\|_1 \geq M$:*

$$\mathbb{E}[t_{0,x}^{(-b)}] \leq \mathbb{E}[t_{0,x}] - b\mathbb{E}[\overline{L}_{0,x}] - D(b)b\|x\|_1. \quad (1.9)$$

Remark 1.9. *In Theorem 1.8, we strengthen very slightly Theorem 5.4 in [5] in the bounded case. Indeed, $\mathbb{E}[\underline{L}_{0,x}]$ in [5] is replaced by $\mathbb{E}[\overline{L}_{0,x}]$ in (1.9). This strengthening is made possible by the fact that Theorem 1.3 gives a result for all geodesics and thus, in particular, for the geodesic of maximal Euclidean length. We focus on the bounded case in Theorem 1.8 since Theorem 5.4 in [5] already contains (1.9) in the unbounded case.*

1.5 Sketch of the proof

Although they share some similarities, the proofs of Theorem 1.3 differ according to whether the support of F is bounded or unbounded. As the proof is easier in the unbounded case, we decide to first give the proof in the unbounded case in Section 3 and then give the proof in the bounded case in Section 4. In both cases, the aim is to prove the following proposition which gives us the main result by a standard re-normalization argument.

Proposition 1.10. *Let $\mathfrak{P} = (\Lambda, u^\Lambda, v^\Lambda, \mathcal{A}^\Lambda)$ be a valid pattern and assume that F is useful and satisfies (1.2). Then there exist $C > 0$ and $D > 0$ such that for all $n \geq 0$, for all x such that $\|x\|_1 = n$,*

$$\mathbb{P}(\exists \text{ a geodesic } \gamma \text{ from } 0 \text{ to } x \text{ such that } N^{\mathfrak{P}}(\gamma) = 0) \leq D e^{-Cn^{\frac{1}{d}}}. \quad (1.10)$$

In what follows we give an informal sketch of the proof of Proposition 1.10. First, assume that we are in the case where geodesics between all points exist with probability one and are unique. We fix 0 and x and we denote by γ the geodesic between 0 and x . We begin by associating to γ a sequence of $q = C\|x\|_1$ disjoint boxes crossed by γ (for a suitable constant $C > 0$). Out of a very low probability event, we can ensure that these boxes are typical in the sense that the passage time are particularly well controlled by μ (see (1.15)). The initial naive intuition is the following.

1. In every such box, the probability that γ takes the pattern is bounded away from 0.
2. Since there are $C\|x\|_1$ boxes, the probability that γ does not take the pattern in any box is exponentially small in $\|x\|_1$.

Let us be slightly more formal. For all $\ell \in \{1, \dots, q\}$, we define the event:

$$\mathcal{M}(\ell) = \{\text{the geodesic from } 0 \text{ to } x \text{ does not take the pattern in the first } \ell \text{ boxes of the sequence}\}.$$

Then, it is sufficient to prove, for all $\ell \geq 1$ and for a constant $\eta > 0$,

$$\mathbb{P}(\mathcal{M}(\ell)) \leq \frac{1}{1 + \eta} \mathbb{P}(\mathcal{M}(\ell - 1)). \quad (1.11)$$

Indeed, by induction and as there are q boxes, this yields

$$\mathbb{P}(\gamma \text{ does not take the pattern}) \leq \mathbb{P}(\mathcal{M}(q)) \leq \left(\frac{1}{1 + \eta}\right)^q$$

and thus the result is established. Since $\mathcal{M}(\ell) \subset \mathcal{M}(\ell - 1)$, (1.11) is equivalent to:

$$\mathbb{P}(\mathcal{M}(\ell - 1) \setminus \mathcal{M}(\ell)) \geq \eta \mathbb{P}(\mathcal{M}(\ell)). \quad (1.12)$$

So it remains to prove (1.12). Here we use a modification argument. The idea is, starting from an initial environment T , to build a new environment T^* – with the same distribution³ as T – by resampling the passage time of some edges of the ℓ -th box. The aim is to get:

$$\mathbb{P}(T^* \in \mathcal{M}(\ell - 1) \setminus \mathcal{M}(\ell) | T) \geq \eta \text{ on } \{T \in \mathcal{M}(\ell)\}, \quad (1.13)$$

where we use the notation $\{T \in \mathcal{M}(j)\}$ to denote "the event $\mathcal{M}(j)$ holds with respect to the environment T ". The inequality (1.12) is a direct consequence of (1.13) by integrating. Thus the aim is to bound from below the probability, conditionally to T , to have the two following properties on the event $\{T \in \mathcal{M}(\ell)\}$:

1. $T^* \notin \mathcal{M}(\ell)$,
2. $T^* \in \mathcal{M}(\ell - 1)$.

In particular, even for the first item, if we denote the ℓ -th box by B , it is not sufficient to prove that, in the environment T^* (that is, after the modification in the ℓ -th box), the geodesic takes the pattern in B . Indeed, we still need B to be the ℓ -th box. Thus, one need to be able to get a global control of how the geodesic is modified in the environment T^* . In particular, one need to control the location of the sequence of boxes and this becomes significantly more difficult when there is no uniqueness of the geodesics and if we want a result valid for all geodesics. To overcome this difficulty, in the definition of the sequence of boxes, we define and use concentric annuli (see Section 3.1).

Comparison with the plan of the proof of Proposition 5.22 in [8]. Let us compare the above strategy with the plan used in [8] to prove Proposition 5.22. Fix x in \mathbb{Z}^d . We shorten geodesic from 0 to x in geodesic. In [8], van den Berg and Kesten also start by associating with some specific geodesic γ some sequence of $q = C\|x\|_1$ typical boxes. By simple geometric arguments, they then get some family \mathcal{B} of boxes such that

$$\mathbb{E}[\text{number of boxes of } \mathcal{B} \text{ which are typical and crossed by } \gamma] \geq c\|x\|_1$$

where c is a positive constant. Fix some box $B \in \mathcal{B}$. Then they also define a new environment T^* by resampling the times of the edges in B . It is then sufficient (this is the technical part of the proof in the bounded case) to prove

$$\mathbb{P}(\text{every geodesic in } T^* \text{ crosses the pattern in } B | \text{in the environment } T, \gamma \text{ crosses } B \text{ and } B \text{ is typical}) \geq \eta$$

for some positive constant $\eta > 0$. In particular, and contrary to what happens in our framework, it is not necessary in this setting to control what happens to geodesic(s) outside the considered box when we resample the times of the edges in the box.

³In fact, the ℓ -th box has to be fixed to get that T and T^* have the same distribution.

Comparison with the plan of the proof of Theorem 2.3 in [1]. In [1], the main difference with the strategy described above is the use of penalized geodesics. Indeed, Andjel and Vares only consider geodesics which do not take edges whose passage time is greater than M and it allows them to get a result on all geodesics from 0 to x thanks to the modification argument. However, it seems difficult to use penalized geodesics with the patterns, that is why we use the strategy of concentric annuli developed in Section 3.1.

1.6 Some tools and notations

In this subsection, we recall some results and fix some notations. First, we denote by \mathbb{N} the set of all non-negative integers, by \mathbb{N}^* the set $\mathbb{N} \setminus \{0\}$, and by \mathbb{R}_+ the set of all $x \in \mathbb{R}$ such that $x \geq 0$.

For a self-avoiding⁴ path $\pi = (x_0, \dots, x_k)$ going from x_0 to x_k , we say that x_i is visited by π before x_j if $i < j$; we say that an edge $\{x_i, x_{i+1}\}$ is visited before an edge $\{x_j, x_{j+1}\}$ if $i < j$. A subpath of π going from x_i to x_j (where $i, j \in \{0, \dots, k\}$ and $i < j$) is the path (x_i, \dots, x_j) and is denoted by π_{x_i, x_j} .

Then, we define different balls in \mathbb{Z}^d or \mathbb{R}^d . Note that all of these balls (or their intersections with \mathbb{Z}^d) are defined as sets of vertices but sometimes, we also want to say that an edge is contained in one of these balls. So, we say that an edge $e = \{u, v\}$ is contained in a ball if u and v are in this ball. For all $c \in \mathbb{Z}^d$ and $r \in \mathbb{R}_+$, we denote

$$\begin{aligned} B_\infty(c, r) &= \{u \in \mathbb{Z}^d : \|u - c\|_\infty \leq r\}, \\ B_1(c, r) &= \{u \in \mathbb{Z}^d : \|u - c\|_1 \leq r\}, \end{aligned}$$

and for $n \in \mathbb{N}^*$, we denote by Γ_n the boundary of $B_1(0, n)$, i.e.

$$\Gamma_n = \{u \in \mathbb{Z}^d : \|u\|_1 = n\}. \quad (1.14)$$

Also for $c \in \mathbb{Z}^d$ and $r \in \mathbb{R}_+$, we denote by $B(c, r)$ the random ball

$$B(c, r) = \{u \in \mathbb{Z}^d : t(c, u) \leq r\}.$$

Then, for x and y in \mathbb{R}^d , we define $t(x, y)$ as $t(x', y')$ where x' is the unique vertex in \mathbb{Z}^d such that $x \in x' + [0, 1)^d$ (similarly for y'). For $c \in \mathbb{Z}^d$ and $r \in \mathbb{R}_+$, we denote by $\tilde{B}(c, r)$ the random ball

$$\tilde{B}(c, r) = \{y \in \mathbb{R}^d : t(c, y) \leq r\}.$$

Let x in \mathbb{R}^d . Thanks to (1.2), we can define

$$\mu(x) = \lim_{n \rightarrow \infty} \frac{t(0, nx)}{n} \text{ a.s.} \quad (1.15)$$

Thanks to the hypothesis (1.2) and since $F(0) < p_c(\mathbb{Z}^d)$, for all $x \in \mathbb{R}^d \setminus \{0\}$, we have $\mu(x) \in (0, \infty)$. Furthermore, μ is a norm on \mathbb{R}^d and describes the first order of approximation of $\tilde{B}(0, r)$ when r goes to infinity. For $c \in \mathbb{Z}^d$ and $r \in \mathbb{R}_+$, we denote

$$B_\mu(c, r) = \{y \in \mathbb{R}^d : \mu(c - y) \leq r\}.$$

Fix $\mathbf{B} = B_\mu(0, 1)$, then the Cox-Durrett shape theorem (Theorem 2.16 in [2]) guarantees that for each $\varepsilon > 0$,

$$\mathbb{P} \left((1 - \varepsilon)\mathbf{B} \subset \frac{\tilde{B}(0, t)}{t} \subset (1 + \varepsilon)\mathbf{B} \text{ for all large } t \right) = 1. \quad (1.16)$$

One can easily prove that the result (1.16) is equivalent to

$$\lim_{\mu(x) \rightarrow \infty} \frac{t(0, x) - \mu(x)}{\mu(x)} = 0 \text{ a.s.} \quad (1.17)$$

Since μ is a norm, we can fix two constants $c_\mu > 0$ and $C_\mu > 0$ such that for all y in \mathbb{R}^d ,

$$c_\mu \|y\|_1 \leq \mu(y) \leq C_\mu \|y\|_1.$$

⁴The definition can be extended to not necessarily self-avoiding paths by saying that a vertex x is visited by π before y if there exists $i_0 \in \{0, \dots, k\}$ such that $x_{i_0} = x$ and for all $j \in \{0, \dots, k\}$, $x_j = y$ implies that $j > i_0$.

Finally, since F is useful, by Lemma 5.5 in [8], there exist $\delta = \delta(F) > 0$ and $D_0 = D_0(F)$ fixed for the remaining of the article such that for all $u, v \in \mathbb{Z}^d$,

$$\mathbb{P}(t(u, v) \leq (t_{\min} + \delta)\|u - v\|_1) \leq e^{-D_0\|u - v\|_1}. \quad (1.18)$$

Notice that, using the Borel-Cantelli lemma with this result, we get that for all $u \in \mathbb{Z}^d$ different from 0,

$$\mu(u) \geq (t_{\min} + \delta)\|u\|_1. \quad (1.19)$$

2 Proofs of generalizations of modification arguments in [5]

2.1 Modification proof for the Euclidean length of geodesics

To prove Theorem 1.7, we begin by defining the valid pattern in three different cases. Recall that k and ℓ are given by the assumptions of this theorem and that $(\varepsilon_i)_{i \in \{1, \dots, d\}}$ are the vectors of the canonical basis

Case where zero is an atom. In addition to the assumptions of Theorem 1.7, we assume that zero is an atom for F . We set $\ell^\Lambda = \max\left(\left\lceil \frac{k}{2} \right\rceil + 1, \ell + 1\right)$ and we define a pattern in $\Lambda = \{-\ell^\Lambda, \dots, \ell^\Lambda\}^d$.

We denote by π^+ the path going from $-\ell^\Lambda \varepsilon_1$ to $\ell^\Lambda \varepsilon_1$ by $2\ell^\Lambda$ steps in the direction ε_1 and by π^{++} the path going from $-\ell^\Lambda \varepsilon_1$ to $(-\ell^\Lambda + 1)\varepsilon_1$ by one step in the direction ε_1 , then to $(-\ell^\Lambda + 1)\varepsilon_1 + \ell\varepsilon_2$ by ℓ steps in the direction ε_2 , then to $(-\ell^\Lambda + k + 1)\varepsilon_1 + \ell\varepsilon_2$ by k steps in the direction ε_1 , then to $(-\ell^\Lambda + k + 1)\varepsilon_1$ by ℓ steps in the direction $-\varepsilon_2$ and then to $\ell^\Lambda \varepsilon_1$ by $2\ell^\Lambda - k - 1$ steps in the direction ε_1 . We define \mathcal{A}^Λ as follows:

- for all $e \in \pi^+ \cup \pi^{++}$, $T(e) = 0$,
- for all $e \in \Lambda$ which is not in $\pi^+ \cup \pi^{++}$, $T(e) > 0$.

Note that \mathcal{A}^Λ has a positive probability. Then, π^+ and π^{++} are the only two optimal paths from u to v entirely contained in the pattern.

Unbounded case. Here, in addition to the assumptions of Theorem 1.7, we assume that zero is not an atom and that the support of F is unbounded. We set ℓ^Λ as in the previous case and we define a pattern in $\Lambda = \{-\ell^\Lambda, \dots, \ell^\Lambda\}^d$. We also build π^+ and π^{++} as in the previous case. Let u' denote the vertex $(-\ell^\Lambda + 1)\varepsilon_1$ and v' the vertex $(-\ell^\Lambda + k + 1)\varepsilon_1$. Then, we index the edges of $\pi_{u', v'}^+$ and the ones of $\pi_{u', v'}^{++}$ in the order in which they are taken by these paths. We respectively denote them by $(e_i^1)_{i \in \{1, \dots, k\}}$

and $(e_i^2)_{i \in \{1, \dots, k + 2\ell\}}$. We fix $M > (2\ell^\Lambda - k)s'_1 + \sum_{j=1}^k s'_j$ and we define \mathcal{A}^Λ as follows:

- for all $e \in \pi_{-\ell^\Lambda \varepsilon_1, u'}^+ \cup \pi_{v', \ell^\Lambda \varepsilon_1}^+$, $T(e) = s'_1$,
- for all $i \in \{1, \dots, k\}$, $T(e_i^1) = s'_i$ and for all $i \in \{1, \dots, k + 2\ell\}$, $T(e_i^2) = r'_i$, thus

$$T(\pi^+) = T(\pi^{++}),$$

- for all $e \in \Lambda$ which are not in π^+ or π^{++} , $T(e) > M$.

Since the support of F is unbounded, $\mathbb{P}(\mathcal{A}^\Lambda)$ is positive. Note that $M > T(\pi^+)$, thus the optimal paths from u to v entirely contained in the pattern can not take other edges than those in $\pi^+ \cup \pi^{++}$. Hence π^+ and π^{++} are the only two optimal paths from u to v entirely contained in the pattern.

Bounded case. In addition to the assumptions of Theorem 1.7, we assume that zero is not an atom and that the support of F is bounded. We set $t_{\max} = \sup(\text{support}(F))$. We denote $a_{\min} = \min(r'_1, \dots, r'_{k+2\ell}, s'_1, \dots, s'_k)$ and $a_{\max} = \max(r'_1, \dots, r'_{k+2\ell}, s'_1, \dots, s'_k)$. Then, there are at least 2ℓ integers $j \in \{1, \dots, k+2\ell\}$ such that $r'_j < a_{\max}$. Indeed, assume that this is not the case. Then, we have

$$\sum_{i=1}^{k+2\ell} r'_i \geq (k+1)a_{\max} > ka_{\max} \geq \sum_{j=1}^k s'_j,$$

and this contradicts (1.6). Thus, even if it means changing the indexes, we can assume that there exists $i_0 \in \{1, \dots, 2\ell\}$ such that $r'_{i_0} < a_{\max}$ and we denote $t_w = a_{\max} - r'_{i_0} > 0$. We fix $\alpha > 0$ an even integer such that:

$$\alpha > \frac{2ka_{\max}}{t_w}. \quad (2.1)$$

We set $k' = \alpha k$, $\ell' = \alpha \ell$ and $\ell^\Lambda = \max\left(\left\lceil \frac{k'}{2} \right\rceil + 1, \ell' + 1\right)$. Now, we define a pattern in $\Lambda = \{-\ell^\Lambda, \dots, \ell^\Lambda\}^d$. Let π^+ be the path going from $u = -\ell^\Lambda \varepsilon_1$ to $v = \ell^\Lambda \varepsilon_1$ by $2\ell^\Lambda$ steps in the direction ε_1 and π^{++} be the path going from u to $u_1 = (-\ell^\Lambda + 1)\varepsilon_1$ by one step in the direction ε_1 , then to $u_2 = (-\ell^\Lambda + 1)\varepsilon_1 + \ell' \varepsilon_2$ by ℓ' steps in the direction ε_2 , then to $u_3 = (-\ell^\Lambda + k' + 1)\varepsilon_1 + \ell' \varepsilon_2$ by k' steps in the direction ε_1 , then to $u_4 = (-\ell^\Lambda + k' + 1)\varepsilon_1$ by ℓ' steps in the direction $-\varepsilon_2$ and then to v by $2\ell^\Lambda - k' - 1$ steps in the direction ε_1 . Note that π^+ and π^{++} share $2\ell^\Lambda - k'$ edges. Then we index the edges of π_{u_1, u_4}^+ , π_{u_1, u_2}^{++} and π_{u_2, u_3}^{++} in the order in which they are taken by these paths. We respectively denote them by $(e_i^1)_{i \in \{1, \dots, k'\}}$, $(e_i^2)_{i \in \{1, \dots, \ell'\}}$ and $(e_i^3)_{i \in \{1, \dots, k'\}}$. Finally, we index the edges in π_{u_3, u_4}^{++} in the reverse order in which they are taken by this path. We denote them by $(e_i^4)_{i \in \{1, \dots, \ell'\}}$. The idea for the event \mathcal{A}^Λ is just to alternate the atoms on every boundary of the rectangle whose vertices are u_1 , u_2 , u_3 and u_4 . It allows us a better control of the time of a path taking both vertices of π_{u_1, u_4}^+ and vertices of π_{u_2, u_3}^{++} . Furthermore, we manage that each edge of π_{u_1, u_2}^{++} has the same passage time as the edge of π_{u_3, u_4}^{++} with which it shares the coordinates in the direction ε_2 (see Figure 1).

So, we define \mathcal{A}^Λ as follows :

- for all $i \in \{1, \dots, k'\}$, $T(e_i^1) = s'_{i[k]}$ and $T(e_i^3) = r'_{2\ell+i[k]}$ where $i[k]$ is the integer in $\{1, \dots, k\}$ such that $i - i[k]$ is divisible by k ,
- for all $i \in \{1, \dots, \ell'\}$, $T(e_i^2) = r'_{i[2\ell]}$ and $T(e_i^4) = r'_{i[2\ell]}$,
- for all $e \in \pi_{u, u_1}^+ \cup \pi_{u_4, v}^+$, $T(e) = a_{\min}$,
- for all other edges $e \in \Lambda$, $T(e) = a_{\max}$.

Note that \mathcal{A}^Λ has a positive probability and that on this event, $T(\pi^+) = T(\pi^{++})$.

Lemma 2.1. *The paths π^+ and π^{++} belong to the family of optimal paths from u to v which are entirely contained in the pattern.*

Proof of Lemma 2.1. Let us begin by introducing some notations. For $i \in \{1, \dots, d\}$ and a path π , $T_i(\pi)$ denotes the sum of the passage times of the edges of π which are in the direction ε_i . So we have $T(\pi) = T_1(\pi) + \dots + T_d(\pi)$. Furthermore, for $j \in \{-\ell^\Lambda, \dots, \ell^\Lambda\}$, we denote by S_2^j the set of edges which can be written $\{x, x + \varepsilon_2\}$ where the second coordinate of x is equal to j .

Now, let us make a remark useful for the proof. For every vertex $w \in \pi_{u_1, u_2}^{++}$, there is no geodesic from u to w taking edges in π_{u_1, u_4}^+ or in π_{u_2, u_3}^{++} . Let us prove it. Assume for a contradiction the existence of such a geodesic. Denote it by π . Then $T_1(\pi_{u, w}^{++}) < T_1(\pi)$. Indeed, π must make at least two steps in the direction ε_1 . The travel time of one of them is at least a_{\min} and the travel time of the other is positive. Furthermore, $T_2(\pi_{u, w}^{++}) \leq T_2(\pi)$. Indeed, for each $j \in \{0, \dots, \ell' - 1\}$, π has to take an edge in S_2^j . By the definition of the pattern, each edge of π_{u_1, u_2}^{++} has the same passage time as the edge of π_{u_3, u_4}^{++} with which it shares the coordinates in the direction ε_2 , and edges in the direction ε_2 which are not in π^{++} have a passage time greater than or equal to a_{\max} . Hence, since $\pi_{u, w}^{++}$ only takes edges in the directions ε_1 and ε_2 , $T(\pi_{u, w}^{++}) < T(\pi)$. Note that, with the same proof, we have that for every vertex $w \in \pi_{u_3, u_4}^{++}$, there is no geodesic from w to v taking edges in π_{u_1, u_4}^+ or in π_{u_2, u_3}^{++} .




Figure 1: Paths π^+ and π^{++} with their passage times in the pattern for the proof of Theorem 1.7 in the bounded case with $d = 2$, $k = 4$, $\ell = 2$ and $\alpha = 4$.

Now, to prove the lemma, assume for a contradiction that there exists an optimal path π entirely contained in Λ such that

$$T(\pi) < T(\pi^+) = T(\pi^{++}). \quad (2.2)$$

Assume that π does not take edges in π_{u_2, u_3}^{++} . Then

$$T(\pi) \geq T_1(\pi) \geq T_1(\pi^+) = T(\pi^+),$$

which contradicts (2.2). Note that the second inequality comes from the fact that edges in the direction ε_1 whose passage time is strictly smaller than those in π^+ must belong to π_{u_2, u_3}^{++} . Hence $T_2(\pi) \geq T_2(\pi^{++})$. Indeed, since π takes edges in π_{u_2, u_3}^{++} , π has to take at least two edges in S_2^j for all $j \in \{0, \dots, \ell' - 1\}$. However, for all $j \in \{0, \dots, \ell' - 1\}$, the two edges whose passage time is the smallest are those in π^{++} . Thus we have $T(\pi) < T(\pi^{++}) = T_1(\pi^{++}) + T_2(\pi^{++})$ and $T_2(\pi) \geq T_2(\pi^{++})$, so $T_1(\pi) < T_1(\pi^{++})$. Hence, π has to take edges in π_{u_1, u_4}^+ .

Then, we order the elements of the set of edges of π_{u_1, u_4}^+ and π_{u_2, u_3}^{++} taken by π in the order in which π visits them. We have just proved that there exists at least one couple of consecutive edges in this set such that one belongs to π_{u_1, u_4}^+ and the other to π_{u_2, u_3}^{++} . We consider the first such couple of edges and we denote the edge in π_{u_1, u_4}^+ by $e^+ = \{u^+, v^+\}$ and the one in π_{u_2, u_3}^{++} by $e^{++} = \{u^{++}, v^{++}\}$ where u^+ (resp. u^{++}) is visited by π before v^+ (resp. v^{++}). There are two cases.

- The first one is that π visits e^+ before e^{++} . The aim is to prove that $T(\pi_{u, u^{++}}) > T(\pi_{u, u^{++}}^{++})$, which contradicts the optimality of π . First, $\pi_{v^+, u^{++}}$ can not take any edge in π_{u_1, u_2}^{++} . Indeed, if it were the case, then $\pi_{u, u^{++}}$ would be a geodesic from u to a vertex of π_{u_1, u_2}^{++} taking an edge in π_{u_1, u_4}^+ , and this would contradict the remark made at the beginning of the proof. The same argument gives us that $\pi_{v^+, u^{++}}$ can not take any edge in π_{u_3, u_4}^{++} . So, $\pi_{v^+, u^{++}}$ has to take ℓ' edges in the direction ε_2 whose time is greater than or equal to a_{\max} , therefore

$$T_2(\pi_{u, u^{++}}) \geq \ell' a_{\max}.$$

By the construction of the pattern, $\pi_{u, u^{++}}^{++}$ saves a time greater than or equal to $\frac{\alpha t_w}{2}$ in the direction

ε_2 compared with $\pi_{u,u^{++}}$, that is

$$T_2(\pi_{u,u^{++}}) \leq \ell' a_{\max} - \frac{\alpha t_w}{2}.$$

Furthermore, for each edge e in $\pi_{u_2,u^{++}}^{++}$, $\pi_{u,u^{++}}$ has to take an edge in the direction ε_1 such that this edge is the edge $e - \ell' \varepsilon_2 \in \pi_{u_1,u_4}^+$ or such that its passage time is equal to a_{\max} . Since $\pi_{u,u^{++}}$ also has to take an edge which has the same coordinates in the direction ε_1 as those of $\{u, u_1\}$, and thus whose passage time is greater than or equal to a_{\min} , we get

$$T_1(\pi_{u,u^{++}}) \geq a_{\min} + \left\lfloor \frac{\|u_2 - u^{++}\|_1}{k} \right\rfloor \sum_{i=1}^k s'_i.$$

Hence,

$$T(\pi_{u,u^{++}}) \geq T_1(\pi_{u,u^{++}}) + T_2(\pi_{u,u^{++}}) \geq a_{\min} + \ell' a_{\max} + \left\lfloor \frac{\|u_2 - u^{++}\|_1}{k} \right\rfloor \sum_{i=1}^k s'_i,$$

and

$$T(\pi_{u,u^{++}}) \leq a_{\min} + \ell' a_{\max} - \frac{\alpha}{2} t_w + \left\lfloor \frac{\|u_2 - u^{++}\|_1}{k} \right\rfloor \sum_{i=1+2\ell}^{k+2\ell} r'_i.$$

Since $\sum_{i=1+2\ell}^{k+2\ell} r'_i \leq \sum_{i=1}^k s'_i$, we have that

$$T(\pi_{u,u^{++}}) - T(\pi_{u,u^{++}}) \leq \sum_{i=1}^k s'_i - \frac{\alpha t_w}{2} \leq k a_{\max} - \frac{\alpha t_w}{2} < 0 \text{ by (2.1).}$$

- The second case is that π visits e^{++} before e^+ and the aim is to prove that $T(\pi_{u,u^+}^+) < T(\pi_{u,u^+})$, which contradicts (2.2). First, π_{v^{++},u^+} can not take any edge in π_{u_3,u_4}^{++} . Indeed, if it were the case, then $\pi_{v^{++},v}$ would be a geodesic from a vertex of π_{u_3,u_4}^{++} to v containing an edge in π_{u_1,u_4}^+ and this would contradict the remark made at the beginning of the proof. So, since for all $j \in \{0, \dots, \ell' - 1\}$, π_{u,u^+} has to take at least two edges in S_2^j , there are at least ℓ' edges in the direction ε_2 taken by π_{u,u^+} with a passage time greater than or equal to a_{\max} . So we have

$$T_2(\pi_{u,u^+}) \geq \ell' a_{\max} + \frac{\alpha}{2} \sum_{i=1}^{2\ell} r'_i \geq \frac{\alpha t_w}{2} + \alpha \sum_{i=1}^{2\ell} r'_i,$$

where the last inequality comes from the fact that $2\ell a_{\max} \geq t_w + \sum_{i=1}^{2\ell} r'_i$. Furthermore, since for each edge in π_{u_2,u^+}^{++} , π_{u,u^+} has to take this edge or an edge whose time is greater than or equal to a_{\max} ,

$$T_1(\pi_{u,u^+}) \geq a_{\min} + \left\lfloor \frac{\|u^+ - u_1\|_1}{k} \right\rfloor \sum_{i=2\ell+1}^{k+2\ell} r'_i.$$

Thus, since $\alpha \geq \left\lfloor \frac{\|u^+ - u_1\|_1}{k} \right\rfloor$, we get

$$T(\pi_{u,u^+}) \geq T_1(\pi_{u,u^+}) + T_2(\pi_{u,u^+}) \geq a_{\min} + \frac{\alpha t_w}{2} + \left\lfloor \frac{\|u^+ - u_1\|_1}{k} \right\rfloor \sum_{i=1}^{k+2\ell} r'_i. \quad (2.3)$$

Furthermore, we have

$$T(\pi_{u,u^+}^+) \leq a_{\min} + \left\lfloor \frac{\|u^+ - u_1\|_1}{k} \right\rfloor \sum_{i=1}^k s'_i. \quad (2.4)$$

Combining (2.3) and (2.4), and using the equality (1.6) yields

$$T(\pi_{u,u^+}) - T(\pi_{u,u^+}^+) \geq \frac{\alpha t_w}{2} - \sum_{i=1}^k s'_i \geq \frac{\alpha t_w}{2} - k a_{\max} > 0 \text{ by (2.1).}$$

□

Conclusion in the three cases.

Proof of Theorem 1.7. For $x \in \mathbb{Z}^d$, we denote by $\pi(x)$ the first self-avoiding geodesic in the lexicographical order⁵ among those who have the minimal number of edges. We say that two patterns are disjoint if they have no vertex in common. We denote by $\mathcal{N}^{\mathfrak{P}}(\pi(x))$ the maximum number of disjoint patterns defined above in the three cases visited by $\pi(x)$. Simple geometric considerations provide a constant $c > 0$ such that for all path π , $\mathcal{N}^{\mathfrak{P}}(\pi) \geq c \mathcal{N}^{\mathfrak{P}}(\pi)$. Further, note that, in each pattern visited by $\pi(x)$, $\pi(x)$ takes the π^+ segment since the π^+ segment belongs to the set of optimal paths entirely contained in the pattern and is the only optimal path for the norm $\|\cdot\|_1$. We can define a self-avoiding path $\widehat{\pi}(x)$ from 0 to x by replacing each π^+ segment of $\pi(x)$ with the π^{++} segment in each disjoint pattern visited by $\pi(x)$. This path $\widehat{\pi}(x)$ has the same passage time and hence both $\pi(x)$ and $\widehat{\pi}(x)$ are geodesics. Note that in the case where zero is not an atom, $\widehat{\pi}$ is obviously self-avoiding (since every geodesic is self avoiding), and in the case where zero is an atom, $\widehat{\pi}$ is self-avoiding since in the patterns, the passage times of all edges which are not in the π^+ or π^{++} segments are strictly positive, and thus $\pi(x)$ can not visit vertices in the π^{++} segments except those which also belong to the corresponding π^+ segment. We have:

$$|\widehat{\pi}(x)| \geq |\pi(x)| + 2\mathcal{N}^{\mathfrak{P}}(\pi(x)).$$

Finally, we get:

$$\overline{L}_{0,x} \geq |\widehat{\pi}(x)| \geq |\pi(x)| + 2\mathcal{N}^{\mathfrak{P}}(\pi(x)) \geq \underline{L}_{0,x} + 2\mathcal{N}^{\mathfrak{P}}(\pi(x)). \quad (2.5)$$

The hypothesis of Theorem 1.3 are satisfied, thus there exist $\alpha > 0$, $\beta_1 > 0$ and $\beta_2 > 0$ such that for all $x \in \mathbb{Z}^d$,

$$\mathbb{P}(\exists \text{ a geodesic } \gamma \text{ from } 0 \text{ to } x \text{ such that } N^{\mathfrak{P}}(\gamma) < \alpha \|x\|_1) \leq \beta_1 e^{-\beta_2 \|x\|_1}.$$

Then, taking $D = 2\alpha c$, we get by (2.5),

$$\mathbb{P}(\overline{L}_{0,x} - \underline{L}_{0,x} \geq D \|x\|_1) \geq \mathbb{P}(\mathcal{N}^{\mathfrak{P}}(\pi(x)) \geq c\alpha \|x\|_1) \geq \mathbb{P}(N^{\mathfrak{P}}(\pi(x)) \geq \alpha \|x\|_1) \geq 1 - \beta_1 e^{-\beta_2 \|x\|_1}.$$

□

2.2 Modification proof for the strict concavity of the expected passage times as a function of the weight shifts

Proof of Theorem 1.8. Recall that we assume that the support of F is bounded and that there are two different positive points in this support. We set $t_{\max} = \sup(\text{support}(F))$. Let $0 < r < s$ be two points in the support of F . Fix $b \in (0, r)$. Applying Lemma 5.5 in [5], we get positive integers k, ℓ fixed for the rest of the proof such that

$$k(s + \delta) < (k + 2\ell)(r - \delta) < (k + 2\ell)(r + \delta) < k(s - \delta) + (2\ell - 1)b \quad (2.6)$$

holds for sufficiently small $\delta > 0$. Fix $\ell_0^\Lambda = \max\left(\left\lceil \frac{k}{2} \right\rceil + 1, \ell + 1\right)$ and an integer ℓ^Λ such that

$$\ell^\Lambda > (2d + 1)\ell_0^\Lambda. \quad (2.7)$$

We define $\Lambda = \{-\ell^\Lambda, \dots, \ell^\Lambda\}^d$, $u^\Lambda = -\ell^\Lambda \varepsilon_1$ and $v^\Lambda = \ell^\Lambda \varepsilon_1$. We take π^+ the path going from u^Λ to v^Λ by $2\ell^\Lambda$ steps in the direction ε_1 , and π^{++} the one going from u^Λ to $u_1 = (-\ell_0^\Lambda + 1)\varepsilon_1$ by $\ell^\Lambda + 1 - \ell_0^\Lambda$ steps in the direction ε_1 , then going to $u_2 = (-\ell_0^\Lambda + 1)\varepsilon_1 + \ell \varepsilon_2$ by ℓ steps in the direction ε_2 , then to

⁵The lexicographical order is based on the directions of the consecutive edges of the geodesics.

$u_3 = (-\ell_0^\Lambda + k + 1)\varepsilon_1 + \ell\varepsilon_2$ by k steps in the direction ε_2 , then to $u_4 = (-\ell_0^\Lambda + k + 1)\varepsilon_1$ by ℓ steps in the direction $-\varepsilon_2$, and then to v^Λ by $\ell^\Lambda + \ell_0^\Lambda - k - 1$ steps in the direction ε_1 .

For a family $(t_e)_{e \in \mathcal{E}_\Lambda}$ of passage times on the edges of Λ and for a path π , we use the abuse of writing $T(\pi)$ to denote $\sum_{e \in \pi} t_e$. For all $\delta \geq 0$, we consider the set $G(\delta)$ of families $(t_e)_{e \in \mathcal{E}_\Lambda}$ of passage times on the edges of Λ which satisfy the following two conditions:

- for all $e \in \pi^{++}$, $t_e \in [r - \delta, r + \delta]$,
- for all other edges e in Λ , $t_e \in [s - \delta, s + \delta]$.

Then, consider the set H of families $(t_e)_{e \in \mathcal{E}_\Lambda}$ such that:

- (P1) π^+ is the unique optimal path from u^Λ to v^Λ among the paths entirely contained in Λ ,
- (P2) for all path π_1 from a vertex w_1 of the boundary of $\{-\ell_0^\Lambda, \dots, \ell_0^\Lambda\}^d$ to a vertex of the boundary of $\{-\ell^\Lambda, \dots, \ell^\Lambda\}^d$, for all w_2 in $\{-\ell_0^\Lambda, \dots, \ell_0^\Lambda\}^d$, for all path π_2 , optimal for the norm $\|\cdot\|_1$, going from w_1 to w_2 , we have $T(\pi_2) < T(\pi_1)$.

Note that for all $\delta > 0$, since r and s belong to the support of F , we have that $\mathbb{P}((T(e))_{e \in \mathcal{E}_\Lambda} \in G(\delta)) > 0$. Then we have $G(0) \subset H$. Indeed, consider a family $(t_e)_{e \in \mathcal{E}_\Lambda} \in G(0)$. So there are only two different passage times in Λ which are r and s . Assume for a contradiction that (P1) does not hold. Then there exists an optimal path π going from u^Λ to v^Λ , different from π^+ . Recall the notation T_i for $i \in \{1, \dots, d\}$ introduced in the proof of Lemma 2.1. Since π^+ is the unique path taking only edges in the direction ε_1 and since there is no passage time equal to zero in Λ , we have $T_1(\pi) < T(\pi)$. Hence

$$T_1(\pi^+) = T(\pi^+) \geq T(\pi) > T_1(\pi).$$

Since the only edges in the direction ε_1 whose passage time is smaller than s are in π_{u_2, u_3}^{++} , π takes an edge of π_{u_2, u_3}^{++} and thus $T_2(\pi) \geq 2\ell r$. Furthermore

$$T_1(\pi) \geq T_1(\pi^+) - k(s - r).$$

Hence we get

$$T(\pi) \geq T_1(\pi) + T_2(\pi) \geq T_1(\pi^+) + 2\ell r - k(s - r) > T_1(\pi^+) = T(\pi^+),$$

where the strict inequality comes from (2.6). Thus, it contradicts the fact that π is an optimal path and (P1) holds.

Now, to prove that (P2) holds, let π_1 be a path from a vertex w_1 of the boundary of $\{-\ell_0^\Lambda, \dots, \ell_0^\Lambda\}^d$ to a vertex of the boundary of $\{-\ell^\Lambda, \dots, \ell^\Lambda\}^d$ and let w_2 a vertex of $\{-\ell_0^\Lambda, \dots, \ell_0^\Lambda\}^d$. Let π_2 be an optimal path for the norm $\|\cdot\|_1$ going from w_1 to w_2 . Then π_1 has to take at least $\ell^\Lambda - \ell_0^\Lambda$ edges whose passage time is equal to s although π_2 takes at most $2d\ell_0^\Lambda$ edges whose passage time is smaller than or equal to s . Thus

$$T(\pi_1) \geq (\ell^\Lambda - \ell_0^\Lambda)s > 2d\ell_0^\Lambda s \geq T(\pi_2),$$

where the strict inequality comes from (2.7). Hence (P2) holds and $G(0) \subset H$.

Furthermore, H is an open set since for a family $(t_e)_{e \in \mathcal{E}_\Lambda}$ to belong to H , it is required that the time of a finite family of paths is strictly smaller than the time of every path of another finite family of paths. Hence, for $\delta > 0$ small enough, we have

$$G(\delta) \subset H. \tag{2.8}$$

Fix $\delta > 0$ such that (2.6) and (2.8) hold. Now, we denote by $\pi(x)$ the first geodesic from 0 to x in the lexicographical order among those of maximal Euclidean length and we denote by $\mathcal{N}^{\mathfrak{P}}(\pi(x))$ the maximum number of disjoint patterns visited by $\pi(x)$. Recall the existence of a constant $c > 0$ small enough such that for all path π , $\mathcal{N}^{\mathfrak{P}}(\pi) \geq c\mathcal{N}^{\mathfrak{P}}(\pi)$. Consider the pattern $\mathfrak{P} = (\Lambda, u^\Lambda, v^\Lambda, \mathcal{A}^\Lambda)$ with $\mathcal{A}^\Lambda = \{(T(e))_{e \in \mathcal{E}_\Lambda} \in G(\delta)\}$. Since $\mathbb{P}(\mathcal{A}^\Lambda) > 0$, we can apply Theorem 1.3. Let $\alpha, \beta_1, \beta_2 > 0$ be the constants given by Theorem 1.3. Then, we have

$$\begin{aligned} \mathbb{E}[\mathcal{N}^{\mathfrak{P}}(\pi(x))] &\geq [c\alpha\|x\|_1]\mathbb{P}(\mathcal{N}^{\mathfrak{P}}(\pi(x)) \geq c\alpha\|x\|_1) \\ &\geq [c\alpha\|x\|_1]\mathbb{P}(\mathcal{N}^{\mathfrak{P}}(\pi(x)) \geq \alpha\|x\|_1) \geq [c\alpha\|x\|_1](1 - \beta_1 e^{-\beta_2\|x\|_1}) \geq C\|x\|_1. \end{aligned}$$

Now, let us follow the end of Stage 3 of the proof of Theorem 5.4 in [5]. By (P1), $\pi(x)$ takes the π^+ segment in each pattern that it takes. Furthermore, by (P2), $\pi(x)$ does not take any edge in the π^{++} segment which is not in the π^+ segment. So, we can define a self-avoiding path $\widehat{\pi}(x)$ from 0 to x by replacing each π^+ segment with the π^{++} segment in each pattern visited by $\pi(x)$. Reduce the weights on each edge e from $T(e)$ to $T^{(-b)}(e) = T(e) - b$. By the definition of the pattern, the $T^{(-b)}$ -passage times of the segments π^+ and π^{++} obey the inequality:

$$T^{(-b)}(\pi^{++}) = T(\pi^{++}) - b|\pi^{++}| < T(\pi^+) + (2\ell - 1)b - b|\pi^{++}| = T^{(-b)}(\pi^+) - b.$$

Then, following the proof of Theorem 5.4 in [5], we get

$$\begin{aligned} t^{(-b)}(0, x) &\leq T^{(-b)}(\widehat{\pi}(x)) < T^{(-b)}(\pi(x)) - b\mathcal{N}^{\mathfrak{P}}(\pi(x)) \\ &= T(\pi(x)) - b|\pi(x)| - b\mathcal{N}^{\mathfrak{P}}(\pi(x)) \\ &= t(0, x) - b\overline{L}_{0,x} - b\mathcal{N}^{\mathfrak{P}}(\pi(x)). \end{aligned} \tag{2.9}$$

Since $b \in (0, t_{\min} + \varepsilon_0)$, $\mathbb{E}[t_{0,x}^{(-b)}]$ is finite. Thus, taking expectation in (2.9), we get the result. \square

3 Unbounded case

3.1 Proof of Proposition 1.10 in the unbounded case

Let $\mathfrak{P} = (\Lambda, u^\Lambda, v^\Lambda, \mathcal{A}^\Lambda)$ be a valid pattern. Let us begin with the definitions of a typical box and of a successful box. To this end, we have to fix some constants.

Boxes and constants. Recall that t_{\min} , ℓ^Λ and δ have been fixed in the introduction. The minimum of the support of F is denoted by t_{\min} , ℓ^Λ denotes the size of the pattern and δ comes from (1.18). Since

$$\lim_{M \rightarrow \infty} \mathbb{P}(\mathcal{A}^\Lambda \text{ is realized and for all edges } e \in \Lambda, T(e) \leq M) = \mathbb{P}(\mathcal{A}^\Lambda) > 0,$$

there exists a positive constant M^Λ fixed for the rest of the proof such that

$$\mathbb{P}(\mathcal{A}^\Lambda \text{ is realized and for all edges } e \in \Lambda, T(e) \leq M^\Lambda) > 0.$$

Even if it means replacing \mathcal{A}^Λ by $\mathcal{A}^\Lambda \cap \{\forall e \in \Lambda, T(e) \leq M^\Lambda\}$, we can assume that

$$\mathcal{A}^\Lambda \subset \{\forall e \in \Lambda, T(e) \leq M^\Lambda\}.$$

We fix

$$\tau^\Lambda = M^\Lambda \|u^\Lambda - v^\Lambda\|_1, \tag{3.1}$$

which is an upper bound for the travel time of an optimal path (for the passage time) going from u^Λ to v^Λ and entirely contained in Λ on the event \mathcal{A}^Λ .

For $i \in \{1, 2, 3\}$, $B_{i,s,N}$ is the ball in \mathbb{Z}^d of radius $r_i N$ for the norm $\|\cdot\|_1$ centered at the point sN where the constants r_i are defined as follows. We fix $r_1 = d$. Denote by K the number of edges in $B_\infty(0, \ell^\Lambda + 3)$. Then, fix r_2 an integer such that

$$r_2 \delta - r_1(t_{\min} + \delta) - K t_{\min} - \tau^\Lambda > 0. \tag{3.2}$$

Let $r_{2,3}$ be a positive real such that

$$B_{2,0,1} \subset B_\mu \left(0, \frac{r_{2,3}}{2}\right) \cap \mathbb{Z}^d,$$

then we fix r_3 an integer such that

$$B_\mu(0, 9r_{2,3}) \cap \mathbb{Z}^d \subset B_{3,0,1}.$$

We use the word "box" to talk about $B_{3,s,N}$. We denote by $\partial B_{i,s,N}$ the boundary of $B_{i,s,N}$, that is the set of points $z \in \mathbb{Z}^d$ such that $\|z - sN\|_1 = r_i N$. For u and v two vertices contained in $B_{3,s,N}$, we denote by $t_{3,s,N}(u, v)$ the minimum of the times of all paths entirely contained in $B_{3,s,N}$ and going from u to v .

Crossed boxes and weakly crossed boxes. We say that a path

- crosses a box $B_{3,s,N}$ if it visits a vertex in $B_{1,s,N}$,
- weakly crosses a box $B_{3,s,N}$ if it visits a vertex in $B_{2,s,N}$.

Paths associated in a box. We say that two paths γ and γ' from 0 to the same vertex x are associated in a box $B_{3,s,N}$ if there exist two distinct vertices s_1 and s_2 such that the following conditions hold:

- γ and γ' visit s_1 and s_2 ,
- $\gamma_{0,s_1} = \gamma'_{0,s_1}$,
- γ_{s_1,s_2} and γ'_{s_1,s_2} are entirely contained in $B_{3,s,N}$,
- $\gamma_{s_2,x} = \gamma'_{s_2,x}$.

In particular, these two paths coincide outside $B_{3,s,N}$.

Typical boxes. We define a sequence $(\nu(N))_{N \in \mathbb{N}^*}$ such that for all $N \in \mathbb{N}^*$, $\nu(N) > M^\Lambda$ and

$$\lim_{N \rightarrow \infty} \mathbb{P} \left(\sum_{e \in B_{2,0,N}} T(e) \geq \nu(N) \right) = 0. \quad (3.3)$$

Note that $F((\nu(N), +\infty)) > 0$ for all $N \in \mathbb{N}^*$ since the support of F is unbounded.

We can now define typical boxes. A box $B_{3,s,N}$ is typical if it verifies the following properties:

(i) $\mathcal{T}(s; N)$ is realized, where $\mathcal{T}(s; N)$ is the following event:

$$\left\{ \sup_{z \in B_{2,s,N}} t_{3,s,N}(Ns, z) \leq r_{2,3}N \right\} \cap \left\{ \inf_{z \in \partial B_{3,s,N}} t_{3,s,N}(Ns, z) \geq 4r_{2,3}N \right\},$$

(ii) every path π entirely contained in $B_{3,s,N}$ from u_π to v_π with $\|u_\pi - v_\pi\|_1 \geq (r_2 - r_1)N$ has a passage time verifying:

$$t(\pi) \geq (t_{\min} + \delta) \|u_\pi - v_\pi\|_1, \quad (3.4)$$

(iii) $\sum_{e \in B_{2,0,N}} T(e) < \nu(N)$.

Lemma 3.1. *We have these three properties about typical boxes.*

1. Let $s \in \mathbb{Z}^d$ and $N \in \mathbb{N}^*$. If $B_{3,s,N}$ is a typical box, for all points u_0 and v_0 in $B_{2,s,N}$, every geodesic from u_0 to v_0 is entirely contained in $B_{3,s,N}$.
2. Let $s \in \mathbb{Z}^d$ and $N \in \mathbb{N}^*$. The typical box property only depends on the passage times of the edges in $B_{3,s,N}$.
3. We have

$$\lim_{N \rightarrow \infty} \mathbb{P}(B_{3,0,N} \text{ is a typical box}) = 1.$$

This lemma guarantees that the properties of a typical box are indeed typical ones and that they are also local ones. Its proof is in Section 3.2. Let us now introduce some further definitions.

Successful boxes. For a fixed $x \in \mathbb{Z}^d$, a box $B_{3,s,N}$ is successful if every geodesic from 0 to x takes a pattern which is entirely contained in $B_{2,s,N}$, i.e. if for every geodesic γ going from 0 to x , there exists $x_\gamma \in \mathbb{Z}^d$ satisfying the condition $(\gamma; \mathfrak{P})$ and such that $B_\infty(x_\gamma, \ell^\Lambda)$ is contained in $B_{2,s,N}$. Note that the notion of successful box depends on some fixed $x \in \mathbb{Z}^d$.

Annuli. Now, we define the annuli mentioned in Section 1.5. Fix

$$r = 2(r_1 + r_3 + 1), \quad (3.5)$$

and for all integers $i \geq 1$, let us define

$$A_{i,N} = \{y \in \mathbb{Z}^d : \|y\|_1 \in [(i-1)rN, irN)\}.$$

For any annulus $A_{i,N}$, we call $\{y \in \mathbb{Z}^d : \|y\|_1 = (i-1)rN\}$ its inner sphere and $\{y \in \mathbb{Z}^d : \|y\|_1 = irN\}$ its outer sphere. Then, we give two definitions about these annuli which are useful in the proof of Lemma 3.3.

- For $i > 1$, we say that a path from 0 to a vertex of \mathbb{Z}^d crosses (resp. weakly crosses) a box $B_{3,s,N}$ in the annulus $A_{i,N}$ if the two following conditions are satisfied:
 - it crosses (resp. weakly crosses) this box before it visits for the first time the outer sphere of $A_{i,N}$,
 - $B_{3,s,N}$ is entirely contained in the annulus, i.e. every vertex of $B_{3,s,N}$ is in $A_{i,N}$.
- We also say that a path takes a pattern in the annulus $A_{i,N}$ if it takes a pattern which is entirely contained in $A_{i,N}$, i.e. if every vertex of this pattern is in $A_{i,N}$. Here, we do not require that the path takes a pattern before it visits the outer sphere of $A_{i,N}$ for the first time.

Note that the choice of r guarantees that every path passing through an annulus has to cross a box in this annulus.

For all integer $p \geq 2$ and all $N \in \mathbb{N}^*$, we denote by $\mathcal{G}^p(N)$ the event on which for all x in the outer sphere of the p -th annulus, every geodesic from 0 to x crosses a typical box in at least $\lfloor \frac{p}{2} \rfloor$ annuli. The following lemma, whose proof is given in Section 3.2, gives us an exponential decrease of the probability of the complement of $\mathcal{G}^p(N)$.

Lemma 3.2. *There exist two positive constants C_1 and D_1 , and an integer $N_0 \geq 1$ such that for all $p \geq 1$ and $N \geq N_0$,*

$$\mathbb{P}(\mathcal{G}^p(N)^c) \leq D_1 e^{-C_1 p^{\frac{1}{d}}}.$$

Setup for the proof of Proposition 1.10. For the rest of the proof, we fix C_1 , D_1 and N_0 given by Lemma 3.2. Recall that K is the number of edges in $B_\infty(0, \ell^\Lambda + 3)$. Then we fix $\delta' > 0$ such that

$$r_2(\delta - \delta') - r_1(t_{\min} + \delta) - K(t_{\min} + \delta') - \tau^\Lambda > 0. \quad (3.6)$$

Note that it is possible since we have taken r_2 large enough (see (3.2)). Then, fix

$$N \geq \max(N_0, \ell^\Lambda + 3), n \geq 2rN \text{ and } x \in \Gamma_n, \quad (3.7)$$

(where Γ_n is defined at (1.14)). Fix $p = \lfloor \frac{n}{rN} \rfloor$ and $q = \lfloor \frac{p}{2} \rfloor$. Note that x belongs to the $(p+1)$ -th annulus.

M -sequences. Let us now define some random sets and variables which are useful for stability questions for the modification argument. Unless otherwise specified, in the remaining of this section, we write geodesic as a shorthand for geodesic from 0 to x .

First, let us associate a sequence of 0 to $p-1$ typical boxes to every geodesic from 0 to x . For a geodesic γ , the deterministic construction is what follows.

Initialize the sequence as an empty sequence. For j from 1 to $p-1$, do:

- let $a_j(\gamma)$ be the index of the first annulus such that γ crosses a typical box in this annulus and such that $a_j(\gamma) > a_{j-1}(\gamma)$ (where $a_0(\gamma) = 1$). If there is no such annulus, then we stop the algorithm.

- Add the first typical box crossed⁶ by γ in the annulus $A_{a_j(\gamma),N}$ to the sequence. Note that, since this typical box is crossed before γ leaves $A_{a_j(\gamma),N}$ by the outer sphere for the first time, the j -th box of the sequence is crossed by γ after the $(j-1)$ -th one.

So, we get a sequence of at most $p-1$ boxes crossed by the geodesic. These boxes are all in different annuli. Furthermore, every box of this sequence is crossed by γ before γ leaves the annulus containing it for the first time by the outer sphere. If the event $\mathcal{G}^p(N)$ occurs, we know that all these sequences have at least q elements. For $j \in \{1, \dots, p-1\}$, we define a set of geodesics Γ^j . A geodesic γ from 0 to x belongs to Γ^j if:

- the length of its sequences defined above is greater than or equal to j ,
- γ does not take the pattern in any annuli $A_{k,N}$ with $k \leq a_j(\gamma)$.

We call the sequences defined above the M -sequences.

Selected geodesic and S^1 -variables. Then, for $j \in \{1, \dots, p-1\}$, if Γ^j is not empty, we define the selected geodesic among the geodesics of Γ^j as the one which minimizes the index of the annulus containing the j -th box of its sequence. If there are several such geodesics, the selected one is the first in the lexicographical order. Then, the random variable S_j^1 is equal to the vertex s such that the box $B_{3,s,N}$ is the j -th box in the M -sequence of the selected geodesic. When j is fixed, we say that the box $B_{3,S_j^1,N}$ is the selected box. Finally, if Γ^j is empty, set $S_j^1 = 0$ and there is no selected geodesic.

S^2 -variables. After a modification in the box of the selected geodesic, we have to select a new box in the new environment T^* defined below. Since we can not guarantee that the selected box in the initial environment is still typical in the new environment, we have to select it in a different way. That is why we introduce the S^2 -variables. Let $j \in \{1, \dots, p-1\}$, we define S_j^2 as follows. For every geodesic $\gamma \in \Gamma^{j-1}$, we define, if it is possible, $S_j^2(\gamma)$ as the vertex s corresponding to the box $B_{3,s,N}$ where $B_{3,s,N}$ is the first successful box crossed by γ in an annulus (in the sense given with the definition of the annuli above). If it is not possible, $S_j^2(\gamma) = 0$. We denote by $a'_j(\gamma)$ the index of the annulus containing $B_{3,S_j^2(\gamma),N}$. Then, S_j^2 is equal to the vertex $S_j^2(\bar{\gamma})$ where $\bar{\gamma}$ satisfies the three following conditions:

- $a'_j(\bar{\gamma}) > 1$,
- for all geodesic $\gamma \in \Gamma^{j-1}$ such that $a'_j(\gamma) \neq 1$, we have $a'_j(\gamma) \geq a'_j(\bar{\gamma})$,
- $\bar{\gamma}$ is the first geodesic in the lexicographical order among the geodesics γ such that $a'_j(\gamma) = a'_j(\bar{\gamma})$.

If it is not possible, $S_j^2 = 0$.

Modification argument. Finally, for $j \in \{1, \dots, p-1\}$, we define $\mathcal{M}(j)$ as the event on which every geodesic from 0 to x has at least j typical boxes in its M -sequence and there exists one geodesic which does not take the pattern in any annuli $A_{k,N}$ with $k \leq a_j(\gamma)$. We also define $\mathcal{M}(0)$ as the event on which there exists a geodesic from 0 to x . Its probability is equal to 1 (see Section 1.1). Now, the aim is to bound from above $\mathbb{P}(\mathcal{M}(q))$ independently of x since we have:

$$\mathbb{P}(\text{there exists a geodesic } \gamma \text{ from 0 to } x \text{ such that } N^{\mathfrak{B}}(\gamma) = 0) \leq \mathbb{P}(T \in \mathcal{G}^p(N)^c) + \mathbb{P}(T \in \mathcal{M}(q)). \quad (3.8)$$

In the sequel, we introduce an independent copy T' of the environment T , the two being defined on the same probability space. It is thus convenient to refer to the considered environment when dealing with the objects defined above. To this aim, we shall use the notation $\{T \in \mathcal{M}(j)\}$ to denote that the event $\mathcal{M}(j)$ holds with respect to the environment T . In other words, $\mathcal{M}(j)$ is now seen as a subset of $(\mathbb{R}_+)^{\mathcal{E}}$, where \mathcal{E} is the set of all the edges. Similarly, for $i \in \{1, 2\}$ we denote by $S_j^i(T')$ the random variables defined above but in the environment T' .

⁶If a path crosses two boxes $B_{3,s_1,N}$ and $B_{3,s_2,N}$, we say that it crosses $B_{3,s_1,N}$ before $B_{3,s_2,N}$ if it visits a vertex of $B_{1,s_1,N}$ before one of $B_{1,s_2,N}$.

Fix $\ell \in \{1, \dots, q\}$. On $\{T \in \mathcal{M}(\ell)\}$, $\Gamma^\ell \neq \emptyset$ and $B_{3, S_\ell^1(T), N}$ is a typical box crossed by the selected geodesic. We get a new environment T^* defined for all edge e by:

$$T^*(e) = \begin{cases} T(e) & \text{if } e \notin B_{2, S_\ell^1(T), N} \\ T'(e) & \text{else.} \end{cases}$$

For y and z in \mathbb{Z}^d , we denote by $t^*(y, z)$ the geodesic time between y and z in the environment T^* . Note that T and T^* do not have the same distribution.

Lemma 3.3. *There exists $\eta = \eta(N) > 0$ such that for all ℓ in $\{1, \dots, q\}$, there exist measurable functions $E_+^* : \mathbb{R}_+^\mathcal{E} \mapsto \mathcal{P}(\mathcal{E})$ and $E_-^* : \mathbb{R}_+^\mathcal{E} \mapsto \mathcal{P}(\mathcal{E})$ such that:*

(i) $E_+^*(T) \cap E_-^*(T) = \emptyset$ and $E_+^*(T) \cup E_-^*(T) \subset B_{2, S_\ell^1(T), N}$,

(ii) on the event $\{T \in \mathcal{M}(\ell)\}$, we have $\mathbb{P}(T' \in \mathcal{B}^*(T) | T) \geq \eta$ where $\{T' \in \mathcal{B}^*(T)\}$ is a shorthand for

$$\{\forall e \in E_+^*(T), T'(e) \geq \nu(N), \forall e \in E_-^*(T), T'(e) \leq t_{\min} + \delta', \theta_{NS_\ell^1(T)} T' \in \mathcal{A}^\Lambda\},$$

(iii) $\{T \in \mathcal{M}(\ell)\} \cap \{T' \in \mathcal{B}^*(T)\} \subset \{T^* \in \mathcal{M}(\ell - 1) \setminus \mathcal{M}(\ell)\}$ and $\|S_\ell^2(T^*) - S_\ell^1(T)\|_1 \leq 2r_3$.

Remark 3.4. *We have $E_+^*(T) \cup E_-^*(T) \cup (NS_\ell^1(T) + \Lambda) = B_{2, S_\ell^1(T), N}$. Note that since $r_1 = d$ and $N \geq \ell^\Lambda + 3$, we have that $(NS_\ell^1(T) + \Lambda) \subset B_{1, S_\ell^1(T), N}$.*

Lemma 3.3 is a consequence of Lemma 3.5 below whose proof is given in Section 3.3. Recall the definition of associated paths given page 15.

Lemma 3.5. *There exists $\eta = \eta(N) > 0$ such that for all ℓ in $\{1, \dots, q\}$, there exist measurable functions $E_+^* : \mathbb{R}_+^\mathcal{E} \mapsto \mathcal{P}(\mathcal{E})$ and $E_-^* : \mathbb{R}_+^\mathcal{E} \mapsto \mathcal{P}(\mathcal{E})$ such that (i) and (ii) of Lemma 3.3 are satisfied and such that if the event $\{T \in \mathcal{M}(\ell)\} \cap \{T' \in \mathcal{B}^*(T)\}$ occurs, then we have the following properties:*

(i) in the environment T^* , every geodesic from 0 to x takes the pattern inside $B_{2, S_\ell^1(T), N}$,

(ii) for all geodesic $\bar{\gamma}^*$ from 0 to x in the environment T^* , there exists a geodesic $\bar{\gamma}$ from 0 to x in the environment T such that $\bar{\gamma}$ and $\bar{\gamma}^*$ are associated in $B_{3, S_\ell^1(T), N}$,

(iii) there exists a geodesic γ^* in the environment T^* from 0 to x such that γ^* and the selected geodesic γ in the environment T are associated in $B_{3, S_\ell^1(T), N}$.

Proof of Lemma 3.3. Let $\ell \in \{1, \dots, q\}$. Consider E_+^* and E_-^* given by Lemma 3.5. Let $s \in \mathbb{Z}^d$ and assume that the event $\{T \in \mathcal{M}(\ell)\} \cap \{S_\ell^1(T) = s\} \cap \{T' \in \mathcal{B}^*(T)\}$ occurs. To prove that the event $\{T^* \in \mathcal{M}(\ell - 1) \setminus \mathcal{M}(\ell)\}$ occurs and that $\|S_\ell^2(T^*) - s\|_1 \leq 2r_3$, it is sufficient to prove that we have the four following points in the environment T^* :

1. every geodesic from 0 to x has at least $\ell - 1$ typical boxes in its M -sequence,
2. there exists a geodesic from 0 to x which does not take the pattern in the annuli until the one containing its $(\ell - 1)$ -th box,
3. every geodesic from 0 to x whose M -sequence contains at least ℓ elements takes the pattern in an annulus whose index is smaller than or equal to the one containing its ℓ -th box,
4. there exists s' such that $S_\ell^2(T^*) = s'$ and $\|s - s'\|_1 \leq 2r_3$.

Let us start with a few remarks. We denote by a_ℓ the index of the annulus which contains $B_{3, S_\ell^1(T), N}$.

- (a) The environments T and T^* coincides outside the box $B_{2, S_\ell^1(T), N}$. As this box is included in the annulus $A_{a_\ell, N}$, the environments T and T^* are the same in all the other annuli. In particular, any box contained in an annulus $A_{i, N}$ for $i \neq a_\ell$ is typical in T if and only if it is typical in T^* .
- (b) Similarly, every path π takes a pattern which is outside the box $B_{3, S_\ell^1(T), N}$ in the environment T if and only if it takes this pattern in the environment T^* . In particular, for any $i \neq a_\ell$, π takes the pattern in the annulus $A_{i, N}$ in the environment T if and only if it takes the pattern in the annulus $A_{i, N}$ in the environment T^* .

- (c) Let $\bar{\gamma}$ and $\bar{\gamma}^*$ be as in item (ii) of Lemma 3.5. Then $\bar{\gamma}$ and $\bar{\gamma}^*$ coincide except maybe for the part between s_1 and s_2 . To sum up:

$$\bar{\gamma}_{0,s_1} = \bar{\gamma}_{0,s_1}^* \text{ and } \bar{\gamma}_{s_2,x} = \bar{\gamma}_{s_2,x}^* \text{ and } \bar{\gamma}_{s_1,s_2} \subset B_{3,S_\ell^1(T),N} \subset A_{a_\ell,N} \text{ and } \bar{\gamma}_{s_1,s_2}^* \subset B_{3,S_\ell^1(T),N} \subset A_{a_\ell,N}. \quad (3.9)$$

Furthermore, by remark (b), we have that for any $i \neq a_\ell$, $\bar{\gamma}$ takes the pattern in the annulus $A_{i,N}$ in the environment T if and only if $\bar{\gamma}^*$ takes the pattern in the annulus $A_{i,N}$ in the environment T^* . The same property holds for the selected geodesic γ and for the associated geodesic γ^* (in the environment T^*) given by item (iii) of Lemma 3.5.

- (d) Let again $\bar{\gamma}$ and $\bar{\gamma}^*$ be as in item (ii) of Lemma 3.5. Let us compare the M -sequence of $\bar{\gamma}$ (which is built in the environment T) with the M -sequence of $\bar{\gamma}^*$ (which is built in the environment T^*). By (a) and (3.9), we get that any box which belongs to the M -sequence of $\bar{\gamma}$, with the possible exception of a box contained in $A_{a_\ell,N}$, also belongs to the M -sequence of $\bar{\gamma}^*$. The same property holds for the selected geodesic γ and for the associated geodesic γ^* (in the environment T^*) given by item (iii) of Lemma 3.5. In particular the first $\ell - 1$ elements of the M -sequence of γ (which is built in the environment T) are the same as the first $\ell - 1$ elements of the M -sequence of γ^* (which is built in the environment T^*).

Let us now proceed to the proof of item 1. We assume $\ell \geq 2$ otherwise there is nothing to prove. Let $\bar{\gamma}^*$ be a geodesic from 0 to x in the environment T^* . Let $\bar{\gamma}$ be the associated geodesic in the environment T given by item (ii) of Lemma 3.5. Since the event $\{T \in \mathcal{M}(\ell)\}$ occurs, the M -sequence (in the environment T) of $\bar{\gamma}$ contains at least ℓ typical boxes. By remark (d) above, the M -sequence (in the environment T^*) of $\bar{\gamma}^*$ contains at least $\ell - 1$ typical boxes.

Let us consider item 2. We can again assume $\ell \geq 2$. Let γ^* be the geodesic given by item (iii) of Lemma 3.5. Recall that γ is the selected geodesic and that $\gamma \in \Gamma^\ell$. In particular, we have the following properties: its M -sequence contains at least ℓ boxes; the ℓ -th box of its M -sequence belongs to $A_{a_\ell,N}$; γ does not take the pattern in any annulus whose index is smaller than or equal to a_ℓ . Therefore, by remark (d) above, the first $\ell - 1$ boxes of the M -sequence of γ and γ^* are the same. By remark (c) above, we conclude that γ^* does not take the pattern (in the environment T^*) in any annulus whose index is smaller than the one containing its $(\ell - 1)$ -th box.

Let us prove item 3. Let $\bar{\gamma}^*$ be such a geodesic. Assume that the ℓ -th box of the M -sequence of $\bar{\gamma}^*$ is in an annulus whose index is strictly smaller than a_ℓ . Let $\bar{\gamma}$ be a geodesic in the environment T given by item (ii) of Lemma 3.5. Assume, aiming at a contradiction, that $\bar{\gamma}^*$ does not take the pattern in an annulus until the one containing its ℓ -th box. By remark (d), the ℓ first boxes of the M -sequences of $\bar{\gamma}^*$ and $\bar{\gamma}$ are the same. By remarks (b) and (c), $\bar{\gamma}$ does not take the pattern until the annulus containing its ℓ -th box. This contradicts the definition of S_ℓ^1 , so it is impossible. Thus the ℓ -th box of the M -sequence of $\bar{\gamma}^*$ is in an annulus whose index is greater than or equal to a_ℓ . By item (i) of Lemma 3.5, $\bar{\gamma}^*$ takes the pattern in the annulus whose index is a_ℓ . Therefore it takes the pattern in an annulus whose index is smaller than or equal to the one containing its ℓ -th box and the third point is satisfied.

Finally, let us prove item 4. Note that since $B_{3,s,N}$ is a successful box, $S_\ell^2(T^*) \neq \emptyset$. There are two steps. First, we prove that $NS_\ell^2(T^*)$ is the center of a box contained in the annulus $A_{a_\ell,N}$ and then we prove that $B_{3,s,N} \cap B_{3,S^2(T^*),N} \neq \emptyset$, which gives the result. Assume that $NS_\ell^2(T^*)$ is not the center of a box contained in $A_{a_\ell,N}$. Let again γ^* be the geodesic given by item (iii) of Lemma 3.5 and recall that γ is the selected geodesic in the environment T . Since $B_{3,s,N}$ is successful, we have $a'_\ell(\gamma) \leq a_\ell$ and thus the definition of $S_\ell^2(T^*)$ implies that $NS_\ell^2(T^*)$ is the center of a box contained in an annulus $A_{k,N}$ such that $k < a_\ell$. So γ^* takes the pattern in this annulus in the environment T^* . By remark (c), γ takes the pattern in this annulus in the environment T^* and then in the environment T by remark (b). This is impossible since γ does not take the pattern in any annuli $A_{k,N}$ with $k \leq a_\ell$.

To conclude, assume, aiming at a contradiction, that $B_{3,s,N} \cap B_{3,S^2(T^*),N} = \emptyset$. It implies that γ^* takes a pattern outside $B_{3,s,N}$ in the environment T^* and the portion of γ^* taking this pattern is a portion of γ_{0,s'_1}^* or $\gamma_{s'_2,x}^*$. By remark (b), the portion of γ^* taking this pattern also takes this pattern in the environment T . By remark (c), it implies that γ takes the pattern in the environment T in the annulus $A_{a_\ell,N}$, which is a contradiction. \square

Now, thanks to Lemma 3.3, we can adapt Lemma 3.8 from [1].

Lemma 3.6. *There exists $\lambda \in (0, 1)$, which does not depend on n and on $x \in \Gamma_n$, such that*

$$\mathbb{P}(T \in \mathcal{M}(q)) \leq \lambda^q.$$

Proof. Let ℓ be in $\{1, \dots, q\}$. For every $s \in \mathbb{Z}^d$, let us consider the environment T_s^* defined for all edge e by:

$$T_s^*(e) = \begin{cases} T(e) & \text{if } e \notin B_{2,s,N} \\ T'(e) & \text{else.} \end{cases}$$

Thus T_s^* and T have the same distribution and on the event $\{T \in \mathcal{M}(\ell)\} \cap \{S_\ell^1(T) = s\}$, $T^* = T_s^*$. So, using this environment and writing with indicator functions the result of Lemma 3.3, we get:

$$\mathbb{1}_{\{T \in \mathcal{M}(\ell)\}} \mathbb{1}_{\{S_\ell^1(T) = s\}} \mathbb{1}_{\{T' \in \mathcal{B}^*(T)\}} \leq \mathbb{1}_{\{T_s^* \in \mathcal{M}(\ell-1) \setminus \mathcal{M}(\ell)\}} \mathbb{1}_{\bigcup_{s' \approx s} \{S_\ell^2(T_s^*) = s'\}},$$

where $s' \approx s$ if $\|s - s'\|_1 \leq 2r_3$. We compute the expectation on both sides. The right side yields

$$\mathbb{P} \left(T_s^* \in \mathcal{M}(\ell-1) \setminus \mathcal{M}(\ell), \bigcup_{s' \approx s} \{S_\ell^2(T_s^*) = s'\} \right) = \mathbb{P} \left(T \in \mathcal{M}(\ell-1) \setminus \mathcal{M}(\ell), \bigcup_{s' \approx s} \{S_\ell^2(T) = s'\} \right).$$

For the left side, we have

$$\mathbb{E} \left[\mathbb{1}_{\{T \in \mathcal{M}(\ell)\}} \mathbb{1}_{\{S_\ell^1(T) = s\}} \mathbb{1}_{\{T' \in \mathcal{B}^*(T)\}} \right] = \mathbb{E} \left[\mathbb{1}_{\{T \in \mathcal{M}(\ell)\}} \mathbb{1}_{\{S_\ell^1(T) = s\}} \mathbb{E} \left[\mathbb{1}_{\{T' \in \mathcal{B}^*(T)\}} \mid T \right] \right].$$

Since on the event $\{T \in \mathcal{M}(\ell)\} \cap \{S_\ell^1(T) = s\}$, we have $\mathbb{P}(T' \in \mathcal{B}^*(T) \mid T) \geq \eta$, the left side is bounded from below by $\eta \mathbb{P}(T \in \mathcal{M}(\ell), S_\ell^1(T) = s)$. Then, by summing on all $s \in \mathbb{Z}^d$ and writing K a constant which bounds from above for all $s' \in \mathbb{Z}^d$ the number of vertices $s \in \mathbb{Z}^d$ such that $s' \approx s$, we get

$$\frac{\eta}{K} \mathbb{P}(T \in \mathcal{M}(\ell)) \leq \mathbb{P}(T \in \mathcal{M}(\ell-1) \setminus \mathcal{M}(\ell)).$$

Now, since $\mathcal{M}(\ell) \subset \mathcal{M}(\ell-1)$,

$$\mathbb{P}(T \in \mathcal{M}(\ell-1) \setminus \mathcal{M}(\ell)) = \mathbb{P}(T \in \mathcal{M}(\ell-1)) - \mathbb{P}(T \in \mathcal{M}(\ell)).$$

Thus,

$$\mathbb{P}(T \in \mathcal{M}(\ell)) \leq \lambda \mathbb{P}(T \in \mathcal{M}(\ell-1)),$$

where $\lambda = \frac{1}{1 + \frac{\eta}{K}} \in (0, 1)$ does not depend on x . Hence, using $\mathbb{P}(T \in \mathcal{M}(0)) = 1$, we get by induction

$$\mathbb{P}(T \in \mathcal{M}(q)) \leq \lambda^q.$$

□

Proof of Proposition 1.10. Recall that N , x (and then n and p) are fixed at (3.7) but that C_1 , D_1 and λ does not depend on x , n and p . Then, by Lemma 3.2 and Lemma 3.6, using the inequality (3.8),

$$\begin{aligned} \mathbb{P}(\text{there exists a geodesic } \gamma \text{ from } 0 \text{ to } x \text{ such that } N^{\mathfrak{P}}(\gamma) = 0) &\leq \mathbb{P}(T \in \mathcal{G}^p(N)^c) + \mathbb{P}(T \in \mathcal{M}(q)) \\ &\leq D_1 e^{-C_1 p^{\frac{1}{d}}} + \lambda \lfloor \frac{q}{2} \rfloor. \end{aligned}$$

As $C_1 > 0$ and $\lambda \in (0, 1)$, and as this inequality holds for any $n \geq 2rN$ and any $x \in \Gamma_n$, we get the existence of two constants $C > 0$ and $D > 0$ such that for all n , for all $x \in \Gamma_n$,

$$\mathbb{P}(\text{there exists a geodesic } \gamma \text{ from } 0 \text{ to } x \text{ such that } N^{\mathfrak{P}}(\gamma) = 0) \leq D \exp(-Cn^{\frac{1}{d}}).$$

□

3.2 Typical boxes crossed by geodesics

Let us first begin with the proof of the lemma stated in the paragraph of typical boxes in Section 3.1.

Proof of Lemma 3.1.

1. Let $B_{3,s,N}$ be a typical box. Then the event $\mathcal{T}(s, N)$ occurs. Let u_0 and v_0 be two vertices in $B_{2,s,N}$. We have

$$t_{3,s,N}(u_0, v_0) \leq 2 \sup_{z \in B_{2,s,N}} t_{3,s,N}(Ns, z) \leq 2r_{2,3}N.$$

Let π_0 be a path from u_0 to v_0 which is not entirely contained in $B_{3,s,N}$. Let z_0 denote the first vertex on the boundary of $B_{3,s,N}$ visited by π_0 . Then

$$\begin{aligned} T(\pi_0) &\geq t_{3,s,N}(u_0, z_0) \geq t_{3,s,N}(z_0, Ns) - t_{3,s,N}(u_0, Ns) \\ &\geq \inf_{z \in \partial B_{3,s,N}} t_{3,s,N}(Ns, z) - \sup_{z \in B_{2,s,N}} t_{3,s,N}(Ns, z) \geq 3r_{2,3}N > 2r_{2,3}N \geq t_{3,s,N}(u_0, v_0). \end{aligned}$$

Hence, every geodesic from u_0 to v_0 has to be entirely contained in $B_{3,s,N}$.

2. The properties (ii) and (iii) only depend on the time of edges in $B_{2,s,N}$. The event $\mathcal{T}(s; N)$ only depends on edges in $B_{3,s,N}$ by the definition of $t_{3,s,N}$.
3. First,

$$\lim_{N \rightarrow \infty} \mathbb{P}(\mathcal{T}(0; N)) = 1.$$

Indeed, by (1.16),

$$\mathbb{P}\left(B_\mu\left(0, \frac{r_{2,3}}{2}N\right) \subset \tilde{B}(0, r_{2,3}N) \text{ for all large } N\right) = 1,$$

and

$$\mathbb{P}(\tilde{B}(0, 4r_{2,3}N) \subset B_\mu(0, 8r_{2,3}N) \text{ for all large } N) = 1.$$

Thus, since $B_{2,0,1} \subset B_\mu\left(0, \frac{r_{2,3}}{2}\right) \cap \mathbb{Z}^d$ and $B_\mu(0, 9r_{2,3}) \cap \mathbb{Z}^d \subset B_{3,0,1}$, almost surely there exists $N_0 \in \mathbb{N}^*$ such that for all $N \geq N_0$,

$$B_{2,0,N} \subset B(0, r_{2,3}N), B(0, 4r_{2,3}N) \subset B_{3,0,N} \text{ and for all } y \in \partial B_{3,0,N}, y \notin B(0, 4r_{2,3}N).$$

So, for all $N \geq N_0$,

$$\sup_{z \in \partial B_{2,0,N}} t_{3,0,N}(0, z) \leq r_{2,3}N \quad \text{and} \quad \inf_{z \in \partial B_{3,0,N}} t_{3,0,N}(0, z) \geq 4r_{2,3}N.$$

Note that, for the first inequality, we use the fact that for all $z \in B_{2,0,N}$, $t_{3,0,N}(0, z) = t(0, z)$ thanks to the first point of Lemma 3.1 proved above.

The probability that (iii) is satisfied by $B_{3,0,N}$ goes to 1 by (3.3). Then, let us prove that the probability that (ii) is satisfied by $B_{3,0,N}$ goes to 1. Let $|B_{3,0,N}|$ denote the number of vertices in $B_{3,0,N}$ and Π_0 denote the set of self-avoiding paths entirely contained in $B_{3,0,N}$. Then, using (3.2), we have that $r_2 > r_1$, and by (1.18),

$$\begin{aligned} &\mathbb{P}(B_{3,0,N} \text{ does not satisfy (ii)}) \\ &\leq \sum_{\substack{u_\pi, v_\pi \in B_{3,0,N} \\ \|u_\pi - v_\pi\|_1 \geq (r_2 - r_1)N}} \mathbb{P}((3.4) \text{ is not satisfied by a path of } \Pi_0 \text{ whose endpoints are } u_\pi \text{ and } v_\pi) \\ &\leq \sum_{\substack{u_\pi, v_\pi \in B_{3,0,N} \\ \|u_\pi - v_\pi\|_1 \geq (r_2 - r_1)N}} \mathbb{P}((3.4) \text{ is not satisfied by a path whose endpoints are } u_\pi \text{ and } v_\pi) \\ &\leq |B_{3,0,N}|^2 e^{-D_0(r_2 - r_1)N} \xrightarrow{N \rightarrow \infty} 0, \end{aligned}$$

since $|B_{3,0,N}|$ is bounded by a polynomial in N .

□

Proof of Lemma 3.2. To begin this proof, one need an upper bound on the Euclidean length of geodesics. Using Theorem 4.6 in [2], we have two positive constants K_1 and C_2 such that for all $y \in \mathbb{Z}^d$,

$$\mathbb{P}(m(y) \geq K_1 \|y\|_1) \leq e^{-C_2 \|y\|_1^{\frac{1}{d}}},$$

where $m(y) = \max\{|\sigma|_e : \sigma \text{ is a geodesic from } 0 \text{ to } y\}$ and where for a path σ , $|\sigma|_e$ means the number of different edges taken by σ . For all $p \in \mathbb{N}^*$, we define the event $\mathcal{N}^p(N)$ on which every geodesic from 0 to the outer sphere of the p -th annulus takes less than $K_1 prN$ distinct edges. Note that $r = 2(r_1 + r_3 + 1)$ is fixed at (3.5) and rN corresponds to the widths of the annuli. Then,

$$\mathbb{P}(\mathcal{N}^p(N)^c) \leq \sum_{y: \|y\|_1 = prN} \mathbb{P}(m(y) \geq K_1 \|y\|_1) \leq (2prN + 1)^d e^{-C_2 (prN)^{\frac{1}{d}}}.$$

Hence, we obtain two positive constants C_3 and D_3 only depending on r , d and F such that for all $p \in \mathbb{N}^*$, for all $N \in \mathbb{N}^*$,

$$\mathbb{P}(\mathcal{N}^p(N)^c) \leq D_3 e^{-C_3 p^{\frac{1}{d}}}.$$

Now, we assume that the event $\mathcal{N}^p(N) \cap \mathcal{G}^p(N)^c$ occurs. So, every geodesic from 0 to the outer sphere of the p -th annulus takes a number of distinct edges which is between prN and $K_1 prN$. Let us consider a re-normalized model. We introduce the meta-cubes

$$B_{s,N}^\infty = \left\{ w \in \mathbb{Z}^d : \left(s - \frac{1}{2} \right) N \leq w < \left(s + \frac{1}{2} \right) N \right\}, \text{ for all } s \in \mathbb{Z}^d,$$

(where $v \leq w$ means $v_i \leq w_i$ for $1 \leq i \leq d$ and $v < w$ means $v_i < w_i$ for $1 \leq i \leq d$.) These meta-cubes form a partition of \mathbb{Z}^d . Furthermore, the meta-cubes and the boxes defined above have the same centers (which are the vertices Ns for $s \in \mathbb{Z}^d$), and for all $s \in \mathbb{Z}^d$, $B_{s,N}^\infty \subset B_{1,s,N}$. So, we can define typical meta-cubes. A meta-cube $B_{s,N}^\infty$ is typical if $B_{3,s,N}$ is a typical box.

For a geodesic γ from 0 to the outer sphere of the p -th annulus, we associate the set of meta-cubes visited by γ , that is

$$\mathfrak{A}(\gamma) = \{B_{s,N}^\infty \mid \gamma \text{ visits at least one vertex of } B_{s,N}^\infty\}.$$

This set can be identify with the subset of the re-normalized graph $N\mathbb{Z}^d$:

$$\mathfrak{A}_v^R(\gamma) = \{sN \mid B_{s,N}^\infty \in \mathfrak{A}(\gamma)\}.$$

Note that, if we consider the set $\mathfrak{A}_e^R(\gamma)$ of edges of $N\mathbb{Z}^d$ linking vertices which are both in $\mathfrak{A}_v^R(\gamma)$, then the pair of sets $(\mathfrak{A}_v^R(\gamma), \mathfrak{A}_e^R(\gamma))$ forms a lattice animal, denoted by $\mathfrak{A}^R(\gamma)$. Recall that a lattice animal \mathfrak{A} in $N\mathbb{Z}^d$ is a finite connected sub-graph of $N\mathbb{Z}^d$ that contains 0. We denote by \mathcal{A}^R the set of lattice animals in $N\mathbb{Z}^d$ associated with a geodesic going from 0 to the outer sphere of the p -th annulus.

Let us bound the size of these lattice animals. By the size of a lattice animal \mathfrak{A}^R , denoted by $|\mathfrak{A}^R|_v$, we mean its number of vertices in the re-normalized model. Recall that, since the event $\mathcal{N}^p(N)$ occurs, every geodesic from 0 to the outer sphere of the p -th annulus takes a number of distinct edges which is between prN and $K_1 prN$. Then, in the meta-cube set $\mathfrak{A}(\gamma)$ associated to such a geodesic γ , since $r_1 = d$ and thanks to the choice of r , there are $p - 1$ meta-cubes associated to boxes crossed by γ in distinct annuli. In particular (considering also the meta-cube centered at the origin), the size of every lattice animal $\mathfrak{A}^R \in \mathcal{A}^R$ is bounded from below by p . For an upper bound, we use the inequality

$$|\gamma|_e \geq N \left(\frac{|\mathfrak{A}^R(\gamma)|_v}{3^d} - 1 \right), \quad (3.10)$$

for all geodesic γ from 0 to the outer sphere of the p -th annulus, where $|\gamma|_e$ still denotes the number of edges taken by γ and where $|\mathfrak{A}^R(\gamma)|_v$ is the number of vertices of $\mathfrak{A}^R(\gamma)$. Let us prove this inequality. Let γ be a geodesic from 0 to the outer sphere of the p -th annulus and denote by $\gamma = (v_0, \dots, v_m)$ the sequence of vertices visited by γ . For all $v \in \mathbb{Z}^d$, denote by $s(v)$ the unique $s \in \mathbb{Z}^d$ such that v belongs to $B_{s,N}^\infty$. We define by induction a strictly increasing sequence i_0, \dots, i_κ by setting $\kappa = 0$ and $i_0 = 0$ and then applying the following algorithm:

- (a) If there exists $i \in \{i_\kappa + 1, \dots, m\}$ such that $s(v_i)$ is at distance at least 2 for the norm $\|\cdot\|_\infty$ from $s(v_{i_\kappa})$, we denote by $i_{\kappa+1}$ the smallest of these i , then we increment κ and go back to (a).
- (b) Otherwise we stop the algorithm.

Then, we necessarily have $3^d(\kappa + 1) \geq |\mathfrak{A}^R(\gamma)|_v$. Furthermore, for all $k \in \{0, \dots, \kappa - 1\}$, we have $\|v_{i_{\kappa+1}} - v_{i_k}\|_1 \geq N$. Hence,

$$|\gamma|_e \geq N\kappa \geq N \left(\frac{|\mathfrak{A}^R(\gamma)|_v}{3^d} - 1 \right),$$

and (3.10) is proved.

Now, using (3.10), writing $K_2 = \lceil 3^d(K_1 r + 1) \rceil$ (which does not depend on p and N), for every lattice animal $\mathfrak{A}^R \in \mathcal{A}^R$, $|\mathfrak{A}^R|_v$ is bounded from above by $K_2 p$. Furthermore, for $j \in \{p, \dots, K_2 p\}$, using (4.24) in [4], we have that

$$|\{\mathfrak{A}^R \in \mathcal{A}^R : |\mathfrak{A}^R|_v = j\}| \leq |\{\text{lattice animals in } \mathbb{Z}^d \text{ of size } j\}| \leq 7^{dj}. \quad (3.11)$$

Let us consider the random variables $(X_\ell^N)_{\ell \in \mathbb{Z}^d}$ such that $X_\ell^N = 1$ if the meta-cube $B_{\ell, N}^\infty$ is typical and $X_\ell^N = 0$ otherwise. By Lemma 3.1, there exists a positive constant K_3 such that X_ℓ^N is independent from the sigma-algebra generated by $\{X_k^N, k \in \mathbb{Z}^d : \|k - \ell\|_1 \geq K_3\}$. Furthermore, also by Lemma 3.1,

$$\lim_{N \rightarrow \infty} \mathbb{P}(X_\ell^N = 1) = 1.$$

Thus, by Corollary 1.4 in [6], there exists $\eta_1 = \eta_1(N) > 0$ such that

$$\eta_1(N) \xrightarrow{N \rightarrow \infty} 0,$$

and there exist i.i.d. random variables $(Y_\ell^N)_{\ell \in \mathbb{Z}^d}$ such that $(X_\ell^N)_\ell \geq (Y_\ell^N)_\ell$ and Y_0^N has a Bernoulli distribution of parameter $(1 - \eta_1(N))$. Finally, we have

$$\mathbb{P}(\mathcal{N}^p(N) \cap \mathcal{G}^p(N)^c) \leq \mathbb{P} \left(\exists \mathfrak{A}^R \in \mathcal{A}^R \text{ such that } p \leq |\mathfrak{A}^R|_v \leq K_2 p \text{ and } \sum_{\ell \in \mathfrak{A}_v^R} X_\ell^N \leq |\mathfrak{A}^R|_v + 1 - \frac{p}{2} \right).$$

Indeed, on $\mathcal{G}^p(N)^c$, there exists a geodesic γ from 0 to the outer sphere of the p -th annulus which crosses a typical box in strictly less than $\lfloor \frac{p}{2} \rfloor$ annuli, and thus there are strictly more than $\lfloor \frac{p}{2} \rfloor - 1$ annuli $A_{i, N}$ with $i > 1$ such that γ does not cross a typical box in them. Furthermore, there are $p - 1$ meta-cubes in $\mathfrak{A}(\gamma)$ such that each of them is associated to a box crossed by γ in one of the $p - 1$ distinct annuli $A_{i, N}$ with $1 < i \leq p$. Thus, there are strictly more than $\lfloor \frac{p}{2} \rfloor - 1$ of these specified meta-cubes which are not typical. Hence the number of typical meta-cubes in $\mathfrak{A}(\gamma)$ is strictly smaller than $|\mathfrak{A}^R|_v - \lfloor \frac{p}{2} \rfloor + 1$. Then, using the random variables $(Y_\ell^N)_{\ell \in \mathbb{Z}^d}$,

$$\begin{aligned} \mathbb{P}(\mathcal{N}^p(N) \cap \mathcal{G}^p(N)^c) &\leq \mathbb{P} \left(\exists \mathfrak{A}^R \in \mathcal{A}^R \text{ such that } p \leq |\mathfrak{A}^R|_v \leq K_2 p \text{ and } \sum_{\ell \in \mathfrak{A}_v^R} Y_\ell^N \leq |\mathfrak{A}^R|_v - \lfloor \frac{p}{2} \rfloor + 1 \right) \\ &\leq \sum_{p \leq j \leq K_2 p} |\{\text{lattice animals in } \mathbb{Z}^d \text{ of size } j\}| \mathbb{P} \left(\text{binomial}(j, \eta_1) \geq \lfloor \frac{p}{2} \rfloor - 1 \right) \\ &\leq \sum_{p \leq j \leq K_2 p} 7^{dj} \mathbb{P} \left(\text{binomial}(j, \eta_1) \geq \lfloor \frac{p}{2} \rfloor - 1 \right) \quad (\text{by (3.11)}) \\ &\leq K_2 p 7^{dK_2} \mathbb{P} \left(\text{binomial}(K_2 p, \eta_1) \geq \lfloor \frac{p}{2} \rfloor - 1 \right). \end{aligned}$$

Then, for $p \geq 4$ and N large enough to have $\eta_1(N) < \frac{1}{4K_2}$, using a Chernov bound for the binomial distribution (see Section 2.2 in [3]), we get

$$\mathbb{P} \left(\text{binomial}(K_2 p, \eta_1) \geq \lfloor \frac{p}{2} \rfloor - 1 \right) \leq \mathbb{P} \left(\text{binomial}(K_2 p, \eta_1) \geq \frac{p}{4} \right) \leq \exp \left(-K_2 p h_{\eta_1} \left(\frac{1}{4K_2} \right) \right),$$

where for $x \in (\eta_1, 1)$,

$$h_{\eta_1}(x) = (1-x) \ln \left(\frac{1-x}{1-\eta_1} \right) + x \ln \left(\frac{x}{\eta_1} \right).$$

Thus, since we can take η_1 as small as we want by taking N large enough,

$$\begin{aligned} \mathbb{P}(\mathcal{N}^p(N) \cap \mathcal{G}^p(N)^c) &\leq K_2 p \left[7^{dK_2} \exp \left(-K_2 h_{\eta_1} \left(\frac{1}{4K_2} \right) \right) \right]^p \\ &\leq K_2 p \exp(-2p) \text{ for } N \text{ large enough} \\ &\leq \exp(-C_4 p). \end{aligned}$$

Finally, we have a constant N_0 such that for all $p \geq 4$, for all $N \geq N_0$,

$$\mathbb{P}(\mathcal{G}^p(N)^c) \leq \mathbb{P}(\mathcal{N}^p(N) \cap \mathcal{G}^p(N)^c) + \mathbb{P}(\mathcal{N}^p(N)^c) \leq e^{-C_4 p} + D_3 e^{-C_3 p^{\frac{1}{d}}}.$$

So, there exist two positive constants C_1 and D_1 such that for all $p \geq 1$, for all $N \geq N_0$,

$$\mathbb{P}(\mathcal{G}^p(N)^c) \leq D_1 e^{-C_1 p^{\frac{1}{d}}}.$$

□

3.3 Modification argument

The aim of this subsection is to prove Lemma 3.5. Let $\ell \in \{1, \dots, q\}$. On $\{T \notin \mathcal{M}(\ell)\}$, we set $E_+^*(T) = \emptyset$ and $E_-^*(T) = \emptyset$. Let s be in \mathbb{Z}^d . We now define E_+^* and E_-^* on the event $\{T \in \mathcal{M}(\ell)\} \cap \{S_\ell^1(T) = s\}$. So assume that this event occurs. On the event $\{T \in \mathcal{M}(\ell)\}$, Γ^ℓ is not empty and thus there is a selected geodesic. We denote this selected geodesic by γ . We define the entry point (resp. the exit point) of a self-avoiding path in a set of vertices as the first (resp. the last) vertex of this path belonging to this set. Let u denote the entry point of γ in $B_{2,s,N}$ and v the exit point.

We call entry point and exit point of the pattern (centered at 0) the endpoints denoted by u^Λ and v^Λ in the introduction. Note that, if a self-avoiding path takes the pattern, its entry and exit points in the set $B_\infty(0, \ell^\Lambda)$ are not necessarily the entry and exit points of the pattern (as it can visit the set before and after taking the pattern).

Here, we want to put the pattern centered at sN . The vertex s being fixed, we keep the notation u^Λ and v^Λ to designate the entry and the exit points of the pattern centered at sN .

Construction of π .

We have the following inclusions:

- $B_\infty(sN, \ell^\Lambda) \subset B_\infty(sN, \ell^\Lambda + 3) \subset B_{1,s,N}$ since $r_1 = d$ and $N \geq \ell^\Lambda + 3$ (see (3.7)),
- $B_{1,s,N} \subset B_{2,s,N}$ since $r_2 > r_1$ by (3.2).

For the modification, we need a path π , constructed in a deterministic way and satisfying several properties, whose existence is guaranteed by the following lemma.

Lemma 3.7. *We can construct a path π in a deterministic way such that :*

- (i) π goes from u to u^Λ without visiting a vertex of $B_\infty(sN, \ell^\Lambda)$, then goes from u^Λ to v^Λ in a shortest way for the norm $\|\cdot\|_1$ (and thus being contained in $B_\infty(sN, \ell^\Lambda)$) and then goes from v^Λ to v without visiting a vertex of $B_\infty(sN, \ell^\Lambda)$,
- (ii) π is entirely contained in $B_{2,s,N}$ and does not have vertices on the boundary of $B_{2,s,N}$ except u and v ,
- (iii) π is self-avoiding,
- (iv) the length of $\pi_{u,u^\Lambda} \cup \pi_{v^\Lambda,v}$ is bounded from above by $2r_2 N + K$, where K is the number of edges in $B_\infty(0, \ell^\Lambda + 3)$.




Figure 2: Example of the construction of π in dimension 2.

The proof of this lemma is given in Appendix A but the idea is to construct two paths, one from u to sN and the other from sN to v which minimize the distance for the norm $\|\cdot\|_1$ and such that the only vertex belonging to both paths is sN . Then, we denote by u_0 the first vertex of $B_{\infty}(sN, \ell^{\Lambda} + 3)$ visited by the path from u to sN and v_0 the last vertex of $B_{\infty}(sN, \ell^{\Lambda} + 3)$ visited by the path from sN to v . We construct two paths entirely contained in $B_{\infty}(sN, \ell^{\Lambda} + 3)$ from u_0 to u^{Λ} and from v^{Λ} to v_0 which do not take vertices of $B_{\infty}(sN, \ell^{\Lambda})$ except u^{Λ} and v^{Λ} and which have no vertices in common and we consider the concatenation of the path from u to u_0 , the one from u_0 to u^{Λ} , a path from u^{Λ} to v^{Λ} in a shortest way, the path from v^{Λ} to v_0 and the one from v_0 to v (see Figure 2).

Let π be the path given by Lemma 3.7.

Definition of E_+^* , E_-^* and \mathcal{B}^* . Define $E_-^*(T)$ as the set of edges e such that $e \in \pi \setminus B_{\infty}(sN, \ell^{\Lambda})$ and $E_+^*(T)$ as the set of edges which are in $B_{2,s,N}$ but which are not in $B_{\infty}(sN, \ell^{\Lambda}) \cup \pi$. Recall that $\{T' \in \mathcal{B}^*(T)\}$ is a shorthand for

$$\{\forall e \in E_+^*(T), T'(e) \geq \nu(N), \forall e \in E_-^*(T), T'(e) \leq r + \delta', \theta_{NS_{\ell}^1(T)} T' \in \mathcal{A}^{\Lambda}\}.$$

Fix $\eta = \tilde{p}^{|B_{2,s,N}|} \mathbb{P}(T \in \mathcal{A}^{\Lambda})$, where $\tilde{p} = \min(F([t_{\min}, t_{\min} + \delta']), F([\nu(N), t_{\max}]))$. Thus, η only depends on F , the pattern and N and we have

$$\mathbb{P}(T' \in \mathcal{B}^*(T)|T) \geq \tilde{p}^{|B_{2,s,N}|} \mathbb{P}(T \in \mathcal{A}^{\Lambda}) = \eta,$$

Consequences of the modification. We denote by γ^* the path $\gamma_{0,u} \cup \pi \cup \gamma_{v,x}$. Note that γ^* is a self-avoiding path.

Lemma 3.8. *We have $T^*(\gamma^*) < T(\gamma)$.*

Proof. We have that $\gamma_{u,v}$ visits at least one vertex in $B_{1,s,N}$. Denote by w the first of these vertices. Then, $\gamma_{u,w}$ and $\gamma_{w,v}$ are two geodesics, both between two vertices in $B_{2,s,N}$. Using item 1 in Lemma 3.1, $\gamma_{u,w}$ and $\gamma_{w,v}$ are entirely contained in $B_{3,s,N}$. Thus, since $B_{3,s,N}$ is a typical box, using (3.4) and the

fact that $\|u - w\|_1 \geq (r_2 - r_1)N$ and $\|v - w\|_1 \geq (r_2 - r_1)N$, we have

$$T(\gamma_{u,v}) \geq 2N(r_2 - r_1)(t_{\min} + \delta).$$

Then, by the construction of π and of $\mathcal{B}^*(T)$,

$$T^*(\pi) \leq (2r_2N + K)(t_{\min} + \delta') + \tau^\Lambda,$$

where τ^Λ is fixed at (3.1). Thus,

$$T(\gamma) - T^*(\gamma^*) \geq 2N(r_2(\delta - \delta') - r_1(t_{\min} + \delta)) - K(t_{\min} + \delta') - \tau^\Lambda.$$

By (3.6) and since $2N \geq 1$, we get $T(\gamma) - T^*(\gamma^*) > 0$. \square

Lemma 3.9. *Let $\bar{\gamma}^*$ be a geodesic from 0 to x in the environment T^* . Then $\bar{\gamma}^*$ weakly crosses the box $B_{3,s,N}$ and the first vertex of $B_{2,s,N}$ visited by $\bar{\gamma}^*$ is u and the last is v . Furthermore, $\bar{\gamma}^*$ takes the pattern in $B_\infty(sN, \ell^\Lambda)$, $\bar{\gamma}_{u,u^\Lambda}^* = \pi_{u,u^\Lambda}$ and $\bar{\gamma}_{v^\Lambda,v}^* = \pi_{v^\Lambda,v}$.*

Proof. Let $\bar{\gamma}^*$ be a geodesic from 0 to x in the environment T^* . By Lemma 3.8, $T^*(\bar{\gamma}^*) < T(\bar{\gamma}^*)$. Thus $\bar{\gamma}^*$ takes an edge of $B_{2,s,N}$ and by item (iii) of the definition of a typical box and since there is no edge whose time has been modified outside $B_{2,s,N}$, $\bar{\gamma}^*$ can not take any edge of time greater than $\nu(N)$ in $B_{2,s,N}$. Indeed, assume that $\bar{\gamma}^*$ takes an edge e such that $T^*(e) \geq \nu(N)$. Then, denoting by $\mathcal{E}(\bar{\gamma}^*)$ the edges of $\bar{\gamma}^*$ and using (3.3),

$$\begin{aligned} T^*(\bar{\gamma}^*) &= \sum_{f \in \mathcal{E}(\bar{\gamma}^*) \cap B_{2,s,N}} T^*(f) + \sum_{f \in \mathcal{E}(\bar{\gamma}^*) \cap B_{2,s,N}^c} T^*(f) \geq \nu(N) + \sum_{f \in \mathcal{E}(\bar{\gamma}^*) \cap B_{2,s,N}^c} T(f) \\ &> \sum_{f \in B_{2,s,N}} T(f) + \sum_{f \in \mathcal{E}(\bar{\gamma}^*) \cap B_{2,s,N}} T(f) \geq \sum_{f \in \mathcal{E}(\bar{\gamma}^*) \cap B_{2,s,N}} T(f) + \sum_{f \in \mathcal{E}(\bar{\gamma}^*) \cap B_{2,s,N}^c} T(f) = T(\bar{\gamma}^*), \end{aligned}$$

which is impossible. Hence, $\bar{\gamma}^*$ has to take edges of π or of the pattern and can not take other edges of $B_{2,s,N}$.

Since π does not visit any vertex on the boundary of $B_{2,s,N}$ except u and v , $\bar{\gamma}^*$ has to visit u and v and to follow π between u and u^Λ and between v^Λ and v . If $\bar{\gamma}_{u^\Lambda,v^\Lambda}^*$ leaves the pattern, it takes an edge whose time is greater than $\nu(N)$, which is impossible. So, $\bar{\gamma}_{u^\Lambda,v^\Lambda}^*$ is a path entirely contained in $B_\infty(sN, \ell^\Lambda)$ and is optimal for the passage time since $\bar{\gamma}^*$ is a geodesic.

To conclude, let us show that u is visited by $\bar{\gamma}^*$ before v . Assume that it is not the case. Then, there exists $\bar{\gamma}_1^*$ a geodesic from 0 to v and $\bar{\gamma}_2^*$ a geodesic from u to x in the environment T^* which does not take any edge in $B_{2,s,N}$. Thus, there are also geodesics in the environment T . Then,

$$T(\bar{\gamma}_1^*) + T(\bar{\gamma}_2^*) \leq t^*(0, x) < t(0, x) \text{ by Lemma 3.8.}$$

By concatenating $\gamma_{0,u}$ and $\bar{\gamma}_2^*$, we obtain a path from 0 to x . Thus,

$$T(\gamma_{0,u}) + T(\bar{\gamma}_2^*) \geq t(0, x).$$

So $T(\bar{\gamma}_1^*) < T(\gamma_{0,u})$, which implies

$$T(\bar{\gamma}_1^*) + T(\gamma_{v,x}) < T(\gamma_{0,u}) + T(\gamma_{v,x}) \leq t(0, x),$$

which is impossible since $\bar{\gamma}_1^* \cup \gamma_{v,x}$ is a path from 0 to x . \square

Using Lemma 3.9, we can prove Lemma 3.5. Indeed, by this previous lemma, every geodesic from 0 to x takes the pattern inside $B_{2,s,N}$ and the first item holds. For the third item, one can check that the concatenation of $\gamma_{0,u}$, π_{u,u^Λ} , one of the optimal paths for the passage time between u^Λ and v^Λ entirely contained in $B_\infty(sN, \ell^\Lambda)$, $\pi_{v^\Lambda,v}$ and $\gamma_{v,x}$ gives a geodesic γ^* which is associated to γ in $B_{3,s,N}$ (with $s_1 = u$ and $s_2 = v$). Finally, let us prove the second item. If $\bar{\gamma}^*$ is a geodesic from 0 to x in the environment T^* , then $\bar{\gamma}_{u,v}^*$ is contained in $B_{2,s,N}$. Furthermore,

$$T(\gamma_{0,u}) = T^*(\gamma_{0,u}) \geq T^*(\bar{\gamma}_{0,u}^*) = T(\bar{\gamma}_{0,u}^*) \geq T(\gamma_{0,u}),$$

so $T(\overline{\gamma}_{0,u}^*) = T(\gamma_{0,u})$ and $\overline{\gamma}_{0,u}^*$ is a geodesic in the environment T . Similarly, $\overline{\gamma}_{v,x}^*$ is a geodesic in the environment T . Hence, we get a not necessarily self-avoiding optimal path for the passage time by considering $\pi' = \overline{\gamma}_{0,u}^* \cup \gamma_{u,v} \cup \overline{\gamma}_{v,x}^*$. We get a geodesic γ' which satisfies the properties of this item by cutting the loops of π' using a standard process⁷.

4 Bounded case

In this section, we consider that the support of F is bounded. In this case, Theorem 1.3 also follows from Proposition 1.10. Our proof of Proposition 1.10 still follows the strategy given in the preceding section, but the modification argument is more involved. Let $\mathfrak{P} = (\Lambda, u^\Lambda, v^\Lambda, \mathcal{A}^\Lambda)$ be a valid pattern. We set $t_{\max} = \sup(\text{support}(F))$. Remark that, because of (1.18), we have $t_{\min} + \delta < t_{\max}$.

4.1 Oriented pattern

The proof in the bounded case uses a modification argument in which we have to connect the pattern to a straight path in a given direction. It is convenient to show the feasibility of this construction before starting the modification. The following lemma, whose proof is in Appendix B, shows that it is indeed feasible by proving that a pattern can be associated to d patterns (with a larger size), each having endpoints aligned in a distinct direction, and each having the original pattern as a sub-pattern. By direction, we mean one of the d directions of the canonical basis which is denoted by $\{\varepsilon_1, \dots, \varepsilon_d\}$.

Lemma 4.1. *There exists $\ell_0 > \ell^\Lambda$ such that the following holds. Set $\Lambda_0 = \{-\ell_0, \dots, \ell_0\}^d$. For all $j \in \{1, \dots, d\}$, there exists a pattern $\mathfrak{P}^j = (\Lambda_0, -\ell_0\varepsilon_j, \ell_0\varepsilon_j, \mathcal{A}_j^{\Lambda_0})$ such that:*

- $\mathbb{P}(\mathcal{A}_j^{\Lambda_0})$ is positive,
- on $\mathcal{A}_j^{\Lambda_0}$, any path from $-\ell_0\varepsilon_j$ to $\ell_0\varepsilon_j$ optimal for the passage time among the paths entirely inside Λ_0 contains a subpath from u^Λ to v^Λ entirely inside Λ ,
- $\mathcal{A}_j^{\Lambda_0} \subset \mathcal{A}^\Lambda$.

We fix ℓ_0, Λ_0 and the patterns \mathfrak{P}_j , for $j = 1, \dots, d$, for the remaining of the proof.

Remark 4.2. *One can check that the proof of this lemma can be adapted to prove that if the two endpoints of the original pattern belong to two different faces of Λ , then the conclusion of Lemma 4.1 is also satisfied and thus we can generalize the notion of valid pattern of Definition 1.1.*

By definition, for all $j \in \{1, \dots, d\}$, $N^{\mathfrak{P}}(\pi) \geq N^{\mathfrak{P}_j}(\pi)$, and therefore

$$N^{\mathfrak{P}}(\pi) \geq \sum_{x \in \mathbb{Z}^d} \mathbb{1}_{\{\text{there exists } j \in \{1, \dots, d\}, x \text{ satisfies the condition } (\pi; \mathfrak{P}_j)\}}.$$

From now on, we forget the original pattern and only consider the oriented patterns \mathfrak{P}_j for all directions $j \in \{1, \dots, d\}$. We talk about oriented pattern when its orientation is specified and we simply say "pattern" when talking about one of the oriented patterns \mathfrak{P}_j . Consistently with these conventions and to lighten notations, we write ℓ^Λ instead of ℓ_0^Λ , Λ instead of Λ_0 and \mathcal{A}_j^Λ instead of $\mathcal{A}_j^{\Lambda_0}$.

Now, several parameters related to the distribution F have to be introduced. First, we fix a positive real ν such that :

- $t_{\min} + \delta \leq \nu \leq t_{\max}$,
- $F([\nu, +\infty)) > 0$,
- the event $\mathcal{A}^\Lambda \cap \{\forall e \in \Lambda, T(e) \leq \nu\}$ has a positive probability.

⁷Note that $\overline{\gamma}_{0,u}^*$, $\gamma_{u,v}$ and $\overline{\gamma}_{v,x}^*$ are three self-avoiding paths and that $\overline{\gamma}_{0,u}^*$ and $\overline{\gamma}_{v,x}^*$ do not have vertices in common. So we can consider s_1 the last vertex belonging to both path $\overline{\gamma}_{0,u}^*$ and $\gamma_{u,v}$ in the order in which they are visited by $\gamma_{u,v}$ and s_2 the first vertex belonging to both paths $\gamma_{s_1,v}$ and $\overline{\gamma}_{v,x}^*$ in the order in which they are visited by $\gamma_{u,v}$. Thus, since $\gamma_{u,v}$ is entirely contained in $B_{3,s,N}$, s_1 and s_2 are two vertices contained in $B_{3,s,N}$ and we can take $\gamma' = \overline{\gamma}_{0,s_1}^* \cup \gamma_{s_1,s_2} \cup \overline{\gamma}_{s_2,x}^*$.

Notice that, if F has an atom, one could have $\nu = t_{\max}$. Even if it means replacing \mathcal{A}_j^Λ by $\mathcal{A}_j^\Lambda \cap \{\forall e \in \Lambda, T(e) \leq \nu\}$, we can assume that

$$\mathcal{A}_j^\Lambda \subset \{\forall e \in \Lambda, T(e) \leq \nu\}.$$

Further, we set $\tau^\Lambda = 2\ell^\Lambda\nu$, which can be interpreted as an upper bound for the passage time of an optimal path from $\ell^\Lambda\varepsilon_j$ to $-\ell^\Lambda\varepsilon_j$ on the event \mathcal{A}_j^Λ . Finally, we denote by T^Λ the constant $K^\Lambda(t_{\max} - t_{\min})$ where K^Λ is the number of edges in an oriented pattern. We will use it as an upper bound for the time that a path can save using edges of a pattern after a modification.

4.2 Proof of Proposition 1.10 in the bounded case

We keep the overall plan of the unbounded case. Unlike in the unbounded case, we cannot use edges of prohibitive time and thus the modification argument is more elaborate here. This section follows the structure of Section 3.1 but the one step modification is replaced by a two-steps modification. To this aim, we slightly change the structure of our boxes and our definition of typical boxes. Let us begin by fixing some constants.

Constants. Note that we keep the notations introduced in Section 1.6 and that τ^Λ and T^Λ are fixed in Section 4.1. In the next two paragraphs, we introduce the constants used in the proof. Before looking at their definitions below, one can keep in mind that we fix $\varepsilon \ll 1$ and $r_1 \ll \frac{1}{\varepsilon} \ll \nabla \ll r_2 \ll r_3 \ll r_4$, where " \ll " means that the ratio is large enough and only depends on the dimension d and on the distribution F .

- We fix $\delta' = \min\left(\frac{\delta}{4}, \frac{\delta}{1+d}\right)$.
- We fix L_1 given by Lemma 4.12 only depending on d and ℓ^Λ , and $L_2 = L_1 + (10+d)\ell^\Lambda$.
- Using Theorem 1.5 with $M = t_{\min} + \delta$, we get two constants $\alpha > 0$ and C , fixed for the rest of the proof, such that for all $n \in \mathbb{N}^*$ and $u, v \in \mathbb{Z}^d$ such that $\|u - v\|_1 = n$,

$$\mathbb{P}\left(\exists \text{ a geodesic } \bar{\gamma} \text{ from } u \text{ to } v \text{ such that } \sum_{e \in \bar{\gamma}} \mathbf{1}_{\{T(e) \geq t_{\min} + \delta\}} \leq \alpha n\right) \leq e^{-Cn}. \quad (4.1)$$

- Fix $\varepsilon > 0$ such that

$$\varepsilon < \min\left(\frac{1}{11}, \frac{\delta}{24C_\mu}\right).$$

- Fix ∇ such that

$$\nabla > \max\left(\frac{4(1+t_{\max})C_\mu}{\varepsilon c_\mu}, 6dL_2C_\mu, \frac{8C_\mu T^\Lambda}{3\delta}, 4C_\mu(2t_{\max} + \tau^\Lambda)\right). \quad (4.2)$$

We give here other lower bounds for ∇ that we need for the sequel and which are consequences of (4.2).

- Using the fact that $c_\mu \leq C_\mu$, we get $\nabla > \frac{4C_\mu}{\varepsilon c_\mu} > \frac{1}{\varepsilon}$.
- From the inequality $\nabla > \frac{4(1+t_{\max})C_\mu}{\varepsilon c_\mu}$, using the fact that $\varepsilon < \frac{1}{11}$, we have $1 - \varepsilon > 1 - 3\varepsilon > 1 - 10\varepsilon > \varepsilon$ and then $\nabla > \frac{1+2t_{\max}}{1-\varepsilon}$, $\nabla > \frac{4(1+t_{\max})}{1-10\varepsilon}$ and $\nabla > \frac{3+2t_{\max}}{1-3\varepsilon}$.
- Since $\varepsilon < \frac{\delta}{24C_\mu}$ and $\nabla > \frac{1}{\varepsilon}$, we get $\nabla > \frac{24C_\mu}{\delta}$.
- Finally, since $\delta - \delta' \geq \frac{3\delta}{4}$, we have from $\nabla > \frac{8C_\mu T^\Lambda}{3\delta}$ that $\nabla > \frac{2C_\mu T^\Lambda}{\delta - \delta'}$.

Boxes. With these constants, we can now define boxes. For $i \in \{1, 2, 3, 4\}$, as in the unbounded case, $B_{i,s,N}$ is the ball of radius r_i for the norm $\|\cdot\|_1$ centered at the point sN with:

- $r_1 = d$,
- r_2 an integer such that

$$r_2 > \max\left(r_1 + \frac{2(\nabla + 2)}{c_\mu}, r_1 + L_1 + \frac{3\nabla}{c_\mu} + \frac{2t_{\max}(1 + (1+d)\ell^\Lambda)}{\nu}\right), \quad (4.3)$$

- r_3 an integer such that

$$r_3 > \frac{7r_2(4t_{\max} + \alpha\delta)}{\alpha\delta},$$

Note that $r_3 \geq r_2 + 1$.

- r_4 an integer such that

$$r_4 > \frac{r_3(t_{\min} + \delta + t_{\max})}{t_{\min} + \delta}. \quad (4.4)$$

Note that $r_4 \geq r_3 + 1$.

We use the word "box" to talk about $B_{4,s,N}$. We denote by $\partial B_{i,s,N}$ the boundary of $B_{i,s,N}$, that is the set of points $z \in \mathbb{Z}^d$ such that $\|z - sN\|_1 = r_i N$.

Crossed boxes and weakly crossed boxes. We say that a path

- crosses a box $B_{4,s,N}$ if it visits a vertex in $B_{1,s,N}$,
- weakly crosses a box $B_{4,s,N}$ if it visits a vertex in $B_{3,s,N}$.

Paths associated in a box. We say that two paths γ and γ' from 0 to the same vertex x are associated in a box $B_{4,s,N}$ if there exist two distinct vertices s_1 and s_2 such that the following conditions hold:

- γ and γ' visit s_1 and s_2 ,
- $\gamma_{0,s_1} = \gamma'_{0,s_1}$,
- γ_{s_1,s_2} and γ'_{s_1,s_2} are entirely contained in $B_{4,s,N}$,
- $\gamma_{s_2,x} = \gamma'_{s_2,x}$.

In particular, these two paths coincide outside $B_{4,s,N}$.

Typical boxes. $B_{4,s,N}$ is called a typical box if it verifies the following properties:

- every geodesic $\gamma_{u,v}$ from u to v entirely contained in $B_{3,s,N}$ with $\|u - v\|_1 \geq N$ has at least $\alpha\|u - v\|_1$ edges whose time is greater than or equal to $t_{\min} + \delta$,
- every path π from u to v entirely contained in $B_{4,s,N}$ with $\|u - v\|_1 \geq N$ has a passage time verifying:

$$t(\pi) \geq (t_{\min} + \delta)\|u - v\|_1, \quad (4.5)$$

- for all u and v in $B_{3,s,N}$, we have

$$(1 - \varepsilon)\mu(u - v) - N \leq t(u, v) \leq (1 + \varepsilon)\mu(u - v) + N.$$

As in the unbounded case, we need properties which are guaranteed with the definition of typical boxes. We state them in the following lemma whose proof is given in Section 4.3.

Lemma 4.3. *We have these three properties about typical boxes.*

- If $B_{4,s,N}$ is a typical box, for all points u_0 and v_0 in $B_{3,s,N}$, every geodesic from u_0 to v_0 is entirely contained in $B_{4,s,N}$.
- The typical box property only depends on the time of the edges in $B_{4,s,N}$.
- We have

$$\lim_{N \rightarrow \infty} \mathbb{P}(B_{4,0,N} \text{ is a typical box}) = 1.$$

Successful boxes. For a fixed $x \in \mathbb{Z}^d$, a box $B_{4,s,N}$ is successful if every geodesic from 0 to x takes a pattern which is entirely contained in $B_{2,s,N}$, i.e. if for every geodesic γ going from 0 to x , there exist $j \in \{1, \dots, d\}$ and $x_\gamma \in \mathbb{Z}^d$ satisfying the condition $(\gamma; \mathfrak{P}^j)$ such that $B_\infty(x_\gamma, \ell^\Lambda)$ is contained in $B_{2,s,N}$.

Annuli. Following the proof in the unbounded case, we define the annuli $A_{i,N}$ with $r = 2(r_1 + r_4 + 1)$ and $\mathcal{G}^p(N)$ as in Section 3.1 but with the definitions of crossed and typical boxes defined here in Section 4.2. The bound on $\mathbb{P}(\mathcal{G}^p(N)^c)$ of Lemma 3.2 also holds here. The proof is exactly the same in this case thanks to Lemma 4.3. For the rest of the proof, we fix C_1 , D_1 and N_0 given by Lemma 3.2.

Modification argument. Fix $K' = T^\Lambda + 2(C_\mu L_1 + t_{\max}(\ell^\Lambda + 1))$. Then, fix

$$N > \max\left(N_0, \frac{12C_\mu K'}{\delta \nabla}\right), \quad n \geq 2rN \text{ and } x \in \Gamma_n, \quad (4.6)$$

(where Γ_n is defined at (1.14)). Fix $p = \lfloor \frac{n}{rN} \rfloor$ and $q = \lfloor \frac{p}{2} \rfloor$. For $j \in \{1, \dots, q\}$, we define Γ^j , S_j^1 , S_j^2 and $\mathcal{M}(j)$ as in Section 3.1 but with the notions of typical and successful boxes defined here in Section 4.2. As in the unbounded case, the aim is to bound from above $\mathbb{P}(T \in \mathcal{M}(q))$ independently of x . For the sequel, we use a two-steps modification. So we introduce two independent copies T' and T'' of the environment T , the three being defined on the same probability space.

Fix $\ell \in \{1, \dots, q\}$. On $\{T \in \mathcal{M}(\ell)\}$, $B_{4,S_\ell^1(T),N}$ is a typical box crossed by the selected geodesic. From this configuration, as a first step, we shall associate a set of edges $E_+^*(T)$ which is contained in $B_{3,S_\ell^1(T),N} \setminus B_{2,S_\ell^1(T),N}$. It corresponds to the edges for which we want to reduce the time. Then, we get a new environment T^* defined for all edge e by:

$$T^*(e) = \begin{cases} T'(e) & \text{if } e \in E_+^*(T) \\ T(e) & \text{else.} \end{cases} \quad (4.7)$$

From this environment, as a second step, we get three new subsets $E_+^{**}(T, T')$, $E_-^{**}(T, T')$ and $E_P^{**}(T, T')$ of edges of $B_{2,S_\ell^1(T),N}$ which are respectively the edges for which we want to reduce the time, to increase the time and the edges of the location where we want to put the pattern. We get a third environment T^{**} defined for all edge e by:

$$T^{**}(e) = \begin{cases} T''(e) & \text{if } e \in E_+^{**}(T, T') \cup E_-^{**}(T, T') \cup E_P^{**}(T, T') \\ T^*(e) & \text{else.} \end{cases} \quad (4.8)$$

Note that T^{**} and T do not have the same distribution. For y and z in \mathbb{Z}^d , we denote by $t^*(y, z)$ (resp. $t^{**}(y, z)$) the geodesic time between y and z in the environment T^* (resp. T^{**}). Similarly, we define for $c \in \mathbb{Z}^d$ and $t \in \mathbb{R}_+$:

$$B^*(c, t) = \{u \in \mathbb{Z}^d : t^*(c, u) \leq t\} \text{ and } B^{**}(c, t) = \{u \in \mathbb{Z}^d : t^{**}(c, u) \leq t\}. \quad (4.9)$$

We formalize this modification in the next lemma and we will describe precisely the construction of E_+^* , E_+^{**} , E_-^{**} and E_P^{**} in the next subsection.

Lemma 4.4. *There exists $\eta = \eta(N)$ such that for all ℓ in $\{1, \dots, q\}$, there exist measurable functions $E_+^* : (\mathbb{R}_+)^{\mathcal{E}} \mapsto \mathcal{P}(\mathcal{E})$, $E_+^{**} : (\mathbb{R}_+^{\mathcal{E}})^2 \mapsto \mathcal{P}(\mathcal{E})$, $E_-^{**} : (\mathbb{R}_+^{\mathcal{E}})^2 \mapsto \mathcal{P}(\mathcal{E})$, $E_P^{**} : (\mathbb{R}_+^{\mathcal{E}})^2 \mapsto \mathcal{P}(\mathcal{E})$ and $\mathcal{O} : (\mathbb{R}_+^{\mathcal{E}})^2 \mapsto \{1, \dots, d\}$ such that:*

(i) $E_+^*(T)$, $E_+^{**}(T, T')$, $E_-^{**}(T, T')$ and $E_P^{**}(T, T')$ are pairwise disjoint and are contained in $B_{3,S_\ell^1(T)}$,

(ii) on the event $\{T \in \mathcal{M}(\ell)\}$, $\mathbb{P}(T' \in \mathcal{B}^*(T) | T) \geq \eta$ and on the event $\{T \in \mathcal{M}(\ell)\} \cap \{T' \in \mathcal{B}^*(T)\}$, $\mathbb{P}(T'' \in \mathcal{B}^{**}(T, T') | T, T') \geq \eta$, where $\{T' \in \mathcal{B}^*(T)\}$ is a shorthand for

$$\{\forall e \in E_+^*(T), T'(e) \leq t_{\min} + \delta'\},$$

and $\{T'' \in \mathcal{B}^{**}(T, T')\}$ is a shorthand for

$$\left\{ \forall e \in E_+^{**}(T, T'), T''(e) \leq t_{\min} + \delta', \forall e \in E_-^{**}(T, T'), T''(e) \geq \nu, \theta_{NS_\ell^1(T)} T'' \in \mathcal{A}_{\mathcal{O}(T, T')}^\Lambda \right\},$$

(iii) $\{T \in \mathcal{M}(\ell)\} \cap \{T' \in \mathcal{B}^*(T)\} \cap \{T'' \in \mathcal{B}^{**}(T, T')\} \subset \{T^{**} \in (\mathcal{M}(\ell-1) \setminus \mathcal{M}(\ell))\}$ and $\|S_\ell^2(T^{**}) - S_\ell^1(T)\|_1 \leq 2r_4$.

The proof of Lemma 4.4 is left to the reader. It is the same as the proof of Lemma 3.3, replacing the use of Lemma 3.5 by the following one.

Lemma 4.5. *There exists $\eta = \eta(N)$ such that for all ℓ in $\{1, \dots, q\}$, there exist measurable functions $E_+^* : (\mathbb{R}_+)^{\mathcal{E}} \mapsto \mathcal{P}(\mathcal{E})$, $E_+^{**} : (\mathbb{R}_+^{\mathcal{E}})^2 \mapsto \mathcal{P}(\mathcal{E})$, $E_-^{**} : (\mathbb{R}_+^{\mathcal{E}})^2 \mapsto \mathcal{P}(\mathcal{E})$, $E_P^{**} : (\mathbb{R}_+^{\mathcal{E}})^2 \mapsto \mathcal{P}(\mathcal{E})$ and $\mathcal{O} : (\mathbb{R}_+^{\mathcal{E}})^2 \mapsto \{1, \dots, d\}$ such that items (i) and (ii) of Lemma 4.4 are satisfied and such that if the event $\{T \in \mathcal{M}(\ell)\} \cap \{T' \in \mathcal{B}^*(T)\} \cap \{T'' \in \mathcal{B}^{**}(T, T')\}$ occurs, then we have the following properties:*

- (i) *in the environment T^{**} , every geodesic from 0 to x takes the pattern inside $B_{2, S_\ell^1(T), N}$,*
- (ii) *for all geodesic $\bar{\gamma}^{**}$ from 0 to x in the environment T^{**} , there exists a geodesic $\bar{\gamma}$ from 0 to x in the environment T such that $\bar{\gamma}$ and $\bar{\gamma}^{**}$ are associated in $B_{4, S_\ell^1(T), N}$,*
- (iii) *there exists a geodesic γ^{**} in the environment T^{**} from 0 to x such that γ^{**} and the selected geodesic γ in the environment T are associated in $B_{4, S_\ell^1(T), N}$.*

The proof of Lemma 4.5 is the aim of Section 4.4. We now conclude the proof of Proposition 1.10 in the bounded case. The following lemma is the counterpart of Lemma 3.6 in this case.

Lemma 4.6. *There exists $\lambda \in (0, 1)$ which does not depend on n and on $x \in \Gamma_n$ such that*

$$\mathbb{P}(T \in \mathcal{M}(q)) \leq \lambda^q.$$

Proof. Let ℓ be in $\{1, \dots, q\}$. For every $s \in \mathbb{Z}^d$ and \mathcal{E}^{**} subset of edges of $B_{3, s, N}$, let us consider the environment $T_{s, \mathcal{E}^{**}}^{**}$ defined for all edge e by:

$$T_{s, \mathcal{E}^{**}}^{**} = \begin{cases} T''(e) & \text{if } e \in \mathcal{E}^{**} \cap B_{2, s, N} \\ T'(e) & \text{if } e \in \mathcal{E}^{**} \cap (B_{3, s, N} \setminus B_{2, s, N}) \\ T(e) & \text{else.} \end{cases}$$

We define $E^{**}(T, T') = E_+^*(T) \cup E_+^{**}(T, T') \cup E_-^{**}(T, T') \cup E_P^{**}(T, T')$. Thus, for every s and \mathcal{E}^{**} , $T_{s, \mathcal{E}^{**}}^{**}$ and T have the same distribution and on the event $\{T \in \mathcal{M}(\ell)\} \cap \{S_\ell^1(T) = s\} \cap \{E^{**}(T, T') = \mathcal{E}^{**}\}$, $T^{**} = T_{s, \mathcal{E}^{**}}^{**}$. Using this environment and writing with indicator functions the result (iii) of Lemma 4.4, we have:

$$\mathbb{1}_{\{T \in \mathcal{M}(\ell)\}} \mathbb{1}_{\{S_\ell^1(T) = s\}} \mathbb{1}_{\{T' \in \mathcal{B}^*(T)\}} \mathbb{1}_{\{E^{**}(T, T') = \mathcal{E}^{**}\}} \mathbb{1}_{\{T'' \in \mathcal{B}^{**}(T, T')\}} \leq \mathbb{1}_{\{T_{s, \mathcal{E}^{**}}^{**} \in \mathcal{M}(\ell-1) \setminus \mathcal{M}(\ell)\}} \mathbb{1}_{\bigcup_{s' \approx s} \{S_\ell^2(T_{s, \mathcal{E}^{**}}^{**}) = s'\}}.$$

We take the expectation on both sides. The right side yields

$$\mathbb{P}\left(T_{s, \mathcal{E}^{**}}^{**} \in \mathcal{M}(\ell-1) \setminus \mathcal{M}(\ell), \bigcup_{s' \approx s} \{S_\ell^2(T_{s, \mathcal{E}^{**}}^{**}) = s'\}\right) = \mathbb{P}\left(T \in \mathcal{M}(\ell-1) \setminus \mathcal{M}(\ell), \bigcup_{s' \approx s} \{S_\ell^2(T) = s'\}\right).$$

For the left side, we have

$$\begin{aligned} & \mathbb{E}\left[\mathbb{1}_{\{T \in \mathcal{M}(\ell)\}} \mathbb{1}_{\{S_\ell^1(T) = s\}} \mathbb{1}_{\{T' \in \mathcal{B}^*(T)\}} \mathbb{1}_{\{E^{**}(T, T') = \mathcal{E}^{**}\}} \mathbb{1}_{\{T'' \in \mathcal{B}^{**}(T, T')\}}\right] \\ &= \mathbb{E}\left[\mathbb{E}\left[\mathbb{E}\left[\mathbb{1}_{\{T \in \mathcal{M}(\ell)\}} \mathbb{1}_{\{S_\ell^1(T) = s\}} \mathbb{1}_{\{T' \in \mathcal{B}^*(T)\}} \mathbb{1}_{\{E^{**}(T, T') = \mathcal{E}^{**}\}} \mathbb{1}_{\{T'' \in \mathcal{B}^{**}(T, T')\}} \middle| T, T'\right] \middle| T\right]\right] \\ &= \mathbb{E}\left[\mathbb{1}_{\{T \in \mathcal{M}(\ell)\}} \mathbb{1}_{\{S_\ell^1(T) = s\}} \mathbb{E}\left[\mathbb{1}_{\{T' \in \mathcal{B}^*(T)\}} \mathbb{1}_{\{E^{**}(T, T') = \mathcal{E}^{**}\}} \mathbb{P}(T'' \in \mathcal{B}^{**}(T, T') | T, T') \middle| T\right]\right]. \end{aligned}$$

Since on the event $\{T \in \mathcal{M}(\ell)\} \cap \{S_\ell^1(T) = s\} \cap \{T' \in \mathcal{B}^*(T)\} \cap \{E^{**}(T, T') = \mathcal{E}^{**}\}$, $\mathbb{P}(T'' \in \mathcal{B}^{**}(T, T') | T, T')$ is bounded from below by η , the left side is bounded from below by

$$\eta \mathbb{E}\left[\mathbb{1}_{\{T \in \mathcal{M}(\ell)\}} \mathbb{1}_{\{S_\ell^1(T) = s\}} \mathbb{E}\left[\mathbb{1}_{\{T' \in \mathcal{B}^*(T)\}} \mathbb{1}_{\{E^{**}(T, T') = \mathcal{E}^{**}\}} \middle| T\right]\right].$$

Then, by summing on all \mathcal{E}^{**} subsets of edges of $B_{3,s,N}$ and writing K a constant which does not depend on s and which bounds from above the number of different subsets of edges of $B_{3,s,N}$, we get

$$\frac{\eta}{K} \mathbb{E} \left[\mathbb{1}_{\{T \in \mathcal{M}(\ell)\}} \mathbb{1}_{\{S_\ell^1(T) = s\}} \mathbb{P}(T' \in \mathcal{B}^*(T) | T) \right] \leq \mathbb{P} \left(T \in \mathcal{M}(\ell-1) \setminus \mathcal{M}(\ell), \bigcup_{s' \approx s} \{S_\ell^2(T) = s'\} \right).$$

Now, on the event $\{T \in \mathcal{M}(\ell)\} \cap \{S_\ell^1(T) = s\}$, $\mathbb{P}(T' \in \mathcal{B}^*(T) | T)$ is bounded from below by η , so the left side is bounded from below by

$$\frac{\eta^2}{K} \mathbb{P}(T \in \mathcal{M}(\ell), S_\ell^1(T) = s).$$

Then, by summing on all $s \in \mathbb{Z}^d$ and writing K' a constant which bounds from above for all $s \in \mathbb{Z}^d$ the number of vertices $s' \in \mathbb{Z}^d$ such that $s' \approx s$, we get

$$\frac{\eta^2}{KK'} \mathbb{P}(T \in \mathcal{M}(\ell)) \leq \mathbb{P}(T \in \mathcal{M}(\ell-1) \setminus \mathcal{M}(\ell)).$$

Now, since $\mathcal{M}(\ell) \subset \mathcal{M}(\ell-1)$,

$$\mathbb{P}(T \in \mathcal{M}(\ell-1) \setminus \mathcal{M}(\ell)) = \mathbb{P}(T \in \mathcal{M}(\ell-1)) - \mathbb{P}(T \in \mathcal{M}(\ell)).$$

Thus,

$$\mathbb{P}(T \in \mathcal{M}(\ell)) \leq \lambda \mathbb{P}(T \in \mathcal{M}(\ell-1)),$$

where $\lambda = \frac{1}{1 + \frac{\eta^2}{KK'}} \in (0, 1)$ does not depend on x . Hence, using $\mathbb{P}(T \in \mathcal{M}(0)) = 1$, we get by induction

$$\mathbb{P}(T \in \mathcal{M}(q)) \leq \lambda^q.$$

□

From Lemma 4.6, the proof of Proposition 1.10 is the same as in the unbounded case.

4.3 Properties of a typical box

In this section, we state and prove the following lemma, which gives us properties of typical boxes useful for the modification argument, and the proof of Lemma 4.3.

Lemma 4.7. *If $B_{4,s,N}$ is a typical box, we have the following properties.*

(i) *For all u and v in $B_{3,s,N}$ with $\|u - v\|_1 \geq K''N$ where $K'' = \frac{1}{\varepsilon c_\mu} > 0$,*

$$(1 - 2\varepsilon)\mu(u - v) \leq t(u, v) \leq (1 + 2\varepsilon)\mu(u - v).$$

(ii) *For all $z \in B_{3,s,N}$, for all $\bar{r} > 0$, for all $N \in \mathbb{N}^*$,*

$$B_{3,s,N} \cap B(z, N\bar{r}) \subset B_{3,s,N} \cap B_\mu \left(z, \frac{N(\bar{r} + 1)}{1 - \varepsilon} \right),$$

and if $\bar{r} \geq \frac{1}{\varepsilon} - 2$,

$$B_{3,s,N} \cap B_\mu \left(z, \frac{N(\bar{r} + 1)}{1 - \varepsilon} \right) \subset B_{3,s,N} \cap B_\mu \left(z, \frac{N\bar{r}}{1 - 2\varepsilon} \right).$$

(iii) *For all $z \in B_{3,s,N}$, for all $\bar{r} > 0$, for all $N \in \mathbb{N}^*$,*

$$B_{3,s,N} \cap B_\mu(z, N\bar{r}) \subset B_{3,s,N} \cap B(z, N((1 + \varepsilon)\bar{r} + 1)),$$

and if $\bar{r} \geq \frac{1}{\varepsilon}$,

$$B_{3,s,N} \cap B(z, N((1 + \varepsilon)\bar{r} + 1)) \subset B_{3,s,N} \cap B(z, (1 + 2\varepsilon)N\bar{r}).$$

Proof. (i) Let u and v be in $B_{3,s,N}$ with $\|u - v\|_1 \geq K''N$. Then, since $B_{4,s,N}$ is a typical box, we have

$$(1 - \varepsilon)\mu(u - v) - N \leq t(u, v) \leq (1 + \varepsilon)\mu(u - v) + N.$$

The requirement on u and v implies

$$\frac{N}{c_\mu} \leq \varepsilon\|u - v\|_1,$$

so $\varepsilon\mu(u - v) \geq N$.

(ii) and (iii) In both cases, it is easy to check that the first inclusion follows from property (iii) of a typical box and an easy computation using (i) shows the second inclusion. \square

Proof of Lemma 4.3.

1. Let $B_{4,s,N}$ be a typical box and u_0 and v_0 two points of $B_{3,s,N}$. Then, taking paths minimizing the distance for the norm $\|\cdot\|_1$ between u_0 and sN and between v_0 and sN , we have $t(u_0, v_0) \leq 2r_3Nt_{\max}$. Then, if a geodesic γ_{u_0, v_0} takes an edge which is not in $B_{4,s,N}$, since $r_4 \geq r_3 + 1$, using the item (ii) of the definition of a typical box leads to

$$T(\gamma_{u_0, v_0}) \geq 2(r_4 - r_3)N(t_{\min} + \delta),$$

which is impossible since $r_4 > \frac{r_3(t_{\min} + \delta + t_{\max})}{t_{\min} + \delta}$. Note that we only use item (ii) of the definition of a typical box to prove this property.

2. It is clear that the first property only depends on the time of edges in $B_{3,s,N}$ and the second only depends on the time of edges in $B_{4,s,N}$. For the third property, by the preceding item, we know that for all points w_1 and w_2 in $B_{3,s,N}$, the knowledge of the time of all edges in $B_{4,s,N}$ allows us to determine $t(w_1, w_2)$, so to know if the two inequalities are satisfied.

3. For each item of the definition of a typical box, we show that the probability that $B_{4,0,N}$ satisfies this item goes to 1. To show that the probability that item (ii) of the definition of a typical box is satisfied goes to 1, we use the same proof as for (ii) in the proof of Lemma 3.1, replacing $r_2 - r_1$ by 1 and $B_{3,s,N}$ by $B_{4,s,N}$. Further, using (4.1) and a similar computation as for item (ii) in the proof of Lemma 3.1, we get:

$$\mathbb{P}(B_{4,0,N} \text{ satisfies (ii)}) \xrightarrow{N \rightarrow \infty} 1.$$

To prove that

$$\mathbb{P}(B_{4,0,N} \text{ satisfies (iii)}) \xrightarrow{N \rightarrow \infty} 1,$$

recall that ε is fixed in Section 4.2 and fix $\rho = \frac{1}{2d((1 + \varepsilon)C_\mu + t_{\max})}$. Let us consider the following property for N large enough to have $\lfloor \rho N \rfloor \neq 0$:

$$\forall u', v' \in \lfloor \rho N \rfloor \mathbb{Z}^d \cap B_{3,0,N}, |t(u', v') - \mu(u' - v')| \leq \varepsilon\mu(u' - v'). \quad (4.10)$$

By (1.17), by stationarity, and since $|\lfloor \rho N \rfloor \mathbb{Z}^d \cap B_{3,0,N}|$ is uniformly bounded in N , we get

$$\mathbb{P}((4.10) \text{ holds}) \xrightarrow{N \rightarrow \infty} 1.$$

Finally, the proof is completed by showing that (4.10) implies that $B_{4,0,N}$ satisfies (iii). Assume (4.10) and let u and v be two vertices in $B_{3,0,N}$. The aim is to show

$$|t(u, v) - \mu(u - v)| \leq \varepsilon\mu(u - v) + N. \quad (4.11)$$

Let $u', v' \in \lfloor \rho N \rfloor \mathbb{Z}^d \cap B_{3,0,N}$ such that $\|u - u'\|_1 \leq d\rho N$ and $\|u - u'\|_1 \leq d\rho N$. Thus,

$$|t(u, v) - \mu(u - v)| \leq |t(u, v) - t(u', v')| + |t(u', v') - \mu(u' - v')| + |\mu(u' - v') - \mu(u - v)|.$$

By (4.10), $|t(u', v') - \mu(u' - v')| \leq \varepsilon\mu(u' - v')$. Furthermore,

$$|\mu(u - v) - \mu(u' - v')| \leq \mu(u - u') + \mu(v - v') \leq (\|u - u'\|_1 + \|v - v'\|_1)C_\mu \leq 2d\rho NC_\mu.$$

Similarly

$$|t(u, v) - t(u', v')| \leq t(u, u') + t(v, v') \leq (\|u - u'\|_1 + \|v - v'\|_1)t_{\max} \leq 2d\rho Nt_{\max},$$

Thus, we get

$$|t(u, v) - \mu(u - v)| \leq \varepsilon\mu(u - v) + 2d\rho N(t_{\max} + (1 + \varepsilon)C_\mu).$$

We get (4.11) thanks to the choice of ρ .

□

4.4 Modification argument

In this section, we prove Lemma 4.5.

4.4.1 Mains ideas of the proof

First ideas. Before proving in detail Lemma 4.5, we give the main ideas of the proof of the modification argument. Fix $x \in \Gamma_n$, $\ell \in \{1, \dots, q\}$ and assume that the event $\{T \in \mathcal{M}(\ell)\}$ occurs. Let us consider γ the selected geodesic, and two vertices u_1, v_1 belonging to γ , chosen in the ℓ -th box of γ , far enough from each other. Since these vertices are in a typical box, the geodesic time between u_1 and v_1 is roughly equal to $\mu(u_1 - v_1)$. We build a path π between u_1 and v_1 which have exactly $\|u_1 - v_1\|_1$ edges (in other words the path is oriented). The idea of the modification is the following:

- we put the pattern in the middle of π ,
- the passage times of the edges of π which are not in the pattern are reduced close to t_{\min} ,
- the passage times of the edges around π are increased closed to t_{\max} .

We denote by T^{**} the modified environment⁸. Let γ^{**} be a geodesic from 0 to x in this environment. Since the modification guarantees that $T^{**}(\gamma) < T(\gamma)$, γ^{**} has to take several edges whose time has been reduced, and thus γ^{**} takes edges of π . To prove that γ^{**} takes the pattern, we begin by showing that γ^{**} takes an edge of π before and after the pattern, which is the aim of Lemma 4.21. So, assume aiming at a contradiction that the first vertex of π taken by γ^{**} is an edge of π after the pattern and denote it by w . Assuming that the effect of the edges of the pattern whose time has been reduced is negligible compared with the size of the path π , we have $T^{**}(\gamma_{0,w}^{**}) \geq t(0, w)$. Furthermore, on the one hand,

$$t(0, w) \leq t(0, u_1) + \underbrace{t(u_1, w)}_{\approx \mu(w - u_1)},$$

since the geodesic time is well approximated by the norm μ since w and u_1 are in a typical box. On the other hand, also using that the geodesic time is well approximated by the norm μ for vertices in a typical box,

$$t(0, w) \geq \underbrace{t(0, v_1)}_{\approx t(0, u_1) + \mu(u_1 - v_1)} - \underbrace{t(v_1, w)}_{\approx \mu(v_1 - w)} \\ \approx \underbrace{t(0, u_1) + \mu(w - u_1)}_{\approx t(0, u_1) + \mu(w - u_1)}$$

Thus $t(0, w) \approx t(0, u_1) + \mu(w - u_1)$ and we get that $T^{**}(\gamma_{0,w}^{**})$ is roughly greater than $t(0, u_1) + \mu(w - u_1)$. But $t^{**}(0, w) \leq t(0, u_1) + T^{**}(\pi_{u_1, w})$ and since the edges of π are closed to t_{\min} , we have $T^{**}(\pi_{u_1, w}) \ll \mu(w - u_1)$. Hence $T^{**}(\gamma_{0,w}^{**}) > t^{**}(0, w)$, which is a contradiction since γ^{**} is a geodesic in the environment T^{**} . Having that γ^{**} visits vertices of π before and after the pattern, using the fact that π is an oriented path, it is not a main difficulty to show that γ^{**} takes the pattern.

⁸The next paragraph will explain the notation $**$.

Two modifications to get associated geodesics. However, it is not sufficient to prove that γ^{**} takes the pattern. With the definition of $\mathcal{M}(\ell)$ given in Section 4.2, we must have a control on the other boxes crossed and the patterns taken by γ^{**} . To this aim, we make the modification described above as a second modification in an environment already modified. A first modification is used to guarantee that there exists a geodesic in the environment T from 0 to x such that this geodesic and γ^{**} are associated in the ℓ -th box of γ . It consists in reducing the time of some edges of γ_{0,u_1} before u_1 and of $\gamma_{v_1,x}$ after v_1 . Thus, in the environment T^* corresponding to this first modification, every geodesic from 0 to x has to take every edge whose time has been reduced, that is some edges of γ_{0,u_1} and of $\gamma_{v_1,x}$. Further, we can guarantee that, after the second modification, in the environment T^{**} , every geodesic also visits edges of γ_{0,u_1} and of $\gamma_{v_1,x}$ by making the time saved by γ during the first modification greater than the geodesic time between u_1 and v_1 (see Lemma 4.18).

Use of balls for the passage time. In the modification described at the beginning of this section, the edges whose time has been reduced can only belong to π or to the pattern but now, it is not the case due to the first modification made before. Thus, even if we prove that a geodesic in the environment obtained after the second modification takes edges of π before and after the pattern, a difficulty appears when we want to prove that it implies that this geodesic takes the pattern. Indeed, we have to justify why it does not exist a shortcut, which does not take the pattern, between a vertex of π before the pattern and one of π after the pattern using edges whose time has been reduced by the first modification. Our idea is to use the balls for the passage time to localize the second modification. We define for $c \in \mathbb{Z}^d$ and $r \in \mathbb{R}_+$:

$$\begin{aligned} B^*(c, r) &= \{u \in \mathbb{Z}^d : t^*(c, u) \leq r\}, \\ B^{**}(c, r) &= \{u \in \mathbb{Z}^d : t^{**}(c, u) \leq r\}, \end{aligned}$$

where t^* (resp. t^{**}) is the geodesic time in the environment T^* (resp. T^{**}) obtained after the first modification (resp. after the second modification). In the second modification, we do not reduce the time of the edges connected to $B^*(0, T^*(\gamma_{0,u_1}))$ and of those connected to $B^*(x, T^*(\gamma_{v_1,x}))$. We prove that there are enough edges of π outside these balls and that the place where we put the pattern is also outside these balls. As a consequence $B^{**}(0, T^*(\gamma_{0,u_1})) = B^*(0, T^*(\gamma_{0,u_1}))$ and $B^{**}(x, T^*(\gamma_{v_1,x})) = B^*(x, T^*(\gamma_{v_1,x}))$. Thus, when a geodesic in the environment T^{**} visits a vertex of π whose time has been reduced, it has left $B^{**}(0, T^*(\gamma_{0,u_1}))$ and it can not take vertices of this ball anymore. The same argument holds for $B^{**}(x, T^*(\gamma_{v_1,x}))$ and it allows us to guarantee that it does not exist shortcuts as those described at the beginning of the paragraph.

4.4.2 First modification

Let us go back to the proof. For the rest of Section 4.4, we fix $\ell \in \{1, \dots, q\}$ and $s \in \mathbb{Z}^d$, and we assume that the event $\{T \in \mathcal{M}(\ell)\} \cap \{S_\ell^1(T) = s\}$ occurs. In particular, $B_{4,s,N}$ is a typical box in the environment T .

As explained in the paragraph above, the first modification is used to get that any geodesic from 0 to x in the new environment goes from 0 to a specific vertex in $B_{4,s,N}$, then follows γ to another specific vertex in $B_{4,s,N}$ and then goes to x . It is useful to get the properties about associated geodesics after the second modification. We denote by γ the selected geodesic in the environment T . Recall the definition of entry and exit points given at the beginning of Section 3.3. Let u denote the entry point of γ in $B_{3,s,N}$ and v the exit point, and let u_0 denote the entry point of γ in $B_{2,s,N}$ and v_0 its exit point. Then, we set

$$E_+^*(T) = \{\text{edges of } B_{3,s,N} \text{ that belong to } \gamma_{u,u_0} \text{ or } \gamma_{v_0,v} \text{ and satisfy } T(e) > t_{\min} + \delta\}. \quad (4.12)$$

We denote the first edge of γ that belongs to $E_+^*(T)$ by e_1 and the last one by e_2 . Moreover, s_1 denote the first vertex of e_1 visited by γ and s_2 the last vertex of e_2 visited by γ . Assume that the event

$$\{T \in \mathcal{M}(\ell)\} \cap \{S_\ell^1(T) = s\} \cap \{T' \in \mathcal{B}^*(T)\} \text{ occurs,}$$

where $\mathcal{B}^*(T)$ is defined in (ii) of Lemma 4.4. Recall the definition of T^* in (4.7). Note that:

- only the time of the edges of $E_+^*(T)$ have been modified,

4.4.3 Construction of π

Here, we shall identify an oriented path π between two vertices of γ in $B_{2,s,N}$. This oriented path is later used to place the pattern. Let c_0 denote the entry point of γ in $B_{1,s,N}$ and recall the definition of ∇ in (4.2). Since by (4.3), $r_2 \geq r_1 + \frac{\nabla}{c_\mu}$, $B_\mu(c_0, N\nabla) \cap \mathbb{Z}^d$ is entirely contained in $B_{2,s,N}$. We introduce

$$u_1 \text{ the entry point of } \gamma \text{ in } B_\mu(c_0, N\nabla) \cap \mathbb{Z}^d \text{ and } v_1 \text{ the exit point.} \quad (4.13)$$

Lemma 4.9. *We have $\mu(u_1 - v_1) \geq N\nabla$ and γ_{u_1, v_1} is contained in $B_{2,s,N}$.*

Remark 4.10. *The idea of the proof is that $\mu(u_1 - v_1)$ is roughly equal to $t(u_1, v_1)$. Furthermore, since γ is a geodesic visiting u_1 , c_0 and v_1 in this order, we have $t(u_1, v_1) = t(u_1, c_0) + t(c_0, v_1)$. So $\mu(u_1 - v_1) \approx \mu(u_1 - c_0) + \mu(v_1 - c_0) \approx 2N\nabla$. We have a sufficient control over the approximations to guarantee a lower bound by $N\nabla$.*

Proof. Using item (ii) of Lemma 4.7 with $z = c_0$ and $\bar{r} = \nabla(1 - \varepsilon) - 1$ leads to

$$B(c_0, N(\nabla(1 - \varepsilon) - 1)) \cap B_{3,s,N} \subset B_\mu(c_0, N\nabla) \cap B_{3,s,N} = B_\mu(c_0, N\nabla) \cap \mathbb{Z}^d. \quad (4.14)$$

Since u_1 is the entry point of a path in $B_\mu(c_0, N\nabla)$, there exists a vertex $\bar{u}_1 \in B_{3,s,N}$ such that $\|\bar{u}_1 - u_1\|_1 = 1$ and $\bar{u}_1 \notin B_\mu(c_0, N\nabla)$. By (4.14), $t(c_0, \bar{u}_1) \geq N(\nabla(1 - \varepsilon) - 1)$, so

$$t(c_0, u_1) \geq t(c_0, \bar{u}_1) - \underbrace{t(\bar{u}_1, u_1)}_{\leq \|\bar{u}_1 - u_1\|_1 t_{\max}}.$$

The same argument holds for $t(c_0, v_1)$. Hence

$$t(u_1, v_1) = t(u_1, c_0) + t(c_0, v_1) \geq 2N(\nabla(1 - \varepsilon) - 1) - 2t_{\max}.$$

Thus, since $\nabla > \frac{3 + 2t_{\max}}{1 - 3\varepsilon}$, by the third item of the definition of a typical box,

$$\mu(u_1 - v_1) \geq \frac{N(2\nabla(1 - \varepsilon) - 3) - 2t_{\max}}{1 + \varepsilon} \geq N\nabla.$$

For the second part of the proof, using the third item of Lemma 4.7 with $z = c_0$ and $\bar{r} = \nabla$ leads to

$$B_\mu(c_0, N\nabla) \cap \mathbb{Z}^d \subset B(c_0, ((1 + \varepsilon)\nabla + 1)N).$$

Then, by (4.3) we have $r_2 > r_1 + \frac{2(\nabla + 2)}{c_\mu}$. So, for all $z \in B_\mu(c_0, 2N(\nabla + 2))$, we have

$$\|z - sN\|_1 \leq \|z - c_0\|_1 + \|c_0 - sN\|_1 \leq \left(\frac{2(\nabla + 2)}{c_\mu} + r_1 \right) N < r_2 N.$$

So, we have $B_\mu(c_0, 2N(\nabla + 2)) \cap \mathbb{Z}^d \subset B_{2,s,N}$ and since $\varepsilon < \frac{1}{3}$, we have

$$B_\mu \left(c_0, N \frac{(1 + \varepsilon)\nabla + 2}{1 - \varepsilon} \right) \subset B_\mu(c_0, 2N(\nabla + 2)).$$

Thus, by the second item of Lemma 4.7 with $z = c_0$ and $\bar{r} = \nabla$, we get

$$B(c_0, ((1 + \varepsilon)\nabla + N)) \subset B_\mu \left(c_0, N \frac{(1 + \varepsilon)\nabla + 2}{1 - \varepsilon} \right) \cap \mathbb{Z}^d \subset B_\mu(c_0, 2N(\nabla + 2)) \cap \mathbb{Z}^d.$$

To sum up,

$$B_\mu(c_0, N\nabla) \cap \mathbb{Z}^d \subset B(c_0, (1 + \varepsilon)N\nabla + N) \subset B_{2,s,N}.$$

Now, assume that γ_{u_1, v_1} visits a vertex which is not in $B_{2,s,N}$. Let denote by z such a vertex and assume for example that z is visited by γ_{u_1, c_0} . Then, we have, thanks to these inclusions, $t(c_0, z) > t(c_0, u_1)$, which is impossible since γ_{u_1, c_0} is a geodesic. \square

Lemma 4.11. *Recall the definition of B^* given at (4.9) and that $T^*(\gamma_{0,u_1}) = t^*(0, u_1)$ and $T^*(\gamma_{v_1,x}) = t^*(v_1, x)$.*

(i) *We have the following inclusions:*

$$B^*(0, T^*(\gamma_{0,u_1})) \subset B(0, t(0, u_1)) \text{ and } B^*(x, T^*(\gamma_{v_1,x})) \subset B(x, t(v_1, x)).$$

(ii) *We have*

$$B^*(0, T^*(\gamma_{0,u_1})) \cap B^*(x, T^*(\gamma_{v_1,x})) = \emptyset.$$

Proof. (i) Let s' be a vertex in $B^*(0, T^*(\gamma_{0,u_1}))$. The aim is to show

$$t(0, s') \leq t(0, u_1).$$

Let $\bar{\gamma}^*$ be a geodesic from 0 to s' in the environment T^* . We denote by s^* the last vertex visited by $\bar{\gamma}^*$ among those visited by γ (note that 0 is such a vertex). First, since $\bar{\gamma}_{s^*,s'}^*$ does not take an edge of γ , $T^*(\bar{\gamma}_{s^*,s'}^*) = T(\bar{\gamma}_{s^*,s'}^*)$. Then, by Lemma 4.9, γ_{u_1,v_1} is entirely contained in $B_{2,s,N}$, so $T^*(\gamma_{u_1,v_1}) = T(\gamma_{u_1,v_1})$. Thus, also by Lemma 4.9, $T^*(\gamma_{u_1,v_1}) > 0$. So $\bar{\gamma}^*$ does not take an edge of $\gamma_{v_1,x}$. Otherwise, since γ is a geodesic in the environment T^* , $t^*(0, s') \geq T^*(\gamma_{0,v_1}) > T^*(\gamma_{0,u_1})$. Hence, the time saved by $\bar{\gamma}^*$ after the modification comes only from the edges of γ_{0,u_1} . So,

$$T(\bar{\gamma}_{0,s^*}^*) - T^*(\bar{\gamma}_{0,s^*}^*) \leq T(\gamma_{0,u_1}) - T^*(\gamma_{0,u_1}). \quad (4.15)$$

Hence,

$$\begin{aligned} t(0, s') &\leq T(\bar{\gamma}_{0,s^*}^*) + T(\bar{\gamma}_{s^*,s'}^*) \\ &\leq T(\gamma_{0,u_1}) - T^*(\gamma_{0,u_1}) + \underbrace{T^*(\bar{\gamma}_{0,s^*}^*) + T^*(\bar{\gamma}_{s^*,s'}^*)}_{=t^*(0,s')} \text{ by (4.15),} \\ &= t^*(0, s') + T(\gamma_{0,u_1}) - T^*(\gamma_{0,u_1}) \\ &\leq T^*(\gamma_{0,u_1}) + T(\gamma_{0,u_1}) - T^*(\gamma_{0,u_1}) \text{ since } s' \in B^*(0, T^*(\gamma_{0,u_1})), \\ &= t(0, u_1), \end{aligned}$$

which proves the inclusion and the same proof gives us the second inclusion.

(ii) Let s' be a point of $B^*(0, T^*(\gamma_{0,u_1})) \cap B^*(x, T^*(\gamma_{v_1,x}))$. Then

$$\begin{aligned} t(0, x) &\leq t(0, s') + t(s', x) \\ &\leq t(0, u_1) + t(v_1, x) \text{ by (i)} \\ &< t(0, x) \end{aligned}$$

since γ is a geodesic visiting 0, u_1 , v_1 and x in this order and since $t(u_1, v_1) > 0$, which is a contradiction. \square

Now, we can make the construction of π , which is the path on which we would like to put the pattern that the geodesics have to take. We begin by two definitions:

- We say that a path between two vertices y_1 and y_2 is oriented if its number of edges is equal to $\|y_1 - y_2\|_1$.
- A step of length ℓ is a path of ℓ consecutive edges in the same direction.

We state the following lemma whose proof is left to the reader and where $[u_1, v_1]$ is the segment in \mathbb{R}^d and $\|\cdot\|_1$ is the norm on \mathbb{R}^d .

Lemma 4.12. *We can construct a path π with a deterministic rule such that:*

- (i) π is an oriented path connecting u_1 and v_1 ,
- (ii) π is the concatenation of steps of length $10\ell^\Lambda$ except in $B_\infty(v_1, 10\ell^\Lambda)$, where π takes steps of length 1,
- (iii) there exists a constant $L_1 \in \mathbb{R}_+$ only depending on ℓ^Λ and d such that for all $z \in \mathbb{Z}^d$ which is in π , the distance for the norm $\|\cdot\|_1$ between z and $[u_1, v_1]$ is bounded by L_1 , and for all $y \in [u_1, v_1]$, the distance for the norm $\|\cdot\|_1$ between y and π (seen as a set of vertices in \mathbb{Z}^d) is bounded by L_1 .

Let π be the path given by Lemma 4.12. We introduce

$$u_2 \text{ the first vertex of } \pi \text{ starting from } v_1 \text{ in } B^*(0, T^*(\gamma_{0, u_1})), \quad (4.16)$$

$$v_2 \text{ the first vertex of } \pi \text{ starting from } u_1 \text{ in } B^*(x, T^*(\gamma_{v_1, x})). \quad (4.17)$$

Remark that $t^*(0, u_2) \leq t^*(0, u_1)$ and $t^*(v_2, x) \leq t^*(v_1, x)$.

Lemma 4.13. *The path π is contained in $B_{2, s, N}$.*

Proof. Let z be a vertex of π . Using Lemma 4.12, there exists $y \in [u_1, v_1]$ such that $\|z - y\|_1 \leq L_1$. The vertices u_1 and v_1 belong to $B_\mu(c_0, N\nabla)$ which is convex, thus $y \in B_\mu(c_0, N\nabla)$. So, we have

$$\begin{aligned} \|z - sN\|_1 &\leq \underbrace{\|z - y\|_1}_{\leq L_1} + \underbrace{\|y - c_0\|_1}_{\leq \frac{\mu(y - c_0)}{c_\mu}} + \underbrace{\|c_0 - sN\|_1}_{\leq r_1 N} \\ &\leq L_1 + \frac{N\nabla}{c_\mu} + r_1 N \text{ since } y \in B_\mu(c_0, N\nabla), \\ &\leq r_2 N \text{ by (4.3),} \end{aligned}$$

which proves that $z \in B_{2, s, N}$. □

4.4.4 Second modification

Now, let us define E_P^{**} , E_+^{**} , E_-^{**} and \mathcal{O} . Let $c_1 \in \mathbb{R}^d$ denote the midpoint of $[u_1, v_1]$ and let us consider the set of all vertices of π_{u_2, v_2} at distance at most L_1 for the norm $\|\cdot\|_1$ from c_1 . This set is not empty, hence we can choose one such vertex with a deterministic rule. Recall that $L_2 = L_1 + (10 + d)\ell^\Lambda$. Since $\nabla \geq 6dL_2C_\mu$ (see (4.2)), we have that the distance for the norm $\|\cdot\|_\infty$ between the chosen vertex and v_1 is greater than $10\ell^\Lambda$. So there exist one or two steps of the path π of length $10\ell^\Lambda$ that contain the chosen vertex. We chose one step (between these one or two steps) with a deterministic rule to put the pattern. We denote the midpoint of this chosen step by c_P . Then,

$$\mathcal{O}(T, T') \text{ is equal to the direction of this step,} \quad (4.18)$$

$$\text{and } E_P^{**}(T, T') = \{\text{edges connecting vertices belonging to } B_\infty(c_P, \ell^\Lambda)\}. \quad (4.19)$$

Note that by the direction of a step, we mean the integer j such that for every distinct vertices z_1 and z_2 of this step, $z_1 - z_2 = \pm\|z_1 - z_2\|_1 \varepsilon_j$. Note also that for all vertex z in $B_\infty(c_P, \ell^\Lambda)$,

$$\|z - c_1\|_1 \leq L_2.$$

We define u_3 and v_3 as the endpoints of the oriented pattern such that π visits u_2 , u_3 , v_3 and v_2 in this order. Recall that for a pattern, the entry and exit points correspond to the endpoints given by the definition of this pattern. Note that, thanks to the construction, u_3 and v_3 are two vertices of the chosen step of length $10\ell^\Lambda$. We denote by γ^π the path composed by the first geodesic in the lexicographic order from 0 to u_2 in the environment T^* , then π_{u_2, v_2} and then the first geodesic in the lexicographic order from v_2 to x in the environment T^* . Then, we have to define $E_+^{**}(T, T')$ and $E_-^{**}(T, T')$:

$$E_+^{**}(T, T') = \{\text{edges of } \pi_{u_2, u_3} \cup \pi_{v_3, v_2} \text{ except the one connected to } u_2 \text{ and the one connected to } v_2\}, \quad (4.20)$$

$$\begin{aligned} E_-^{**}(T, T') &= \{\text{edges } e \text{ in } B_{2, s, N} \text{ such that } T(e) < \nu \text{ and which are not} \\ &\text{in } E_P^{**}(T, T'), \pi_{u_2, u_3} \cup \pi_{v_3, v_2}, B^*(0, T^*(\gamma_{0, u_1})) \text{ or } B^*(x, T^*(\gamma_{v_1, x}))\}. \end{aligned} \quad (4.21)$$




Figure 4: The construction of π , of the vertices $u_1, u_2, u_3, v_1, v_2, v_3$ and of the pattern-location in dimension 2.

For all the sequel, we assume that the event

$$\{T \in \mathcal{M}(\ell)\} \cap \{S_\ell^1(T) = s\} \cap \{T' \in \mathcal{B}^*(T)\} \cap \{T'' \in \mathcal{B}^{**}(T, T')\} \text{ occurs.}$$

Recall the definition of the environment T^{**} given at (4.8) and of $\mathcal{B}^{**}(T, T')$ given in Lemma 4.4. In particular,

- for all $e \in E_+^*(T)$, $T^{**}(e) = T^*(e) < T(e)$,
- for all $e \in E_+^{**}(T, T')$, $T^{**}(e) \leq t_{\min} + \delta'$,
- for all $e \in E_-^{**}(T, T')$, $T^{**}(e) \geq \nu > T(e) = T^*(e)$,
- since by Lemma 4.13, π is contained in $B_{2,s,N}$, we have that $E_+^{**}(T, T')$, $E_-^{**}(T, T')$ and $E_P^{**}(T, T')$ are pairwise disjoint and are contained in $B_{2,s,N}$. Thus, since $E_+^*(T)$ is contained in $B_{3,s,N} \setminus B_{2,s,N}$, item (i) of Lemma 4.4 is satisfied.

In what follows, when we talk about edges whose time has been reduced, it means the edges e such that $T^{**}(e) < T(e)$. Before proving item (ii) of Lemma 4.4, we state the following lemma which completes the vision of the sets $E_+^{**}(T, T')$, $E_-^{**}(T, T')$ and $E_P^{**}(T, T')$. Recall the definition of B^{**} given at (4.9).

Lemma 4.14. *We have $B^{**}(0, T^*(\gamma_{0,u_1})) = B^*(0, T^*(\gamma_{0,u_1}))$, $B^{**}(x, T^*(\gamma_{v_1,x})) = B^*(x, T^*(\gamma_{v_1,x}))$ and there is no edge of $E_P^{**}(T, T')$ in any of these balls.*

Proof. We begin by proving that there is no edge of $E_P^{**}(T, T')$ in $B^*(0, T^*(\gamma_{0,u_1}))$. To this aim, we prove that there is no vertex of $B_\infty(c_P, \ell^\Lambda)$ in this ball. Let z be a vertex of $B_\infty(c_P, \ell^\Lambda)$. The idea is the following. Since γ is a geodesic in the environment T^* , since the time of the edges of γ_{u_1, v_1} is not modified by the first modification and since $B_{4,s,N}$ is a typical box in the environment T , for every w vertex of γ_{u_1, v_1} ,

$$t^*(0, w) = t^*(0, u_1) + t^*(u_1, w) = t^*(0, u_1) + t(u_1, w) \approx t^*(0, u_1) + \mu(u_1 - w).$$

We can not guarantee that z is a vertex of γ_{u_1, v_1} but using a similar argument:

$$t^*(0, z) \geq t^*(0, v_1) - t^*(v_1, z) = t^*(0, u_1) + t^*(u_1, v_1) - t^*(v_1, z),$$

since γ is a geodesic in the environment T^* . But

$$t^*(u_1, v_1) = t(u_1, v_1) \approx 2N\nabla \text{ and } t^*(v_1, z) \leq t^*(v_1, c_1) + t^*(c_1, z) \approx N\nabla,$$

since $t^*(c_1, z)$ is negligible compared to $N\nabla$, which gives that $t^*(0, z)$ is roughly greater than $t^*(0, u_1) + N\nabla$.

Now, we make the proof rigorous. Since $z \in B_\infty(c_P, \ell^\Lambda)$, we have $z \in B_\mu(c_1, C_\mu L_2)$. Since by (4.2), $\nabla \geq 2C_\mu L_2$, $z \in B_\mu\left(c_1, \frac{N\nabla}{2}\right)$. Furthermore, since c_1 is the midpoint of $[u_1, v_1]$ and $\mu(u_1 - v_1) \leq 2N\nabla$, we have $\mu(v_1 - c_1) \leq N\nabla$. Thus, since $\nabla \geq \frac{1}{\varepsilon}$ (by (4.2)), using the third item of Lemma 4.7 with $z = v_1$ and $\bar{r} = \frac{3\nabla}{2}$, we have

$$z \in B_\mu\left(v_1, \frac{3N\nabla}{2}\right) \cap \mathbb{Z}^d \subset B\left(v_1, \frac{3N\nabla(1+2\varepsilon)}{2}\right).$$

Hence, using the lower bound for $t(u_1, v_1)$ of the proof of Lemma 4.9 and since by (4.2), $\nabla > \frac{4(1+t_{\max})}{1-10\varepsilon}$, we have

$$t(0, v_1) - t(0, u_1) = t(u_1, v_1) \geq 2N(\nabla(1-\varepsilon) - 1) - 2t_{\max} > \frac{3N\nabla(1+2\varepsilon)}{2},$$

and thus

$$t(0, z) \geq t(0, v_1) - t(v_1, z) \geq t(0, v_1) - \frac{3N\nabla(1+2\varepsilon)}{2} > t(0, u_1).$$

The first item of Lemma 4.11 allows us to conclude.

Then, let us prove the first equality. The proof of the second one is the same. The inclusion $B^*(0, T^*(\gamma_{0, u_1})) \subset B^{**}(0, T^*(\gamma_{0, u_1}))$ is easy to check. Let us take $z \in B^*(0, T^*(\gamma_{0, u_1}))$ and $\bar{\gamma}^*$ a geodesic from 0 to z in the environment T^* . Then $\bar{\gamma}^*$ is entirely contained in $B^*(0, T^*(\gamma_{0, u_1}))$ and there are no edges of $E_-^{**}(T, T')$ or $E_P^{**}(T, T')$ in $B^*(0, T^*(\gamma_{0, u_1}))$, so

$$t^{**}(0, z) \leq T^{**}(\bar{\gamma}^*) \leq T^*(\bar{\gamma}^*) \leq T^*(\gamma_{0, u_1}).$$

For the other inclusion, assume that there exists a vertex $z \in B^{**}(0, T^*(\gamma_{0, u_1})) \setminus B^*(0, T^*(\gamma_{0, u_1}))$ and let $\bar{\gamma}^{**}$ be a geodesic from 0 to z in the environment T^{**} . Let ω be the first vertex of $\bar{\gamma}^{**}$ which is not in $B^*(0, T^*(\gamma_{0, u_1}))$. By construction, there is no edge of $\bar{\gamma}_{0, \omega}^{**}$ in $E_+^{**}(T, T')$ or $E_P^{**}(T, T')$ and thus:

$$t^{**}(0, z) \geq t^{**}(0, \omega) \geq t^*(0, \omega) > T^*(\gamma_{0, u_1}),$$

which is a contradiction. \square

Now, to get item (ii) of Lemma 4.4, fix $\eta = \min_{j \in \{1, \dots, d\}} \mathbb{P}(T \in \mathcal{A}_j^\Lambda) \tilde{p}^{|B_{3, s, N}|}$, where

$$\tilde{p} = \min(F([t_{\min}, t_{\min} + \delta']), F([\nu(N), t_{\max}])) > 0.$$

Thus, η only depends on F , the pattern and N and we have that

$$\mathbb{P}(T' \in \mathcal{B}^*(T) | T) \geq \tilde{p}^{|B_{3, s, N}|} \geq \eta$$

and

$$\mathbb{P}(T'' \in \mathcal{B}^{**}(T, T') | T, T') \geq \min_{j \in \{1, \dots, d\}} \mathbb{P}(T \in \mathcal{A}_j^\Lambda) \tilde{p}^{|B_{3, s, N}|} = \eta.$$

We end this section with two lemmas, one giving a lower bound on the time saved by the geodesics from 0 to x thanks to the second modification and the other on the distance between u_1 and any vertex of π_{v_3, v_2} .

Lemma 4.15. *For all $N \in \mathbb{N}^*$, we have*

$$T^*(\gamma) - T^{**}(\gamma^\pi) \geq \frac{N\nabla}{2C_\mu}(\delta - \delta') > T^\Lambda.$$

Proof. This result is an easy consequence of Lemma 4.9. Since $\nabla \geq C_\mu$ (by (4.2)) and all edges of γ_{u_1, v_1} are contained in $B_{2, s, N}$, which implies that there is no edge of γ_{u_1, v_1} whose time has been reduced between the environment T and T^* ,

$$T^*(\gamma) \geq T^*(\gamma_{0, u_1}) + \|u_1 - v_1\|_1(t_{\min} + \delta) + T^*(\gamma_{v_1, x}),$$

where we used (i) of the definition of a typical box. Further

$$T^{**}(\gamma^\pi) \leq T^{**}(\gamma_{0, u_2}^\pi) + \|u_2 - v_2\|_1(t_{\min} + \delta') + T^{**}(\gamma_{v_2, x}^\pi) + 2t_{\max} + \tau^\Lambda,$$

where the term $2t_{\max}$ is an upper bound for the time for both, the edge connecting u_2 to $E_+^{**}(T, T')$ and the one connecting v_2 to $E_+^{**}(T, T')$, and the term τ^Λ is an upper bound for the time collected by γ^π in $E_P^{**}(T, T')$. Since we have $T^{**}(\gamma_{0, u_2}^\pi) = T^*(\gamma_{0, u_2}^\pi) \leq T^*(\gamma_{0, u_1})$, $T^{**}(\gamma_{v_2, x}^\pi) = T^*(\gamma_{v_2, x}^\pi) \leq T^*(\gamma_{v_1, x})$, $\|u_2 - v_2\|_1 \leq \|u_1 - v_1\|_1$ since π is an oriented path, and $\nabla \geq 4C_\mu(2t_{\max} + \tau^\Lambda)$ (by (4.2)), and using Lemma 4.9, we obtain for all $N \in \mathbb{N}^*$,

$$T^*(\gamma) - T^{**}(\gamma^\pi) \geq \|u_1 - v_1\|_1(\delta - \delta') - \tau^\Lambda - 2t_{\max} \geq \frac{N\nabla}{2C_\mu}(\delta - \delta').$$

Finally, since by (4.2), $\nabla > \frac{2C_\mu T^\Lambda}{\delta - \delta'}$ we have the result. \square

Lemma 4.16. *For all $N \in \mathbb{N}^*$, for all w vertex of π_{v_3, v_2} , we have*

$$\|u_1 - w\|_1 \geq \frac{\|u_1 - v_1\|_1}{3}.$$

Proof. Let us note that since π is an oriented path, for all w vertex of $\pi_{u_2, u_3} \cup \pi_{v_3, v_2}$ after $E_P^{**}(T, T')$, we have $\|u_1 - w\|_1 \geq \|u_1 - v_3\|_1$. So

$$\|u_1 - w\|_1 \geq \|u_1 - c_1\|_1 - \|c_1 - v_3\|_1 \geq \frac{1}{2}\|u_1 - v_1\|_1 - L_2.$$

Then, since by (4.2), $\nabla \geq 6L_2C_\mu$, we have $\frac{N\nabla}{6C_\mu} \geq L_2$ and using Lemma 4.9 leads to the result. \square

4.4.5 Proof of the second and third items of Lemma 4.5

Recall that we assume that

$$\{T \in \mathcal{M}(\ell)\} \cap \{S_\ell^1(T) = s\} \cap \{T' \in \mathcal{B}^*(T)\} \cap \{T'' \in \mathcal{B}^{**}(T, T')\} \text{ occurs.}$$

The aim of this section is to prove the following properties (which are the second and third items of Lemma 4.5):

- (ii) for all geodesic $\bar{\gamma}^{**}$ from 0 to x in the environment T^{**} , there exists a geodesic $\bar{\gamma}$ from 0 to x in the environment T such that $\bar{\gamma}$ and $\bar{\gamma}^{**}$ are associated in $B_{4, S_\ell^1(T), N}$,
- (iii) there exists a geodesic γ^{**} in the environment T^{**} from 0 to x such that γ^{**} and the selected geodesic γ in the environment T are associated in $B_{4, S_\ell^1(T), N}$.

To prove this, we use the following sequence of lemmas.

Lemma 4.17. *Every geodesic from 0 to x in the environment T^{**} takes at least one edge of $\pi_{u_2, u_3} \cup \pi_{v_3, v_2}$.*

Proof. Let γ^{**} be a geodesic from 0 to x in the environment T^{**} . Since γ is a geodesic in the environment T^* , we have the following inequalities

$$T^{**}(\gamma^{**}) \leq T^{**}(\gamma^\pi) < T^*(\gamma) \leq T^*(\gamma^{**}).$$

So, using Lemma 4.15,

$$T^*(\gamma^{**}) - T^{**}(\gamma^{**}) > T^\Lambda.$$

It means that γ^{**} has to take at least one edge whose time has been reduced during the second modification which is not in $E_P^{**}(T, T')$. Hence, since the only edges which are not in $E_P^{**}(T, T')$ whose time has been reduced are edges of $\pi_{u_2, u_3} \cup \pi_{v_3, v_2}$, the result follows. \square

Lemma 4.18. *Every geodesic from 0 to x in the environment T^{**} takes at least one edge of γ whose time has been reduced before taking its first edge of $B_{2,s,N}$, and takes at least one edge of γ whose time has been reduced after taking its last edge of $B_{2,s,N}$.*

Proof. The idea is to use that the time saved by the geodesics from 0 to x after the first modification is much greater than the geodesic time between any two vertices of $B_{2,s,N}$ in any environment. Let γ^{**} be a geodesic from 0 to x in the environment T^{**} . Let u^{**} be the first vertex in $B_{2,s,N}$ that γ^{**} visits. Its existence is guaranteed by Lemma 4.17. The aim of the proof is to show

$$T^{**}(\gamma_{0,u^{**}}^{**}) < T(\gamma_{0,u^{**}}^{**}).$$

Indeed, the definition of u^{**} and the fact that the only edges whose time has been reduced which are in $B_{3,s,N}$ but not in $B_{2,s,N}$ are edges of γ gives us the result. Recall that u_0 is the entry point of γ in $B_{2,s,N}$. First, since γ^{**} is a geodesic in the environment T^{**} ,

$$T^{**}(\gamma_{0,u^{**}}^{**}) \leq T^{**}(\gamma_{0,u_0}) + 2r_2 N t_{\max}.$$

Then, using the first item of Lemma 4.8, we obtain

$$T^{**}(\gamma_{0,u^{**}}^{**}) \leq T(\gamma_{0,u_0}) + 2r_2 N t_{\max} - \alpha(r_3 - r_2)N(\delta - \delta').$$

Finally, using the fact that γ is a geodesic in the environment T leads to

$$T^{**}(\gamma_{0,u^{**}}^{**}) \leq T(\gamma_{0,u^{**}}^{**}) + 4r_2 N t_{\max} - \alpha(r_3 - r_2)N(\delta - \delta').$$

To conclude, it is sufficient to observe that the condition $r_3 > \frac{7r_2(4t_{\max} + \alpha\delta)}{\alpha\delta}$ implies that $\alpha(r_3 - r_2)N(\delta - \delta') - 4r_2 N t_{\max} > 0$, so we have the desired strict inequality. The same proof gives us the second part of the lemma. \square

Lemma 4.19. *Let γ^{**} be a geodesic from 0 to x in the environment T^{**} . Then the first edge of γ^{**} whose time has been reduced is e_1 and the last is e_2 . Moreover, the first vertex of e_1 taking by γ^{**} is s_1 and the last of e_2 is s_2 .*

Proof. Let γ^{**} be a geodesic from 0 to x in the environment T^{**} . Let z^{**} denote the last vertex visited by γ^{**} before it takes for the first time an edge of γ whose time has been reduced. We know by the construction and by Lemma 4.18 that z^{**} is a vertex visited by γ_{u,u_0} or $\gamma_{v_0,v}$, and thus it is a vertex visited by γ_{u,u_1} or $\gamma_{v_1,v}$. Let us prove that it is a vertex visited by γ_{u,u_1} . Assume, aiming at a contradiction that z^{**} is a vertex visited by $\gamma_{v_1,v}$.

On the one hand, by Lemma 4.18, $\gamma_{0,z^{**}}^{**}$ does not take edges in $B_{2,s,N}$ and since by Lemma 4.13, π is contained in $B_{2,s,N}$, $\gamma_{0,z^{**}}^{**}$ does not take any edge of $\pi_{u_2,u_3} \cup \pi_{v_3,v_2}$.

On the other hand, all edges of $\gamma_{v_1,x}$ are in $B^*(x, T^*(\gamma_{v_1,x}))$. So, if z^{**} is visited by $\gamma_{v_1,x}$, we have $z^{**} \in B^*(x, T^*(\gamma_{v_1,x}))$. But $B^{**}(x, T^*(\gamma_{v_1,x})) = B^*(x, T^*(\gamma_{v_1,x}))$ by Lemma 4.14. Since γ^{**} is a geodesic in the environment T^{**} , it implies that $\gamma_{z^{**},x}^{**}$ is entirely contained in $B^*(x, T^*(\gamma_{v_1,x}))$. Thus $\gamma_{z^{**},x}^{**}$ does not take any edge of $\pi_{u_2,u_3} \cup \pi_{v_3,v_2}$ since there is no edge of $\pi_{u_2,u_3} \cup \pi_{v_3,v_2}$ in $B^*(x, T^*(\gamma_{v_1,x}))$.

Combining these two conclusions, we get that γ^{**} does not visit any edge of $\pi_{u_2,u_3} \cup \pi_{v_3,v_2}$, which contradicts Lemma 4.17. So z^{**} is a vertex visited by γ_{u,u_1} . Knowing this, we can complete the proof. By the definition of z^{**} and by Lemma 4.18, we have

$$T(\gamma_{0,z^{**}}^{**}) = T^{**}(\gamma_{0,z^{**}}^{**}).$$

Since γ^{**} is a geodesic,

$$T^{**}(\gamma_{0,z^{**}}^{**}) \leq T^{**}(\gamma_{0,z^{**}}^{**}).$$

Now, let us assume z^{**} is not s_1 . Then, by the definition of s_1 , z^{**} is visited by γ_{u,u_1} after s_1 and thus,

$$T^{**}(\gamma_{0,z^{**}}^{**}) < T(\gamma_{0,z^{**}}^{**}).$$

Combining these three inequalities yields

$$T(\gamma_{0,z^{**}}^{**}) < T(\gamma_{0,z^{**}}^{**}),$$

which is impossible because $\gamma_{0,z^{**}}$ is a geodesic in the environment T . So, the result is proved and the same proof leads to the second part of this lemma. \square

We can now prove the two properties stated at the beginning of this subsection. For item (ii), let $\bar{\gamma}^{**}$ be a geodesic from 0 to x in the environment T^{**} . By Lemma 4.18 and Lemma 4.19, like γ , $\bar{\gamma}^{**}$ visits s_1 and before that $\bar{\gamma}^{**}$ does not visit any edge whose time has been changed when replacing T by T^{**} . Hence,

$$T(\gamma_{0,s_1}) = T^{**}(\gamma_{0,s_1}) \geq T^{**}(\bar{\gamma}_{0,s_1}^{**}) = T(\bar{\gamma}_{0,s_1}^{**}) \geq T(\gamma_{0,s_1}). \quad (4.22)$$

This proves that $T(\gamma_{0,s_1}) = T(\bar{\gamma}_{0,s_1}^{**})$ and $\bar{\gamma}_{0,s_1}^{**}$ is a geodesic in the environment T . Similarly, $\bar{\gamma}_{s_2,x}^{**}$ is a geodesic in the environment T and the path $\bar{\gamma} = \bar{\gamma}_{0,s_1}^{**} \cup \gamma_{s_1,s_2} \cup \bar{\gamma}_{s_2,x}^{**}$ is a geodesic in the environment T that satisfies (ii) in Lemma 4.5 if we prove that it is a self-avoiding path and it is contained in $B_{4,s,N}$. Assume, aiming at a contradiction that $\bar{\gamma}_{0,s_1}^{**}$ visits a vertex of γ_{s_1,s_2} which is not s_1 and denote it by s_3 . Then $\bar{\gamma}_{0,s_1}^{**} \cup \gamma_{s_1,x}$ is an optimal path for the passage time in the environment T since $\bar{\gamma}_{0,s_1}^{**}$ is a geodesic in this environment. It implies that $\bar{\gamma}_{s_3,s_1}^{**} \cup \gamma_{s_1,s_3}$ has at least one edge which is e_1 and that $T(\bar{\gamma}_{s_3,s_1}^{**} \cup \gamma_{s_1,s_3}) = 0$, which is impossible since $T(e_1) > 0$. The same proof gives that $\bar{\gamma}_{s_2,x}^{**}$ does not visit a vertex of γ_{s_1,s_2} and thus $\bar{\gamma}$ is self-avoiding. To get item (ii), it remains to prove that $\bar{\gamma}_{s_1,s_2}^{**}$ is contained in $B_{4,s,N}$. This comes from the fact that, in any environment, the geodesic time between two vertices of $B_{3,s,N}$ is bounded by $2r_3 t_{\max}$, and, since the edges in $B_{4,s,N} \setminus B_{3,s,N}$ have the same times in the environments T or T^{**} , by property (i) of a typical box, the geodesic time to reach a vertex outside $B_{4,s,N}$ and to come back in $B_{3,s,N}$ is bounded from below by $2(r_4 - r_3)(t_{\min} + \delta)$. The condition on r_4 insures that $(r_4 - r_3)(t_{\min} + \delta) > r_3 t_{\max}$.

For item (iii), observe that from (4.22), we also get $T^{**}(\bar{\gamma}_{0,s_1}^{**}) = T^{**}(\gamma_{0,s_1})$ and $T^{**}(\bar{\gamma}_{s_2,x}^{**}) = T^{**}(\gamma_{s_2,x})$. Thus, the path $\gamma_{0,s_1} \cup \bar{\gamma}_{s_1,s_2}^{**} \cup \gamma_{s_2,x}$ is an optimal path for the passage time in the environment T^{**} . If it is not self-avoiding, we get a geodesic that satisfies the requirement of (iii) in Lemma 4.5 by cutting its loops with the same process as in the proof of Lemma 3.5 in the unbounded case. If we denote by s'_1 and s'_2 the two vertices such that $\gamma_{0,s'_1} \cup \bar{\gamma}_{s'_1,s'_2}^{**} \cup \gamma_{s'_2,x}$ is a geodesic in the environment T^{**} obtained by cutting the loops, we have to justify that $\bar{\gamma}_{s'_1,s'_2}^{**}$ is entirely contained in $B_{4,s,N}$. We know that γ_{s_1,s_2} is entirely contained in $B_{4,s,N}$. Let us show that $\gamma_{s'_1,s'_2}$ is also entirely contained in $B_{4,s,N}$, the proof for γ_{s_2,s'_2} is the same. Since $\gamma_{0,s_1} \cup \bar{\gamma}_{s_1,s_2}^{**} \cup \gamma_{s_2,x}$ is an optimal path for the passage time in the environment T^{**} , we have $T^{**}(\bar{\gamma}_{s_1,s'_1}^{**}) + T^{**}(\gamma_{s'_1,s_1}) = 0$ and thus in particular $T^{**}(\gamma_{s'_1,s_1}) = 0$. Since s_1 belongs to $B_{3,s,N}$, if $\gamma_{s'_1,s_1}$ visits a vertex outside $B_{4,s,N}$, there exist two vertices z and z' of $\gamma_{s'_1,s_1}$ such that $z \in \partial B_{3,s,N}$, $z' \in \partial B_{4,s,N}$ and $\gamma_{z,z'}$ except for z is contained in $B_{4,s,N} \setminus B_{3,s,N}$. So $T^{**}(\gamma_{s'_1,s_1}) = 0$ implies that $T^{**}(\gamma_{z,z'}) = 0$ but $T(\gamma_{z,z'}) = T^{**}(\gamma_{z,z'}) = 0$ since the time of the edges of $B_{4,s,N} \setminus B_{3,s,N}$ has not been changed. It makes a contradiction with (4.5) since $r_4 - r_3 \geq 1$ by (4.4).

4.4.6 Every geodesic takes the pattern.

Recall that we assume that

$$\{T \in \mathcal{M}(\ell)\} \cap \{S_\ell^1(T) = s\} \cap \{T' \in \mathcal{B}^*(T)\} \cap \{T'' \in \mathcal{B}^{**}(T, T')\} \text{ occurs.}$$

The aim of this last subsection is to show that every geodesic in the environment T^{**} takes the pattern in $E_P^{**}(T, T')$. The proof is decomposed in two steps. The first step is to show that every geodesic takes an edge of π_{u_2, u_3} and an edge of π_{v_3, v_2} . The second step is to show that every geodesic verifying this property takes the pattern in $E_P^{**}(T, T')$. We begin with a technical lemma.

Lemma 4.20. *For all w vertex of $\pi_{u_2, u_3} \cup \pi_{v_3, v_2}$,*

$$|\mu(u_1 - w) + \mu(v_1 - w) - \mu(u_1 - v_1)| \leq 2C_\mu L_1.$$

Proof. Let w be a vertex of $\pi_{u_2, u_3} \cup \pi_{v_3, v_2}$. Then, by the construction of π , there exists a $\bar{w} \in [u_1, v_1] \subset \mathbb{R}^d$ such that $\|w - \bar{w}\|_1 \leq L_1$. We have

$$\mu(u_1 - v_1) = \mu(u_1 - \bar{w}) + \mu(v_1 - \bar{w}).$$

Then

$$\begin{aligned} |\mu(u_1 - w) - \mu(u_1 - v_1) + \mu(v_1 - w)| &\leq |\mu(u_1 - w) - \mu(u_1 - \bar{w})| + |\mu(v_1 - w) - \mu(v_1 - \bar{w})| \\ &\leq 2C_\mu \|w - \bar{w}\|_1 \\ &\leq 2C_\mu L_1. \end{aligned}$$

□

Lemma 4.21. *Let γ^{**} be a geodesic from 0 to x in the environment T^{**} . Then γ^{**} visits a vertex of π_{u_2, u_3} and one of π_{v_3, v_2} . More precisely, the first vertex of γ^{**} that belongs to $\pi_{u_2, u_3} \cup \pi_{v_3, v_2}$ belongs to π_{u_2, u_3} and the last to belongs to π_{v_3, v_2} .*

Proof. Let γ^{**} be a geodesic from 0 to x in the environment T^{**} . By Lemma 4.17, there exists at least one vertex of $\pi_{u_2, u_3} \cup \pi_{v_3, v_2}$ visited by γ^{**} . Let w be the first vertex of γ^{**} that belongs to $\pi_{u_2, u_3} \cup \pi_{v_3, v_2}$ and assume that w belongs to π_{v_3, v_2} . Note that $\gamma_{0, w}^{**}$ does not visit any other vertex of $\pi_{u_2, u_3} \cup \pi_{v_3, v_2}$. The aim of the proof is to show that

$$T^{**}(\gamma_{0, w}^{\pi}) < T^{**}(\gamma_{0, w}^{**}),$$

which is impossible since γ^{**} is a geodesic in the environment T^{**} . We start with

$$\begin{aligned} T^*(\gamma_{0, w}^{**}) &\geq t^*(0, w) \\ &\geq t^*(0, v_1) - t^*(v_1, w) \\ &= t^*(0, u_1) + t^*(u_1, v_1) - t^*(v_1, w), \end{aligned}$$

since, by Lemma 4.8, γ is a geodesic in the environment T^* . By construction, there is no edge of γ_{u_1, v_1} whose time has been changed at the first modification, thus $t^*(u_1, v_1) = t(u_1, v_1)$. Furthermore, since there is no edge whose time has been increased at the first modification, $t^*(v_1, w) \leq t(v_1, w)$, so

$$T^*(\gamma_{0, w}^{**}) \geq t^*(0, u_1) + t(u_1, v_1) - t(v_1, w).$$

We now want to bound from below $T^{**}(\gamma_{0, w}^{**})$. The only edges whose time has been reduced at the second modification are among those of $\pi_{u_2, u_3} \cup \pi_{v_3, v_2}$ and of $E_{P'}^{**}(T, T')$. So, since $\gamma_{0, w}^{**}$ does not take any edge of $\pi_{u_2, u_3} \cup \pi_{v_3, v_2}$, it can only save time taking edges of $E_{P'}^{**}(T, T')$. So,

$$T^{**}(\gamma_{0, w}^{**}) \geq t^*(0, u_1) + t(u_1, v_1) - t(v_1, w) - T^\Lambda.$$

Then, using the definition of a typical box, Lemma 4.20 and the inequality $t^*(0, u_1) \geq t^*(0, u_2)$ (which comes from the definition of u_2) leads to

$$T^{**}(\gamma_{0, w}^{**}) \geq t^*(0, u_2) + \mu(u_1 - w) - \varepsilon(\mu(u_1 - v_1) + \mu(v_1 - w)) - 2N - 2C_\mu L_1 - T^\Lambda.$$

On the other hand, note that, π being an oriented path, $\|u_1 - w\|_1 \geq \|u_2 - w\|_1$, so, using the knowledge of T^{**} on edges of π ,

$$\begin{aligned} T^{**}(\gamma_{0, w}^{\pi}) &= T^{**}(\gamma_{0, u_2}^{\pi}) + T^{**}(\gamma_{u_2, w}^{\pi}) \\ &\leq t^*(0, u_2) + 2t_{\max} + 2t_{\max}\ell^\Lambda + (t_{\min} + \delta')\|u_1 - w\|_1. \end{aligned}$$

To conclude, let us show that we have the inequality

$$\begin{aligned} t^*(0, u_2) + 2t_{\max} + 2t_{\max}\ell^\Lambda + (t_{\min} + \delta')\|u_1 - w\|_1 \\ < t^*(0, u_2) + \mu(u_1 - w) - \varepsilon(\mu(u_1 - v_1) + \mu(v_1 - w)) - 2N - 2C_\mu L_1 - T^\Lambda. \end{aligned} \quad (4.23)$$

First, combining Lemma 4.9 and Lemma 4.16 leads to

$$\|u_1 - w\|_1 \geq \frac{N\nabla}{3C_\mu}. \quad (4.24)$$

Then, by (1.19), we have

$$\mu(u_1 - w) \geq (t_{\min} + \delta)\|u_1 - w\|_1.$$

Recall that $K' = T^\Lambda + 2(C_\mu L_1 + t_{\max} + t_{\max}\ell^\Lambda)$, it is sufficient to have

$$\delta' < \delta - \varepsilon C_\mu \frac{\|u_1 - v_1\|_1 + \|v_1 - w\|_1}{\|u_1 - w\|_1} - \frac{2N}{\|u_1 - w\|_1} - \frac{K'}{\|u_1 - w\|_1}.$$

Then, by Lemma 4.16, $\frac{\|u_1 - v_1\|_1 + \|v_1 - w\|_1}{\|u_1 - w\|_1} \leq 6$. So, since $\varepsilon < \frac{\delta}{24C_\mu}$, we have

$$\varepsilon C_\mu \frac{\|u_1 - v_1\|_1 + \|v_1 - w\|_1}{\|u_1 - w\|_1} < \frac{\delta}{4}.$$

By (4.24) and since $1 < \frac{\delta \nabla}{24C_\mu}$ (by (4.2)) and $N > \frac{12C_\mu K'}{\delta \nabla}$ (by (4.6)), the condition $\delta' \leq \frac{\delta}{4}$ gives us (4.23). Hence $T^{**}(\gamma_{0,w}^\pi) < T^{**}(\gamma_{0,w}^{**})$. \square

Finally, let us prove the following lemma which completes the proof of Lemma 4.5.

Lemma 4.22. *Any geodesic from 0 to x takes the pattern at the pattern-location $E_P^{**}(T, T')$.*

Proof. Let γ^{**} be a geodesic from 0 to x in the environment T^{**} . By Lemma 4.21, γ^{**} visits a vertex of π_{u_2, u_3} and one of π_{v_3, v_2} . As a consequence, there exist a vertex u_4 of π_{u_2, u_3} and a vertex v_4 of π_{v_3, v_2} such that γ^{**} goes from u_4 to v_4 without taking edges of $\pi_{u_2, u_3} \cup \pi_{v_3, v_2}$. Let us remember that for all edge e of $E_P^{**}(T, T')$, we have $T^{**}(e) \leq \nu$. We prove successive properties.

- *The edges of γ_{u_4, v_4}^{**} which are not in $E_P^{**}(T, T')$ have a passage time greater than or equal to ν .*

Since there is no edge of γ_{u_4, v_4}^{**} in $B^{**}(0, T^*(\gamma_{0, u_1}))$, $B^{**}(x, T^*(\gamma_{v_1, x}))$ or $\pi_{u_2, u_3} \cup \pi_{v_3, v_2}$, it is sufficient to prove that γ_{u_4, v_4}^{**} is entirely contained in $B_{2, s, N}$. By convexity, we have that all points of $[u_1, v_1]$ are contained in $B_\mu(c_0, N\nabla)$, so by Lemma 4.12, $\|u_4 - c_0\|_1 \leq \frac{N\nabla}{c_\mu} + L_1$, and thus,

$$\|u_4 - sN\|_1 \leq N \left(\frac{\nabla}{c_\mu} + r_1 \right) + L_1.$$

So, if γ_{u_4, v_4}^{**} is not entirely contained in $B_{2, s, N}$, the number of edges whose time is greater than or equal to ν that γ_{u_4, v_4}^{**} has to travel to leave $B_{2, s, N}$ is bounded from below by $\left(r_2 - \frac{\nabla}{c_\mu} - r_1 \right) N - 2d\ell^\Lambda - L_1$, and we get

$$T^{**}(\gamma_{u_4, v_4}^{**}) \geq \left(\left(r_2 - \frac{\nabla}{c_\mu} - r_1 \right) N - 2d\ell^\Lambda - L_1 \right) \nu.$$

But

$$T^{**}(\gamma_{u_4, v_4}^\pi) \leq \frac{2N\nabla}{c_\mu} (t_{\min} + \delta') + 2t_{\max}(1 + \ell^\Lambda),$$

and since $N \geq 1$, $\frac{t_{\max}}{\nu} \geq 1$, $\frac{t_{\min} + \delta}{\nu} \leq 1$ and by (4.3),

$$r_2 > r_1 + L_1 + \frac{3\nabla}{c_\mu} + \frac{2t_{\max}}{\nu} (1 + (1+d)\ell^\Lambda),$$

we have $T^{**}(\gamma_{u_4, v_4}^\pi) < T^{**}(\gamma_{u_4, v_4}^{**})$, which is impossible since γ_{u_4, v_4}^{**} is a geodesic in the environment T^{**} .

- *We have that $\|u_4 - u_3\|_1 \leq 4\ell^\Lambda$ and $\|v_4 - v_3\|_1 \leq 4\ell^\Lambda$.*

Assume that it is not the case. Let us show that $T^{**}(\gamma_{u_4, v_4}^{**}) - T^{**}(\pi_{u_4, v_4}) > 0$, which is a contradiction since γ_{u_4, v_4}^{**} is a geodesic in the environment T^{**} . To this aim, let us compare the time that can save each path compared with the other in any direction. Recall the notation introduced in the proof of Lemma 2.1: for $i \in \{1, \dots, d\}$ and a path $\tilde{\pi}$, $T_i^{**}(\tilde{\pi})$ denotes the sum of the passage times of the edges of $\tilde{\pi}$ which are in the direction ε_i . In the direction $\mathcal{O}(T, T')$, since $\|u_4 - u_3\|_1 > 4\ell^\Lambda$ or $\|v_4 - v_3\|_1 > 4\ell^\Lambda$, and by the construction, we have that

$$T_{\mathcal{O}(T, T')}^{**}(\gamma_{u_4, v_4}^{**}) - T_{\mathcal{O}(T, T')}^{**}(\pi_{u_4, v_4}) \geq 4\ell^\Lambda \nu - 4\ell^\Lambda (t_{\min} + \delta') + 2\ell^\Lambda t_{\min} - 2\ell^\Lambda \nu,$$

where the term $2\ell^\Lambda t_{\min} - 2\ell^\Lambda \nu$ comes from the time potentially saved by γ_{u_4, v_4}^{**} by taking edges of $E_P^{**}(T, T')$. Then, in any other direction, γ_{u_4, v_4}^{**} can save a time lower than or equal to $2\ell^\Lambda \delta'$ compared with π_{u_4, v_4} thanks to the edges in $E_P^{**}(T, T')$. Hence,

$$\begin{aligned} T^{**}(\gamma_{u_4, v_4}^{**}) - T^{**}(\pi_{u_4, v_4}) &\geq 2\ell^\Lambda \nu + 2\ell^\Lambda t_{\min} - 4\ell^\Lambda (t_{\min} + \delta') - 2(d-1)\ell^\Lambda \delta' \\ &\geq 2\ell^\Lambda \nu - 2\ell^\Lambda t_{\min} - (4\ell^\Lambda + 2(d-1)\ell^\Lambda)\delta' > 0 \end{aligned}$$

since $t_{\min} + (1+d)\delta' < t_{\min} + \delta \leq \nu$.

- We have that $u_4 = u_3$ and $v_4 = v_3$.

Assume that it is not the case. First, assume that γ_{u_4, v_4}^{**} does not take any edge of $E_P^{**}(T, T')$. According to the first property, $T^{**}(\gamma_{u_4, v_4}^{**}) \geq \|u_4 - v_4\|_1 \nu$. Then, since π is an oriented path,

$$T^{**}(\gamma_{u_4, v_4}^\pi) \leq (\|u_4 - v_4\|_1 - 2\ell^\Lambda)(t_{\min} + \delta') + 2\ell^\Lambda \nu.$$

Since $u_4 \neq u_3$ or $v_4 \neq v_3$, we have that $\|u_4 - v_4\|_1 > 2\ell^\Lambda$ and we obtain

$$T^{**}(\gamma_{u_4, v_4}^\pi) < T^{**}(\gamma_{u_4, v_4}^{**}),$$

which is a contradiction since γ_{u_4, v_4}^{**} is a geodesic in the environment T^{**} . So γ_{u_4, v_4}^{**} takes an edge of $E_P^{**}(T, T')$. Let u_0^{**} be the first entry point of γ_{u_4, v_4}^{**} in $E_P^{**}(T, T')$ and consider the path π^{**} following π_{u_4, u_3} , then going from u_3 to u_0^{**} in one of the shortest way for the norm $\|\cdot\|_1$ and then following $\gamma_{u_0^{**}, v_4}^{**}$. Then the number of edges of $\pi_{u_4, u_0^{**}}^{**}$ is lower than or equal to the number of edges of $\gamma_{u_4, u_0^{**}}^{**}$, for all $e \in \pi_{u_4, u_0^{**}}^{**}$, $T^{**}(e) \leq \nu$, there exists $e' \in \pi_{u_4, u_0^{**}}^{**}$ such that $T^{**}(e') \leq t_{\min} + \delta'$ and for all $e \in \gamma_{u_4, u_0^{**}}^{**}$, $T^{**}(e) \geq \nu$. So we have $T^{**}(\pi^{**}) < T^{**}(\gamma_{u_4, v_4}^{**})$, which is impossible since γ^{**} is a geodesic in the environment T^{**} . The same proof gives $v_4 = v_3$.

- γ^{**} takes the pattern at the pattern-location $E_P^{**}(T, T')$.

Assume that γ_{u_4, v_4}^{**} is not entirely contained in $E_P^{**}(T, T')$. Let v_0^{**} be the first exit point from $E_P^{**}(T, T')$ of γ_{u_4, v_4}^{**} and u_1^{**} the first entry point after v_0^{**} . Let us consider the shortcut π^{**} going from v_0^{**} to u_1^{**} in one of the shortest way for the norm $\|\cdot\|_1$. Then let $v_{0,+}^{**}$ denote the first vertex visited by γ^{**} after v_0^{**} , then

$$\|u_1^{**} - v_{0,+}^{**}\|_1 = \|u_1^{**} - v_0^{**}\|_1 + 1.$$

Indeed, we have that $\|u_1^{**} - v_{0,+}^{**}\|_1 - \|u_1^{**} - v_0^{**}\|_1$ is equal to 1 or -1 , and if it is equal to -1 , it implies that $v_{0,+}^{**}$ is in $E_P^{**}(T, T')$, which is impossible. So, $\gamma_{v_0^{**}, u_1^{**}}^{**}$ has strictly more edges than π^{**} . Furthermore, all edges of $\gamma_{v_0^{**}, u_1^{**}}^{**}$ have a time greater than or equal to ν although all edges of π^{**} have a time lower than or equal to ν . So, $T^{**}(\pi^{**}) < T^{**}(\gamma_{v_0^{**}, u_1^{**}}^{**})$, which is a contradiction since $\gamma_{v_0^{**}, u_1^{**}}^{**}$ is a geodesic in the environment T^{**} .

Thus γ_{u_4, v_4}^{**} is a path entirely contained in $E_P^{**}(T, T')$, going from u_3 to v_3 and with an optimal time. So, we have the result. □

A Construction of the path π for the modification in the unbounded case

Recall that, here, u^Λ and v^Λ are the vertices defined at the beginning of Section 3.3. The aim is to prove that we can construct a path π in a deterministic way such that:

- π goes from u to u^Λ without visiting a vertex of $B_\infty(sN, \ell^\Lambda)$, then going from u^Λ to v^Λ in a shortest way for the norm $\|\cdot\|_1$ (and thus being contained in $B_\infty(sN, \ell^\Lambda)$) and then goes from v^Λ to v without visiting a vertex of $B_\infty(sN, \ell^\Lambda)$,




Figure 5: Example of the construction of the portion of π which is contained in $B_\infty(sN, \ell^\Lambda + 3) \setminus B_\infty(sN, \ell^\Lambda)$ in dimension 2. On the left, this is an example of the case where $\|u_0 - v^\Lambda\|_1 > 3$ or $\|v_0 - u^\Lambda\|_1 > 3$ and on the right the other case.

- (ii) π is entirely contained in $B_{2,s,N}$ and does not have vertices on the boundary of $B_{2,s,N}$ except u and v ,
- (iii) π is self-avoiding,
- (iv) the length of $\pi_{u,u^\Lambda} \cup \pi_{v^\Lambda,v}$ is bounded from above by $2r_2N + K$, where K is the number of edges in $B_\infty(0, \ell^\Lambda + 3)$.

As it is said in Section 3.3, we want to construct a path from u to sN and a path from sN to v which have no vertex in common except sN and such that their lengths are bounded from above by r_2N . To get them, we use the following lemma whose proof is left to the reader.

Lemma A.1. *Let $m \in \mathbb{N}^*$ and x, y two vertices of \mathbb{Z}^d such that $\|x\|_1 = \|y\|_1 = m$ and $x \neq y$. Then we can build in a deterministic way two paths π_x and π_y linking respectively x and y to 0 and such that their length is equal to m , they have only 0 as a common vertex and they have respectively only x and y as vertices whose norm $\|\cdot\|_1$ is greater than or equal to m .*

Using this lemma and replacing 0 by sN using a translation, we get two paths linking u to sN and v to sN with the stated properties. Recall that $B_\infty(sN, \ell^\Lambda + 3) \subset B_{2,s,N}$ and let u_0 (resp. v_0) denote the first vertex in $B_\infty(sN, \ell^\Lambda + 3)$ visited by the path going from u to sN (resp. the one going from v to sN). Then we get two paths π_{u,u_0} and $\pi_{v_0,v}$ respectively from u to u_0 and from v_0 to v both constructed in a deterministic way such that π_{u,u_0} and $\pi_{v_0,v}$ do not have any vertex in common, are entirely contained in $B_{2,s,N}$, have only u or v as points on the boundary of $B_{2,s,N}$, and their lengths are bounded from above by $r_2N - (\ell^\Lambda + 3)$.

Then we build two paths π_{u_0,u^Λ} and π_{v^Λ,v_0} respectively from u_0 to u^Λ and from v^Λ to v_0 contained in $B_\infty(sN, \ell^\Lambda + 3) \setminus B_\infty(sN, \ell^\Lambda)$ except for u^Λ and v^Λ , such that they do not have any vertex in common. By the definition of K , this implies that the sum of their lengths is bounded from above by K (recall that K is the number of edges in $B_\infty(0, \ell^\Lambda + 3)$). To get these paths, assume first that $\|u_0 - v^\Lambda\|_1 > 3$ or $\|v_0 - u^\Lambda\|_1 > 3$. Assume that we have $\|v_0 - u^\Lambda\|_1 > 3$, the other case being the same. We begin by considering a path $\tilde{\pi}$ going in a shortest way from v_0 to a vertex of the boundary of $B_\infty(sN, \ell^\Lambda + 2)$, denoted by v_0' . This path has at most d edges. Then, let $u^{\Lambda,0}$ be a vertex on the boundary of $B_\infty(sN, \ell^\Lambda + 3)$ such

that $\|u^\Lambda - u^{\Lambda,0}\|_1 = 3$. We get π_{u_0, u^Λ} by going from u_0 to $u^{\Lambda,0}$ in a shortest way on the boundary of $B_\infty(sN, \ell^\Lambda + 3)$ avoiding all vertices of $\tilde{\pi}$, and then by going from $u^{\Lambda,0}$ to u^Λ by three steps. This is possible since $\|v_0 - u^\Lambda\|_1 > 3$. To get π_{v_0, v^Λ} , we begin by following $\tilde{\pi}$. Let $v^{\Lambda,0}$ be a vertex on the boundary of $B_\infty(sN, \ell^\Lambda + 2)$ such that $\|v^\Lambda - v^{\Lambda,0}\|_1 = 2$. Then, π_{v_0, v^Λ} goes from v_0 to $v^{\Lambda,0}$ in a shortest way on the boundary of $B_\infty(sN, \ell^\Lambda + 2)$ avoiding the unique vertex of π_{u_0, u^Λ} belonging to $B_\infty(sN, \ell^\Lambda + 2)$. Finally, π_{v_0, v^Λ} goes from $v^{\Lambda,0}$ to v^Λ by two steps (see the left side of Figure 5 for an example of this construction in dimension 2).

Now, if $\|u_0 - v^\Lambda\|_1 = 3$ and $\|v_0 - u^\Lambda\|_1 = 3$, the construction has to be slightly different. From u_0 , π_{u_0, u^Λ} goes to a vertex u_0'' belonging to the boundary of $B_\infty(sN, \ell^\Lambda + 3)$ and such that $\|u_0'' - v_0\|_1 = 1$ in a shortest way on the boundary of $B_\infty(sN, \ell^\Lambda + 3)$. After, it makes one step to go to the boundary of $B_\infty(sN, \ell^\Lambda + 2)$ and goes to u^Λ in a shortest way by taking only one vertex in $B_\infty(sN, \ell^\Lambda + 1)$. Then, π_{v_0, v^Λ} goes on the boundary of $B_\infty(sN, \ell^\Lambda + 3)$ to a vertex v_0'' belonging to the boundary of $B_\infty(sN, \ell^\Lambda + 3)$ and such that $\|v_0'' - u_0\|_1 = 1$ by avoiding every vertex which belongs to π_{u_0, u^Λ} , and then makes two steps to go to the boundary of $B_\infty(sN, \ell^\Lambda + 1)$ and goes to v^Λ in a shortest way (see the right side of Figure 5 for an example of this construction in dimension 2). In this case, we also have that the sum of their lengths is bounded from above by K .

Finally, π is the path obtained by concatenating π_{u, u_0} , π_{u_0, u^Λ} , a path going from u^Λ to v^Λ in a shortest way for the norm $\|\cdot\|_1$, π_{v^Λ, v_0} and $\pi_{v_0, v}$ in this order. We have that π is a self-avoiding path contained in $B_{2, s, N}$ and has only u and v on the boundary of $B_{2, s, N}$.

B Overlapping pattern

Proof of Lemma 4.1. Recall that ν is fixed at the beginning of Section 4 and that $\delta' = \min\left(\frac{\delta}{8}, \frac{\delta}{1+d}\right)$.

Then, we fix a positive real ν_0 such that:

- $t_{\min} + \delta \leq \nu_0 \leq \nu \leq t_{\max}$,
- the event $\mathcal{A}^\Lambda \cap \{\forall e \in \Lambda, t_e \leq \nu_0\}$ has a positive probability,
- $F([\nu_0, \nu]) > 0$.

Notice that, if F has an atom, one could have $\nu_0 = \nu$, or even $\nu_0 = \nu = t_{\max}$. Then, fix

$$\ell_1 > \frac{2\ell^\Lambda(1+d)(\nu_0 - t_{\min})}{\nu_0 - t_{\min} - \delta'}, \quad (\text{B.1})$$

$$\text{and } \ell_0 > \frac{(\ell^\Lambda + 1)(\nu_0 - t_{\min}) + 2(2\ell_1 + \ell^\Lambda)(t_{\min} + \delta') + 2\ell^\Lambda\nu_0}{\nu_0 - t_{\min} - 2\delta'}. \quad (\text{B.2})$$

Let $j \in \{1, \dots, d\}$. Let us construct the overlapping pattern in $\Lambda_0 = \{-\ell_0, \dots, \ell_0\}^d$ with endpoints $\ell_0\varepsilon_j$ and $-\ell_0\varepsilon_j$. We denote u_3 and v_3 the endpoints of the original pattern, which are respectively $\ell^\Lambda\varepsilon_1$ and $-\ell^\Lambda\varepsilon_1$.

The first step is the construction of a path $\tilde{\pi}$ between $\ell_0\varepsilon_j$ and $-\ell_0\varepsilon_j$. From $\ell_0\varepsilon_j$, the path goes to $(\ell_0 - 1)\varepsilon_j$ by one step in the j -th direction, then goes to $u_1 = (\ell^\Lambda + \ell_1)\varepsilon_1 + (\ell_0 - 1)\varepsilon_j$ by $(\ell^\Lambda + \ell_1)$ steps in the first direction, then to $u_2 = (\ell^\Lambda + \ell_1)\varepsilon_1$ by $(\ell_0 - 1)$ steps in the j -th direction and finally to u_3 by ℓ_1 steps in the first direction. Then, it goes from u_3 to v_3 by $2\ell^\Lambda$ steps in the direction ε_1 and we get $\tilde{\pi}_{v_3, -\ell^\Lambda\varepsilon_j}$ by the same construction as $\tilde{\pi}_{\ell^\Lambda\varepsilon_j, u_3}$. Set $v_1 = -(\ell^\Lambda + \ell_1)\varepsilon_1 - (\ell_0 - 1)\varepsilon_j$ and $v_2 = -(\ell^\Lambda + \ell_1)\varepsilon_1$ (see Figure 6 for an example of the construction of $\tilde{\pi}$).

The event (whose probability is positive) of the overlapping pattern, denoted by $\mathcal{A}_j^{\Lambda_0}$ is the following:

- for all $e \in \tilde{\pi}_{\ell_0\varepsilon_j, u_3}$ and $e \in \tilde{\pi}_{v_3, -\ell_0\varepsilon_j}$, $T(e) < t_{\min} + \delta'$,
- the event $\mathcal{A}^\Lambda \cap \{\forall e \in \Lambda, T(e) \leq \nu_0\}$ is realized,
- for all e in $\Lambda_0 \setminus (\tilde{\pi} \cup \{-\ell^\Lambda, \dots, \ell^\Lambda\}^d)$, $\nu_0 \leq T(e) \leq \nu$.




Figure 6: Example of the construction of $\tilde{\pi}$ in an overlapping pattern in dimension 2.

On this event, let $\bar{\gamma}$ be one of the fastest path from $\ell_0\varepsilon_j$ to $-\ell_0\varepsilon_j$ among the path entirely contained in Λ_0 and let us show that $\bar{\gamma}$ visits u^Λ and v^Λ and that $\bar{\gamma}_{u^\Lambda, v^\Lambda}$ is entirely contained in $\Lambda = \{-\ell^\Lambda, \dots, \ell^\Lambda\}^d$. We proceed by proving successive properties.

- There exist one vertex a_0 of $\tilde{\pi}_{u_1, u_2}$ and one vertex b_0 of $\tilde{\pi}_{v_2, v_1}$ visited by $\bar{\gamma}$. Further, $\bar{\gamma}_{\ell_0\varepsilon_j, a_0} = \tilde{\pi}_{\ell_0\varepsilon_j, a_0}$ and $\bar{\gamma}_{b_0, -\ell_0\varepsilon_j} = \tilde{\pi}_{b_0, -\ell_0\varepsilon_j}$.

Let us assume that $\bar{\gamma}$ does not visit any vertex of $\tilde{\pi}_{u_1, u_2}$, the other case being the same. The path $\bar{\gamma}$ has to take at least $\ell_0 - 1 - \ell^\Lambda$ edges connecting vertices such that at least one of them has its j -th coordinate strictly between ℓ^Λ and $\ell_0 - 1$. The only edges in this set whose passage time is smaller than ν_0 are those of $\tilde{\pi}_{u_1, u_2}$. Hence

$$T(\bar{\gamma}) \geq (\ell_0 - 1 - \ell^\Lambda)\nu_0 + (\ell_0 + \ell^\Lambda + 1)t_{\min}.$$

But, thanks to our construction, we have

$$T(\tilde{\pi}) \leq (2\ell_0 + 2(2\ell_1 + \ell^\Lambda))(t_{\min} + \delta') + 2\ell^\Lambda\nu_0.$$

So, (B.2) leads to $T(\tilde{\pi}) < T(\bar{\gamma})$, which is impossible.

The fact that $\bar{\gamma}_{\ell_0\varepsilon_j, a_0} = \tilde{\pi}_{\ell_0\varepsilon_j, a_0}$ follows from the fact that $\tilde{\pi}_{\ell_0\varepsilon_j, a_0}$ is an oriented path and from the fact that $t_{\min} + \delta' < \nu_0$.

- If a_1 (resp. b_1) is a vertex of $\tilde{\pi}_{u_2, u_3}$ (resp. $\tilde{\pi}_{v_3, v_2}$) visited by $\bar{\gamma}$, then $\bar{\gamma}_{\ell_0\varepsilon_j, a_1} = \tilde{\pi}_{\ell_0\varepsilon_j, a_1}$ (resp. $\bar{\gamma}_{b_1, -\ell_0\varepsilon_j} = \tilde{\pi}_{b_1, -\ell_0\varepsilon_j}$).

Indeed, let a_1 be such a vertex and let a_0 be the vertex of the preceding property. We only have to prove that $\bar{\gamma}_{a_0, a_1} = \tilde{\pi}_{a_0, a_1}$ and it also follows from the fact that $\tilde{\pi}_{a_0, a_1}$ is an oriented path.

Among the vertices visited by $\bar{\gamma}$, we denote by a (resp. b) the last vertex of $\tilde{\pi}_{\ell_0\varepsilon_j, u_3}$ (resp. the first vertex of $\tilde{\pi}_{v_3, -\ell_0\varepsilon_j}$). Then, $\bar{\gamma}_{a, b}$ does not visit any vertex of $\tilde{\pi}_{\ell_0\varepsilon_j, u_3} \cup \tilde{\pi}_{v_3, -\ell_0\varepsilon_j}$ (except a and b). Indeed, by the two properties above, we have that $\bar{\gamma}_{\ell_0\varepsilon_j, a} = \tilde{\pi}_{\ell_0\varepsilon_j, a}$ and $\bar{\gamma}_{b, -\ell_0\varepsilon_j} = \tilde{\pi}_{b, -\ell_0\varepsilon_j}$. Even if $t_{\min} = 0$

and $\bar{\gamma}$ is not self-avoiding, since $\nu_0 > 0$ and since the edges going to a vertex of $\tilde{\pi}_{\ell_0 \varepsilon_j, a} \cup \tilde{\pi}_{b, -\ell_0 \varepsilon_j}$ whose time is smaller than $t_{\min} + \delta'$ are already visited by $\bar{\gamma} \setminus \bar{\gamma}_{a,b}$, it is not possible that $\bar{\gamma}_{a,b}$ visits a vertex of $\tilde{\pi}_{\ell_0 \varepsilon_j, a} \cup \tilde{\pi}_{b, -\ell_0 \varepsilon_j}$ (except a and b). Then, $\bar{\gamma}_{a,b}$ can not visit a vertex of $\tilde{\pi}_{a, u_3} \cup \tilde{\pi}_{v_3, b}$ thanks to the definition of a and b .

- The vertex a (resp. b) belongs to $\tilde{\pi}_{u_2, u_3}$ (resp. $\tilde{\pi}_{v_3, v_2}$).

Assume that a or b does not satisfy this property. Then $\|a - u_3\|_1 \geq \ell_1$ or $\|b - v_3\|_1 \geq \ell_1$. Since outside the pattern, $\tilde{\pi}_{a,b}$ is oriented and since $\|u_3 - v_3\|_1 \leq 2\ell^\Lambda$,

$$T(\tilde{\pi}_{a,b}) \leq (\|a - b\|_1 - 2\ell^\Lambda)(t_{\min} + \delta') + 2\ell^\Lambda \nu_0.$$

Then, since the edges whose time is smaller than ν_0 taken by $\bar{\gamma}_{a,b}$ are those in Λ ,

$$T(\bar{\gamma}_{a,b}) \geq (\|a - b\|_1 - 2d\ell^\Lambda)\nu_0 + 2d\ell^\Lambda t_{\min}.$$

Using (B.1), we get $T(\bar{\gamma}_{a,b}) > T(\tilde{\pi}_{a,b})$, which is impossible.

- We have that $a = u_3$ and $b = v_3$.

We have

$$T(\tilde{\pi}_{a,b}) \leq (\|a - b\|_1 - 2\ell^\Lambda)(t_{\min} + \delta') + 2\ell^\Lambda \nu_0,$$

and thus $\bar{\gamma}_{a,b}$ takes at least one edge of Λ otherwise $T(\bar{\gamma}_{a,b}) \geq (\|a - b\|_1 + 2\ell^\Lambda)\nu_0 > T(\tilde{\pi}_{a,b})$ since $\ell^\Lambda > 0$. Now, assume that $a \neq u_3$, the other case being the same. Let u_0 be the first entry point of $\bar{\gamma}_{a,b}$ in Λ and let consider the path π following $\tilde{\pi}_{a, u_3}$, then going from u_3 to u_0 in a shortest way and then following $\bar{\gamma}_{u_0, b}$. Then, the number of edges of π_{a, u_0} is lower than or equal to the number of edges of $\bar{\gamma}_{a, u_0}$, for all $e \in \pi_{a, u_0}$, $T(e) \leq \nu_0$, there exists $e' \in \pi_{a, u_0}$ such that $T(e') \leq t_{\min} + \delta'$ and for all $e \in \bar{\gamma}_{a, u_0}$, $T(e) \geq \nu_0$. So, we have $T(\pi) < T(\bar{\gamma}_{a,b})$ which is impossible since $\bar{\gamma}$ is an optimal path among paths entirely contained in Λ_0 .

- $\bar{\gamma}_{a,b}$ takes the original pattern.

Assume that $\bar{\gamma}_{a,b}$ is not entirely contained in Λ . Let v_0 be the first exit point from Λ of $\bar{\gamma}_{a,b}$ and u_0 the first entry point after v_0 . Let us consider the shortcut π going from v_0 to u_0 in a shortest way. Then, using the same argument as in the proof of Lemma 4.22, we have that $\bar{\gamma}_{v_0, u_0}$ has strictly more edges than π . Furthermore, all edges of $\bar{\gamma}_{v_0, u_0}$ have a time greater than or equal to ν_0 although all edges of π have a time lower than or equal to ν_0 . So, $T(\pi) < T(\bar{\gamma}_{v_0, u_0})$, which is impossible since $\bar{\gamma}$ is an optimal path among paths entirely contained in Λ_0 . Since $\bar{\gamma}_{a,b}$ is a path entirely contained in Λ , going from u_3 to v_3 and with an optimal time, so we have the result. □

References

- [1] Enrique D. Andjel and Maria E. Vares. First passage percolation and escape strategies. *Random Structures and Algorithms*, 47(3):414–423, October 2015.
- [2] A. Auffinger, M. Damron, and J. Hanson. *50 years of first-passage percolation*. American Mathematical Society, 2017.
- [3] S. Boucheron, G. Lugosi, and P. Massart. *Concentration Inequalities: A Nonasymptotic Theory of Independence*. OUP Oxford, 2013.
- [4] Geoffrey Grimmett. *Percolation*, volume 321 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, second edition, 1999.
- [5] Arjun Krishnan, Firas Rassoul-Agha, and Timo Seppäläinen. Geodesic length and shifted weights in first-passage percolation. *arXiv preprint arXiv:2101.12324*, 2021.

- [6] T. M. Liggett, R. H. Schonmann, and A. M. Stacey. Domination by product measures. *The Annals of Probability*, 25(1):71 – 95, 1997.
- [7] Shuta Nakajima. On properties of optimal paths in first-passage percolation. *Journal of Statistical Physics*, 174(2):259–275, Oct 2018.
- [8] J. van den Berg and H. Kesten. Inequalities for the Time Constant in First-Passage Percolation. *The Annals of Applied Probability*, 3(1):56 – 80, 1993.