



HAL
open science

A note on adiabatic time evolution and quasi-static processes in translation-invariant quantum systems

Vojkan Jakšić, Claude-Alain Pillet, Clément Tauber

► To cite this version:

Vojkan Jakšić, Claude-Alain Pillet, Clément Tauber. A note on adiabatic time evolution and quasi-static processes in translation-invariant quantum systems. *Annales Henri Poincaré*, 2023, 10.1007/s00023-023-01282-5 . hal-03633713

HAL Id: hal-03633713

<https://hal.science/hal-03633713>

Submitted on 2 Feb 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

A note on adiabatic time evolution and quasi-static processes in translation-invariant quantum systems

Vojkan Jakšić¹, Claude-Alain Pillet^{*2}, and Clément Tauber³

¹Department of Mathematics and Statistics, McGill University, 805 Sherbrooke Street West, Montreal, QC, H3A 2K6, Canada

²Aix Marseille Univ, Université de Toulon, CNRS, CPT, Marseille, France

³Institut de Recherche Mathématique Avancé, UMR 7501 Université de Strasbourg et CNRS, 7 rue René-Descartes, 67000 Strasbourg, France

Dedicated to the memory of Krzysztof Gawędzki

Abstract. We study the slowly varying, non-autonomous quantum dynamics of a translation invariant spin or fermion system on the lattice \mathbb{Z}^d . This system is assumed to be initially in thermal equilibrium, and we consider realizations of quasi-static processes in the adiabatic limit. By combining the Gibbs variational principle with the notion of quantum weak Gibbs states introduced in [JPT23], we establish a number of general structural results regarding such realizations. In particular, we show that such a quasi-static process is incompatible with the property of approach to equilibrium studied in this previous work.

1 Introduction

This paper is a direct continuation of [JPT23]. Although its topic and implications are different, we will rely heavily on the technical tools and conceptual framework introduced in this earlier work. In particular, in what follows and without further saying, we will use terminology, notation, and results stated in the introductory section of [JPT23].¹

The set of quantum weak Gibbs states $\text{WG}(\Phi)$ for the spin/fermion interaction $\Phi \in \mathcal{B}^r$ is introduced in Section 2.2 of [JPT23]. An immediate consequence of our definition of weak Gibbs states is the existence of the specific relative entropy and the validity of the formula

$$s(\nu|\omega) := \lim_{\Lambda \uparrow \mathbb{Z}^d} \frac{S(\nu_\Lambda|\omega_\Lambda)}{|\Lambda|} = -s(\nu) + \nu(E_\Phi) + P(\Phi), \quad (1)$$

for any $\nu \in \mathcal{A}(\mathfrak{A})$ and $\omega \in \text{WG}(\Phi)$. The current technology allowed us to prove the identification $\mathcal{S}_{\text{eq}}(\Phi) = \text{WG}(\Phi)$ in dimension $d = 1$ for finite range interactions, and for any d and $\Phi \in \mathcal{B}^r$ in the high temperature regime. Since this identification is expected to hold much more generally, we have introduced the notion of *regular pair* $(\omega, \Phi) \in \mathcal{A}(\mathfrak{A}) \times \mathcal{B}^r$ for which (1) holds for all $\nu \in \mathcal{A}(\mathfrak{A})$. This notion, combined with the Gibbs variational principle, has led to several

¹The general references in [JPT23] related to the mathematical theory of algebraic quantum statistical mechanics apply to this work as well.

structural results about the fundamental problem of Approach to Equilibrium—the zeroth law of thermodynamics—in quantum statistical mechanics.

In this work, starting with the same general ingredients, we study whether quasi-static transitions of extended spin or fermion systems can be described by slowly varying, non-autonomous translation invariant interactions. The emerging structural theory in particular yields that a realization of such a quasi-static transition is possible only if the specific entropy is constant along the state trajectory. Combined with the results of [JPT23], this gives that, in the adiabatic limit, quasi-static transitions are incompatible with the approach to equilibrium in the translation invariant setting of algebraic quantum statistical mechanics.

The paper is organized as follows. In Sections 2 and 3 we briefly review the basic adiabatic theorems of quantum mechanics and quantum statistical mechanics. For a more complete review, historical perspective, and additional references, we refer the reader to [BFJP].² We introduce our translation invariant framework in Section 4.1. Our main results are stated and proved in Section 4.2. They are compared with the main result of [JPT23] in Section 4.3. Similar to [JPT23], most of our proofs are technically simple. The proofs of two technically more involved results, Propositions 3.2 and 4.1, are postponed to Section 5. For the reader’s convenience, Appendix A collects the notations inherited from [JPT23].

Acknowledgments This work was supported by the French *Agence Nationale de la Recherche*, grant NONSTOPS (ANR-17-CE40-0006-01, ANR-17-CE40-0006-02, ANR-17-CE40-0006-03) and the CY Initiative of Excellence through the grant Investissements d’Avenir ANR-16-IDEX-0008. It was partly developed during VJ’s stay at the CY Advanced Studies, whose support is gratefully acknowledged. VJ also acknowledges the support of NSERC. The authors wish to thank Martin Fraas for useful discussions.

2 Adiabatic theorems in quantum mechanics

The adiabatic theorem with a gap. The formulation and the proof of the adiabatic theorem in quantum mechanics go back to the seminal work of Born and Fock [BF28]. The first modern proof, in functional analytic framework, is due to Kato [Kat50]. The method of the proof introduced there played an important role in all the future developments of the subject. For refinements of Kato’s result under the gap assumption made in [Kat50], see [ASY87, JP91, Nen93].

Let H be a (possibly unbounded) self-adjoint operator on the Hilbert space \mathcal{H} and

$$V : [0, 1] \rightarrow \mathcal{B}(\mathcal{H}) \tag{2}$$

a continuous map taking its values in the bounded self-adjoint operators on \mathcal{H} . For $T > 0$, we consider the non-autonomous Schrödinger equation

$$i\partial_t \psi(t) = (H + V(t/T))\psi(t), \quad \psi(s) = f, \quad s, t \in [0, T].$$

For any $f \in \text{Dom}(H)$ this equation has a unique solution $[0, T] \ni t \mapsto \psi(t) \in \text{Dom}(H)$, which can be written as $\psi(t) = U_T^{s/T \rightarrow t/T} f$, where $[0, 1] \times [0, 1] \ni (\sigma, \tau) \mapsto U_T^{\sigma \rightarrow \tau} \in \mathcal{B}(\mathcal{H})$ is a strongly continuous map taking its values in the unitary operators on \mathcal{H} and satisfying $U_T^{\sigma \rightarrow \sigma} = I$, $U_T^{\sigma \rightarrow \tau} \text{Dom}(H) \subset \text{Dom}(H)$, and $U_T^{\tau \rightarrow \nu} U_T^{\sigma \rightarrow \tau} = U_T^{\sigma \rightarrow \nu}$ for all $\sigma, \tau, \nu \in [0, 1]$. U_T is the propagator associated to the time-dependent Hamiltonian $[0, 1] \ni \tau \mapsto T(H + V(\tau))$.

²In particular, we have not attempted here to formulate these results in a technically optimal setting.

For each $\tau \in [0, 1]$, let $E(\tau)$ be an eigenvalue of finite multiplicity of $H(\tau) = H + V(\tau)$ which is uniformly separated from the remaining part of its spectrum, i.e.,

$$\inf_{\tau \in [0,1]} \text{dist}(E(\tau), \text{sp}(H(\tau)) \setminus \{E(\tau)\}) > 0, \quad (3)$$

and denote by $P(\tau)$ the orthogonal projection onto the associated eigenspace.

Theorem 2.1. [Kat50] *Suppose, in addition to (3), that the map (2) is C^2 . Then, as $T \uparrow \infty$,*

$$\sup_{\tau \in [0,1]} \|(I - P(\tau))U_T^{0-\tau}P(0)\| = O(T^{-1}).$$

The adiabatic theorem without a gap. This fundamental refinement of Theorem 2.1 goes back to Avron and Elgart [AE99], who dispense with the gap assumption (3); for an important technical comment on their result see [Teu01]. The setting is the same as in Theorem 2.1, except that (3) is replaced by the following assumption:

Assumption (AD) There exists a C^2 map $[0, 1] \ni \tau \mapsto P(\tau) \in \mathcal{B}(\mathcal{H})$ such that, for Lebesgue a.e. $\tau \in [0, 1]$, $P(\tau)$ is the orthogonal projection onto the eigenspace of a finite multiplicity eigenvalue of $H(\tau)$.

Theorem 2.2. [AE99, Teu01] *Suppose that (AD) holds. Then*

$$\lim_{T \rightarrow \infty} \sup_{\tau \in [0,1]} \|(I - P(\tau))U_T^{0-\tau}P(0)\| = 0.$$

3 Adiabatic theorems in quantum statistical mechanics

The isothermal adiabatic theorem for local perturbations. Let (\mathcal{O}, α) be a C^* -dynamical system. Denote by δ its generator, $\alpha^t = e^{t\delta}$, and by ω an α -KMS state at inverse temperature $\beta > 0$. To a self-adjoint $V \in \mathcal{O}$ we associate the perturbed dynamics $\alpha_V^t = e^{t\delta_V}$, where $\delta_V = \delta + i[V, \cdot]$, and the perturbed (α_V, β) -KMS state ω_V .

Let

$$[0, 1] \ni \tau \mapsto V(\tau) \in \mathcal{O} \quad (4)$$

be a continuous map such that $V(\tau)$ is self-adjoint for all τ . For $T > 0$, the unique solution of the Cauchy problem for the non-autonomous Heisenberg equation

$$\partial_t \gamma^t(A) = \gamma^t \circ \delta_{V(t/T)}(A), \quad \gamma^s(A) = A \in \text{Dom}(\delta), \quad s, t \in [0, T],$$

is given by $\gamma^t = \alpha_T^{s/T \rightarrow t/T}$, where $[0, 1] \times [0, 1] \ni (\sigma, \tau) \mapsto \alpha_T^{\sigma \rightarrow \tau}$ is a strongly continuous two-parameter family of $*$ -automorphisms of \mathcal{O} satisfying the relation $\alpha_T^{\sigma \rightarrow \tau} \circ \alpha_T^{\tau \rightarrow \nu} = \alpha_T^{\sigma \rightarrow \nu}$ for any $\sigma, \tau, \nu \in [0, 1]$. For $0 \leq \sigma \leq \tau \leq 1$, it has the norm-convergent expansion³

$$\alpha_T^{\sigma \rightarrow \tau}(A) = \sum_{n=0}^{\infty} T^n \int_{\sigma \leq \sigma_1 \leq \dots \leq \sigma_n \leq \tau} i[\alpha^{(\sigma_1 - \sigma)T}(V(\sigma_1)), i[\dots, i[\alpha^{(\sigma_n - \sigma)T}(V(\sigma_n)), \alpha^{(\tau - \sigma)T}(A)] \dots]] d\sigma_1 \dots d\sigma_n.$$

Recall the property of Return to Equilibrium reviewed in [JPT23, Section 1.1], where the reader can also find references to this well-studied topic.

³It is understood that the zeroth term in this expansion is $\alpha^{(\tau - \sigma)T}(A)$. A similar expression holds for $0 \leq \tau \leq \sigma \leq 1$.

Theorem 3.1. *Suppose that the map (4) is C^2 . If, for Lebesgue a.e. $\tau \in [0, 1]$, the quantum dynamical system $(\mathcal{C}, \alpha_{V(\tau)}, \omega_{V(\tau)})$ has the property of return to equilibrium, then*

$$\lim_{T \rightarrow \infty} \sup_{\tau \in [0, 1]} \|\omega_{V(0)} \circ \alpha_T^{0 \rightarrow \tau} - \omega_{V(\tau)}\| = 0. \quad (5)$$

This result goes back to [ASF07, JP]; see also [JP14]. The proof is a simple combination of the Avron–Elgart gapless adiabatic theorem 2.2 and Araki’s perturbation theory of the modular structure.

For an in depth discussion of Theorem 3.1 and its thermodynamical implications we refer the reader to [BFJP]. Here we recall three results that will be of relevance in what follows; see also [JP14, Section 5] for a related discussion.

The first is the following entropy balance equation:

Proposition 3.2. *Suppose that the map (4) is C^1 and denote by \dot{V} its derivative. Then⁴*

$$S\left(\omega_{V(0)} \circ \alpha_T^{0 \rightarrow \tau} \middle| \omega_{V(\tau)}\right) = \beta \int_0^\tau (\omega_{V(0)} \circ \alpha_T^{0 \rightarrow \sigma} - \omega_{V(\sigma)}) (\dot{V}(\sigma)) d\sigma$$

holds for all $\tau \in [0, 1]$.

For the reader’s convenience, a proof of this proposition is given in Section 5.1.

The entropy balance equation yields the estimate

$$\sup_{\tau \in [0, 1]} S\left(\omega_{V(0)} \circ \alpha_T^{0 \rightarrow \tau} \middle| \omega_{V(\tau)}\right) \leq \beta \sup_{\tau \in [0, 1]} \|\omega_{V(0)} \circ \alpha_T^{0 \rightarrow \tau} - \omega_{V(\tau)}\| \sup_{\tau \in [0, 1]} \|\dot{V}(\tau)\|,$$

and the validity of (5) implies that the adiabatic theorem for local perturbations also holds in the entropic sense:

$$\lim_{T \rightarrow \infty} \sup_{\tau \in [0, 1]} S\left(\omega_{V(0)} \circ \alpha_T^{0 \rightarrow \tau} \middle| \omega_{V(\tau)}\right) = 0. \quad (6)$$

On the other hand, the Pinsker–Csiszàr inequality [OP93, Proposition 5.23], $\|\omega - \nu\|^2 \leq 2S(\omega|\nu)$, gives that (6) implies (5). In summary:

Theorem 3.3. *Suppose that the map (4) is C^1 . Then, the following statements are equivalent:*

(i)

$$\lim_{T \rightarrow \infty} \sup_{\tau \in [0, 1]} S\left(\omega_{V(0)} \circ \alpha_T^{0 \rightarrow \tau} \middle| \omega_{V(\tau)}\right) = 0.$$

(ii)

$$\lim_{T \rightarrow \infty} \sup_{\tau \in [0, 1]} \|\omega_{V(0)} \circ \alpha_T^{0 \rightarrow \tau} - \omega_{V(\tau)}\| = 0.$$

Starting with Proposition 3.2, a similar argument yields the third result which is also of independent interest.

Theorem 3.4. *Suppose that the map (4) is C^1 . Then, the following statements are equivalent:*

(i) *For Lebesgue a.e. $\tau \in [0, 1]$,*

$$\lim_{T \rightarrow \infty} \omega_{V(0)} \circ \alpha_T^{0 \rightarrow \tau} (\dot{V}(\tau)) = \omega_{V(\tau)} (\dot{V}(\tau)).$$

⁴ $S(\cdot|\cdot)$ is the relative entropy functional, with the sign and ordering convention of [JPT23, Section 1.1].

(ii) For all $\tau \in [0, 1]$,

$$\lim_{T \rightarrow \infty} S\left(\omega_{V(0)} \circ \alpha_T^{0 \rightarrow \tau} \Big| \omega_{V(\tau)}\right) = 0.$$

(iii) For all $\tau \in [0, 1]$,

$$\lim_{T \rightarrow \infty} \|\omega_{V(0)} \circ \alpha_T^{0 \rightarrow \tau} - \omega_{V(\tau)}\| = 0.$$

In Section 4.2 we extend Theorem 3.4 to translation invariant many body systems.

The adiabatic theorem for gapped ground states of quantum spin systems. A very general extension of Theorem 2.1 to many body quantum spin systems was established in [BDRF17, BDRF18]. Since this result plays no role in our work, we will not discuss it further. The low temperature adiabatic theory of lattice fermion systems was studied in the recent work [GLMP22]. The relation between our work and [GLMP22] remains to be studied further.

4 Structural theory

4.1 Setup

As in [JPT23], we will work with quantum spin systems on the lattice \mathbb{Z}^d and set the inverse temperature to $\beta = 1$. All the results and proofs directly extend to the fermionic case. The notation and terminology are the same as in [JPT23]; see Appendix A below for a brief summary. Given an interaction $\Phi \in \mathcal{B}^r$, the associated local Hamiltonian in a finite cube Λ is

$$H_\Lambda(\Phi) = \sum_{X \subset \Lambda} \Phi(X).$$

The C^* -dynamics it generates on \mathfrak{A} is

$$\alpha_{\Phi, \Lambda}^t(A) = e^{itH_\Lambda(\Phi)} A e^{-itH_\Lambda(\Phi)},$$

and its thermodynamic limit

$$\alpha_\Phi^t(A) = \lim_{\Lambda \uparrow \mathbb{Z}^d} \alpha_{\Phi, \Lambda}^t(A),$$

yields a C^* -dynamics commuting with the natural action of the translation group $\mathbb{Z}^d \ni x \mapsto \varphi^x$.

We consider time-dependent translation invariant interactions described by continuous maps

$$\Phi : [0, 1] \ni \tau \mapsto \Phi_\tau \in \mathcal{B}^r.$$

In analogy with Section 3, the local time-dependent Hamiltonians $H_\Lambda(\Phi_\tau)$ generate a non-autonomous C^* -dynamics $\alpha_{\Phi, \Lambda}^{\sigma \rightarrow \tau}$ on \mathfrak{A} , uniquely characterized by the Cauchy problem for the Heisenberg equation of motion

$$\partial_\tau \alpha_{\Phi, \Lambda}^{\sigma \rightarrow \tau}(A) = \alpha_{\Phi, \Lambda}^{\sigma \rightarrow \tau}(i[H_\Lambda(\Phi_\tau), A]), \quad \alpha_{\Phi, \Lambda}^{\sigma \rightarrow \sigma}(A) = A \in \mathfrak{A}, \quad \sigma, \tau \in [0, 1].$$

The following proposition summarizes the basic properties of its thermodynamic limit needed in this paper. The reader should consult [NSY19] for a more detailed and more general discussion of non-autonomous C^* -dynamics.

Proposition 4.1. *Let $\Phi : [0, 1] \rightarrow \mathcal{B}^r$ be a continuous, time-dependent, translation invariant interaction.*

(1) *For all $\sigma, \tau \in [0, 1]$ and $A \in \mathfrak{A}$, the limit*

$$\alpha_{\Phi}^{\sigma \rightarrow \tau}(A) = \lim_{\Lambda \uparrow \mathbb{Z}^d} \alpha_{\Phi, \Lambda}^{\sigma \rightarrow \tau}(A) \quad (7)$$

exists and defines a two-parameter family of $$ -automorphisms of \mathfrak{A} such that*

$$\alpha_{\Phi}^{\sigma \rightarrow \tau} \circ \alpha_{\Phi}^{\tau \rightarrow \nu} = \alpha_{\Phi}^{\sigma \rightarrow \nu}, \quad \alpha_{\Phi}^{\sigma \rightarrow \sigma} = \text{Id},$$

for all $\sigma, \tau, \nu \in [0, 1]$.

(2) *α_{Φ} commutes with the group action of \mathbb{Z}^d , i.e.,*

$$\alpha_{\Phi}^{\sigma \rightarrow \tau} \circ \varphi^x = \varphi^x \circ \alpha_{\Phi}^{\sigma \rightarrow \tau},$$

for all $\sigma, \tau \in [0, 1]$ and $x \in \mathbb{Z}^d$.

(3) *The map $[0, 1] \ni \sigma, \tau \mapsto \alpha_{\Phi}^{\sigma \rightarrow \tau} \in \text{Aut}(\mathfrak{A})$ is strongly continuous. Moreover, for all $\sigma, \tau \in]0, 1[$ and any $A \in \mathfrak{A}_{\text{loc}}$,*

$$\partial_{\tau} \alpha_{\Phi}^{\sigma \rightarrow \tau}(A) = \alpha_{\Phi}^{\sigma \rightarrow \tau} \circ \delta_{\Phi_{\tau}}(A),$$

where $\delta_{\Phi_{\tau}}$ denotes the map defined by

$$\delta_{\Phi_{\tau}}(A) = \sum_{X \cap \text{supp}(A) \neq \emptyset} i[\Phi_{\tau}(X), A].$$

(4) *For $\sigma, \tau \in [0, 1]$ and integer $N \geq 1$, set $\xi_N = (\tau - \sigma)/N$ and $v_k = \sigma + k\xi_N$. Then, for $A \in \mathfrak{A}$, one has*

$$\alpha_{\Phi}^{\sigma \rightarrow \tau}(A) = \lim_{N \rightarrow \infty} \alpha_{\Phi_{v_0}}^{\xi_N} \circ \dots \circ \alpha_{\Phi_{v_{N-1}}}^{\xi_N}(A).$$

A proof is given in Section 5.2. The family $\{\alpha_{\Phi}^{\sigma \rightarrow \tau}\}_{\sigma, \tau \in [0, 1]}$ defines the non-autonomous C^* -dynamics generated by the time-dependent interaction Φ .

The basic result on the constancy of specific entropy along the state trajectory, [LR68, Theorem 5], extends to time-dependent interactions.

Proposition 4.2. *For any $\nu \in \mathcal{S}_1(\mathfrak{A})$ and $\sigma, \tau \in [0, 1]$,*

$$s(\nu) = s(\nu \circ \alpha_{\Phi}^{\sigma \rightarrow \tau}).$$

Proof. Invoking reversibility, it suffices to prove that, for all $\sigma, \tau \in [0, 1]$, one has

$$s(\nu) \leq s(\nu \circ \alpha_{\Phi}^{\sigma \rightarrow \tau}). \quad (8)$$

Part (4) of Proposition 4.1 gives

$$s(\nu \circ \alpha_{\Phi}^{\sigma \rightarrow \tau}) = s\left(\mathbf{w}^* - \lim_{N \rightarrow \infty} \nu \circ \alpha_{\Phi_{v_0}}^{\xi_N} \circ \dots \circ \alpha_{\Phi_{v_{N-1}}}^{\xi_N}\right),$$

and by [LR68, Theorem 5], we have

$$s(\nu \circ \alpha_{\Phi_{v_0}}^{\xi_N} \circ \dots \circ \alpha_{\Phi_{v_{N-1}}}^{\xi_N}) = s(\nu).$$

The upper-semicontinuity of specific entropy yields (8). \square

4.2 Main results

In this section we associate to a time-dependent interaction Ψ a family of instantaneous equilibrium states. More precisely, we will denote by $\llbracket \nu, \Psi \rrbracket$ a continuous map

$$[0, 1] \ni \tau \mapsto (\nu_\tau, \Psi_\tau) \in \mathcal{S}_1(\mathfrak{A}) \times \mathcal{B}^r$$

such that $\nu_\tau \in \mathcal{S}_{\text{eq}}(\Psi_\tau)$ for all $\tau \in [0, 1]$. We will consider the non-autonomous time evolution on \mathfrak{A} defined by the Cauchy problem

$$\partial_t \gamma^t(A) = \gamma^t \circ \delta_{\Psi_{it}}(A), \quad \gamma^s(A) = A \in \mathfrak{A}_{\text{loc}}, \quad s, t \in [0, T],$$

in the adiabatic limit $T \rightarrow \infty$. A rescaling of the time variables s and t gives

$$\gamma^t = \alpha_{T\Psi}^{s/T \rightarrow t/T},$$

where $\{\alpha_{T\Psi}^{\sigma \rightarrow \tau}\}_{\sigma, \tau \in [0, 1]}$ is the two-parameter family of $*$ -automorphisms generated by the rescaled interaction $T\Psi$, as described in Proposition 4.1.

Definition 4.3.

- (1) A regular pair $(\nu, \Psi) \in \mathcal{S}_1(\mathfrak{A}) \times \mathcal{B}^r$ is *uniquely regular* whenever $\mathcal{S}_{\text{eq}}(\Psi) = \{\nu\}$. A path $\llbracket \nu, \Psi \rrbracket$ is uniquely regular whenever (ν_τ, Ψ_τ) is uniquely regular for all $\tau \in [0, 1]$.
- (2) We say that $\llbracket \nu, \Psi \rrbracket$ satisfies the *adiabatic theorem* if

$$\text{w}^* - \lim_{T \rightarrow \infty} \nu_0 \circ \alpha_{T\Psi}^{0 \rightarrow 1} = \nu_1,$$

and the *path adiabatic theorem* if, for all $\tau \in [0, 1]$,

$$\text{w}^* - \lim_{T \rightarrow \infty} \nu_0 \circ \alpha_{T\Psi}^{0 \rightarrow \tau} = \nu_\tau.$$

- (3) We say that $\llbracket \nu, \Psi \rrbracket$ satisfies the *entropic adiabatic theorem* if

$$\lim_{T \rightarrow \infty} s(\nu_0 \circ \alpha_{T\Psi}^{0 \rightarrow 1} | \nu_1) = 0,$$

and the *entropic path adiabatic theorem* if, for all $\tau \in [0, 1]$,

$$\lim_{T \rightarrow \infty} s(\nu_0 \circ \alpha_{T\Psi}^{0 \rightarrow \tau} | \nu_\tau) = 0.$$

In the fermionic case, gauge-invariance implies that the adiabatic theorem can hold only if $\nu_0(E_N) = \nu_1(E_N)$. Similarly, the path adiabatic theorem can hold only if $\nu_0(E_N) = \nu_\tau(E_N)$ for all $\tau \in [0, 1]$. In what follows we will discuss only the quantum spin case, and leave the elementary reformulations needed to accommodate the fermionic setting to the interested reader.

Proposition 4.4. *Suppose that (ν_1, Ψ_1) is a uniquely regular pair. Then the following statements are equivalent:*

- (i) $s(\nu_0) = s(\nu_1)$ and the adiabatic theorem holds for $\llbracket \nu, \Psi \rrbracket$.
- (ii) $s(\nu_0) = s(\nu_1)$ and $\lim_{T \rightarrow \infty} \nu_0 \circ \alpha_{T\Psi}^{0 \rightarrow 1}(E_{\Psi_1}) = \nu_1(E_{\Psi_1})$.
- (iii) The entropic adiabatic theorem holds for $\llbracket \nu, \Psi \rrbracket$.

Proof. Note that $\nu_0 \circ \alpha_{T\Psi}^{0 \rightarrow 1} \in \mathcal{S}_1(\mathfrak{A})$ by Part (2) of Proposition 4.1.

(i) \Rightarrow (ii). Follows directly from Definition 4.3.

(ii) \Rightarrow (iii). The regularity of (ν_1, Ψ_1) gives

$$s(\nu_0 \circ \alpha_{T\Psi}^{0 \rightarrow 1} | \nu_1) = -s(\nu_0 \circ \alpha_{T\Psi}^{0 \rightarrow 1}) + \nu_0 \circ \alpha_{T\Psi}^{0 \rightarrow 1}(E_{\Psi_1}) + P(\Psi_1). \quad (9)$$

The assumption $s(\nu_0) = s(\nu_1)$ and Proposition 4.2 yield that $s(\nu_0 \circ \alpha_{T\Psi}^{0 \rightarrow 1}) = s(\nu_1)$, and so

$$\lim_{T \rightarrow \infty} s(\nu_0 \circ \alpha_{T\Psi}^{0 \rightarrow 1} | \nu_1) = -s(\nu_1) + \nu_1(E_{\Psi_1}) + P(\Psi_1) = 0,$$

where we used that $\nu_1 \in \mathcal{S}_{\text{eq}}(\Psi_1)$.

(iii) \Rightarrow (i). Let ν_+ be a weak*-limit point of the net $(\nu_0 \circ \alpha_{T\Psi}^{0 \rightarrow 1})_{T>0}$ as $T \rightarrow \infty$. The regularity of (ν_1, Ψ_1) implies that the map $\mathcal{S}_1(\mathfrak{A}) \ni \omega \mapsto s(\omega | \nu_1)$ is lower-semicontinuous, and so

$$0 \leq s(\nu_+ | \nu_1) \leq \liminf_{T \rightarrow \infty} s(\nu_0 \circ \alpha_{T\Psi}^{0 \rightarrow 1} | \nu_1) = 0.$$

Since $\nu_+ \in \mathcal{S}_1(\mathfrak{A})$, we deduce from

$$0 = s(\nu_+ | \nu_1) = -s(\nu_+) + \nu_+(E_{\Psi_1}) + P(\Psi_1)$$

that $\nu_+ \in \mathcal{S}_{\text{eq}}(\Psi_1) = \{\nu_1\}$. It follows that $\nu_0 \circ \alpha_{T\Psi}^{0 \rightarrow 1} \rightarrow \nu_1$ as $T \rightarrow \infty$.

To prove that $s(\nu_0) = s(\nu_1)$, note that the formula (9) and Proposition 4.2 yield

$$s(\nu_0 \circ \alpha_{T\Psi}^{0 \rightarrow 1} | \nu_1) = -s(\nu_0) + \nu_0 \circ \alpha_{T\Psi}^{0 \rightarrow 1}(E_{\Psi_1}) + P(\Psi_1).$$

Hence,

$$0 = \lim_{T \rightarrow \infty} s(\nu_0 \circ \alpha_{T\Psi}^{0 \rightarrow 1} | \nu_1) = -s(\nu_0) + \nu_1(E_{\Psi_1}) + P(\Psi_1) = s(\nu_1) - s(\nu_0),$$

where we used that $\nu_1 \in \mathcal{S}_{\text{eq}}(\Psi_1)$. □

An immediate consequence of the last result is:

Theorem 4.5. *Suppose that the path $\llbracket \nu, \Psi \rrbracket$ is uniquely regular. Then the following statements are equivalent:*

- (i) $s(\nu_0) = s(\nu_\tau)$ for all $\tau \in [0, 1]$ and the path adiabatic theorem holds for $\llbracket \nu, \Psi \rrbracket$.
- (ii) $s(\nu_0) = s(\nu_\tau)$ and $\lim_{T \rightarrow \infty} \nu_0 \circ \alpha_{T\Psi}^{0 \rightarrow \tau}(E_{\Psi_\tau}) = \nu_\tau(E_{\Psi_\tau})$ for all $\tau \in [0, 1]$.
- (iii) The entropic path adiabatic theorem holds for $\llbracket \nu, \Psi \rrbracket$.

Our next result is:

Theorem 4.6. *Suppose that the path $\llbracket \nu, \Psi \rrbracket$ is uniquely regular. If $\Psi \in C^1([0, 1], \mathcal{B}^r)$, then the following statements are equivalent:*

- (i) For Lebesgue a.e. $\tau \in [0, 1]$,

$$\lim_{T \rightarrow \infty} \nu_0 \circ \alpha_{T\Psi}^{0 \rightarrow \tau}(\partial_\tau E_{\Psi_\tau}) = \nu_\tau(\partial_\tau E_{\Psi_\tau}).$$

- (ii) The entropic path adiabatic theorem holds for $\llbracket \nu, \Psi \rrbracket$.

Proof. (ii) \Rightarrow (i) follows from Theorem 4.5.

(i) \Rightarrow (ii). The map $[0, 1] \ni \tau \mapsto E_{\Psi_\tau}$ is continuously differentiable by assumption. Moreover, it follows from Part (3) of Proposition 4.1 that for $A \in \mathfrak{A}_{\text{loc}}$,

$$\alpha_{T\Psi}^{\sigma \rightarrow \tau}(A) = A + T \int_{\sigma}^{\tau} \alpha_{T\Psi}^{\sigma \rightarrow \nu} \circ \delta_{\Psi_\nu}(A) d\nu.$$

By the proof of Part (2) of [JPT23, Theorem 2.8], for any $\tau, \nu \in [0, 1]$ one has $E_{\Psi_\tau} \in \text{Dom}(\delta_{\Psi_\nu})$ and

$$\omega \circ \delta_{\Psi_\tau}(E_{\Psi_\tau}) = 0 \text{ for all } \omega \in \mathcal{S}_1(\mathfrak{A}). \quad (10)$$

Moreover, there exist a sequence $(E_{\Psi_{\tau,n}})_{n \in \mathbb{N}}$ in $\mathfrak{A}_{\text{loc}}$ such that⁵ $\|\delta_{\Psi_\nu}(E_{\Psi_{\tau,n}})\| \leq 2\|\Psi\|_\tau^2$, and

$$\lim_{n \rightarrow \infty} E_{\Psi_{\tau,n}} = E_{\Psi_\tau}, \quad \lim_{n \rightarrow \infty} \delta_{\Psi_\nu}(E_{\Psi_{\tau,n}}) = \delta_{\Psi_\nu}(E_{\Psi_\tau}).$$

Hence, the dominated convergence theorem yields

$$\alpha_{T\Psi}^{\sigma \rightarrow \tau}(E_{\Psi_\tau}) = E_{\Psi_\tau} + T \int_{\sigma}^{\tau} \alpha_{T\Psi}^{\sigma \rightarrow \nu} \circ \delta_{\Psi_\nu}(E_{\Psi_\tau}) d\nu,$$

from which we deduce

$$\partial_\tau \nu_0 \circ \alpha_{T\Psi}^{0 \rightarrow \tau}(E_{\Psi_\tau}) = T \nu_0 \circ \alpha_{T\Psi}^{0 \rightarrow \tau} \circ \delta_{\Psi_\tau}(E_{\Psi_\tau}) + \nu_0 \circ \alpha_{T\Psi}^{0 \rightarrow \tau}(\partial_\tau E_{\Psi_\tau}).$$

Since $\nu_0 \circ \alpha_{T\Psi}^{0 \rightarrow \tau} \in \mathcal{S}_1(\mathfrak{A})$ by Part (2) of Proposition 4.1, it follows from (10) that

$$\nu_0 \circ \alpha_{T\Psi}^{0 \rightarrow \tau}(E_{\Psi_\tau}) - \nu_0(E_{\Psi_0}) = \int_0^\tau \nu_0 \circ \alpha_{T\Psi}^{0 \rightarrow \sigma}(\partial_\sigma E_{\Psi_\sigma}) d\sigma.$$

Taking the limit $T \rightarrow \infty$, it follows from our hypotheses and the dominated convergence theorem that

$$\lim_{T \rightarrow \infty} \nu_0 \circ \alpha_{\Psi, T}^{0 \rightarrow \tau}(E_{\Psi_\tau}) - \nu_0(E_{\Psi_0}) = \int_0^\tau \nu_\sigma(\partial_\sigma E_{\Psi_\sigma}) d\sigma. \quad (11)$$

The uniqueness of the equilibrium state for Ψ_τ implies that the pressure $\mathcal{B}^r \ni \Phi \mapsto P(\Phi)$ is differentiable at Ψ_τ , with

$$P(\Psi_\tau + \Phi) - P(\Psi_\tau) + \nu_\tau(E_\Phi) = o(\|\Phi\|_r)$$

for $\Phi \in \mathcal{B}^r$ [LR68, Theorem 3]. Invoking again the differentiability of $\tau \mapsto E_{\Psi_\tau}$ yields

$$\partial_\tau P(\Psi_\tau) = -\nu_\tau(\partial_\tau E_{\Psi_\tau}). \quad (12)$$

Combining Relations (12) and (11) we derive

$$\lim_{T \rightarrow \infty} \nu_0 \circ \alpha_{\Psi, T}^{0 \rightarrow \tau}(E_{\Psi_\tau}) + P(\Psi_\tau) = \nu_0(E_{\Psi_0}) + P(\Psi_0).$$

Now, since (ν_τ, Ψ_τ) is a regular pair, we have

$$s(\nu_0 \circ \alpha_{\Psi, T}^{0 \rightarrow \tau} | \nu_\tau) = -s(\nu_0) + \nu_0 \circ \alpha_{\Psi, T}^{0 \rightarrow \tau}(E_{\Psi_\tau}) + P(\Psi_\tau),$$

and so

$$\lim_{T \rightarrow \infty} s(\nu_0 \circ \alpha_{\Psi, T}^{0 \rightarrow \tau} | \nu_\tau) = -s(\nu_0) + \nu_0(E_{\Psi_0}) + P(\Psi_0) = 0.$$

□

⁵The norm $\|\Psi\|_r$ of a time-dependent interaction Ψ is defined in (21).

The following example illustrates the last result. Let $\Phi_0, \Phi_1 \in \mathcal{B}^r$, and let $\lambda \in C^1([0, 1])$ be such that $\lambda(0) = 0, \lambda(1) = 1$. Set

$$\Psi_\tau = \Phi_0 + \lambda(\tau)(\Phi_1 - \Phi_0),$$

and assume that the path $\llbracket \nu, \Psi \rrbracket$ is uniquely regular. Then the entropic path adiabatic theorem holds for $\llbracket \nu, \Psi \rrbracket$ iff, for all $\tau \in [0, 1]$ with $\partial_\tau \lambda(\tau) \neq 0$,

$$\lim_{T \rightarrow \infty} \nu_0 \circ \alpha_{T\Psi}^{0 \rightarrow \tau}(E_{\Phi_1 - \Phi_0}) = \nu_\tau(E_{\Phi_1 - \Phi_0}).$$

We finish this section with the translation invariant version Theorem 3.4. This result is direct a consequence Theorems 4.5 and 4.6 and we state it for completeness reason.

Theorem 4.7. *Suppose that the path $\llbracket \nu, \Psi \rrbracket$ is uniquely regular. If $\Psi \in C^1([0, 1], \mathcal{B}^r)$, then the following statements are equivalent:*

(i) *For Lebesgue a.e. $\tau \in [0, 1]$,*

$$\lim_{T \rightarrow \infty} \nu_0 \circ \alpha_{T\Psi}^{0 \rightarrow \tau}(\partial_\tau E_{\Psi_\tau}) = \nu_\tau(\partial_\tau E_{\Psi_\tau}).$$

(ii) *The entropic path adiabatic theorem holds for $\llbracket \nu, \Psi \rrbracket$.*

(iii) *The path adiabatic theorem holds for $\llbracket \nu, \Psi \rrbracket$.*

Moreover, any of the above statements implies

(iv) *For all $\tau \in [0, 1]$,*

$$s(\nu_\tau) = s(\nu_0).$$

4.3 Adiabaticity and the approach to equilibrium

Theorem 3.1 points to a close relation between the adiabatic theorem for local perturbations of KMS states and return to equilibrium. In view of the formal analogy between Proposition 3.4 and Theorem 4.7, one may expect a similar relation between the adiabatic theorem for translation invariant interactions and approach to equilibrium. Our last result shows that this is not the case.

We first recall the framework of approach to equilibrium developed in [JPT23]. Let $\omega \in \mathcal{S}_1(\mathfrak{A})$ and $\Phi \in \mathcal{B}^r$. Set

$$\bar{\omega}_T = \frac{1}{T} \int_0^T \omega \circ \tau_\Phi^t dt.$$

We denote by $\mathcal{S}_+(\omega, \Phi)$ the set of weak*-limit points of the net $(\bar{\omega}_T)_{T>0}$ as $T \rightarrow \infty$. Approach to equilibrium holds in this setting whenever $\mathcal{S}_+(\omega, \Phi) \cap \mathcal{S}_{\text{eq}}(\Phi) \neq \emptyset$.

Theorem 4.8. *Let (ν_0, Φ_0) and (ν_1, Φ_1) be two uniquely regular pairs and suppose that Φ_0 and Φ_1 are not physically equivalent. Then the following statements are mutually exclusive.*

(i) $\nu_1 \in \mathcal{S}_+(\nu_0, \Phi_1)$, i.e., *approach to equilibrium holds for $(\mathfrak{A}, \alpha_{\Phi_1}, \nu_0)$.*

(ii) *There exists a uniquely regular path $\llbracket \nu, \Psi \rrbracket$, with $\Psi \in C^1([0, 1], \mathcal{B}^r)$, connecting (ν_0, Φ_0) to (ν_1, Φ_1) and satisfying the path adiabatic theorem.*

Proof. Approach to equilibrium has been characterized in terms of a strict increase of specific entropy. According to [JPT23, Theorem 2.20], if $v_1 \in \mathcal{S}_+(v_0, \Phi_1)$, then $s(v_0) < s(v_1)$ since Φ_0 and Φ_1 are not physically equivalent. By Theorem 4.7, this fact is incompatible with the existence of the path $[[v, \Psi]]$.

Conversely, Theorem 4.7 shows that the existence of the path $[[v, \Psi]]$ implies the equality $s(v_0) = s(v_1)$. By [JPT23, Theorem 2.20], if $v_1 \in \mathcal{S}_+(v_0, \Phi_1)$, then $s(v_0) = s(v_1)$ implies $v_0 = v_1$ and that Φ_0 and Φ_1 are physically equivalent, which contradicts the assumption of the theorem. \square

5 Proofs of Propositions 3.2 and 4.1

5.1 Proof of Proposition 3.2

Our standing assumption is that the map $V \in C^1([0, 1], \mathcal{O})$ takes its value in the self-adjoint elements of \mathcal{O} , \dot{V} denoting its derivative. Denote by $[0, 1] \ni s, t \mapsto \alpha_T^{s \rightarrow t}$ the 2-parameter family of $*$ -automorphisms of \mathcal{O} uniquely determined by the Cauchy problem

$$\frac{1}{T} \partial_t \alpha_T^{s \rightarrow t}(A) = \alpha_T^{s \rightarrow t} \circ \delta_{V(t)}(A), \quad \alpha_T^{s \rightarrow s}(A) = A \in \text{Dom}(\delta), \quad s, t \in [0, 1].$$

We start with:

Lemma 5.1. *For $s, t \in [0, 1]$, one has*

$$\alpha_T^{s \rightarrow t} = \gamma_T^{s \rightarrow t} \circ \alpha_{V(t)}^{(t-s)T}, \quad (13)$$

where $\gamma_T^{s \rightarrow t}$ is the inner $*$ -automorphism of \mathcal{O} given by $\gamma_T^{s \rightarrow t}(A) = \Gamma_T^{s \rightarrow t} A \Gamma_T^{s \rightarrow t*}$, and the 2-parameter family of unitaries Γ_T is the solution of the Cauchy problem

$$\frac{i}{T} \partial_t \Gamma_T^{s \rightarrow t} = \Gamma_T^{s \rightarrow t} \int_s^t \alpha_{V(t)}^{(t-r)T} (\dot{V}(t)) dr, \quad \Gamma_T^{s \rightarrow s} = \mathbb{1} \quad s, t \in [0, 1]. \quad (14)$$

Moreover, one has $\Gamma_T^{s \rightarrow t} \in \text{Dom}(\delta)$ for all $s, t \in [0, 1]$.

Proof. Fix $T > 0$, $s \in [0, 1]$, and set

$$\tilde{H}_t = T \int_s^t \alpha_{V(t)}^{(t-r)T} (\dot{V}(t)) dr,$$

so that

$$\partial_t \gamma_T^{s \rightarrow t}(A) = \gamma_T^{s \rightarrow t}(-i[\tilde{H}_t, A]). \quad (15)$$

For $A \in \text{Dom}(\delta)$, one has $\partial_u \delta_{V(u)}(A) = i[\dot{V}(u), A]$, and Duhamel's formula yields

$$\partial_u \alpha_{V(u)}^{(t-s)T}(A) = T \int_s^t \alpha_{V(u)}^{(r-s)T} (i[\dot{V}(u), \alpha_{V(u)}^{(t-r)T}(A)]) dr,$$

so that

$$\partial_u \alpha_{V(u)}^{(t-s)T}(A) \Big|_{u=t} = i[\tilde{H}_t, \alpha_{V(t)}^{(t-s)T}(A)],$$

and hence

$$\partial_t \alpha_{V(t)}^{(t-s)T}(A) = T \alpha_{V(t)}^{(t-s)T} \circ \delta_{V(t)}(A) + i[\tilde{H}_t, \alpha_{V(t)}^{(t-s)T}(A)]. \quad (16)$$

Combining Relations (15) and (16), we derive

$$\begin{aligned} \partial_t \gamma_T^{s \rightarrow t} \circ \alpha_{V(t)}^{(t-s)T}(A) &= \gamma_T^{s \rightarrow t}(-i[\tilde{H}_t, \alpha_{V(t)}^{(t-s)T}(A)]) + \gamma_T^{s \rightarrow t}(T \alpha_{V(t)}^{(t-s)T} \circ \delta_{V(t)}(A) + i[\tilde{H}_t, \alpha_{V(t)}^{(t-s)T}(A)]) \\ &= T \gamma_T^{s \rightarrow t} \circ \alpha_{V(t)}^{(t-s)T} \circ \delta_{V(t)}(A). \end{aligned}$$

Since obviously $\gamma_T^{s \rightarrow t} \circ \alpha_{V(t)}^{(s-s)T}(A) = A$, Relation (13) holds.

To prove the last statement, for fixed $s, t \in [0, 1]$ let the sequence $(Q_n)_{n \in \mathbb{N}} \subset \text{Dom}(\delta_{V(t)})$ be such that $\lim_{n \rightarrow \infty} Q_n = \dot{V}(t)$. Setting

$$H_n = T \int_s^t \alpha_{V(t)}^{(r-s)T}(Q_n) dr,$$

and observing that

$$T \int_s^t \alpha_{V(t)}^{(r-s)T}(\delta_{V(t)}(Q_n)) dr = \int_s^t \partial_r \alpha_{V(t)}^{(r-s)T}(Q_n) dr = \alpha_{V(t)}^{(t-s)T}(Q_n) - Q_n,$$

we derive

$$\delta_{V(t)}(H_n) = \alpha_{V(t)}^{(t-s)T}(Q_n) - Q_n.$$

From the obvious facts that

$$\lim_{n \rightarrow \infty} H_n = \tilde{H}_t, \quad \lim_{n \rightarrow \infty} \delta_{V(t)}(H_n) = \alpha_{V(t)}^{(t-s)T}(\dot{V}(t)) - \dot{V}(t),$$

the closedness of $\delta_{V(t)}$ allows us to conclude that $\tilde{H}_t \in \text{Dom}(\delta_{V(t)}) = \text{Dom}(\delta)$ and that

$$\delta_{V(t)}(\tilde{H}_t) = \alpha_{V(t)}^{(t-s)T}(\dot{V}(t)) - \dot{V}(t). \quad (17)$$

Writing the solution to the Cauchy problem (14) as the uniformly convergent Dyson series

$$\Gamma_T^{s \rightarrow t} = \sum_{n \geq 0} (-i)^n \int_{s \leq t_1 \leq \dots \leq t_n \leq t} \tilde{H}_{t_1} \cdots \tilde{H}_{t_n} dt_1 \cdots dt_n,$$

one deduces $\Gamma_T^{s \rightarrow t} \in \text{Dom}(\delta)$ for all $s, t \in [0, 1]$ and that

$$\delta(\Gamma_T^{s \rightarrow t}) = \sum_{n \geq 0} (-i)^n \sum_{k=1}^n \int_{s \leq t_1 \leq \dots \leq t_n \leq t} \tilde{H}_{t_1} \cdots \delta(\tilde{H}_{t_k}) \cdots \tilde{H}_{t_n} dt_1 \cdots dt_n,$$

where the series on right hand side converges uniformly. \square

We now proceed with the proof of Proposition 3.2.

The previous lemma gives

$$S(\omega_{V(s)} \circ \alpha_T^{s \rightarrow t} | \omega_{V(t)}) = S(\omega_{V(s)} \circ \gamma_T^{s \rightarrow t} \circ \alpha_{V(t)}^{(t-s)T} | \omega_{V(t)}) = S(\omega_{V(s)} \circ \gamma_T^{s \rightarrow t} | \omega_{V(t)}),$$

and invoking [JP03, Theorem 1.1]⁶, we can write

$$S(\omega_{V(s)} \circ \alpha_T^{s \rightarrow t} | \omega_{V(t)}) = S(\omega_{V(s)} | \omega_{V(t)}) - i\beta \omega_{V(s)}(\Gamma_T^{s \rightarrow t} \delta_{V(t)}(\Gamma_T^{s \rightarrow t*})). \quad (18)$$

To deal with the first term on the right-hand side, we invoke Araki's perturbation theory (see, e.g., [DJP03, Theorem 5.1]). Setting $\nu = \omega_{V(s)}$ and $W_t = V(t) - V(s)$, we have

$$S(\omega_{V(s)} | \omega_{V(t)}) = S(\nu | \nu_{W_t}) = \beta \nu(W_t) + \log \langle \Omega_\nu, e^{-\beta(L + \pi_\nu(W_t))} \Omega_\nu \rangle,$$

where $(\mathcal{H}_\nu, \pi_\nu, \Omega_\nu)$ denotes the GNS representation of \mathcal{O} induced by ν , and L is the standard Liouvillean generating the unitary implementation of the C^* -dynamics $\alpha_{V(s)}$ on \mathcal{H}_ν . Taking the derivative w.r.t. t of the logarithmic term and using the facts that $W_s = 0$ and $L\Omega_\nu = 0$ gives

$$\log \langle \Omega_\nu, e^{-\beta(L + \pi_\nu(W_t))} \Omega_\nu \rangle = \int_s^t \partial_u \log \langle \Omega_\nu, e^{-\beta(L + \pi_\nu(W_u))} \Omega_\nu \rangle du.$$

⁶We note that the definition of relative entropy used in [JP03, DJP03] differs in its sign with the one used here.

Duhamel's formula further yields

$$\begin{aligned}\partial_u \log \langle \Omega_v, e^{-\beta(L+\pi_v(W_u))} \Omega_v \rangle &= - \int_0^\beta \frac{\langle \Omega_v, e^{-\gamma(L+\pi_v(W_u))} \pi_v(\dot{V}(u)) e^{-(\beta-\gamma)(L+\pi_v(W_u))} \Omega_v \rangle}{\|\Psi_u\|^2} d\gamma \\ &= - \int_{-\beta/2}^{\beta/2} \frac{\langle \Psi_u, e^{-\gamma(L+\pi_v(W_u))} \pi_v(\dot{V}(u)) e^{\gamma(L+\pi_v(W_u))} \Psi_u \rangle}{\|\Psi_u\|^2} d\gamma,\end{aligned}$$

where $\Psi_u = e^{-\beta(L+\pi_v(W_u))/2} \Omega_v$ is, up to normalization, the vector representative of $\omega_{V(u)}$ in \mathcal{H}_v . It follows that

$$\partial_u \log \langle \Omega_v, e^{-\beta(L+\pi_v(W_u))} \Omega_v \rangle = - \int_{-\beta/2}^{\beta/2} \omega_{V(u)} \left(\alpha_{V(u)}^{i\gamma}(\dot{V}(u)) \right) d\gamma = -\beta \omega_{V(u)}(\dot{V}(u)),$$

and hence

$$S(\omega_{V(s)} | \omega_{V(t)}) = \beta \omega_{V(s)}(V(t) - V(s)) - \beta \int_s^t \omega_{V(u)}(\dot{V}(u)) du = -\beta \int_s^t (\omega_{V(u)} - \omega_{V(s)})(\dot{V}(u)) du.$$

To deal with the second term on the right-hand side of (18), we note that

$$i\Gamma_T^{s \rightarrow t} \delta_{V(t)}(\Gamma_T^{s \rightarrow t*}) = \int_s^t i\partial_u \Gamma_T^{s \rightarrow u} \delta_{V(u)}(\Gamma_T^{s \rightarrow u*}) du,$$

and so, using Relation (17), we have that

$$\begin{aligned}i\partial_u \Gamma_T^{s \rightarrow u} \delta_{V(u)}(\Gamma_T^{s \rightarrow u*}) &= \Gamma_T^{s \rightarrow u} \tilde{H}_u \delta_{V(u)}(\Gamma_T^{s \rightarrow u*}) - \Gamma_T^{s \rightarrow u} \delta_{V(u)}(\tilde{H}_u \Gamma_T^{s \rightarrow u*}) - \Gamma_T^{s \rightarrow u} [\dot{V}(u), \Gamma_T^{s \rightarrow u*}] \\ &= -\Gamma_T^{s \rightarrow u} (\delta_{V(u)}(\tilde{H}_u) + \dot{V}(u)) \Gamma_T^{s \rightarrow u*} + \dot{V}(u) \\ &= -\Gamma_T^{s \rightarrow u} \alpha_{V(u)}^{(u-s)T}(\dot{V}(u)) \Gamma_T^{s \rightarrow u*} + \dot{V}(u) \\ &= -\alpha_T^{s \rightarrow u}(\dot{V}(u)) + \dot{V}(u).\end{aligned}$$

It follows that

$$-i\beta \omega_{V(s)}(\Gamma_T^{s \rightarrow t} \delta_{V(t)}(\Gamma_T^{s \rightarrow t*})) = \beta \int_s^t (\omega_{V(s)} \circ \alpha_T^{s \rightarrow u}(\dot{V}(u)) - \omega_{V(s)}(\dot{V}(u))) du,$$

and finally, Relation (18) gives

$$S(\omega_{V(s)} \circ \alpha_T^{s \rightarrow t} | \omega_{V(t)}) = \beta \int_s^t (\omega_{V(s)} \circ \alpha_T^{s \rightarrow u}(\dot{V}(u)) - \omega_{V(u)}(\dot{V}(u))) du.$$

5.2 Proof of Proposition 4.1

We start with some preliminary observations. By [BR81, Theorem 6.2.4], the time-dependent derivation $[0, 1] \ni \tau \mapsto \delta_{\Phi_\tau}$ defined on $\mathfrak{A}_{\text{loc}}$ by

$$\delta_{\Phi_\tau}(A) = \sum_{X \cap \text{supp}(A) \neq \emptyset} i[\Phi_\tau(X), A],$$

is closable and its closure, which we shall denote by the same symbol, generates the frozen C^* -dynamics

$$\mathbb{R} \ni t \mapsto \alpha_{\Phi_\tau}^t = e^{t\delta_{\Phi_\tau}}.$$

The translation invariance of the interaction Φ_τ implies that $\varphi^x(\text{Dom}(\delta_{\Phi_\tau})) = \text{Dom}(\delta_{\Phi_\tau})$ and that

$$\delta_{\Phi_\tau} \circ \varphi^x = \varphi^x \circ \delta_{\Phi_\tau}, \quad (19)$$

for all $x \in \mathbb{Z}^d$.

Note that, for $A \in \mathfrak{A}_{\text{loc}}$, one has

$$\delta_{\Phi_\tau}(A) = \lim_{\Lambda \uparrow \mathbb{Z}^d} \delta_{\Phi_\tau, \Lambda}(A),$$

where the convergence is in norm and

$$\delta_{\Phi_\tau, \Lambda}(A) = \sum_{X \subset \Lambda} i[\Phi_\tau(X), A] = i[H_\Lambda(\Phi_\tau), A],$$

is such that $\text{supp}(\delta_{\Phi_\tau, \Lambda}(A)) \subset \text{supp}(A) \cup \Lambda$, i.e., $\delta_{\Phi_\tau, \Lambda}(\mathfrak{A}_{\text{loc}}) \subset \mathfrak{A}_{\text{loc}}$. It follows that the non-autonomous C^* -dynamics generated by the time-dependent local Hamiltonian $H_\Lambda(\Phi_\tau)$ is given by the norm-convergent Dyson expansion

$$\alpha_{\Phi, \Lambda}^{\sigma \rightarrow \tau}(A) = \begin{cases} A + \sum_{n \geq 1} \int_{\Delta_n(\sigma, \tau)} \delta_{\Phi_{\tau_1}, \Lambda} \circ \cdots \circ \delta_{\Phi_{\tau_n}, \Lambda}(A) d\tau_1 \cdots d\tau_n, & \text{for } 0 \leq \sigma \leq \tau \leq 1; \\ A + \sum_{n \geq 1} \int_{\Delta_n(\sigma, \tau)} \delta_{\Phi_{\tau_n}, \Lambda} \circ \cdots \circ \delta_{\Phi_{\tau_1}, \Lambda}(A) d\tau_1 \cdots d\tau_n, & \text{for } 0 \leq \tau \leq \sigma \leq 1; \end{cases} \quad (20)$$

for $A \in \mathfrak{A}_{\text{loc}}$, where $\Delta_n(\sigma, \tau) = \{(\tau_1, \dots, \tau_n) \mid 0 \leq \min(\sigma, \tau) \leq \tau_1 \leq \cdots \leq \tau_n \leq \max(\sigma, \tau)\}$.

In the following, the norm of a time-dependent continuous interaction $[0, 1] \ni \tau \mapsto \Phi(\tau) \in \mathfrak{B}^r$ is taken to be

$$\|\Phi\|_r = \sup_{\tau \in [0, 1]} \|\Phi_\tau\|_r. \quad (21)$$

(1) A trivial extension of [Isr79, Lemma III.3.5] gives that, for $A \in \mathfrak{A}_{\text{loc}}$ and $\tau_1, \dots, \tau_n \in [0, 1]$,

$$\sum_{X_1, \dots, X_n \in \mathfrak{F}} \|i[\Phi_{\tau_1}(X_1), i[\Phi_{\tau_2}(X_2), \dots, i[\Phi_{\tau_n}(X_n), A] \cdots]]\| \leq \|A\| e^{r|\text{supp}(A)|} \left(\frac{2\|\Phi\|_r}{r}\right)^n n!.$$

Hence,

$$\sup_{\tau_1, \dots, \tau_n \in [0, 1]} \|\delta_{\Phi_{\tau_1}} \circ \cdots \circ \delta_{\Phi_{\tau_n}}(A)\| \leq \|A\| e^{r|\text{supp}(A)|} \left(\frac{2\|\Phi\|_r}{r}\right)^n n!, \quad (22)$$

and the same estimate holds with $\delta_{\Phi_{\tau_k}}$ replaced by $\delta_{\Phi_{\tau_k}, \Lambda}$. Thus, taking the thermodynamic limit in (20) yields that, for $|\tau - \sigma| < \epsilon = r/2\|\Phi\|_r$,

$$\lim_{\Lambda \uparrow \mathbb{Z}^d} \alpha_{\Phi, \Lambda}^{\sigma \rightarrow \tau}(A) = \begin{cases} A + \sum_{n \geq 1} \int_{\Delta_n(\sigma, \tau)} \delta_{\Phi_{\tau_1}} \circ \cdots \circ \delta_{\Phi_{\tau_n}}(A) d\tau_1 \cdots d\tau_n, & \text{for } 0 \leq \sigma \leq \tau \leq 1; \\ A + \sum_{n \geq 1} \int_{\Delta_n(\sigma, \tau)} \delta_{\Phi_{\tau_n}} \circ \cdots \circ \delta_{\Phi_{\tau_1}}(A) d\tau_1 \cdots d\tau_n, & \text{for } 0 \leq \tau \leq \sigma \leq 1; \end{cases} \quad (23)$$

the limit being uniform for $\tau - \sigma$ in compact subsets of $] -\epsilon, \epsilon[$. Since the maps $\alpha_{\Phi, \Lambda}^{\sigma \rightarrow \tau} : \mathfrak{A}_{\text{loc}} \rightarrow \mathfrak{A}$ are isometric, their limit $\alpha_{\Phi}^{\sigma \rightarrow \tau}$ is norm continuous, and hence uniquely extends by continuity to an isometry on \mathfrak{A} . It follows that (7) holds for all $A \in \mathfrak{A}$ and $\sigma, \tau \in [0, 1]$ satisfying $|\tau - \sigma| < \epsilon$. Moreover, as norm-limits of $*$ -morphisms, the maps $\alpha_{\Phi}^{\sigma \rightarrow \tau}$ are themselves $*$ -morphisms. For $\sigma, \tau, \nu \in [0, 1]$ such that $\max(|\tau - \sigma|, |\sigma - \nu|) < \epsilon$, it follows from the continuity of $\alpha_{\Phi}^{\sigma \rightarrow \tau}$ that

$$\lim_{\Lambda \uparrow \mathbb{Z}^d} \alpha_{\Phi}^{\sigma \rightarrow \tau} \circ \alpha_{\Phi, \Lambda}^{\tau \rightarrow \nu}(A) = \alpha_{\Phi}^{\sigma \rightarrow \tau} \circ \alpha_{\Phi}^{\tau \rightarrow \nu}(A),$$

for $A \in \mathfrak{A}_{\text{loc}}$. Writing

$$\alpha_{\Phi}^{\sigma \rightarrow \tau} \circ \alpha_{\Phi, \Lambda}^{\tau \rightarrow \nu}(A) = \alpha_{\Phi, \Lambda}^{\sigma \rightarrow \nu}(A) + (\alpha_{\Phi}^{\sigma \rightarrow \tau} - \alpha_{\Phi, \Lambda}^{\sigma \rightarrow \tau}) \circ \alpha_{\Phi}^{\tau \rightarrow \nu}(A) + (\alpha_{\Phi}^{\sigma \rightarrow \tau} - \alpha_{\Phi, \Lambda}^{\sigma \rightarrow \tau}) \circ (\alpha_{\Phi, \Lambda}^{\tau \rightarrow \nu} - \alpha_{\Phi}^{\tau \rightarrow \nu})(A),$$

and using that

$$\begin{aligned} \lim_{\Lambda \uparrow \mathbb{Z}^d} (\alpha_{\Phi}^{\sigma \rightarrow \tau} - \alpha_{\Phi, \Lambda}^{\sigma \rightarrow \tau}) \circ \alpha_{\Phi}^{\tau \rightarrow \nu}(A) &= 0, \\ \lim_{\Lambda \uparrow \mathbb{Z}^d} \|(\alpha_{\Phi}^{\sigma \rightarrow \tau} - \alpha_{\Phi, \Lambda}^{\sigma \rightarrow \tau}) \circ (\alpha_{\Phi, \Lambda}^{\tau \rightarrow \nu} - \alpha_{\Phi}^{\tau \rightarrow \nu})(A)\| &\leq 2 \lim_{\Lambda \uparrow \mathbb{Z}^d} \|\alpha_{\Phi, \Lambda}^{\tau \rightarrow \nu}(A) - \alpha_{\Phi}^{\tau \rightarrow \nu}(A)\| = 0, \end{aligned}$$

we derive

$$\alpha_{\Phi}^{\sigma \rightarrow \tau} \circ \alpha_{\Phi}^{\tau \rightarrow \nu}(A) = \alpha_{\Phi}^{\sigma \rightarrow \nu}(A), \quad (24)$$

provided that $|\nu - \sigma| < \epsilon$. This identity extends by continuity to all $A \in \mathfrak{A}$, and allows to extend the family of *-morphisms $(\alpha_{\Phi}^{\sigma \rightarrow \tau})_{|\tau - \sigma| < \epsilon}$ to arbitrary $\sigma, \tau \in [0, 1]$ by setting

$$\alpha_{\Phi}^{\sigma \rightarrow \tau} = \alpha_{\Phi}^{\sigma \rightarrow \tau_1} \circ \alpha_{\Phi}^{\tau_1 \rightarrow \tau_2} \circ \dots \circ \alpha_{\Phi}^{\tau_{n-1} \rightarrow \tau_n} \circ \alpha_{\Phi}^{\tau_n \rightarrow \tau}, \quad (25)$$

where $\tau_0 = \sigma, \tau_1, \dots, \tau_{n+1} = \tau$ are such that $\max_k |\tau_{k+1} - \tau_k| < \epsilon$. One easily checks that the left-hand side of this expression does not depend on the choice of the subdivisions τ_1, \dots, τ_n on the right-hand side. By construction, this extension satisfies (24) for all $\sigma, \tau, \nu \in [0, 1]$. From the telescopic expansion based on (25),

$$\alpha_{\Phi}^{\sigma \rightarrow \tau}(A) - \alpha_{\Phi, \Lambda}^{\sigma \rightarrow \tau}(A) = \sum_{k=0}^n \alpha_{\Phi, \Lambda}^{\sigma \rightarrow \tau_1} \circ \dots \circ \alpha_{\Phi, \Lambda}^{\tau_{k-1} \rightarrow \tau_k} \circ (\alpha_{\Phi}^{\tau_k \rightarrow \tau_{k+1}} - \alpha_{\Phi, \Lambda}^{\tau_k \rightarrow \tau_{k+1}}) \circ \alpha_{\Phi}^{\tau_{k+1} \rightarrow \tau_{k+2}} \circ \dots \circ \alpha_{\Phi}^{\tau_n \rightarrow \tau}(A),$$

we derive

$$\|\alpha_{\Phi}^{\sigma \rightarrow \tau}(A) - \alpha_{\Phi, \Lambda}^{\sigma \rightarrow \tau}(A)\| \leq \sum_{k=0}^n \|(\alpha_{\Phi}^{\tau_k \rightarrow \tau_{k+1}} - \alpha_{\Phi, \Lambda}^{\tau_k \rightarrow \tau_{k+1}})(A_k)\|,$$

with $A_k = \alpha_{\Phi}^{\tau_{k+1} \rightarrow \tau_{k+2}} \circ \dots \circ \alpha_{\Phi}^{\tau_n \rightarrow \tau}(A)$. It follows that (7) holds for all $A \in \mathfrak{A}$ and all $\sigma, \tau \in [0, 1]$. Note that (24) implies that $\alpha_{\Phi}^{\sigma \rightarrow \tau} \circ \alpha_{\Phi}^{\tau \rightarrow \sigma} = \alpha_{\Phi}^{\sigma \rightarrow \sigma} = \text{Id}$, which shows that the maps $\alpha_{\Phi}^{\sigma \rightarrow \tau}$ are *-automorphisms of \mathfrak{A} .

(2) The translation invariance $\alpha_{\Phi}^{\sigma \rightarrow \tau} \circ \varphi^x = \varphi^x \circ \alpha_{\Phi}^{\sigma \rightarrow \tau}$ follows immediately from Relation (19).

(3) For $A \in \mathfrak{A}_{\text{loc}}$ and $|\tau - \sigma| < \epsilon/2$, combining the Dyson expansion (23) with the estimate (22), we get

$$\|\alpha_{\Phi}^{\sigma \rightarrow \tau}(A) - A\| \leq 2\epsilon^{-1} \|A\| e^{d|\text{supp}(A)|} |\tau - \sigma|.$$

Since $\alpha_{\Phi}^{\sigma \rightarrow \tau}$ is an isometry and $\mathfrak{A}_{\text{loc}}$ is dense in \mathfrak{A} , we conclude that the map $(\sigma, \tau) \mapsto \alpha_{\Phi}^{\sigma \rightarrow \tau}$ is strongly continuous on the diagonal $(\tau, \tau) \in [0, 1] \times [0, 1]$. For $(\sigma_0, \tau_0) \in [0, 1] \times [0, 1]$, Property (24) gives

$$\alpha_{\Phi}^{\sigma \rightarrow \tau}(A) - \alpha_{\Phi}^{\sigma_0 \rightarrow \tau_0}(A) = \alpha_{\Phi}^{\sigma \rightarrow \tau_0}(\alpha_{\Phi}^{\tau_0 \rightarrow \tau}(A) - A) + (\alpha_{\Phi}^{\sigma \rightarrow \sigma_0}(B) - B),$$

with $B = \alpha_{\Phi}^{\sigma_0 \rightarrow \tau_0}(A)$, which shows strong continuity at (σ_0, τ_0) .

From (24) and the Dyson expansion (23), we further derive that

$$\partial_{\tau} \alpha_{\Phi}^{\sigma \rightarrow \tau}(A) = \lim_{\nu \rightarrow 0} \frac{1}{\nu} (\alpha_{\Phi}^{\sigma \rightarrow \tau + \nu}(A) - \alpha_{\Phi}^{\sigma \rightarrow \tau}(A)) = \lim_{\nu \rightarrow 0} \frac{1}{\nu} \alpha_{\Phi}^{\sigma \rightarrow \tau} (\alpha_{\Phi}^{\tau \rightarrow \tau + \nu}(A) - A) = \alpha_{\Phi}^{\sigma \rightarrow \tau} \circ \delta_{\Phi_{\tau}}(A),$$

holds in norm for arbitrary $\sigma, \tau \in [0, 1]$ and $A \in \mathfrak{A}_{\text{loc}}$.

(4) We will consider the case $\sigma < \tau$, the opposite case is similar. We also observe that Part (1) extends to time-dependent piecewise continuous interactions Φ . Indeed, the continuity was only tacitly used to justify the Riemann integration in the Dyson expansions (20) and (23).

Let $\Phi, \Psi : [0, 1] \rightarrow \mathcal{B}^r$ be piecewise continuous and set $\bar{\epsilon} = r/2 \max(\|\Phi\|_r, \|\Psi\|_r)$. The telescopic expansion

$$\delta_{\Phi_{\tau_1}} \circ \dots \circ \delta_{\Phi_{\tau_n}} - \delta_{\Psi_{\tau_1}} \circ \dots \circ \delta_{\Psi_{\tau_n}} = \sum_{k=1}^n \delta_{\Phi_{\tau_1}} \circ \dots \circ \delta_{\Phi_{\tau_{k-1}}} \circ \delta_{(\Phi-\Psi)_{\tau_k}} \circ \delta_{\Psi_{\tau_{k+1}}} \circ \dots \circ \delta_{\Psi_{\tau_n}},$$

combined with the estimate (22), gives

$$\sup_{\tau_1, \dots, \tau_n \in [0, 1]} \|\delta_{\Phi_{\tau_1}} \circ \dots \circ \delta_{\Phi_{\tau_n}}(A) - \delta_{\Psi_{\tau_1}} \circ \dots \circ \delta_{\Psi_{\tau_n}}(A)\| \leq 2n \|A\| e^{r|\text{supp}(A)|} \bar{\epsilon}^{-n+1} \frac{\|\Phi - \Psi\|_r}{r} n!,$$

for $A \in \mathfrak{A}_{\text{loc}}$. Thus, it follows from the Dyson expansion (23) that

$$\sup_{|\tau - \sigma| \leq \bar{\epsilon}/2} \|\alpha_{\Phi}^{\sigma \rightarrow \tau}(A) - \alpha_{\Psi}^{\sigma \rightarrow \tau}(A)\| \leq \frac{4\bar{\epsilon}}{r} \|A\| e^{r|\text{supp}(A)|} \|\Phi - \Psi\|_r.$$

By density/continuity, we derive that

$$\lim_{\Psi \rightarrow \Phi} \|\alpha_{\Phi}^{\sigma \rightarrow \tau}(A) - \alpha_{\Psi}^{\sigma \rightarrow \tau}(A)\| = 0 \quad (26)$$

for all $A \in \mathfrak{A}$ and $|\tau - \sigma| \leq \bar{\epsilon}/2$. Finally, a telescopic expansion analogous to (25) and with $\max_k (|\tau_{k+1} - \tau_k|) < \bar{\epsilon}/2$ gives

$$\|\alpha_{\Phi}^{\sigma \rightarrow \tau}(A) - \alpha_{\Psi}^{\sigma \rightarrow \tau}(A)\| \leq \sum_{k=0}^n \|(\alpha_{\Phi}^{\tau_k \rightarrow \tau_{k+1}} - \alpha_{\Psi}^{\tau_k \rightarrow \tau_{k+1}})(A_k)\|,$$

with $A_k = \alpha_{\Phi}^{\tau_{k+1} \rightarrow \tau_{k+2}} \circ \dots \circ \alpha_{\Phi}^{\tau_n \rightarrow \tau}(A)$, from which we conclude that (26) holds for all $A \in \mathfrak{A}$ and all $\sigma, \tau \in [0, 1]$.

To complete the proof, given the continuous time-dependent interaction Φ , set

$$\Phi_t^{(N)} = 1_{[0, v_0]}(t) \Phi(t) + \sum_{k=0}^{N-1} 1_{[v_k, v_{k+1}]}(t) \Phi_{v_k} + 1_{[v_N, 1]}(t) \Phi_t,$$

where 1_I denotes the indicator function of the set I . Since

$$\sup_{t \in [0, 1]} \|\Phi_t - \Phi_t^{(N)}\|_r \leq \sup_{|t-s| < \xi_N} \|\Phi_t - \Phi_s\|_r,$$

the uniform continuity of the map $[0, 1] \ni \tau \mapsto \Phi_{\tau} \in \mathcal{B}^r$ implies

$$\lim_{N \rightarrow \infty} \|\Phi - \Phi^{(N)}\|_r = 0.$$

Thus, the claim follows from (26) and the fact that

$$\alpha_{\Phi^{(N)}}^{\sigma \rightarrow \tau}(A) = \alpha_{\Phi_{v_0}}^{\xi_N} \circ \dots \circ \alpha_{\Phi_{v_{N-1}}}^{\xi_N}(A).$$

A Glossary of terms

More details can be found in [JPT23].

- \mathfrak{A} : the C^* -algebra of a spin system on \mathbb{Z}^d , or equivalently, the gauge-invariant sector of the fermionic CAR algebra on $\ell^2(\mathbb{Z}^d)$.

- \mathcal{F} : the set of all finite subsets of \mathbb{Z}^d .
- Φ : an interaction, namely a translation invariant family $\{\Phi(X)\}_{X \in \mathcal{F}}$ of self-adjoint elements of \mathfrak{A} with $\text{supp}(\Phi) = X$.
- \mathcal{B}^r : the Banach space of interactions satisfying $\|\Phi\|_r = \sum_{X \ni 0} e^{r(|X|-1)} \|\Phi(X)\| < \infty$, ($r > 0$).
- $\mathbb{Z}^d \ni x \mapsto \varphi^x$: the group action of \mathbb{Z}^d on \mathfrak{A} .
- $\mathcal{S}_1(\mathfrak{A})$: the set of translation invariant states on \mathfrak{A} .
- $s(\nu)$: the specific entropy of a state $\nu \in \mathcal{S}_1(\mathfrak{A})$.
- $s(\nu|\omega)$: the specific relative entropy of two states $\nu, \omega \in \mathcal{S}_1(\mathfrak{A})$.
- $P(\Phi)$: the pressure of the interaction Φ .
- $E_\Phi = \sum_{X \ni 0} |X|^{-1} \Phi(X)$, so that $\nu(E_\Phi)$ is the expected specific energy of the interaction Φ in the state $\nu \in \mathcal{S}_1(\mathfrak{A})$.
- $\mathcal{S}_{\text{eq}}(\Phi) = \{\nu \in \mathcal{S}_1(\mathfrak{A}) \mid P(\Phi) = s(\nu) - \nu(E_\Phi)\}$: the set of equilibrium states for Φ (from the Gibbs variational principle).
- $\text{WG}(\Phi)$: the set of weak Gibbs states for Φ , see [JPT23, Section 2.2].
- α_Φ : the Heisenberg dynamics generated by the (time-independent) interaction Φ .
- Physical equivalence: two interactions $\Phi, \Psi \in \mathcal{B}^r$ are physically equivalent iff $\alpha_\Phi = \alpha_\Psi$, see [JPT23, Theorem 2.7].

References

- [AE99] AVRON, J. E. and ELGART, A.: Adiabatic theorem without a gap condition. *Commun. Math. Phys.* **203**, 445–463 (1999), [DOI:10.1007/s002200050620].
- [ASF07] ABOU SALEM, W. K. and FRÖHLICH, J.: Status of the fundamental laws of thermodynamics. *J. Stat. Phys.* **126**, 1045–1068 (2007), [DOI:10.1007/s10955-006-9222-8].
- [ASY87] AVRON, J. E., SEILER, R. and YAFFE, L. G.: Adiabatic theorems and applications to the quantum Hall effect. *Commun. Math. Phys.* **110**, 33–49 (1987), [DOI:10.1007/bf01209015].
- [BDRF17] BACHMANN, S., DE ROECK, W. and FRAAS, M.: Adiabatic theorem for quantum spin systems. *Phys. Rev. Lett.* **119**, 060201 (2017), [DOI:10.1103/PhysRevLett.119.060201].
- [BDRF18] ———: The adiabatic theorem and linear response theory for extended quantum systems. *Commun. Math. Phys.* **361**, 997–1027 (2018), [DOI:10.1007/s00220-018-3117-9].
- [BF28] BORN, M. and FOCK, V.: Beweis des Adiabatenatzes. *Z. Phys.* **51**, 165–169 (1928), [DOI:10.1007/bf01343193].
- [BFJP] BENOIST, T., FRAAS, M., JAKŠIĆ, V. and PILLET, C.-A.: Adiabatic theorem in quantum statistical mechanics. In preparation.

- [BR81] BRATTELI, O. and ROBINSON, D. W.: *Operator Algebras and Quantum-Statistical Mechanics. II*. Texts and Monographs in Physics, Springer-Verlag, New York-Berlin, 1981, [DOI:10.1007/978-3-662-03444-6].
- [DJP03] DEREZIŃSKI, J., JAKŠIĆ, V. and PILLET, C.-A.: Perturbation theory of W^* -dynamics, Liouvilleans and KMS-states. *Rev. Math. Phys.* **15**, 447–489 (2003), [DOI:10.1142/S0129055X03001679].
- [GLMP22] GREENBLATT, R. L., LANGE, M., MARCELLI, G. and PORTA, M.: Adiabatic evolution of low-temperature many-body systems. Preprint (2022), [DOI:10.48550/arXiv.2211.16836].
- [Isr79] ISRAEL, R. B.: *Convexity in the Theory of Lattice Gases*. Princeton Series in Physics, Princeton University Press, Princeton, N.J., 1979.
- [JP] JAKŠIĆ, V. and PILLET, C.-A.: Adiabatic theorem for KMS states. Unpublished.
- [JP91] JOYE, A. and PFISTER, C.-E.: Exponentially small adiabatic invariant for the Schrödinger equation. *Commun. Math. Phys.* **140**, 15–41 (1991), [DOI:10.1007/bf02099288].
- [JP03] JAKŠIĆ, V. and PILLET, C.-A.: A note on the entropy production formula. In *Advances in differential equations and mathematical physics (Birmingham, AL, 2002)*, *Contemp. Math.*, vol. 327, Amer. Math. Soc., Providence, RI, 2003, p. 175–180, [DOI:10.1090/conm/327/05813].
- [JP14] ———: A note on the Landauer principle in quantum statistical mechanics. *J. Math. Phys.* **55**, 075210, 21 (2014), [DOI:10.1063/1.4884475].
- [JPT23] JAKŠIĆ, V., PILLET, C.-A. and TAUBER, C.: Approach to equilibrium in translation invariant quantum systems: some structural results. Preprint (2023), [DOI:10.48550/arXiv.2204.00440].
- [Kat50] KATO, T.: On the adiabatic theorem of quantum mechanics. *J. Phys. Soc. Jpn.* **5**, 435–439 (1950), [DOI:10.1143/JPSJ.5.435].
- [LR68] LANFORD, O. E. and ROBINSON, D. W.: Statistical mechanics of quantum spin systems. III. *Commun. Math. Phys.* **9**, 327–338 (1968), [DOI:10.1007/bf01654286].
- [Nen93] NENCIU, G.: Linear adiabatic theory. Exponential estimates. *Commun. Math. Phys.* **152**, 479–496 (1993), [DOI:10.1007/bf02096616].
- [NSY19] NACHTERGAELE, B., SIMS, R. and YOUNG, A.: Quasi-locality bounds for quantum lattice systems. I. Lieb-Robinson bounds, quasi-local maps, and spectral flow automorphisms. *J. Math. Phys.* **60**, 061101 (2019), [DOI:10.1063/1.5095769].
- [OP93] OHYA, M. and PETZ, D.: *Quantum Entropy and its Use*. Texts and Monographs in Physics, Springer-Verlag, Berlin, 1993, [DOI:10.1007/978-3-642-57997-4].
- [Teu01] TEUFEL, S.: A note on the adiabatic theorem without gap condition. *Lett. Math. Phys.* **58**, 261–266 (2001), [DOI:10.1023/A:1014556511004].