



Random System of lines in the Euclidean plane E2

Giuseppe Caristi

► To cite this version:

Giuseppe Caristi. Random System of lines in the Euclidean plane E2. Annales de l'ISUP, 2009, LIII (1), pp.53-58. hal-03634193

HAL Id: hal-03634193

<https://hal.science/hal-03634193v1>

Submitted on 7 Apr 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

*Pub. Inst. Stat. Univ. Paris
LIII, fasc. 1, 2009, 53 à 58*

Random system of lines in the Euclidean plane E_2

Giuseppe Caristi

Abstract. In this paper we consider a random variable arising from a problem of geometrical intersection between a fixed convex body K and a system of random lines in E_2 .

M.S.C. 2000: 60D05,52A22.

Key words: Geometric probability, stochastic geometry, random sets, random convex sets, integral geometry.

1 Introduction

Let E_2 be the Euclidean plane and let K be a convex non empty and bounded domain of area S_K and with boundary ∂K of length L . We consider a family F of random, uniformly distributed n -lines $\{G_1, \dots, G_n\}$ with $n \geq 2$. We assume that if $G_h, G_k \in F$, then $G_h \cap G_k \neq \emptyset$. It is possible that this points belongs to K or not. In this way we have a random variable $X_{(n,K)}$. In this paper we give the following result

Theorem 1. *The expression of the mean value $\mathbf{E}(X_{(n,K)})$, the k -moments $\mathbf{E}(X_{(n,K)}^k)$ and the variance $\sigma^2(X_{(n,K)})$ of the random variable $X_{(n,K)}$ can be calculated as follows*

$$\mathbf{E}(X_{(n,K)}) = \alpha \pi \frac{S_K}{L^2}, \quad \mathbf{E}(X_{(n,K)}^k) = \left[\sum_{J_1+...+J_\alpha=k} \frac{k!}{J_1! \dots J_\alpha!} \right] \frac{\pi S_K}{L^2},$$

where

$$\alpha = \frac{n(n-1)}{2},$$

and k is a positive integer.

Other results about the computation of the variance are investigated in [2] and an extension in the 3-dimensional Euclidean space of the same problem is studied in [6].

2 Main Results

Let N be the set of natural numbers, $n \geq 2$ a fixed natural integer, $\{G_1, \dots, G_n\}$ and K as in introduction. We can state the following

Theorem 2. *Let us consider the random variable $X_{(n,K)}$. Then*

$$\mathbf{E}(X_{(n,K)}) = \alpha \pi \frac{S_K}{L^2},$$

where $\alpha = \frac{n(n-1)}{2}$, L is the length of ∂K and S_K is the area of the domain defined by K .

Proof. It is easy to see that, denoting with L the length of ∂K and dG the elementary measure of the lines in the Euclidean plane E_2 , we have

$$\int_{\{G \cap K \neq \emptyset\}} dG = L.$$

Since G_1, \dots, G_n are stochastically independent, we get

$$\int_{\{G \cap K \neq \emptyset\}} dG_1 \wedge \dots \wedge dG_n = L^n.$$

If we consider the lines G_1, \dots, G_n , then the intersection points might belong to K or not. Hence we obtain a random variable which we denote by $X_{(n,K)}$.

In order to compute the variance, we have the following integral

$$\int_{\{G \cap K \neq \emptyset\}} X_{(n,K)} dG_1 \wedge \dots \wedge dG_n$$

We define the application $\epsilon_{hk} = 1$ if $G_h \cap G_k \in K$ (with $h \neq k = 1, \dots, n$) and zero otherwise. Then

$$X_{(n,K)} = \sum_{h,k=1} \epsilon_{hk}.$$

Further, let us consider

$$I_2 := \int_{\{G_h, G_k \cap K \neq \emptyset\}} \epsilon_{hk} dG_h \wedge dG_k.$$

If $(G_h \cap E_k) \in K$, then we have $\epsilon_{hk} = 1$. We denote with λ_k the chord intercepted by G_k on K (and its length).

We have

$$\int_{\{G_h, G_k \cap K \neq \emptyset\}} \epsilon_{hk} dG_h \wedge dG_k = \int_{\{G_k \cap K \neq \emptyset\}} \left(\int_{\{G_h \cap K \neq \emptyset\}} dG_h \right) dG_k,$$

but it is well known that

$$\int_{\{G_h \cap \lambda_k \neq \emptyset\}} dG_h = \lambda_k,$$

$$\int_{\{G_k \cap K \neq \emptyset\}} \lambda_k dG_k = \pi S_K.$$

Moreover,

$$\begin{aligned} \int_{\{G \cap K \neq \emptyset\}} X_{(n,K)} dG_1 \wedge \dots \wedge dG_n &= \int_{\{G \cap K \neq \emptyset\}} \sum_{h,k=1} \epsilon_{hk} dG_1 \wedge \dots \wedge dG_n = \\ &\pi S_K L^{n-2} + \dots + \pi S_K L^{n-2}. \end{aligned}$$

Taking into account that the number of the different sets $\{G_h, G_k\}$ is (the binomial coefficient)

$$\alpha = \frac{n(n-1)}{2},$$

we have

$$\int_{\{G \cap K \neq \emptyset\}} X_{(n,K)} dG_1 \wedge \dots \wedge dG_n = \alpha \pi S_K L^{n-2}.$$

By definition,

$$\mathbf{E}(X_{(n,K)}) := \frac{\int_{\{G \cap K \neq \emptyset\}} X_{(n,K)} dG_1 \wedge \dots \wedge dG_n}{\int_{\{G \cap K \neq \emptyset\}} dG_1 \wedge \dots \wedge dG_n},$$

and hence $\mathbf{E}(X_{(n,K)}^2) = \alpha \pi \frac{S_K}{L^2}$. □

Now we compute

$$\mathbf{E}(X_{(n,K)}^2) = \frac{\int_{\{G \cap K \neq \emptyset\}} X_{(n,K)}^2 dG_1 \wedge \dots \wedge dG_n}{\int_{\{G \cap K \neq \emptyset\}} dG_1 \wedge \dots \wedge dG_n}.$$

We put

$$J = \int_{\{G \cap K \neq \emptyset\}} X_{(n,K)}^2 dG_1 \wedge \dots \wedge dG_n$$

We can prove that

$$X_{(n,K)}^2 = \left(\sum_{h,k=1} \epsilon_{hk} \right)^2,$$

and then

$$X_{(n,K)}^2 = \sum_{h,k=1} \epsilon_{hk}^2 + 2 \sum_{(h,k) \neq (s,n)} \epsilon_{hk} \epsilon_{sn}.$$

With this observations we can compute the integral

$$J = \int_{\{G \cap K \neq \emptyset\}} \left(\sum_{h,k=1} \epsilon_{hk}^2 + 2 \sum_{(h,k) \neq (s,n)} \epsilon_{hk} \epsilon_{sn} \right) dG_1 \wedge \dots \wedge dG_n.$$

Considering that

$$\int_{\{G \cap K \neq \emptyset\}} \epsilon_{hk}^2 dG_h \wedge dG_k = \int_{\{G_k \cap K \neq \emptyset\}} \left(\int_{\{G_h \cap \lambda_k \neq \emptyset\}} dG_h \right) dG_k,$$

where λ_k is the chord intercepted by G_k on K , we obtain

$$J = \pi S_K L^{n-2} + \dots + \pi S_K L^{n-2} + 2\pi S_K L^{n-2} + \dots + 2\pi S_K L^{n-2}.$$

Taking into account that the number of different sets $\{G_h, G_k\}$ is α and that G_1, \dots, G_2 are independent, we infer

$$\mathbf{E}(X_{(n,K)}^2) = \frac{\pi S_K \alpha L^{n-2}}{L^n} = \frac{\pi S_K \alpha^2}{L^2}.$$

We obtain

Theorem 3. Let us consider the random variable $X_{(n,K)}$. Hence

$$\sigma^2(X_{(n,K)}) = \frac{\pi S_K}{L^2} \left(1 - \frac{\pi S_K}{L^2}\right) \alpha^2,$$

where $\alpha = \frac{n(n-1)}{2}$, L is the length of ∂K and S_K is the area of the domain defined by K .

Moreover, we note that

$$X_{(n,K)}^k = \left(\sum_{h,k=1}^{\alpha} \epsilon_{hk} \right)^k,$$

implies

$$\mathbf{E}(X_{(n,K)}^k) = \left[\sum_{J_1+\dots+J_\alpha=k} \frac{k!}{J_1! \dots J_\alpha!} \right] \frac{\pi S_K}{L^2}.$$

3 Applications

1. As first case we consider in the plane a square Q of side a . We can compute the following values

$$\mathbf{E}(X_{(n,Q)}) = \frac{\alpha \pi}{16} \approx 0,196345\alpha.$$

$$\mathbf{E}(X_{(n,Q)}^k) = \left[\sum_{J_1+\dots+J_\alpha=k} \frac{k!}{J_1! \dots J_\alpha!} \right] \frac{\pi}{16}.$$

and the variance is

$$\sigma^2 X_{(n,Q)} = \frac{\pi}{16} \left(1 - \frac{\pi}{16}\right) \alpha^2 \approx 0,15779\alpha^2.$$

2. Taking in plane a rectangle R of sides a and b we have

$$\mathbf{E}(X_{(n,R)}) = \frac{\alpha\pi ab}{4(a+b)^2},$$

$$\mathbf{E}(X_{(n,R)}^k) = \left[\sum_{J_1+\dots+J_\alpha=k} \frac{k!}{J_1! \dots J_\alpha!} \right] \frac{\pi}{4(a+b)^2},$$

and the variance

$$\sigma^2 X_{(n,Q)} = \frac{\pi}{4(a+b)^2} \left(1 - \frac{\pi}{4(a+b)^2} \right) \alpha^2.$$

3. Let C be a circle of radius δ . We have

$$\mathbf{E}(X_{(n,C)}^k) = \frac{\alpha}{4},$$

$$\mathbf{E}(X_{(n,C)}^k) = \left[\sum_{J_1+\dots+J_\alpha=k} \frac{k!}{J_1! \dots J_\alpha!} \right] \frac{\alpha}{4}$$

and the variance is

$$\sigma^2(X_{(n,C)}) = \frac{\pi}{4} \left(1 - \frac{\alpha}{4} \right) \alpha^2 \approx 0,16855\alpha^2.$$

4. As last case we consider an equilateral triangle T of side a , obtaining,

$$\mathbf{E}(X_{(n,T)}^k) = \frac{\alpha\pi\sqrt{3}}{18} \approx 0,3023\alpha,$$

$$\mathbf{E}(X_{(n,T)}^k) = \left[\sum_{J_1+\dots+J_\alpha=k} \frac{k!}{J_1! \dots J_\alpha!} \right] \frac{\pi\sqrt{3}}{18},$$

and for the variance the expression

$$\sigma^2(X_{(n,T)}) = \frac{\pi\sqrt{3}}{18} \left(1 - \frac{\pi\sqrt{3}}{18} \right) \alpha^2 \approx 0,2109\alpha^2.$$

Remark 4. We observe that in examples 1, 3, 4 the functions are independent of the dimension of the convex body.

References

- [1] W. Blaschke, *Vorlesungen über Integralgeometrie*, III Auflage, V.E.B. Deutscher Verlag der Wiss., Berlin 1955.
- [2] G. Caristi and G. Molica Bisci, *On the variance associated to a family of ovaloids in the Euclidean Space E_3* , Bollettino U.M.I. (8) 10-B (2007), 87-98.

- [3] E. Czuber, *Zur Theorie der geometrischen Wahrscheinlichkeiten*, Sitz. Akad. Wiss. Wien 90 (1884), 719-742.
- [4] R. Deltheil, *Probabilités géométriques*, Gauthier-Villars, Paris 1926.
- [5] A. Duma and M. Stoka, *Probabilità geometriche per sistemi di piani nello spazio euclideo E_3* , Rend. Sem. Mat. Messina 7 (2000), 5-11.
- [6] G. Molica Bisci, *Random systems of planes in the Euclidean space E_3* , Atti Acc. Scienze di Torino 2005, 43-48.
- [7] H. Poincaré, *Calcul des probabilités*, Ed. 2, Carré, Paris, 1912.

Author's address:

Giuseppe Caristi
 University of Messina,
 75 Via dei Verdi, 98122 Messina, Italy.
 E-mail: gcaristi@unime.it

3 Applications