



**HAL**  
open science

# Problems of Buffon type with multiple intersections for lattices of isoscele trapeziums

Giuseppe Caristi

► **To cite this version:**

Giuseppe Caristi. Problems of Buffon type with multiple intersections for lattices of isoscele trapeziums. *Annales de l'ISUP*, 2007, LI (3), pp.17-26. hal-03635419

**HAL Id: hal-03635419**

**<https://hal.science/hal-03635419v1>**

Submitted on 8 Apr 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Problems of Buffon type with multiple intersections for lattices of isoscele trapeziums

Giuseppe Caristi

## Abstract

In this paper we study problems of Buffon type with multiple intersections for a lattice of isoscele trapeziums, having angles  $\frac{\pi}{3}$  and  $\frac{2\pi}{3}$ , and body test a line segment.

AMS2000 subject Classification: Geometric Probability, stochastic geometry, random sets, random convex sets and integral geometry.

AMS Classification: 60D05, 52A22.

## 1 Preliminaries

In the two works [1] and [2] are studied problems of Buffon type with multiple intersections for lattices of parallelograms and of equilater triangles taking as test body a line segment of constant length  $s$ . For the same test body we consider a lattice  $\mathfrak{R}$  with elementary tile a trapezium  $\mathfrak{T}$  of parallels sides  $A$  and  $A - B$  and of sides  $B$  and  $B$ , that form an angle equal to  $\frac{\pi}{3}$  with the side  $A$ .




Fig.1

We want to determine the probability of multiple intersections of the test body  $s$  with the sides of lattice  $\mathfrak{R}$ , that is the probability  $p_1, p_2, \dots$  that the test body intersects respective one time, two times, ... the sides of lattice.

Because of the different relations between the geometric objects of the trapezium  $\mathfrak{S}$ , that is between sides  $A$ ,  $B$ , the heights  $h = \frac{B\sqrt{3}}{2}$  and  $H = \frac{A\sqrt{3}}{2}$ , and the diagonal  $\delta = \sqrt{A^2 - AB + B^2}$  we must consider the following cases:

Case 1:  $B \leq A \leq \frac{2B}{\sqrt{3}}$ . In this case we have:

$$A - B \leq h \leq H \leq B < \delta < A;$$

Case 2:  $\frac{2B}{\sqrt{3}} \leq A \leq \left(1 + \frac{\sqrt{3}}{2}\right)B$ . In this case we have:

$$A - B \leq h \leq B \leq H < \delta < A;$$

Case 3:  $\left(1 + \frac{\sqrt{3}}{2}\right)B \leq A \leq 2B$ . In this case we have:

$$h \leq A - B \leq B \leq H \leq \delta < A;$$

Case 4:  $2B \leq A \leq (4 + 2\sqrt{3})B$ . In this case we have:

$$h \leq B \leq A - B \leq H \leq \delta < A;$$

Case 5:  $(4 + 2\sqrt{3})B \leq A$ . In this case we have:

$$h \leq B \leq H \leq A - B < \delta < A.$$

The length needle  $s$  is small (in the given sense in [3]) respectively if and only if  $l \leq A - B$  in the cases 1 and 2 and  $l \leq B\frac{\sqrt{3}}{2}$  in the cases 3, 4 and 5.

## 2 The length needle

We consider that  $l$  is small respect to  $\mathfrak{R}$ . Under this hypothesis we don't have to distinguish between the 5 previous cases. The test body can intersect one time, two times or three times the sides of lattice; therefore we want to calculate the probabilities  $p_1, p_2, p_3$ .

**Theorem 1** *Let  $s$  a line segment of length  $l$  small regarding the lattice  $\mathfrak{R}$ , uniformly distributed in a bounded region of the Euclidean plane. The probabilities that  $s$  intersects one time, two times or three times the sides of  $\mathfrak{R}$  are respectively*

$$p_1 = \frac{4}{\pi\sqrt{3}} \cdot \frac{2A+B}{2A-B} \cdot \frac{l}{B} - \left( \frac{7}{2\pi\sqrt{3}} + \frac{5}{18} \right) \cdot \frac{l^2}{(2A-B)B}; \quad (1)$$

$$p_2 = \left( \frac{4}{9} + \frac{2}{\pi\sqrt{3}} \right) \cdot \frac{l^2}{(2A-B)B} \quad (2)$$

$$p_3 = \left( \frac{1}{\pi\sqrt{3}} - \frac{1}{9} \right) \cdot \frac{l^2}{(2A-B)B}. \quad (3)$$

**Proof.** Let  $\varphi$  the angle formed from  $s$  with the the straight parallels of the lattice, we observe that for reasons of simmetry is sufficient to consider the direction of  $\varphi$  in the interval  $[0, \frac{\pi}{2}]$ . Let  $M$  the set of the length segments  $l$  that have the barycentre  $m$  in the trapezium  $\mathfrak{S}$  and the arbitrary direction  $\varphi \in [0, \frac{\pi}{2}]$ . For  $k = 0, 1, 2, 3$  let  $\aleph_k$  set of segments  $s$  length  $l$  that have the barycentre  $m$  in the trapezium  $\mathfrak{S}$ , the direction  $\varphi$  such that che  $s$  intersects  $\aleph$   $k$  times. Therefore  $M = \aleph_1 \cup \aleph_2 \cup \aleph_3$  is a three-dimensional set and if  $\varphi$  is the direction  $s$ , then  $s \in \aleph_k$  if and only if the centre  $m$  of  $s$  is founded in a set of  $\mathfrak{S}$ , that will be denote by  $\mathfrak{S}_k(\varphi)$ . If  $\mu$  is the Lebesgue measure in the Euclidean space, then Stoka's formula gives to us

$$p_k = \frac{\mu(\aleph_k)}{\mu(M)} = \frac{\int_0^{\frac{\pi}{2}} \text{Area } \mathfrak{S}_k(\varphi) d\varphi}{\frac{\pi}{2} \text{Area } \mathfrak{S}} = \frac{4 \int_0^{\frac{\pi}{2}} \text{Area } \mathfrak{S}_k(\varphi) d\varphi}{\pi (AB\sqrt{3} - B^2\frac{\sqrt{3}}{2})}, \quad (4)$$

hence  $\text{Area } \mathfrak{S} = \frac{1}{2}(2A - B) \cdot h = \frac{AB\sqrt{3}}{2} - \frac{B^2\sqrt{3}}{4}$ .

For  $\varphi \in [0, \frac{\pi}{3}[$  the set  $\mathfrak{S}_3(\varphi)$  is empty; the set  $\mathfrak{S}_2(\varphi)$  is the union of two parallelograms and of two trapezioms with height  $\frac{1}{2} \sin \varphi$  and sides  $\frac{l}{\sqrt{3}} \sin(\frac{\pi}{3} - \varphi) =: l_1(\varphi)$ ,  $\frac{l}{\sqrt{3}} \sin(\frac{\pi}{3} + \varphi) =: l_2(\varphi)$ , like in Fig. 2.




Fig.2

Then we obtain:

$$\begin{aligned} \text{Area } \mathfrak{S}_2(\varphi) &= \left( l_1(\varphi) + l_2(\varphi) + 2 \cdot \frac{l_1(\varphi) + l_2(\varphi)}{2} \right) \frac{l}{2} \sin \varphi = \\ &= 2 \frac{l}{\sqrt{3}} \left( \sin\left(\frac{\pi}{3} - \varphi\right) + \sin\left(\frac{\pi}{3} + \varphi\right) \right) \frac{l}{2} \sin \varphi = l^2 \sin \varphi \cos \varphi. \end{aligned} \quad (5)$$

The set  $\mathfrak{S}_1(\varphi)$  is composed from four parallelograms; the two "horizontal" have the height  $\frac{1}{2} \sin \varphi$  and horizontal sides  $A - 2l_2(\varphi)$  respectively  $A - B -$

$2l_1(\varphi)$ ; the two "oblique" have the oblique sides  $B - \frac{2}{\sqrt{3}}l \sin \varphi$  and the height  $\frac{\sqrt{3}}{2}l_1(\varphi)$  respectively  $\frac{\sqrt{3}}{2}l_2(\varphi)$ . Then:

$$\begin{aligned} \text{Area } \mathfrak{S}_1(\varphi) &= 2A - B - 2(l_1(\varphi) + l_2(\varphi)) \frac{l}{2} \sin \varphi + \\ &\quad \frac{\sqrt{3}}{2} \left( B - \frac{2l}{\sqrt{3}} \sin \varphi \right) (l_1(\varphi) + l_2(\varphi)) = \\ &\quad Al \sin \varphi + Bl \sin \left( \frac{\pi}{3} - \varphi \right) - 2l^2 \sin \varphi \cos \varphi. \end{aligned} \quad (6)$$

For  $\varphi \in [0, \frac{\pi}{3}[$  and  $\varphi \in ]\frac{\pi}{3}, \frac{\pi}{2}]$  we have two different geometric situation as we can see in Fig. 2 and 3.




Fig.3

For  $\varphi \in ]\frac{\pi}{3}, \frac{\pi}{2}]$  the set  $\mathfrak{S}_3(\varphi)$  is composed by a little equilateral triangle with side  $l_1^*(\varphi) = \frac{l}{\sqrt{3}} \sin(\varphi - \frac{\pi}{3})$ , by a trapezium that have the sides  $l \cos \varphi$  e  $l_2(\varphi) = \frac{l}{\sqrt{3}} \sin(\varphi + \frac{\pi}{3})$ , and height  $\frac{l}{2} \sin(\varphi - \frac{\pi}{3})$ , and by two triangle that entirely forms a parallelogram with sides  $l_2(\varphi)$  and  $l_1^*(\varphi)$  having height  $\frac{l}{2} \sin(\varphi - \frac{\pi}{3})$ .

$$\begin{aligned} \text{Area } \mathfrak{S}_3(\varphi) &= \frac{l^2}{4\sqrt{3}} \sin^2 \left( \varphi - \frac{\pi}{3} \right) + \frac{l^2}{4} \sin \left( \varphi - \frac{\pi}{3} \right) \left[ \cos \varphi + \frac{1}{\sqrt{3}} \sin \left( \varphi + \frac{\pi}{3} \right) \right] + \\ &\quad \frac{l^2}{2\sqrt{3}} \sin \left( \varphi - \frac{\pi}{3} \right) \sin \left( \varphi + \frac{\pi}{3} \right) = \frac{l^2}{4\sqrt{3}} \sin \left( \varphi - \frac{\pi}{3} \right) \sin \left( \varphi - \frac{\pi}{3} \right) + \\ &\quad \frac{l^2}{4\sqrt{3}} \sin \left( \varphi - \frac{\pi}{3} \right) \sin \left( \varphi + \frac{\pi}{3} \right) + \frac{l^2}{2\sqrt{3}} \sin \left( \varphi - \frac{\pi}{3} \right) \sin \left( \varphi + \frac{\pi}{3} \right) = \\ &\quad \frac{l^2}{\sqrt{3}} \sin \left( \varphi - \frac{\pi}{3} \right) \sin \left( \varphi + \frac{\pi}{3} \right) = -\frac{l^2}{2\sqrt{3}} \left[ \frac{1}{2} + \cos 2\varphi \right]. \end{aligned} \quad (7)$$

The set  $\mathfrak{S}_2(\varphi)$  is the union of an equilateral of side  $l \cos \varphi$ , a trapezium with parallel sides  $l_1^*(\varphi)$  and  $2l_1^*(\varphi)$  and height  $\frac{1}{2} \sin(\varphi - \frac{\pi}{3})$ , of an equilateral triangle with side  $l_2(\varphi)$ , and of a pentagon that with a little equilateral triangle with side  $l_1^*(\varphi)$  they form a parallelogram with sides  $l_2(\varphi)$ ,  $l_3(\varphi) := \frac{l}{\sqrt{3}} \sin \varphi$  and angles  $\frac{\pi}{3}$ ,  $\frac{2\pi}{3}$ .

In the figure 3 we see again that the set  $\mathfrak{S}_1(\varphi)$  is the union of a pentagon and of three parallelograms with angles  $\frac{\pi}{3}$ ,  $\frac{2\pi}{3}$ . The sides of the parallelogram are respectively  $l_1^*(\varphi)$  and  $B - 2l_3(\varphi)$ ,  $A - B$  and  $\frac{l}{2}$ , and  $l_2(\varphi)$  and  $B - 2l_3(\varphi)$ . The pentagon is the difference between a trapezium with parallel sides  $A - [l_1^*(\varphi) + l_2(\varphi)]$  and  $A - 2[l_1^*(\varphi) + l_2(\varphi)]$  and height  $\frac{l}{2} \sin \varphi$  and a little equilateral triangle with sides  $l_1^*(\varphi)$ . Consequently:

$$\begin{aligned} \text{Area } \mathfrak{S}_2(\varphi) &= \left[ \frac{\sqrt{3}}{4} \cos^2 \varphi + \frac{1}{\sqrt{3}} \sin^2 \left( \varphi - \frac{\pi}{3} \right) + \frac{1}{4\sqrt{3}} \sin^2 \left( \varphi + \frac{\pi}{3} \right) \right. \\ &\quad \left. + \frac{1}{2\sqrt{3}} \sin \varphi \sin \left( \varphi + \frac{\pi}{3} \right) - \frac{1}{4\sqrt{3}} \sin^2 \left( \varphi - \frac{\pi}{3} \right) \right] l^2 = \\ &\quad \left[ \frac{1}{\sqrt{3}} + \frac{\sqrt{3}}{8} \cos 2\varphi - \frac{\sqrt{3}}{8} \cos \left( 2\varphi - \frac{2\pi}{3} \right) - \right. \\ &\quad \left. \frac{1}{8\sqrt{3}} \cos \left( 2\varphi + \frac{2\pi}{3} \right) - \frac{1}{4\sqrt{3}} \cos \left( 2\varphi + \frac{\pi}{3} \right) \right] l^2 \quad (8) \end{aligned}$$

and

$$\begin{aligned} \text{Area } \mathfrak{S}_1(\varphi) &= (A - B) \frac{l}{2} \sin \varphi + \\ &\quad \left( B \frac{\sqrt{3}}{2} - l \sin \varphi \right) \left( \frac{l}{\sqrt{3}} \sin \left( \varphi - \frac{\pi}{3} \right) + \frac{l}{\sqrt{3}} \sin \left( \varphi + \frac{\pi}{3} \right) \right) + \\ &\quad \left[ A - \frac{3}{2} \left( \frac{l}{\sqrt{3}} \sin \left( \varphi - \frac{\pi}{3} \right) + \frac{l}{\sqrt{3}} \sin \left( \varphi + \frac{\pi}{3} \right) \right) \right] \frac{l}{2} \sin \varphi - (l \cos \varphi)^2 \frac{\sqrt{3}}{4} = \\ &\quad Al \sin \varphi - \frac{7}{4\sqrt{3}} l^2 \sin^2 \varphi - \frac{\sqrt{3}}{4} l^2 \cos^2 \varphi = Al \sin \varphi + \frac{\cos 2\varphi}{2\sqrt{3}} l^2 - \frac{5l^2}{4\sqrt{3}}. \quad (9) \end{aligned}$$

We calculate the integrals of these areas:

$$\begin{aligned} \int_0^{\frac{\pi}{2}} \text{Area } \mathfrak{S}_1(\varphi) d\varphi &= \int_0^{\frac{\pi}{3}} \text{Area } \mathfrak{S}_1(\varphi) d\varphi + \int_{\frac{\pi}{3}}^{\frac{\pi}{2}} \text{Area } \mathfrak{S}_1(\varphi) d\varphi = \\ &\quad \left( -Al \cos \varphi + Bl \cos \left( \frac{\pi}{3} - \varphi \right) - l^2 \sin^2 \varphi \right) \Big|_0^{\frac{\pi}{3}} + \left( -Al \cos \varphi + \frac{\sin 2\varphi}{4\sqrt{3}} l^2 - \frac{5l^2}{4\sqrt{3}} \varphi \right) \Big|_{\frac{\pi}{3}}^{\frac{\pi}{2}} = \\ &\quad Al + \frac{1}{2} Bl - \left( \frac{7}{8} + \frac{5\pi}{24\sqrt{3}} \right) l^2. \quad (10) \end{aligned}$$

$$\begin{aligned}
\int_0^{\frac{\pi}{2}} \text{Area } \mathfrak{S}_2(\varphi) d\varphi &= \int_0^{\frac{\pi}{3}} \text{Area } \mathfrak{S}_2(\varphi) d\varphi + \int_{\frac{\pi}{3}}^{\frac{\pi}{2}} \text{Area } \mathfrak{S}_2(\varphi) d\varphi = \\
&= \frac{l^2}{2} \sin^2 \varphi \Big|_0^{\frac{\pi}{3}} + \frac{\pi}{6\sqrt{3}} l^2 + l^2 \left[ \left( \frac{\sqrt{3}}{16} \sin 2\varphi \right) \Big|_{\frac{\pi}{3}}^{\frac{\pi}{2}} - \right. \\
&\quad \left. \frac{\sqrt{3}}{16} \sin \left( 2\varphi - \frac{2\pi}{3} \right) \Big|_{\frac{\pi}{3}}^{\frac{\pi}{2}} - \frac{1}{16\sqrt{3}} \sin \left( 2\varphi + \frac{2\pi}{3} \right) \Big|_{\frac{\pi}{3}}^{\frac{\pi}{2}} - \right. \\
&\quad \left. \left| \frac{1}{8\sqrt{3}} \sin \left( 2\varphi + \frac{\pi}{3} \right) \Big|_{\frac{\pi}{3}}^{\frac{\pi}{2}} \right] = \left( \frac{\pi}{6\sqrt{3}} + \frac{1}{4} \right) l^2, \tag{11}
\end{aligned}$$

$$\begin{aligned}
\int_0^{\frac{\pi}{2}} \text{Area } \mathfrak{S}_3(\varphi) d\varphi &= \int_{\frac{\pi}{3}}^{\frac{\pi}{2}} \text{Area } \mathfrak{S}_3(\varphi) d\varphi = \\
&= -\frac{l^2}{2\sqrt{3}} \left[ \frac{\pi}{12} - \frac{\sqrt{3}}{4} \right] = \left( \frac{1}{8} - \frac{\pi}{24\sqrt{3}} \right) l^2. \tag{12}
\end{aligned}$$

Considering the formulas (4), (10), (11) and (12) we obtain the probabilities (1), (2) e (3). ■

### 3 The not small needle

Now we consider that the needle of length  $l$  is not little respect to  $\mathfrak{R}$ , but minor or equal to  $A$ . For each of the five previous cases we must disintinguish others five undercases, for example in the case 1 the undercases are:  $A - B \leq l \leq h$ ;  $h \leq l \leq H$ ;  $H \leq l \leq B$ ;  $B \leq l \leq \delta$ ;  $\delta \leq l \leq A$ . The cases 1 and 2 have got two identical undercases ( $A - B \leq l \leq h$  e  $\delta \leq l \leq A$ ) and the cases 2 and 3 have got three identical undercases ( $B \leq l \leq H$ ,  $H \leq l \leq \delta$  e  $\delta \leq l \leq A$ ), and so on. We observe that for  $l \leq A$  exactly 16 undercases are different. In order to simplify we will make the calculations on the undercase  $A - B \leq l \leq h$ . In this case we must distinguish between the following situations:

$$i) h \leq 2(A - B); \quad ii) 2(A - B) \leq h.$$

In this last situation two possibilities exist:

$$ii)_1 A - B \leq l \leq 2(A - B); \quad ii)_2 2(A - B) \leq l \leq h.$$

Now we consider the situations *i*) and *ii*)<sub>1</sub>. The interval  $[0, \frac{\pi}{2}]$  is the union of  $[0, \varphi_0]$ ,  $[\varphi_0, \frac{\pi}{3}]$  and  $[\frac{\pi}{3}, \frac{\pi}{2}]$ , where  $\varphi_0 \in [0, \frac{\pi}{3}]$  is defined by the relation

$$\sin\left(\frac{\pi}{3} - \varphi_0\right) = \frac{\sqrt{3}}{2} \cdot \frac{A - B}{l}.$$

So for  $\varphi \in [\varphi_0, \frac{\pi}{3}] \cup [\frac{\pi}{3}, \frac{\pi}{2}]$  we can be useful for the calculations of the areas maked for the little case, it remain to determine *Area*  $\mathfrak{S}_1(\varphi)$ , *Area*  $\mathfrak{S}_2(\varphi)$  and *Area*  $\mathfrak{S}_3(\varphi)$  when  $\varphi \in [0, \varphi_0]$ , and for this we observe the Fig. 4:




Fig.4

*Area*  $\mathfrak{S}_3(\varphi)$  is the area of parallelogram with sides  $\frac{2l}{\sqrt{3}} \sin\left(\frac{\pi}{3} - \varphi\right) - (A - B)$ ,  $\frac{l}{\sqrt{3}} \sin \varphi$  and angles  $\frac{\pi}{3}$ ,  $\frac{2\pi}{3}$ ; therefore we have:

$$\begin{aligned} \text{Area } \mathfrak{S}_3(\varphi) &= \frac{l}{2} \sin \varphi \left[ \frac{2l}{\sqrt{3}} \sin\left(\frac{\pi}{3} - \varphi\right) - (A - B) \right] = \\ &= \frac{l^2}{2\sqrt{3}} \cos\left(\frac{\pi}{3} - 2\varphi\right) - \frac{l^2}{4\sqrt{3}} - \frac{l(A - B)}{2} \sin \varphi, \\ \int_0^{\varphi_0} \text{Area } \mathfrak{S}_3(\varphi) d\varphi &= \frac{l(A - B)}{2} (\cos \varphi_0 - 1) + \\ &+ \frac{l^2}{8} - \frac{l^2}{4\sqrt{3}} \left( \varphi_0 + \sin\left(\frac{\pi}{3} - 2\varphi_0\right) \right). \end{aligned} \quad (13)$$



From (7) and (13) we obtain

$$\int_0^{\frac{\pi}{2}} \text{Area } \mathfrak{S}_3(\varphi) d\varphi = \left( \int_0^{\varphi_0} + \int_{\frac{\pi}{3}}^{\frac{\pi}{2}} \right) \text{Area } \mathfrak{S}_3(\varphi) d\varphi = \frac{l(A-B)}{2} (\cos \varphi_0 - 1) + \frac{l^2}{8} - \frac{l^2}{4\sqrt{3}} \left( \varphi_0 + \sin \left( \frac{\pi}{3} - 2\varphi_0 \right) \right) + l^2 \left( \frac{1}{8} - \frac{\pi}{24\sqrt{3}} \right) = \frac{l(A-B)}{2} (\cos \varphi_0 - 1) + \frac{l^2}{4} - \frac{l^2}{4\sqrt{3}} \left( \frac{\pi}{6} + \varphi_0 + \sin \left( \frac{\pi}{3} - 2\varphi_0 \right) \right). \quad (14)$$

$\mathfrak{S}_2(\varphi)$  is composed from two parallelograms with sides that are respectively  $\frac{l}{\sqrt{3}} \sin \left( \varphi + \frac{\pi}{3} \right)$  and  $A - B - \frac{l}{\sqrt{3}} \sin \left( \frac{\pi}{3} - \varphi \right)$  and the same correspondent height of length  $\frac{l}{2} \sin \varphi$ , from two trapeziums with parallel sides respectively  $\frac{l}{\sqrt{3}} \sin \left( \varphi + \frac{\pi}{3} \right)$ ,  $\frac{l}{\sqrt{3}} \sin \left( \varphi - \frac{\pi}{3} \right)$  and  $A - B + \frac{l}{\sqrt{3}} \sin \left( \varphi + \frac{\pi}{3} \right) - \frac{2l}{\sqrt{3}} \sin \left( \frac{\pi}{3} - \varphi \right)$ ,  $A - B - \frac{l}{\sqrt{3}} \sin \left( \frac{\pi}{3} - \varphi \right)$  and the same correspondent height of length  $\frac{l}{2} \sin \varphi$ , and from an equilateral triangle with side  $\frac{2l}{\sqrt{3}} \sin \left( \frac{\pi}{3} - \varphi \right) - (A - B)$ . It follows that

$$\begin{aligned} \text{Area } \mathfrak{S}_2(\varphi) &= \left[ \frac{l}{\sqrt{3}} \sin \left( \frac{\pi}{3} + \varphi \right) + (A - B) - \frac{l}{\sqrt{3}} \sin \left( \frac{\pi}{3} - \varphi \right) + \frac{l}{2\sqrt{3}} \sin \left( \varphi + \frac{\pi}{3} \right) \right. \\ &+ \left. \frac{l}{2\sqrt{3}} \sin \left( \frac{\pi}{3} - \varphi \right) + (A - B) + \frac{l}{2\sqrt{3}} \sin \left( \varphi + \frac{\pi}{3} \right) - \frac{3l}{2\sqrt{3}} \sin \left( \frac{\pi}{3} - \varphi \right) \right] \frac{l}{2} \sin \varphi + \\ &\frac{\sqrt{3}}{4} \left[ \frac{2l}{\sqrt{3}} \sin \left( \frac{\pi}{3} - \varphi \right) - (A - B) \right]^2 = \frac{l^2}{\sqrt{3}} \left[ \sin \left( \varphi + \frac{\pi}{3} \right) - \sin \left( \frac{\pi}{3} - \varphi \right) \right] \sin \varphi + \\ &(A - B) l \left[ \sin \varphi - \sin \left( \frac{\pi}{3} - \varphi \right) \right] + \frac{\sqrt{3}}{4} (A - B)^2 + \frac{l^2}{\sqrt{3}} \sin^2 \left( \frac{\pi}{3} - \varphi \right) = \\ &\frac{\sqrt{3}}{4} (A - B)^2 + \sqrt{3} (A - B) l \sin \left( \varphi - \frac{\pi}{6} \right) + \frac{l^2}{\sqrt{3}} \left[ 1 - \frac{1}{2} \cos \left( \frac{\pi}{3} - 2\varphi \right) \right], \\ &\int_0^{\frac{\pi}{2}} \mathfrak{S}_2(\varphi) d\varphi = \left| \frac{\sqrt{3}}{4} (A - B)^2 \varphi_0 - \sqrt{3} (A - B) l \cos \left( \varphi - \frac{\pi}{6} \right) \right|_0^{\varphi_0} + \\ &\frac{l^2}{\sqrt{3}} \varphi_0 + \frac{l^2}{4\sqrt{3}} \sin \left( \frac{\pi}{3} - 2\varphi_0 \right) \Big|_0^{\varphi_0} + \frac{l^2}{2} \sin^2 \varphi \Big|_{\varphi_0}^{\frac{\pi}{3}} + \frac{l^2 \pi}{6\sqrt{3}} + \left[ \frac{\sqrt{3}}{16} \sin 2\varphi \Big|_{\frac{\pi}{3}}^{\frac{\pi}{2}} - \right. \\ &\left. \frac{\sqrt{3}}{16} \sin \left( 2\varphi - \frac{2\pi}{3} \right) \Big|_{\frac{\pi}{3}}^{\frac{\pi}{2}} - \frac{1}{16} \sin \left( 2\varphi + \frac{2\pi}{3} \right) \Big|_{\frac{\pi}{3}}^{\frac{\pi}{2}} + \frac{1}{8\sqrt{3}} \sin \left( 2\varphi + \frac{\pi}{3} \right) \Big| + \right. \\ &\left. \frac{1}{8\sqrt{3}} \sin \left( 2\varphi + \frac{\pi}{3} \right) \Big|_{\frac{\pi}{3}}^{\frac{\pi}{2}} \right] l = \frac{\sqrt{3}}{4} (A - B)^2 \varphi_0 + \sqrt{3} (A - B) l \left[ \frac{\sqrt{3}}{2} - \cos \left( \varphi_0 - \frac{\pi}{6} \right) \right] + \end{aligned}$$

$$l^2 \left[ \frac{\varphi_0}{\sqrt{3}} + \frac{\pi}{6\sqrt{3}} + \frac{1}{4\sqrt{3}} \sin \left( \frac{\pi}{3} - 2\varphi_0 \right) - \frac{1}{2} \sin^2 \varphi_0 + \frac{1}{8} \right]. \quad (15)$$

Finally the set  $\mathfrak{S}_1(\varphi)$  if  $\varphi \in [0, \varphi_0]$  is the union of three polygons: an horizontal parallelogram with sides  $A - 2l_2(\varphi)$  and  $\frac{l}{\sqrt{3}} \sin \varphi$ , having angles  $\frac{\pi}{3}$  and  $\frac{2\pi}{3}$ , and two oblique parallelograms; of eachone of the parallelograms we eliminate an equilateral triangle of side  $\frac{2l}{\sqrt{3}} \sin \left( \frac{\pi}{3} - \varphi \right) - (A - B)$ . The horizontal sides of this parallelogram are  $l_1(\varphi)$  and  $l_2(\varphi)$ ; The vertical height have length  $B \frac{\sqrt{3}}{2} - l \sin \varphi$ .

Hence for  $\varphi \in [0, \varphi_0]$  we have

$$\begin{aligned} \text{Area } \mathfrak{S}_1(\varphi) &= \left[ A - \frac{2l}{\sqrt{3}} \sin \left( \varphi + \frac{\pi}{3} \right) \right] \frac{l}{2} \sin \varphi + \\ &\quad \left( B \frac{\sqrt{3}}{2} - l \sin \varphi \right) \left( \sin \left( \varphi + \frac{\pi}{3} \right) + \sin \left( \frac{\pi}{3} - \varphi \right) \right) \frac{l}{\sqrt{3}} - \\ &\quad 2 \cdot \frac{\sqrt{3}}{4} \left[ \frac{2l}{\sqrt{3}} \sin \left( \frac{\pi}{3} - \varphi \right) - (A - B) \right]^2 = \frac{Al}{2} \sin \varphi + \frac{\sqrt{3}}{2} Bl \cos \varphi - \\ &\quad \frac{\sqrt{3}}{2} (A - B)^2 + 2(A - B) l \sin \left( \frac{\pi}{3} - \varphi \right) - \\ &\quad l^2 \left[ \frac{5}{4\sqrt{3}} + \frac{1}{2\sqrt{3}} \cos \left( \frac{\pi}{3} - 2\varphi \right) \right], \\ \int_0^{\varphi_0} \text{Area } \mathfrak{S}_1(\varphi) d\varphi &= \frac{Al}{2} (1 - \cos \varphi_0) + \frac{\sqrt{3}}{2} Bl \sin \varphi_0 - \frac{\sqrt{3}}{2} (A - B)^2 \varphi_0 + \\ &\quad 2(A - B) l \left[ \cos \left( \frac{\pi}{3} - \varphi_0 \right) - \frac{1}{2} \right] - l^2 \left[ \frac{5}{4\sqrt{3}} \varphi_0 + \frac{1}{8} - \frac{1}{4\sqrt{3}} \sin \left( \frac{\pi}{3} - 2\varphi_0 \right) \right] \end{aligned}$$

and therefore it is had that:

$$\begin{aligned} \int_0^{\frac{\pi}{2}} \text{Area } \mathfrak{S}_1(\varphi) d\varphi &= \frac{Al}{2} (1 + \cos \varphi_0) + Bl \left( 1 - \frac{1}{2} \cos \varphi_0 \right) - \frac{\sqrt{3}}{2} (A - B)^2 \varphi_0 + \\ &\quad 2(A - B) l \left[ \cos \left( \frac{\pi}{3} - \varphi_0 \right) - \frac{1}{2} \right] - \\ &\quad l^2 \left[ \frac{5}{4\sqrt{3}} \varphi_0 + \frac{5\pi}{24\sqrt{3}} + \cos^2 \varphi_0 - \frac{1}{4\sqrt{3}} \sin \left( \frac{\pi}{3} - 2\varphi_0 \right) \right]. \quad (16) \end{aligned}$$

Form the formulas (14), (15), (16) and (4) follows:

**Theorem 2** Let  $s$  a line segment of length  $l$ , that respect to the dimension of lattice  $\mathfrak{R}$  satisfies the relations  $A - B \leq l \leq h \leq 2(A - B)$  or  $A - B \leq l \leq 2(A - B) \leq h$ . Let  $\varphi_0 \in [0, \frac{\pi}{3}]$  is the only angle with the property  $\sin(\frac{\pi}{3} - \varphi_0) = \frac{A-B}{l} \cdot \frac{\sqrt{3}}{2}$ . The line segment  $s$  is uniformly distributed in a bounded region of the Euclidean plane, then the probability that  $s$  intersects one time, two times or three times the sides of lattice  $\mathfrak{R}$  are respectively

$$p_1 = \frac{4}{\pi\sqrt{3}(A-\frac{B}{2})B} \left[ \frac{Al}{2} (1 + \cos \varphi_0) + Bl \left( 1 - \frac{1}{2} \cos \varphi_0 \right) - \frac{\sqrt{3}}{2} (A - B)^2 \varphi_0 + 2(A - B)l \left( \cos \left( \frac{\pi}{3} - \varphi_0 \right) - \frac{1}{2} \right) - l^2 \left( \frac{5\varphi_0}{4\sqrt{3}} + \frac{5\pi}{24\sqrt{3}} + \cos^2 \varphi_0 - \frac{1}{4\sqrt{3}} \sin \left( \frac{\pi}{3} - 2\varphi_0 \right) \right) \right], \quad (17)$$

$$p_2 = \frac{4}{\pi\sqrt{3}(A-\frac{B}{2})B} \left[ \frac{\sqrt{3}}{4} (A - B)^2 \varphi_0 + \sqrt{3}(A - B)l \left( \frac{\sqrt{3}}{2} - \cos \left( \varphi_0 - \frac{\pi}{6} \right) \right) + l^2 \left( \frac{\varphi_0}{\sqrt{3}} + \frac{\pi}{6\sqrt{3}} + \frac{1}{4\sqrt{3}} \sin \left( \frac{\pi}{3} - 2\varphi_0 \right) - \frac{1}{2} \sin^2 \varphi_0 + \frac{1}{8} \right) \right], \quad (18)$$

$$p_3 = \frac{4}{\pi\sqrt{3}(A-\frac{B}{2})B} \left[ \frac{l(A-B)}{2} (\cos \varphi_0 - 1) + \frac{l^2}{4} - \frac{l^2}{4\sqrt{3}} \left( \frac{\pi}{6} + \varphi_0 + \sin \left( \frac{\pi}{3} - 2\varphi_0 \right) \right) \right]. \quad (19)$$

## References

- [1] Duma, A., Stoka M.I., Problems of Buffon type with multiple intersections for lattices of parallelograms, Rend. Circ. Mate. Palermo, serie II, 2006, pp. 241-248.
- [2] Duma, A., Stoka M.I., Problems of Buffon Type with multiple intersections for lattices of equilateral triangles, preprint.
- [3] Duma, A., Problems of Buffon type for "non-small" needles, Rend. Circ. Mat. Palermo, serie II, 1999, pp. 23-40.
- [4] Stoka, M.I., Probabilités géométrique de type Buffon dans le plan euclidean, Atti Acc. Sci. Torino, vol. 10, (1975-1976), pp. 53-59.

Giuseppe Caristi

Department D.E.S.Ma.S. "V. Pareto"

Faculty of Economics, University of Messina - ITALY

Via dei Verdi 75, 98122 Messina.

E-mail address: gcaristi@dipmat.unime.it