



Microbiota of the Digestive Glands and Extrapallial Fluids of Clams Evolve Differently Over Time Depending on the Intertidal Position

Clément Offret, Olivier Gauthier, Garance Despréaux, Adeline Bidault, Charlotte Corporeau, Philippe Miner, Bruno Petton, Fabrice Pernet, Caroline Fabioux, Christine Paillard, et al.

► To cite this version:

Clément Offret, Olivier Gauthier, Garance Despréaux, Adeline Bidault, Charlotte Corporeau, et al.. Microbiota of the Digestive Glands and Extrapallial Fluids of Clams Evolve Differently Over Time Depending on the Intertidal Position. Microbial ecology, 2022, 85 (1), pp.288-297. 10.1007/s00248-022-01959-0 . hal-03637412

HAL Id: hal-03637412

<https://hal.science/hal-03637412>

Submitted on 20 Nov 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Microbiota of The Digestive Glands and Extrapallial Fluids of Clams Evolve Differently Over Time Depending On the Intertidal Position

Clément Offret (✉ clement.offret@univ-brest.fr)

Institut Universitaire Européen de la Mer <https://orcid.org/0000-0001-5786-2929>

Olivier Gauthier

Institut Universitaire Européen de la Mer

Garance Desprésaux

Institut Universitaire Européen de la Mer

Adeline Bidault

Institut Universitaire Européen de la Mer

Charlotte Corporeau

Univ Brest

Philippe Miner

Univ Brest

Bruno Petton

Univ Brest

Fabrice Pernet

Univ Brest

Caroline Fabioux

Institut Universitaire Européen de la Mer

Christine Paillard

Institut Universitaire Européen de la Mer

Gwenaëlle Le Blay

Institut Universitaire Européen de la Mer

Research Article

Keywords: Clam, Microbiota, Digestive Gland, Extrapallial Fluids, Temporal variations

Posted Date: October 20th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-976571/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Microbial Ecology on January 22nd, 2022.

See the published version at <https://doi.org/10.1007/s00248-022-01959-0>.

Abstract

The Manila clam (*Ruditapes philippinarum*) is the second most exploited bivalve in the world but remains threatened by diseases and global changes. Their associated microbiota play a key role in their fitness and acclimation capacities. This study aimed at better understanding the behavior of clam digestive glands and extrapallial fluids microbiota at small, but contrasting spatial and temporal scales. Results showed that environmental variations impacted clam microbiota differently according to the considered tissue. Each clam tissue presented its own microbiota, and showed different dynamics according to the intertidal position and sampling period. Extrapallial fluids microbiota was modified more rapidly than digestive glands microbiota, for clams placed on the upper and lower intertidal position, respectively. Clam tissues could be considered as different microhabitats for bacteria as they presented different responses to small-scale temporal and spatial variabilities in natural conditions. These differences underlined a more stringent environmental filter capacity of the digestive glands.

1. Introduction

Host-associated microbiota are organized into trophic groups of microorganisms that interact with each other and the organ they colonize. These microorganisms come from the parents (vertical transmission) or from the immediate environment (horizontal transmission). The colonized tissues offer particular physicochemical conditions, and act as micro-habitats. Both the host and its associated microbiota contribute to the metabolic specificity of these organs. As a result, each tissue is colonized by specific microbiota communities that are both complex and dynamic [1]. Associated microorganisms contribute to varying degrees to many metabolic processes and gene expressions that can have major impacts on host nutrition, detoxification, growth, immune system development and resistance to infections [2–4]. They probably play a key role in hosts acclimation to different ecological conditions in fluctuating environments as coastal zones.

It has been shown that marine host-associated microbiota may be influenced by different biotic and abiotic factors (season, location, host age, health and genotype) [5–8]. Being filter-feeders, bivalves interact with numerous microorganisms coming from the environment. They maintain tissue-specific microbiota that differ from that of their surrounding environment [9], indicating that both organisms and tissues select the bacteria they host filters [10, 11]. Nonetheless, these host-associated microbiota may vary spatially and temporally reflecting seasonal changes and geographic location [12–16]. Intrinsic factors, like life stage, genetic, host tissue or host taxonomy may also induce different variations in terms of microbiota composition and dynamic [17–19]. However, the mechanisms underlying the interactions in bivalve microbiota and how these ones are likely to modify their host acclimation to a changing environment are still largely unknown.

The Manila clam (*Ruditapes philippinarum*), is a sediment burrower that mainly ingests benthic microalgae and sedimented phytoplankton [20]. Its microbiota is influenced by environmental factors such as season [21], contaminants [22], and intertidal position [11]. They also differ among heart, gonad,

mantle, hemolymph, gills or gut [21, 23], indicating an organ-specific microbiota. This suggests that factors may have different effects on microbiota dynamics depending on the considered tissues, and so differently affect their functions.

Almost nothing is known about clam's extrapallial fluids microbiota, although this compartment plays an important role in immunity and calcification processes [24], which could be influenced by microbial changes. Indeed, it has been previously shown that microbiota from extrapallial fluids of Eastern oyster (*Crassostrea virginica*) is influenced by location and season [10].

Microbiota is important in adaptation to fluctuating environment like the intertidal zone, and that bivalves are good models for that. To highlight specific responses of tissue-associated microbiota in the face of environmental small-scale variations, we investigated the dynamics of clam microbiota during the spring revival, a period that is characterized by marked environmental changes impacting the host species [25]. Animals were deployed at two tidal levels, and were sampled in February, March and April. Two tissues that are differently exposed to the environment were compared (with and without depuration), the extrapallial fluids (EF), which are at the interface between the animal and the environment, and the digestive glands (DG), an internal organ that is exposed to what is ingested. This study will provide a better understanding of the dynamic and scales of variation of microbiota associated with the DG and EF of healthy clams in natural conditions.

2. Experimental Procedures

2.1. Experimental design

The Manilla clams *Ruditapes philippinarum* were hatched in April 2016 and were provided by a commercial exploitation (SATMAR, France). In October 2017, clams were placed at the site IFREMER Br08 located in the Bay of Brest at Pointe du Château (48° 20' 06.19" N, 4° 19' 06.37" W, Brittany, France) at two different intertidal levels corresponding to 20% (L20) and 56% (L56) of exondation time (Fig.1) in two duplicated mesh bags.

Healthy clams (i.e. without signs of Brown Ring Disease) were collected from the two intertidal levels over three consecutive days (to avoid sampling bias) in T1 (February 2018, 19.6 ± 3.4 mm), T2 (March 2018, 18.1 ± 2.4 mm) and T3 (April 2018, 19.5 ± 2.8 mm). For each sampling day, sediments were sampled in triplicate at each level, next to the clams, and seawater was collected in triplicate 2h before low tide. The seawater samples (1L) were successively passed through 8 (SW8) and 0.22 μ m (SW022) polycarbonate filters (Whatman, USA) that were kept at -80°C until DNA extraction and sequencing. For each day, clams (n=10) were collected by level for bacterial microbiota analyses, and half of them (n=5) were directly dissected for tissue sampling, while others (n=5) were transferred to the laboratory and grouped by day and intertidal level for depuration process in order to empty the DG and reduce environmental microorganisms [26]. After 14 days in tanks (30 L) fed with flow-through (3 L / min) filtered seawater (10 and 5 μ m sand filters and UV treatment before two 1 μ m filters, and a second UV treatment), tissues were

dissected and collected before DNA extraction. Tanks were cleaned every day to avoid biofilm formation and bacterial proliferation. No feed was added.

2.2. Bacterial genomic DNA sequencing

Microbiota analyses were performed from digestive glands (DG) and extrapallial fluids (EF) of both depurated and non-depurated clams, as detailed by Offret *et al.*, [11]. Briefly, DG and EF were aseptically dissected from clams directly after sampling or after depuration. Bacterial genomic DNA from both clam tissues and environmental samples were extracted using the PowerLyzer PowerSoil DNA Isolation Kit (Qiagen, USA). For each sample, the DNA was amplified by PCR targeting the variable V3V4 region coding for the bacterial 16S rRNA gene using the 341F (5'-CCTACGGGNGGCWGCAG -3') and 805R (5'-GACTACHVGGGTATCTAATCC-3') primers [27]. Finally, paired-end sequencing with a 300-pb read length was performed at McGill University (G  nome Quebec Innovation Centre, Montr  al, Qc, Canada) on a MiSeq system (Illumina).

2.3. Bacterial microbiota analysis

Sequences were processed with the Galaxy-supported FROGS pipeline (« Find, Rapidly, OTUs with Galaxy Solution »). This pipeline was developed in order to analyze large sets of amplicon sequences and produce abundance tables of Operational Taxonomic Units (OTUs) with their taxonomic affiliation [28].

Firstly, bacterial 16S rDNA sequences of samples were pre-process. Paired-end reads were merged using Flash [29] with 10% mismatch tolerated. Primers and adapters were removed using cutadapt [30]. Clustering was achieved using the Swarm method with an aggregation distance ($d=3$) [31]. Afterwards, chimeras were detected and removed with VSEARCH (*de novo* UCHIME method) before applying a cross-sample validation step [28, 32, 33]. Next, a relative abundance filter was applied on OTUs with a threshold of 0.005% (of global relative abundance), as recommended by Bokulich *et al.*, [34]. Taxonomic affiliation was done on this new data by blastn+ [35], using the Silva 132 16S database [36] formatted to include taxonomic levels up to species [28]. For analysis, cyanobacterial and eukaryotic sequences (chloroplast and mitochondrial 16S rRNA gene from algal cells) were removed.

2.4. Statistical analyses

2.4.1. Relative abundance and α -diversity

The relative abundance of the whole bacterial microbiota communities was assessed at the order level at each period [8]. Microbiota α -diversity was assessed at the OTUs level and the rarefied richness and Shannon indices were computed. The *period*, *level*, *tissue* and *depuration* effects (and their interactions) were tested on clam's α -diversity indices. For the environmental samples, the effects of *period* and *level* (and their interaction) were tested on the sediment and the effects *period* and *filter* (and their interaction) on the seawater. For each model, the effects were tested using a linear model on Box-Cox transformed data and a Generalized Linear Model (GLM) on non-transformed data. The linear model was either a two-

way analysis of variance (ANOVA) or an analysis of covariance (ANCOVA) with the number of sequences as a co-variable. Finally, the models with the greater adjusted R^2 were kept.

2.4.2. β -diversity

Microbial communities β -diversity was also assessed at the OTUs level. On the one hand, the OTUs abundances tables were transformed using the Box-Cox-chord transformation with exponent 0 which corresponds to a log-chord transformation [37]. Then, the variation in microbiota composition and structure between all samples was visualized with a PCA of these transformed OTUs abundances. The effects of *period*, *level* and their interactions were tested using PERMANOVA [38] when group dispersions were homogeneous. Multivariate homogeneity of group dispersions was assessed for the DG, EF, sediments and seawater samples separately. All analyses were carried out using R3.6.2 [39], with functions from the *phyloseq* [40], *vegan* [41], *emmeans* [42], *RAM* [43] and *VIM* [44] packages.

3. Results

3.1. Sequences analysis information

The total number of sequences before processing was 13 406 675. After processing, 7 369 012 sequences were kept (*i.e.* 54.97 %). After removing sequences affiliated to chloroplasts and samples with a number of sequences reads inferior to 3 000, 432 samples were kept: 160 DG's samples, 177 EF's samples, 41 sediment's samples and 54 seawater's samples (27 SW8 and 27 SW022). The swarm clustering produced 1 255 different OTUs, which were identified in GD (1152), EF (1 201), sediments (935) and seawater (1 057 and 972 into SW8 and SW022 fractions, respectively). The mean number of sequences reads by sample was $16\,063 \pm 332$ for all samples: $15\,297 \pm 478$ for the DG, $19\,251 \pm 446$ for the EF, $5\,487 \pm 203$ for the sediments, $15\,917 \pm 851$ for the seawater ($10\,924 \pm 601$ for SW8 and $20\,910 \pm 824$ for SW022).

3.2. α -diversity comparison for clam tissues and environmental samples

The effect of the number of sequence reads was significant for observed richness ($F = 7.6$, $P < 0.01$) justifying the use of a Rarefied richness for describing α -diversity patterns. Models applied to clam's tissues microbiota α -diversity indexes (rarefied richness and Shannon indices) highlighted two main factors: *Depuration* and *Tissue* (including the seawater fractions), followed by the *Period* (Fig.2). The *Depuration* induced a significant decrease of Rarefied richness ($F = 4.7$, $P < 0.01$) and Shannon indices ($F = 62.2$, $P < 0.001$), while *Tissue* differences results in an average EF α -diversity that were always significantly higher than those for the GD (Rarefied richness: $F = 29.4$, $P < 0.001$; Shannon: $F = 14.6$, $P < 0.001$). The two seawater fractions exhibited two clearly different α -diversity, where the bacterial communities from SW8 fraction were always greater than those from the SW022 one (Rarefied richness: $F = 211.9$, $P < 0.001$). Nonetheless, on average α -diversity of microbiota associated with the two clam tissues was always significantly lower than those of bacterial communities from sediment and SW8 (post-hoc Tukey test, $P < 0.001$).

The *Period* factor was significant for all models, except for the Shannon indices of sediment ($F = 1.4$, $P > 0.05$). Overall regarding both indices, average α -diversity of bacterial communities from non-depurated DG and seawater samples, significantly decreased during the studied period, notably between T1 and T3 (post-hoc Tukey test, $P < 0.001$). In the other hand, non-depurated EF and sediment alpha diversities significantly increased between T1 and T3.

3.3. Order level bacterial diversity from clam tissues and environmental samples

Based on the 30 most abundant bacterial orders, DG, EF, sediment and seawater fractions showed clearly distinct bacterial communities in terms of both abundance and taxa composition (Fig.3). The two fractions of seawater (SW8 and SW022) were dominated by the orders *Rhodobacterales* (55-70 % in SW022 and 20-30 % in SW8) and *Flavobacteriales* (12-15% in SW022 and 10-22 % in SW8). However, their composition differed for less abundant orders (inferior to 6% of total abundance). Indeed, SW022's bacterial community was mostly composed of *Oceanospirillales* and *Cellvibrionales*, whereas SW8's bacterial community was more diverse with *Cellvibrionales*, *Desulfobacterales*, *Campylobacterales*, *Pirellulales*, *Rhizobiales*, *Verrucomicrobiales*, *Clostridiales* and *Myxococcales* present in low abundance (inferior to 5%). Sediment bacterial communities were mostly composed of numerous few abundant orders, accounting for more than 25% of the total abundance, while the most abundant orders were mainly affiliated to *Desulfobacterales* (13-19%), *Actinomarinales* (6-8%), *Campylobacterales* (5-7%), *Microtrichales* (5-7%) and *Rhizobiales* (4-6%).

The non-depurated clams, DG microbiota was mostly composed in a decreasing order of contribution, *Rickettsiales*, *Mycoplasmatales*, *Diplorickettsiales*, *Spirochaetales*, *Pirellulales*, *Clostridiales*, *Oceanospirillales* and *Flavobacteriales*. DG microbiota from clams sampled in T1 was characterized by a lower relative abundance of *Rickettsiales* and a higher relative abundance of *Spirochaetales* and *Clostridiales*, while higher relative abundances of *Campylobacterales* and *Flavobacteriales* were observed respectively in T2 and T3. One of the most abundant order found in DG microbiota, the *Diplorickettsiales*, drastically decreased in T3 compared to T1 and T2. In depurated clams, the DG microbiota was globally composed of the same orders as non-depurated clams, but in different abundances. The main changes were observed for *Spirochaetales*, increasing from around 6% to 20% between the non-depurated and depurated clams, while *Pirellulales* abundances were divided by 7 in depurated clams.

Although it harbored some taxa (at the order level) in common, EF microbiota of non-depurated clams was clearly different from that of the DG's. It was mainly composed of *Oceanospirillales*, *Spirochaetales*, *Rickettsiales*, *Flavobacteriales*, *Sneathiellales*, *Campylobacterales*, and also *Rhodobacterales*, *Francisellales*, *Verrucomicrobiales*, *Cellvibrionales* and *Alteromonadales*, those last clearly contributing to the differences with DG microbiota. In T1, the EF microbiota was characterized by a higher relative abundance of *Rickettsiales* and *Oceanospirillales* and a lower abundance of *Alteromonadales*, *Sneathiellales* and *Rhodobacterales*. A higher relative abundance of *Campylobacterales* and *Flavobacteriales* were observed in T2 and T3, respectively. In EF microbiota from depurated clams, the same orders were globally found than in non-depurated clams but in different abundances, notably for

Alteromonadales, *Francisellales* and *Pseudomonadales* which were present in much higher abundances, whereas *Flavobacteriales* and *Rhodobacterales* were reduced.

3.4. Factors influencing Beta diversity

From this point on, results are those of analyses conducted at the OTU's level. PCA (Fig.4) allow clearly discriminating sample matrix (clams vs sediment and seawater on PC1), seawater fractions (SW8 vs SW022 on PC2) and clam tissues (EF vs DG on PC2). The SW8 bacterial community was more similar to that of sediments, whereas the one from SW022 was more similar to that of the EF.

Homogeneity of variances was previously tested for each sample grouping, revealing that multivariate dispersion was not homogenous (Fig.5) for both DG ($F = 17.333$, $P < 0.001$, 999 permutations) and EF tissues ($F = 6.893$, $P = 0.001$, 999 permutations). Using the *betadisper* function, the DG microbiota had a higher heterogeneity of group dispersions than EF. The heterogeneity of dispersions occurred between the depurated samples of L56 in T3 and all the groups in T1 (Fig. 5A). It also occurred between the non-depurated samples of L20 in T2 and all the groups in T2. The dispersions were also heterogeneous between T1 and T3. In the case of the EF's microbiota (Fig. 5B), the heterogeneity of dispersions implied three groups of clams: the depurated at L20 in T1, the non-depurated at L56 in T2 and all the depurated in T3. The L20 levels of T2 and T3 seemed more similar to T1 (L20 and L56) than to L56. For the depurated clams, all clusters were overlapping, indicating a more similar community over the three periods. However, L56 and L20 of the three periods tended to pull away from each other and the centroid, indicating differences between levels.

Regarding the DG bacterial microbiota (Fig.5a), a high heterogeneity of dispersion, which increased between T1 and T3, was observed. However, it did not prevent us from seeing the dynamic of the microbiota. A clear distinction between depurated and non-depurated clam's microbiota was observed along the first axis of the PCA for both levels (L20 & L56) in T1 and T2, which was much less the case in T3. For the non-depurated clams, the DG bacterial microbiota in T1 and T2 were very close regardless of the levels, no period or level effects were observed. It is only in T3 that the DG bacterial microbiota detached (especially for L20) from T1 and T2, and that a slight level effect appeared. For the depurated clams, the DG microbiota were more similar between the three periods. Moreover, in that case the level effect was more pronounced in T2 than in T3.

For the EF bacterial microbiota (Fig.5b), compared to the DG microbiota, the distinction between depurated and non-depurated was clearer along the first axis, as clusters did not overlap in the PCA regardless of levels and periods. The EF microbiota behaved differently from the DG ones. In T1, no level effect was visible (as for the DG), but in T2 and T3 the level effect becomes more important than the period effect, with a clear distinction between levels regardless of the time period. The L56 levels were clearly distinct from the other ones in both T2 and T3.

The PCA representing the sediments communities (Fig.5c) showed a level effect but no period effect. Only the L56 samples in T1 appeared to have a distinct microbiota from other L56 samples. Multivariate

analysis of variance on the Box-Cox transformed and standardized environmental data reveals a period effect ($F = 1.77$, $P = 0.025$, $R^2 adj = 0.069$), the level ($F = 9.25$, $P = 0.001$, $R^2 adj = 0.18$) and their interaction ($F = 1.59$, $P = 0.041$, $R^2 adj = 0.062$). The seawater's bacterial community (Fig.5d) was clearly different between the three periods and the two filters. The heterogeneity of dispersion between the groups did not allow a multivariate analysis of variance ($F = 3.6108$, $P = 0.037$, 999 permutations).

4. Discussion

In this study, we investigated the diversity and the temporal dynamic of the DG and EF microbiota from Manila clams placed on two intertidal levels in the foreshore during spring revival. Results brought more information regarding specificity of clam tissue microbiota, while both small spatial and temporal scales significantly contributed to the shaping of the DG and EF microbiota. The observed differences were attributed in a decreasing order to the depuration process, the tissue, the sampling period and finally to the intertidal level. Furthermore, different dynamics were clearly observed depending on the tissue and intertidal level.

4.1. The environmental contribution to the clam holobiont

Regarding clam microbiota, both DG and EF were clearly different from the environment, although the bacterial communities in the surrounding seawater and sediments contributed to the microbiota shaping. Indeed, EF microbiota was much closer to environmental bacterial communities than DG microbiota, especially from the 0.22–8 μm seawater fraction, suggesting a higher proximity between EF and seawater (free and small particles associated bacteria) than with the digestive glands. OTUs present in the clams DG microbiota, such as *Rickettsiales*, *Mycoplasmatales*, *Diplorickettsiales*, *Spirochaetales*, *Oceanospirillales* and *Flavobacteriales*, as commonly found in clams DG [11, 22], with *Mycoplasmatales* and *Rickettsiales* attributed to the core members [22]. EF microbiota harbored a high abundance of taxa orders such as *Rhodobacterales*, similar to those found in seawater but not in the DG. This is probably the result of a direct exchanges between seawater and EF, maybe through the pallial line, possibly associated with less stringent EF physical and chemical conditions for the seawater bacteria than in the DG. To the author's knowledge, EF microbiota of Manila clams has never been described so far. However, *Rickettsiales* and *Flavobacteriales* were already found in high abundance in EF microbiota from another bivalve species, the Eastern Oyster *Crassostrea virginica* [10].

4.2. Tissue-specific of clam microbiota dynamics

This study revealed a different response of the two-tissue microbiota over time. Even if both DG and EF microbiota showed a strong temporal variability in terms of α -diversity, EF's α -diversity significantly increased over time in their natural environment, whereas that of the DG's decreased. Indeed, organ microbiota are shaped by the organ's physiological role, notably digestion and shell mineralization for DG and EF, respectively (Wilbur and Saleuddin, 1983; Allam and Paillard, 1998). On the one hand, the Manila clam's different organs are known to be bacterial niches with unique features [21], and so they could be considered as different micro-habitats. On the other hand, this could be linked to environmental

variations, such as temperature and dissolved oxygen which have already been shown to impact EF microbial composition of Pacific Oysters [10]. Regarding DG microbiota, a higher α -diversity has been described from hepatopancreas microbiota of the Manila clams sampled in winter than in summer [22]. We hypothesize that the observed decrease of bacterial α -diversity in clam DG, could be caused by presence of other water associated microorganisms such as micro-eukaryotes.

In addition to the temporal heterogeneity, a tidal level effect has also been observed for both DG and EF, but their dynamics were very different. The greater effect of the intertidal level on EF microbiota in T2 and T3 may be explained by a greater heterogeneity of environmental conditions in L56 since the immersion period was shorter. Microbiota from bivalve fluids seemed to be particularly sensitive to the sampling zone as it was shown for hemolymph's microbiota from the Pacific Oyster [47]. These results seem to hint that EF microbiota are more impacted by environmental variations when they are placed in the upper part of the foreshore. On the contrary, DG microbiota were more impacted by the period when they were placed in the lower part of the foreshore (L20), which was under a greater influence of sea water and feeding sources. So, depending on the considered tissue, microbiota dynamics vary with intertidal position.

4.3. GD are more stringent environmental filters than EF

Analysis of β -diversity components (data not shown) also revealed distinct variations between the two-tissue microbiota. EF bacterial microbiota differences between individuals were mostly driven by species replacement for the all three sampling periods. It is thus hypothesized that EF reached maximum niche occupancy. On the other hand, DG bacterial microbiota β -diversity was mostly driven by richness difference, except for clams sampled in T1. These results highlight the possibility that OTUs richness might be more constrained in EF and less so in DG.

In addition to the fact that α -diversity was higher in the EF microbiota, depuration process more affected EF than GD may illustrate that EF microbiota is composed of more transient microorganisms coming from the surrounding sea water. The analysis of β -diversity revealed a higher dispersion in DG microbiota compared to EF microbiota, illustrating a higher within-tissue diversity in the DG. All these results revealed that DG is a potentially more stringent environmental filter than EF.

In conclusion, this study has shown that environmental variations may have a strong influence on clam microbiota, through impact on both digestive gland and extrapallial fluids associated bacteria. DG and EF presented their own microbiota specificity in terms of abundance, richness and dynamics. Furthermore, clam's tissue could be considered as distinct habitats from the bacterial OTUs point of view, as they presented different responses to either small-scale temporal and spatial variabilities or to a depuration process, illustrating that their selection processes differ. These differences underlined that DG could be a more stringent micro-habitat probably linked to its particular physico-chemical conditions. This study helps to have a better understanding of the Manila Clam's microbiota dynamic in the face of environmental variation, and especially of the EF microbiota which is still very poorly known.

Declarations

Declarations: Not applicable.

Funding: This work was supported by the ISblue project, Interdisciplinary graduate school for the blue planet (grant number ANR-17-EURE-0015) and co-funded by a grant from the French government under the program 'Investissements d'Avenir'. The Region Bretagne SAD (2017, Stratégie d'Attractivité Durable) contributed to this study through the postdoctoral fellowship of Clement Offret. This work was also supported by the HORIZON2020 project 'Preventing and mitigating farmed bivalve disease' VIVALDI (grant number 678589).

Conflict of interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Availability of data and material: The sequence data have been deposited in the NCBI database under BioProject ID PRJNA748558.

Authors' contributions: C.O., G.D., G.L.B. and O.G. wrote the manuscript, and all authors commented on the final version. Experiments, microbiota analyses, results interpretation, C.O.; microbiota analyses, results interpretation, G.D.; statistical analyses, O.G. and GD; Project and experimental design, oyster and clam dissections, A.B., P.M., B.P., C.C., F.P., C.F., Project and experimental design, oyster and clam dissections and funding acquisition C.P.; Project supervision and experimental design, field monitoring, oyster and clam dissections, results interpretation and funding acquisition, G.L.B.

Acknowledgements

The authors wish to thank Bruno Petton, Elodie Fleury, Christophe Lambert, Nelly Le Goic, Morgan Perennou, Christine Dubreuil, Valerian Le Roy, Claudie Quere, Amelie Segarra, Alexandra Rahmani, Elyne Dugeny, Morgan Smith, Lizenn Delisle, Veronique Loiseau and Amandine Morot for their help with sampling preparation and dissections of bivalves, and Dominique Hervio for her help with water filtration. Thanks to Stephane Pouvreau and the Marine environmental station database of Daoulas bay for their essential help (<https://www.seanoe.org/data/00314/42493/>).

References

1. Weissenbach J, Sghir A (2016) Microbiotes and Metagenomics. *Medecine sciences: M/S* 32:937–943
2. Rawls JF, Samuel BS, Gordon JI (2004) Gnotobiotic zebrafish reveal evolutionarily conserved responses to the gut microbiota. *Proceedings of the National Academy of Sciences* 101:4596–4601
3. Rosenberg E, Koren O, Reshef L et al (2007) The role of microorganisms in coral health, disease and evolution. *Nat Rev Microbiol* 5:355–362. <https://doi.org/10.1038/nrmicro1635>

4. Dishaw LJ, Cannon JP, Litman GW, Parker W (2014) Immune-directed support of rich microbial communities in the gut has ancient roots. *Developmental Comparative Immunology* 47:36–51
5. Gil-Agudelo DL, Myers C, Smith GW, Kim K (2006) Changes in the microbial communities associated with *Gorgonia ventalina* during aspergillosis infection. *Dis Aquat Organ* 69:89–94
6. Koren O, Rosenberg E (2006) Bacteria associated with mucus and tissues of the coral *Oculina patagonica* in summer and winter. *Appl Environ Microbiol* 72:5254–5259
7. Littman RA, Willis BL, Pfeffer C, Bourne DG (2009) Diversities of coral-associated bacteria differ with location, but not species, for three acroporid corals on the Great Barrier Reef. *FEMS Microbiol Ecol* 68:152–163
8. Chen C-P, Tseng C-H, Chen CA, Tang S-L (2011) The dynamics of microbial partnerships in the coral *Isopora palifera*. *The ISME Journal* 5:728–740
9. Harris JM (1993) The presence, nature, and role of gut microflora in aquatic invertebrates: A synthesis. *Microb Ecol* 25:195–231. <https://doi.org/10.1007/BF00171889>
10. Sakowski EG (2015) The microbiome of the eastern oyster, *Crassostrea virginica*, in health and disease. Thesis, University of Delaware
11. Offret C, Paulino S, Gauthier O et al (2020) The marine intertidal zone shapes oyster and clam digestive bacterial microbiota. *FEMS Microbiol Ecol* 96:. <https://doi.org/10.1093/femsec/fiaa078>
12. Lokmer A, Wegner KM (2014) Hemolymph microbiome of Pacific oysters in response to temperature, temperature stress and infection. *ISME J* 9:670–682. <https://doi.org/10.1038/ismej.2014.160>
13. Cúcio C, Engelen AH, Costa R, Muyzer G (2016) Rhizosphere microbiomes of European seagrasses are selected by the plant, but are not species specific. *Frontiers in microbiology* 7:440
14. Sharp KH, Pratte ZA, Kerwin AH et al (2017) Season, but not symbiont state, drives microbiome structure in the temperate coral *Astrangia poculata*. *Microbiome* 5:120
15. Woo S, Yang S-H, Chen H-J et al (2017) Geographical variations in bacterial communities associated with soft coral *Scleronephthya gracillimum*. *PLOS ONE* 12:e0183663. <https://doi.org/10.1371/journal.pone.0183663>
16. Vijayan N, Lema KA, Nedved BT, Hadfield MG (2019) Microbiomes of the polychaete *Hydroides elegans* (Polychaeta: Serpulidae) across its life-history stages. *Mar Biol* 166:19. <https://doi.org/10.1007/s00227-019-3465-9>
17. Trabal Fernández N, Mazón-Suástegui JM, Vázquez-Juárez R et al (2014) Changes in the composition and diversity of the bacterial microbiota associated with oysters (*Crassostrea corteziensis*, *Crassostrea gigas* and *Crassostrea sikamea*) during commercial production. *FEMS Microbiol Ecol* 88:69–83. <https://doi.org/10.1111/1574-6941.12270>
18. Pierce ML, Ward JE (2019) Gut Microbiomes of the Eastern Oyster (*Crassostrea virginica*) and the Blue Mussel (*Mytilus edulis*): Temporal Variation and the Influence of Marine Aggregate-Associated. Microbial Communities *mSphere* 4:e00730–e00719. [/msphere/4/6/mSphere730-19.atom. https://doi.org/10.1128/mSphere.00730-19](https://doi.org/10.1128/mSphere.00730-19)

19. King WL, Siboni N, Kahlke T et al (2020) Regional and oyster microenvironmental scale heterogeneity in the Pacific oyster bacterial community. *FEMS Microbiol Ecol* 96:.
<https://doi.org/10.1093/femsec/fiaa054>
20. Simons AL, Churches N, Nuzhdin S (2018) High turnover of faecal microbiome from algal feedstock experimental manipulations in the Pacific oyster (*Crassostrea gigas*). *Microb Biotechnol* 11:848–858. <https://doi.org/10.1111/1751-7915.13277>
21. Meisterhans G, Raymond N, Girault E et al (2016) Structure of Manila Clam (*Ruditapes philippinarum*) Microbiota at the Organ Scale in Contrasting Sets of Individuals. *Microb Ecol* 71:194–206. <https://doi.org/10.1007/s00248-015-0662-z>
22. Milan M, Carraro L, Fariselli P et al (2018) Microbiota and environmental stress: how pollution affects microbial communities in Manila clams. *Aquat Toxicol* 194:195–207.
<https://doi.org/10.1016/j.aquatox.2017.11.019>
23. Leite L, Jude-Lemeilleur F, Raymond N et al (2017) Phylogenetic diversity and functional characterization of the Manila clam microbiota: a culture-based approach. *Environ Sci Pollut Res* 24:21721–21732. <https://doi.org/10.1007/s11356-017-9838-z>
24. Allam B, Pales Espinosa E (2016) Bivalve immunity and response to infections: Are we looking at the right place? *Fish Shellfish Immunol* 53:4–12. <https://doi.org/10.1016/j.fsi.2016.03.037>
25. Lavaud R, Artigaud S, Le Grand F et al (2018) New insights into the seasonal feeding ecology of *Pecten maximus* using pigments, fatty acids and sterols analyses. *Mar Ecol Prog Ser* 590:109–129
26. Lee R, Lovatelli A, Ababouch L (2008) Bivalve depuration: fundamental and practical aspects. Food and Agriculture Organization of the United Nations
27. Herlemann DP, Labrenz M, Jürgens K et al (2011) Transitions in bacterial communities along the 2000 km salinity gradient of the Baltic Sea. *ISME J* 5:1571–1579.
<https://doi.org/10.1038/ismej.2011.41>
28. Escudié F, Auer L, Bernard M et al (2018) FROGS: Find, Rapidly, OTUs with Galaxy Solution. *Bioinformatics* 34:1287–1294. <https://doi.org/10.1093/bioinformatics/btx791>
29. Magoč T, Salzberg SL (2011) FLASH: fast length adjustment of short reads to improve genome assemblies. *Bioinformatics* 27:2957–2963
30. Martin M (2011) Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet journal* 17:10–12
31. Mahé F, Rognes T, Quince C et al (2014) Swarm: robust and fast clustering method for amplicon-based studies. *PeerJ* 2:e593
32. Edgar RC, Haas BJ, Clemente JC et al (2011) UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics* 27:2194–2200
33. Rognes T, Flouri T, Nichols B et al (2016) VSEARCH: a versatile open source tool for metagenomics. *PeerJ* 4:e2584. <https://doi.org/10.7717/peerj.2584>

34. Bokulich NA, Subramanian S, Faith JJ et al (2013) Quality-filtering vastly improves diversity estimates from Illumina amplicon sequencing. *Nat Methods* 10:57–59.
<https://doi.org/10.1038/nmeth.2276>
35. Camacho C, Coulouris G, Avagyan V et al (2009) BLAST+: architecture and applications. *BMC Bioinform* 10:421
36. Quast C, Pruesse E, Yilmaz P et al (2012) The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic acids research* 41:D590–D596
37. Legendre P, Borcard D (2018) Box–Cox-chord transformations for community composition data prior to beta diversity analysis. *Ecography* 41:1820–1824
38. McArdle BH, Anderson MJ (2001) Fitting Multivariate Models to Community Data: A Comment on Distance-Based Redundancy Analysis. *Ecology* 82:290–297. [https://doi.org/10.1890/0012-9658\(2001\)082\[0290:FMMTCD\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2001)082[0290:FMMTCD]2.0.CO;2)
39. Team RC (2018) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. 2012. URL <http://www.R-project.org>
40. McMurdie PJ, Holmes S (2013) phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. *PloS one* 8:e61217
41. Oksanen J, Blanchet FG, Friendly M et al (2019) vegan: Community Ecology Package
42. Lenth R, Buerkner P, Herve M et al (2020) emmeans: Estimated Marginal Means. aka Least-Squares Means
43. Chen W, Simpson J, Levesque CA (2016) RAM: R for amplicon-sequencing-based microbial-ecology. R package version 1
44. Kowarik A, Templ M (2016) Imputation with the R Package VIM. *J Stat Softw* 74:1–16
45. WILBUR KM, Saleuddin ASM (1983) Shell formation. In: *The mollusca*. Elsevier, pp 235–287
46. Allam B, Paillard C (1998) Defense factors in clam extrapallial fluids. *Diseases of Aquatic Organisms* 33:123–128. <https://doi.org/10.3354/dao033123>
47. Lokmer A, Goedknecht MA, Thielges DW et al (2016) Spatial and Temporal Dynamics of Pacific Oyster Hemolymph Microbiota across Multiple Scales. *Front Microbiol* 7:..
<https://doi.org/10.3389/fmicb.2016.01367>

Figures




Figure 1

Representation of experimental design detailing the deployment of clams on the two intertidal levels (L20 and L56 corresponding to 20% and 56% of exondation time, respectively).




Figure 2

α -diversity indices (a Rarefied richness, b Shannon) of clam tissues microbiota and bacterial communities from sediment and seawater along the three-sampling periods, T1 (blue), T2 (yellow) and T3 (red). Samples were divided according to the significant factors (Tissue, Depuration, Period). Asterix indicate significant pairwise comparisons between the three periods (*, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$, post-hoc Tukey test).




Figure 3

Relative abundance of the 30 most abundant bacterial orders in digestive glands (DG) and extrapallial fluids (EF) in non-depurated (ND) and depurated (D) clams, and found in sediments (Sed), 8 μ m (SW8) and 0.22 μ m (SW022) seawater fractions, sampled at T1, T2 or T3.




Figure 4

Distance biplot of the principal component analysis of the log-chord transformed OTUs abundance of the bacterial communities of the DG (black), EF (orange), sediments (brown), SW8 (blue) and SW022 (yellow). The two axes represent 25.3% of total variance.




Figure 5

Distance biplots of the principal component analysis of the log-chord transformed OTUs abundance of the bacterial communities of the DG (a), EF (b), Sediment (c) and Seawater (d). Symbol colors and shapes in legend separating the sampling period combined to intertidal level, and the depuration process or seawater fractions.