



HAL
open science

Interaction of picolinamide fungicide primary metabolites UK-2A and CAS -649 with the cytochrome bc 1 complex Qi site: mutation effects and modeling in *Saccharomyces cerevisiae*

David Young, Nick Wang, Brigitte Meunier

► **To cite this version:**

David Young, Nick Wang, Brigitte Meunier. Interaction of picolinamide fungicide primary metabolites UK-2A and CAS -649 with the cytochrome bc 1 complex Qi site: mutation effects and modeling in *Saccharomyces cerevisiae*. *Pest Management Science*, In press, 10.1002/ps.6893 . hal-03643114

HAL Id: hal-03643114

<https://hal.science/hal-03643114v1>

Submitted on 15 Apr 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Title: Interaction of picolinamide fungicide primary metabolites UK-2A and CAS-649 with the**
2 **cytochrome *bc*₁ complex Qi site: mutation effects and modeling in *Saccharomyces cerevisiae***

3

4 **Running title: Effects of Qi site mutations on UK-2A and CAS-649**

5

6 **David H. Young,^{a,*} Brigitte Meunier^b and Nick X. Wang^a**

7

8 **Abstract**

9

10 **BACKGROUND:** Fenpicoxamid and florylpicoxamid are picolinamide fungicides targeting the Qi site of
11 the cytochrome *bc*₁ complex, *via* their primary metabolites UK-2A and CAS-649, respectively. We
12 explore binding interactions and resistance mechanisms for picolinamides, antimycin A and ilicicolin H in
13 yeast by testing effects of cytochrome *b* amino acid changes on fungicide sensitivity and interpreting
14 results using molecular docking.

15

16 **RESULTS:** Effects of amino acid changes on sensitivity to UK-2A and CAS-649 were similar, with highest
17 resistance associated with exchanges involving G37 and substitutions N31K and L198F. These changes,
18 as well as K228M, also affected antimycin A, while ilicicolin H was affected by changes at G37 and L198,
19 as well as Q22E. N31 substitution patterns suggest that a lysine at position 31 introduces an electrostatic
20 interaction with neighboring D229, causing disruption of a key salt-bridge interaction with
21 picolinamides. Changes involving G37 and L198 imply resistance primarily through steric interference.
22 G37 changes also showed differences between CAS-649 and UK-2A or antimycin A with respect to
23 branched versus unbranched amino acids. N31K and substitution of G37 by large amino acids reduced
24 growth rate substantially while L198 substitutions showed little effect on growth.

25

26 **CONCLUSION:** Binding of UK-2A and CAS-649 at the Qi site involves similar interactions such that general
27 cross-resistance between fenpicoxamid and florylpicoxamid is anticipated in target pathogens. Some
28 resistance mutations reduced growth rate and could carry a fitness penalty in pathogens. However,
29 certain changes involving G37 and L198 carry little or no growth penalty and may pose the greatest risk
30 for resistance development in the field.

31

32

33 **Keywords:** UK-2A; CAS-649; picolinamide fungicide; antimycin A; ilicicolin H; resistance mutation

34

35

36 **Correspondence to: David H Young, Corteva Agriscience, 9330 Zionsville Road, Indianapolis, IN 46268,*
37 *USA. E-mail: david.young@corteva.com*

38 ^a *Corteva Agriscience, 9330 Zionsville Road, Indianapolis, IN 46268, USA*

39 ^b *Université Paris-Saclay, CEA, CNRS, Institute for Integrative Biology of the Cell (I2BC), 91198, Gif-sur-*
40 *Yvette, France*

41

42 1 INTRODUCTION

43 Fenpicoxamid (Fig. 1) was recently introduced as a new fungicide which acts at the Qi site of the
44 cytochrome bc_1 complex (complex III) in the mitochondrial electron transport chain.¹ As the first Qil
45 fungicide with efficacy against Ascomycete pathogens, fenpicoxamid provides a new mode of action for
46 control of Septoria leaf blotch (*Zymoseptoria tritici*) and other diseases in cereals, as well as black
47 Sigatoka (*Mycosphaerella fijiensis*) in bananas. Chemically, fenpicoxamid belongs to a new
48 “picolinamide” class of fungicides and is derived from the natural product UK-2A,² a highly potent Qi site
49 inhibitor.^{3,4} UK-2A is structurally related to the Qi site inhibitor antimycin A and is produced by
50 fermentation of *Streptomyces* s. 517-02.¹ Conversion of UK-2A to fenpicoxamid is achieved by addition
51 of an isobutyryloxymethyl ether substituent. This modification to the hydroxyl substituent on the
52 picolinamide head group increases stability on the plant surface, and results in greatly improved disease
53 control.¹ Within the plant or pathogen, fenpicoxamid is then converted back to UK-2A, as the
54 fungicidally active species.

55
56 Exploration of structure-activity relationships for UK-2A analogs enabled the design of a fully synthetic
57 new Qil fungicide florylpicoxamid (Fig. 1), which is currently in commercial development and lacks the
58 complex 9-member macrocycle of UK-2A.⁵ The primary metabolite CAS-649 is responsible for activity of
59 florylpicoxamid, and is generated *in vivo* by removal of the acetyl substituent on the picolinamide head.⁵
60 An important feature of florylpicoxamid is its strong efficacy against a broad range of Ascomycetes,
61 including *Botrytis*, *Colletotrichum* and *Alternaria* species, as well as powdery mildew pathogens.⁶ This
62 will expand the use of picolinamide fungicides to many additional markets and provide a new mode of
63 action for controlling important diseases in row crops, fruits and vegetables.

64

65 Fungicides which target the bc_1 complex act at either the Qo or Qi sites that are formed by the central
66 subunit of the multisubunit complex, namely cytochrome *b* (cyt *b*). They are referred to as QoI and QiI
67 fungicides, respectively. Resistance to QoIs and QiIs is most commonly caused by cyt *b* changes at their
68 respective target sites, such that resistance mutations at the Qo site do not reduce efficacy of QiI
69 fungicides, and vice-versa.^{7,8} In the specific case of picolinamide fungicides, the G143A change
70 responsible for strobilurin-resistance does not reduce efficacy of fenpicoxamid and florylpicoxamid in *Z.*
71 *tritici*,^{1,6,7} and there is some evidence for negatively correlated cross-resistance.⁸ Strobilurins represent
72 the largest class of bc_1 complex inhibitors, which when first introduced were believed to be at low risk
73 for resistance development because the target protein, cyt *b*, is encoded by a mitochondrial gene.
74 Multiple applications per season and use in solo applications rather than mixtures likely contributed to a
75 rapid development of resistance.⁹ The mutation G143A is responsible for efficacy loss in most cases of
76 strobilurin resistance, while the F129L mutation confers lower levels of field resistance in *Phakopsora*
77 *pachyrhizi* (soybean rust), *Pyrenophora teres* (net blotch in barley), *Alternaria solani* (potato/tomato
78 early blight) and *Pythium aphanidermatum* (Pythium blight of turfgrass).^{10,11} Effects of these mutations
79 have been explained by their impacts on binding interactions at the Qo site in cyt *b* molecular docking
80 experiments.¹² The QiI fungicides cyazofamid and amisulbrom were launched in 2001 and 2003,
81 respectively, and are used to control Oomycete diseases. Cyt *b* modifications which cause resistance
82 have been reported in *Plasmopara viticola* (grape downy mildew), involving the insertions E203-DE-
83 V204 and E203-VE-V204, as well as substitution L201S.¹³ Another Oomycete fungicide, ametoctradin, is
84 believed to bind to both Qo and Qi sites¹⁴ and the Qi site resistance mutation S34L has been reported.¹⁵
85
86 Baseline sensitivity monitoring for fenpicoxamid in *Z. tritici* populations has shown a relatively narrow
87 range of sensitivities and no detectable resistance.⁹ In the laboratory, however, mutants resistant to
88 fenpicoxamid and UK-2A have been isolated in yeast and *Z. tritici*: amino acid substitutions N31K, G37C

89 and L198F at the Qi site conferred resistance in *Saccharomyces cerevisiae*,⁷ and the mutation G37V was
90 identified in lab mutants of *Zymoseptoria tritici* by selection in the presence of fenpicoxamid using an
91 experimental evolution approach.⁸ Target site changes conferring resistance to fenpicoxamid have not
92 yet been reported in field populations. However, the aforementioned examples of resistance to
93 commercialized bc_1 complex inhibitor fungicides, as well as lab studies with fenpicoxamid, suggest that
94 target site resistance to both fenpicoxamid and florypicoxamid must be anticipated and appropriate
95 resistance management practices implemented. To preserve its effectiveness, fenpicoxamid should be
96 applied only once per season and in mixture with another effective mode of action. Suitable resistance
97 management strategies will also be employed for florypicoxamid when it is commercialized, as
98 appropriate for the targeted disease and crop.

99

100 The isolation of resistant mutants in yeast, or modification of binding sites through introduction of
101 specific amino acid changes, combined with crystallographic and modeling studies has been a successful
102 strategy in understanding binding site interactions for a wide variety of bc_1 complex inhibitors.¹⁶⁻¹⁹
103 Moreover, *S. cerevisiae* is the only fungal species currently amenable to introduction of specific amino
104 acid substitutions in the mitochondrially encoded *cyt b*. This allows proposed inhibitor docking poses to
105 be evaluated experimentally, resulting in more robust binding site models. In addition, mechanisms
106 responsible for target site resistance can be explored by analyzing effects of amino acid exchange. In our
107 current study, we use yeast as a model system to compare binding site interactions and resistance
108 mechanisms for UK-2A and CAS-649, as well as two other natural product Qi site inhibitors antimycin A
109 and ilicicolin H, providing insights into similarities and differences between these structurally different
110 ligands.

111

112 The ability of amino acid changes to disrupt target site interactions without significant fitness penalty
113 will likely determine the identity of resistance mutations which might emerge in pathogen populations
114 exposed to picolinamide fungicides. Consequently, similarities and differences between UK-2A and CAS-
115 649 with respect to effects of mutations on sensitivity are of practical importance in assessing the
116 likelihood of cross-resistance in pathogens on certain crops where both fenpicoxamid and
117 florylpicoxamid might be registered, and in providing recommendations for resistance management.
118

119 **2 MATERIALS AND METHODS**

120 **2.1 Cytochrome *b* mutants**

121 A wild-type diploid strain of *S. cerevisiae* containing an intron-free version of the cytochrome *b* gene was
122 used as the parental strain. Mutants containing the G37C, L198F and N31K changes were isolated by
123 selection on fenpicoxamid or antimycin A.⁷ All other mutants were generated by site-directed
124 mutagenesis and mitochondrial transformation by methods described previously.^{20,21}

125

126 **2.2 Fungitoxicity assays**

127 Compounds were dissolved in DMSO and 3-fold dilutions in DMSO were prepared by serial dilution.
128 Aliquots (1 μ L) of each dilution were transferred to wells of flat-bottomed 96-well microtiter plates, with
129 three replicate wells for each treatment, and DMSO alone in control wells. Wells then received 100 μ L of
130 YPEth broth (10 g of yeast extract, 20 g of Bacto peptone and 2% (v/v) of ethanol per liter). Cells taken
131 from overnight cultures grown on YPD agar (Becton, Dickinson and Co., Sparks, MD, USA) at 30 °C were
132 suspended in YPEth broth at a cell density of 2×10^5 cells mL⁻¹. Wells were inoculated with 100 μ L of cell
133 suspension and the plates were incubated at 30 °C for 45 h. After mixing well contents to provide
134 homogeneous cell suspension, growth was measured using a NEPHELOstar Galaxy plate reader (BMG
135 Labtech, Cary, NC, USA) and 50% effective concentration (EC50) values for growth inhibition were
136 determined using the CALCUSYN program (Biosoft, Cambridge, UK). Mutants were tested in two or more
137 independent experiments and average EC50 values \pm SEM determined across all replicates.

138

139 **2.3 Molecular docking of ligands at the *S. cerevisiae* Qi site**

140 The crystal structure of the *S. cerevisiae* cytochrome *bc*₁ complex (PDB: 1EZV)²² was used as the protein
141 target. The structure was first parameterized and then refined in the QuickPrep of MOE (CCG Inc.,
142 Montreal, Canada) using Amber14:EHT force field. After the protein system had been well prepared, UK-

143 2A, CAS-649, antimycin A and ilicicolin H were docked into the Qi site using the Induce Fit docking
144 method with MMFF94x force field in MOE. Each docking run generated 30 docking poses. The most
145 likely binding pose for each molecule was selected based on measured target site mutation effects and
146 docking score.

147

148 **2.4 Evaluation of kinetic growth rate**

149 Cells from overnight cultures grown on YPD medium at 30 °C were suspended in YPEth broth and
150 adjusted to $A_{600} = 0.3$ in a 1 cm cuvette. Cell suspensions (100 μ L) were added to wells of a 96-well flat-
151 bottom tissue culture plate (Cat. No. 655180, Greiner Bio-One, Monroe, NC, USA) with 8 replicate wells
152 per strain. The plate was sealed with Breathe-Easy™ sealing film (Cat. No. BEM-1, Diversified Biotech,
153 Dedham, MA, USA). Growth curves were measured in a Cytation 3 instrument (BioTek Instruments,
154 Winooski, VT, USA) at 30 °C using continuous double orbital shaking at 425 cpm with 3 mm amplitude
155 and measurements taken at 1 h intervals. Specific growth rates were determined during exponential
156 growth over the 6 - 11 h period after start of the experiment and used to calculate doubling time. All
157 experiments were repeated twice with 8 replicates per experiment and growth rate \pm SD determined
158 across all replicates.

159

160 **3 RESULTS AND DISCUSSION**

161 **3.1 Qi site mutants**

162 The Qi site mutant strains used in the study were selected based on various criteria (Table 1). Different
163 amino acid substitutions at N31, G37 and L198 were designed to explore mechanisms responsible for
164 reduced sensitivity of N31K, G37C/V and L198F, which confer resistance to UK-2A,^{7,8} to characterize
165 effects of these changes on CAS-649, antimycin A and illicicolin H, and evaluate natural variants found in
166 fungal pathogens. Mutations Y16A and M221F were studied to investigate proposed roles in ligand
167 binding of Y16²³ and F221,³³ a natural variant in *Z. tritici*. Several additional mutations known to confer
168 reduced sensitivity to other Qi site ligands were also included to evaluate potential cross-resistance to
169 UK-2A and CAS-649.

170

171 **3.2 Effects of mutations on inhibitor sensitivity**

172 Despite considerable structural differences between UK-2A and CAS-649, the effects of mutations at the
173 Qi site on sensitivity were quite similar overall for the two molecules (Figs. 2 and 3). Among the various
174 amino acid changes examined, those associated with high resistance factors for the two molecules
175 involved various G37 changes, and substitutions N31K and L198F. The same changes affected sensitivity
176 to antimycin A, although resistance factors were lower (Fig. 4). Sensitivity to antimycin A was also
177 reduced by mutation K228M, which did not affect UK-2A or CAS-649. The similarity in mutation effects
178 for UK-2A and CAS-649 in yeast suggests that cross-resistance between fenpicoxamid and
179 florylpicoxamid is most likely to be observed in plant pathogens should target site resistance develop in
180 the field.

181

182 Illicicolin H is structurally different from UK-2A, antimycin A and CAS-649, and is known to be affected by
183 a number of Qi site mutations including S20I/L/T, Q22E/T, G37C/D/S, L198F and M221E/Q.^{19,25,26,30} The

184 strong resistance observed in our study for Q22E, G37C/S and L198F (Fig. 5) is in agreement with
185 previous work. M221Q conferred only a low level of resistance, consistent with the weak effect reported
186 for this mutation at the enzyme level.²⁶ Whereas cross-resistance between illicicolin H, antimycin A and
187 picolinamides was seen for L198F and G37 substitutions, differences between illicicolin H and the other
188 ligands were apparent for N31K, which did not confer resistance to illicicolin H, and for Q22E which
189 strongly reduced sensitivity to illicicolin H but did not affect the other inhibitors.

190

191 **3.3 Binding poses for UK-2A, CAS-649 and antimycin A**

192 The preferred docking poses for UK-2A and CAS-649 show the flat, polar picolinamide head located deep
193 within the binding pocket and the bulky hydrophobic tail portions extending towards the outside of
194 pocket (Figs. 6 and 7). For the head moiety, both molecules use the protonated pyridine N to form a
195 strong salt bridge with the deprotonated carboxylic side chain of D229. As reported previously, this pose
196 for UK-2A is strongly supported by analysis of yeast mutation effects on activity of analogs with different
197 head groups and the importance of the pyridyl N for activity.⁷ Different binding interactions are
198 employed by antimycin A (Fig. 8), which instead forms two hydrogen bonds with D229 using the
199 hydroxyl and formamide substituents of the phenyl head. The carbonyl of the formylamino group also
200 forms a H-bond network via a water molecule with K228 and N31. UK-2A, CAS-649 and antimycin A
201 share the same amide linker which forms an intramolecular hydrogen bond between the amide N and
202 the OH substituent on the aryl ring to stabilize the co-planar conformation of the head. The tail portions
203 of UK-2A and antimycin A share the same bislactone core but differ with respect to side chain
204 substituents. UK-2A has 2-methyl propanoate and benzyl at the C-7 and C-8 positions, respectively,
205 whereas antimycin A has 3-methyl butanoate and a hexyl moiety. Binding interactions for the tail
206 portions of UK-2A, CAS-649 and antimycin A are primarily hydrophobic and the ligands are near residues
207 G37 and L198, which can explain altered sensitivity in response to substitution of these residues.

208 In a *Z. tritici* binding model, an alternative binding orientation with the picolinamide head of UK-2A
209 flipped by 180° had been postulated primarily to explain high potency towards *Z. tritici* of certain
210 analogs with alternative head groups which lack the ability to form a salt bridge, yet are highly active
211 against *Z. tritici*.³³ In this pose, the OH substituent on the picolinamide head binds to D229, rather than
212 the pyridyl N, and this pose is enabled by pi-pi stacking with F221 as a key interaction. Since in yeast the
213 pi-pi stacking interaction is not feasible due to a methionine instead of a phenylalanine at position 221,
214 we used the yeast system to compare these two amino acids in terms of their impact on sensitivity to
215 UK-2A and CAS-649, based on the assumption that an increase in sensitivity should be observed if pi-pi
216 stacking with F221 was enabled to mimic the *Z. tritici* binding pocket. The fact that the M221F
217 substitution did not alter sensitivity to UK-2A and CAS-649 (Figs. 2 and 3) argues against the suggested
218 pi-pi interaction with F221 in *Z. tritici* as a key interaction for analogs with the picolinamide head. A
219 critical role for F221 in binding is also not supported by sequence analysis of other pathogens since
220 some pathogens which are highly sensitive to picolinamides, such as *Erysiphe necator*, *Blumeria graminis*
221 and *Puccinia triticina*, have M221 like yeast, or I221 in the case of *Phakopsora pachyrhizi* (Table 2). While
222 the potency against *Z. tritici* of picolinamide analogs with alternative head groups that lack the ability to
223 form a salt bridge may indeed involve an alternative pose enabled by pi-pi stacking with F221, for
224 picolinamides like UK-2A and CAS-649 the pose involving the salt bridge interaction between the pyridyl
225 N and D229 is the most likely.

226 Another amino acid which has been proposed to play a possible role in binding of UK-2A is Y16,²³ a
227 residue which is highly conserved across fungi. In the binding pocket this residue lies in close proximity
228 to the benzyl substituent at C8 of the bislactone ring with the possibility of participation in a pi-pi
229 stacking interaction. Although CAS-649 and UK-2A adopt a similar binding pose, pi-pi stacking with Y16
230 would be less likely for CAS-649 because the branched *p*-fluoro phenyl substituent of CAS-649 is farther
231 from Y16 than the benzyl of UK-2A. To test the possible role for Y16 in binding, we evaluated the Y16A

232 mutation since introduction of an alanine would negate any pi-pi stacking interaction. In fact, the Y16A
233 change did not reduce sensitivity to UK-2A and CAS-649, but instead increased sensitivity, particularly
234 for CAS-649 (Figs. 2 and 3). These results suggest that interaction with Y16 is not required for binding of
235 picolinamides.

236

237 **3.4 Binding pose for ilicicolin H**

238 The most likely binding pose of ilicicolin H (Fig. 9) overlaps those of UK-2A, CAS-649 and antimycin A but
239 involves different ligand-protein interactions. A key interaction involves H-bonding of the phenol head
240 to Q22. The high level of resistance conferred by the Q22E mutation can be explained by loss of this
241 interaction. In contrast to Q22E, the Q22M change reduced sensitivity to ilicicolin H only slightly
242 suggesting that a methionine can substitute for glutamine in H-bonding. The hydroxyl group in the
243 middle ring of ilicicolin H forms an intramolecular hydrogen bond with the nearby ketone group to
244 position the bulky fused ring portion between the G37 and L198 residues. As for picolinamides and
245 antimycin A, proximity to these residues explains the mutation effects resulting from amino acid
246 substitution. Unlike the other ligands, no involvement of N31 or D229 is apparent in this binding pose.
247 Our binding pose differs from an earlier model in *Candida albicans* in which a key interaction of the
248 phenol moiety with N31 was proposed.³⁰ However, such an interaction is not consistent with the lack of
249 effect of N31K on sensitivity (Fig. 5). Ilicicolin H is positioned very close to both G37 and L198, with
250 closest distances of 3.6 and 3.5 Å, respectively. This close proximity suggests that replacement of these
251 amino acids by larger residues would interfere with binding due to steric effects and likely explains the
252 high levels of resistance associated with G37 changes and L198F (Fig. 5).

253

254 **3.5 N31 Mutations**

255 Since changes involving N31, G37 and L198 are the primary ones associated with reduced sensitivity to
256 UK-2A and CAS-649, we explored the mechanisms responsible for resistance by comparing the effects of
257 various substitutions at these positions. We also examined effects of these changes on growth rate in
258 the hope of gaining insight into potential fitness penalties, which could determine mutations most likely
259 to appear in pathogens in nature in response to fungicide selection pressure.

260

261 In the crystal structure of the bc_1 complex from *Bos taurus* with bound antimycin A³⁵ and in our yeast
262 models for binding of UK-2A and CAS-649, there is no direct interaction between the ligands and N31
263 (N32 in *B. taurus*). According to the bovine structure, N31 participates in an extensive H-bonding chain
264 involving other amino acids and water molecules: one water molecule forms a link between N31, K228
265 and I26, while a second water links N31, S34 and D229.³⁵ The latter residue has been proposed to bind
266 to the benzoquinone moiety of coenzyme Q6 and act as a proton donor during reduction of the
267 substrate.²² Exchange of N31 by other amino acids or substitution of S34 should disrupt this H-bonding
268 network and would likely impact D229, potentially interfering with its key role in binding of picolinamide
269 fungicides and antimycin A (Figs. 6 - 8). However, in the yeast crystal structure with the bound natural
270 substrate coenzyme Q6 (ubiquinone 6),²² N31 does not interact with D229 via water. It is not clear
271 whether the apparent differences in binding of N31 to D229 via water between the bovine structure
272 with bound antimycin A and the yeast structure with bound coenzyme Q6 are a consequence of
273 different induced fit conformational changes during binding of antimycin A versus coenzyme Q6, the
274 different protein sources, or methods for preparation of crystals. It is significant that resistance to UK-
275 2A, CAS-649 and antimycin A was associated with introduction of the basic lysine side chain and not with
276 acidic (aspartate), polar neutral (serine) or glycine residues, or with the S34L exchange (Figs. 2-4). If
277 resistance from substitution of N31 was due to disruption of a H-bonding network, these other
278 mutations would also be expected to confer resistance. A more likely explanation is that introduction of

279 a lysine at position 31 affects D229 by electrostatic lysine-aspartate attraction and prevents or weakens
280 the critical ligand binding interactions involving D229. In Ascomycete and Basidiomycete pathogens, N31
281 is conserved (Table 2), and in yeast its replacement by other amino acids resulted in markedly lower
282 growth rates (Table 3). It is possible that this reflects a loss in efficiency of the bc_1 complex resulting
283 from an effect, via the H-bonding network mentioned above, on the nearby D229 through interference
284 with its proposed catalytic function in substrate reduction. These considerations suggests that the N31K
285 mutation, although conferring a high level of resistance in yeast, might carry a considerable fitness
286 penalty if it were to occur in field populations of plant pathogens.

287

288 **3.6 G37 Mutations**

289 Among the G37 modifications explored, a high level of resistance to both UK-2A and CAS-649 was found
290 for G37V/C/I/L, while G37M conferred a moderate level of resistance (Figs. 2 and 3). G37V/C/I/L also
291 conferred resistance to antimycin A (Fig. 4) although resistance factors (RFs) were lower than for
292 picolinamides. The data suggest a rough correlation between RFs and amino acid size, consistent with
293 steric effects on binding since the highest RFs were found for G37V/C/I/L and the smallest for G37A/S.
294 However, branching at the β carbon atom of amino acids also impacted the level of resistance in the
295 case of UK-2A and antimycin A, since the branched amino acids valine and isoleucine produced
296 significantly higher RFs than unbranched residues of comparable size, cysteine and leucine. Interestingly,
297 the differential effect of branched and unbranched amino acids observed for these natural products,
298 which have the same 9-membered macrocyclic scaffold, was not apparent for CAS-649. These data
299 suggest that subtle differences in binding interactions may exist between UK-2A and CAS-649 in the
300 region near G37. In the case of ilicicolin H, the exchange of G37 by other amino acids resulted in even
301 higher RFs than observed for the other ligands, suggesting even more stringent steric limitations in this
302 region of the binding site. The G37M exchange presented a curious exception to the general correlation

303 between level of resistance and residue size. Although methionine was the largest amino acid among
304 those examined, the G37M change resulted in lower RFs against all ligands than was observed for
305 substitution of G37 by various smaller amino acids. The reason for this is unknown but these data
306 suggest that the G37M change may alter the binding pocket in such a way as to modulate the effect of
307 size alone on ligand binding.

308

309 Pathogens which are most sensitive to fenpicoxamid and florylpicoxamid have a glycine or an alanine in
310 position 37 (Table 2). This appears consistent with the relatively small effect of the yeast G37A mutation
311 on sensitivity. *Fusarium* spp. are relatively insensitive to fenpicoxamid and florylpicoxamid (unpublished
312 results, Corteva Agriscience) and contain L37 and M198. In yeast, the change L198M showed little effect
313 on sensitivity to all ligands and no reduction in growth rate. However, G37L caused very high resistance
314 to the picolinamides, suggesting that the L37 variant may explain insensitivity of *Fusarium* spp. Yeast
315 mutant G37L+L198M was designed to mimic the *Fusarium* Qi site more closely, and showed interesting
316 differences from the G37L mutant in terms of inhibitor sensitivity and growth rate. For G37L+L198M,
317 resistance to UK-2A and antimycin A was lower than for the G37L mutant. However, this was not the
318 case for CAS-649 and ilicicolin H which had equivalent RFs for both the single and double mutant.
319 Moreover, in the double mutant, the L198M change compensated partially for a large reduction in
320 growth rate associated with the G37L change. One can speculate that occurrence of these two variations
321 together in *Fusarium* spp. may overcome effects of an otherwise deleterious G37L change, if it occurred
322 alone.

323

324 Mutations involving G37 impacted growth rate to varying degrees (Table 3). Substitution of glycine by
325 relatively small amino acids, alanine and serine, did not affect growth rate. Among mutants which
326 showed strong resistance to picolinamides, G37S showed no reduction in growth rate, G37C showed a

327 slight reduction in growth rate, while growth of G37V/I/M was substantially reduced. It is interesting to
328 note that G37C was the mutation identified in yeast through selection in the presence of fenpicoxamid⁷
329 although G37V/I mutants have higher resistance factors. Similarly, for ilicicolin H G37V/I mutants had
330 higher resistance factors than G37S. Yet G37S/D were the mutants isolated by selection on ilicicolin.²⁵ It
331 is likely that the smaller fitness penalties in terms of growth rate for G37C/S favored their isolation in
332 these cases. By contrast, in *Z. tritici* the only mutants obtained by selection on fenpicoxamid using
333 experimental evolution contained the G37V mutation.⁸ This mutation, which produced a high level of
334 resistance to UK-2A, caused a lower efficiency of *bc*₁ complex in a cell-free *Z. tritici* mitochondrial assay
335 but, unlike in yeast, did not affect the *in vitro* growth rate. These examples highlight the fact that the
336 particular mutant(s) selected in the laboratory or field may differ depending on the organism, and is
337 dependent on both the associated level of resistance and effect on growth or fitness.

338

339 3.7 L198 Mutations

340 The L198F mutation was found previously by selection in the presence of fenpicoxamid⁷ and ilicicolin
341 H,²⁵ and showed cross-resistance to antimycin A. This mutation also conferred resistance to CAS-649
342 (Fig. 3). Binding poses indicate a similar distance of 3.8 or 3.9 Å between L198 and these various ligands
343 (Figs. 6 – 9). L198F introduces the largest amino acid among the variants tested and showed the greatest
344 mutation effect against all compounds. Other changes, involving smaller amino acids, had little or no
345 effect on sensitivity to the picolinamides and antimycin A and relatively small effects on ilicicolin H.
346 These data suggest that L198F confers resistance due to a steric clash with the ligands

347

348 For the L198F mutant, RFs for UK-2A and CAS-649 were lower than for some G37 mutants: G37V/I/L in
349 the case of UK-2A, and G37I/L in the case of CAS-649. However, in contrast to those changes involving
350 G37, substitution of L198F did not reduce growth rate (Table 3). Occurrence of F198 as a natural variant

351 in *Rhizoctonia solani* has been suggested as a possible explanation for the low sensitivity of this
352 pathogens to fenpicoxamid.³⁶ The combination of a fairly high level of resistance and little impact on
353 growth in the yeast model highlights L198F as a mutation of interest which potentially could lead to field
354 resistance to picolinamide fungicides in plant pathogens.

355

356 **5 CONCLUSIONS**

357 Comparison of picolinamides with antimycin A and ilicicolin H showed both similarities and differences
358 in mutation effects which could be explained by overlapping binding poses. For UK-2A and CAS-649,
359 mutation effects were similar and molecular docking at the Qi site indicated common binding
360 interactions. This suggests that cross-resistance between fenpicoxamid and florypicoxamid is likely in
361 plant pathogens should target site resistance develop in the field. Florypicoxamid is intended for use to
362 control diseases on many crops not targeted by fenpicoxamid, such that cross-resistance would have no
363 practical implications. However, the likelihood of cross-resistance is an important consideration for
364 resistance management in the case of crops like wheat and bananas where both compounds might be
365 used in future years. Certain changes involving G37 and substitution L198F were associated with high
366 levels of resistance and little or no effect on growth rate, as well as insensitivity in some species due to
367 natural variants. Notwithstanding obvious limitations in use of a yeast model to predict resistance and
368 disease-associated fitness in a different organism, our results highlight these residues as ones most likely
369 to be involved in target site resistance to picolinamide fungicides in pathogens under selection pressure
370 in the field.

371

372 **ACKNOWLEDGMENTS**

373 The authors would like to thank the many colleagues at Corteva Agriscience involved in the discovery
374 and development of florylpicoxamid. We would also like to acknowledge Meiji Seika Pharma Co. Ltd. for
375 their collaboration on the development of fenpicoxamid.

376

377 **SUPPORTING INFORMATION**

378 Supporting information may be found in the online version of this article.

379

380 **REFERENCES**

- 381 1. Owen WJ, Yao C, Myung K, Kemmitt G, Leader A, Meyer KG, Bowling AJ, Slanec T and Kramer VJ,
382 Biological characterization of fenpicoxamid, a new fungicide with utility in cereals and other
383 crops. *Pest Manag Sci* **73**:2005-2016 (2017).
- 384 2. Ueki M, Abe K, Hanafi K, Shibata K, Tanaka T and Taniguchi M, UK-2A, B, C and D, novel
385 antifungal antibiotics from *Streptomyces* sp. 517-02. I. Fermentation, isolation and biological
386 properties. *J Antibiot* **49**:639-643 (1996).
- 387 3. Ueki M and Taniguchi M, The mode of action of UK-2A and UK-3A, novel antifungal antibiotics
388 from *Streptomyces* sp. 517-02. *J Antibiot* **50**:1052-1057 (1997).
- 389 4. Machida K, Takimoto H, Miyoshi H and Taniguchi M, UK-2A, B, C and D, novel antifungal
390 antibiotics from *Streptomyces* sp. 517-02. V. Inhibition mechanism of bovine heart
391 mitochondrial cytochrome *bc*₁ by the novel antibiotic UK-2A. *J Antibiot* **52**:748-753 (1999).
- 392 5. Meyer KG, Bravo-Altamirano K, Herrick J, Loy BA, Yao C, Nugent B, Buchan Z, Daeuble JF,
393 Heemstra R, Jones DM, Wilmot J, Lu Y, DeKorver K, DeLorbe J and Rigoli J, Discovery of
394 florylpicoxamid, a mimic of a macrocyclic natural product. *Bioorg Med Chem* **50**: Article 116455
395 (2021).


- 396 6. Yao C, Meyer KG, Gallup C, Bowling AJ, Hufnagl A, Myung K, Lutz J, Slanec T, Pence HE, Delgado J
397 and Wang NX, Florylpicoxamid, a new picolinamide fungicide with broad spectrum activity. *Pest*
398 *Manag Sci* **77**: 4483-4496 (2021).
- 399 7. Young DH, Wang NX, Meyer ST, and Avila-Adame C, Characterization of the mechanism of action
400 of the fungicide fenpicoxamid and its metabolite UK-2A. *Pest Management Science* **74**: 489-498
401 (2018).
- 402 8. Fouché G, Michel T, Lalève A, Wang NX, Young DH, Meunier B, Debieu D, Fillinger S and Walker
403 A-S. Directed evolution predicts cytochrome *b* G37V target site modification as probable
404 adaptive mechanism towards the Qil fungicide fenpicoxamid in *Zymoseptoria tritici*. *Environ*
405 *Microbiol* <https://doi.org/10.1111/1462-2920.15760> (2021).
- 406 9. Young D, Kemmitt G and Leader A, Fenpicoxamid provides a new target site for control of
407 Septoria leaf blotch in cereals, in *Modern Fungicides and Antifungal Compounds IX*, ed. by
408 Deising HB, Fraaije B, Mehl A, Oerke EC, Sierotzki H and Stammler G. DPG Verlag, Braunschweig,
409 Germany, pp. 279-284 (2020).
- 410 10. Sierotzki H, Respiration inhibitors: complex III, in *Fungicide Resistance in Plant Pathogens*, ed. by
411 Ishii H and Hollomon DW. Springer, Tokyo, pp. 119-143 (2015).
- 412 11. Klosowski AC, May De Mio LL, Miessner S, Rodrigues R and Stammler G, Detection of the F129L
413 mutation in the cytochrome *b* gene in *Phakopsora pachyrhizi*. *Pest Manag Sci* **72**: 1211-1215
414 (2016).
- 415 12. Link T, Iwata M, Bjorkman J, Van der Spoel D, Stocker A and Iwata S, Molecular modeling of
416 inhibitors at Qi and Qo sites in cytochrome *bc₁* complex, in *Chemistry of Crop Protection*, ed. by
417 Voss G and Ramos G. Wiley-VCH, Weinheim, Germany, pp. 110-127 (2003).

- 418 13. R4P, Note Technique Commune Resistances 2020 Maladies de la Vigne: Mildiou, Oidium,
419 Pourriture Grise, Black-Rot. Available: [https://www.vignevin.com/wp-](https://www.vignevin.com/wp-content/uploads/2020/01/Note-technique-commune-vigne-2020-Vdef.pdf)
420 [content/uploads/2020/01/Note-technique-commune-vigne-2020-Vdef.pdf](https://www.vignevin.com/wp-content/uploads/2020/01/Note-technique-commune-vigne-2020-Vdef.pdf) [6 January 2020].
- 421 14. Dreinert A, Wolf A, Mentzel T, Meunier B and Fehr M, The cytochrome *bc1* complex inhibitor
422 Ametoctradin has an unusual binding mode. *Biochim Biophys Acta* **1859**: 567-576 (2018).
- 423 15. Fontaine S, Remuson F, Caddoux L and Barres B, Investigation of the sensitivity of *Plasmopara*
424 *viticola* to amisulbrom and ametoctradin in French vineyards using bioassays and molecular
425 tools. *Pest Manag Sci* **75**:2115-2123 (2019).
- 426 16. Gao X, Wen X, Esser L, Quinn B, Yu L, Yu C-A and Xia D, Structural basis for the quinone reduction
427 in the *bc₁* complex: a comparative analysis of crystal structures of mitochondrial cytochrome *bc₁*
428 with bound substrate and inhibitors at the Q_i site. *Biochemistry* **42**:9067-9080 (2003).
- 429 17. Esser L, Quinn B, Li Y-F, Zhang M, Elberry M, Yu L, Yu C-A and Xia D, Crystallographic studies of
430 quinol oxidation site inhibitors for the cytochrome *bc₁* complex. *J Mol Biol* **341**: 281-302 (2004).
- 431 18. Fisher N and Meunier B, Molecular basis of resistance to cytochrome *bc₁* inhibitors. *FEMS Yeast*
432 *Res* **8**: 183-192 (2008).
- 433 19. Mounkoro P, Michel T, Benhachemi R, Surpateanu G, Iorga BI, Fisher N and Meunier B,
434 Mitochondrial complex III Q_i-site inhibitor resistance mutations found in laboratory selected
435 mutants and field isolates. *Pest Manag Sci* **75**: 2107-2114 (2019).
- 436 20. Hill P, Kessl K, Fisher N, Meshnick S, Trumpower BL and Meunier B, Recapitulation in
437 *Saccharomyces cerevisiae* of cytochrome *b* mutations conferring resistance to atovaquone in
438 *Pneumocystis jiroveci*. *Antimicrob Agents Chemother* **47**:2725-2731 (2003).
- 439 21. Fisher N, Brown AC, Sexton G, Cook A, Windass J and Meunier B, Modeling the Q_o site of crop
440 pathogens in *Saccharomyces cerevisiae* cytochrome *b*. *Eur J Biochem* **271**:2264-2271 (2004).

- 441 22. Hunte C, Koepke J, Lange C, Rossmann T and Michel H, Structure at 2.3 Å resolution of the
442 cytochrome *bc*₁ complex from the yeast *Saccharomyces cerevisiae* co-crystallized with an
443 antibody Fv fragment. *Structure* **8**:669-684 (2000).
- 444 23. Owen WJ, Meyer KG, Meyer ST, Li F, Slanec TJ, Wang NX and Yao C, Synthesis and biological
445 activity of analogs of the antifungal antibiotic UK-2A. II. Impact of modifications to the
446 macrocycle benzyl position. *Pest Manag Sci* **75**:1831-1846 (2019).
- 447 24. Di Rago J-P, Perea J and Colson A-M, DNA sequence analysis of diuron-resistant mutations in the
448 mitochondrial cytochrome *b* gene of *Saccharomyces cerevisiae*. *FEBS Lett* **208**:208-210 (1986).
- 449 25. Ding MG, Di Rago JP and Trumpower BL, Investigating the Q_n site of the cytochrome *bc*₁
450 complex in *Saccharomyces cerevisiae* with mutants resistant to ilicicolin H, a novel Q_n site
451 inhibitor. *J Biol Chem* **281**:36036-36043 (2006).
- 452 26. Rotsaert FAJ, Ding MG and Trumpower BL, Differential efficacy of inhibition of mitochondrial
453 and bacterial cytochrome *bc*₁ complexes by center N inhibitors antimycin, ilicicolin H and
454 funiculosin. *Biochim Biophys Acta* **1777**:211-219 (2008).
- 455 27. Di Rago JP and Colson A-M, Molecular basis for resistance to antimycin and diuron, Q-cycle
456 inhibitors acting at the Q_i site in the mitochondrial ubiquinol-cytochrome *c* reductase in
457 *Saccharomyces cerevisiae*. *J Biol Chem* **263**:12564-12570 (1988).
- 458 28. Brasseur G and Brivet-Chevillotte P, Characterization of mutations in the mitochondrial
459 cytochrome *b* gene of *Saccharomyces cerevisiae* affecting the quinone reductase site (Q_N). *Eur J*
460 *Biochem* **230**:1118-1124 (1995).
- 461 29. Lukens AK, Heidebrecht Jr RW, Mulrooney C, Beaudoin JA, Comer E, Duvall JR, Fitzgerald ME,
462 Masi D, Galinsky K, Scherer CA, Palmer M, Munoz B, Foley M, Schreiber SL, Wiegand RC and
463 Wirth DF, Diversity-oriented synthesis probe targets *Plasmodium falciparum* cytochrome *b*

- 464 ubiquinone reduction site and synergizes with oxidation site inhibitors. *J Infect Dis* **211**:1097-
465 1103 (2015).
- 466 30. Singh SB, Liu W, Li X, Chen T, Shafiee A, Card D, Abruzzo G, Flattery A, Gill C, Thompson JR,
467 Rosenbach M, Dreikorn S, Hornak V, Meinz M, Kurtz M, Kelly R and Onishi JC, Antifungal
468 spectrum, in vivo efficacy, and structure-relationship of ilicicolin H. *ACS Med Chem Lett* **3**: 814-
469 817 (2012).
- 470 31. Di Rago J-P, Perea J and Colson A-M, Isolation and RNA sequence analysis of cytochrome *b*
471 mutants resistant to funiculosin, a center I inhibitor of the mitochondrial ubiquinol-cytochrome
472 *c* reductase in *Saccharomyces cerevisiae*. *FEBS Lett* **263**:93-98 (1990).
- 473 32. Brasseur G, Saribas AS and Daldal F, A compilation of mutations located in the cytochrome *b*
474 subunit of the bacterial and mitochondrial *bc*₁ complex. *Biochim Biophys Acta* **1275**:61-69
475 (1996).
- 476 33. Owen WJ, Meyer KG, Slanec TJ, Wang NX, Meyer ST, Niyaz NM, Rogers RB, Bravo-Altamirano K,
477 Herrick JL and Yao C, Synthesis and biological activity of analogs of the antifungal antibiotic UK-
478 2A. I. Impact of picolinamide ring replacement. *Pest Manag Sci* **75**:413-426 (2019).
- 479 34. Doggett JS, Nilsen A, Forquer I, Wegmann KW, Jones-Brando L, Yolken RH, Bordon C, Charman
480 SA, Katneni K, Schultz T, Burrows JN, Hinrichs DJ, Meunier B, Carruthers VB and Riscoe MK,
481 Endochin-like quinolones are highly efficacious against acute and latent experimental
482 toxoplasmosis. *Proc Natl Acad Sci* **109**:15936-15941 (2012).
- 483 35. Huang L, Cobessi D, Tung EY and Berry EA, Binding of the respiratory chain inhibitor antimycin to
484 the mitochondrial *bc*₁ complex: a new crystal structure reveals an altered intramolecular
485 hydrogen-bonding pattern. *J Mol Biol* **351**:573-597 (2005).
- 486 36. Avila-Adame C, Slanec T, Yao C and Wang N, *In vitro* insensitivity of *Rhizoctonia solani* to the
487 natural product UK-2A associated with L198F variant at the Qi target site of cytochrome *b*. APS

488 Annual Meeting, Cleveland, OH, 2019. Available:
489 <https://apsnet.confex.com/apsnet/2019/meetingapp.cgi/Paper/13802> [5 August 2019].
490



491

492


493

494

495

Figure 1. Structures of picolinamide fungicides fenpicoxamid and florypicoxamid, their primary metabolites UK-2A and CAS-649, antimycin A and illicolin H.

496




497

498 **Figure 2.** Effect of Qi site amino acid substitutions on sensitivity to UK-2A. Changes involving N31, G37
 499 and L198 are highlighted in orange, green and blue, respectively. Resistance factor values represent
 500 mean values, with SEM indicated by error bars.

501

502




503

504 **Figure 3.** Effect of Qi site amino acid substitutions on sensitivity to CAS-649. Changes involving N31, G37

505 and L198 are highlighted in orange, green and blue, respectively. Resistance factor values represent

506 mean values, with SEM indicated by error bars.

507




508

509 **Figure 4.** Effect of Qi site amino acid substitutions on sensitivity to antimycin A. Changes involving N31,

510 G37, L198 and K228 are highlighted in orange, green, blue and pink, respectively. Resistance factor

511 values represent mean values, with SEM indicated by error bars.

512



513


514 **Figure 5.** Effect of Qi site amino acid substitutions on sensitivity to ilicicolin H. Changes involving Q22,

515 G37 and L198 are highlighted in yellow, green and blue, respectively. Resistance factor values represent

516 mean values, with SEM indicated by error bars. Asterisks indicate RF > indicated value due to <50%

517 inhibition at highest test concentration of 10 $\mu\text{g mL}^{-1}$.

518



519


520

521 **Figure 6.** Docking pose of UK-2A bound to the Qi site of cytochrome *b*, showing the salt bridge

522 interaction with D229. Red arrows indicate the distances (Å) between the ligand and residues G37 and

523 L198.

524



525


526

527 **Figure 7.** Docking pose of CAS-649 bound to the Qi site of cytochrome *b*, showing the salt bridge

528 interaction with D229. Red arrows indicate the distances (Å) between the ligand and residues G37 and

529 L198.

530



531


532

533 **Figure 8.** Docking pose of antimycin A bound to the Qi site of cytochrome *b*, showing the H-bond

534 interactions with D229 and K228. Red arrows indicate the distances (Å) between the ligand and residues

535 G37 and L198.

536



537

538

539 **Figure 9.** Docking pose of ilicicolin H bound to the Qi site of cytochrome *b*, showing the H-bond

540 interaction with Q22. Red arrows indicate the distances (Å) between the ligand and residues G37 and

541 L198.

542

543 **Table 1.** Yeast mutants used in the current study containing amino acid substitutions in cytochrome *b*
 544 and relevance of mutations

545

Mutation	Relevance
Y16A	Y16 was hypothesized to participate in UK-2A binding ²³
I17F	Resistance to diuron and HQNO in <i>S. cerevisiae</i> ²⁴
Q22E/M	Q22E confers resistance to ilicicolin H in <i>S. cerevisiae</i> ^{25,26}
N31K	N31K confers resistance in <i>S. cerevisiae</i> to UK-2A, ⁷ antimycin A, diuron, and HQNO ^{24,27}
N31G/S/D	Exploration of substitution at N31
G33A	Resistance to funiculosin and HQNO in <i>S. cerevisiae</i> ²⁸
S34A/L	S34L confers resistance to ametocradin in <i>Plasmopara viticola</i> ¹⁵
G37A	Resistance to ML238, BRD6923 in <i>Plasmodium falciparum</i> ²⁹
G37S	Resistance to ilicicolin H in <i>S. cerevisiae</i> ²⁵
G37V	Resistance to UK-2A in <i>Z. tritici</i> , ⁸ antimycin A in <i>S. cerevisiae</i> , ²⁷ ML238, BRD6923 and antimycin A in <i>Plasmodium falciparum</i> ²⁹
G37C	Resistance to UK-2A in <i>S. cerevisiae</i> ⁷ and ilicicolin H in <i>Candida albicans</i> ³⁰
G37I/M	Exploration of substitution at G37
G37L	Exploration of substitution at G37 and natural variant in <i>Fusarium</i> spp.
G37L+L198M	Natural variant in <i>Fusarium</i> spp.
L38F	Proximity to G37 and natural variant in <i>Z. tritici</i>
L198F	Resistance to UK-2A, antimycin A, ⁷ funiculosin ³¹ and ilicicolin H ²⁵ in <i>S. cerevisiae</i>
L198G/A/V/C/M	Exploration of substitution at L198
L201S	Resistance to cyazofamid in <i>Plasmopara viticola</i> ¹³
H204Y, S206T/V	Resistance to HQNO in <i>S. cerevisiae</i> ³²
M221F	F221 is a natural variant in <i>Z. tritici</i> and was hypothesized to participate in UK-2A binding ³³
M221Q	Resistance to HQNO, ²⁸ ELQ 271 ³⁴ and ilicicolin H ²⁶ in <i>S. cerevisiae</i>
F225L	Resistance to diuron in <i>S. cerevisiae</i> ²⁷
K228M	Resistance to antimycin A in <i>S. cerevisiae</i> ²⁷

546

547 **Table 2.** Amino acids at positions 16, 31, 37, 198 and 221 in various plant pathogens based on yeast
 548 numbering

Organism	NCBI Accession #	Amino Acid				
		16	31	37	198	221
<i>Saccharomyces cerevisiae</i>	YP_009144709	Y	N	G	L	M
<i>Zyoseptoria tritici</i> ^a	AAP81933	Y	N	G	L	F
<i>Pseudocercospora fijiensis</i> ^a	YP_009681541	Y	N	G	L	F
<i>Erysiphe necator</i> ^a	QQY98156	Y	N	G	L	M
<i>Cercospora beticola</i> ^a	AFQ23210	Y	N	G	L	F
<i>Puccinia triticina</i> ^a	YP_009680272	Y	N	G	L	M
<i>Phaeosphaeria nodorum</i> ^a	YP_001427390	Y	N	G	L	F
<i>Phakopsora pachyrhizi</i> ^a	ACT15476	Y	N	G	L	I
<i>Botrytis cinerea</i> ^a	ACL50595	Y	N	A	L	F
<i>Magnaporthe grisea</i> ^a	AAO91631	Y	N	A	L	F
<i>Blumeria graminis f. sp. tritici</i> ^a	QQY98353	Y	N	A	L	M
<i>Alternaria alternata</i>	QGA30432	Y	N	A	L	F
<i>Venturia inaequalis</i> ^a	AAB95255	Y	N	A	L	F
<i>Colletotrichum acutatum</i>	YP_009145212	Y	N	A	L	F
<i>Colletotrichum gloeosporioides</i>	AQT39566	Y	N	A	L	F
<i>Colletotrichum graminicola</i>	EFQ24843	Y	N	A	L	F
<i>Monilinia laxa</i>	ADF43785	Y	N	A	L	F
<i>Monilinia fructigena</i>	ADK35189	Y	N	A	L	F
<i>Monilinia fructicola</i>	ACT75643	Y	N	A	L	F
<i>Rhynchosporium commune</i> ^a	YP_008965385	Y	N	A	L	F
<i>Fusarium solani</i>	YP_005088108	Y	N	L	M	M
<i>Fusarium graminearum</i>	DAC73963	Y	N	L	M	M
<i>Fusarium oxysporum</i>	QJQ35616	Y	N	L	M	M
<i>Rhizoctonia solani</i>	YP_008082056	Y	N	G	F	F

549 ^aHighly sensitive to fenpicoxamid and/or florypicoxamid.⁶

550 **Table 3.** Effects of amino acid change at N31, G37 and L198 on doubling time

Change	Doubling time \pm SD (h) ^a
None	4.33 \pm 0.67 (n = 88)
N31G	6.21 \pm 0.75 (n = 16)**
N31S	7.14 \pm 1.27 (n = 16)**
N31D	5.84 \pm 1.04 (n = 16)**
N31K	7.10 \pm 0.66 (n = 16)**
G37A	4.22 \pm 0.38 (n = 16)
G37S	4.14 \pm 0.39 (n = 16)
G37V	7.36 \pm 1.49 (n = 16)**
G37C	4.52 \pm 0.60 (n = 16)
G37L	13.44 \pm 3.50 (n = 16)**
G37L+L198M	8.26 \pm 2.78 (n = 16)**
G37I	11.60 \pm 2.47 (n = 16)**
G37M	7.78 \pm 1.37 (n = 16)**
L198G	4.76 \pm 0.69 (n = 16)*
L198A	4.53 \pm 0.38 (n = 16)
L198V	4.25 \pm 0.37 (n = 16)
L198C	3.74 \pm 0.19 (n = 16)**
L198M	3.70 \pm 0.20 (n = 16)**
L198F	3.91 \pm 0.33 (n = 16)*

551 ^aOne and two asterisks indicate a statistically significant difference in doubling time from the parent
552 strain at P < 0.05 and P < 0.001, respectively (t-test).

553