



HAL
open science

A genetic and chemical studies of six Oedipodinae species (Orthoptera: Caelifera: Acrididae) from Algeria.

Zina Sofrane, Simon Dupont, Saleheddine Doumandji, Anne-Geneviève Bagnères

► To cite this version:

Zina Sofrane, Simon Dupont, Saleheddine Doumandji, Anne-Geneviève Bagnères. A genetic and chemical studies of six Oedipodinae species (Orthoptera: Caelifera: Acrididae) from Algeria.. *Revista de la Sociedad Entomológica Argentina*, 2022, 81 (1), pp.29-40. hal-03645967v1

HAL Id: hal-03645967

<https://hal.science/hal-03645967v1>

Submitted on 19 Apr 2022 (v1), last revised 5 May 2022 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A genetic and chemical study of six Oedipodinae species (Orthoptera: Caelifera: Acrididae) from Algeria

Sofrane Z.^{1,2}, Dupont S.², Doumandj S.³ & Bagnères A.-G.^{2,4}

¹ *Department of Animal Biology and Physiology. Faculty of Natural and Life Sciences, Ferhat Abbas University Sétif 1, El-Bez 2 campus, 19000, Sétif, Algeria*

² *Institut de Recherche sur la Biologie de l’Insecte, UMR7261, CNRS-University of Tours, Parc Grandmont, 37200, Tours, France*

³ *Research Laboratory of Plant Protection, Zoology. National Higher School of Agronomy, El- Harrach, 16004, Algiers, Algeria*

⁴ *CEFE, CNRS, University of Montpellier, University Paul Valéry Montpellier 3, EPHE, IRD, 34293, Montpellier, France*

Estudios genéticos y químicos de seis especies de Oedipodinae (Orthoptera: Caelifera: Acrididae) de Argelia

RESUMEN. La subfamilia Oedipodinae es el grupo de Acrididae más rico en especies y numéricamente más abundante en Argelia. Aquí, se utilizaron datos genéticos y químicos para caracterizar la identidad de seis especies: *Oedipoda miniata mauritanica* Lucas, 1849, *Acrotylus patruelis* (Herrich-Shaeffer, 1838), *A. insubricus insubricus* (Scopoli, 1786), *Sphingonotus (Neosphingonotus) azurescens* (Rambur, 1838), *Sphingonotus (Neosphingonotus) finotianus* (Saussure, 1885) y *Sphingoderus carinatus* (Saussure, 1888). Se recolectaron individuos de dos sitios diferentes en la región de Sétif, al noreste de Argelia. Primero, los especímenes se clasificaron en función de sus rasgos morfológicos. En segundo lugar, se realizó un análisis filogenético utilizando datos de códigos de barras de ADN (gen COI). En tercer lugar, se analizaron los hidrocarburos cuticulares de las muestras. El análisis filogenético reveló claras

distinciones entre las especies. El análisis quimiotaconómico destacó las similitudes de perfil entre *S. azurescens* y *S. finotianus* y la diferencia de perfil entre *A. patruelis* y *A. i. insubricus*. El perfil de hidrocarburos cuticulares de *S. carinatus* fue diferente de los de *S. azurescens* y *S. finotianus* por *n*-alcanos y trimetilalcanos con cadenas de carbono mucho más cortas. Los datos genéticos y químicos obtenidos han ayudado a aclarar las diferencias dentro y entre los géneros. También pudieron distinguir claramente entre las seis especies.

PALABRAS CLAVE. Orthoptera, Oedipodinae. Código de barras de ADN. Filogenia. Quimiotaconomía.

ABSTRACT. The Oedipodinae subfamily is the most species-rich and numerically abundant group of Acrididae in Algeria. Here, genetic and chemical data were used to characterize the identities of six species: *Oedipoda miniata mauritanica* (Lucas, 1849); *Acrotylus patruelis* (Herrich-Shaeffer, 1838); *A. insubricus insubricus* (Scopoli, 1786); *Sphingonotus (Neosphingonotus) azurescens* (Rambur, 1838); *Sphingonotus (Neosphingonotus) finotianus* (Saussure, 1885); and *Sphingoderus carinatus* (Saussure, 1888). Individuals were collected from two different sites in the Sétif region of northeastern Algeria. First, specimens were classified based on their morphological traits. Second, phylogenetic analysis was performed using DNA barcoding data (COI gene). Third, the specimens' cuticular hydrocarbons (CHCs) were analyzed. The phylogenetic analysis revealed clear distinctions among the species. The chemotaxonomic analysis highlighted CHC profile similarities between *S. azurescens* and *S. finotianus* and CHC profile differences between *A. patruelis* and *A. i. insubricus*. Compared to *S. azurescens* and *S. finotianus*, *S. carinatus* had *n*-alkanes and trimethylalkanes with much shorter carbon chains. The genetic and chemical data obtained here have helped clarify differences within and between genera. They were also able to clearly distinguish among the six species.

KEYWORDS. Orthoptera. Oedipodinae. DNA barcoding. Phylogeny. Chemotaxonomy.

INTRODUCTION

In living organisms in general, and in acrididians in particular, species have traditionally been identified based mainly on morphological criteria. However, although a species' morphology may be unique, its taxonomic identity comprises multiple facets. Indeed, according to Morin (1980), taxonomic identity is determined by several interdependent factors: morphology, genetics, physiology, and behavior. In 2005, Dayrat indicated that identification based exclusively on morphology (i.e., resulting in morphospecies) could lead to valid classifications but that the resulting species should be treated as hypothetical taxa whose identities required confirmation through different types of approaches and data. He proposed using a broader toolkit, including molecular methods and population genetics analysis, in lieu of relying exclusively on classical taxonomic techniques. Padial et al. (2010) suggested that characters can be classified by type: biochemical, molecular, morphological, behavioral, or ecological. Moreover, they emphasized that, although traditional methods may remain useful in many cases, taxonomy must become more interdisciplinary and adopt new approaches to define species if it wishes to become a modern discipline. More recently, chemotaxonomy has been used to separate out species that do not display clear morphological differences (Clément et al. 2001; Wicker & Bagnères, 2020).

In the Acridoidea superfamily (Orthoptera, Caelifera), a single family predominates: Acrididae. Within Acrididae the subfamily Oedipodinae is largely composed of species with brightly colored hind wings that are only visible during flight. When their wings are closed, individuals are camouflaged within the environment. In several studies carried out in Algeria (notably by Mekkioui & Mesli, 2010; Sofrane & Doumandji, 2016), it was found that Oedipodinae was the best inventoried taxonomic group because of its high levels of abundance. Similar results were obtained in the Maghreb (Usmani, 2008). In addition, it has been shown that many species belong to Oedipodinae based on the list of Acridoidea for northwestern Africa in general and for Algeria in particular; this list was established by Louveaux & Ben Halima (1987) and Louveaux et al. (2020). The current taxonomic approach for this group relies generally on morphological traits; however, it lacks the accuracy to unequivocally identify species. There is thus a need to develop complementary approaches in which additional, non-morphological traits are employed in species identification for this group.

Intensive research on Oedipodinae phylogeny and biogeography was carried out by Hochkirch & Husemann (2008), who examined representatives of the tribe Sphingonotini collected from the Canary Islands. Based on a molecular phylogeny, the researchers proposed synonymizing the genera *Wernerella*, *Pseudosphingonotus*, and *Neosphingonotus* with the genus *Sphingonotus*. Additional research in this vein was performed by Husemann et al. (2012) using two tribes of Oedipodinae (Bryodemini and Sphingonotini). Their findings illustrated that 1) external morphological characters are often unreliable when drawing conclusions about taxonomic relationships and 2) genetic data, such as DNA sequences, are more useful when defining higher-level taxonomic groups. Using the mitochondrial cytochrome oxidase I (COI) gene, Moussi et al. (2018) found that DNA barcoding is a valuable tool for identifying and delimiting species in many Oedipodinae genera. At present, COI-based barcoding is the most widespread method.

In addition, chemotaxonomy has been used by a number of researchers to study invertebrates, especially insects (e.g., Formicidae: Lucas et al., 2002; Blattidae and Termitidae: Said et al., 2005; Rhinotermitidae: Clément et al., 2001; and Culicidae: Amira et al., 2013). However, the chemical profiles of Acrididae in general and of Oedipodinae in particular, remain poorly described when compared to what has been done in other insect groups (e.g., Lockey & Oraha, 1990 and Grunshaw et al., 1990). In the above research, one of the most examined chemotaxonomic characters is the cuticular hydrocarbons (CHC) profile (see the review by Bagnères & Wicker-Thomas, 2010). The cuticle has two main layers: a thin, non-chitinous outer epicuticle and a thicker, chitinous inner procuticle. The epicuticle is waxy and composed of lipids comprising various hydrocarbons; the latter can be saturated (alkanes with saturated linear chains [n-alkanes]), methylated (methylalkanes), or unsaturated with double bonds (e.g., monoenes, dienes) (Lockey, 1988). According to Allen (1998), because epicuticular lipids reduce cuticular transpiration, they play a vital role in enabling arthropods to thrive in terrestrial environments. According to Bagnères & Wicker-Thomas (2010), many species and their populations can be identified based on their CHC profiles. Furthermore, several studies have shown that CHC profiles can also be used to differentiate between the sexes within species (e.g., mosquitoes: Amira et al., 2013).

With the aim to building a broader kit of taxonomic tools for identifying members of Oedipodinae, this study focused on six grasshopper species belonging to four Oedipodinae genera to assess differences within and among genera. Our specific research objectives were as follows: 1) to characterize species identity and relatedness using DNA barcoding and 2) to explore similarity in CHCs profiles within and among species.

MATERIAL AND METHODS

Sample collection

We collected 111 specimens from two sites in the Sétif region of northeastern Algeria. The first, El-Ourecia, is a wasteland (36° 14'43.5" N; 05° 24' 05.9" E; elevation 1040 m). Soil salinity is very low; the electrical conductivity (EC) level (1/5) is 0.182 deciSiemens per meter (dS/m). The second, Bazer Sakhra, is a sebkha (36° 04'18" N, 05° 39'56" E; elevation 915 m). Soil salinity is high (EC [1/15] = 18.43 dS/m). All specimens were frozen at -20 °C until their use in the analyses. Taxonomic identification was carried out using taxonomic keys that employ external morphology (Chopard (1943), Mestre (1988), Defaut (2006), Belmann & Luquet (2009), Fontana et al. (2019), and Louveaux et al. (2020)). Six species were identified: *Oedipoda miniata mauritanica* (Lucas 1849); *Acrotylus patruelis* (Herrich-Shaeffer 1838); *Acrotylus insubricus insubricus* (Scopoli 1786); *Sphingonotus (Neosphingonotus) azurescens* (Rambur 1838) (see specific discussion later); *Sphingonotus (Neosphingonotus) finotianus* (Saussure 1885); and *S. carinatus*. Fifty-one individuals were used in the DNA barcoding analysis, where each species was represented by at least seven specimens. The only exception was *A. patruelis*, for which we only had a single specimen because it is rare in the study area (see Appendix I for details). All the grasshoppers (n = 111) were used in the chemotaxonomic analyses.

DNA barcoding

Genomic DNA was extracted using the Wizard® Genomic DNA Purification Kit (Promega, Madison, WI, USA) in accordance with the manufacturer's instructions. The antennae were used or, in their absence, the legs. Prior to the extraction, the samples

were stored in 96% alcohol at 4 °C. The barcoding region of the COI gene (650 bp) was amplified using the modified primers described in Moulton et al. (2010): MDG-F (5'TYTCAACWAAAYCAYAARGAYATYGG-3') and MDG-R (5'TADACTTCWGGRTGWCCRAARAATCA-3') for *Oedipoda miniata mauritanica*, *Sphingonotus (Neosphingonotus) azurescens*, *Sphingonotus (Neosphingonotus) finotianus*, and *S. carinatus*. For *Acrotylus patruelis* and *A. insubricus insubricus*, COI-F and COI-R were used instead (Husmann et al., 2012). The PCR reaction mixture had a total volume of 40 µl and contained 2 ng of DNA template, 1 × Qiagen Multiplex PCR Master Mix, and 2 µM of each primer. PCR amplification was performed using an ESCO Swift Maxi[®] Thermocycler. The initial denaturation step at 95 °C (15 min) was followed by 35 iterations of the following cycle: 94 °C (30 s), 52 °C (1 min 30 sec), and 72 °C (2 min); there was a final extension step at 72 °C (10 min). The PCR products were sequenced by the GenoScreen Institute (<http://www.genoscreen.fr>) using BigDye 3.1 and a 96-Capillary 3730 xl DNA Analyzer (Applied Biosystems, Foster City, CA, USA). Sequences were inspected, corrected, and aligned using Geneious[®] Pro v. 5.6.6 (<http://www.geneious.com/>). They were then deposited in GenBank.

Phylogenetic analysis

The 51 aligned COI sequences were analyzed using neighbor-joining (NJ), maximum-likelihood (ML), and Bayesian inference (BI) methods. The NJ method was applied using PHYLO_WIN (Galtier et al., 1996). The ML method was applied using PhyML (Guindon & Gascuel, 2003). MrAIC was used to find the appropriate sequence evolution model for the data (GTR + G) (Nylander, 2004). BI was carried out using MrBayes v. 3.1.2 (Huelsenbeck & Ronquist, 2001). The program was run for 15 million generations. No prior assumptions were made about the topology. All the sequences were analyzed using the COI reference sequences available for Oedipodinae in GenBank and BOLD (<https://www.boldsystems.org/>). These reference sequences are indicated in the phylogenetic tree (GenBank or BOLD accession code followed by species name). Two sequences were used for the outgroup: one from *Chortophaga viridifasciata* (JQ513034) and one from *Lilaea fuliginosa* (KM243529).

Analysis of cuticular hydrocarbons

CHCs were extracted from all the specimens using pentane (Blomquist et al., 1984); solvent quantity varied between 2 and 5 ml because it depended on specimen size. Immersion time was 10 min, and, during this period, the mixture was agitated several times. The resulting extract was evaporated under nitrogen flow. Between 40–80 μl of 10^{-5} g/ml of *n*-eicosane was added as an internal standard to calibrate the gas chromatograph (GC) profiles; the exact quantity again depended on specimen size. Then, 2 μl of the extract was injected into a GC (Agilent Technologies 6850, Santa Clara, CA, USA) equipped with a flame ionization detector (FID) and an apolar HP-1 capillary column (Agilent, 100% methylsiloxane; 30 m x 320 μm x 0.25 μm). The flame was supplied with air (450 ml/min) and hydrogen (40 ml/min). The carrier gas was helium at a pressure of 1.2 ml/min. The temperature of the detector as well as that of the injector was 250 °C. The oven's initial temperature was 70 °C. Temperature first climbed by 30 °C/min until it reached 150 °C; it then climbed by 5 °C/min to reach a final temperature of 320 °C. The injection occurred in splitless mode (30 sec), and the GC analysis cycle lasted 62 min. The CHCs were identified using a GC-mass spectrometer (MS) (Autosystem XL coupled with a TurboMass 5.4; Perkin Elmer, Waltham, MA, USA); an electron ionization setting of 70 eV was employed. The parameters of the GC cycle were the same as those described above. The mass spectra were identified using known mass spectra from the literature (Lockey 1988; review in Blomquist & Bagnères, 2010).

The relative percentages of the CHCs were calculated by performing GC analysis on the individual specimens; the cuticular mixtures were identified by performing GC-MS analysis on pools of specimens representing the same species. Principal component analysis (PCA) and hierarchical cluster analysis were carried out using the relative percentages of CHCs. An analysis of variance (ANOVA) was performed to test for intraspecific differences between the sexes (SPSS v. 18).

RESULTS

Phylogenetic results

The three phylogenetic methods yielded trees with congruent topologies that contained the same three major clades. Consequently, we have chosen to present only the BI tree (Fig. 1). The first clade was formed by all the specimens of *O. miniata mauritanica*. The second clade contained two well-supported monophyletic groups that corresponded to the two *Acrotylus* species. The third clade comprised the vast majority of the specimens representing the two *Sphingonotus* species as well as the specimens for *S. carinatus*. In the first clade (i.e., corresponding to *O. m. mauritanica*), four haplotypes were present that corresponded to four variable nucleotide sites. In the second clade, there were two groups. The first group contained a single haplotype (AA10) that was associated with *A. partuelis*; the second group contained six haplotypes (corresponding to eight variable nucleotide sites) that were associated with *A. insubricus insubricus*. In the third clade, there were three well-supported monophyletic groups that corresponded to the three species. The first group was composed of eight specimens divided into two haplotypes (corresponding to a single variable nucleotide site); this group was associated with the BOLD reference sequence for *Sphingonotus lucasii* (OBAL053-18). The second group was composed of nine specimens displaying a single haplotype that were associated with *S. (Neosphingonotus) finotianus*. The third group consisted of eight specimens divided into two haplotypes (corresponding to a single variable nucleotide site); this group was associated with *S. carinatus*.

Analysis of cuticular hydrocarbons

Using the GC and GC-MS analyses, CHC profiles were obtained for the six study species (Fig. 2); they contained four CHC classes: *n*-alkanes, monomethylalkanes, dimethylalkanes, and trimethylalkanes. No unsaturated CHCs were seen (Table I). Within species, individuals displayed certain similarities in compound classes, although relative compound quantities differed (Fig. 3).

In *Oedipoda miniata mauritanica* and *S. azurescens*, *n*-alkanes were more common in females; in *S. finotianus*, *S. carinatus*, and *Acrotylus insubricus insubricus*, they were more common in males. Monomethylalkanes were also generally more common in females. However, in *S. azurescens* and *S. carinatus*, they were more common in males. There were only minor differences in dimethylalkanes between the sexes. That said,

dimethylalkanes were far more prominent in females in *A. i. insubricus* and *S. carinatus*. Across all six species, trimethylalkanes were generally more common in males. The only exceptions were seen in *S. carinatus* (more common in females) and in *A. i. insubricus* (no sex-specific difference) (Fig. 4).

In general, in *Oedipoda miniata mauritanica*, *Acrotylus patruelis* and *A. insubricus insubricus*, the most prominent CHC classes were the monomethylalkanes and *n*-alkanes. The latter were most prominent in *S. azurescens*, *S. finotianus*, and *S. carinatus*.

Comparing relative CHC levels among species, *n*-alkanes were more common in *S. azurescens*; monomethylalkanes were more common in *Acrotylus patruelis*; and dimethylalkanes were more common in *S. carinatus*; and trimethylalkanes were more common in *S. finotianus*.

A PCA was carried out using 64 variables, which corresponded to the CHC profiles of all the specimens (i.e., the GC-MS data (Table I), and the identities of the six study species. Taken together, axis 1 and axis 2 explained 53.18% of the total variance (33.04% and 20.14%, respectively; Fig. 5). Three clusters formed largely along taxonomic lines: one with individuals from *Oedipoda miniata mauritanica*; another with individuals from both *Acrotylus insubricus insubricus* and *A. patruelis*; and a third with individuals from *S. azurescens*, *S. finotianus*, and *S. carinatus*. Furthermore, *O. m. mauritanica* had a very distinct profile, characterized by monomethylalkanes, dimethylalkanes, and trimethylalkanes with much shorter carbon chains. The clustering of *S. azurescens*, *S. finotianus*, and *S. carinatus* can be explained by the similarities in their CHCs classes, namely the mono- and dimethylalkanes. (Table II).

Sex-specific differences in CHCs profiles were seen in *Oedipoda mauritanica* and *S. carinatus* (ANOVA: $p < 0.05$) but not in *S. azurescens*, *S. finotianus*, *Acrotylus insubricus insubricus*, or *A. patruelis*.

DISCUSSION

There is a need to expand the taxonomic toolkit for identifying species in Oedipodinae. To this end, we explored the ability of a combined approach—using both genetic and chemical data—to differentiate among six study species.

Phylogenetic results

The COI-based phylogenetic analysis generally confirmed our morphology-based classifications. Indeed, the sole member of the second clade in the phylogenetic tree was identified as *Acrotylus patruelis*. Thus, the difference between *A. patruelis* and *A. insubricus* was clear from the barcoding data.

Individuals that were morphologically identified as belonging to *S. azurescens*, *S. finotianus*, and *S. carinatus* occurred in three well-supported monophyletic groups in the third clade. Consequently, questions are raised about the *S. azurescens* (JQ513072) and *S. lucasii* (OBAL053-18) reference sequences—are those species assignments actually correct? Our specimens that grouped genetically with the *S. lucasii* reference sequence were morphologically classified as *S. azurescens*. The latter species is identified based on the following traits. It has a pronotum that is nearly smooth, weakly rough, and finely punctuated. The pronotum's median carina is prominent in the prozone but reduced to a single line in the metazone. The species' posterior femurs have a dark inner side with two clear bands. The upper carina of the posterior femur descends in its apical half. The posterior tibiae are yellowish white with a bluish tinge. The tegmina have a broad brown spot in their basal third and a narrow brown stripe in their middle. The wings have a broad brown fascia that varies in length; it starts at the middle of the anterior wing edge and closely follows the edge. Given that our specimens clearly displayed these traits, we assumed that they were indeed *S. azurescens*.

According to Chopard (1943), *S. azurescens* has been confused with neighboring species, notably with *S. lucasii*. Louveaux et al. (2020) reiterated that confusion between these species may occur and that their distribution ranges need to be confirmed. The species appear to occur in different regions of North Africa (e.g., northern and southern Algeria). They are geographically close.

Moussi et al. (2018) stated that there was a match between the molecular and morphological classifications of many Oedipodinae species; however, *Sphingonotus*

remained a problematic taxon in their phylogenetic analysis. These authors interpreted this result as showing that COI-based species delimitation remains difficult, at least for most Sphingonotini genera, perhaps because of nuclear copies of mitochondrial sequences (pseudogenes), even though DNA barcoding is a valuable tool for identifying species in these genera. They suggested additional genetic data may be required to achieve better levels of resolution. In conclusion, we assume here definitely that our eight specimens as *S. azurescens* rather than as *S. lucasii*.

Analysis of cuticular hydrocarbons

The length of the CHC carbon chains observed in this study are consistent with those seen in previous research on insects. According to Nelson & Blomquist (1995), this length ranges approximately from 21 to 50 carbons, and the shorter chain compounds are volatile. In general, the number of methylated compounds was high, and there was a remarkable absence of unsaturated products (alkenes). These results agree with those obtained for *Locusta migratoria cinerascens* (Genin et al., 1986).

The cluster analysis revealed species specific variation in CHC profiles. Some past studies (Nielsen et al., 1999; Buczkowski et al., 2005; Parkarsh et al., 2008; Bontonou et al., 2013) have found that variation in CHCs can be linked to factors such as age, diet, habitat, and diverse environmental characteristics, while other studies have seen no such correlation (Toolson, 1982; Vander Meer et al., 1989; Dahbi et al., 1996).

Consequently, CHCs could contribute to creating a species-specific chemical signature. The PCA results highlighted the profile similarities among *S. azurescens*, *S. finotianus*, and *S. carinatus*. The three species formed a single cluster that was distinct from the cluster formed by *Oedipoda miniata mauritanica* and from the cluster formed by *Acrotylus insubricus insubricus* and *A. patruelis*. These results show that, although there were similarities, the CHC profiles were nonetheless relatively species specific, notably in their number of compounds. The clusters arising from the chemical data generally corresponded to the clusters arising from the genetic data.

There was a significant sex-specific difference in relative CHC percentages in *S. carinatus* and *Oedipoda miniata mauritanica* but not in *S. azurescens*, *S. finotianus*, *Acrotylus insubricus insubricus*, or *A. patruelis*. However, it could be that more volatile compounds not analyzed here are also at play. Sexual selection could be involved

(Thomas & Simmons, 2008), given that sexual dimorphism in CHC profiles was seen in only some species and that profile variation was seen within genera. An explanation could be that hydrocarbons are involved in reproduction, acting as sex pheromones as they do in other insect taxa, such as mosquitoes (Amira et al., 2013), *Drosophila* (Bontonou et al, 2013), *Blattella germanica* (Morakchi et al., 2010), and cricket *Teleogryllus oceanicus* (Thomas & Simmons, 2008).

In this study, we successfully explored intraspecific diversity within *Acrotylus insubricus* and used our combined taxonomic approach to distinguish between the two *Acrotylus* species. For *Sphingonotus finotianus* and *S. carinatus*, we increased the taxonomic specificity and showed that the species display clear genetic and chemical differences. We also helped flesh out the genetic and chemical taxonomy of the two other study species.

The genetic and chemical data we obtained have helped clarify differences within and between genera. They were also able to clearly distinguish between the six study species. Our findings indicate that such multifaceted analyses could make a significant contribution by complementing current morphotaxonomic techniques and thus help clarify species systematics.

ACKNOWLEDGMENTS

This study was funded by grants from the French and Algerian Ministries of Foreign Affairs (PROFAS Program) obtained during ZS's PhD and by CNRS grants to AGBU, who hosted ZS in her lab. We would like to thank JP Christidès for his technical assistance. The authors also thank Jessica Pearce-Duvet for her English editing services.

LITERATURE CITED

- Allen, G. (1998) Water-proofing properties of cuticular lipids. *Integrative and Comparative Biology*, **38**(3), 471-482.
- Amira, K., Boudjelida, H., Farine, J.P., Djeghader. N.E.H. (2013). Caractérisation des hydrocarbures cuticulaires du moustique *Culex pipiens*. *Microbiologie et Hygiène Alimentaire*, **25**, 112-115.

- Bagnères, A.-G. & Wicker-Thomas, C. (2010) Chemical taxonomy with hydrocarbons. *Insect hydrocarbons: biology, biochemistry and chemical ecology*. Cambridge (UK) (ed. Blomquist, G.J., Bagnères, A.-G.), pp. 121-162. Cambridge University Press.
- Belmann, H. & Luquet, G. (2009) *Guide des sauterelles, grillons et criquets d'Europe Occidentale*. 2nded. Delachaux et Niestlé, Paris.
- Blomquist, G.J., Adams, T.S. & Dillwith, J.W. (1984) Induction of female sex pheromone production in male houseflies by ovary implants or 20-hydroxyecdysone. *Journal of Insect Physiology*, **30**, 295-302.
- Blomquist, G.J. & Bagnères, A.-G. (2010) *Insect Hydrocarbons: Biology, Biochemistry and Chemical Ecology*. 1st ed. Cambridge (UK): Cambridge University Press.
- Bontonou, G., Denis, B. & Wicker-Thomas, C. (2013) Interaction between temperature and male pheromone in sexual isolation in *Drosophila melanogaster*. *Journal of Evolutionary Biology*, **26**, 2008-2020.
- Buczowski, G., Kumar, R., Suib, S.L. & Silverman, J. (2005) Diet-related modification of cuticular hydrocarbon profiles of the Argentine ant, *Linepithema humile*, diminishes intercolony aggression. *Journal of Chemical Ecology*, **31**, 829-843.
- Chopard, L. (1943) *Faune de l'empire français, Orthoptéroïdes de l'Afrique du nord*. I. 1st ed., Librairie Larose, Paris, France.
- Clément, J.L., Bagnères, A.-G., Uva, P., Wilfert, L., Quintana, A., Reinhard, J. & Dronnet, S. (2001) Biosystematics of *Reticulitermes termites* in Europe. Morphological, chemical, molecular data. *Insectes Sociaux*, **48**, 202-215.
- Dahbi, A., Cerdá, X., Hefetz, A. & Lenoir, A. (1996) Social closure, aggressive behaviour, and cuticular hydrocarbon profiles in the polydomous ant *Cataglyphis iberica* (Hymenoptera, Formicidae). *Journal of Chemical Ecology*, **22**, 2173-2186.
- Dayrat, B. (2005) Towards integrative taxonomy. *Biological Journal of the Linnean Society*, **85**, 407-415.
- Defaut, B. (2006) Révision préliminaire des *Oedipoda* ouest-paléarctiques (Caelifera, Acrididae, Oedipodinae). *Matériaux Orthoptériques et Entomocénétiques*, **11**, 23-48.
- Fontana, P., Massa, B. & Buzzetti, F.M. (2019) A new rare species of *Oedipoda* Latreille, 1829 (Orthoptera: Acrididae) from South Italy. *Zootaxa*, **4614**(1), 50-60.
- Galtier, N., Gouy, M. & Gautier, C. (1996) Seaview and Phylo Win: two graphic tools for sequence alignment and molecular phylogeny. *Computer Applications in the Biosciences*, **12**, 543-548.
- Genin, E., Jullien, R., Pere, F. & Fuzeau-Braesch, S. (1986) Cuticular hydrocarbons of gregarious and solitary locusts (*Locusta migratoria cinerascens*). *Journal of Chemical Ecology*, **12**, 1213-1238.
- Grunshaw, J.P., Guermouche, H., Guermouche, S., Jago, N.D., Jullien, R., Knowles, E. & Perez, F. (1990) Chemical taxonomic studies of cuticular hydrocarbons in locusts of the *Schistocerca americana complex* (Acrididae: Cyrtacanthacridinae) chemical relationships between New world and Old world species. *Journal of Chemical Ecology*, **16**(10), 2835-2858.

- Guindon, S. & Gascuel, O. (2003) A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Systematic Biology*, **52**, 696-704.
- Hebert, P.D. N., Cywinska, A., Ball, S. & Landdewaard, J.R. (2003) Biological identifications through DNA barcodes. *Proceedings of the Royal Society of London, Series B* **270**, 313-322.
- Hochkirch, A. & Husemann, M. (2008) A Review of the Canarian Spthingonotini with description of a new species from Fuerteventura (Orthoptera: Acridae: Oedipodinae). *Zoological Studies*, **47(4)**, 495-506.
- Huelsenbeck, J.P. & Ronquist, F. (2001) MrBAYES Bayesian inference of phylogenetic trees. *Bioinformatics*, **17**, 754-755.
- Husemann, M., Namkung, S., Habel, J.C., Danley, P.D. & Hochkirch, A. (2012) Phylogenetic analyses of band-winged grasshoppers (Orthoptera, Acrididae, Oedipodinae) reveal convergence of wing morphology. *Zoologica Scripta*, **41**, 515-526.
- Lockey, K.H. (1988) Lipids of the insect cuticle: origin, composition and function. *Comparative Biochemistry and Physiology* **89**, 595-645.
- Lockey, K.H. & Oraha, V.S. (1990) Cuticular lipids of adult *Locusta migratory migratorioides* (R and F), *Schistocerca gregaria* (Forsk.) (Acrididae) and other orthoptera species. II. Hydrocarbons. *Comparative Biochemistry and Physiology*, **95B**, 721-74.
- Louveaux, A. & Ben Halima, T. (1987) Catalogue des orthoptères Acridoidea d'Afrique du nord-ouest. *Bulletin de la Société Entomologique Fr*, **91(3-4)**, 73-87.
- Louveaux, A., Amedegnato, C., Poulains, S. & Desutter-Grandcolas, L. (2020) *Orthoptères Acridomorpha de l'Afrique du Nord-ouest*. Version [2.1]. <http://acrinwafrica.mnhn.fr/>
- Lucas, C., Fresneau, D., Kolmer, K., Heinze, J., Delabie, J.H.C. & Pho, D. B. (2012) A multidisciplinary approach to discriminating different taxa in the species complex *Pachycondyla villosa* (Formicidae). *Biological Journal of the Linnean Society*, 2002, **75**, 249-259.
- Mekkioui, A. & Mesli, L. (2010) Etude préliminaire des Orthoptères Caelifères de deux stations dans la région de Hafir (Monts de Tlemcen, Ouest Algérien). *Matériaux Orthoptériques et Entomocénétiques*, **14(2009)**, 53-60.
- Mestre, J. (1988) *Les acridiens des formations herbeuses d'Afrique de l'Ouest*. 2 nd ed. Cirad-Prifas, Montpellier, France.
- Morakchi, S., Aribi, N., Farine, J.P. & Soltani, N. (2010) Analyse des hydrocarbures cuticulaires chez *Blattella germanica*. In *Actes de la CIFE VI, Travaux de l'Institut Scientifique, Série Zoologie*, Rabat, 2010, N° 47, Tome I, 141-145.
- Morin, E. (1980) *La méthode, la vie de la vie*. II. 1st ed. Le seuil, Paris, France (in French).
- Moulton, M.J., Song, H. & Whiting, M.F. (2010) Assessing the effects of primer specificity on eliminating numt coamplification in DNA barcoding: a case study from Orthoptera (Arthropoda: Insecta). *Molecular Ecology Resources*, **10**, 615-627.

- Moussi, A., Dey, L.S., Petit, D., Abba, A., Klessner, R. & Husemann, M. (2018) First genetic data for band winged grasshoppers (Orthoptera: Acrididae: Oedipodinae) of the Biskra region of Algeria with new records for the country. *African Zoology*, **53**(1), 31-40.
- Nelson, D.R. & Blomquist, G.J. (1995). Insect waxes. In *Waxes: Chemistry, Molecular Biology and Functions*. Ed. R. J. Hamilton. Dundee, Scotland: *The Oily Press*, pp. 1-90.
- Nielsen, J., Boomsma, J.J., Oldham, N.J., Petersen, H.C. & Morgan, E.D. (1999) Colony-level and season specific variation in cuticular hydrocarbon profiles of individual workers in the ant *Formica truncorum*. *Insectes Sociaux*, **46**, 58-65.
- Nylander, J.A.A. (2004) *Evolutionary Biology. MrAIC.pl*. Program distributed by the author, Centre, Uppsala University.
- Padial, J.M., Miralles, A., De la Riva, I. & Vences, M., (2010) The integrative future of taxonomy. *Frontiers in Zoology*, **7**, 1-14.
- Parkash, R., Kalra, B. & Sharma, V. (2008) Changes in cuticular lipids, water loss and desiccation resistance in a tropical drosophilid; Analysis of variation between and within populations. *Fly*, **2**, 189-197.
- Said, I., Costagliola, G., Leoncini, I. & Rivault C. (2005) Cuticular hydrocarbon profiles and aggregation in four *Periplaneta* species (Insecta: Dictyoptera). *Journal of insect physiology*, **51**(9), 995-1003,
- Schumakov, E.M. (1963): Acridoidea of Afghanistan and Iran. *URSS Academy of Sciences*, 493-248.
- Sofrane, Z. & Doumandji, S. (2016) Contribution à l'étude de la biodiversité des Caelifera (Orthoptera) et des Tettigoniidae (Orthoptera, Ensifera) dans la région de Sétif (Nord- Est algérien). In: *Proceedings of 10th Maghreb Conference on Marine Sciences and 5th France-Maghreb Congress of Zoology and Ichthyology* (2016), Fèz, Morocco. p.87.
- Thomas M.L. & Simmons L.W. (2008) Sexual dimorphism in cuticular hydrocarbons of the Australian field cricket *Teleogryllus oceanicus* (Orthoptera: Gryllidae) *Journal of Insect Physiology*, **54**, 1081-1089.
- Toolson, E.C. (1982) Effects of rearing temperature on cuticle permeability and epicuticular composition in *Drosophila pseudoobscura*. *Journal Experimental Zoology*, **222**, 249-253.
- Usmani, M.K. (2008) Studies on Acridoidea (Orthoptera) with some new records from Fezzan, Libya. *Zootaxa*, **1946**, 1-41.
- Vander Meer, R.K., Saliwanchik, D. & Lavine, B. (1989) Temporal changes in colony cuticular hydrocarbon patterns of *Solenopsis invicta*: implications for nestmate recognition. *Journal of Chemical Ecology*, **15**, 2115-2125.

Table captions

Table I. List of cuticular hydrocarbons identified in *Oedipoda miniata mauritanica* (Omin), *Acrotylus insubricus insubricus* (Ains), *Acrotylus patruelis* (Apat), *Sphingonotus azurescens* (Sazu), *Sphingonotus finotianus* (Sfin), and *Sphingoderus carinatus* (Scar) collected in Sétif. The abbreviations are as follows: t, present in trace amounts; -, absent; and X, unknown.

Table II. Carbon chain length range for the cuticular hydrocarbon classes seen in *Oedipoda miniata mauritanica* (Omin), *Acrotylus insubricus insubricus* (Ains), *Acrotylus patruelis* (Apat), *Sphingonotus azurescens* (Sazu), *S. finotianus* (Sfin), and *Sphingoderus carinatus* (Scar).

Appendix I. Details about the study species specimens: *Oedipoda miniata mauritanica* (Omin), *Acrotylus insubricus insubricus* (Ains), *Acrotylus patruelis* (Apat), *Sphingonotus azurescens* (Sazu), *Sphingoderus carinatus* (Scar), and *Sphingonotus finotianus* (Sfin). Sites: A = El-Ourecia, B = Bazer Sakhra. Abbreviations: morph = morphology-based classification; CHCs = characterization of cuticular hydrocarbon profile; COI = COI gene barcoding; + = used; - = not used.

Figure captions

Fig. 1. Phylogenetic tree built using COI gene sequences and Bayesian inference. The branch support values are Bayesian posterior probabilities. The reference sequences are labeled (Genbank or BOLD accession code followed by species name). The brackets on the right side of the tree indicate the species identity of the specimens.

Fig. 2. Total ion chromatograms (GC-MS) for *Oedipoda miniata mauritanica* (A), *Acrotylus insubricus insubricus* (B), *Acrotylus patruelis* (C), *Sphingonotus azurescens* (D), *Sphingonotus finotianus* (E), and *Sphingoderus carinatus* (F).

Fig. 3. Hierarchical cluster analysis of the CHC profiles of *Oedipoda miniata mauritanica* (A), *Acrotylus insubricus insubricus* (B), *Sphingonotus azurescens* (C), *Sphingonotus finotianus* (D), and *Sphingoderus carinatus* (E).

Fig. 4. Relative percentages of the different CHC classes observed in *Oedipoda miniata mauritanica* (Omin), *Acrotylus insubricus insubricus* (Ains), *Acrotylus patruelis* (Apat), *Sphingonotus azurescens* (Sazu), *Sphingoderus carinatus* (Scar), and *Sphingonotus finotianus* (Sfin).

Fig. 5 Principal component analysis of the CHC profiles of *Oedipoda miniata mauritanica* (Omin), *Acrotylus insubricus insubricus* (Ains), *Acrotylus patruelis* (Apat), *Sphingonotus azurescens* (Sazu), *Sphingoderus carinatus* (Scar), and *Sphingonotus finotianus* (Sfin). The results for males (1) and females (2) are shown.

Compound identity	Peak number per species					
	Omin	Ains	Apat	Sazu	Sfin	Scar
<i>n</i> -C23	1	t	-	1	1	-
X	-	-	-	2	2	-
3-MeC23	2	-	-	-	-	-
<i>n</i> -C24	3	t	-	3	3	t
4/2-MeC24	4	-	-	-	-	-
<i>n</i> -C25	5	t	-	4	4	2
11-MeC25	6	-	-	5	5	t
diMeC25 (+5MeC25)		-	-	6	6	t
4-MeC25	7	-	-	-	-	
3-MeC25	8	-	-	7	7	t
<i>n</i> -C26	9	t	-	8	8	3
12-MeC26+X	T	-	-	t	t	t
4/2-MeC26	10	t	-	10	t	3a
<i>n</i> -C27	11	1	1	11	11	4
13-+11-MeC27	12	t	-	12	12	5
5-MeC27	13	-	-	13	13	5a
diMeC27		t	-	14	14	5b
4/2-MeC27	14	-	-	-	-	-
3-Me27	15	t	-	15	15	5c
<i>n</i> -C28	16	2	2	17	17	6
12-MeC28 +X	t	t	-	18	t	t
<i>n</i> -C29	18	4	4	20	20	10
15-+13-+11-MeC29	19	5	t	21	21	11

Table I. List of cuticular hydrocarbons identified in *Oedipoda miniata mauritanica* (Omin), *Acrotylus insubricus insubricus* (Ains), *Acrotylus patruelis* (Apat), *Sphingonotus azurescens* (Sazu), *Sphingonotus finotianus* (Sfin), and *Sphingoderus carinatus* (Scar) collected in Sétif. The abbreviations are as follows: t, present in trace amounts; -, absent; and X, unknown.

Compound identity	Peak number per species					
	Omin	Ains	Apat	Sazu	Sfin	Scar
7-MeC29	t	t	–	22	22	11a
5-MeC29	t	t	–	23	23	11b
diMeC29	20	t	-	24	24	11c
4/2-MeC29		6	6			
3-MeC29	21	7	t	25	25	12
X				26	26	
<i>n</i> -C30	22	8	8	27	27	13
12-MeC30+X	23	t	-	28	28	t
4/2-MeC30	24	9	9	29	29	14
X	-	10	-	-	-	-
<i>n</i> -C31	25	11	11	30	30	18
4,12,16-triMeC30	-	12	t	-	-	-
15-+13-+11-MeC31 + X	26	13	t	31	31	19
9+7-MeC31	26a	T	t	t	t	-
5-MeC31	26b	T	–	t	32	t
x,y-diMeC31	27	15	t	32	32a	t
4/2-MeC31+X	28	16	16	-	-	-
3-MeC31	29	t	t	33	33	t
w,z-diMeC31	-	-	-	34	34	t
<i>n</i> -C32	30	18	18	35	35	22
X	t	t	19	-	-	-
14-+12-+11-+10-MeC32	-	20	-	36	36	23
14-+12-+11-+10-MeC32	-	20	-	36	36	23
6-MeC32	-	t	t	-	-	-
diMeC32	-	21	t	-	-	-
4/2-MeC32	31	22	22	37	37	t
3-MeC32		t	–		38	t

Table I. (continued)

Compound identity	Peaknumber per species					
	Omin	Ains	Apat	Sazu	Sfin	Scar
X		t	-		38a	
<i>n</i> -C33	32	25	25	39	39	27
triMeC32	-	26	t	-	-	-
9-+7-MeC33		t	t	40a		30a
13,17+13,19-+13,21-diMeC33	-	29	29	-	-	-
11,17+11,19-+11,21-diMeC33	34	-	-	41	41	31
4/2-MeC33	-	t	30	-	-	-
7,15-diMeC33	-	-	-	41b	41a	32
13,17,21-triMeC33+3MeC33	35	-	-	42	42	32a
<i>n</i> -C34	36	31	31	43	43	33
triMeC33	-	t	t	-	-	-
X	36b	-	-	-	-	t
X	37	-	-	-	-	-
X	37b	-	-	-	-	-
X	38	-	-	-	-	-
13-+12-+11-MeC34	-	t	t	44	44	35
diMeC34	-	32	t	45	45	37
8,12,16-triMeC34	-	-	33	-	-	-
3-MeC34	-	-	-	-	t	-
4/2-MeC34	-	33	t			
X	-	34	t	t	-	
X	-	35	t	-	-	-
<i>n</i> -C35	39	36	36	46	46	40a
triMeC34+X	t	37	-	-	-	t
17-+15-+13-+11-MeC35	40	t	t	47	47	42
X	-	39	t	-	-	-
diMeC35	41	40	40	48	48	44

Table I (1). (continued)

Compound identity	Peaknumber per species					
	Omin	Ains	Apat	Sazu	Sfin	Scar
X	-	-	-	49	-	t
triMeC35	42	-	-	50	49+50	45
X		41	t	-	-	t
<i>n</i> -C36	42a	t	t	51a	51	47
14-+12-MeC36+X	t	-	-	52	52	48
dimeC36	-	-	-	53	53	49
trimethylC36	-	-	t	-	-	-
N	-	-	-	-	53a	t
<i>n</i> -C37	45	t	t	54	t	52a
19-+17-+15-+13-+11-MeC37	46	-	-	55	55	54
15,21+17,21-diMeC37	47	-	-	56	56	55
triMeC37	48	-	-	57	57	56
<i>n</i> -C38	t	-	-	58	t	58
14-+12-MeC38	49	-	-	59	t	59
diMeC38		-	-	60	t	60
<i>n</i> -C39	t	-	-	61	t	61a
19-+11-Me C39	50	-	-	62	62	62
diMeC29	-	-	-	63	63	63
triMeC29	-	-	-	64	64	-

Table I (2). (continued)

Species	<i>n</i> -alkanes	monomethylalkanes	dimethylalkanes	trimethylalkanes
Omin	23–39	23–39	29–39	35–39
Ains	23–37	26–35	27–35	30–34
Apat	27–37	28–35	31–35	30–36
Sazu	23–39	25–39	25–39	33–39
Sfin	23–39	25–39	25–39	33–39
Scar	24–39	25–39	25–39	33–37

Table II. Carbon chain length range for the cuticular hydrocarbon classes seen in *Oedipoda miniata mauritanica* (Omin), *Acrotylus insubricus insubricus* (Ains), *Acrotylus patruelis* (Apat), *Sphingonotus azurescens* (Sazu), *S. finotianus* (Sfin), and *Sphingoderus carinatus* (Scar).

Code	Site	Sex	Life stage	Collection date	Morph	CHCs	COI	Accession Number
AOm1	A	♂	Adult	02.11.2012	Omin	+	+	MK251045
BOm2	B	♂	Adult	16.11.2012	Omin	+	+	MK251037
BOm3	B	♂	Adult	16.11.2012	Omin	+	-	
BOm4	B	♂	Adult	09.11.2012	Omin	+	-	
BOm5	B	♀	Adult	13.10.2012	Omin	+	-	
BOm6	B	♂	Adult	01.11.2012	Omin	+	-	
BOm7	B	♂	Adult	29.06.2012	Omin	+	-	
AOm8	A	♂	Adult	14.09.2012	Omin	+	-	
BOm9	B	♀	Adult	01.11.2012	Omin	+	+	MK251043
BOm10	B	♂	Adult	01.11.2012	Omin	+	-	
AOm11	A	♂	Adult	05.10.2012	Omin	+	+	MK251042
AOm12	A	♀	Adult	05.10.2012	Omin	+	+	MK251039
BOm13	B	♀	Adult	28.09.2012	Omin	+	+	MK251044
AOm14	A	♀	Adult	22.06.2012	Omin	+	-	
AOm15	A	♀	Adult	23.11.2012	Omin	+	-	
AOm16	A	♀	Adult	23.11.2012	Omin	+	+	MK251038
AOm17	A	♂	Adult	08.09.2012	Omin	+	+	MK251040
BOm18	B	♀	Adult	09.11.2012	Omin	+	+	MK251041
AOm19	A	♀	Adult	06.07.2012	Omin	+	-	
AOm20	A	♀	Adult	23.11.2012	Omin	+	-	
AOm21	A	♂	Adult	07.08.2012	Omin	+	-	

Appendix I. Details about the study species specimens: *Oedipoda miniata mauritanica* (Omin), *Acrotylus insubricus insubricus* (Ains), *Acrotylus patruelis* (Apat), *Sphingonotus azurescens* (Sazu), *Sphingoderus carinatus* (Scar), and *Sphingonotus finotianus* (Sfin). Sites: A = El-Ourecia, B = Bazer Sakhra. Abbreviations: morph = morphology-based classification; CHCs = characterization of cuticular hydrocarbon profile; COI = COI gene barcoding; + = used; - = not used.

Code	Site	Sex	Life stage	Collection date	Morph	CHCs	COI	Accession Number
AA1	A	♀	Adult	19.03.2012	Ains	+	+	MK251036
AA2	A	♂	Adult	02.11.2012	Ains	+	-	
AA3	A	♂	Adult	23.11.2012	Ains	+	-	
AA4	A	♂	Adult	16.11.2012	Ains	+	+	MK251021
BA5	B	♂	Adult	23.11.2012	Ains	+	+	MK251023
AA6	A	♀	Adult	19.10.2012	Ains	+	+	MK251022
AA7	A	♀	Adult	09.11.2012	Ains	+	-	
BA8	B	♀	Adult	09.11.2012	Ains	+	-	
BA9	B	♂	Adult	18.05.2012	Ains	+	-	
AA10	A	♀	Adult	16.11.2012	Apat	+	+	MK251020
BA11	B	♀	Adult	23.11.2012	Ains	+	+	MK251034
BA12	B	♂	Adult	09.11.2012	Ains	+	-	
BA13	B	♀	Adult	01.11.2012	Ains	+	+	MK251025
AA14	A	♂	Adult	02.11.2012	Ains	+	+	MK251026
BA15	B	♀	Adult	01.11.2012	Ains	+	+	MK251028
AA16	A	♀	Adult	19.03.2012	Ains	+	+	MK251027
BA17	B	♀	Adult	01.11.2012	Ains	+	+	MK251029
BA18	B	♂	Adult	09.11.2012	Ains	+	+	MK251030
BA19	B	♀	Adult	09.11.2012	Ains	+	+	MK251035
AA20	A	♀	Adult	09.11.2012	Ains	+	-	
BA21	B	♀	Adult	28.09.2012	Ains	+	-	
AA22	A	♂	Adult	16.11.2012	Ains	+	+	MK251031
AA23	A	♂	Adult	16.11.2012	Ains	+	+	MK251032
AA24	A	♀	Adult	16.11.2012	Ains	+	-	
AA25	A	♀	Adult	02.11.2012	Ains	+	+	MK251033
AA26	A	♂	Adult	19.10.2012	Ains	+	+	MK251024
Bsm1	B	♀	Adult	31.08.2012	Sazu	+	+	MK251009

Appendix I (continued)

Code	Site	Sex	Life stage	Collection date	Morph	CHCs	COI	Accession Number
ASm2	A	♂	Adult	26.07.2012	Sazu	+	-	
ASm3	A	♀	Adult	19.10.2012	Sazu	+	-	
ASm4	A	♀	Adult	05.10.2012	Sazu	+	+	MK251003
ASm5	A	♀	Adult	17.08.2012	Sazu	+	-	
Bsm6	B	♀	Adult	21.09.2012	Sazu	+	+	MK251006
Bsm7	B	♀	Adult	09.11.2012	Sazu	+	+	MK251007
ASm8	A	♂	Adult	14.09.2012	Sazu	+	+	MK251004
ASm9	A	♀	Adult	07.08.2012	Sazu	+	+	MK251005
Bsm10	B	♂	Adult	13.10.2012	Sazu	+	+	MK251010
Bsm11	B	♀	Adult	31.08.2012	Sazu	+	-	
Bsm12	B	♂	Adult	21.09.2012	Sazu	+	-	
ASm13	B	♀	Adult	07.08.2012	Sazu	+	+	MK251008
Bsm14	B	♀	Adult	21.09.2012	Sazu	+	-	
ASm15	B	♀	Adult	14.09.2012	Sazu	+	-	
Bsm16	B	♀	Adult	21.09.2012	Sazu	+	-	
ASm17	A	♀	Adult	05.10.2012	Sazu	+	-	
ASm18	A	♀	Adult	17.08.2012	Sazu	+	-	
ASf1	A	♂	Adult	19.10.2012	Sfin	+	+	MK251013
ASf2	A	♀	Adult	09.11.2012	Sfin	+	-	
ASf3	A	♀	Adult	14.09.2012	Sfin	+	-	
ASf4	A	♂	Adult	23.11.2012	Sfin	+	-	
ASf5	A	♀	Adult	09.11.2012	Sfin	+	-	
ASf6	A	♀	Adult	26.07.2012	Sfin	+	+	MK251012
ASf8	A	♀	Adult	09.11.2012	Sfin	+	+	MK251018
ASf9	A	♀	Adult	26.07.2012	Sfin	+	+	MK251016
ASf10	A	♀	Adult	08.09.2012	Sfin	+	+	MK251011
ASf11	A	♀	Adult	14.09.2012	Sfin	+	-	

Appendix I (1) (continued)

Code	Site	Sex	Life stage	Collection date	Morph	CHCs	COI	Accession Number
ASf12	A	♀	Adult	14.09.2012	Sfin	+	-	
ASf13	A	♀	Adult	19.10.2012	Sfin	+	-	
ASf14	A	♂	Adult	08.09.2012	Sfin	+	+	MK251017
ASf15	A	♀	Adult	05.10.2012	Sfin	+	-	
ASf16	A	♂	Adult	14.09.2012	Sfin	+	-	
ASf18	A	♂	Adult	02.11.2012	Sfin	+	-	
ASf19	A	♂	Adult	16.11.2012	Sfin	+	-	
ASf20	A	♂	Adult	09.11.2012	Sfin	+	-	
ASc9	A	♀	Adult	02.11.2012	Sfin	+	+	MK251014
ASc13	A	♂	Adult	23.11.2012	Sfin	+	+	MK251019
ASc14	A	♀	Adult	26.07.2012	Sfin	+	+	MK251015
BSf7	B	♂	Adult	31.08.2012	Scar	+	+	MK251000
BSf17	B	♂	Adult	31.08.2012	Scar	+	+	MK250999
BSc1	B	♀	Adult	31.08.2012	Scar	+	+	MK250995
ASc2	A	♀	Adult	07.08.2012	Scar	+	-	
BSc3	B	♀	Adult	01.08.2012	Scar	+	-	
BSc4	B	♀	Adult	01.08.2012	Scar	+	-	
ASc5	A	♀	Adult	23.11.2012	Scar	+	-	
BSc6	B	♀	Adult	31.08.2012	Scar	+	-	
BSc7	B	♀	Adult	15.06.2012	Scar	+	-	
BSc8	B	♀	Adult	31.08.2012	Scar	+	+	MK251001
BSc10	B	♂	Adult	01.11.2012	Scar	+	-	
BSc11	B	♂	Adult	31.08.2012	Scar	+	+	MK250996
BSc12	B	♂	Adult	15.06.2012	Scar	+	+	MK250997
BSc15	B	♂	Adult	13.10.2012	Scar	+	+	MK251002
ASc16	A	♀	Adult	23.11.2012	Scar	+	-	
ASc17	A	♂	Adult	09.11.2012	Scar	+	-	

Appendix I (2). (continued)

Code	Site	Sex	Life stage	Collection date	Morph	CHCs	COI	Accession Number
BSc18	B	♂	Adult	31.08.2012	Scar	+	+	MK250998
BSc19	B	♀	Adult	21.09.2012	Scar	+	-	
BSc20	B	♀	Adult	01.08.2012	Scar	+	-	
ASc21	A	♀	Adult	05.10.2012	Scar	+	-	
BSc22	B	♂	Adult	01.08.2012	Scar	+	-	
ASc23	A	♀	Adult	23.11.2012	Scar	+	-	
ASc24	A	♀	Adult	23.11.2012	Scar	+	-	
BSc25	B	♂	Adult	19.10.2012	Scar	+	-	
BSc26	B	♀	Adult	16.11.2012	Scar	+	-	

Appendix I (3) (continued)

Fig. 1




Fig. 1. Phylogenetic tree built using COI gene sequences and Bayesian inference. The branch support values are Bayesian posterior probabilities. The reference sequences are labeled (Genbank or BOLD accession code followed by species name). The brackets on the right side of the tree indicate the species identity of the specimens.

Fig. 2




Fig. 3




Fig. 4




Fig. 5

