



Detectability of nontrivial topologies

M. Kunz, N. Aghanim, A. Riazuelo, O. Forni

► To cite this version:

M. Kunz, N. Aghanim, A. Riazuelo, O. Forni. Detectability of nontrivial topologies. Physical Review D, 2008, 77, <10.1103/PhysRevD.77.023525>. <hal-03646509>

HAL Id: hal-03646509

<https://hal.science/hal-03646509v1>

Submitted on 2 May 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Detectability of nontrivial topologies

M. Kunz,^{1,*} N. Aghanim,^{2,†} A. Riazuelo,^{3,‡} and O. Forni^{4,§}

¹*Département de Physique Théorique, Université de Genève, 24 quai Ernest Ansermet, CH-1211 Geneva 4, Switzerland*

²*IAS, CNRS & Univ. Paris-Sud, Bât. 121, F-91405, Orsay Cedex, France*

³*Institut d'Astrophysique de Paris, UMR7095 CNRS, Université Pierre & Marie Curie, 98 bis boulevard Arago, 75014 Paris, France*

⁴*CESR, 9 Avenue du Colonel Roche, B.P. 44346, F-31028 Toulouse, Cedex 4, France*

(Received 7 May 2007; revised manuscript received 7 December 2007; published 23 January 2008)

We study how the uncertainty in the cosmological parameters impacts on the detection of topological signals, focussing on three cubic torus universes and using three tests: the information content, the S/N statistic, and the Bayesian evidence. We find, within the concordance cosmological model, that 3D torus universes with a size of $\sim 29 \text{ Gpc}^3$ or larger cannot be detected. For the toroidal models that can be detected, the detection significance is primarily influenced by Ω_Λ , which enters both in the noise amplitude due to the Integrated Sachs-Wolfe effect and in the size of the causal horizon which limits the accessible fundamental domain. On large angular scales $\ell < 40$, only Ω_Λ significantly alters the detection for all three estimators considered here.

DOI: [10.1103/PhysRevD.77.023525](https://doi.org/10.1103/PhysRevD.77.023525)

PACS numbers: 98.80.Es, 04.20.Gz, 98.70.Vc, 98.80.Bp

I. INTRODUCTION

One of the fundamental questions of cosmology is whether the universe is infinite or finite, a question which is especially relevant in the context of string theory where several dimensions may be compact. With the move towards precision cosmology, it has become possible to study the topology of the universe up to the causal horizon which fundamentally limits our view of the world.

Many tests have recently been developed to detect and constrain the topology of the universe, see Refs. [1,2] and references therein. Observationally a nontrivial topology can be detected directly with the observed map of cosmic microwave background (CMB) temperature and polarization fluctuations through matched and correlated circles in the CMB [3–8]. The topology can also be searched for in the expansion of the CMB map in terms of spherical harmonics, $T(x) = \sum_{\ell,m} a_{\ell m} Y_{\ell m}(x)$, where x are the pixels and both the temperature field and the $a_{\ell m}$ are random variables. The presence of a nontrivial topology introduces preferred directions, which breaks global isotropy and induces correlations between off-diagonal elements of the two-point correlation matrix $\langle a_{\ell m} a_{\ell' m'}^* \rangle$. Methods to find such off-diagonal correlations and to put optimal constraints on the size of the universe were proposed in Ref. [9]. They are based on matching the measured correlations to a given correlation matrix through a correlation coefficient, as well as a maximum likelihood method and Bayesian model comparison.

Recently the focus was purely on trying to detect any nontrivial topologies. However, to determine reliable limits, it is also necessary to understand all effects which can

affect the detection. The integrated Sachs-Wolfe (ISW) effect was identified to be an important noise for nontrivial topology searches since COBE [10,11]. For this purpose, we study the impact on the detection of topological signals of changing the cosmological parameters in the ranges allowed by present data. This should be seen as a step towards a rigorous search for the presence of signs of nontrivial topologies in the CMB. Eventually, such an analysis should account for other contributions like foreground emissions as well as sky cuts.

One additional reason to perform such a detailed analysis comes from statistics. The potentially most direct way to quantify the probability that our universe is infinite is to compute the model probability with Bayesian evidence. To do this, one needs to integrate over all parameters, which means that one needs to know the behavior of the likelihood for a topology under changes of the parameters.

This paper is organized as follows: we start by reviewing our notation and the likelihood-based probes of topology. We then study the main physical effects that are relevant for topology studies. Finally, we discuss how changes in the cosmological parameters affect our detection ability, and what this means for realistic universes where the parameters are not known with absolute certainty.

II. METHODOLOGY

In this section, all the simulations (as described in Refs. [12,13]) are based on a flat Λ CDM model with $\Omega_\Lambda = 0.7$, a reduced Hubble parameter of $h = 0.01 H_0 \times (1 \text{ Km s}^{-1} \text{ Mpc}^{-1})^{-1} = 0.67$, a Harrison-Zel'dovich initial power spectrum ($n_s = 1$) and a baryon density of $\Omega_b h^2 = 0.019$. With this choice of cosmological parameters we find a Hubble radius $c/H_0 \approx 4.8 \text{ Gpc}$, while the radius of the particle horizon is $R_h \approx 15.6 \text{ Gpc}$. In this paper we limit ourselves to cubic tori, denoted as T[X, X, X] where X is the size of the fundamental domain, in units

*Martin.Kunz@physics.unige.ch

†Nabila.Aghanim@ias.u-psud.fr

‡riazuelo@iap.fr

§Olivier.Forni@cesr.fr

of the Hubble radius. As an example, $T[4, 4, 4]$ is a cubic torus of size $(19.3\text{Gpc})^3$. The volume of such a torus is nearly half that of the observable universe. The diameter of the particle horizon is about $6.5c/H_0$ (cf. Sec. III A). We will focus on three different sizes, $T[2, 2, 2]$, $T[4, 4, 4]$, and $T[6, 6, 6]$, smaller universes being already ruled out and larger sizes having very little effect on the CMB. We combine the ℓ and m indices of the spherical harmonic coefficients to a single index $s \equiv \ell(\ell + 1) + m$ and mix both notations frequently.

In this paper we use the likelihood-based methods of Ref. [9], which we review here quickly. The CMB likelihood is taken to be a multivariate Gaussian distribution with covariance matrix $\mathcal{B}_{ss'} = \langle a_s^* a_{s'} \rangle$ and zero mean, $\langle a_s \rangle = 0$. This covariance matrix provides the full statistical description of the CMB for a given topology. The likelihood function then is

$$\mathcal{L}(\mathcal{B}) \equiv p(a_{\ell m}|\mathcal{B}) \propto \frac{1}{\sqrt{|\mathcal{B}|}} \exp\left[-\frac{1}{2} \sum_{s,s'} a_s^* \mathcal{B}_{ss'}^{-1} a_{s'}\right], \quad (1)$$

where $|\mathcal{B}|$ is the determinant of the matrix \mathcal{B} . Any further model assumptions like the values of the cosmological parameters are implicitly included in the choice of \mathcal{B} .

Often it is easier to consider the logarithm of the likelihood, $\ln \mathcal{L} = -1/2(\ln|\mathcal{B}| + \chi^2 + s_{\max} \ln(2\pi))$ where we have defined

$$\chi^2 \equiv \sum_{s,s'} a_s^* \mathcal{B}_{ss'}^{-1} a_{s'}. \quad (2)$$

Writing $\langle a_s^* a_{s'} \rangle = \mathcal{A}_{ss'}$ for the expectation value of the two-point function of the “sky” $a_{\ell m}$, we can compute the expectation value of the χ^2 ,

$$\langle \chi^2 \rangle = \sum_{ss'} \langle a_s^* a_{s'} \rangle \mathcal{B}_{ss'}^{-1} = \text{tr}(\mathcal{A} \mathcal{B}^{-1}), \quad (3)$$

where we have used the Hermiticity of the correlation matrices. For the two special cases of an infinite universe and one for which $\mathcal{A} = \mathcal{B}$ (i.e. the measured $a_{\ell m}$ are drawn from the template covariance matrix \mathcal{B}), we find

$$\langle \chi^2 \rangle_{\infty} = \sum_s C_s \mathcal{B}_{ss}^{-1}, \quad (4)$$

$$\langle \chi^2 \rangle_{\mathcal{B}} = \text{tr}(1) = s_{\max}. \quad (5)$$

As the $a_{\ell m}$ are Gaussian random variables, we expect to find that χ^2 is distributed with a χ^2 -like distribution. The general expression for the variance is rather cumbersome, but for the two special cases we find

$$\sigma_{\mathcal{B}}^2 \equiv \langle (\chi^2)^2 \rangle_{\mathcal{B}} - \langle \chi^2 \rangle_{\mathcal{B}}^2 = 2s_{\max} \quad (6)$$

and

$$\sigma_{\infty}^2 = 2 \sum_{ss'} C_s C_{s'} |\mathcal{B}_{ss'}^{-1}|^2. \quad (7)$$

To improve the stability of the algorithm against misesti-

mates of the power spectrum we whiten the $a_{\ell m}$ by dividing out the power spectrum

$$C_{\ell} = \frac{1}{2\ell + 1} \sum_m |a_{\ell m}|^2, \quad (8)$$

via the prescription

$$\mathcal{B}_{ss'} \rightarrow \frac{\mathcal{B}_{ss'}}{\sqrt{C_s C_{s'}}}, \quad (9)$$

$$a_{\ell m} \rightarrow \frac{a_{\ell m}}{\sqrt{C_{\ell}}}. \quad (10)$$

The power spectrum of the $a_{\ell m}$ has to be estimated from the data, which changes their distribution. We found in Ref. [9] that this tends to increase the sensitivity of the estimators discussed below.

A. Information content

As discussed in Ref. [9] it is possible to study directly the information content in the CMB sky. For Gaussian fluctuations, the infinite universe contains a maximum of information (maximal entropy), while a finite (“small”) universe is less random. Mathematically this is expressed through the nonzero correlations in the off-diagonal part of the two-point function which make it easier to predict the temperature distribution (at least in theory).

A commonly used measure of information content is given by the Kullback-Leibler (KL) divergence between two probability distributions p and q ,

$$D_{\text{KL}}(p||q) \equiv \int p \log\left(\frac{p}{q}\right). \quad (11)$$

This quantity measures the relative information entropy between the two distributions. It is often used in information theory. For example, if information distributed according to q is transmitted with a coding based on p then $D_{\text{KL}}(p||q)$ describes how much bandwidth is wasted because of the wrong coding (the base of the logarithm is equivalent to a choice of units in which the KL divergence is expressed). It is also possible to view the KL divergence as a regularized entropy for continuous random variables, as it satisfies the Gibbs inequality, $D_{\text{KL}}(p||q) \geq 0$ with equality only for $p = q$. Here we use the latter interpretation.

In our case we are dealing with Gaussian probability distributions characterized by zero mean and covariance matrices \mathcal{A} and \mathcal{B} , for which the KL divergence can be evaluated directly,

$$\begin{aligned} & \int p(\mathcal{A}) \log \frac{p(\mathcal{A})}{p(\mathcal{B})} \\ &= \frac{1}{2} [\log|\mathcal{B}| - \log|\mathcal{A}| - \text{tr}(1 - \mathcal{B}^{-1} \mathcal{A})]. \end{aligned} \quad (12)$$

Unfortunately the KL divergence is not symmetric in its




FIG. 1 (color online). The information content as a function of the size of the fundamental domain and of ℓ_{\max} : The left panel shows H and right panel the increase per ℓ , ΔH , for (from the top downwards) $T[2, 2, 2]$ [solid (red) line], $T[4, 4, 4]$ [dashed (blue) line], and $T[6, 6, 6]$ [dotted (green) line].

arguments, making its interpretation somewhat difficult. It is also not rotationally invariant, so that it depends on the relative orientation of the coordinate systems in which the correlation matrices are expressed. We will therefore only consider a special case where we take $p(\mathcal{B})$ to be the distribution of an infinite universe. In this case the KL divergence is the relative entropy to the topology described by the correlation matrix \mathcal{A} . This is both rotationally invariant (as the infinite universe is isotropic) and has a well-defined interpretation. It is also very easy to compute: For the infinite universe $\mathcal{B} = \mathcal{B}^{-1} = 1$. The whitening enforces $\text{tr}(\mathcal{A}) = s_{\max} = \text{tr}(1)$, so that the relative entropy of \mathcal{B} is just

$$H(\mathcal{A}) \equiv D_{\text{KL}}(\mathcal{A}||1) = -\frac{1}{2}\log|\mathcal{A}|. \quad (13)$$

This quantity describes how much less information is present in CMB data of the (easier to predict) finite universe, as compared to the information of the CMB of an infinite universe. This gives an *upper limit* on the chance to detect a given topology.

We plot in Fig. 1 both information content H as a function of ℓ_{\max} (left panel) as well as the increase of H per ℓ (right panel) for the three topological models $T[2, 2, 2]$ [solid (red) line], $T[4, 4, 4]$ [dashed (blue) line] and $T[6, 6, 6]$ [dotted (green) line], respectively. The information content decreases as expected for topologies closer to an infinite universe ($T[6, 6, 6]$ in our case).

B. Measuring topology

While the previous subsection deals with a purely theoretical measure of detectability, namely, the information content in a given map or matrix, we apply in this section to two actual detection algorithms discussed in Ref. [9]. For both we use likelihood, Eq. (1), or its logarithm χ^2 , Eq. (2). We exploit the fact that the likelihood will be higher if the observed map and template agree.

For the first analysis we compute the correlation between an observed map and a template and we search the global minimum of χ^2 over all rotations of the $a_{\ell m}$ (corresponding to relative orientations of the observed map and template). In order to quantify possible false detections of the topological signal, we give an estimate of the null hypothesis, that is the possibility for an infinite universe to be taken for a universe with a nontrivial topology. We use the following procedure: For each given topology ($T[2, 2, 2]$, $T[4, 4, 4]$, $T[6, 6, 6]$) we fix the template and generate several thousand random maps both from the chosen template and for an infinite universe. We then use the mean and standard deviation for both distributions to compute a signal-to-noise statistic,

$$S/N(\mathcal{B}, \chi^2) = \frac{|\langle \chi^2 \rangle_{\infty} - \langle \chi^2 \rangle_{\mathcal{B}}|}{\sqrt{\sigma(\chi^2)_{\infty}^2 + \sigma(\chi^2)_{\mathcal{B}}^2}}. \quad (14)$$

This corresponds to the distance between the means in units of the errors added in quadrature. We show results in Fig. 2, with the same line styles (colors) as in Fig. 1.




FIG. 2 (color online). S/N for (from the top downwards) $T[2, 2, 2]$ [solid (red) line], $T[4, 4, 4]$ [dashed (blue) line], and $T[6, 6, 6]$ [dotted (green) line].

Again, we see that larger universes are more difficult to detect. The $T[6, 6, 6]$ universe is barely detectable, while the others can be detected easily for $\ell_{\max} > 16$.

The above test is somewhat ad-hoc so that we compute additionally the probability that the measured $a_{\ell m}$ are actually drawn from the distribution described by the correlation matrix \mathcal{B} . This is often called the Bayesian evidence in astrophysical applications. Formally, the probability of a topology (specified here via \mathcal{B} , which additionally depends on the cosmological parameters θ) is $p(\mathcal{B}|a_{\ell m}, \theta)$, which we can connect to the likelihood, Eq. (1) through Bayes theorem,

$$p(\mathcal{B}|a_{\ell m}, \theta) = p(a_{\ell m}|\mathcal{B}, \theta) \frac{p(\mathcal{B})}{p(a_{\ell m})} \propto \mathcal{L}(\mathcal{B})p(\mathcal{B}). \quad (15)$$

We will be interested in ratios of these probabilities for different \mathcal{B} (describing different topologies). As we have no *a priori* knowledge of the topology, we will take $p(\mathcal{B})$ to be constant. However, the numerical value of $\mathcal{L}(\mathcal{B})$ depends on the orientation of the template, so that we need to marginalize over the angles (see Ref. [9] for more details, and [14–16] for other examples of the use of Bayesian evidence in topology). Unfortunately the likelihood is very strongly peaked in the vicinity of correct alignments. The numerical integration is difficult, even using adaptive schemes.

Having computed $p(\mathcal{B}|a_{\ell m})$ we define a relative evidence for a topology as the ratio of this probability to the one for an infinite universe,




FIG. 3 (color online). Evidence for (from the top downwards) $T[2, 2, 2]$ [solid (red) line], $T[4, 4, 4]$ [dashed (blue) line], and $T[6, 6, 6]$ [dotted (green) line] relative to an infinite universe, for the full-sky reconstructed $a_{\ell m}$ of [17].

$$\Delta \log E(\mathcal{B}) = \log p(\mathcal{B}|a_{\ell m}) - \log p(1|a_{\ell m}). \quad (16)$$

This gives the relative probability that the $a_{\ell m}$ are due to a universe described by the topology and cosmology encoded in \mathcal{B} , compared to the probability that they are due to an infinite universe. Figure 3 shows the strong dependence of the evidence on the size of fundamental domain; the figure looks somewhat similar to the one of H . For this figure we used the $a_{\ell m}$ of the full-sky reconstruction by [17], and we fixed the cosmological parameters of the templates to their fiducial values (the central values of Table I). One can further eliminate the dependence on the cosmological parameters by marginalizing (integrating) over them as well,

$$\begin{aligned} p(\mathcal{B}|a_{\ell m}) &= \int d\theta p(\mathcal{B}, \theta|a_{\ell m}) \\ &= \int d\theta p(\mathcal{B}|a_{\ell m}, \theta) p(\theta|a_{\ell m}). \end{aligned} \quad (17)$$

We see that the integral of Eq. (15) over θ is weighted by the posterior of the parameters, $p(\theta|a_{\ell m})$. One of the motivations for this paper is to study the dependence of $\Delta \log E(\mathcal{B})$ on the cosmological parameters and to under-

TABLE I. Parameter values investigated.

h	$\{0.678, 0.724, 0.770\}$
n_s	$\{0.916, 0.950, 0.982\}$
Ω_Λ	$\{0.50, 0.70, 0.75, 0.80, 0.90\}$

stand which integrals are necessary. Finally, by also marginalizing over a class of topologies (e.g. the X of cubic tori $T[X, X, X]$) and using the actually measured $a_{\ell m}$, we can compute the probability that our universe has that kind of topology.

III. PHYSICAL EFFECTS AT LARGE SCALES

The shape of the universe leaves specific imprints on large scale CMB anisotropies at multipoles lower than $\ell \sim 60$ that can be detected via the correlation matrix of the $a_{\ell m}$ coefficients. Such scales suffer many problems. First, it is at these scales that cosmic variance $\Delta C_\ell = \sqrt{2/(2\ell + 1)}C_\ell$ is the largest. Then, at those scales several physical processes affect the CMB signal which may change our ability to detect nontrivial topologies and to measure their properties. In the following we will focus on the ISW effect, in addition to the reionisation and to the relative size of the particle horizon. Other contributions at low ℓ could come from gravitational waves or defects. But their contribution to the $a_{\ell m}$ should be mostly uncorrelated with the topological signal and thus act as an additional noise. Present analyses [18,19] have restricted the contributions of primordial tensor modes and defects to be subdominant. In the following, we will thus consider them as a negligible source of “noise” for measurements of the topology.

In the next section, we will vary the relevant cosmological parameters in the range allowed by their current posterior distribution [from the WMAP3 (Wilkinson Microwave Anisotropy Probe 3-year) data] and study their influence on the detection of the topological signal.

A. Relative size of fundamental domain and particle horizon

As discussed in the introduction, we fix the size of fundamental domain in units of c/H_0 . Another key scale is the distance to the last scattering surface which is very close to the size of the particle horizon. In the following, we will not discriminate between these two. As light rays move on geodesics with $ds^2 = 0$, we can compute the particle horizon today in comoving coordinates from

$$\eta_0 = \int_0^{t_0} \frac{cdt}{a(t)} = c \int_0^1 \frac{da}{a^2 H(a)}. \quad (18)$$

The quantity relevant to us is the diameter of the particle horizon D (the “size of the causal domain”), which, in units of c/H_0 is

$$D = \frac{2a_0 H_0 \eta_0}{c} = 2 \int_0^1 dx (\omega_r h^{-2} + \Omega_m x + \Omega_\Lambda x^4)^{-1/2}, \quad (19)$$

where the new integration variable is the normalized scale factor $x \equiv a/a_0$, and we have assumed flat spacelike sections and dark energy in the form of a pure cosmological

constant. The energy density in radiation $\omega_r \equiv \Omega_r h^2$ is fixed very precisely by the temperature of the CMB (assuming massless neutrini) to 4.2×10^{-5} . Its presence changes the value of the above integral roughly by $\sqrt{\Omega_r} \approx 0.01$, i.e. the residual influence of the Hubble constant is of the order of 1% only. Since we assume that the universe is flat, we have $\Omega_m = 1 - \Omega_r - \Omega_\Lambda$. In reality, the important scale is rather the distance to the last scattering surface, so that we should set the lower limit of integration to $x_{\text{LSS}} \approx 1/1090$. This modifies the result by about 2% but does not change the conclusions.

Obviously, the strength of the topological signal depends (among other things) on the ratio of the two scales. A universe that is much smaller than the causal domain leaves a strong imprint in the CMB, while one that is much larger will be impossible to distinguish from a truly infinite universe. The value of D corresponds roughly to the largest X that we can hope to detect. We show in Figs. 1–3 how the topology becomes easier to detect as the fundamental domain becomes smaller (for a fixed size of the particle horizon).

For a fixed size of the fundamental domain, the parameter affecting the ratio of the two scales is Ω_Λ as it changes D , see Fig. 4. If the matter density is very large, $\Omega_m \approx 1$, then the size of causal domain corresponds roughly to the size of a $T[4, 4, 4]$ universe (in a purely matter-dominated universe, the horizon size equals twice the Hubble radius, hence $D = 4$). On the other hand, the causal domain of a flat low-density universe can have an over 1000 times larger volume.




FIG. 4. Diameter of the particle horizon, $D = 2a_0\eta_0 H_0/c$, for a flat universe and as a function of Ω_Λ . This corresponds roughly to the maximal fundamental domain size X that we can hope to detect.

The importance of the correct orientation of the fundamental domain was already discussed in Ref. [9]. We found that the orientation is one of the critical parameters and needs to be taken into account in any analysis. This can be done by averaging quantities over the orientation (as is done for a Bayesian study), or by taking the maximum over all orientations (with the matched filter, or also when looking for correlated circles).

B. ISW effect

The CMB photons are influenced by the change in the gravitational potential through which they must pass. The effects arising from time-variable metric perturbations are generally known as the integrated Sachs-Wolfe effect [20] in the linear regime and go by the names of Rees-Sciama effect [21] and the effect from moving halos [22] in the nonlinear regime. The ISW effect depends on the time derivative of the gravitational potential. The anisotropies due to the ISW effect thus depend on the parameters of the background cosmology and are also tightly coupled to the clustering and the spatial and temporal evolution of the intervening structures. The most important contribution for topological studies is the late ISW effect due to the decay of the gravitational potential [23,24]. This happens in a low matter density universe at the onset of dark energy (or spatial curvature) domination, when the increased rate of expansion of the universe reduces the amplitude of gravitational potential.

The power spectrum of the ISW can be written as a function of the power spectrum of the potential $P_\Phi(k)$ like

$$C_\ell = (4\pi)^2 \int dk k^2 P_\Phi(k) \left[\int_{\eta_{\text{LSS}}}^{\eta_0} 2\dot{F}(\eta) j_\ell(kr) d\eta \right]^2, \quad (20)$$

where the last scattering surface subscript stands for the epoch of last scattering, $j_\ell(x)$ is the spherical Bessel function, and r is the comoving distance. $F(k, \eta) = G/x$ is the growth rate of the potential, where x is the (normalized) scale factor and G is the linear growth factor. The potential is related to the matter density contrast δ via the Poisson equation, $k^2\Phi = -\frac{3}{2}a_0^2\Omega_m H_0^2 \delta/x$. In the matter-dominated era, the density contrast grows as x and therefore the gravitational potential is constant. This is no longer the case in the dark-energy dominated era.

The ISW effect is seen mainly in the lowest ℓ -values of the power spectrum where the cosmic variance is large. This makes its direct measurement very difficult. Since the time evolution of the potential that gives rise to the ISW effect may also be probed by observations of the large scale structure, one expects the ISW to be correlated with tracers of the large scale structure (see, e.g. Ref. [25]). This could conceivably help to remove it at least partially. We show in Fig. 5 the information content H of our standard T[4, 4, 4] model [dashed (blue) line] and compare it to a fictitious case where the ISW effect was turned off [solid (red) line]; the anisotropies are then only generated at last scattering.




FIG. 5 (color online). Influence of the ISW effect on the information content: The dashed (blue) line shows the H for the T[4, 4, 4] model while the solid (red) line shows the information content for the same model but without ISW effect. The difference is very important.

The ISW is clearly a very strong effect that needs to be taken into account correctly, as shown by [10,11]. We also checked its influence on the detection algorithms discussed in Sec. II B and found the same result.

C. Reionisation

At reionisation, CMB photons scatter off free electrons, altering the CMB temperature anisotropies and inducing a polarized signal. However the relevant effects in the case of topological studies are mostly those induced by reionisation on the CMB temperature anisotropies. At reionisation, the photon-electron interactions randomize the directions of propagation of the fraction of rescattered photons. As a result, the small scale primary CMB anisotropies are suppressed and the power in the acoustic peaks is damped by a factor $\exp(-2\tau)$, where τ is the Thomson optical depth. Scales approaching the horizon at reionization are barely affected by the photon rescattering, and the Doppler induced anisotropies are much smaller than the ISW contribution at the same scales [26]. Therefore, we do not expect reionization to alter much the detection of nontrivial universe signatures at small ℓ s. We show its effect on the detectability in Fig. 6 by comparing a universe with no reionization [solid (red) line] and a universe with a large optical depth, compatible with the WMAP1 constraints [$\tau = 0.166$, dashed (blue) line]. The induced changes are negligible. The latest WMAP results suggest a significantly lower optical depth; therefore, the effect of reionization is




FIG. 6 (color online). Influence of reionization on the information content: The dashed (blue) line shows the H for the T[4, 4, 4] model while the solid (red) line shows the information content for the same model but without reionization. The difference is negligible.

even smaller, and we do not investigate further the variations of τ in the study.

IV. SENSITIVITY TO COSMOLOGICAL PARAMETERS

We choose a fiducial model motivated by the WMAP3 + SNLS results for a flat Λ CDM model [18], with $H_0 = 72.4 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $n_s = 0.95$, $\Omega_\Lambda = 0.75$, $\Omega_b h^2 = 0.0234$, and $\tau = 0.085$. We then explore the effect of the cosmological parameters on the detection of the topological signal by varying in turn H_0 , n_s , and Ω_Λ (keeping all other parameters fixed). Table I shows the range of variation, corresponding to roughly the two-sigma limits of WMAP3 (with the exception of Ω_Λ where we also investigate other values). We do not expect the normalization A_s or the baryon density $\Omega_b h^2$ to play an important role. In Sec. III C we further saw that there is only a small difference between $\tau = 0$ and the fiducial case. We conclude that variations of τ are also uncritical for measuring topology.

We apply our three estimators (the information content, the S/N statistic, and the Bayesian evidence) to the 3D torus universes T[2, 2, 2], T[4, 4, 4], and T[6, 6, 6] in the concordance cosmological model. We find, in an ideal configuration (no foreground contamination and no instrumental effects) but accounting for the noise introduced by the ISW, that T[6, 6, 6] universe is barely detected. Larger toroidal universes will thus be impossible to distinguish

from an infinite universe. We therefore limit ourselves to showing the results of varying cosmological parameters for the T[4, 4, 4] case only since it is more realistic than T[2, 2, 2].

We show in the top panel of Fig. 7 how the information content H depends on the three values of n_s (cf Table I). The other two panels show how such a variation of n_s affects the detection of a T[4, 4, 4] topology based on our fiducial model. To this end we simulated maps with the




FIG. 7 (color online). For T[4, 4, 4], the effect of varying n_s : The top panel shows the information content, for the fiducial model [solid (black) line, $n_s = 0.950$] as well as $n_s = 0.916$ [dashed (red) line], and $n_s = 0.982$ [dotted (blue) line]. The middle panel (S/N) and bottom panel (evidence) show how the detection changes when testing simulated maps against the fiducial model. The (black) solid line uses a map derived from the fiducial model, while the (red) dashed and (blue) dotted lines are for the smaller and larger n_s , respectively.

different values of n_S (and all other parameters fixed to the fiducial value) and then tested them against our fiducial model, applying the algorithms of Sec. II B. We find that the information content does not vary significantly. Also the detections measured by S/N (middle panel) and the Bayesian evidence (bottom panel) do not change for $\ell_{\max} < 10$ with only small changes at higher resolutions (within the statistical fluctuations induced by the map realizations). At least for the relatively small range of

scales explored in this study ($\ell < 40$) the whitening of the spectrum suffices to ensure that the tilt has only marginal impact on the topological signatures. We conclude that n_S can safely be fixed to the central value in Table I.

The reduced Hubble parameter h intervenes both in the size of the particle horizon (Sec. III A) and in the ISW effect (Sec. III B). We therefore expect some dependence on its value. Repeating the same analyses as for n_S , we find that the results depend even less on h than on n_S , see




FIG. 8 (color online). The effect of varying h for T[4, 4, 4]: The top panel shows the information content, for the fiducial model [solid (black) line, $h = 0.724$] as well as $h = 0.678$ [dashed (red) line], and $h = 0.770$ [dotted (blue) line]. The middle panel (S/N) and bottom panel (evidence) show how the detection changes when testing simulated maps against the fiducial model. The (black) solid line uses a map derived from the fiducial model, while the (red) dashed and (blue) dotted lines are for the smaller and larger h , respectively.




FIG. 9 (color online). The effect of varying Ω_Λ for T[4, 4, 4]: The top panel shows the information content, for the fiducial model [solid (black) line, $\Omega_\Lambda = 0.75$] as well as $\Omega_\Lambda = 0.80$ [dashed (red) line], $\Omega_\Lambda = 0.90$ [dotted-dashed (magenta) line], $\Omega_\Lambda = 0.7$ [dotted (blue) line], and $\Omega_\Lambda = 0.50$ [long-dashed (cyan) line]. The middle panel (S/N) and bottom panel (evidence) show how the detection changes when testing simulated maps against the fiducial model. The (black) solid line uses a map derived from the fiducial model. The line styles of the other models are the same as those of the information content (top panel).

Fig. 8. Clearly, h can also be fixed to the central value in Table I.

We follow the same procedure as above, varying Ω_Λ only, for the values given in Table I. In the top panel of Fig. 9 we plot the information content for these models, with the largest Ω_Λ leading to the highest H . Ω_Λ intervenes both in the causal domain and in the ISW amplitude. The larger Ω_Λ , the larger the causal domain, but also the bigger the ISW contribution. Its effect on the detection of topology is thus a combination of both dependencies. The graph suggests that the effect due to the size of the fundamental domain dominates over the increase in “ISW noise.” As before, we then investigate how the detection changes if map and template have a different Ω_Λ . In contrast to the two previous parameters, we find that there is a strong variation in the detectability. We only have a high S/N if map and template agree (black solid line in the middle panel). The maps with the next higher [dashed (red) line] and lower [dotted (blue) line] values of Ω_Λ are already below a significant detection limit, and the other cases lie even lower. The evidence shows the same behavior, with the fiducial template leading to a much higher model probability than the others. A full analysis requires one to explore all acceptable values of the cosmological parameters. Given that the topological signal remains nearly constant for all parameters except Ω_Λ , it suffices to vary only Ω_Λ within the 2σ region allowed by WMAP3 + SNLS.

V. CONCLUSIONS

We have studied T[2, 2, 2], T[4, 4, 4], and T[6, 6, 6] cubic torus universes, investigating how the detection probability changes as a function of the cosmological parameters. Within the current 2σ ranges and for $\ell_{\max} < 40$, we find that the optical depth and thus reionization as well as the tilt of the primordial power spectrum have negligible effect, as does the Hubble constant. The cosmological constant in turn changes the topological signal

strongly. It enters in the two most relevant effects for studying topology: the ISW which is not correlated with the topological signal and so acts as an additional noise and the size of the causal domain D (which is obviously the key quantity limiting our ability to detect a given topology). Topological defects and tensor modes also contribute as additional noise but they are observationally constrained to a lower amplitude than the ISW effect and can be neglected for now. An additional important parameter is the relative orientation of the fundamental domain [9]. The methods we use in the present study take this already into account so that we do not discuss it in detail.

When searching for traces of a nontrivial topology, we have to test *a priori* templates for all cosmological models, varying the full set of cosmological parameters, which would require computing templates for a large number of possible models. We have shown here that, given the constraints from WMAP3 + SNLS and the range of scales that we explore ($\ell < 40$), only Ω_Λ is able to affect significantly the detections. Templates for several values of Ω_Λ are thus required for a complete analysis, but the other parameters can be held fixed at their maximum likelihood values, significantly reducing the complexity of a full analysis.

As a byproduct, we also investigated what is the largest 3D torus that can be detected, using WMAP3 data. We find that only torus models with a volume smaller than $\sim 29 \text{ Gpc}^3$ can be detected in the current concordance model.

ACKNOWLEDGMENTS

M. K. thanks Ruth Durrer for interesting discussions and acknowledges financial support from the Swiss NSF. The authors acknowledge partial support from Programme National de Cosmologie. N.A. and O.F. thank the University of Geneva and M. K. thanks IAS for hospitality. We acknowledge the use of HEALPix [27] to manipulate the CMB maps.

-
- [1] M. Lachièze-Rey and J.-P. Luminet, Phys. Rep. **254**, 135 (1995).
 - [2] J. Levin, Phys. Rep. **365**, 251 (2002).
 - [3] N. J. Cornish *et al.*, Phys. Rev. Lett. **92**, 201302 (2004).
 - [4] B. Roukema *et al.*, Astron. Astrophys. **423**, 821 (2004).
 - [5] J.-P. Luminet, Phys. World **18**, 22 (2005).
 - [6] N. J. Cornish, D. Spergel, and G. Starkmann, Classical Quantum Gravity **15**, 2657 (1998).
 - [7] J. G. Cresswell *et al.*, Phys. Rev. D **73**, 041302 (2006).
 - [8] A. Riazuelo *et al.*, arXiv:astro-ph/0601433.
 - [9] M. Kunz *et al.*, Phys. Rev. D **73**, 023511 (2006).
 - [10] N. J. Cornish and D. Spergel, Phys. Rev. D **62**, 087304 (2000).
 - [11] K. T. Inoue, Classical Quantum Gravity **18**, 1967 (2001).
 - [12] A. Riazuelo *et al.*, Phys. Rev. D **69**, 103514 (2004).
 - [13] A. Riazuelo *et al.*, Phys. Rev. D **69**, 103518 (2004).
 - [14] K. T. Inoue and N. Sugiyama, Phys. Rev. D **67**, 043003 (2003).
 - [15] A. Niarchou and A. Jaffe, AIP Conf. Proc. **848**, 774 (2006).
 - [16] A. Niarchou and A. Jaffe, Phys. Rev. Lett. **99**, 081302 (2007).
 - [17] H. K. Eriksen *et al.*, Astrophys. J. **612**, 633 (2004).
 - [18] D. N. Spergel *et al.*, Astrophys. J. Suppl. Ser. **170**, 377 (2007).

- [19] N. Bevis *et al.*, arXiv:astro-ph/0702223 [Phys. Rev. Lett. (to be published)].
- [20] R. K. Sachs and A. M. Wolfe, *Astrophys. J.* **147**, 73 (1967).
- [21] M. J. Rees and D. W. Sciama, *Nature (London)* **217**, 511 (1968).
- [22] N. Aghanim *et al.*, *Astron. Astrophys.* **334**, 409 (1998).
- [23] L. A. Kofman and A. A. Starobinsky, *Sov. Astron. Lett.* **11**, 271 (1985).
- [24] M. Kamionkowski and D. N. Spergel, *Astrophys. J.* **432**, 7 (1994).
- [25] R. G. Crittenden and N. Turok, *Phys. Rev. Lett.* **76**, 575 (1996).
- [26] W. Hu and S. Dodelson, *Annu. Rev. Astron. Astrophys.* **40**, 171 (2002).
- [27] K. M. Górski, É. Hivon, and B. D. Wandelt, arXiv:astro-ph/9812350; <http://www.eso.org/science/healpix/>.