



HAL
open science

Commentary: The Dynamics of Aerotaxis in a Simple Eukaryotic Model

Jean-Paul Rieu, Olivier Cochet-Escartin, Christophe Anjard, Mete Demircigil,
Vincent Calvez

► **To cite this version:**

Jean-Paul Rieu, Olivier Cochet-Escartin, Christophe Anjard, Mete Demircigil, Vincent Calvez. Commentary: The Dynamics of Aerotaxis in a Simple Eukaryotic Model. *Frontiers in Cell and Developmental Biology*, 2022, 10, pp.844812. 10.3389/fcell.2022.844812 . hal-03650465

HAL Id: hal-03650465

<https://hal.science/hal-03650465v1>

Submitted on 25 Apr 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Commentary: The Dynamics of Aerotaxis in a Simple Eukaryotic Model

Jean-Paul Rieu^{1*}, Olivier Cochet-Escartin¹, Christophe Anjard¹, Mete Demircigil² and Vincent Calvez²

¹Institut Lumière Matière, UMR5306, Université Lyon 1-CNRS, Université de Lyon, Villeurbanne, France, ²Institut Camille Jordan, UMR5208, Université Lyon 1-CNRS, Université de Lyon, Villeurbanne, France

Keywords: Self-generated gradients, Aerotaxis, Collective migration, Oxygen sensing, Dictyostelium discoideum

A Commentary on

The Dynamics of Aerotaxis in a Simple Eukaryotic Model

by Biondo, M., Panuzzo, C., Ali, S. M., Bozzaro, S., Osella, M., Bracco, E., and Pergolizzi, B. (2021). *Front. Cell Dev. Biol.* 9:720623. doi: 10.3389/fcell.2021.720623

We read with interest the article by Biondo et al. (2021) in *Frontiers in Cell and Developmental Biology*, “*The Dynamics of Aerotaxis in a Simple Eukaryotic Model*.” Reproducing the confinement assay we published in eLife earlier this year (Cochet-Escartin et al., 2021) with the same cell line, they found the same emergent behavior, i.e., the propagation of a ring of cells, which they named corona, from a dense, confined colony through the self-generation of oxygen gradients by cell consumption. The authors claimed that cell division plays no role in the phenomenon, whereas in our study, we insisted on its important role.

This message is wrong. In this commentary, we first clarify that ring formation is independent on cell division but that ring propagation over long times depends on it. Second, we discuss the possible experimental biases that may have led the authors to this conclusion.

Cell division is not necessary for ring formation but is necessary for its sustained propagation.

Biondo et al. observed exactly the same collective phenotype as us for the confined colony in a starving buffer that prevents cell division. A ring forms internally, but as soon as it reaches the colony edge (at ~6 h), it stops and cells aggregate (compare Movie 3 of Biondo et al. with our movie M6 and Figure 5; Supplementary Figure 2, Cochet-Escartin et al.). In contrast, in a nutrient medium, the ring propagates far away from the initial colony for days (see **Figure 1A** below). Biondo et al. neither commented on this fundamental difference between the two conditions nor on our model that demonstrates that cell division is necessary to maintain ring propagation even if it contributes little to the expansion speed (Figures 5A,B and Eq. 6, Cochet-Escartin et al.). Independently of any model, a simple mass balance equation for the total cell number N with N_B cells in the bulk region (core) and N_R cells in the ring region invalidates Biondo’s assertion that division plays no role:

$$N(t) = N_B(t) + N_R(t) = \rho_B \pi (R(t) - L)^2 + \rho_R 2\pi R(t)L \quad (1)$$

Using the experimental observations (Figures 1D,E, Supplementary Figure S3B in Cochet-Escartin et al., **Figures 1B–D** below) that the ring width L and density ρ_R and bulk density ρ_B are constant, and that the ring radius R is expanding at constant speed $R(t) = R_0 + \sigma t$, we predict that $N_B(t)$ increases faster with time (i.e., as $R^2 \sim t^2$) than $N_R(t)$ (i.e., as $LR \sim t$). Experimentally, up to 30 h, $N_B(t)$ increases faster than linearly with time while N_R increases linearly (**Figure 1C**). Initially $N_B/N_R = 1$, but after 24 h, $N_B/N_R = 1.8$, and after 47 h, $N_B/N_R = 2.8$ (**Figure 1C**). Hence, N_B largely contributes to the overall cell number increase $N(t)$. By comparison, Biondo et al. assume a constant $N_R(t)$, and they do not consider bulk cells at all.

OPEN ACCESS

Edited by:

Verena Ruprecht,
Centre for Genomic Regulation (CRG),
Spain

Reviewed by:

Jean Clairambault,
Institut National de Recherche en
Informatique et en Automatique
(INRIA), France

*Correspondence:

Jean-Paul Rieu
jean-paul.rieu@univ-lyon1.fr

Specialty section:

This article was submitted to
Molecular and Cellular Pathology,
a section of the journal
*Frontiers in Cell and Developmental
Biology*


Received: 28 December 2021

Accepted: 20 January 2022

Published: 02 March 2022

Citation:

Rieu J-P, Cochet-Escartin O, Anjard C,
Demircigil M and Calvez V (2022)
Commentary: The Dynamics of
Aerotaxis in a Simple
Eukaryotic Model.
Front. Cell Dev. Biol. 10:844812.
doi: 10.3389/fcell.2022.844812



Cell divisions hold in the ring. Our confined colony grows slower than exponentially (see solid black line with a typical 8 h doubling time (d’Alessandro et al., 2016) in **Figure 1C**), but it grows (i.e., $N(47\text{ h})/N(0\text{ h}) = 8.5$). In Cochet-Escartin et al., we propose a go-or-grow model where aerotaxis holds at low O_2 and cell division at high O_2 . The threshold is around 1% O_2 as estimated by direct aerotaxis investigations using microfluidic devices (Cochet-Escartin et al.) and from literature values for cell division in hypoxic conditions (Schiavo and Bisson, 1989; West et al., 2007). Such value corresponds to the O_2 level measured in the ring (Cochet-Escartin et al.). Hence, divisions occur mostly in the ring, but ring cells are constantly transferred to the bulk to maintain a constant ρ_B while R is increasing. This transfer occurs in our models (see Figure 7 of Cochet-Escartin et al.), but perhaps it was not sufficiently supported by data. In **Figures 1E,F**, we present manually tracked trajectories in the ring frame. A few ones displayed with a green or purple color enter or escape the outward ring position, canceling any ring-to-front flux. Far more trajectories are directed backward (i.e., ring to bulk, in blue). Interestingly, the measured flux of such a cell transfer, $\varphi_{R \rightarrow B} = 12\ \text{cells}/\text{mm}/\text{h}$, explains fairly well the bulk cell number increase using the following equation:

$$dN_B = 2\pi R \varphi_{R \rightarrow B} dt \quad (2)$$

The fit is displayed in **Figure 1C**.

Biondo et al. may have caught a transient regime only. Biondo et al. measured 3% and 5% O_2 in the bulk and ring regions, respectively (Supplementary Figure S1). Above 2% O_2 , aerotaxis should not hold (Cochet-Escartin et al.); the division

rate is fairly the same as in normoxic conditions (Schiavo and Bisson, 1989; West et al., 2007). A possible reason for this discrepancy is that O_2 is overestimated. Their measurements were performed with a commercial sensing film that is not compatible with transmission microscopy, contrary to the technology we developed in Cochet-Escartin et al. They may have a different confinement on plastic (their usual experimental condition) than on the sensing film. A loose (resp. tight) confinement may generate a higher (resp. lower) O_2 value under the colony. They also made colonies with a huge amount of cells (50,000 instead of 1,000 and 2,000 in our case). As the self-generated O_2 field depends on the consumption of every cell, we expect a huge degree of hypoxia. Finally, they never reached a stationary expansion regime due to the large initial excess of inner cells. That excess density slowly decreases with time as visible on their kymograph. We have actually simulated a moderate bulk cell excess in our work (Figure 4 of Cochet-Escartin et al.) which is also transiently visible at 3 h in **Figure 1B**. Such an inner cell mass transfer has to be taken into account to establish a correct mass balance equation, and the only $L(t)R(t)$ quantity tested by Biondo et al. is clearly not sufficient to draw a conclusion on cell divisions.

AUTHOR CONTRIBUTIONS

All authors listed have made a substantial, direct and intellectual contribution to the work, and approved it for publication.

FUNDING

This study was supported by the CNRS - Mission pour les Initiatives Transverses et Interdisciplinaires' "Défis Modélisation du vivant - 2019", by the European Research

Council (ERC) under the European Union's Horizon 2020 research and innovation program (grant agreement No 865711 to VC) and by the International Human Frontier Science Program Organization, Grant Number RGP0051/2021 (to J-PR).

REFERENCES

- Biondo, M., Panuzzo, C., Ali, S. M., Bozzaro, S., Osella, M., Bracco, E., et al. (2021). The Dynamics of Aerotaxis in a Simple Eukaryotic Model. *Front. Cel Dev. Biol.* 9, 720623. doi:10.3389/fcell.2021.720623
- Cochet-Escartin, O., Demircigil, M., Hirose, S., Allais, B., Gonzalo, P., Mikaelian, I., et al. (2021). Hypoxia Triggers Collective Aerotactic Migration in *Dictyostelium discoideum*. *eLife* 10, e64731. doi:10.7554/eLife.64731
- d'Alessandro, J., Mas, L., Aubry, L., Rieu, J. P., Rivière, C., and Anjard, C. (2018). Collective Regulation of Cell Motility Using an Accurate Density-Sensing System. *J. R. Soc. Interf.* 15, 45. doi:10.1098/rsif.2018.0006
- Schiavo, G., and Bisson, R. (1989). Oxygen Influences the Subunit Structure of Cytochrome C Oxidase in the Slime Mold *Dictyostelium discoideum*. *J. Biol. Chem.* 264, 7129–7134. doi:10.1016/s0021-9258(18)83211-2
- West, C. M., van der Wel, H., and Wang, Z. A. (2007). Prolyl 4-hydroxylase-1 Mediates O₂ Signaling during Development of *Dictyostelium*. *Development* 134, 3349–3358. doi:10.1242/dev.000893

Conflict of Interest: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2022 Rieu, Cochet-Escartin, Anjard, Demircigil and Calvez. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.