



HAL
open science

Flavoprotein Photochemistry: Fundamental Processes and Photocatalytic Perspectives

Bo Zhuang, Ursula Liebl, Marten H. Vos

► **To cite this version:**

Bo Zhuang, Ursula Liebl, Marten H. Vos. Flavoprotein Photochemistry: Fundamental Processes and Photocatalytic Perspectives. *Journal of Physical Chemistry B*, 2022, 126, pp.3199-3207. 10.1021/acs.jpcc.2c00969 . hal-03650652

HAL Id: hal-03650652

<https://hal.science/hal-03650652>

Submitted on 4 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Flavoprotein photochemistry: fundamental processes and photocatalytic perspectives

*Bo Zhuang, Ursula Liebl and Marten H. Vos**

LOB, CNRS, INSERM, Ecole Polytechnique, Institut Polytechnique de Paris, 91128 Palaiseau
Cedex, France

*Phone: +33169335066, marten.vos@polytechnique.edu


ABSTRACT

Flavins are highly versatile redox-active and colored cofactors in a large variety of proteins. These do include photoenzymes and photoreceptors, although the vast majority performs non-light-driven physiological functions. Nevertheless electron transfer between flavins and specific nearby amino acid residues (in particular tyrosine, tryptophan, and presumably histidine and arginine) takes place upon excitation of flavin in many flavoproteins. For oxidized flavoproteins these reactions potentially have a photoprotective role. In this Perspective, we outline work on the characterization of early reaction intermediates not only in the relatively well-studied resting oxidized forms, but also in the fully reduced and the intrinsically unstable semi-reduced forms, where ultrafast photooxidation of flavin was recently demonstrated. Along different lines, flavoprotein-based novel photocatalysts for biotechnological applications are presently emerging, employing both substrate photooxidation and photoreduction strategies. Deep insight into the fundamental flavin photochemical reactions may help in guiding and optimizing their development and in the exploration of novel photocatalytic approaches.

Introduction

Colored biomolecules are ubiquitous in natural systems. Some of them, such as chlorophyll and retinal, are exclusively employed in photobiological processes. Others, such as heme-proteins and many metal center-containing proteins, do perform photochemistry, although not for known photobiological functions, and their spectroscopic properties can be used to follow biochemical reactions. Flavins form a highly versatile group of chromophores found in a large variety of enzymes and photoreceptors. In most of these systems they do not perform photobiological functions, but in some systems they do. The versatility of these systems makes them interesting starting points for the bioengineering of novel photocatalysts. In this Perspective, we will outline recent progress in the understanding of fundamental photochemical reactions in flavoproteins, and discuss how this insight may allow the exploration of novel photocatalytic approaches.

Scheme 1. Molecular structure of flavins



Flavins (Fl) are derivatives of vitamin B₂ (riboflavin, Rfl) and are accommodated in proteins essentially in the form of flavin adenine dinucleotide (FAD) or flavin mononucleotide (FMN); see Scheme 1. The isoalloxazine ring system is at the heart of the reactivity of flavins and

of their spectroscopic properties. Flavins can adopt three different redox forms with various protonation states, leading to at least five physiologically relevant states: the oxidized form Fl_{ox} , the semi-reduced forms $\text{Fl}^{\bullet-}$ and FlH^{\bullet} and the fully reduced forms FlH^- and FlH_2 (Fig. 1A). Interacting with a manifold of substrates, as well as with other proteins, these properties allow flavoproteins to partake in a large variety of biochemical electron (one- and two-electron) and proton transfer reactions, including those that couple one- and two-electron reactions and bifurcation reactions¹⁻². The flavin color depends on the redox state (for the neutral states Fl_{ox} , FlH^{\bullet} and FlH_2 , respectively yellow, blue and virtually transparent in the visible). Hence the flavin absorption spectrum (with moderate extinction coefficients of up to $\sim 13,000 \text{ M}^{-1}\text{cm}^{-1}$ in the visible range, see Fig. 1B) is widely used to monitor its redox states during biochemical reactions, for instance in rapid mixing experiments with external reactants.




Fig. 1 A. Five redox and protonation states of flavins involved in the functional processes of flavoproteins. Flavin is represented by the isoalloxazine moiety. **B.** Steady-state absorption spectra of flavins in different redox and protonation states in glucose oxidase from *Aspergillus niger*. The spectrum of FADH[•] is reproduced from ref. ³, and normalized with respect to the spectrum of FAD_{ox}. Adapted with permission from Ref. ⁴. Copyright 2022 the Authors.

Photochemistry of oxidized flavoproteins and photoprotection

Flavin excited state formation by photon absorption can lead to fluorescence, and indeed fluorescence has been reported for most redox states, with the possible exception of the Fl⁻ state⁵. The best-studied state is the Fl_{ox} state, which is the resting state in the vast majority of flavoproteins. In solution, Rfl_{ox} and FMN_{ox} are highly fluorescent with lifetimes of the singlet excited state ¹Fl* in the range of a few nanoseconds; for FAD_{ox} there is a fraction ('open configuration') that is also highly fluorescent and a fraction in which the isoalloxazine ring is stacked with the adenine moiety ('closed configuration') leading to a lifetime of a few picoseconds⁶⁻⁸ (see also Fig. 4). In proteins ¹Fl fluorescence is very often strongly quenched (although flavin fluorescence can still be observed spectroscopically and in microscopy of living materials in a useful⁹ or perturbing¹⁰ way). In many cases this quenching is due to a flavin photoreduction process involving close-lying aromatic amino acids: the radical states Fl⁻X^{•+}, in which X is either tryptophan (TrpH) or tyrosine (TyrOH), lie energetically above the neutral ground state, but below the singlet excited state ¹Fl*. The driving force for tryptophan and tyrosine photo-oxidation is estimated at ~1.2 eV¹¹ and 0.7-1.0 eV¹²⁻¹³, respectively. The electron transfer (ET) reaction towards the flavin thus competes with the intrinsic ¹Fl* decay. The rate of such a reaction depends on the configuration of the donor-acceptor pair, which determines the electronic

coupling and the energetics involved¹⁴. Using a rough estimate based on distance and average protein density only¹⁵, direct electron donors located up to at most ~1 nm can efficiently compete, in the case of barrierless ET, with the nanosecond rates of intrinsic excited-state decay to the singlet or triplet ground state. In many cases tyrosines and/or tryptophanes are located much closer, up to near-van der Waals interaction with the flavin, and ET reactions with time constants down to the order of hundreds of femtoseconds regime have been documented¹⁶⁻²⁴. The rates of such ET reactions are a sensitive probe of the interaction between the flavin cofactor and the amino acid electron donor. This feature can be exploited to study configurational heterogeneity and exchange rates between configurations in the active site of flexible flavoenzymes^{13, 25-26}.

The presence of these intrinsic amino acid reactants incorporated in the protein thus prevents long-lived singlet flavin excited states (Fig. 1A). As long-lived (intrinsically microseconds) triplet states ^3Fl can be formed from $^1\text{Fl}^*$ by intersystem crossing on the timescale of ~10 ns²⁷⁻²⁸, and hence with quite sizeable quantum yields²⁹, in this way the build-up of this state is also impeded. Ground state molecular oxygen ($^3\text{O}_2$) can quench such states yielding $^1\text{O}_2$ ³⁰ and other reactive oxygen species (ROS)³¹. Therefore it has been put forward that the photoprotective effect of the presence of ET-quenchers of the $^1\text{Fl}^*$ state yields a strong selection pressure in the evolution of flavoenzymes³².




Fig. 2 **A.** Simplified scheme of photochemical reactions involving oxidized flavins. Excitation of the ^1Fl ground state leads to population of the $^1\text{Fl}^*$ state that can decay back to the ground state through fluorescence, to the triplet state $^3\text{Fl}^*$ by intersystem crossing, or to a charge separated radical pair state $\text{Fl}^- \text{X}^{++}$, in which X is an amino acid electron donor (tryptophan or tyrosine). The triplet state can decay back to the singlet ground state via interaction with molecular oxygen ($^3\text{O}_2$) yielding highly reactive singlet oxygen ($^1\text{O}_2$) and other ROS. **B, C, D and E.** Closest tryptophan or tyrosine residues in the flavin environment in four characteristic flavoproteins: the photoenzyme Fatty Acid Photodecarboxylase from *Chlorella variabilis*, PDB entry: 6ZH7 (B) and the non-photoenzyme glucose oxidase from *Aspergillus niger*, PDB entry: 1CF3 (C), photoreceptors LOV domain protein from *Rhodobacter sphaeroides*, PDB entry: 4HJ3 (D) and cryptochrome from *Drosophila melanogaster*, PDB entry: 4JZY (E). In the latter, Trp 420 and Trp 397 are the first two constituents of the Trp triad ET chain³³ (see text). Distances are indicated in Å.

In most flavoproteins, photoinduced radical states rapidly decay by charge recombination, typically on the picosecond timescale, thus eventually thermally dissipating the photon energy. These closed reaction cycles thus do not lead to long-lived photochemical products, but rather help avoiding them. On the other hand, it is interesting to note that there is also a limited class of flavoproteins that support functional photobiochemical reactions. In such systems, competitive

quenching processes could be detrimental. An interesting system is the recently discovered photoenzyme Fatty Acid Photodecarboxylase (FAP)³⁴. In FAP, a cascade of reactions is initiated by ET, in ~ 300 ps, from the fatty acid substrate to the excited flavin cofactor³⁵. ET from closeby tyrosine or tryptophan residues to the flavin would compete with this functional reaction. Inspection of the crystal structure (Fig. 1B) reveals that indeed the nearest aromatic residues are located over 8 Å (minimal ring-to-ring distance), much further than the carboxyl group of the substrate electron donor, and much further than the closest aromatic residues found typically in non-photoactive proteins (cf. glucose oxidase (GOX) structure, from the same GMC oxidase family of flavoenzymes as FAP, in Fig. 1C). Indeed, in FAP, replacement of the closest tyrosine residue, Tyr466, by ET-inert phenylalanine, does not enhance the activity³⁵. Interestingly, in the absence of substrate, the enzyme is quite unstable and sensitive to light. In this case $^3\text{FAD}^*$ is generated to a considerable extent by light absorption, which is likely, through ROS production, at the origin of the instability. We note that the instability has also been proposed to arise from harmful photooxidation of nearby residues³⁶, but this proposal is inconsistent with the sheer lack of close-by oxidizable residues described above, and the corresponding long-lived fluorescence in the absence of substrates.

In this context, another interesting flavoprotein performing a completely different functional photochemical process is the light-oxygen-voltage (LOV) domain from plant phototropin photosensors, which contains an FMN cofactor. Here, upon formation of the triplet state $^3\text{FMN}^*$ with high yield by intersystem crossing from the photo-generated singlet excited state $^1\text{FMN}^*$ in ~ 2 ns³⁷, a flavin-cysteine adduct is formed on the microsecond timescale³⁸ that is a precursor of more remote structural changes³⁹. Therefore, in this system the triplet state is part of the signaling pathway, and quenching of the singlet excited state by ET reactions competing with

decay to the triplet state would be unfavorable. Indeed, tyrosine and tryptophan residues are located at $> 9 \text{ \AA}$ from the isoalloxazine ring⁴⁰ (Fig. 1D) precluding quenching of the singlet state. It should be noted that in these systems under certain circumstances long-distance ET from tryptophan to the $^3\text{FMN}^*$ state can compete with intrinsic $^3\text{FMN}^*$ decay occurring on the microsecond timescale⁴⁰. Based on the reactivity of their photogenerated triplets, variants of LOV domains are employed as optogenetic tools⁴¹. When ET in the triplet states can in addition be sufficiently reduced, they can also be tailored for use as light-induced generators of singlet oxygen (singlet oxygen generator, SOG) and other ROS^{31, 42} using the mechanism described above.

As mentioned above, tryptophan and tyrosine are the only documented amino acid residues known to act as direct electron donor to photoexcited flavins. Interestingly, the involvement of tryptophan as electron donor was first explored not in a system with oxidized flavin, but in a photoenzyme that stabilizes the neutral radical FADH^\bullet state, DNA photolyase. In this protein, a chain of three tryptophan residues is located between the flavin cofactor and the solvent. Excitation of FADH^\bullet leads to a long-lived, and catalytically active, FADH^- . Here the initial ET step occurs in 30 ps and the primary $\text{FADH}^- \text{TrpH}^{+\bullet}$ photoproduct is stabilized by subsequent ultrafast hopping along the chain and deprotonation of the solvent-exposed tryptophan yielding $\text{FADH}^- \text{Trp}^{\bullet}$ ⁴³⁻⁴⁵. To disentangle these steps, fully spectrally and polarization resolved transient absorption experiments were exploited along with the fact that the tryptophan radical species (weakly) absorb in the visible range, and solution model spectra for them are available from pulse radiolysis experiments. Similar approaches have been used for the homologous class of cryptochrome blue light sensors (Fig. 1E), in which the resting state of the flavin is the FAD_{ox} state that has a higher-lying lowest excited state, allowing ~ 2 orders of magnitude faster ($\sim 400 \text{ fs}$) initial charge separation towards the $\text{FAD}^\bullet \text{TrpH}^{+\bullet}$ state^{16, 46}, and subsequent stabilization of the radical through the triple tryptophan

chain or variations thereof involving additional tryptophans or tyrosines. These stabilized radical pairs are thought to serve as intermediates in the signal-transmission chain for light adaptation in a wide variety of ways, the precise nature of the more down-stream events still remaining uncertain in many cryptochromes. There are also increasingly strong indications that cryptochromes in the retinal of migratory birds are involved in their orientation sensing through an (earth) magnetic field effect on the spin conversion yields based on light-induced radical pair formation⁴⁷. The latter possibility has inspired the development of *de novo* designed protein-riboflavin complexes in which the yield of neutral radicals involving flavin and tryptophan is magneto-sensitive, albeit at levels above the earth magnetic field⁴⁸. Recent ultrafast spectroscopic approaches indicate that in these systems, other than in cryptochromes, the charge separation generating the initial radical pairs occurs on a much slower timescale, and strongly involves the riboflavin triplet excited state¹¹.

The studies on native proteins discussed in the preceding paragraph made use of the distinct TrpH^{++} and Trp^{\bullet} spectral properties in the visible region (the latter is blue-shifted by ~ 60 nm) that have been well characterized in solution using pulse radiolysis and photolysis approaches. The $\text{p}K_a$ of TrpH^{++} is ~ 4 , hence at physiological pH this species will eventually deprotonate, but it can be long-lived enough (hundreds of nanoseconds in DNA photolyase⁴³) to be characterized as separate intermediate. For tyrosine radicals the situation is different. Whereas the TyrO^{\bullet} spectrum has been well characterized in solution, the TyrOH^{++} species had not been described, in general agreement with its estimated very low $\text{p}K_a$ (~ -2). The question whether this latter species can be generated as a separate intermediate was solved recently using the bacterial flavoproteins, methyl transferase TrmFO and ferredoxin-NAD(P)⁺ oxidoreductase (FNR, see Fig. 3), harboring a tyrosine close to the active site, where the primary photoproduct could be assigned to $\text{FAD}^{\bullet-}\text{TyrOH}^{++}$ based on its spectral properties, substantially shifted with respect to those of TyrO^{\bullet} both in the visible^{19, 21-22}

and in the infrared⁴⁹ spectral region. In these systems, the $\text{FAD}^{\cdot-}\text{TyrOH}^{\cdot+}$ state is formed on the timescale of ~ 100 fs – 1 ps and recombines within a few picoseconds. In GOX, where both tryptophan and tyrosine had previously been considered as primary electron donors^{12, 20, 50-51}, using the newly identified $\text{TyrOH}^{\cdot+}$ spectrum, $\text{FAD}^{\cdot-}\text{TyrOH}^{\cdot+}$ was also found to form in ~ 1 ps⁵². In competition with charge recombination and re-reduction of tyrosine by tryptophan, some deprotonation of the tyrosine cation radical by proton transfer to the flavin then takes place in a few picoseconds, yielding $\text{FADH}^{\cdot}\text{TyrO}^{\cdot}$. These studies thus imply that TyrO^{\cdot} , an important intermediate in a number of biochemical reactions⁵³, does not necessarily have to be formed by strictly concerted electron and proton transfer. Thus, whereas the photochemistry in these flavoproteins themselves has no catalytic role, dissection of the mechanism does allow shining light on potential mechanisms in systems where short-lived intermediates cannot be accumulated.




Fig. 3 Transient absorption spectra, measured 3 ps after excitation of FAD_{ox} , of wild-type (WT) and the Y50W and Y50G variants of FNR from *Bacillus subtilis*, corresponding to the difference spectra of $\text{FAD}^{\cdot-}\text{TyrOH}^{\cdot+}$, $\text{FAD}^{\cdot-}\text{TrpH}^{\cdot+}$, and $^1\text{FAD}^*$, respectively, with FAD_{ox} . The negative signals around 450 nm correspond to bleaching of the FAD_{ox} ground state. The positive signals above 500 nm are dominated by $\text{TyrOH}^{\cdot+}$, $\text{TrpH}^{\cdot+}$ and $^1\text{FAD}^*$, respectively, and the negative band around 560 nm in Y50G corresponds to stimulated emission from $^1\text{FAD}^*$. Reprinted with permission from Ref. ²². Copyright 2022 American Chemical Society.

As a side remark, these approaches are facilitated by the fact that in the visible range the extinction of the oxidized and semi-reduced flavin species (maxima $\sim 5,000$ – $13,000$ $\text{M}^{-1}\text{cm}^{-1}$, Fig.

1) is of the same order of magnitude as the one of the tryptophan and tyrosine radicals ($\sim 2,000$ – $15,000 \text{ M}^{-1}\text{cm}^{-1}$), as illustrated in Fig. 3. Studying such amino acid radicals generated by much stronger absorbing chromophore cofactors, such as chlorophylls or hemes (maxima up to $\sim 100,000$ and $200,000 \text{ M}^{-1}\text{cm}^{-1}$, respectively), would lead to small signals that risk being buried in much larger chromophore spectral changes. Thus flavoproteins, due to their moderate extinction in the visible range, are systems of choice to study such radicals by transient optical spectroscopy.

In the context of photoinduced ET to Fl_{ox}^* , it is interesting to note that in the flavoenzyme choline oxidase, a low-yield highly fluorescent species with an unusual fluorescence excitation spectrum has been assigned to a flavin-histidyl adduct⁵⁴, suggested to be formed in a light-induced way via photooxidation of the initially neutral, singly protonated close-lying histidine residue. Investigation of this hypothesis by time-resolved spectroscopic methods may shed light on such a mechanism, which should compete with cyclic photoinduced photooxidation of close-lying tyrosine and tryptophan residues occurring in $\sim 300 \text{ fs}$ ⁴. Interestingly, highly fluorescent flavin species cross-linked to the protein have also been reported in TrmFO⁵⁵⁻⁵⁶, and photo-transformation of such an adduct has been observed⁵⁶. Investigation of the photochemical processes occurring in complexes of flavins and protein residues is yet a virtually unexplored domain that may unearth new photocatalytic roles of flavins.




Fig. 4 Fluorescence decay on the picosecond timescale of free FAD_{ox} and FADH^- in solution (A) and of $\text{FAD}^{\bullet-}$, FAD_{ox} and FADH^- in GOX. For FAD_{ox} the decay on the timescale of a few picosecond is assigned ET to the isoalloxazine ring system, in solution from adenine to which it is stacked in a fraction of the flavins, in GOX from closely tyrosine. For FADH^- in solution the decay is assigned to a conical intersection with the ground state reached by conformational flexibility of the isoalloxazine ring system. This conical intersection is thought to be more difficult to reach in a constrained protein environment, as in GOX. Panel B is reprinted with permission from Ref. ⁴. Copyright 2022 the Authors.

Photochemistry of fully reduced flavoproteins

Photochemistry of flavin forms other than the oxidized resting state is interesting from a fundamental point of view, as well as for developing potential applications in using flavoprotein variants as photocatalysts (see below). Yet, studies of these forms remain scarce. As alluded to above, the excited-state and photoreduction properties of neutral semi-reduced radicals have been

studied mostly in the larger framework of the photoactivation of DNA photolyase^{43-45, 57-58}. Fully reduced (neutral or anionic) flavins have very long been known to be virtually non-fluorescent in solution, but can have long-lived emissive excited states when their conformational dynamics are restricted by the protein environment or low temperatures⁵⁹⁻⁶¹. This feature has been interpreted as a result of deactivation of the excited state via a conical intersection with the ground state that can be reached through butterfly bending motions of the isoalloxazine ring system. In confined environments such motions are hindered, leading to long-lived (up to nanoseconds) excited states. Such excited states are known to have a functional role in the photoenzyme DNA photolyase, where FADH^-* is the precursor of ET to the substrate, photodamaged DNA. The latter process occurs in ~ 200 ps, an order of magnitude faster than the intrinsic FADH^-* decay in the absence of substrate.⁴⁴ Photochemical processes of fully reduced flavins are also thought to play a role in bio-engineered photocatalysts (see below); therefore a basic understanding of the competing excited-state deactivation processes is warranted. For instance, using transient absorption spectroscopy, the decay of FADH^- in the methyl transferase flavoenzyme ThyX has been interpreted in terms of the rigidity of the active site⁶². As fluorescence represents the most direct measure of the excited state, we are currently engaged in studying excited-state decays of fully reduced flavins in well-characterized model protein systems, as well as in aqueous solution, using higher time resolution and more sensitive fluorescence detection systems than available in the classical studies. Fig. 4 gives an example comparing fluorescence decay of FADH^- in GOX and in solution. A promising perspective of a system that is worth further exploring in this regard is FNR. It has been demonstrated that in some FNR, the lowest electronic transition band of the fully reduced flavins is significantly red-shifted by ~ 40 nm compared with the commonly known absorption spectra of FADH^- or FADH_2 .⁶³ It also appears that in FNR, the presence of this red-shifted band correlates

with the reactivity of the fully reduced enzyme toward NADP^+ . Investigating the origin of this lower-lying electronic transition can provide useful insight into the electronic structures and surrounding protein environments of the fully reduced flavins in FNR, which may provide a better understanding of the function of this important class of flavoenzymes.

Photochemistry of semi-reduced flavoproteins

Semi-reduced flavin species play an important role as reaction intermediates in many biochemical reactions. Yet they are intrinsically quite unstable and can be accumulated for steady-state investigation only for a limited number of cases. The most emblematic example concerns the neutral semi-reduced FADH^\bullet reaction intermediate form, which is the resting state of the isolated DNA-repair photoenzyme DNA photolyase, although its catalytically competent form is FADH^- . As alluded to above, the excited-state and photoreduction properties of neutral semi-reduced radicals have been studied mostly in the larger framework of the photoactivation mechanism of this enzyme^{43-45, 57-58}, a second, substrate-independent, photoreaction that in DNA photolyase converts FADH^\bullet to FADH^- .




Fig. 5 Decay associated spectra (DAS) obtained from transient absorption measurements of $\text{FAD}^{\cdot-}$ in GOX upon excitation at 390 nm and 520 nm. The steady-state FAD_{ox} minus $\text{FAD}^{\cdot-}$ spectrum difference spectrum is shown (dashed line) for comparison (cf. Fig. 1B). Reprinted with permission from Ref. ⁴. Copyright 2022 the Authors.

The photophysical properties of the semi-reduced anionic radical species, $\text{Fl}^{\cdot-}$, have been even more scarcely studied^{46, 64}, and photochemical products have been reported only this year⁴. By photochemical or chemical reduction of Fl_{ox} under anaerobic conditions, $\text{Fl}^{\cdot-}$ can be stabilized to a substantial extent in some systems that harbor positively charged residues close to the flavin cofactors, in particular in flavoprotein oxidases where such residues may serve to maintain $\text{O}_2^{\cdot-}$ radical intermediates in the active site. Our recent study on a number of such oxidases shows that the $\text{Fl}^{\cdot-*}$ excited state is extremely short-lived (<100 fs, Fig. 4)⁴ to a point that time-resolved fluorescence could not be observed; previous reports on $\text{Fl}^{\cdot-*}$ fluorescence^{46, 64} may have been complicated by admixture of minor quantities of other fluorescent forms of the flavin. Remarkably, in all flavin oxidases (including the GOX model system) transient absorption spectroscopy witnessed that Fl_{ox} was formed quasi-instantaneously with high yield, demonstrating photooxidation of $\text{Fl}^{\cdot-}$ (Fig. 5).⁴ Thus, the inverse of the ubiquitous photoreduction reaction of Fl_{ox} also occurs (Fig. 6). The electron acceptor in these oxidases is likely to be a positively charged protonated residue, probably histidine or arginine, located in near-van der Waals distance from the flavin. The driving force for $\text{Fl}^{\cdot-*} \text{HisH}_2^+ \rightarrow \text{Fl}_{\text{ox}}\text{HisH}_2^{\cdot}$ ET was estimated at $\sim 0.2\text{eV}$ from theoretical considerations⁴. The mechanism of this ultrafast, yet presumably low-driving force, electron transfer reaction can likely not be described in terms of classical Marcus theory with the high activation energies expected with reorganization energies in the often assumed 0.7–1.0 eV range¹⁴⁻¹⁵. Future theoretical and experimental studies will be required to investigate whether for instance two-state strong coupling, near adiabatic transitions and/or direct charge transfer optical

transitions may play a role. Also, identification of systems that stabilize Fl^- in the absence of a nearby positively charged acceptor (at present these are not available) may allow spectral characterization of the $\text{Fl}^{\cdot-}$ precursor state.




Fig. 6 **A.** General reaction pathways for photoinduced ET in flavoproteins that include the proposed cycle of the photooxidation of Fl^- . X stands for the electron-donating residue (tryptophan or tyrosine) and Y^+ represents the electron-accepting residue (presumably histidine or arginine) in the protein active site. [Red] and [Ox] represent external reductants and oxidants. An exchange mechanism between reduced free flavins and protein-bound Fl_{ox} has also been proposed to explain the accumulation of protein-bound Fl^- under light illumination in the presence of hole scavenger⁶⁵, which is not included in the present scheme. **B.** Active site in the crystal structure of GOX from *Aspergillus niger* (PDB entry: 1CF3). The presence of electron-donating tyrosines and tryptophans, as well as electron-

accepting histidines in close proximity of the flavin ring allows both photoreduction of FAD_{ox} and photooxidation of $\text{FAD}^{\bullet-}$ to occur. Distances are indicated in Å.

In the five investigated systems in the aforementioned study⁴, the resultant photoproduct state (presumably $\text{Fl}_{\text{ox}}\text{HisH}_2^{\bullet}$ in GOX) fully reverts (or ‘re-separates’) to the initial dipolar state ($\text{Fl}^{\bullet-}\text{HisH}_2^+$ in GOX) where the radical is on the flavin moiety in 10–20 ps depending on the system. An interesting perspective from a more applied viewpoint would be the use of genetic engineering approaches for creating pathways in these systems that will allow hopping of the radical away from the flavin, thus stabilizing the photoproduct, and ultimately photoreduction of external reactants.

Photobiocatalysis perspectives

Flavoproteins play an important role in the strongly developing field of photobiocatalysis, where the light sensitivity of -often engineered- proteins is employed for catalyzing chemical reactions in a sustainable way.⁶⁶⁻⁶⁷ Indeed, in two of the only three known families of naturally occurring photoenzymes, the catalytic reaction is initiated by absorption of a photon by a flavin cofactor. These are DNA photolyase, where catalysis is initiated by light-induced electron transfer from fully reduced flavin to the pre-bound substrate⁴⁴, and FAP, where catalysis is initiated by light-induced electron transfer, in ~300 ps, from the pre-bound substrate to oxidized flavin³⁴. In particular for FAP, in the few years since its discovery in 2017, a strong world-wide activity has developed to explore variants with altered substrate-binding site residues produced by directed or semi-random mutagenesis strategies. In particular efforts have been made to adapt this enzyme to

catalyze decarboxylation of a variety of substrates, including molecules much smaller than the medium chain-length lipids of the native enzyme; furthermore decarboxylation at multiple positions of the same substrate was investigated⁶⁸. Herewith efforts have mainly been directed towards optimizing the substrate positioning in the active site. Yet another important issue is the low photostability of the enzyme in the absence of substrate. As outlined above, this feature is likely due to the lack of photoprotective photoreduction excited-state decay processes that many non-photoactive flavoenzymes display. To optimize practical applicability it may therefore be useful to also (co)-engineer intrinsic flavin photoreduction pathways, introducing tyrosine or tryptophan residues allowing ET to occur in the hundreds of picosecond to nanosecond range. In this way these new pathways would moderately compete with the catalytic pathway in the presence of substrate, allowing substrate-dependent balances between photocatalytic efficiencies and photostability.

Light-activated flavoenzymes that are not naturally light-driven are also being used as a starting point for developing photocatalysts for a variety of chemical reactions. In particular, ene-reductases such as Old Yellow Enzyme that harbor a versatile substrate-binding site have recently been explored to this aim by the Hyster group⁶⁹⁻⁷². In these systems, photoreduction of alternative substrates by one-electron transfer from excited fully reduced flavin was shown to initiate a variety of reactions, including stereoselective cyclization and reductive breaking of double bonds. In native and near-native enzyme systems the photochemical quantum yield of these reactions is often very low, possibly due to wasteful (from a catalytic point of view) early back reactions. Extensive screening of genetically modified variants has allowed the enhancement of yields up to the several-percent range⁷¹ (still well below the >50% quantum yields observed for native photoenzymes³⁴,

⁴⁴), at least in part by modifying the substrate-flavin interaction so as to optimize the competition between forward reactive reactions and back reactions.

In ene-reductases, photocatalytic reduction of acrylamides has been shown to occur from a state where the flavin is in a well-defined, fluorescent, fully-reduced FMNH⁻ form⁷² and the reaction is initiated analogously to substrate photoreduction in DNA photolyase. On the other hand, addition of substrates for photo-cyclization to the fully reduced enzyme leads to a strong absorption increase in the blue spectral range that is assigned to a charge-transfer transition allowing direct photoreduction of the substrate⁶⁹⁻⁷⁰. Intriguingly, the spectral properties of this band are very similar to the ones of the semi-reduced flavin anion FI⁻ (Fig. 1B) and the spectral properties of the initial photoproduct also show resemblance to those assigned to fully oxidized flavin in our recent photochemical study of flavoproteins containing semi-reduced flavin anion radicals⁴. Further theoretical and experimental investigations into such systems may help to better characterize the electronic properties of the starting and intermediate states and ultimately devise strategies to further improve their photocatalytic properties.

Photooxidation of substrates starting from oxidized flavoenzymes may also occur in non-native photoenzymes. In the dark, the bacterial monooxygenase PqsL does sustain activity with reduced 1-methylnicotinamide as a hydride donor, but not with NAD(P)H³². However, under light illumination the product can be formed with NAD(P)H. Based on time-resolved experiments under continuous illumination a scheme for this finding has been proposed that involves ET from this nicotinamide to FAD_{ox}* in two subsequent light-dependent steps. Each step yields FAD*⁻ that is subsequently rapidly transformed to FADH*, ultimately leading to full reduction of the flavin. Consistently, as the nearest tyrosine or tryptophan residue is located 7 Å from the flavin ring⁷³, ET from intra-protein donors is expected to occur on the timescale of tens of picoseconds at fastest¹⁵,

which leaves room for efficient competition of photoreduction by NAD(P)H. Further understanding and verification of the photoactivity of Pqsl, which may be extendible to other photocatalytic processes, can benefit from further characterization using high time resolution spectroscopic approaches.

Concluding remarks

Altogether, recent fundamental spectroscopic and modeling work has highlighted the extreme diversity of photochemical redox processes in flavoproteins. At the same time, a variety of applied approaches are emerging in which flavoenzymes are sought to perform ‘green chemistry’ photocatalytic processes that differ from their native physiological function, sometimes using pseudo-random optimization strategies. It is foreseen that insight in the fundamental reactions that underlie the photocatalytic reactions will guide their development along directed pathways, as well as allow novel photocatalytic approaches to be explored.

Acknowledgments.

This work was partly supported by ANR SNAPsHOTs to M.H.V. B.Z. thanks the China Scholarship Council for providing a Ph.D. scholarship.

References.

1. Buckel, W.; Thauer, R. K., Flavin-Based Electron Bifurcation, A New Mechanism of Biological Energy Coupling. *Chem. Rev.* **2018**, *118*, 3862-3886.

2. Duan, H. D.; Khan, S. A.; Miller, A.-F., Photogeneration and reactivity of flavin anionic semiquinone in a bifurcating electron transfer flavoprotein. *Biochim. Biophys. Acta* **2021**, *1862*, 148415.
3. Massey, V.; Palmer, G., On the existence of spectrally distinct classes of flavoprotein semiquinones. A new method for the quantitative production of flavoprotein semiquinones. *Biochemistry* **1966**, *5*, 3181–3189.
4. Zhuang, B.; Ramodiharilafy, R.; Liebl, U.; Aleksandrov, A.; Vos, M. H., Ultrafast Photooxidation of Protein-Bound Anionic Flavin Radicals. *Proc. Natl. Acad. Sci. U.S.A.* **2022**, *119*, e2118924119.
5. Su, D.; Kabir, M. P.; Orozco-Gonzalez, Y.; Gozem, S.; Gadda, G., Fluorescence Properties of Flavin Semiquinone Radicals in Nitronate Monooxygenase. *ChemBioChem* **2019**, *20*, 1646-1652.
6. Brazard, J.; Usman, A.; Lacomat, F.; Ley, C.; Martin, M. M.; Plaza, P., New Insights into the Ultrafast Photophysics of Oxidized and Reduced FAD in Solution. *J. Phys. Chem. A* **2011**, *115*, 3251-3262.
7. Stanley, R. J., Advances in flavin and flavoprotein optical spectroscopy. *Antioxid. Redox Signal.* **2001**, *3*, 847-866.
8. van den Berg, P. A. W.; Feenstra, K. A.; Mark, A. E.; Berendsen, H. J. C.; Visser, A. J. W. G., Dynamic Conformations of Flavin Adenine Dinucleotide: Simulated Molecular Dynamics of the Flavin Cofactor Related to the Time-Resolved Fluorescence Characteristics. *J. Phys. Chem. B* **2002**, *106*, 8858-8869.
9. Chen, A. X.; Conti, T. F.; Hom, G. L.; Greenlee, T. E.; Raimondi, R.; Briskin, I. N.; Rich, C. A.; Kampani, R.; Engel, R.; Sharma, S., et al., Functional imaging of mitochondria in retinal diseases using flavoprotein fluorescence. *Eye* **2021**, *35*, 74-92.
10. Engel, R.; Van Haastert, P. J. M.; Visser, A. J. W. G., Spectral characterization of *Dictyostelium* autofluorescence. *Micr. Res. Tech.* **2006**, *69*, 168-174.
11. Bialas, C.; Barnard, D. T.; Auman, D. B.; McBride, R. A.; Jarocho, L. E.; Hore, P. J.; Dutton, P. L.; Stanley, R. J.; Moser, C. C., Ultrafast flavin/tryptophan radical pair kinetics in a magnetically sensitive artificial protein. *Phys. Chem. Chem. Phys.* **2019**, *21*, 13453-13461.
12. Mataga, N.; Chosrowjan, H.; Shibata, Y.; Tanaka, F.; Nishina, Y.; Shiga, K., Dynamics and Mechanisms of Ultrafast Fluorescence Quenching Reactions of Flavin Chromophores in Protein Nanospace. *J. Phys. Chem. B* **2000**, *104*, 10667-10677.
13. Yang, H.; Luo, G.; Karnchanaphanurach, P.; Louie, T.-M.; Rech, I.; Cova, S.; Xun, L.; Xie, X. S., Protein Conformational Dynamics Probed by Single-Molecule Electron Transfer. *Science* **2003**, *302*, 262-266.
14. Marcus, R. A.; Sutin, N., Electron transfers in chemistry and biology. *Biochim. Biophys. Acta* **1985**, *811*, 265-322.
15. Page, C. C.; Moser, C. C.; Chen, X.; Dutton, P. L., Natural engineering principles of electron tunnelling in biological oxidation-reduction. *Nature* **1999**, *402*, 47-52.
16. Brazard, J.; Usman, A.; Lacomat, F.; Ley, C.; Martin, M. M.; Plaza, P.; Mony, L.; Heijde, M.; Zabulon, G.; Bowler, C., Spectro-Temporal Characterization of the Photoactivation Mechanism of Two New Oxidized Cryptochrome/Photolyase Photoreceptors. *J. Am. Chem. Soc.* **2010**, *132*, 4935-4945.
17. Chosrowjan, H.; Taniguchi, S.; Mataga, N.; Tanaka, F.; Todoroki, D.; Kitamura, M., Ultrafast fluorescence dynamics of FMN-binding protein from *Desulfovibrio vulgaris* (Miyazaki F) and its site-directed mutated proteins. *Chem. Phys. Lett.* **2008**, *462*, 121-124.

18. Chosrowjan, H.; Taniguchi, S.; Wongnate, T.; Sucharitakul, J.; Chaiyen, P.; Tanaka, F., Conformational heterogeneity in pyranose 2-oxidase from *Trametes multicolor* revealed by ultrafast fluorescence dynamics. *J. Photochem. Photobiol. A* **2012**, *234*, 44-48.
19. Dozova, N.; Lacombat, F.; Bou-Nader, C.; Hamdane, D.; Plaza, P., Ultrafast photoinduced flavin dynamics in the unusual active site of the tRNA methyltransferase TrmFO. *Phys. Chem. Chem. Phys.* **2019**, *21*, 8743-8756.
20. Mataga, N.; Chosrowjan, H.; Shibata, Y.; Tanaka, F., Ultrafast fluorescence quenching dynamics of flavin chromophores in protein nanospace. *J. Phys. Chem. B* **1998**, *102*, 7081-7084.
21. Nag, L.; Sournia, P.; Myllykallio, H.; Liebl, U.; Vos, M. H., Identification of the TyrOH⁺ Radical Cation in the Flavoenzyme TrmFO. *J. Am. Chem. Soc.* **2017**, *139*, 11500-11505.
22. Zhuang, B.; Seo, D.; Aleksandrov, A.; Vos, M. H., Characterization of Light-Induced Short-Lived Interacting Radicals in the Active Site of Flavoprotein Ferredoxin-NADP⁺ Oxidoreductase. *J. Am. Chem. Soc.* **2021**, *143* 2457-2768.
23. Staudt, H.; Oesterhelt, D.; Grininger, M.; Wachtveitl, J., Ultrafast Excited-state Deactivation of Flavins Bound to Dodecin. *J. Biol. Chem.* **2012**, *287*, 17637-17644.
24. Kudisch, B.; Oblinsky, D. G.; Black, M. J.; Zieleniewska, A.; Emmanuel, M. A.; Rumbles, G.; Hyster, T. K.; Scholes, G. D., Active-Site Environmental Factors Customize the Photophysics of Photoenzymatic Old Yellow Enzymes. *J. Phys. Chem. B* **2020**, *124*, 11236-11249.
25. Laptanok, S. P.; Bouzahir-Sima, L.; Lambry, J.-C.; Myllykallio, H.; Liebl, U.; Vos, M. H., Ultrafast real time visualization of the active site flexibility of the flavoenzyme thymidylate synthase ThyX. *Proc. Natl. Acad. Sci. U.S.A.* **2013**, *110*, 8924-8929.
26. Zhuang, B.; Nag, L.; Sournia, P.; Croitoru, A.; Ramodiharilafy, R.; Lambry, J.-C.; Myllykallio, H.; Aleksandrov, A.; Liebl, U.; Vos, M. H., Photochemical processes in flavoenzymes as a probe for active site dynamics: TrmFO of *Thermus thermophilus*. *Photochem. Photobiol. Sci.* **2021**, *20*, 663-670.
27. Sichula, V.; Kucheryavy, P.; Khatmullin, R.; Hu, Y.; Mirzakulova, E.; Vyas, S.; Manzer, S. F.; Hadad, C. M.; Glusac, K. D., Electronic Properties of N(5)-Ethyl Flavinium Ion. *J. Phys. Chem. A* **2010**, *114*, 12138-12147.
28. Li, G.; Glusac, K. D., Light-Triggered Proton and Electron Transfer in Flavin Cofactors. *J. Phys. Chem. A* **2008**, *112*, 4573-4583.
29. Ahmad, I.; Vaid, F. H. M., Photochemistry of Flavins in Aqueous and Organic Solvents. In *Flavins: Photochemistry and Photobiology*, Silva, E.; Edwards, A. M., Eds. The Royal Society of Chemistry: 2006; Vol. 6, pp 13-40.
30. Baier, J.; Maisch, T.; Maier, M.; Engel, E.; Landthaler, M.; Baumler, W., Singlet Oxygen Generation by UVA Light Exposure of Endogenous Photosensitizers. *Biophys. J.* **2006**, *91*, 1452-1459.
31. Barnett, M. E.; Baran, T. M.; Foster, T. H.; Wojtovich, A. P., Quantification of light-induced miniSOG superoxide production using the selective marker, 2-hydroxyethidium. *Free Radic. Biol. Med.* **2018**, *116*, 134-140.
32. Ernst, S.; Rovida, S.; Mattevi, A.; Fetzner, S.; Drees, S. L., Photoinduced monooxygenation involving NAD(P)H-FAD sequential single-electron transfer. *Nature Comm.* **2020**, *11*, 2600.

33. Ozturk, N.; Selby, C. P.; Zhong, D. P.; Sancar, A., Mechanism of Photosignaling by *Drosophila* Cryptochrome ROLE OF THE REDOX STATUS OF THE FLAVIN CHROMOPHORE*. *J. Biol. Chem.* **2014**, *289*, 4634-4642.
34. Sorigué, D.; Légeret, B.; Cuiné, S.; Blangy, S.; Moulin, S.; Billon, E.; Richaud, P.; Brugière, S.; Couté, Y.; Nurizzo, D., et al., An algal photoenzyme converts fatty acids to hydrocarbons. *Science* **2017**, *357*, 903-907.
35. Sorigué, D.; Hadjidemetriou, K.; Blangy, S.; Gotthard, G.; Bonvalet, A.; Coquelle, N.; Samire, P.; Aleksandrov, A.; Antonucci, L.; Benachir, A., et al., Mechanism and dynamics of fatty acid photodecarboxylase. *Science* **2021**, *372*, eabd5687.
36. Wu, Y.; Paul, C. E.; Hollmann, F., Stabilisation of the Fatty Acid Decarboxylase from *Chlorella variabilis* by Caprylic Acid. *ChemBioChem* **2021**, *22*, 2420-2423.
37. Kennis, J. T. M.; Crosson, S.; Gauden, M.; van Stokkum, I. H. M.; Moffat, K.; van Grondelle, R., Primary Reactions of the LOV2 Domain of Phototropin, a Plant Blue-Light Photoreceptor. *Biochemistry* **2003**, *42*, 3385-3392.
38. Swartz, T. E.; Corchnoy, S. B.; Christie, J. M.; Lewis, J. W.; Szundi, I.; Briggs, W. R.; Bogomolni, R. A., The Photocycle of a Flavin-binding Domain of the Blue Light Photoreceptor Phototropin. *J. Biol. Chem.* **2001**, *276*, 36493-36500.
39. Iuliano, J. N.; Collado, J. T.; Gil, A. A.; Ravindran, P. T.; Lukacs, A.; Shin, S.; Woroniecka, H. A.; Adamczyk, K.; Aramini, J. M.; Edupuganti, U. R., et al., Unraveling the Mechanism of a LOV Domain Optogenetic Sensor: A Glutamine Lever Induces Unfolding of the α Helix. *ACS Chem. Biol.* **2020**, *15*, 2752-2765.
40. Kopka, B.; Magerl, K.; Savitsky, A.; Davari, M. D.; Röllen, K.; Bocola, M.; Dick, B.; Schwaneberg, U.; Jaeger, K.-E.; Krauss, U., Electron transfer pathways in a light, oxygen, voltage (LOV) protein devoid of the photoactive cysteine. *Sci. Rep.* **2017**, *7*, 13346.
41. Christie, J. M.; Gawthorne, J.; Young, G.; Fraser, N. J.; Roe, A. J., LOV to BLUF: Flavoprotein Contributions to the Optogenetic Toolkit. *Molecular Plant* **2012**, *5*, 533-544.
42. Torra, J.; Lafaye, C.; Signor, L.; Aumonier, S.; Flors, C.; Shu, X.; Nonell, S.; Gotthard, G.; Royant, A., Tailing miniSOG: structural bases of the complex photophysics of a flavin-binding singlet oxygen photosensitizing protein. *Sci. Rep.* **2019**, *9*, 2428.
43. Aubert, C.; Vos, M. H.; Mathis, P.; Eker, A. P. M.; Brettel, K., Intra-protein radical transfer during photoactivation of DNA photolyase. *Nature* **2000**, *405*, 586-590.
44. Brettel, K.; Byrdin, M.; Vos, M. H., Ultrafast light-induced processes in DNA-photolyase and its substrate-bound complex. In *Ultrafast Dynamics at the Nanoscale: Biomolecules and Supramolecular Assemblies*, Haacke, S.; Burghardt, I., Eds. Pan Stanford: Singapore, 2016; pp 65-90.
45. Lukacs, A.; Eker, A. P. M.; Byrdin, M.; Brettel, K.; Vos, M. H., Electron hopping through the 15 Å triple tryptophan molecular wire in DNA photolyase occurs within 30 ps. *J. Am. Chem. Soc.* **2008**, *130*, 14394-14395.
46. Kao, Y.-T.; Tan, C.; Song, S.-H.; Öztürk, N.; Li, J.; Wang, L.; Sancar, A.; Zhong, D., Ultrafast Dynamics and Anionic Active States of the Flavin Cofactor in Cryptochrome and Photolyase. *J. Am. Chem. Soc.* **2008**, *130*, 7695-7701.
47. Xu, J.; Jarocho, L. E.; Zollitsch, T.; Konowalczyk, M.; Henbest, K. B.; Richert, S.; Golesworthy, M. J.; Schmidt, J.; Déjean, V.; Sowood, D. J. C., et al., Magnetic sensitivity of cryptochrome 4 from a migratory songbird. *Nature* **2021**, *594*, 535-540.
48. Bialas, C.; Jarocho, L. E.; Henbest, K. B.; Zollitsch, T. M.; Kodali, G.; Timmel, C. R.; Mackenzie, S. R.; Dutton, P. L.; Moser, C. C.; Hore, P. J., Engineering an Artificial Flavoprotein Magnetosensor. *Journal of the American Chemical Society* **2016**, *138*, 16584-16587.

49. Pirisi, K.; Nag, L.; Fekete, Z.; Iuliano, J. N.; Tollentino Collado, J.; Clark, I. P.; Pécsi, I.; Sournia, P.; Liebl, U.; Greetham, G. M., et al., Identification of the vibrational marker of tyrosine cation radical using ultrafast transient infrared spectroscopy of flavoprotein systems. *Photochem. Photobiol. Sci.* **2021**, *20*, 369–378.
50. Fujiwara, A.; Mizutani, Y., Photoinduced electron transfer in glucose oxidase: a picosecond time-resolved ultraviolet resonance Raman study. *J. Raman Spectrosc.* **2008**, *39*, 1600-1605.
51. Zhong, D.; Zewail, A. H., Femtosecond dynamics of flavoproteins: charge separation and recombination in riboflavin (vitamin B2)-binding protein and in glucose oxidase enzyme. *Proc. Natl. Acad. Sci. U.S.A.* **2001**, *98*, 11867-11872.
52. Nag, L.; Lukacs, A.; Vos, M. H., Short-Lived Radical Intermediates in the Photochemistry of Glucose Oxidase. *ChemPhysChem* **2019**, *20*, 1793-1798.
53. Winkler, J. R.; Gray, H. B., Long-Range Electron Tunneling. *J. Am. Chem. Soc.* **2014**, *136*, 2930-2939.
54. Su, D.; Smitherman, C.; Gadda, G., A Metastable Photoinduced Protein–Flavin Adduct in Choline Oxidase, an Enzyme Not Involved in Light-Dependent Processes. *J. Phys. Chem. B* **2020**, *124*, 3936-3943.
55. Hamdane, D.; Guérineau, V.; Un, S.; Golinelli-Pimpaneau, B., A Catalytic Intermediate and Several Flavin Redox States Stabilized by Folate-Dependent tRNA Methyltransferase from *Bacillus subtilis*. *Biochemistry* **2011**, *50*, 5208-5219.
56. Sournia, P. La méthylation flavine-dépendante d'acides nucléiques : aspects évolutifs, métaboliques, biochimiques et spectroscopiques. PhD Thesis, Ecole Polytechnique, 2016.
57. Pan, J.; Byrdin, M.; Aubert, C.; Eker, A. P. M.; Brettel, K.; Vos, M. H., Excited-state properties of flavin radicals in flavoproteins: femtosecond spectroscopy of DNA photolyase, glucose oxidase and flavodoxin. *J. Phys. Chem. B* **2004**, *108*, 10160-10167.
58. Wang, H.; Saxena, C.; Quan, D.; Sancar, A.; Zhong, D., Femtosecond Dynamics of Flavin Cofactor in DNA Photolyase: Radical Reduction, Local Solvation, and Charge Recombination. *J. Phys. Chem. B* **2005**, *109*, 1329-1333.
59. Enescu, M.; Lindqvist, L.; Soep, B., Excited-state dynamics of fully reduced flavins and flavoenzymes studied at subpicosecond time resolution. *Photochem. Photobiol.* **1998**, *68*, 150-6.
60. Visser, A. J. W. G.; Ghisla, S.; Massey, V.; Müller, F.; Veeger, C., Fluorescence Properties of Reduced Flavins and Flavoproteins. *Eur. J. Biochem.* **1979**, *101*, 13-21.
61. Kao, Y.-T.; Saxena, C.; He, T.-F.; Guo, L.; Wang, L.; Sancar, A.; Zhong, D., Ultrafast Dynamics of Flavins in Five Redox States. *J. Am. Chem. Soc.* **2008**, *130*, 13132-13139.
62. Dozova, N.; Lacomat, F.; Lombard, M.; Hamdane, D.; Plaza, P., Ultrafast dynamics of fully reduced flavin in catalytic structures of thymidylate synthase ThyX. *Phys. Chem. Chem. Phys.* **2021**, *23*, 22692-22702.
63. Seo, D.; Muraki, N.; Kurisu, G., Kinetic and structural insight into a role of the re-face Tyr328 residue of the homodimer type ferredoxin-NADP⁺ oxidoreductase from *Rhodospseudomonas palustris* in the reaction with NADP⁺/NADPH. *Biochim. Biophys. Acta* **2020**, *1861*, 148140.
64. Ghanem, M.; Fan, F.; Francis, K.; Gadda, G., Spectroscopic and Kinetic Properties of Recombinant Choline Oxidase from *Arthrobacter globiformis*. *Biochemistry* **2003**, *42*, 15179-15188.
65. Massey, V.; Stankovich, M.; Hemmerich, P., Light-mediated reduction of flavoproteins with flavins as catalysts. *Biochemistry* **1978**, *17*, 1-8.

66. Meyer, L.-E.; Eser, B. E.; Kara, S., Coupling light with biocatalysis for sustainable synthesis—very recent developments and future perspectives. *Curr. Opin. Green Sust. Chem.* **2021**, *31*, 100496.
67. Schmermund, L.; Jurkaš, V.; Özgen, F. F.; Barone, G. D.; Büchenschütz, H. C.; Winkler, C. K.; Schmidt, S.; Kourist, R.; Kroutil, W., Photo-Biocatalysis: Biotransformations in the Presence of Light. *ACS Catal.* **2019**, *9*, 4115-4144.
68. Hedison, T. M.; Heyes, D. J.; Scrutton, N. S., Making molecules with photodecarboxylases: A great start or a false dawn? *Curr. Res. Chem. Biol.* **2022**, *2*, 100017.
69. Biegasiewicz, K. F.; Cooper, S. J.; Gao, X.; Oblinsky, D. G.; Kim, J. H.; Garfinkle, S. E.; Joyce, L. A.; Sandoval, B. A.; Scholes, G. D.; Hyster, T. K., Photoexcitation of flavoenzymes enables a stereoselective radical cyclization. *Science* **2019**, *364*, 1166-1169.
70. Gao, X.; Turek-Herman, J. R.; Choi, Y. J.; Cohen, R. D.; Hyster, T. K., Photoenzymatic Synthesis of α -Tertiary Amines by Engineered Flavin-Dependent “Ene”-Reductases. *J. Am. Chem. Soc.* **2021**, *143*, 19643-19647.
71. Nicholls, B. T.; Oblinsky, D. G.; Kurtoic, S. I.; Grosheva, D.; Ye, Y.; Scholes, G. D.; Hyster, T. K., Engineering a Non-Natural Photoenzyme for Improved Photon Efficiency. *Angew. Chem. Int. Ed.* **2022**, *61*, e202113842.
72. Sandoval, B. A.; Clayman, P. D.; Oblinsky, D. G.; Oh, S.; Nakano, Y.; Bird, M.; Scholes, G. D.; Hyster, T. K., Photoenzymatic Reductions Enabled by Direct Excitation of Flavin-Dependent “Ene”-Reductases. *J. Am. Chem. Soc.* **2021**, *143*, 1735-1739.
73. Drees, S. L.; Ernst, S.; Belviso, B. D.; Jagmann, N.; Hennecke, U.; Fetzner, S., PqsL uses reduced flavin to produce 2-hydroxylaminobenzoylacetate, a preferred PqsBC substrate in alkyl quinolone biosynthesis in *Pseudomonas aeruginosa*. *J. Biol. Chem.* **2018**, *293*, 9345-9357.

Biographies

Bo Zhuang obtained his Bachelor's degree in Material Chemistry from Southwest University, China, and Master's degree in Applied Chemistry from Osaka University, Japan. He is currently working toward his PhD in Biophysics at Ecole Polytechnique. His PhD thesis is mainly focusing on the photochemistry of protein-bound flavin species, using experimental and computational techniques to reveal the underlying mechanisms.

Ursula Liebl received her PhD in Biology from the University of Regensburg, Germany, in 1993. After post-doctoral stays at the University of Pennsylvania, Philadelphia, USA and in France, she joined the faculty of Biology of the Ecole Polytechnique, before becoming CNRS staff scientist. One focus of her current research interests concerns deciphering microbial methylation processes involving flavo-enzymes, using molecular biological and biochemical approaches.

Marten Vos received his PhD in Biophysics from the University of Leiden, the Netherlands, in 1990. After postdoctoral work at CEA Saclay and Ecole Polytechnique-ENSTA, Palaiseau he became a CNRS staff scientist working subsequently at the Laboratory for Applied Optics and the Laboratory for Optics and Biosciences at Ecole Polytechnique in Palaiseau. His research interest mainly concerns the application of optical spectroscopic techniques to understand ultrafast processes in proteins.

TOC graphics

