



HAL
open science

Gadd45g is required for timely Sry expression independently of RSPO1 activity

Nick Warr, Pam Siggers, Joel May, Nicolas Chalon, Madeleine Pope, Sara Wells, Marie-Christine Chaboissier, Andy Greenfield

► **To cite this version:**

Nick Warr, Pam Siggers, Joel May, Nicolas Chalon, Madeleine Pope, et al.. Gadd45g is required for timely Sry expression independently of RSPO1 activity. *Reproduction* [Cambridge], 2022, 163, pp.333-340. 10.1530/REP-21-0443 . hal-03650971

HAL Id: hal-03650971

<https://hal.science/hal-03650971v1>

Submitted on 25 Apr 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Gadd45g is required for timely *Sry* expression independently of RSPO1 activity

Nick Warr¹, Pam Siggers¹, Joel May^{1,†}, Nicolas Chalon^{1,‡}, Madeleine Pope¹, Sara Wells², Marie-Christine Chaboissier³ and Andy Greenfield^{1,§}

¹Mammalian Genetics Unit, Medical Research Council, Harwell Institute, Oxfordshire, UK, ²The Mary Lyon Centre, Medical Research Council, Harwell Institute, Oxfordshire, UK and ³Université Côte d'Azur, Inserm, CNRS, Institut de Biologie Valrose (iBV), Nice, France

Correspondence should be addressed to A Greenfield; Email: andy.greenfield@wrh.ox.ac.uk

†(J May is now at Center For Advanced Molecular Diagnostics, Brigham and Women's Hospital, Boston, Massachusetts, USA)

‡(N Chalon is now at de Duve Institute, Université Catholique de Louvain, Brussels, Belgium)

§(A Greenfield is now at Nuffield Department of Women's & Reproductive Health, Institute of Reproductive Sciences, University of Oxford, Oxford, UK)

Abstract

Sex determination in mammals is controlled by the dominance of either pro-testis (SRY-SOX9-FGF9) or pro-ovary (RSPO1-WNT4-FOXL2) genetic pathways during early gonad development in XY and XX embryos, respectively. We have previously shown that early, robust expression of mouse *Sry* is dependent on the nuclear protein GADD45g. In the absence of GADD45g, XY gonadal sex reversal occurs, associated with a major reduction of *Sry* levels at 11.5 dpc. Here, we probe the relationship between *Gadd45g* and *Sry* further, using gain- and loss-of-function genetics. First, we show that transgenic *Gadd45g* overexpression can elevate *Sry* expression levels at 11.5 dpc in the B6.Y^{POS} model of sex reversal, resulting in phenotypic rescue. We then show that the zygosity of pro-ovarian *Rspo1* is critical for the degree of gonadal sex reversal observed in both B6.Y^{POS} and *Gadd45g*-deficient XY gonads, in contrast to that of *Foxl2*. Phenotypic rescue of sex reversal is observed in XY gonads lacking both *Gadd45g* and *Rspo1*, but this is not associated with rescue of *Sry* expression levels at 11.5 dpc. Instead, *Sox9* levels are rescued by around 12.5 dpc. We conclude that *Gadd45g* is absolutely required for timely expression of *Sry* in XY gonads, independently of RSPO1-mediated WNT signalling, and discuss these data in light of our understanding of antagonistic interactions between the pro-testis and pro-ovary pathways.

Reproduction (2022) 163 333–340

Introduction

Mammalian sexual development involves the commitment of the bipotential gonadal primordium to a testicular or ovarian fate (Greenfield 2013). In mice, testis determination is dependent on the timely expression of the Y-linked gene *Sry* in supporting cell precursors of the developing XY gonad (Bullejos & Koopman 2001). SRY protein, a transcription factor, acts through regulatory elements of the *Sox9* gene in order to effect its up-regulation (Sekido & Lovell-Badge 2008, Gonen *et al.* 2017, 2018). Subsequently, SOX9, another transcription factor, binds target genes in order to direct the differentiation of Sertoli cells, which orchestrate morphogenetic processes required for testis development (Cool *et al.* 2012, Rahmoun *et al.* 2017). In XX gonads, the absence of SRY leads to distinct pro-ovarian pathways of gene expression gaining dominance; these comprise canonical WNT signalling,

especially RSPO1- and WNT4-dependent signals, and the activity of the pioneer transcription factor FOXL2 (Pannetier *et al.* 2016).

The establishment of such sexually dimorphic pathways of gene expression is dependent on the mutually antagonistic interactions between them in early gonad development (Greenfield 2015). Any significant reduction or delay in *Sry* expression during a critical phase of testis determination can result in XY gonadal sex reversal (Hiramatsu *et al.* 2009, Carre *et al.* 2018), due to a failure of such inhibition. Moreover, mutually antagonistic molecular interactions exist at the heart of the sex-determining mechanism: the ovarian determinant RSPO1 inhibits the activity of the testis-determining factor ZNRF3, which itself inhibits canonical WNT signalling required for ovary development (Harris *et al.* 2018).

A number of gene mutations have been reported that disrupt *Sry* regulators and these can cause testis



determination phenotypes (reviewed in Carre & Greenfield 2014, Larney *et al.* 2014, Nef *et al.* 2019). While such disrupted genes were assumed to act directly on *Sry* expression in a positive manner, it has recently been shown that one inhibitor of the pro-ovarian canonical WNT signalling pathway is required for the establishment of robust *Sry* expression (Garcia-Moreno *et al.* 2019). CBX2, a component of the polycomb repressive complex1 that binds H3K27me3 to mediate gene silencing, is a known factor required for normal *Sry* expression (Kato-Fukui *et al.* 2012). However, *Sry* expression is re-established in fetal XY gonads of embryos lacking both CBX2 and WNT4, suggesting that CBX2 acts primarily in a negative fashion to oppose the influence of WNT4/beta-catenin (Garcia-Moreno *et al.* 2019). Moreover, these data suggest that elimination of pro-ovarian WNT4/beta-catenin signals in the early XY gonad is a prerequisite of establishing a molecular environment in which early, robust *Sry* expression can occur.

We have previously reported a series of loss- and gain-of-function studies that reveal a role for the mitogen-activated protein kinase (MAPK) signalling pathway in testis determination via its effect on *Sry* expression. Loss of the kinase MAP3K4, or its regulator GADD45g, results in embryonic XY gonadal sex reversal associated with reduced *Sry* expression around 11.5 dpc on the C57BL/6J (B6/J) mouse strain (Bogani *et al.* 2009, Warr *et al.* 2012). Spatiotemporal profiling suggests that *Sry* expression eventually recovers to near-normal levels in both *Map3k4*- and *Gadd45g*-deficient XY gonads, but too late to affect the developmental fate of gonadal somatic cells, which acquire an ovarian fate as a consequence (Warr *et al.* 2012). Moreover, overexpression of *Map3k4* from a functional BAC transgene has been shown to rescue an *Sry* deficit in the classical mouse sex reversal mutation *T*-associated sex reversal (*Tas*) by re-establishing robust levels of *Sry* (Warr *et al.* 2012, 2014).

Here, we extend our analysis of the role of GADD45g in sex determination by probing its relationship with *Sry* further, using loss- and gain-of-function mouse lines. Using a previously reported BAC transgene (Warr *et al.* 2018), we show that transgenic overexpression of *Gadd45g* rescues the B6.Y^{POS} model of XY gonadal sex reversal by promoting *Sry*^{POS} expression. Exploiting this transgenic rescue, we then generated B6.Y^{POS} fetuses that lack one copy of either *Rspo1* or *Foxl2* in order to assess the relative contribution of these two main pro-ovarian genes to XY fetal ovary development. B6.Y^{POS}, *Rspo1*^{-/+} heterozygous fetuses at 14.5 dpc exhibit substantial rescue of gonadal sex reversal defects, with testicular tissue forming throughout much of the gonad. By contrast, B6.Y^{POS}, *Foxl2*^{-/+} heterozygotes, as well as homozygotes, show only minimal evidence of testicular tissue development. We then assessed whether loss of *Rspo1* operates to rescue sex reversal by rescuing *Sry*

expression. We show that gonads in XY fetuses lacking both *Gadd45g* and *Rspo1* develop as ovotestes with substantial testicular tissue formation at 14.5 dpc. However, this rescue of gonadal sex reversal in the double knockout is not associated with any rescue of the levels of *Sry* expression at 11.5 dpc, with only minimal *Sry* expression detected, as in XY *Gadd45g*-deficient gonads. We conclude that GADD45g is absolutely required for timely *Sry* expression independently of RSPO1 activity and that, therefore, loss of RSPO1 promotes testis determination in the absence of GADD45g through positive impacts downstream of initiation of timely *Sry* expression.

Materials and methods

Mouse strains and breeding

Breeding of B6.Y^{POS} mice has been previously described (Livermore *et al.* 2020). BAC transgenic lines harbouring functional *Map3k4* and *Gadd45g* have been previously described (Warr *et al.* 2014, 2018). *Gadd45g* knockout (KO) mice, *Foxl2* KO mice (a kind gift from Robin Lovell-Badge and Mathias Treier) and *Rspo1* KO mice have also previously been described (Hoffmeyer *et al.* 2001, Schmidt *et al.* 2004, Chassot *et al.* 2008). Mice were bred and maintained on the C57BL/6J background. All animal experiments were approved by the Animal Welfare and Ethical Review Body at MRC Harwell. Mice used were bred under a license from the UK Home Office (PPL 70/8898). Mice were housed in individually ventilated cages in a specific pathogen-free environment. Further details of micro- and macroenvironmental conditions are available on request.

Generation of embryos

Noon on the day of the copulatory plug was counted as 0.5 days *post coitum* (dpc). Embryos collected at 11.5dpc were staged accurately based on the number of tail somites (ts).

Whole-mount *in situ* hybridization

Whole-mount *in situ* hybridization (WMISH) analysis of embryonic tissues was performed as previously described (Grimmond *et al.* 2000, Cox *et al.* 2006). Probes for *Sox9* (Wright *et al.* 1995) and *Stra8* (Bogani *et al.* 2009, Warr *et al.* 2012), *Sry* (Bullejos & Koopman 2001) and *3b-Hsd* (Siggers *et al.* 2002) have been previously described.

Immunostaining

Gonads were fixed overnight in 4% PFA and processed for wax sectioning. Sections were dewaxed in xylene and antigen retrieval was achieved using Declere solution (Cell Marque, Rocklin, CA, USA). The following primary (1:100) and secondary (1:200) antibodies were used: anti-DDX (Abcam ab13840) and anti-AMH (Santa Cruz sc6886), donkey-anti-rabbit Alexa Fluor 594 (Invitrogen A21207) and donkey-

anti-goat Alexa Fluor 488 (Invitrogen A11055). Images were captured using a Zeiss LSM 710 upright confocal microscope and Zen software.

Quantitative reverse transcription-PCR

RNA was extracted using RNAeasy micro kit (Qiagen, 74004) according to the manufacturer’s instructions. cDNA was synthesized from approximately 100 ng total RNA (one pair of sub-dissected gonads) using random primers (high capacity cDNA RT kit, Applied Biosystems, 4368814). Relative cDNA levels were analysed by quantitative PCR using an Applied Biosystems 7500 real-time PCR machine. The TaqMan-MGB-FAM gene expression assays (Applied Biosystems) used were: *Rps29* (Mm02342448_gH) and *Sry* (Mm00441712_s1).

Results

We have shown previously that a bacterial artificial chromosome (BAC) encoding *Gadd45g* is capable of rescuing sex determination defects in a *Gadd45g* knockout, indicating its functionality (Warr et al. 2018). We first sought to assess the impact of *Gadd45g* overexpression in a classical model of XY gonadal sex reversal, the B6.Y^{POS} mouse (Eicher et al. 1982), by crossing the *Gadd45g* BAC transgene onto this strain. Testis determination defects in B6.Y^{POS} are associated with dysregulation of *Sry* expression (Bullejos & Koopman 2005, Livermore et al. 2020), presumably due to some functional mismatch between the *Sry*^{POS} allele and autosomal/X-linked gene variants in B6/J mice. We recently reported an autosomal modifier that protects against B6.Y^{POS} sex reversal (Livermore et al. 2020): all experiments described here used mice that lack this modifier. We first examined B6.Y^{POS} transgenic gonads at 14.5 dpc using WMISH with *Sox9* and *Stra8* (Fig. 1). Control B6.Y^{POS} gonads were most commonly ovaries that lacked the Sertoli cell marker *Sox9* (Fig. 1A). However, all transgenic gonads had a testicular morphology, with testis cords extending almost to the poles of the gonad, like B6.Y^{B6} controls. Some polar expressions of *Stra8*, a marker of germ cell meiotic entry that is strongly expressed in XX and B6.Y^{POS} controls, in transgenic gonads indicates that rescue of sex reversal is not absolutely complete (Fig. 1B). However, this does not prevent the development of fertile males in B6.Y^{POS}, *Gadd45g* BAC transgenic mice. Such males were routinely bred to B6 females to generate breeding stock and embryos for experimentation. When a functional BAC transgene encoding the kinase *Map3k4* (Warr et al. 2014), which interacts with *Gadd45g* in testis determination (Warr et al. 2012), was bred onto B6.Y^{POS}, the same rescue of fetal gonadal sex reversal was observed (Supplementary Fig. 1, see section on supplementary materials given at the end of this article). But our focus here is the role of *Gadd45g*.

We then carefully examined the expression of *Sry* and *Sox9* at 11.5 dpc, the sex-determining stage of




Figure 1 Rescue of fetal B6.Y^{POS} gonadal sex reversal by BAC transgenic overexpression of *Gadd45g* through positive impact on *Sry*. Whole-mount *in situ* hybridisation (WMISH) of XX wild-type, XY (B6.Y^{B6}) wildtype, B6.Y^{POS} and transgenic B6.Y^{POS} gonads (left to right) at 14.5 dpc with *Sox9* (A) and *Stra8* (B) probes. WMISH analysis of fetal gonads with the same genotypes at 11.5 dpc (16 ts, *Sry*, C) and *Sox9* (19ts, D). Quantitative reverse transcription-PCR analysis of *Sry* expression (E) confirms WMISH data, with significant reduction in *Sry* levels in B6.XY^{POS} (compared to B6.XY^{B6} controls) being rescued to near-control levels by transgene expression (B6.XY^{POS} Tg). ***P* ≤ 0.01 (two-tailed *t*-test); B6=C57BL6/J.

gonad development, in control and *Gadd45g* transgenic B6.Y^{POS} gonads (Fig. 1C, D and E). We confirmed that *Sry* expression was reduced at the 16 tail somite (ts) stage in B6.Y^{POS} gonads when compared to B6.Y^{B6} controls (Fig. 1C), consistent with earlier reports. Similarly, and presumably as a consequence, *Sox9* expression was greatly reduced at these early stages (Fig. 1D). However, in the presence of the *Gadd45g* BAC transgene, *Sry* expression levels were higher, compared with those detected in B6.Y^{B6} controls (Fig. 1C). These semi-quantitative WMISH data on *Sry* expression were confirmed by quantitative RT-PCR analysis, suggesting that WMISH is a reliable method for detecting its expression (Fig. 1E). *Sox9* expression also recovered, though not to levels comparable to B6.Y^{B6} controls, perhaps due to the presence of the SRY^{POS} protein isoform (Fig. 1D). These data suggest that overexpression of *Gadd45g* can have a positive impact on *Sry*^{POS} expression. However, they do not reveal whether this positive impact reflects an absolute requirement for GADD45g in timely expression of *Sry*, or whether, like CBX2 (Garcia-Moreno et al. 2019), this requirement is dependent on levels of canonical WNT signalling.

To test the role of canonical WNT signalling in the context of testis determination defects on B6/J, we used a loss-of-function *Rspo1* allele (Chassot et al. 2008) to assess its impact on B6.Y^{POS} gonadal sex reversal. As a control, we also examined the impact of a *Foxl2* null allele (Schmidt et al. 2004), thereby assessing the role of the two major pro-ovarian (granulosa cell) pathways in XY fetal ovary development. B6.Y^{POS} adults lacking a single copy of *Rspo1* (*Rspo1*^{-/+}) were scored as males at weaning and at necropsy (12 weeks of age) were found to contain testes of reduced size compared to B6.Y^{B6} controls, but with no overt changes in morphology or anatomy (Fig. 2A, B and Supplementary Fig. 2). Examination of fetal gonads at 14.5 dpc in B6.Y^{POS}, *Rspo1*^{-/+} animals by WMISH revealed substantial rescue of gonadal sex reversal, with extensive *Sox9*-positive testis cord formation throughout much of the gonad (Fig. 2C), which lacked *Stra8* expression apart from at the poles (Fig. 2D). B6.Y^{POS}, *Rspo1*^{-/-} homozygous fetal gonads at 14.5 dpc also had testis cords and completely lacked *Stra8* expression, but *Sox9* expression was reduced and the testes were smaller than B6.Y^{POS}, *Rspo1*^{-/+} counterparts (Fig. 2C and D). Control gonads (B6.Y^{POS}, *Rspo1*^{+/+}) were




Figure 2 Rescue of B6.Y^{POS} gonadal sex reversal by genetic ablation of *Rspo1*. (A) Images of adult testes (11.5 weeks) from control (XY^{B6}) and B6.Y^{POS}, *Rspo1*^{-/+} heterozygous (+/R) animals; (B) Testis weights of the animals from the same genotypic cohorts (per terminal body weight). WMISH of XX WT, XY (B6.Y^{B6}) WT, B6.Y^{POS} and B6.Y^{POS} embryos lacking one (+/R) or two (R/R) copies of *Rspo1* (left to right) at 14.5 dpc with *Sox9* (C) and *Stra8* (D) probes; WMISH of XX WT, XY (B6.Y^{B6}) WT, B6.Y^{POS} and B6.Y^{POS} embryos lacking one (+/F) or two (F/F) copies of *Foxl2* (left to right) at 14.5 dpc with *Sox9* probe (E); WMISH of XX WT, XY (B6.Y^{B6}) WT, B6.Y^{POS} and B6.Y^{POS} embryos lacking one copy (+/R1) of *Rspo1* (left to right) with *Sox9* probe at 12.5 dpc (F) and 11.5 dpc (G); and with an *Sry* probe at 11.5 dpc (H). ****P* ≤ 0.001 (two-tailed *t*-test); B6 = C57BL6/J.

uniformly ovarian by contrast (Fig. 2C and D). Equivalent examination of B6.Y^{POS}, Foxl2^{-/-} fetal gonads 14.5 dpc revealed no significant rescue of XY gonadal sex reversal, with only very modest levels of Sox9-positive testicular tissue (Fig. 2E). The rescue was a little more pronounced in B6.Y^{POS}, Foxl2^{-/-} gonads, with a small amount of testicular tissue forming centrally, but still not to the extent of B6.Y^{POS}, Rspo1^{-/+} gonads (Fig. 2E). From this, we conclude that RSPO1-mediated signalling, and canonical WNT signalling by implication, is the dominant regulator of granulosa cell fate specification during B6.Y^{POS} sex reversal, and XY ovary development more generally. By comparison, FOXL2 plays a relatively minor role.

We then examined rescue of gonadal sex reversal in B6.Y^{POS}, Rspo1^{-/+} fetuses at earlier stages. Examination of Sox9 at 12.5 dpc revealed robust expression throughout the gonad, in contrast to B6.Y^{POS}, Rspo1^{+/+} controls, although without significant testis cord formation, suggesting a delay in testis determination (Fig. 2F). However, neither Sox9 (Fig. 2G) nor Sry (Fig. 2H) were up-regulated at 11.5 dpc in B6.Y^{POS}, Rspo1^{-/+} fetuses relative to in B6.Y^{POS}, Rspo1^{+/+} controls. This suggests that

the mechanism of rescue caused by haploinsufficiency of Rspo1 differs from that caused by transgenic overexpression of Gadd45g (compare with Fig. 1).

We then exploited this role for Rspo1 in XY ovary development by testing whether GADD45g regulates Sry expression in an RSPO1-dependent fashion by generating Gadd45g null mice (Gadd45g^{-/-}) that also lacked one or two copies of Rspo1. We first examined XY gonad development at 14.5 dpc in Gadd45g^{-/-}, Rspo1^{-/+} (GhomRhet) and Gadd45g^{-/-}, Rspo1^{-/-} (double knockout, DKO) fetuses, and controls (Fig. 3A, B, C, D, E and F). Examination of Sertoli cell (Fig. 3A), Leydig cell (Fig. 3B) and meiotic germ cell (Fig. 3C) development using cell lineage markers revealed a substantial degree of rescue of sex in both XY GhomRhet and DKO fetal gonads. Both types of rescued gonads exhibited Sox9- and AMH-positive Sertoli cells and DDX4-positive germ cells in reasonably well-formed testis cords throughout much of the organ (Fig. 3D, E and F), although this was variable from mutant to mutant. Rescue was generally more complete in XY DKO gonads; however, as with B6.Y^{POS}, Rspo1^{-/-} gonads, these were smaller than GhomRhet




Figure 3 Rescue of gonadal sex reversal in mice lacking *Gadd45g* by genetic ablation of *Rspo1*. WMISH of XX and XY^{B6} WT (+/+), XY *Gadd45g*^{-/-} (G/G), XY *Gadd45g*^{-/-}, *Rspo1*^{+/-} (G/G, +/R) and XY *Gadd45g*^{-/-}, *Rspo1*^{-/-} (DKO) fetal gonads at 14.5 dpc with Sox9 (A), Leydig cell marker *3b-Hsd* (B) and *Stra8* (C) probes; immunostaining of XY^{B6} +/+ (D), G/G, +/R (E) and DKO (F) gonadal sections at 14.5 dpc with anti-DDX4 (magenta) and anti-AMH (green) antibodies; adult testes (10 weeks) from XY +/+ (G) and XY G/G, +/R (H) males with weights (per terminal body weight) (I); WMISH of XX and XY^{B6} +/+, XY G/G, +/R and XY DKO fetal gonads with Sox9 at 12.5 (J) and 11.5 dpc (K) and Sry at 11.5 dpc (17 ts) (L). *****P* ≤ 0.0001 (two-tailed t-test); B6=C57BL6/J.

gonads (Fig. 3). XY *Gadd45g*^{-/-} gonads at the same stage were ovaries with no *Sox9* expression or Leydig cells (*3b-Hsd*-negative), but exhibiting robust *Stra8* expression (Fig. 3A, B and C). In contrast to XY *Gadd45g*^{-/-} adult mice, which are phenotypic females, XY GhomRhet mice were fully masculinized males with hypoplastic testes (Fig. 3G, H, I and Supplementary Fig. 2).

Finally, we examined *Sox9* and *Sry* expression in control, GhomRhet and XY DKO fetal gonads (Fig. 3J, K and L). At 12.5 dpc, *Sox9* expression was robust in GhomRhet and XY DKO gonads compared to *Gadd45g*^{-/-} controls (Fig. 3J). However, at 11.5 dpc (19 ts), no *Sox9* expression was detectable in any gonads lacking *Gadd45g* following standard WMISH (Fig. 3K) and *Sry* expression was also minimal at 11.5 dpc (17 ts) in the same mutant gonads, compared to robust expression in B6.Y^{B6} control gonads at this stage (Fig. 3L). We conclude that timely *Sry* expression requires GADD45g function irrespective of the zygosity, and thus levels of expression, of *Rspo1*.

Discussion

The study reported here further underlines the intimate positive relationship between *Gadd45g* and *Sry* expression and function. We have previously shown that the temporal expression profile of each gene is remarkably similar, with initiation in the centre of the developing gonad at around 10.5 dpc, prior to overt gonad differentiation, and cessation of expression once this is completed, at 13.5 dpc (Warr *et al.* 2012). The regulation of *Sry* expression by GADD45g is suggested by these profiles, especially given that expression of the latter precedes that of the former (Warr *et al.* 2012). Moreover, the expression of both is restricted to supporting cell precursors (Warr *et al.* 2012, Stevant *et al.* 2019). The spatiotemporal expression profile of these genes is, therefore, indicative of a role in testis determination, now clearly established for *Gadd45g* by a number of genetic ablation studies (Gierl *et al.* 2012, Warr *et al.* 2012, Johnen *et al.* 2013). We and others have reported a role for GADD45g in activation of MAPK signalling, in sex determination (Gierl *et al.* 2012, Warr *et al.* 2012) and other contexts (Lu *et al.* 2001). However, additional molecular functions cannot be excluded: for example, other members of the GADD45 family are implicated in epigenomic functions on chromatin (Barreto *et al.* 2007, Schmitz *et al.* 2009, Niehrs & Schafer 2012, Schafer *et al.* 2013), so much remains to be elucidated concerning the role of GADD45g in sex determination.

We report testicular development in fetal XY gonads lacking both *Gadd45g* and *Rspo1* (XY DKO), i.e. rescue of XY gonadal sex reversal in *Gadd45g*-deficient fetal gonads by genetic ablation of *Rspo1*, in contrast to the essentially ovarian development of XY gonads lacking both *Gadd45g* and *Foxl2* (see Supplementary Fig. 3). This indicates that *Rspo1*, and downstream canonical

WNT signals, are the primary determinant of ovarian tissue development in an XY *Gadd45g*-deficient context. The relatively insignificant role for *Foxl2* in this genetic context may reflect the later expression of this gene during fetal gonad development in mice (Auguste *et al.* 2011, Niu & Spradling 2020). The absence of any significant increase in *Sry* expression levels in XY DKO gonads at 11.5 dpc, when compared to XY *Gadd45g*-deficient gonads, indicates an absolute requirement for GADD45g in directing early, robust expression of *Sry*, even in the absence of the pro-WNT/pro-ovarian gene *Rspo1*. This observation is in contrast to the situation reported for XY gonads lacking both *Cbx2* and *Wnt4* (Garcia-Moreno *et al.* 2019). CBX2, also required for normal *Sry* expression (Katoh-Fukui *et al.* 2012), acts by epigenomic inhibition of WNT gene expression and in its absence XY gonadal sex reversal is observed. Rescue of sex reversal is observed upon additional genetic ablation of *Wnt4*, but rescue in *Cbx2*^{-/-}, *Wnt4*^{-/-} XY DKO gonads is associated with restoration of *Sry* expression levels at 11.5 dpc (Garcia-Moreno *et al.* 2019). It is unclear why similar rescue of *Sry* expression is not observed in the XY DKO reported here. It may reflect differences in background strain or *Sry* detection methods; alternatively, it may reflect differing impacts on canonical WNT signalling caused by loss of *Rspo1* and *Wnt4*, or, more likely, an essential and positive role for GADD45g in *Sry* regulation not replicated by the essentially negative, inhibitory role of CBX2 in opposing WNT signalling.

It is also interesting to note that significant rescue of gonadal sex reversal due to the absence of *Gadd45g* occurs when one or two functional copies of *Rspo1* are removed, suggesting that dosage of this pro-WNT gene is important in XY fetal ovary development. This is notable given that ablation of *Rspo1* does not significantly disrupt fetal XX ovary development, i.e. does not result in the appearance of testicular tissue in the fetal gonad, as is also the case following the loss of either *Wnt4* or *Foxl2* (Pannetier *et al.* 2016). This is presumably due to the absence of *Sry* in the XX context and/or functional redundancy with other pro-ovarian (anti-testis) factors. Indeed, partial rescue of XY fetal testis development in other examples of a double knockout involving a pro-testis and pro-ovarian gene pair (Lavery *et al.* 2012, Nicol & Yao 2015, Garcia-Moreno *et al.* 2019) is likely to be due to the presence of *Sry*. It is our conclusion that in the absence of *Gadd45g* and pro-WNT *Rspo1*, a testis-determining gene, perhaps *Sox9* or another pro-testis gene (Nicol & Yao 2015, Richardson *et al.* 2020), can eventually respond satisfactorily to SRY even when there is a delay in *Sry* reaching functional levels. It is also worth noting that loss of both copies of *Rspo1* results in an XY DKO (XY *Gadd45g*^{-/-}, *Rspo1*^{-/-}) gonad with testicular morphology of reduced size and with less prominent *Sox9* expression than the XY GhomRhet (*Gadd45g*^{-/-}, *Rspo1*^{+/+}) counterpart. This presumably reflects the role of RSPO1 and WNT signalling in cell

proliferation in the early gonad (Jeays-Ward *et al.* 2004). The impact on cell proliferation caused by loss of *Rspo1* results in the development of hypoplastic XY gonads with a reduced number of Sertoli cells (Gregoire *et al.* 2018). This impact of the loss of RSPO1 on Sertoli cell differentiation may also explain the less robust formation of testis cords, and associated reduced *Sox9* expression, in B6.Y^{POS} testes lacking *Rspo1* when compared to their heterozygous counterparts (see Fig. 2C).

In conclusion, we show that the requirement for GADD45g to initiate timely expression of *Sry* during testis determination is not contingent on the presence of RSPO1-dependent signals i.e. WNT signals. This observation is consistent with the model involving an essential role for GADD45g in direct activation of *Sry* transcription through MAPK signals (Gierl *et al.* 2012, Warr *et al.* 2012), but it does not rule out additional, perhaps negative, roles for *Gadd45g* in testis determination. The fact that rescue of B6.Y^{POS} sex reversal is more complete when *Gadd45g* is overexpressed by transgenesis, compared to the loss of *Rspo1*, possibly reflects other roles for GADD45g in promoting pro-testis gene expression or opposing pro-ovary genes.

Supplementary materials

This is linked to the online version of the paper at <https://doi.org/10.1530/REP-21-0443>.

Declaration of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

This research was supported by support to A G (MC_U142684167) through core funding at the Mammalian Genetics Unit, MRC Harwell Institute.

Author contribution statement

N W, P S, N C, J M, M P and S W performed the study and analysed data. M C and A G conceived the study. A G wrote the manuscript.

Acknowledgements

The authors thank all staff in Ward 5 of the Mary Lyon Centre at MRC Harwell for animal husbandry, especially Jackie Harrison, Michelle Sandell, Gemma Atkins, Anju Paudyal and Michelle Stewart. We thank Robin Lovell-Badge (The Francis Crick Institute, London, UK) and Mathias Treier (Max-Delbrück Center for Molecular Medicine, Berlin, Germany) for kindly providing *Foxl2* KO mice.

<https://rep.bioscientifica.com>

References

- Auguste A, Chassot AA, Gregoire EP, Renault L, Pannetier M, Treier M, Pailhoux E & Chaboissier MC 2011 Loss of R-spondin1 and Foxl2 amplifies female-to-male sex reversal in XX mice. *Sexual Development* 5 304–317. (<https://doi.org/10.1159/000334517>)
- Barreto G, Schafer A, Marhold J, Stach D, Swaminathan SK, Handa V, Doderlein G, Maltry N, Wu W, Lyko F *et al.* 2007 Gadd45a promotes epigenetic gene activation by repair-mediated DNA demethylation. *Nature* 445 671–675. (<https://doi.org/10.1038/nature05515>)
- Bogani D, Siggers P, Brixey R, Warr N, Beddow S, Edwards J, Williams D, Wilhelm D, Koopman P, Flavell RA *et al.* 2009 Loss of mitogen-activated protein kinase kinase kinase 4 (MAP3K4) reveals a requirement for MAPK signalling in mouse sex determination. *PLoS Biology* 7 e1000196. (<https://doi.org/10.1371/journal.pbio.1000196>)
- Bullejos M & Koopman P 2001 Spatially dynamic expression of Sry in mouse genital ridges. *Developmental Dynamics* 221 201–205. (<https://doi.org/10.1002/dvdy.1134>)
- Bullejos M & Koopman P 2005 Delayed Sry and Sox9 expression in developing mouse gonads underlies B6-Y(DOM) sex reversal. *Developmental Biology* 278 473–481. (<https://doi.org/10.1016/j.ydbio.2004.11.030>)
- Carre GA & Greenfield A 2014 Characterising novel pathways in testis determination using mouse genetics. *Sexual Development* 8 199–207. (<https://doi.org/10.1159/000358402>)
- Carre GA, Siggers P, Xipolita M, Brindle P, Lutz B, Wells S & Greenfield A 2018 Loss of p300 and CBP disrupts histone acetylation at the mouse Sry promoter and causes XY gonadal sex reversal. *Human Molecular Genetics* 27 190–198. (<https://doi.org/10.1093/hmg/ddx398>)
- Chassot AA, Ranc F, Gregoire EP, Roepers-Gajadien HL, Taketo MM, Camerino G, De Rooij DG, Schedl A & Chaboissier MC 2008 Activation of beta-catenin signaling by Rspo1 controls differentiation of the mammalian ovary. *Human Molecular Genetics* 17 1264–1277. (<https://doi.org/10.1093/hmg/ddn016>)
- Cool J, Defalco T & Capel B 2012 Testis formation in the fetal mouse: dynamic and complex de novo tubulogenesis. *Wiley Interdisciplinary Reviews: Developmental Biology* 1 847–859. (<https://doi.org/10.1002/wdev.62>)
- Cox S, Smith L, Bogani D, Cheeseman M, Siggers P & Greenfield A 2006 Sexually dimorphic expression of secreted frizzled-related (SFRP) genes in the developing mouse Mullerian duct. *Molecular Reproduction and Development* 73 1008–1016. (<https://doi.org/10.1002/mrd.20507>)
- Eicher EM, Washburn LL, Whitney JB & Morrow KE 1982 *Mus poschiavinus* Y chromosome in the C57BL/6J murine genome causes sex reversal. *Science* 217 535–537. (<https://doi.org/10.1126/science.7089579>)
- Garcia-Moreno SA, Lin YT, Futtner CR, Salamone IM, Capel B & Maatouk DM 2019 CBX2 is required to stabilize the testis pathway by repressing Wnt signaling. *PLoS Genetics* 15 e1007895. (<https://doi.org/10.1371/journal.pgen.1007895>)
- Gierl MS, Gruhn WH, Von Seggern A, Maltry N & Niehrs C 2012 GADD45G functions in male sex determination by promoting p38 signaling and Sry expression. *Developmental Cell* 23 1032–1042. (<https://doi.org/10.1016/j.devcel.2012.09.014>)
- Gonen N, Quinn A, O'Neill HC, Koopman P & Lovell-Badge R 2017 Normal levels of Sox9 expression in the developing mouse testis depend on the TES/TESCO enhancer, but this does not act alone. *PLoS Genetics* 13 e1006520. (<https://doi.org/10.1371/journal.pgen.1006520>)
- Gonen N, Futtner CR, Wood S, Garcia-Moreno SA, Salamone IM, Samson SC, Sekido R, Poulat F, Maatouk DM & Lovell-Badge R 2018 Sex reversal following deletion of a single distal enhancer of Sox9. *Science* 360 1469–1473. (<https://doi.org/10.1126/science.aas9408>)
- Greenfield A 2013 Sexual development. In *Textbook of Clinical Embryology*. Eds K Coward & D Wells. Cambridge University Press.
- Greenfield A 2015 Understanding sex determination in the mouse: genetics, epigenetics and the story of mutual antagonisms. *Journal of Genetics* 94 585–590. (<https://doi.org/10.1007/s12041-015-0565-2>)
- Gregoire EP, Stevant I, Chassot AA, Martin L, Lachambre S, Mondin M, De Rooij DG, Nef S & Chaboissier MC 2018 NRG1 signalling regulates the establishment of Sertoli cell stock in the mouse testis. *Molecular and Cellular Endocrinology* 478 17–31. (<https://doi.org/10.1016/j.mce.2018.07.004>)
- Grimmond S, Van Hateren N, Siggers P, Arkell R, Larder R, Soares MB, De Fatima Bonaldo M, Smith L, Tymowska-Lalanne Z, Wells C *et al.* 2000 Sexually dimorphic expression of protease nexin-1 and vanin-1

- in the developing mouse gonad prior to overt differentiation suggests a role in mammalian sexual development. *Human Molecular Genetics* **9** 1553–1560. (<https://doi.org/10.1093/hmg/9.10.1553>)
- Harris A, Siggers P, Corrochano S, Warr N, Sagar D, Grimes DT, Suzuki M, Burdine RD, Cong F, Koo BK *et al.* 2018 ZNRF3 functions in mammalian sex determination by inhibiting canonical WNT signaling. *PNAS* **115** 5474–5479. (<https://doi.org/10.1073/pnas.1801223115>)
- Hiramatsu R, Matoba S, Kanai-Azuma M, Tsunekawa N, Katoh-Fukui Y, Kurohmaru M, Morohashi KI, Wilhelm D, Koopman P & Kanai Y 2009 A critical time window of Sry action in gonadal sex determination in mice. *Development* **136** 129–138. (<https://doi.org/10.1242/dev.029587>)
- Hoffmeyer A, Piekorz R, Moriggl R & Ihle JN 2001 Gadd45gamma is dispensable for normal mouse development and T-cell proliferation. *Molecular and Cellular Biology* **21** 3137–3143. (<https://doi.org/10.1128/MCB.21.9.3137-3143.2001>)
- Jeays-Ward K, Dandonneau M & Swain A 2004 Wnt4 is required for proper male as well as female sexual development. *Developmental Biology* **276** 431–440. (<https://doi.org/10.1016/j.ydbio.2004.08.049>)
- Johnen H, Gonzalez-Silva L, Carramolino L, Flores JM, Torres M & Salvador JM 2013 Gadd45g is essential for primary sex determination, male fertility and testis development. *PLoS ONE* **8** e58751. (<https://doi.org/10.1371/journal.pone.0058751>)
- Katoh-Fukui Y, Miyabayashi K, Komatsu T, Owaki A, Baba T, Shima Y, Kidokoro T, Kanai Y, Schedl A, Wilhelm D *et al.* 2012 Cbx2, a polycomb group gene, is required for Sry gene expression in mice. *Endocrinology* **153** 913–924. (<https://doi.org/10.1210/en.2011-1055>)
- Larney C, Bailey TL & Koopman P 2014 Switching on sex: transcriptional regulation of the testis-determining gene Sry. *Development* **141** 2195–2205. (<https://doi.org/10.1242/dev.107052>)
- Lavery R, Chassot AA, Pauper E, Gregoire EP, Klopfenstein M, De Rooij DG, Mark M, Schedl A, Ghyselincx NB & Chaboissier MC 2012 Testicular differentiation occurs in absence of R-spondin1 and Sox9 in mouse sex reversals. *PLoS Genetics* **8** e1003170. (<https://doi.org/10.1371/journal.pgen.1003170>)
- Livermore C, Simon M, Reeves R, Stevant I, Nef S, Pope M, Mallon AM, Wells S, Warr N & Greenfield A 2020 Protection against XY gonadal sex reversal by a variant region on mouse chromosome 13. *Genetics* **214** 467–477. (<https://doi.org/10.1534/genetics.119.302786>)
- Lu B, Yu H, Chow C, Li B, Zheng W, Davis RJ & Flavell RA 2001 GADD45gamma mediates the activation of the p38 and JNK MAP kinase pathways and cytokine production in effector TH1 cells. *Immunity* **14** 583–590. ([https://doi.org/10.1016/s1074-7613\(01\)00141-8](https://doi.org/10.1016/s1074-7613(01)00141-8))
- Nef S, Stevant I & Greenfield A 2019 Characterizing the bipotential mammalian gonad. *Current Topics in Developmental Biology* **134** 167–194. (<https://doi.org/10.1016/bs.ctdb.2019.01.002>)
- Nicol B & Yao HH 2015 Gonadal identity in the absence of pro-testis factor SOX9 and pro-ovary factor beta-catenin in mice. *Biology of Reproduction* **93** 35. (<https://doi.org/10.1095/biolreprod.115.131276>)
- Niehrs C & Schafer A 2012 Active DNA demethylation by Gadd45 and DNA repair. *Trends in Cell Biology* **22** 220–227. (<https://doi.org/10.1016/j.tcb.2012.01.002>)
- Niu W & Spradling AC 2020 Two distinct pathways of pregranulosa cell differentiation support follicle formation in the mouse ovary. *PNAS* **117** 20015–20026. (<https://doi.org/10.1073/pnas.2005570117>)
- Pannetier M, Chassot AA, Chaboissier MC & Pailhoux E 2016 Involvement of FOXL2 and RSPO1 in ovarian determination, development, and maintenance in mammals. *Sexual Development* **10** 167–184. (<https://doi.org/10.1159/000448667>)
- Rahmoun M, Lavery R, Laurent-Chaballier S, Bellora N, Philip GK, Rossitto M, Symon A, Pailhoux E, Cammas F, Chung J *et al.* 2017 In mammalian foetal testes, SOX9 regulates expression of its target genes by binding to genomic regions with conserved signatures. *Nucleic Acids Research* **45** 7191–7211. (<https://doi.org/10.1093/nar/gkx328>)
- Richardson N, Gillot I, Gregoire EP, Youssef SA, De Rooij D, De Bruin A, De Cian MC & Chaboissier MC 2020 Sox8 and Sox9 act redundantly for ovarian-to-testicular fate reprogramming in the absence of R-spondin1 in mouse sex reversals. *eLife* **9** e53972. (<https://doi.org/10.7554/eLife.53972>)
- Schafer A, Karaulanov E, Stapf U, Doderlein G & Niehrs C 2013 Ing1 functions in DNA demethylation by directing Gadd45a to H3K4me3. *Genes and Development* **27** 261–273. (<https://doi.org/10.1101/gad.186916.112>)
- Schmidt D, Ovitt CE, Anlag K, Fehsenfeld S, Gredsted L, Treier AC & Treier M 2004 The murine winged-helix transcription factor Foxl2 is required for granulosa cell differentiation and ovary maintenance. *Development* **131** 933–942. (<https://doi.org/10.1242/dev.00969>)
- Schmitz KM, Schmitt N, Hoffmann-Rohrer U, Schafer A, Grummt I & Mayer C 2009 TAF12 recruits Gadd45a and the nucleotide excision repair complex to the promoter of rRNA genes leading to active DNA demethylation. *Molecular Cell* **33** 344–353. (<https://doi.org/10.1016/j.molcel.2009.01.015>)
- Sekido R & Lovell-Badge R 2008 Sex determination involves synergistic action of SRY and SF1 on a specific Sox9 enhancer. *Nature* **453** 930–934. (<https://doi.org/10.1038/nature06944>)
- Siggers P, Smith L & Greenfield A 2002 Sexually dimorphic expression of Gata-2 during mouse gonad development. *Mechanisms of Development* **111** 159–162. ([https://doi.org/10.1016/s0925-4773\(01\)00602-5](https://doi.org/10.1016/s0925-4773(01)00602-5))
- Stevant I, Kuhne F, Greenfield A, Chaboissier MC, Dermitzakis ET & Nef S 2019 Dissecting cell lineage specification and sex fate determination in gonadal somatic cells using single-cell transcriptomics. *Cell Reports* **26** 3272.e3–3283.e3. (<https://doi.org/10.1016/j.celrep.2019.02.069>)
- Warr N, Carre GA, Siggers P, Faleato JV, Brixey R, Pope M, Bogani D, Childers M, Wells S, Scudamore CL *et al.* 2012 Gadd45gamma and Map3k4 interactions regulate mouse testis determination via p38 MAPK-mediated control of Sry expression. *Developmental Cell* **23** 1020–1031. (<https://doi.org/10.1016/j.devcel.2012.09.016>)
- Warr N, Siggers P, Carre GA, Bogani D, Brixey R, Akiyoshi M, Tachibana M, Teboul L, Wells S, Sanderson J *et al.* 2014 Transgenic expression of Map3k4 rescues T-associated sex reversal (Tas) in mice. *Human Molecular Genetics* **23** 3035–3044. (<https://doi.org/10.1093/hmg/ddu020>)
- Warr N, May J, Teboul L, Suzuki T, Asami M, Perry ACF, Wells S & Greenfield A 2018 Characterisation and use of a functional Gadd45g bacterial artificial chromosome. *Scientific Reports* **8** 17318. (<https://doi.org/10.1038/s41598-018-35458-5>)
- Wright E, Hargrave MR, Christiansen J, Cooper L, Kun J, Evans T, Gangadharan U, Greenfield A & Koopman P 1995 The Sry-related gene Sox-9 is expressed during chondrogenesis in mouse embryos. *Nature Genetics* **9** 15–20. (<https://doi.org/10.1038/ng0195-15>)

Received 4 November 2021

First decision 26 January 2022

Revised manuscript received 3 March 2022

Accepted 22 March 2022