

## Evidence that pain sensitivity is rhythmic in humans, mainly driven by the endogenous circadian system and little by sleep

I Daguet, V Raverot, D Bouhassira, Claude Gronfier

### ▶ To cite this version:

I Daguet, V Raverot, D Bouhassira, Claude Gronfier. Evidence that pain sensitivity is rhythmic in humans, mainly driven by the endogenous circadian system and little by sleep. 2022. hal-03651567v1

## HAL Id: hal-03651567 https://hal.science/hal-03651567v1

Preprint submitted on 25 Apr 2022 (v1), last revised 9 Sep 2022 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1	Evidence that pain sensitivity is rhythmic in humans, mainly driven
2	by the endogenous circadian system and little by sleep
3	
4	
5	Daguet I <sup>1</sup> , Raverot V <sup>2</sup> , Bouhassira D <sup>3</sup> , and Gronfier C <sup>1*</sup>
6	
7	<sup>1</sup> Lyon Neuroscience Research Center (CRNL), Neurocampus, Waking Team, Inserm UMRS 1028, CNRS
8	UMR 5292, Université Claude Bernard Lyon 1, Université de Lyon, 69000, Lyon, France
9	<sup>2</sup> Centre de Biologie et de Pathologie Est, Groupement Hospitalier Est, Hospices Civils de Lyon, 69677,
10	Bron cedex, France
11	<sup>3</sup> Inserm U987, APHP, UVSQ, Paris-Saclay Université, CHU Ambroise Paré, 92100,
12	Boulogne-Billancourt, France
13	
14	
15	
16	* Corresponding author:
17	Claude Gronfier, PhD, HDR, https://orcid.org/0000-0002-6549-799X
18	claude.gronfier@inserm.fr

#### 19 Abstract

20 Pain intensity has been reported to fluctuate during the day in some experimental and clinical conditions, but 21 the mechanisms underlying these fluctuations are unknown. Although the circadian timing system is known to 22 regulate a wide range of physiological functions, its implication in pain regulation is unknown. We show here, 23 using highly controlled laboratory constant routine conditions, that pain sensitivity is rhythmic over the 24-24 hours and strongly controlled by the endogenous circadian timing system. We find that pain sensitivity follows 25 a sinusoidal circadian rhythmicity, with a maximum in the middle of the night and a minimum in the afternoon. 26 We also find a weak homeostatic control of pain sensitivity, with a linear increase over the 34 hours of prolonged 27 wakefulness, which parallels that of sleep pressure. Using mathematical modelling, we describe that the 28 circadian system accounts for 80% of the full magnitude of pain sensitivity over the 24 hours, and that sleep-29 related processes account for only 20%. This result reveals that nocturnal analgesia is predominantly induced 30 by the circadian system and has been wrongly attributed only to sleep. Our findings highlight the need to 31 consider the time of day in pain assessment, and suggest that personalized circadian medicine may be a 32 promising approach to pain management.

33

#### 34 Significance statement

We discovered that sensitivity to pain is rhythmic in healthy humans, that sensitivity is maximal at night and minimal in the afternoon. Contrarily to the current thinking that sleep is the best painkiller, we find that the 24h rhythmicity of sensitivity to pain is mainly controlled by a biological circadian clock in our body, and very little by our sleep. Our article reveals the neurobiological mechanisms involved in driving the rhythmicity of pain perception in humans, with the main time-piece located in the brain (the suprachiasmatic nuclei in the hypothalamus). Our findings challenge the current vision of pain physiology, and reveal the need to consider time-of-day and internal biological time for pain evaluation and pain management.

#### 42 Introduction

Pain intensity has been reported to fluctuate during the day in a number of clinical conditions (3). The cyclic nature of some headaches (4, 5) and the diurnal variation of pain related to osteoarthritis are classical clinical observations (6, 7). The mechanisms underlying these fluctuations, however, are unknown. In particular, it remains unclear whether such daily variations are related to the internal circadian clock, or to behavioral or environmental factors, such as the sleep/wake cycle or the rest-activity cycle.

Pain has two main interconnected components: a sensory-discriminative component (location, quality, duration, intensity etc.) and an emotional component (unpleasantness, anxiety, motivation, etc.) (8). This multidimensional nociceptive response involves the activation of numerous subcortical and cortical regions of the brain (e.g. somatosensory cortices, insula, thalamus, prefrontal cortex), often referred to as the "pain matrix" (9). These structures are known to be regulated by the sleep/wake cycle or the circadian clock (10–12), but it remains unclear whether pain sensitivity is rhythmic and how it is regulated.

54 The circadian timekeeping system plays a key role in physiology by regulating the rhythmicity of numerous 55 functions, from gene expression to cortical activity and behavioral functions (10, 11, 13-16). It is, therefore, 56 also likely to be involved in pain perception. The surprising lack of knowledge about the rhythmicity of pain 57 sensitivity may result from the impact of timing on pain perception rarely having been taken into account (3), 58 and the use of inappropriate protocols for the exploration of pain rhythmicity from a neurobiological and 59 mechanistic point of view. The experimental studies performed to date to investigate pain sensitivity changes 60 during the day in healthy individuals have reported conflicting results (17–22). In both experimental and clinical 61 studies, the limited number of measurements and their timing (mostly during the daytime) made it impossible 62 to demonstrate unequivocally the existence of a 24-hour rhythmicity in pain sensation. It is also impossible to determine the origin of any rhythmicity in pain from these studies, because neither of the two types of highly 63 64 controlled laboratory protocols (constant-routine and forced-desynchrony paradigms) capable of separating 65 endogenous and exogenous rhythms (23, 24) were used. Endogenous rhythms are controlled by the central 66 biological clock located in the suprachiasmatic nuclei (SCN) of the hypothalamus, and exogenous rhythms 67 depend on behavioral or environmental changes, such as the sleep/wake cycle, the dark/light cycle, or the 68 rest/activity cycle. In real-life conditions, endogenous and exogenous influences are expressed simultaneously, 69 making it impossible to attribute rhythmicity to one or the other. In this study, we aimed to determine whether

sensitivity to heat pain displays rhythmicity over the 24-h day, and to assess the precise contribution of the circadian clock and sleep-related processes, by systematically assessing pain sensation and gold-standard markers of circadian rhythmicity in highly controlled constant-routine conditions.

73

#### 74 **Results**

#### 75 Pain rhythmicity is regulated by homeostatic and circadian processes

76 Twelve healthy men aged  $22.7 \pm 3.3$  years (mean  $\pm$  SEM) participated in a 56-hour experimental protocol 77 (Figure 1) including a 34-hour highly controlled constant routine (CR) designed to unmask endogenous 78 rhythmicity (enforced wakefulness, constant posture, low physical and cognitive activity, constant dim light, 79 equicaloric snacks every hour)(23). We assessed the effect of time-of-day on pain sensitivity, by measuring heat 80 pain every two hours during the 34 h of constant routine. In accordance with the current view that two main 81 processes regulate sleep (25), and in agreement with studies showing that physiological functions (such as 82 executive functions (26)) and cortical brain responses (measured by EEG (16) and fMRI (10)) are influenced 83 by both sleep pressure and the circadian timing system, we then modeled the effect of time on pain with an 84 additive mathematical model including a linear component (sleep-related homeostatic drive - process S) and a 85 sinusoidal component (circadian drive - process C)(15).

86

87



**Figure 1.** Overview of the experimental protocol. After a day of habituation (day 1) and an 8-h sleep episode, participants were subjected to a 34-hour constant routine (CR: days 2 and 3). Melatonin levels were assessed hourly (blue stars); pain sensitivity, temperature, heart rate and RMSSD were evaluated every two hours (red circles). Participants arrived at about 10:00 on day 1 (down arrow) and left the laboratory at about 18:00 on day 3 (up arrow). Gray rectangles represent wakefulness in dim light (~ 0.5 lux) and black rectangles represent scheduled sleep in darkness.

#### 89 Pain sensitivity increases with sleep debt

We probed subjective pain (visual analog scale ratings) in response to two-second heat stimuli (42 °C, 44 °C and 46 °C) every two hours over the entire 34-hour constant routine (Figure 2A, 2B and 2C; all  $R^2 > 0.72$ ). A



Figure 2. Mean pain intensities in response to 2-second heat stimuli at 42 °C, 44 °C and 46 °C are rhythmic across the 34-h constant routine protocol (n = 12). Dark bars correspond to the average timing of habitual sleep episodes (biological night). Circadian time 0 corresponds to dim light melatonin onset (DLMOn, mean  $\approx 21:30$ ). A-C. Combined models (sum of linear and sinusoidal components) applied to normalized data (mean  $\pm$  SEM) for stimuli at 42 °C (A. R<sup>2</sup> = 0.72), 44 °C (B. R<sup>2</sup> = 0.92) and 46 °C (C. R<sup>2</sup> = 0.92). D-F. Linear components for stimuli at 42 °C (D. R<sup>2</sup> = 0.10; p = 0.23), 44 °C (E. R<sup>2</sup> = 0.73; p < 0.0001) and 46 °C (F. R<sup>2</sup> = 0.81; p < 0.00001). Pain sensitivity increases with time spent awake for stimuli at 44 °C and 46 °C. G-I. Sinusoidal components for stimuli at 42 °C (G. R<sup>2</sup> = 0.70), 44 °C (H. R<sup>2</sup> = 0.90) and 46 °C (I. R<sup>2</sup> = 0.86). Pain sensitivity follows a circadian rhythm, with maximal pain at 3:30 (42 °C and 44 °C) or 3:00 (46 °C).

92 linear component was observed for the stimuli at 44 °C and 46 °C (Figure 2E and 2F; all p < 0.0001; all R<sup>2</sup> > 93 0.73), but not for the less painful stimuli at 42 °C (Figure 2D; p = 0.23;  $R^2 = 0.10$ ). Our results thus confirm the 94 known relationship between sleep deprivation and greater pain sensitivity (27–29), but suggest that this 95 relationship may not apply to low levels of pain. As the participants were in a constant state of wakefulness 96 during the CR, the linear component of our model translates the effect of sleep debt and reflects homeostatic 97 sleep pressure. The slope of the linear regression line increased with stimulation temperature (Supplementary 98 Figure 1), so the largest changes in amplitude were observed for stimuli at 46 °C, which caused a change in pain 99 level of 1/10 on the visual analog scale (VAS). As pain responses were measured at three arbitrary temperatures, 100 we then used a modeling approach (classically used in pharmacology and photobiology) to extract an overall 101 pain sensitivity value (Figure 3). The mathematically modelled sigmoidal intensity response curve (based on 102 the combined results obtained at 42, 44 and 46 °C) yielded sensitivity values ( $ET_{50}$ ) that confirmed the results 103 reported above; a linear increase in sensitivity to pain with time awake (lower  $ET_{50}$  values) (Figure 3C;  $R^2 =$ 104 0.81; p < 0.01).

105

#### 106 Pain sensitivity is driven by the circadian timing system, with maximal pain felt during the night

107 Subjective measurements of pain in response to two-second thermal stimuli (42 °C, 44 °C and 46 °C) revealed 108 that pain sensitivity was influenced not only by sleep pressure, but also by the circadian timing system (Figure 109 2A, 2B and 2C; all  $R^2 > 0.72$ ). Indeed, independently of the effect of sleep pressure, a sinusoidal component in 110 our model strongly accounted for changes in pain sensitivity across the 34 hours of constant routine, with a pain 111 sensitivity peak between 3:00 and 4:30 for both the responses to graded stimuli (Figure 2G, 2H and 2I) and heat 112 pain thresholds (Supplementary Figure 2C). These results were confirmed by the modeling of a sigmoidal 113 intensity response curve, which also showed a strong circadian rhythmicity of pain sensitivity (Figure 3D;  $R^2 =$ 114 (0.93) and a pain peak in the middle of the night (at 4:30). Interestingly, the lack of circadian rhythmicity for 115 warm non-painful stimuli (Supplementary Figure 3C;  $R^2 = 0.13$ ) suggests that the rhythmicity of pain sensitivity 116 is specific to pain and is not related to a general rhythmicity of thermal sensitivity. These results provide the 117 first evidence, to our knowledge, of a circadian rhythmicity of pain sensitivity in humans.

118



Figure 3. Mean pain sensitivity (ET<sub>50</sub>) is rhythmic across the 34-h constant routine protocol (n = 12). A. Intensity response curves calculated on the 6 measures obtained at 42, 44, and 46 °C over two consecutive 2-hour segments (9 curves; all R<sup>2</sup> between 0.68 and 0.99). B. Combined model (sum of linear and sinusoidal components) applied to raw ET<sub>50</sub> values (R<sup>2</sup> = 0.96). C. Linear component (R<sup>2</sup> = 0.81; p < 0.01). ET<sub>50s</sub> decrease and pain sensitivity increases with time spent awake. D. Sinusoidal component (R<sup>2</sup> = 0.93). Pain sensitivity follows a circadian rhythm with maximal pain at 4:30. B, C and D. Dark bars correspond to average habitual sleep episodes (biological night). Circadian time 0 corresponds to DLMOn (mean DLMOn  $\approx 21:30$ ).

# 119 Changes in pain sensitivity over the 24-hour day are mostly induced by the circadian system rather than a lack 120 of sleep

We investigated the relative contributions of sleep and circadian drives to pain sensitivity, by calculating the mean changes in both these components and expressing them relatively to the total amplitude over 24 hours (Supplementary Figure 4). We found that the circadian system accounted for 80 % of the full magnitude of pain sensitivity changes over 24 hours, the remaining 20 % being accounted for by the homeostatic component. Surprisingly, the decrease in pain sensitivity attributable to sleep at night was very small.

#### 126 *Phase relationships between the circadian components of pain modulation and interoceptive responses*

127 Having identified a circadian drive for pain, we investigated whether the rhythm of pain sensitivity displayed 128 phase relationships with interoceptive responses. Using cross-correlation analyzes, we identified a clear phase 129 opposition ( $\sim$ 12-h lag) between the rhythms of pain sensitivity and core body temperature (Figure 4A), with the 130 acrophase of pain (at 3:30) occurring at about the same time as the nadir of core body temperature (at 3:00). We 131 also found that pain sensitivity peaked 1.5 hours after endogenous melatonin secretion (at 2:00) (Figure 4B). 132 Autonomic nervous system responses displayed strong circadian rhythmicity, with a nadir of vagal activity 133 (minimal heart rate) and a peak of parasympathetic activity (maximal RMSSD) at 2:00, preceding the pain 134 sensitivity peak by 1.5 hours (Figure 4C and 4D).



Figure 4. Phase relationships between circadian components of pain sensitivity and temperature (A), melatonin (B), heart rate (C) and parasympathetic activity (D) across the 34-hour constant routine protocol. Dark bars correspond to average habitual sleep episodes (biological night). Circadian time 0 corresponds to DLMOn (mean DLMOn  $\approx 21:30$ ). All curves represent the sine component of the modeled parameter. All panels. Circadian rhythm of VAS pain intensity scores in response to a two-second stimulation at 44 °C, with a sensitivity peak at 3:30 (green curve;  $R^2 = 0.90$ ). A. Circadian rhythm of baseline body temperature, with a minimal core body temperature at 3:00 (blue curve;  $R^2 = 0.97$ ). B. Circadian rhythm of melatonin secretion (pg/mL), with a secretion peak at 2:00 (yellow curve;  $R^2 = 0.98$ ). C. Circadian rhythm of heart rate, with a minimal heart rate at 2:00 (red curve;  $R^2 = 0.99$ ). D. Circadian rhythm of MSDD (parasympathetic activity), with an activity peak at 2:00 (purple curve;  $R^2 = 0.89$ ).

#### 135 Discussion

This is the first highly controlled laboratory study specifically designed to investigate pain rhythmicity and its underlying mechanisms in healthy individuals. Our results unequivocally demonstrate that pain sensitivity is endogenously driven by the circadian timing system, and that sleep and sleep deprivation have a much weaker influence on pain sensitivity than previously thought.

140 A limited number of peer-reviewed studies have systematically investigated the rhythmicity of pain perception 141 in healthy individuals. Careful analysis reveals that published results are equivocal, some studies showing no 142 rhythmicity, others reporting maximal sensitivity either during the day or during the night (17–21) A recent 143 modelling work, using pooled datasets from four experimental studies, proposed a sinusoidal model of pain 144 sensitivity very similar to ours, with a peak sensitivity close to midnight (43). However, because the model was 145 built on data obtained from different populations and protocols, and collected during either sleep, wake, rest, 146 activity, light, or dark conditions, both the phase (timing) and the origin of this rhythmicity in pain sensitivity 147 cannot be attributed to any underlying timing mechanism, neither circadian, nor sleep-related. Overall, although 148 often claimed by the authors, none of the previous studies we have analyzed has demonstrated that pain 149 perception was circadian, i.e. of endogenous origin.

150 By contrast, our results, showing a strong sinusoidal oscillation of pain sensitivity in a constant routine protocol, 151 i.e. in the absence of rhythmic influences and times cues, provide unequivocal evidence that the rhythmicity of 152 pain sensitivity is driven from within, by the endogenous circadian timing system, and does not result from 153 influences evoked the light-dark cycle, the rest-activity cycle, or the sleep-wake cycle. Indeed, if pain sensitivity 154 were to be regulated exclusively by the sleep/wake cycle, as previously thought, we would have observed a 155 peak in pain sensitivity at the end of our 34-hour experimental constant-routine day and not in the middle of it 156 (after 20 hours) as we did. The very observation that the cyclicity of pain sensitivity is driven by the circadian 157 system, independently from the sleep/wake cycle or any other environmental cycle, demonstrates that both the 158 rhythmicity and its specific timing (its phase) are fundamental requirements in humans. Contrarily to the widely 159 held view that pain sensitivity is driven by the sleep-wake cycle (decreasing during sleep and increasing during 160 the day), our quantification that the circadian oscillation accounts for 80% of the full magnitude of pain 161 sensitivity over the 24-h, and that sleep deprivation accounts for only 20% of it, reveals that sleep and sleep

162 deprivation have in fact a very modest effect on pain. We also conclude that nocturnal analgesia is 163 predominantly due to the circadian system, that it has wrongly been attributed only to sleep in previous studies. 164 The pathways linking the circadian timekeeping system to pain perception cannot be inferred from this study. 165 but the suprachiasmatic nucleus is undoubtedly the starting point, and the subcortical and cortical regions of the 166 brain (e.g. somatosensory cortices, insula, thalamus, prefrontal cortex), often referred to as the "pain matrix" 167 (9) are likely to be involved, given that they have been shown to be regulated by the sleep/wake cycle or the 168 circadian clock (10–12). Our study suggests an interoceptive regulation of pain, where the circadian pacemaker 169 is likely to be central. Pain is traditionally regarded as an exteroceptive response depending on both the 170 somatosensory and emotional systems, however, it has been suggested that it may also be part of the 171 interoceptive system, relating to the condition of the body (8, 44). The interoceptive responses underlying the 172 maintenance of the internal environment of the body are organized in a hierarchical manner. They involve a 173 number of extensively connected physiological systems, so any change in one interoceptive function is usually 174 associated with changes in one or several other interoceptive functions. Our data are consistent with this view 175 as they show that, like other interoceptive functions, pain is driven by a time-specific circadian rhythm that is 176 directly related to the rhythmicity of other functions. The phase opposition we find between pain sensitivity and 177 core body temperature (CBT) suggests an interaction between thermoregulation and nociception (45), both of 178 which are components of the interoceptive system (8, 44). The phase relationships observed between the 179 rhythms of the sympathetic and parasympathetic systems (as assessed by cardiovascular measurements) and 180 pain are also consistent with this hypothesis and suggest the existence of strong interactions between the 181 nociceptive pathways and the autonomic nervous system (ANS) (46-48). The circadian timing system may, via 182 the SCN, serve as a key interface between pain and other interoceptive functions. The mechanisms underlying 183 these interactions are unclear, but, interestingly, our data suggest that they are probably not mediated by 184 melatonin. Melatonin, a nocturnal hormone released by the pineal gland, is generally reported to induce 185 antinociceptive effects (49–51). Such effects are not consistent with the temporal relationship between peak 186 pain sensitivity and peak endogenous melatonin secretion reported here, which instead suggests a 187 pronociceptive effect of melatonin. None of these mechanisms can be validated on the basis of our results as 188 we describe only temporal relationships between time series, but they could all be relatively easily tested 189 experimentally to determine their causality. Alternatively, the circadian rhythmicity of pain may be accounted

190 for by direct control of the nociceptive network (or the cognitive/emotional structures) by the SCN. In this 191 regulatory model of pain regulation, the circadian system may be responsible for controlling the precise timing 192 of nociception (8). As the thalamus is a key player in the nociceptive pathway and projections from the SCN to 193 the anterior paraventricular thalamus have been identified (52), pain sensitivity may be directly modulated by 194 this brain structure over the course of the 24-hour day. Multiple other pathways could be involved. Using the 195 same highly controlled experimental conditions we employed here, a study showed that  $\sim 15\%$  of all identified 196 metabolites in plasma and saliva are under circadian control in humans (Dallmann 2012). These include 197 metabolites involved in pain pathways, and recently identified metabolites of neuroinflammation specifically 198 found elevated in patients with neuropathic pain compared to those without neuropathic pain (Pfyffer 2020). 199 Whether those metabolites are involved in all clinical conditions of pain or in experimentally induced pain is 200 unknown, but overlapping the human circadian metabolome and our results allows to propose that the circadian 201 system regulates pain sensitivity though multiple pathways, both in normal and pathological situations.

202 The influence of sleep and sleep deprivation on pain sensitivity is modest in terms of its impact on the full 203 magnitude of pain sensitivity over the 24-h, but it is not negligible. The linear increase in pain sensitivity that 204 we find during enforced wakefulness, after mathematically removing the circadian component, confirms that 205 pain sensitivity does increase with time spent awake and reveals that it is under the influence of an independent 206 (from the circadian system) homeostatic drive, possibly related to that involved in the buildup of sleep pressure 207 from waketime to bedtime (25). This finding is consistent with the studies we previously discussed (17-21, 43) 208 and with the classically described interaction between pain and sleep (3, 30-33), whereby pain sensitivity 209 appears to be driven by the sleep/wake cycle, with pain perception low in the morning after a night of good-210 quality sleep, increasing during the day to reach a peak before bedtime, and then decreasing during sleep (29). 211 In the absence of sleep (after one night of total sleep deprivation), pain sensitivity has been shown to be higher 212 than it was at the same time on the previous day (27, 28), highlighting that there is an analysic effect of sleep 213 and/or a hyperalgesic effect of sleep deprivation. This sleep drive is usually considered to explain why sleep disorders, such as insomnia, are associated with an exacerbation of clinical pain (29, 34). The reciprocal 214 interactions between sleep homeostasis and pain may result from functional changes in the interconnected sleep 215 216 and pain systems. Consistent with this hypothesis, sleep loss is associated with an increase in the activation of 217 somatosensory brain areas induced by painful stimuli, potentially reflecting an amplification of neuronal

218 responses in the cortical nociceptive systems and/or a disinhibition of normal thalamocortical pain signaling 219 (33). In addition, sleep deprivation blunts activity in areas of the brain involved in endogenous pain modulation, 220 such as the striatum and insular cortex (33). The specific mechanisms underlying the interactions between pain 221 and sleep remain unknown, but may involve sleep-promoting factors, such as adenosine (35). Adenosine 222 accumulates with increasing homeostatic sleep pressure during wakefulness, reaching high levels at the end of 223 the day (36, 37), and then declining during sleep (38). In addition to its role in the sleep/wake cycle, adenosine 224 is also involved in the nociceptive system and may play an anti- or pronociceptive role, depending on the 225 receptors activated (39, 40). Thus, the hyperalgesic effect of constant wakefulness reported here may be at least 226 partly due to adenosine accumulation, leading to A1B receptor activation (37). Obviously, other mediators, such 227 as cytokines, which also play a role in both pain (41) and sleep regulation (42), may be involved in the sleep-228 related modulation of pain sensitivity.

229 This study has a number of limitations. First, our protocol was conducted under non-ecological and highly 230 controlled laboratory conditions, which were nevertheless absolutely essential to dissect out the rhythmic and 231 endogenous elements of pain sensitivity. Pain sensation may be different in real-life conditions, but the 232 endogenous mechanisms controlling pain sensitivity are expected to be the same. The modest influence of sleep 233 deprivation on pain sensitivity that our model finds, may also be different in real-life conditions. Indeed, prior 234 to their experimental session in the laboratory, our participants underwent 3 weeks of sleep monitoring, during 235 which time they slept on average 8 hours per night, and ensured they were sleep satiated upon arrival. In real 236 life conditions, where sleep deprivation is common in our societies, the strength of sleep-related drive may be 237 higher than in our conditions. This does not invalidate our model, but asks for its careful interpretation in 238 different conditions (Prayag et al. 2021) and also for its evaluation in conditions of sleep deprivation. Second, 239 pain intensity was evaluated in healthy participants, with an experimental heat pain paradigm. It is conceivable 240 that sleep pressure and the circadian timing system have the same effect on any type of pain, but our results 241 cannot be directly extrapolated to clinical populations. Third, the population examined in this study consisted 242 exclusively of men. Circadian physiology is very similar in men and women, with only minor differences, such 243 as a slightly larger amplitude (53, 54), and a slightly shorter period (55) in women, but our results should not 244 be extrapolated to premenopausal women, in whom the menstrual cycle may modulate both the homeostatic 245 and circadian drives of pain sensitivity.

246 In conclusion, our results reveal the neurobiological mechanisms driving the rhythmicity of pain perception in 247 humans, with the main driving brain structure located in the suprachiasmatic nuclei of the hypothalamus. We show 248 that pain sensitivity is controlled by two superimposed processes: a strong circadian component and a modest 249 homeostatic sleep-related component. This finding may have clinical implications, as dysregulations of the 250 circadian system have been implicated in a number of diseases with major consequences for health (13). Such 251 alterations may also be involved in the pathophysiology of some chronic pain syndromes, as suggested for 252 cluster headaches, for example (56). The existence of a circadian rhythmicity in pain suggests that the efficacy 253 of pain management could be optimized using circadian medicine (1, 2). With this approach, analgesic 254 treatments could be administered according to the each patient's internal time (circadian time) rather than 255 according to a uniform timing schedule mostly based on pragmatic considerations (57, 58). Such circadian 256 approaches have already proved effective in cancer treatment (59), but have not been systematically evaluated 257 for the treatment of pain. Individually timed medication could improve chronic pain management and greatly 258 improve patients' quality of life, not only by improving treatment efficacy but also by reducing the adverse 259 effects of painkillers, including those pejorative to sleep and circadian physiology.

260

#### 261 Methods

#### 262 <u>Participants</u>

263 Twelve healthy men (20 - 29 years old, mean age =  $22.7 \pm 3.3$  years; BMI =  $21.8 \pm 3.1$  kg/m<sup>2</sup>) were included in 264 this study. Neurological, psychiatric and sleep disorders were excluded by clinical examination and 265 psychological questionnaires (Pittsburg Sleep Quality Index Questionnaire and Beck Depression Inventory)(60, 266 61). Participants had an intermediate chronotype (Horne and Ostberg Chronotype Questionnaire score between 267 31-69)(62) and had not done any shift work, or experienced transmeridian travel during the previous three 268 months. Participants had normal visual acuity (Landolt Ring Test and Monoyer scale), contrast vision 269 (Functional Acuity Contrast Test) and color vision (Farnworth D-15 and Ishihara Color Test). All experimental 270 procedures were carried out in accordance with the Declaration of Helsinki. The study was approved by the 271 local research ethics committee (CPP Lyon Sud-Est II) and participants provided written informed consent for 272 participation.

273

#### 274 <u>Study design</u>

275 Participants were asked to maintain a regular sleep/wake schedule (bedtimes and waketimes within  $\pm$  30 minutes 276 of self-targeted times) for an average of three weeks before admission to the laboratory, with verification by 277 wrist activity and light exposure recordings (ActTrust, Condor Instruments, São Paulo, Brazil). Subjects were 278 then admitted to the laboratory for a 56-hour experimental protocol (Figure 1), in which they were kept in an 279 environment free from external time cues (clocks, television, smartphones, internet, visitors, sunlight etc.). 280 Subjects maintained contact with staff members specifically trained to avoid communicating time-of-day 281 information or the nature of the experimental conditions to the subjects. Participants arrived at about 10:00 on 282 the first day. They were allowed to familiarize themselves with the laboratory environment, low light levels (< 283 0.5 lux), equipment, and measurements. Lunch and dinner were served at about 12:30 and 19:00. A series of 284 measurements were then performed until bedtime (participant's habitual bedtime), and an 8-hour sleep episode 285 was scheduled (constant darkness; recumbent position). This was followed by a 34-hour constant-routine 286 protocol beginning at the participant's usual waketime on day 2, and ending on day 3 (18:00 on average). 287 Habitual bedtimes were determined on the basis of sleep times averaged over the seven days preceding the 288 laboratory segment of the protocol. Average bedtime was 23:45 and average waketime was 8:00.

289

#### 290 <u>Constant routine protocol</u>

291 A constant routine (CR) paradigm was used to reveal the endogenous circadian rhythmicity of various 292 parameters. The CR was conducted under constant environmental conditions, to eliminate, or distribute across 293 the circadian cycle, the physiological responses evoked by environmental or behavioral stimuli (i.e. sleeping, 294 eating, changes in posture, light intensity variations)(23, 63). In practical terms, participants were asked to 295 remain awake for 34 hours (starting at their habitual waketime), with minimal physical activity, while lying in 296 a semi-recumbent (45°) posture in bed. This posture was also maintained for the collection of urine samples 297 and bowel movements. Room temperature (mean =  $23 \degree C \pm 0.6$  (SD)) and ambient very dim halogen light levels 298 were kept constant. Light intensity was homogeneous in the room (< 0.5 lux at the participant's eye level in all 299 directions of gaze). Participants were given small equicaloric snacks and fluids at hourly intervals, to maintain 300 an equal nutritional caloric intake and stable hydration over the circadian cycle. Caloric requirements were 301 calculated on the basis of basal metabolic rate determined with the Wilmore nomogram and were adjusted

302	upward by a 7 % activity factor(64, 65). Fluid intake was calculated for each subject, to account for the sedentary
303	nature of the CR(65). A member of the study staff remained in the room with the participant at all times during
304	the CR, to monitor wakefulness and to ensure compliance with the study procedures.

305

#### 306 *Heat and pain evaluation*

Thermal stimuli were applied to the forearm with a Peltier-type thermode  $(30 \times 35 \text{ mm})$  connected to a thermotest device (Somedic AB, Stockholm, Sweden). Heat detection and pain thresholds were determined according to the method of limits (mean of three measurements).

Thermode temperature was gradually increased from a baseline temperature of 32 °C, at a rate of 1 °C/s, and participants were asked to stop the increase in temperature when they started to feel a warm sensation (detection threshold) or a pain sensation (pain threshold). At this point, the temperature returned to baseline at a rate of 1 °C/s. A minimum interval of 20 s was respected between each threshold measurement. If participants had not pressed the button by the time the maximum temperature (50 °C) was reached, the stimulation was stopped and the maximum temperature was recorded as the threshold value.

The pain induced by graded thermal stimuli was assessed with a 100-mm visual analog scale (VAS). All 316 317 participants received stimulation with three pseudorandomized heat stimuli (42 °C, 44 °C and 46 °C). For each 318 stimulus, participants were asked to rate the intensity of the pain on a VAS, extending from "no pain" to 319 "maximal imaginable pain". For each stimulation, the thermode temperature gradually rose from baseline 320 temperature (32 °C) at a rate of 1 °C/s. Once the target temperature was reached, it was maintained for 2 s and 321 the temperature then returned to baseline. Stimuli were separated by an interval of at least 45 s. Pain sensitization 322 was prevented by applying the thermode to adjacent regions of the forearm, never using the same site for 323 consecutive stimuli.

For more precise assessments of pain sensitivity than could be achieved with the responses to arbitrary temperatures, intensity response curves were calculated (Figure 3). This is a better approach to the assessment of sensitivity, as it can be used to determine the half maximal effective temperature, or  $ET_{50}$ , corresponding to the stimulation temperature required to induce 50 % of the maximal response (pain intensity of 5/10).

- 328
- 329

330 The data were modeled with a sigmoidal function:

331 
$$f(temperature) = max + \frac{min - max}{1 + \left(\frac{temperature}{ET_{50}}\right)^{hillslope}}$$

As the VAS is a bounded scale, minimum (min) and maximum (max) pain scores were set at 0 and 10, respectively. Hillslope, the slope of the curve, and  $ET_{50}$  were left free. The statistical power of the modeling approach was increased by calculating sigmoidal fits over 4-hour time epochs, corresponding to two evaluations of pain sensitivity for each of the three stimuli (42 °C, 44 °C and 46 °C), providing six points on the regression curve (Supplementary Figure 5). The  $ET_{50}$  values were extracted from each of the nine sigmoidal regressions (see formula above; Figure 3A) and plotted over time (Figure 3B).

338

#### 339 <u>Body temperature</u>

340 Core body temperature was measured every 2 h, with an ear thermometer (Braun Thermoscan Pro 6000, Welch

Allyn, New York, USA). Body temperature was measured within 2-3 seconds, with a precision of 0.2 °C.

342

#### 343 <u>Electrocardiogram</u>

344 An electrocardiogram (ECG) was recorded with two adhesive skin electrodes (BlueSensor N, Ambu, Ballerup, 345 Denmark) positioned on the sternum and the lateral thorax (RA, LL, respectively, Fontaine bipolar precordial 346 leads). The signal was recorded at 256 Hz, with a Vitaport 4 digital recorder (Temec Instruments, Kerkrade, 347 The Netherlands), to assess autonomic nervous system activity. Heart rate (HR) and heart-rate variability (HRV) 348 were analyzed on the basis of the bipolar ECG signal. R-wave peak detection was performed over 10-second 349 windows during a 4.5-minute baseline resting episode. For interval analysis, data were resampled at a rate of 10 350 Hz. RMSSD was determined to estimate the vagally mediated changes reflected in HRV(66, 67). It was not 351 possible to obtain ECG data for the first participant, for technical reasons, so ECG analysis was performed for 352 11 participants.

353

#### 354 <u>Melatonin</u>

355 Saliva was collected hourly, with cotton swabs placed directly in the mouth of the participant (Salivettes,

356 Sarstedt, Nümbrecht, Germany). Samples were stored at -20 °C until centrifugation and assay. Melatonin levels

were measured with an in-house radioimmunoassay  $^{125}I$  (RIA). This assay was based on a competition technique. The radioactive signal, reflecting the amount of  $^{125}I$ -labeled melatonin, was therefore inversely proportional to the concentration of melatonin in the sample. The sensitivity of the assay was 1.5 pg/mL. The inter-assay coefficients of variation for high (18.5 pg/mL) and low (10 pg/mL) melatonin-concentration controls were 19 % and 22 % respectively, and the mean intra-assay coefficient of variation was below 10 %. We determined the circadian melatonin profile of each participant over a 24-hour day, by applying a three-harmonic regression individually to the raw data collected during the CR (days 2 and 3)(68, 69). The model equation was:

364 
$$f(time) = mesor + amplitude1 \times \cos\left(2\pi \times \frac{time}{tau} + phase1\right) + amplitude2 \times \cos\left(2\pi \times \frac{time}{\frac{tau}{2}} + phase2\right) + dental dent$$

365 
$$amplitude3 \times \cos\left(2\pi \times \frac{time}{tau} + phase3\right)$$

366

In the model, Tau (the circadian period) was constrained between 23.5 and 24.5 h; mesor, amplitudes (1 to 3)
and phases (1 to 3) were set free.

369 The dim light melatonin onset (DLMOn), corresponding to the circadian phase, was calculated for each 370 participant. DLMOn was defined as the time at which the ascending phase of the melatonin profile crossed the 371 25 % threshold of the peak-to-trough amplitude of the fitted curve. Due to technical problems with some saliva 372 samples, the full 24-hour melatonin profile could not be obtained for two participants. For one of these 373 participants, DLMOn was calculated on the basis of melatonin levels during the habituation day (day 1), rather 374 than during the CR, for which we could not determine melatonin concentrations. For the second participant, in 375 the absence of melatonin concentration data (flat profile below the limit of quantification of the assay), DLMOn 376 was estimated from the mean phase angle calculated between habitual bedtime and DLMOn (calculated from 377 data published by Gronfier et al., 2004)(68).

378

#### 379 <u>Statistics</u>

Outliers were identified on the basis of normalized data (*z*-scores) and were excluded from subsequent analyses (outlier.test, R, Version 3.6.1 - 2019-07-05, R Foundation for Statistical Computing, Vienna, Austria). We reduced inter-individual variability, by normalizing all data (except melatonin concentrations) by calculating individual *z*-scores and smoothing them with a moving average (calculated on 3 points). The endogenous

circadian phase was taken into account for each participant, by aligning the data with the onset of melatonin secretion (DLMOn). As DLMOn occurred at different times in different participants, individual melatonin onset values were set to 0 (DLMOn = circadian time 0), and all measurement times are expressed relative to melatonin onset. We modeled the effects of time on the responses observed during the 34-hour constant routine, using an additive model including a linear component (homeostatic, process S) and a sinusoidal component (circadian, process C). The equation of the combined model was:

390

$$f(time) = y0 + a \times time + mesor + amplitude \times \cos\left(2\pi \times \frac{time}{tau} + phase\right)$$

Tau (circadian period) was constrained between 23.5 and 24.5 hours(70, 71), whereas all other parameters were left free. Once the parameters of the combined model had been defined, process S and process C were modeled separately. The homeostatic component (process S) was regressed against the linear component of the model:  $f(time) = y0 + a \times time$ . The circadian rhythmicity (process C) of the data was regressed against the sinusoidal component of the model:  $f(time) = mesor + amplitude \times \cos (2\pi \times \frac{time}{tau} + phase)$ . Statistics were calculated with R (Version 3.6.1 - 2019-07-05, R Foundation for Statistical Computing, Vienna, Austria). Results were considered significant if p < 0.05. Unless otherwise stated, results are expressed as means

398 ± SEM.

#### 399 Author contributions

400	The experiment was conceived by CG, and designed by CG, DB and ID. Data were collected by ID and CG.
401	Melatonin assays were conducted by VR. Data were analyzed and interpreted by ID and CG. ID, CG and DB
402	wrote the manuscript, and VR revised the manuscript. All authors agree to be accountable for all aspects of the
403	work.
404	
405	Acknowledgments
406	We wish to thank all the volunteers who participated in this study. We also wish to thank the staff and students,
407	and especially Pauline Kirchhoff who participated in data collection and analysis. Special thanks also go to Dr
408	Alain Nicolas, who conducted the medical and physical examinations.
409	
410	Conflicts of interest
411	The authors declare that the research was conducted in the absence of any commercial or financial relationships
412	that could be construed as a potential conflict of interest.
413	
414	Funding
415	This work was supported by fundings from "Societé Française de Recherche et Médecine du Sommeil"
416	(SFRMS) and "Société Française d'Etude et de Traitement de la Douleur" (SFETD) to ID, and grants from the
417	French National Research Agency (ANR-12-TECS-0013-01 and ANR-16-IDEX-0005) to CG. ID was
418	supported by a doctoral fellowship from the French "Ministère de l'Enseignement Supérieur et de la Recherche"
419	

#### 420 References

- 421 1. S. Panda, The arrival of circadian medicine. *Nat Rev Endocrinol* 15, 67–69 (2019).
- 422 2. Peeples L, Medicine's secret ingredient it's in the timing. *Nature* 556 (7701), 290–292 (2018).
- 423 3. V. Palada, I. Gilron, B. Canlon, C. I. Svensson, E. Kalso, The circadian clock at the intercept of
  424 sleep and pain. *Pain* (2019) https://doi.org/10.1097/j.pain.00000000001786.
- 425 4. A. W. Fox, R. L. Davis, Migraine chronobiology. *Headache* **38**, 436–441 (1998).
- S. Gori, *et al.*, Sleep quality, chronotypes and preferential timing of attacks in migraine without aura. *J Headache Pain* 6, 258–260 (2005).
- A. N. Bellamy, R. B. Sothern, J. Campbell, W. W. Buchanan, Circadian rhythm in pain, stiffness, and manual dexterity in rheumatoid arthritis: relation between discomfort and disability. *Annals of the Rheumatic Diseases* 50, 243–248 (1991).
- 431 7. I. C. Kowanko, M. S. Knapp, R. Pownall, A. J. Swannell, Domiciliary self-measurement in the
  432 rheumatoid arthritis and the demonstration of circadian rhythmicity. *Ann. Rheum. Dis.* 41, 453–
  433 455 (1982).
- 434 8. A. D. Craig, How do you feel? Interoception: the sense of the physiological condition of the
  435 body. *Nat Rev Neurosci* 3, 655–666 (2002).
- 436 9. A. V. Apkarian, M. C. Bushnell, R.-D. Treede, J.-K. Zubieta, Human brain mechanisms of pain
  437 perception and regulation in health and disease. *European Journal of Pain* 9, 463–463 (2005).
- 438 10. V. Muto, *et al.*, Local modulation of human brain responses by circadian rhythmicity and sleep
  439 debt. *Science* 353, 687–690 (2016).
- 440 11. J. Q. M. Ly, *et al.*, Circadian regulation of human cortical excitability. *Nat Commun* 7, 11828
  441 (2016).
- S. A. Brown, A. Azzi, "Peripheral Circadian Oscillators in Mammals" in *Circadian Clocks*, Handbook of Experimental Pharmacology., A. Kramer, M. Merrow, Eds. (Springer Berlin Heidelberg, 2013), pp. 45–66.
- T. Roenneberg, M. Merrow, The Circadian Clock and Human Health. *Current Biology* 26, R432–R443 (2016).
- 447 14. A. Patke, M. W. Young, S. Axelrod, Molecular mechanisms and physiological importance of
  448 circadian rhythms. *Nat Rev Mol Cell Biol* 21, 67–84 (2020).
- I. Daguet, D. Bouhassira, C. Gronfier, Baseline Pupil Diameter Is Not a Reliable Biomarker of
   Subjective Sleepiness. *Frontiers in Neurology* 10 (2019).
- 451 16. C. Cajochen, J. K. Wyatt, C. A. Czeisler, D. J. Dijk, Separation of circadian and wake duration452 dependent modulation of EEG activation during wakefulness. *Neuroscience* 114, 1047–1060
  453 (2002).

- J. Aviram, T. Shochat, D. Pud, Pain Perception in Healthy Young Men Is Modified by TimeOf-Day and Is Modality Dependent. *Pain Medicine* 16, 1137–1144 (2015).
- 456 18. H. Gobel, P. Cordes, Circadian Variation of Pain Sensitivity in Pericranial Musculature.
   457 *Headache* 30, 418–422 (1990).
- H. J. Koch, C. Raschka, Diurnal Variation of Pain Perception in Young Volunteers Using the
   Tourniquet Pain Model. *Chronobiology International* 21, 171–173 (2004).
- 460 20. F. Strian, S. Lautenbacher, G. Galfe, R. Hölzl, Diurnal variations in pain perception and thermal
  461 sensitivity. *Pain* 36, 125–131 (1989).
- 462 21. C. G. Bachmann, *et al.*, Diurnal time course of heat pain perception in healthy humans.
   463 *Neuroscience Letters* 489, 122–125 (2011).
- W. P. Chapman, C. M. Jones, Variations in cutaneous and visceral pain sensitivity in normal subjects. *J. Clin. Invest.* 23, 81–91 (1944).
- 466 23. J. F. Duffy, D.-J. Dijk, Getting through to circadian oscillators: why use constant routines? *J. Biol. Rhythms* 17, 4–13 (2002).
- 468 24. N. Kleitman, *Sleep and Wakefulness*, Rev. ed (Chicago : University of Chicago Press, 1987).
- 469 25. A. A. Borbély, S. Daan, A. Wirz-Justice, T. Deboer, The two-process model of sleep
  470 regulation: a reappraisal. *Journal of Sleep Research* 25, 131–143 (2016).
- P. Sagaspe, *et al.*, Influence of Age, Circadian and Homeostatic Processes on Inhibitory Motor
  Control: A Go/Nogo Task Study. *PLoS ONE* 7, e39410 (2012).
- 473 27. R. A. Larson, J. R. Carter, Total sleep deprivation and pain perception during cold noxious
  474 stimuli in humans. *Scandinavian Journal of Pain* 13, 12–16 (2016).
- 475 28. S. H. Onen, A. Alloui, A. Gross, A. Eschallier, C. Dubray, The effects of total sleep
  476 deprivation, selective sleep interruption and sleep recovery on pain tolerance thresholds in
  477 healthy subjects. *J Sleep Res* 10, 35–42 (2001).
- Y. Wei, T. F. Blanken, E. J. W. Van Someren, Insomnia Really Hurts: Effect of a Bad Night's
  Sleep on Pain Increases With Insomnia Severity. *Front Psychiatry* 9, 377 (2018).
- 480 30. B. Faraut, *et al.*, Napping Reverses Increased Pain Sensitivity Due to Sleep Restriction. *PLOS*481 *ONE* 10, e0117425 (2015).
- 482 31. A. Herrero Babiloni, *et al.*, Sleep and pain: recent insights, mechanisms, and future directions in
  483 the investigation of this relationship. *J Neural Transm* (2019) https://doi.org/10.1007/s00702484 019-02067-z (October 8, 2019).
- 485 32. G. Moscou-Jackson, P. H. Finan, C. M. Campbell, J. M. Smyth, J. A. Haythornthwaite, The
  486 Effect of Sleep Continuity on Pain in Adults With Sickle Cell Disease. *The Journal of Pain* 16, 587–593 (2015).

- 488 33. A. J. Krause, A. A. Prather, T. D. Wager, M. A. Lindquist, M. P. Walker, The Pain of Sleep
  489 Loss: A Brain Characterization in Humans. *The Journal of Neuroscience* 39, 2291–2300
  490 (2019).
- 491 34. M. D. Cheatle, *et al.*, Assessing and Managing Sleep Disturbance in Patients with Chronic Pain.
   492 *Anesthesiol Clin* 34, 379–393 (2016).
- 493 35. M. F. Bear, B. Connors, M. Paradiso, "Brain Rhythms and Sleep" in *Neuroscience: Exploring*494 *the Brain*, (Wolters Kluwer, 2016), pp. 645–684.
- 495 36. M. Iovino, *et al.*, Vigilance States: Central Neural Pathways, Neurotransmitters And
  496 Neurohormone. *Endocr Metab Immune Disord Drug Targets* (2018)
  497 https://doi.org/10.2174/1871530318666180816115720.
- 498 37. H.-P. Landolt, Sleep homeostasis: A role for adenosine in humans? *Biochemical Pharmacology*499 75, 2070–2079 (2008).
- Sol 38. C. B. Saper, T. E. Scammell, J. Lu, Hypothalamic regulation of sleep and circadian rhythms.
   *Nature* 437, 1257–1263 (2005).
- Solari Sciences 7, 30 (2017).
   Solari Sciences 7, 30 (2017).
- 40. S. Ferré, *et al.*, Adenosine A2A receptors in ventral striatum, hypothalamus and nociceptive circuitry. *Progress in Neurobiology* **83**, 332–347 (2007).
- K. A. Sluka, D. J. Clauw, Neurobiology of fibromyalgia and chronic widespread pain.
   *Neuroscience* 338, 114–129 (2016).
- 508 42. J. Krueger, The Role of Cytokines in Sleep Regulation. CPD 14, 3408–3416 (2008).
- J. Crodelle, S. H. Piltz, M. H. Hagenauer, V. Booth, Modeling the daily rhythm of human pain
  processing in the dorsal horn. *PLoS Comput Biol* 15, e1007106 (2019).
- 44. A. D. Craig, A new view of pain as a homeostatic emotion. *Trends Neurosci.* 26, 303–307
  (2003).
- 45. P. Alfonsi, F. Adam, D. Bouhassira, Thermoregulation and pain perception: Evidence for a homoeostatic (interoceptive) dimension of pain. *Eur J Pain* 20, 138–148 (2016).
- 46. P. Cortelli, G. Giannini, V. Favoni, S. Cevoli, G. Pierangeli, Nociception and autonomic
  nervous system. *Neurol Sci* 34, 41–46 (2013).
- 517 47. T. Schlereth, F. Birklein, The Sympathetic Nervous System and Pain. *Neuromol Med* 10, 141–
  518 147 (2008).
- 519 48. E. E. Benarroch, Pain-autonomic interactions. *Neurol Sci* 27, s130–s133 (2006).
- W.-W. Chen, X. Zhang, W.-J. Huang, Pain control by melatonin: Physiological and
   pharmacological effects. *Experimental and Therapeutic Medicine* 12, 1963–1968 (2016).

- 50. M. Wilhelmsen, I. Amirian, R. J. Reiter, J. Rosenberg, I. Gögenur, Analgesic effects of
  melatonin: a review of current evidence from experimental and clinical studies: Analgesic
  effects of melatonin. *Journal of Pineal Research* 51, 270–277 (2011).
- 525 51. C. Zhu, *et al.*, Exogenous melatonin in the treatment of pain: a systematic review and meta-526 analysis. *Oncotarget* **8** (2017).
- 527 52. L. P. Morin, Neuroanatomy of the extended circadian rhythm system. *Experimental Neurology*528 243, 4–20 (2013).
- 53. S. W. Cain, *et al.*, Sex Differences in Phase Angle of Entrainment and Melatonin Amplitude in
  Humans. *Journal of Biological Rhythms* 25, 288–296 (2010).
- 54. P. J. Gunn, B. Middleton, S. K. Davies, V. L. Revell, D. J. Skene, Sex differences in the
  circadian profiles of melatonin and cortisol in plasma and urine matrices under constant routine
  conditions. *Chronobiology International* 33, 39–50 (2016).
- 534 55. J. F. Duffy, *et al.*, Sex difference in the near-24-hour intrinsic period of the human circadian 535 timing system. *Proceedings of the National Academy of Sciences* **108**, 15602–15608 (2011).
- 536 56. J. Hoffmann, A. May, Diagnosis, pathophysiology, and management of cluster headache. *The Lancet Neurology* 17, 75–83 (2018).
- 538 57. M. D. Ruben, D. F. Smith, G. A. FitzGerald, J. B. Hogenesch, Dosing time matters. *Science* 539 365, 547–549 (2019).
- 540 58. M. D. Ruben, *et al.*, A large-scale study reveals 24-h operational rhythms in hospital treatment.
   541 *Proc Natl Acad Sci USA*, 201909557 (2019).
- 542 59. A. Ballesta, P. F. Innominato, R. Dallmann, D. A. Rand, F. A. Lévi, Systems
  543 Chronotherapeutics. *Pharmacological Reviews* 69, 161–199 (2017).
- 60. A. T. Beck, C. H. Ward, M. Mendelson, J. Mock, J. Erbaugh, An inventory for measuring
  depression. *Arch. Gen. Psychiatry* 4, 561–571 (1961).
- 546 61. D. J. Buysse, C. F. Reynolds, T. H. Monk, S. R. Berman, D. J. Kupfer, The Pittsburgh Sleep
  547 Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Res* 28, 193–
  548 213 (1989).
- 549 62. J. A. Horne, O. Ostberg, A self-assessment questionnaire to determine morningness550 eveningness in human circadian rhythms. *Int J Chronobiol* 4, 97–110 (1976).
- 551 63. J. N. Mills, D. S. Minors, J. M. Waterhouse, Adaptation to abrupt time shifts of the oscillator (s)
  552 controlling human circadian rhythms. *The Journal of physiology* 285, 455–470 (1978).
- 64. C. M. Jung, *et al.*, Energy expenditure during sleep, sleep deprivation and sleep following sleep
  deprivation in adult humans. *The Journal of Physiology* 589, 235–244 (2011).
- 555 65. M. D. Mifflin, *et al.*, A new predictive equation for resting energy expenditure in healthy 556 individuals. *Am J Clin Nutr* **51**, 241–247 (1990).

- 557 66. F. Shaffer, J. P. Ginsberg, An Overview of Heart Rate Variability Metrics and Norms. *Front.* 558 *Public Health* 5, 258 (2017).
- Task Force of the European Society of Cardiology and the North American Society of Pacing
  and Electrophysiology, Heart rate variability. Standards of measurement, physiological
  interpretation, and clinical use. *Eur. Heart J.* 17, 354–381 (1996).
- 68. C. Gronfier, Efficacy of a single sequence of intermittent bright light pulses for delaying
  circadian phase in humans. *AJP: Endocrinology and Metabolism* 287, E174–E181 (2004).
- 564 69. E. N. Brown, C. A. Czeisler, The Statistical Analysis of Circadian Phase and Amplitude in
  565 Constant-Routine Core-Temperature Data. *J Biol Rhythms* 7, 177–202 (1992).
- J. F. Duffy, *et al.*, Sex difference in the near-24-hour intrinsic period of the human circadian timing system. *Proc. Natl. Acad. Sci. U.S.A.* 108 Suppl 3, 15602–15608 (2011).
- 568 71. C. Gronfier, K. P. Wright, R. E. Kronauer, C. A. Czeisler, Entrainment of the human circadian
  569 pacemaker to longer-than-24-h days. *Proceedings of the National Academy of Sciences* 104,
  570 9081–9086 (2007).
- 571

572





582

583 Supplementary Figure 1. Mean pain intensity in response to heat stimuli at 42 °C, 44 °C and 46 °C, across 584 the 34-hour constant routine protocol (n = 12). Dark bars correspond to average habitual sleep episodes 585 (biological night). Circadian time 0 corresponds to DLMOn (mean DLMOn  $\approx$  21:30). Combined models (sum 586 of linear and sinusoidal components) of pain sensitivity in response to heat stimuli at 42 °C (light blue curve), 587 44 °C (blue curve) and 46 °C (dark blue curve). The amplitude changes (between peak and through) in pain 588 intensity for 42 °C, 44 °C and 46 °C are of 0.7, 0.8, and 1 VAS respectively.

- 589
- 590
- 591
- 592
- 593



Supplementary Figure 2. Mean heat pain thresholds, across the 34-h constant routine protocol (n = 12). Dark bars correspond to average habitual sleep episodes (biological night). Circadian time 0 corresponds to DLMOn (mean DLMOn  $\approx$ 21:30). A. Combined model (sum of linear and sinusoidal components) applied to normalized data (mean  $\pm$  SEM; R<sup>2</sup> = 0.69). B. Linear component (R<sup>2</sup> = 0.53; p < 0.01). Pain sensitivity decreases with time spent awake. C. Sinusoidal component (R<sup>2</sup> = 0.57). Pain sensitivity follows a circadian rhythm with maximal pain at 4:30.



Supplementary Figure 3. Mean warm detection thresholds across the 34-h constant routine protocol (n = 12). Dark bars correspond to average habitual sleep episodes (biological night). Circadian time 0 corresponds to DLMOn (mean DLMOn  $\approx$ 21:30). A. Combined model (sum of linear and sinusoidal components) applied to normalised data (mean  $\pm$  SEM; R<sup>2</sup> = 0.70). B. Linear component (R<sup>2</sup> = 0 68; p < 0.0001). Heat sensitivity decreases with time spent awake. C. No sinusoidal component is found (R<sup>2</sup> = 0.13).









#### 674

575 Supplementary Figure 5. Pain Intensity Response Curve (IRC) at circadian time -9 (~ 12:30). Sigmoidal 576 regression, calculated on a 4-hour time epoch (10:30 – 14:30), corresponding to 2 evaluations of pain sensitivity 577 to each of the 3 heat stimulations (42 °C, 44 °C and 46 °C), providing 6 points for the regression. The  $ET_{50}$ 578 value, corresponding to a stimulus inducing a pain of 5/10, is extracted from the sigmoidal regression (here  $ET_{50}$ 579 = 1.66 log[temperature]).

- 680
- 681

#### 682

#### 683 Time-of-day mechanisms of heat detection and pain thresholds

684 Heat pain thresholds and warm detection thresholds were measured every 2 hours throughout the whole 34-hour 685 constant routine (Supplementary Figures 2 and 3; all  $R^2 > 0.69$ ). The significant linear trend observed for heat 686 pain thresholds (Supplementary Figure 2; p < 0.01;  $R^2 = 0.53$ ), suggests a decrease in pain sensitivity with sleep pressure. This result might reflect a deterioration of cognitivo-motor performances (slower reaction times) 687 688 associated with sleep pressure (1,2). The fact that this effect was not specific to the heat pain threshold, since a 689 similar relationship was seen for the warm detection threshold (Supplementary Figure 3;  $R^2 = 0.68$ ; p < 0.0001), 690 is consistent with this hypothesis. The strong circadian rhythm of pain sensitivity (with a peak at 4:30), assessed 691 through heat pain threshold measures, confirms the results found with graded heat stimuli (and presented in the 692 main article). The lack of circadian rhythmicity for warm non-painful stimuli (Supplementary Figure 3C;  $R^2 =$ 693 (0.13) suggests that the rhythmicity of pain sensitivity is specific to pain and is not related to a general 694 rhythmicity of thermal sensitivity.

695 696

#### 697 <u>References</u>

- Krause, A. J., Simon, E. B., Mander, B. A., Greer, S. M., Saletin, J. M., Goldstein-Piekarski, A. N., & Walker,
  M. P. (2017). The sleep-deprived human brain. Nature Reviews Neuroscience, 18(7), 404–418. https://doi.org/10.1038/nrn.2017.55
- 2. Schmidt, C., Collette, F., Cajochen, C., & Peigneux, P. (2007). A time to think: Circadian rhythms in human cognition. Cognitive Neuropsychology, 24(7), 755–789. https://doi.org/10.1080/02643290701754158
- 703