



HAL
open science

Numerical approximation of the stochastic Navier-Stokes equations through artificial compressibility

Jad Doghman

► **To cite this version:**

Jad Doghman. Numerical approximation of the stochastic Navier-Stokes equations through artificial compressibility: Numerical analysis of the stochastic Navier-Stokes. 2023. hal-03653870v2

HAL Id: hal-03653870

<https://hal.science/hal-03653870v2>

Preprint submitted on 12 Apr 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Numerical approximation of the stochastic Navier-Stokes equations through artificial compressibility

Jad Doghman*

CNRS, Fédération de Mathématiques de CentraleSupélec FR 3487, Univ. Paris-Saclay, CentraleSupélec, 91190 Gif-sur-Yvette, France

Abstract

A constructive numerical approximation of the two-dimensional unsteady stochastic Navier-Stokes equations of an incompressible fluid is proposed via a pseudo-compressibility technique involving a penalty parameter ε . Space and time are discretized through a finite element approximation and an Euler method. The convergence analysis of the suggested numerical scheme is investigated throughout this paper. It is based on a local monotonicity property permitting the convergence toward the unique strong solution of the stochastic Navier-Stokes equations to occur within the originally introduced probability space. Justified optimal conditions are imposed on the parameter ε to ensure convergence within the best rate.

Keywords: stochastic Navier-Stokes, multiplicative noise, cylindrical Wiener process, Penalty method, finite element, Euler method

2020 MSC: 76D05, 65M12, 35Q35, 35Q30, 60H15, 60H35

Funding and acknowledgement

Funding: The author is supported by a public grant as part of the Investissement d'avenir project [ANR-11-LABX-0056-LMH, LabEx LMH], and is part of the SIMALIN project [ANR-19-CE40-0016] of the French National Research Agency.

Acknowledgement: The author would like to thank Prof. Ludovic Goudenège (Ecole CentraleSupélec) for his valuable comments.

1 Introduction

The first thought that springs to mind when it comes to the numerical simulation of the Navier-Stokes equations (NSEs) is the complexity of the occurring situation, which can be represented by turbulent

*Email address: jad.doghman@centralesupelec.fr

behaviors and physical processes by which energy becomes not only unavailable but irrecoverable in any form. The notorious NSEs are widely-known for their essential role in modeling phenomena that emerge from aeronautical science, thermo-hydraulics, ocean dynamics, and so on. They read in this chapter's context:

$$\begin{cases} \frac{\partial v}{\partial t} - \nu \Delta v + [v \cdot \nabla]v + \nabla p = f + g(v) \frac{\partial W}{\partial t}, \\ \operatorname{div}(v) = 0, \\ v(0, \cdot) = v_0, \end{cases} \quad (1.1)$$

with $v = v(\omega, t, x)$ being the fluid velocity, $p = p(\omega, t, x)$ is the pressure, $f = f(\omega, t, x)$ embodies an external force, g represents the diffusion coefficient, and the positive constant ν designates the fluid kinematic viscosity. The quantity W is regarded as a cylindrical Wiener process.

The present paper deals with numerical approximations of the two-dimensional incompressible NSEs driven by a multiplicative noise, equipped with homogeneous Dirichlet boundary conditions, within a bounded polygonal domain of \mathbb{R}^2 . Since the construction of divergence-free subspaces is not an effortless task (see for instance [2, 9, 19]), the attention will be turned toward a variant of the underlying equations involving a pseudo-compressibility method, avoiding divergence-free fields. This variant possesses the unique strong solution of the stochastic NSEs when passing to the limit (in ε), under a few assumptions. To be more accurate, the model which will undergo the discretization later on satisfies:

$$\begin{cases} \frac{\partial v^\varepsilon}{\partial t} - \nu \Delta v^\varepsilon + [v^\varepsilon \cdot \nabla]v^\varepsilon + \frac{1}{2}[\operatorname{div}(v^\varepsilon)]v^\varepsilon + \nabla p^\varepsilon = f + g(v^\varepsilon) \frac{\partial W}{\partial t}, \\ \varepsilon \frac{\partial p^\varepsilon}{\partial t} + \operatorname{div}(v^\varepsilon) = 0, \\ (v^\varepsilon(0, \cdot), p^\varepsilon(0, \cdot)) = (v_0, \pi_0), \end{cases} \quad (1.2)$$

where v^ε and p^ε are the associated fluid velocity and pressure, respectively. The constant $\varepsilon > 0$ is called the penalty parameter and it represents a small scale that will eventually tend to zero with the other discretization parameters to recover a solution to equations (1.1), and (v_0, π_0) is the initial condition. Observe that system (1.2) has a supplementary initial datum $p^\varepsilon(0, \cdot)$ denoted by π_0 because it does not need to depend on ε . The notation π_0 shall not be considered as a quantity arising from the pressure of equations (1.1) either; namely $\pi_0 \neq p(0)$. The supplementary term $\frac{1}{2}[\operatorname{div}(v^\varepsilon)]v^\varepsilon$ ensures the well-posedness of the model (1.2), which is why it cannot be taken out. Notice that alternative configurations (also known as penalty methods) might have been possible, especially for the mass conservation equation of problem (1.2). For instance,

$$\begin{aligned} \varepsilon p^\varepsilon + \operatorname{div}(v^\varepsilon) &= 0, \\ \varepsilon \Delta p^\varepsilon - \operatorname{div}(v^\varepsilon) &= 0 \quad \text{with} \quad \frac{\partial p^\varepsilon}{\partial n} = 0, \\ \varepsilon \Delta \partial_t p^\varepsilon - \operatorname{div}(v^\varepsilon) &= 0 \quad \text{with} \quad \frac{\partial}{\partial n}(\partial_t p^\varepsilon) = 0, \quad \text{and} \quad p^\varepsilon(0, \cdot) = \pi_0. \end{aligned}$$

The reader may refer to [22], [23], [24], and [26] for thorough deterministic studies of the above mentioned techniques, including the one considered here. The convergence rate of the Stokes problem driven by a multiplicative noise and subject to an artificial compressibility was conducted in [10] where optimal rates are obtained.

The mass conservation equation in problem (1.2) returns, in terms of regularity, good a priori estimates for the pressure p^ε (see [17, Proposition 3.1]), which may be taken advantage of during the convergence rate analysis. In point of fact, the pressure's lack of time-regularity in equations (1.1) (see for instance [15, Theorem 4.1]) has a negative effect on the convergence rate of those equations, which appears through the time convergence rate $O(\Delta t^{-1})$, as it was illustrated in [5, Corollary 4.2].

Problem (1.2) was theoretically investigated in [17] where the authors conducted the existence and uniqueness properties of the associated solution. The proof technique therein consists of the local

monotonicity property of the sum of the Stokes operator and the nonlinear term. A discrete version of this method will be considered in the present paper in order to demonstrate the convergence of the proposed numerical scheme and to avoid the Skorokhod theorem as well.

Finite element analysis of system (1.2) will be carried out hereafter, allowing the space variables to be discretized across the domain D . The proposed approximate finite element spaces for the velocity vector v^ε and the pressure field p^ε consist of continuous piecewise polynomials whose degrees can be chosen arbitrarily without any constraint, unlike the case of a saddle point problem where a discrete inf-sup condition must be imposed, leading to restrictive choices. Time discretization relies on the Euler method and is offered in two options: linear and nonlinear (Algorithms 1 and 2). As broadly known, an implicit numerical scheme gathers more stability properties than an explicit version. This appears in both Algorithms 1 and 2, especially regarding the initial datum's regularity. In contrast, when it comes to iterates' uniqueness, explicit numerical schemes for stochastic partial differential equations perform better than implicit ones. Finally, in order for the proposed numerical scheme to convergence toward the unique strong solution of equations (1.1), the spatial and temporal discretization parameters along with the scale ε should vanish at the same time.

Unlike article [24] where a numerical scheme for the deterministic version of equations (1.2) is investigated, there will be no need for $\Delta t/\varepsilon$ to converge toward 0 when $\Delta t, \varepsilon \rightarrow 0$, with Δt being the time discretization step size, thanks to the finite element method and the used demonstration technique herein. According to the artificial compressibility method that has been chosen here, the supplementary term $\varepsilon \partial_t p^\varepsilon$ allows the pressure to gain time-regularity that is not traditional for the incompressible Navier-Stokes equations (1.1). This extra regularity is usually linked to the penalty parameter when studying the convergence rate. For instance, penalizing by $\varepsilon \Delta p^\varepsilon$ instead of $\varepsilon \partial_t p^\varepsilon$ imposes that $h^2/\sqrt{\varepsilon}$ should go to 0 as $\varepsilon, h \rightarrow 0$, with h being the space discretization step size, as illustrated in [10, Theorem 5.9] for the stochastic time-dependent Stokes problem.

This paper is split into five sections and is organized as follows. Section 2 provides the adequate preliminaries and configurations, including the required assumptions, solutions' definitions to problems (1.1), (1.2), and the numerical scheme. Section 3 is devoted to giving the main theorem of this paper. Solvability, stability, and convergence of the numerical approximation are given in Section 4 along with a linear version of the proposed numerical scheme. This same section grants a small analysis scope concerned with the best choice of the scale ε in terms of the discretization parameters regarding numerical schemes with saddle point aspects. Section 5 supplies the reader with pieces of evidence through numerical experiments and comparisons with other schemes. The last section concludes all the work in this paper.

2 Notations, materials and algorithm

Let $T > 0$ be a finishing time. Given a bounded polygonal domain $D \subset \mathbb{R}^2$ (for simplicity's sake), denote by ∂D its boundary, and by $\vec{n}: \partial D \rightarrow \mathbb{R}^2$ its corresponding unit outward normal vector field. Function spaces in the Navier-Stokes framework are commonly denoted by \mathbb{H} and \mathbb{V} and are defined by

$$\begin{aligned} \mathcal{V} &:= \left\{ z \in [C_c^\infty(D)]^2 \mid \operatorname{div}(z) = 0 \text{ in } D \right\}, \\ \mathbb{H} &:= \left\{ z \in (L^2(D))^2 \mid \operatorname{div}(z) = 0 \text{ a.e. in } D, z \cdot \vec{n} = 0 \text{ a.e. on } \partial D \right\}, \\ \mathbb{V} &:= \left\{ z \in (H_0^1(D))^2 \mid \operatorname{div}(z) = 0 \text{ a.e. in } D \right\}, \end{aligned}$$

where $C_c^\infty(D)$ denotes the space of $C^\infty(D)$ functions with compact support. The vector spaces will be henceforth indicated by blackboard bold letters for clarity's sake (e.g. $\mathbb{H}^1 = (H^1(D))^2$). The inner product of the Lebesgue space \mathbb{L}^2 and the duality product between \mathbb{H}_0^1 and \mathbb{H}^{-1} are denoted by (\cdot, \cdot) and $\langle \cdot, \cdot \rangle$, respectively. The parameter ε of equation (1.2) satisfies all this paper long the condition $\varepsilon \leq 1$, the Gelfand triple $(\mathbb{H}_0^1, \mathbb{L}^2, \mathbb{H}^{-1})$ will solely be employed, and the trilinear form

$$\hat{b}(u, v, w) := ([u \cdot \nabla]v, w) + \frac{1}{2} ([div(u)]v, w)$$

will be linked to equation (1.2). Two operators can be associated with \hat{b} ; the trilinear form $b(u, v, w) := ([u \cdot \nabla]v, w)$ that arises from the NSEs and the bilinear operator $\hat{B}: \mathbb{H}_0^1 \times \mathbb{H}_0^1 \rightarrow \mathbb{H}^{-1}$ which reads: $\langle \hat{B}(u, v), w \rangle = \hat{b}(u, v, w)$, for all $u, v, w \in \mathbb{H}_0^1$. The upcoming proposition lists a few properties of the trilinear form \hat{b} (cf. [22]).

Proposition 2.1 (i) $\hat{b}: \mathbb{H}_0^1 \times \mathbb{H}_0^1 \times \mathbb{H}_0^1 \rightarrow \mathbb{R}$ is continuous.

(ii) $\hat{b}(u, v, v) = 0$ for all $u, v \in \mathbb{H}_0^1$.

(iii) $|\hat{b}(u, v, w)| \leq C_D \|u\|_{\mathbb{L}^2}^{\frac{1}{2}} \|\nabla u\|_{\mathbb{L}^2}^{\frac{1}{2}} \|\nabla v\|_{\mathbb{L}^2} \|\nabla w\|_{\mathbb{L}^2}$, for all $u, v, w \in \mathbb{H}_0^1$.

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space endowed with a filtration $(\mathcal{F}_t)_{0 \leq t \leq T}$ such that \mathcal{F}_0 contains all the null sets and $\mathcal{F}_t = \bigcap_{s>t} \mathcal{F}_s$. Let K be a separable Hilbert space equipped with a complete orthonormal basis $\{w_k, k \geq 1\}$. The noise W will be considered hereafter as a K -valued cylindrical Wiener process and it is defined by

$$W(t, x) = \sum_{k \geq 1} \beta^k(t) w_k(x),$$

where $\{\beta^k(\cdot), k \geq 1\}$ is a sequence of independent and identically distributed real-valued Brownian motions. With that said, the required assumptions are listed below.

Assumptions

(S₁) For $p \in [2, +\infty)$, $v_0 \in L^{2p}(\Omega; \mathbb{L}^2)$ and $\pi_0 \in L^{2p}(\Omega; L_0^2(D))$ are \mathcal{F}_0 -measurable.

(S₂) For $p \in [1, +\infty)$, $f \in L^{2p}(\Omega; L^2(0, T; \mathbb{H}^{-1}))$ and $g \in L^2(\Omega; L^2(0, T; \mathcal{L}_2(K, \mathbb{L}^2)))$ satisfies

$$\begin{aligned} \|g(u) - g(v)\|_{\mathcal{L}_2(K, \mathbb{L}^2)} &\leq L_g \|u - v\|_{\mathbb{L}^2}, \quad \forall u, v \in \mathbb{L}^2, \\ \|g(u)\|_{\mathcal{L}_2(K, \mathbb{L}^2)} &\leq K_1 + K_2 \|u\|_{\mathbb{L}^2}, \quad \forall u \in \mathbb{L}^2, \end{aligned}$$

for some positive time-independent constants K_1, K_2, L_g such that $L_g \leq \sqrt{\frac{\nu}{2C_P^2}}$, where C_P is the Poincaré constant.

Throughout this paper, the writing $x \lesssim y$ designates $x \leq cy$ for a universal constant $c \geq 0$, the constant C_D may vary from one calculation to another; however, it will depend only on the domain D , and finally the symbol $\mathcal{L}_2(X, Y)$ refers to the space of Hilbert-Schmidt operators from X to Y , where X and Y are two Hilbert spaces.

2.1 Concept of solutions

According to [17], a solution to equations (1.2) satisfies the following definition.

Definition 2.1 Let $T > 0$ and $\varepsilon > 0$ be given. For a filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{0 \leq t \leq T}, \mathbb{P})$, a stochastic process $\{(v^\varepsilon(t), p^\varepsilon(t)), t \in [0, T]\}$ is said to be a strong solution to equations (1.2) under assumptions (S_1) - (S_2) if it belongs to $L^2(\Omega; C([0, T]; \mathbb{L}^2) \cap L^2(0, T; \mathbb{H}_0^1)) \times L^2(\Omega; C([0, T]; L_0^2(D)))$, and it satisfies for all $t \in [0, T]$, \mathbb{P} -a.s.

$$\begin{cases} (v^\varepsilon(t), \varphi) + \nu \int_0^t (\nabla v^\varepsilon(s), \nabla \varphi) ds + \int_0^t \hat{b}(v^\varepsilon(s), v^\varepsilon(s), \varphi) ds - \int_0^t (p^\varepsilon(s), \operatorname{div}(\varphi)) ds \\ = (v_0, \varphi) + \int_0^t \langle f(s), \varphi \rangle ds + \left(\int_0^t g(v^\varepsilon(s)) dW(s), \varphi \right), \quad \forall \varphi \in \mathbb{H}_0^1, \\ (\varepsilon p^\varepsilon(t) - \varepsilon \pi_0, q) + \int_0^t (\operatorname{div}(v^\varepsilon(s)), q) ds = 0, \quad \forall q \in L^2(D), \end{cases}$$

along with the energy inequality

$$\mathbb{E} \left[\sup_{0 \leq t \leq T} (\|v^\varepsilon(t)\|_{\mathbb{L}^2}^p + \varepsilon \|p^\varepsilon(t)\|_{\mathbb{L}^2}^p) e^{-\delta t} + p\nu \int_0^T \|\nabla v^\varepsilon(t)\|_{\mathbb{L}^2}^2 \|v^\varepsilon(t)\|_{\mathbb{L}^2}^{p-2} e^{-\delta t} dt \right] \leq C,$$

for all $p \in [2, +\infty)$, $\delta > 0$, and for some constant $C > 0$ depending on $\delta, p, T, v_0, \pi_0, f, K_1, K_2$ and ε .

On the other hand, a solution to problem (1.1) in 2D can be defined as follows.

Definition 2.2 Assume (S_1) - (S_2) and let $T > 0$. A stochastic process $\{v(t), t \in [0, T]\}$ on a given filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{0 \leq t \leq T}, \mathbb{P})$ is a strong solution to equations (1.1) if it belongs to $L^2(\Omega; C([0, T]; \mathbb{H}) \cap L^2(0, T; \mathbb{V}))$, and it fulfills for all $0 \leq t \leq T$, \mathbb{P} -a.s.

$$\begin{aligned} & (v(t), \varphi) + \nu \int_0^t (\nabla v(s), \nabla \varphi) ds + \int_0^t ([v(s) \cdot \nabla]v(s), \varphi) ds \\ & = (v_0, \varphi) + \int_0^t \langle f(s), \varphi \rangle ds + \left(\int_0^t g(v(s)) dW(s), \varphi \right), \quad \forall \varphi \in \mathbb{V}. \end{aligned}$$

2.2 Discretization

The time interval $[0, T]$ will be decomposed into $M \in \mathbb{N} \setminus \{0\}$ subintervals with equidistant nodes $\{t_\ell\}_{\ell=0}^M =: I_k$ for simplicity's sake. The corresponding step is denoted by $k := \frac{T}{M}$.

The spatial domain D , which is assumed to be convex, bounded and polygonal, will be covered by a quasi-uniform triangulation \mathcal{T}_h , with h being the diameters' maximum of all triangles. Let \mathbb{H}_h be a subspace of \mathbb{H}_0^1 consisting of $[C(\bar{D})]^2$ -valued piecewise polynomials over \mathcal{T}_h , and fulfilling for all $m \geq 2$:

$$\inf_{v_h \in \mathbb{H}_h} \{ \|v - v_h\|_{\mathbb{L}^2} + h \|\nabla(v - v_h)\|_{\mathbb{L}^2} \} \leq Ch^m \|v\|_{\mathbb{H}^m}, \quad \forall v \in \mathbb{H}_0^1 \cap \mathbb{H}^m. \quad (2.1)$$

The quasi-uniformity of \mathcal{T}_h permits the inverse inequality (cf. [3, Lemma 4.5.3]):

$$\|v_h\|_{\mathbb{H}^\ell} \leq \mathcal{C} h^{m-\ell} \|v_h\|_{\mathbb{H}^m}, \quad \forall v_h \in \mathbb{H}_h, \quad \forall 0 \leq m \leq \ell, \quad (2.2)$$

for some $\mathcal{C} > 0$ independent of h . Let L_h be a subspace of $L_0^2(D) := \{q \in L^2(D) \mid \int_D q dx = 0\}$ consisting of $C(\bar{D})$ piecewise polynomial functions over \mathcal{T}_h , and satisfying for all $m \geq 1$:

$$\inf_{p_h \in L_h} \|p - p_h\|_{\mathbb{L}^2} \leq Ch^m \|p\|_{\mathbb{H}^m}, \quad \forall p \in L_0^2(D) \cap H^m(D). \quad (2.3)$$

For $(v, p) \in \mathbb{L}^2 \times L^2(D)$, the associated orthogonal projections are denoted by $\Pi_h: \mathbb{L}^2 \rightarrow \mathbb{H}_h$ and $\rho_h: L^2(D) \rightarrow L_h$ and are defined by the following identities, respectively:

$$(v - \Pi_h v, \varphi_h) = 0, \quad \forall \varphi_h \in \mathbb{H}_h \quad \text{and} \quad (p - \rho_h p, q_h) = 0, \quad \forall q_h \in L_h. \quad (2.4)$$

Thanks to the pseudo-compressibility method which is provided by equations (1.2), the finite element pair (\mathbb{H}_h, L_h) is not forced to satisfy the discrete LBB condition.

For the sake of clarity, the notations φ^+ and φ^- will designate throughout this paper piecewise constant functions with respect to time. For instance, $\varphi^+(t) := \varphi^m, \forall t \in (t_{m-1}, t_m]$ and $\varphi^-(t) := \varphi^{m-1}, \forall t \in [t_{m-1}, t_m)$ for the a given sequence $\{\varphi^m\}_m$. The discrete derivation with respect to time will also intervene later on. For this purpose, the below proposition (cf. [4, Appendix B]) lists a few associated properties.

Proposition 2.2 *Given a sequence $\{\varphi^m\}_m$, the discrete derivative is defined by $d_t \varphi^m = \frac{\varphi^m - \varphi^{m-1}}{k}$, for all $m \in \{1, \dots, M\}$, and it fulfills the following assertions:*

- (i) $d_t(\varphi^+ \psi^+) = \varphi^+ d_t \psi^+ + \psi^- d_t \varphi^+$,
- (ii) $\int_0^T \varphi^+ d_t \psi^+ dt = \varphi^+(T) \psi^+(T) - \varphi^-(0) \psi^-(0) - \int_0^T (d_t \varphi^+) \psi^- dt$,
- (iii) $d_t e^{\varphi^+} = e^{\varphi^-} d_t \varphi^+ + e^{\delta} \frac{(\varphi^+ - \varphi^-)^2}{2k}$, for some $\delta \in (\varphi^-, \varphi^+)$.

Relying on Definition 2.1 and the space-time discretization, the numerical scheme which will be studied throughout the rest of this paper is given by:

Algorithm 1 *Let $m \in \{1, \dots, M\}$ and $(v_h^0, p_h^0) \in \mathbb{H}_h \times L_h$ be a starting point. For a given $(V_\varepsilon^{m-1}, \Pi_\varepsilon^{m-1}) \in \mathbb{H}_h \times L_h$ such that $(V_\varepsilon^0, \Pi_\varepsilon^0) := (v_h^0, p_h^0)$, find $(V_\varepsilon^m, \Pi_\varepsilon^m) \in \mathbb{H}_h \times L_h$ that satisfies*

$$\begin{cases} (V_\varepsilon^m - V_\varepsilon^{m-1}, \varphi_h) + k\nu(\nabla V_\varepsilon^m, \nabla \varphi_h) + k\hat{b}(V_\varepsilon^m, V_\varepsilon^m, \varphi_h) - k(\Pi_\varepsilon^m, \text{div}(\varphi_h)) \\ \quad = k\langle f^m, \varphi_h \rangle + (g(V_\varepsilon^{m-1})\Delta_m W, \varphi_h), \quad \forall \varphi_h \in \mathbb{H}_h, \\ \frac{\varepsilon}{k}(\Pi_\varepsilon^m - \Pi_\varepsilon^{m-1}, q_h) + (\text{div}(V_\varepsilon^m), q_h) = 0, \quad \forall q_h \in L_h, \end{cases}$$

where for all $m \in \{1, \dots, M\}$, $f^m := \frac{1}{k} \int_{t_{m-1}}^{t_m} f(t) dt$ and $\Delta_m W := W(t_m) - W(t_{m-1})$.

The initial datum (v_h^0, p_h^0) of Algorithm 1 is required to be uniformly bounded in $\mathbb{L}^2 \times L_0^2(D)$ with respect to h . To this end, it suffices to consider $v_h^0 = \Pi_h v_0$ and $p_h^0 = \rho_h \pi_0$ because both projectors Π_h and ρ_h are stable in L^2 (cf. [8]):

$$\|\Pi_h u\|_{\mathbb{L}^2} \leq \|u\|_{\mathbb{L}^2}, \quad \forall u \in \mathbb{L}^2 \quad \text{and} \quad \|\rho_h q\|_{L^2} \leq \|q\|_{L^2}, \quad \forall q \in L^2. \quad (2.5)$$

Owing to [25, Lemma III.4.5], it holds that

$$k \sum_{m=1}^M \|f^m\|_{\mathbb{H}^{-1}}^2 \leq \int_0^T \|f(t)\|_{\mathbb{H}^{-1}}^2 dt. \quad (2.6)$$

3 Main result

Theorem 3.1 *For $T > 0$, let $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{0 \leq t \leq T}, \mathbb{P})$ be a filtered probability space, $D \subset \mathbb{R}^2$ be a polygonal domain, and assumptions (S_1) - (S_2) be satisfied. Given a positive integer M , define the discretization step size $k := \frac{T}{M}$ such that $k \in (0, 1)$ and I_k forms a uniform partition of the time interval $[0, T]$. Let $\varepsilon \in (0, 1)$ be a given penalty parameter, and $h \in (0, 1)$ be the space discretization step size such that the triangulation \mathcal{T}_h is quasi-uniform. Define the finite element triple $(\mathbb{H}_h, L_h, (v_h^0, p_h^0))$ such that the initial datum (v_h^0, p_h^0) belongs to (\mathbb{H}_h, L_h) . Then, the following results are true:*

- *For a given triple $(k, h, \varepsilon) \in (0, 1)^3$, there exists a solution $\{(V_\varepsilon^m, \Pi_\varepsilon^m)\}_{m=1}^M$ to Algorithm 1 satisfying Lemmata 4.1, 4.2, and 4.3.*
- *For a family $\{k, h, \varepsilon\}$ of parameters fulfilling $k, h, \varepsilon \rightarrow 0$ simultaneously, such that the initial datum $v_h^0 \rightarrow v_0$ as $h \rightarrow 0$ in $L^2(\Omega; \mathbb{L}^2)$, the solution $\{(V_\varepsilon^m, \Pi_\varepsilon^m)\}$ of Algorithm 1 converges toward the unique strong solution of the stochastic Navier-Stokes equations (1.1) in the sense of Definition 2.2.*

All this paper long, the penalty parameter ε is meant to be a vanishing scale, just as the discretization parameters k and h . The passage to the limit in ε , h , and k will be simultaneous, meaning that none of the mentioned parameters should vanish on its own.

The convergence of Algorithm 1 can also be investigated with a fixed non-vanishing ε to obtain a solution to equations (1.2) in the sense of Definition 2.1. Then, one can take advantage of [17, Proposition 4.1] to gain a solution to equations (1.1) in the sense of Definition 2.2. However, this idea is beyond the scope of this paper.

4 Discussion

4.1 Existence and uniqueness of solutions

This section is devoted to giving existence and uniqueness properties to the discrete stochastic process $\{(V_\varepsilon^m, \Pi_\varepsilon^m)\}_{m=1}^M$. The solvability of Algorithm 1 and the measurability of its iterates are handled first in the following lemma.

Lemma 4.1 *Let $T > 0$ be fixed. Under assumptions (S_1) - (S_2) , Algorithm 1 has at least one discrete solution. Moreover, for all $m \in \{1, \dots, M\}$, the processes $V_\varepsilon^m: \Omega \rightarrow \mathbb{H}_h$ and $\Pi_\varepsilon^m: \Omega \rightarrow L_h$ are \mathcal{F}_{t_m} -measurable.*

Proof: The solvability of Algorithm 1 can be proven by induction. Indeed, assume that iterates V_ε^ℓ and Π_ε^ℓ exist for all $\ell \in \{1, \dots, m-1\}$. The existence of $(V_\varepsilon^m, \Pi_\varepsilon^m)$ is therefore the target. To this end, let $E := \mathbb{H}_0^1 \times L_0^2$, and $B_\varepsilon: E \rightarrow E$ be define by

$$(B_\varepsilon(u, p), (v, q))_{\mathbb{L}^2 \times L^2} = (u - V_\varepsilon^{m-1}(\omega), v) + k\nu (\nabla u, \nabla v) + k\hat{b}(u, u, v) - k(p, \operatorname{div}(v)) - k\langle f^m, v \rangle - (g(V_\varepsilon^{m-1}(\omega))\Delta_m W(\omega), v) + \varepsilon(p - \Pi_\varepsilon^{m-1}(\omega), q) + k(\operatorname{div}(u), q),$$

for all $(u, p), (v, q) \in E$, and for almost all $\omega \in \Omega$. The symbol $(\cdot, \cdot)_{\mathbb{L}^2 \times L^2}$ denotes the $\mathbb{L}^2 \times L^2(D)$ -inner product. Thanks to Proposition 2.1-(i), the continuity of B_ε can be tackled easily. Through the application

of Proposition 2.1-(ii), the Poincaré and Young inequalities, estimate (2.6), and assumption (S_2) , one obtains

$$\begin{aligned} (B_\varepsilon(u, p), (u, p))_{\mathbb{L}^2 \times L^2} &\geq \frac{1}{2} \|u\|_{\mathbb{L}^2}^2 - \frac{1}{2} \|V_\varepsilon^{m-1}\|_{\mathbb{L}^2}^2 + k\nu \|\nabla u\|_{\mathbb{L}^2}^2 - k \|f^m\|_{\mathbb{H}^{-1}} \|u\|_{\mathbb{H}^1} \\ &- \|g(V_\varepsilon^{m-1})\|_{\mathcal{L}_2(K, \mathbb{L}^2)} \|\Delta_m W\|_K \|u\|_{\mathbb{L}^2} + \frac{\varepsilon}{2} \|p\|_{L^2}^2 - \frac{\varepsilon}{2} \|\Pi_\varepsilon^{m-1}\|_{L^2}^2 \geq \frac{1}{4} \|u\|_{\mathbb{L}^2}^2 + \frac{\varepsilon}{2} \|p\|_{L^2}^2 \\ &- \frac{1}{2} \|V_\varepsilon^{m-1}\|_{\mathbb{L}^2}^2 - \frac{\varepsilon}{2} \|\Pi_\varepsilon^{m-1}\|_{L^2}^2 - \frac{C_D^2}{4\nu} \|f\|_{L^2(0, T; \mathbb{H}^{-1})}^2 - (K_1 + K_2 \|V_\varepsilon^{m-1}\|_{\mathbb{L}^2})^2 \|\Delta_m W\|_K^2 \geq 0, \end{aligned}$$

for all $(u, p) \in E_h(\omega) := \left\{ (v, q) \in \mathbb{H}_h \times L_h \mid \|v\|_{\mathbb{L}^2} \geq \sqrt{S(\omega)}, \|q\|_{L^2} \geq \|\Pi_\varepsilon^{m-1}(\omega)\|_{L^2} \right\}$, where $S(\omega) := 2\|V_\varepsilon^{m-1}(\omega)\|_{\mathbb{L}^2}^2 + \frac{C_D^2}{\nu} \|f\|_{L^2(0, T; \mathbb{H}^{-1})}^2 + 4(K_1 + K_2 \|V_\varepsilon^{m-1}(\omega)\|_{\mathbb{L}^2})^2 \|\Delta_m W\|_K^2$. Both $S(\omega)$ and $\|\Pi_\varepsilon^{m-1}(\omega)\|_{L^2}$ are \mathbb{P} -a.s. finite, thanks to the induction supposition. With that said, the Brouwer fixed point theorem [13, Corollary IV.1.1] implies the existence of at least one $(u_\omega, p_\omega) \in \mathbb{H}_h \times L_h$ such that $B_\varepsilon(u_\omega, p_\omega) = (0, 0)$, $\|u_\omega\|_{\mathbb{L}^2} \leq \sqrt{S(\omega)}$ and $\|p_\omega\|_{L^2} \leq \|\Pi_\varepsilon^{m-1}\|_{L^2}$. Therewith, it suffices to set $(V_\varepsilon^m, \Pi_\varepsilon^m) = (u_\omega, p_\omega)$. On the other hand, the measurability of $\{(V_\varepsilon^m, \Pi_\varepsilon^m)\}_{m=1}^M$ can also be demonstrated by induction. The idea consists in expressing the newly obtained iterates (u_ω, p_ω) in terms of the existing ones. This can be done through a universally Borel-measurable selector function $\sigma: \mathbb{H}_h \times L_h \times K \rightarrow \mathbb{H}_h \times L_h$. For instance, $(u_\omega, p_\omega) = \sigma(V_\varepsilon^{m-1}, \Pi_\varepsilon^{m-1}, \Delta_m W)$, and the \mathcal{F}_{t_m} -measurability arises from the Wiener increment $\Delta_m W$. The reader may refer to [7, Page 744] for a detailed approach. \square

Lemma 4.1 dealt with the existence of a discrete solution which might not be unique. In point of fact, uniqueness in the whole probability set Ω does not seem to hold due to the nonlinearity interaction. Also, a contraction argument does not perform well in the discrete settings because the discrete time-derivative of an exponential function leads to a supplementary term that blocks the demonstration (see Proposition 2.2-(iii)). However, it can be proven that iterates' uniqueness holds true in a sample subset of Ω as demonstrated in the following lemma.

Lemma 4.2 *Assume (S_1) - (S_2) and let $\delta > 0$ be a small constant. Solutions $\{(V_\varepsilon^m, \Pi_\varepsilon^m)\}_{m=1}^M$ to Algorithm 1 are \mathbb{P} -almost surely unique within either of the following probability subsets:*

$$\begin{aligned} (i) \quad \Omega_\delta^1 &:= \left\{ \omega \in \Omega \mid k \sum_{m=1}^M \|V_\varepsilon^m\|_{\mathbb{L}^4}^4 \leq \frac{1}{\delta} \right\} \text{ provided that } \frac{1}{\nu^3 \delta} \leq 4c_0^3, \\ (ii) \quad \Omega_\delta^2 &:= \left\{ \omega \in \Omega \mid \max_{1 \leq m \leq M} \|V_\varepsilon^m\|_{\mathbb{L}^2}^4 \leq \frac{1}{\delta} \right\} \text{ provided that } \frac{k}{\nu^3 \delta h^2} \leq \frac{2c_0^3}{\mathcal{C}^2}, \end{aligned}$$

for some universal constant $c_0 \in (0, 3^{-1} 2^{\frac{2}{3}}]$. Furthermore, $\mathbb{P}(\Omega_\delta^1) \geq 1 - \delta \mathbb{E} \left[k \sum_{m=1}^M \|V_\varepsilon^m\|_{\mathbb{L}^4}^4 \right]$ and

$$\mathbb{P}(\Omega_\delta^2) \geq 1 - \delta \mathbb{E} \left[\max_{1 \leq m \leq M} \|V_\varepsilon^m\|_{\mathbb{L}^2}^4 \right].$$

Proof: Assume that $\{(V_\varepsilon^m, \Pi_\varepsilon^m)\}_{m=1}^M$ and $\{(U_\varepsilon^m, P_\varepsilon^m)\}_{m=1}^M$ are solutions to Algorithm 1 starting from the same initial condition (v_h^0, p_h^0) . For all $m \in \{0, 1, \dots, M\}$, let $Z_\varepsilon^m := V_\varepsilon^m - U_\varepsilon^m$ and $Q_\varepsilon^m := \Pi_\varepsilon^m - P_\varepsilon^m$. Then, iterates $\{(Z_\varepsilon^m, Q_\varepsilon^m)\}_{m=1}^M$ satisfy for all $m \in \{1, \dots, M\}$ and \mathbb{P} -a.s. the following equations

$$\begin{cases} (Z_\varepsilon^m - Z_\varepsilon^{m-1}, \varphi_h) + k\nu (\nabla Z_\varepsilon^m, \nabla \varphi_h) + k(\hat{B}(V_\varepsilon^m, V_\varepsilon^m) - \hat{B}(U_\varepsilon^m, U_\varepsilon^m), Z_\varepsilon^m) \\ \quad - k(Q_\varepsilon^m, \operatorname{div}(\varphi_h)) = ([g(V_\varepsilon^{m-1}) - g(U_\varepsilon^{m-1})] \Delta_m W, \varphi_h), \quad \forall \varphi_h \in \mathbb{H}_h, \\ \frac{\varepsilon}{k} (Q_\varepsilon^m - Q_\varepsilon^{m-1}, q_h) + (\operatorname{div}(Z_\varepsilon^m), q_h) = 0, \quad \forall q_h \in L_h. \end{cases} \quad (4.1)$$

Observe that the stochastic term in equation (4.1) can be eliminated if one had $V_\varepsilon^{m-1} = U_\varepsilon^{m-1}$. Since $U_\varepsilon^0 = V_\varepsilon^0 = v_h^0$, an induction argument seems to be legitimate. Indeed, for $m = 1$ and $(\varphi_h, q_h) = (Z_\varepsilon^1, Q_\varepsilon^1)$, equation (4.1) turns into

$$\begin{aligned} & \|Z_\varepsilon^1\|_{\mathbb{L}^2}^2 + \varepsilon \|Q_\varepsilon^1\|_{L^2}^2 + k\nu \|\nabla Z_\varepsilon^1\|_{\mathbb{L}^2}^2 = k \langle \hat{B}(U_\varepsilon^1, U_\varepsilon^1) - \hat{B}(V_\varepsilon^1, V_\varepsilon^1), Z_\varepsilon^1 \rangle \\ & \leq 2k \|\nabla Z_\varepsilon^1\|_{\mathbb{L}^2}^{\frac{3}{2}} \|Z_\varepsilon^1\|_{\mathbb{L}^2}^{\frac{1}{2}} \|V_\varepsilon^1\|_{\mathbb{L}^4} \leq 3 \cdot 2^{-\frac{2}{3}} c_0 k \nu \|\nabla Z_\varepsilon^1\|_{\mathbb{L}^2}^2 + \frac{k}{4\nu^3 c_0^3} \|Z_\varepsilon^1\|_{\mathbb{L}^2}^2 \|V_\varepsilon^1\|_{\mathbb{L}^4}^4, \end{aligned} \quad (4.2)$$

where the first inequality employs the estimate $|\langle \hat{B}(u, u) - \hat{B}(v, v), z \rangle| \leq 2 \|\nabla(u - v)\|_{\mathbb{L}^2}^{3/2} \|u - v\|_{\mathbb{L}^2}^{1/2} \|z\|_{\mathbb{L}^4}$ for all $u, v, z \in \mathbb{H}_0^1$ (see the proof in [17, Lemma 2.3]), and the second inequality uses Young's inequality for some constant $c_0 \in (0, 3^{-1}2^{\frac{2}{3}}]$. Subsequently, equation (4.2) becomes

$$(1 - k4^{-1}\nu^{-3}c_0^{-3}\|V_\varepsilon^1\|_{\mathbb{L}^4}^4) \|Z_\varepsilon^1\|_{\mathbb{L}^2}^2 + \varepsilon \|Q_\varepsilon^1\|_{L^2}^2 \leq 0. \quad (4.3)$$

One way to obtain uniqueness is by multiplying equation (4.3) by the indicator function $\mathbb{1}_{\Omega_\delta^1}$ which grants $(1 - 4^{-1}\nu^{-3}c_0^{-3}\delta^{-1})\mathbb{1}_{\Omega_\delta^1} \|Z_\varepsilon^1\|_{\mathbb{L}^2}^2 + \varepsilon \mathbb{1}_{\Omega_\delta^1} \|Q_\varepsilon^1\|_{L^2}^2 \leq 0$. It follows that $Z_\varepsilon^1 = Q_\varepsilon^1 = 0$ a.e. in D and \mathbb{P} -a.s. in Ω_δ^1 provided that the coefficient of $\|Z_\varepsilon^1\|_{\mathbb{L}^2}^2$ is positive. The second way for uniqueness consists in multiplying equation (4.3) by $\mathbb{1}_{\Omega_\delta^2}$ after employing the inverse estimate (2.2). That is, $\|V_\varepsilon^1\|_{\mathbb{L}^4}^4 \leq 2\|V_\varepsilon^1\|_{\mathbb{L}^2}^2 \|\nabla V_\varepsilon^1\|_{\mathbb{L}^2}^2 \leq 2\mathcal{C}^2 h^{-2} \|V_\varepsilon^1\|_{\mathbb{L}^2}^4$, where the first inequality is due to Ladyzhenskaya (see [14, Lemma I.1]). Therefore, equation (4.3) turns into $(1 - 2^{-1}\nu^{-3}c_0^{-3}\mathcal{C}^2\delta^{-1}h^{-2}k)\mathbb{1}_{\Omega_\delta^2} \|Z_\varepsilon^1\|_{\mathbb{L}^2}^2 + \varepsilon \mathbb{1}_{\Omega_\delta^2} \|Q_\varepsilon^1\|_{L^2}^2 \leq 0$ which implies $Z_\varepsilon^1 = Q_\varepsilon^1 = 0$ a.e. in D and \mathbb{P} -a.s. in Ω_δ^2 provided the coefficient of $\|Z_\varepsilon^1\|_{\mathbb{L}^2}^2$ is positive. With that being said, it suffices to assume that $Z_\varepsilon^{m-1} = Q_\varepsilon^{m-1} = 0$ a.e. in D , \mathbb{P} -a.s. in either Ω_δ^1 or Ω_δ^2 , and re-apply the same technique to obtain a similar result for the rank m . Finally, estimates of $\mathbb{P}(\Omega_\delta^1)$ and $\mathbb{P}(\Omega_\delta^2)$ derive from the Markov inequality. \square

Remark 4.1 *Picking between Ω_δ^1 and Ω_δ^2 in Lemma 4.2 depends on the choice of the viscosity ν . Observe that the condition $\frac{1}{\nu^3\delta} \leq 4c_0^3$ does not allow δ to be small when ν is tiny. Therewith, choosing ν large (resp. small) corresponds to Ω_δ^1 (resp. Ω_δ^2). Moreover, lower bounds associated with $\mathbb{P}(\Omega_\delta^1)$ and $\mathbb{P}(\Omega_\delta^2)$ in Lemma 4.2 are finite as illustrated in Lemma 4.3. It is worth mentioning that $\mathbb{E} \left[k \sum_{m=1}^M \|V_\varepsilon^m\|_{\mathbb{L}^4}^4 \right] \lesssim \mathbb{E} \left[\max_{1 \leq m \leq M} \|V_\varepsilon^m\|_{\mathbb{L}^2}^4 \right]^{\frac{1}{2}} \mathbb{E} \left[\left(k \sum_{m=1}^M \|\nabla V_\varepsilon^m\|_{\mathbb{L}^2}^2 \right)^2 \right]^{\frac{1}{2}}$.*

4.2 A priori bounds and convergence

The first part of this section is dedicated to achieving stability of Algorithm 1, whose convergence toward the unique solution of equations (1.1) is handled in the second part.

4.2.1 A priori bounds

Lemma 4.3 *Let $p \in [2, +\infty) \cap \mathbb{N}$ be fixed and assumptions (S_1) - (S_2) be satisfied. Then, iterates $\{(V_\varepsilon^m, \Pi_\varepsilon^m)\}_{m=1}^M$ of Algorithm 1 fulfill the following estimates:*

$$\begin{aligned} (i) \quad & \mathbb{E} \left[\max_{1 \leq m \leq M} \|V_\varepsilon^m\|_{\mathbb{L}^2}^2 + k\nu \sum_{m=1}^M \|\nabla V_\varepsilon^m\|_{\mathbb{L}^2}^2 + \sum_{m=1}^M \|V_\varepsilon^m - V_\varepsilon^{m-1}\|_{\mathbb{L}^2}^2 \right] \leq C_1, \\ (ii) \quad & \mathbb{E} \left[\max_{1 \leq m \leq M} \|\Pi_\varepsilon^m\|_{L^2}^2 + \sum_{m=1}^M \|\Pi_\varepsilon^m - \Pi_\varepsilon^{m-1}\|_{L^2}^2 \right] \leq \frac{C_1}{\varepsilon}, \end{aligned}$$

$$(iii) \mathbb{E} \left[\max_{1 \leq m \leq M} \|V_\varepsilon^m\|_{\mathbb{L}^2}^{2p} + \left(k\nu \sum_{m=1}^M \|\nabla V_\varepsilon^m\|_{\mathbb{L}^2}^2 \right)^{2p-1} + \left(\sum_{m=1}^M \|V_\varepsilon^m - V_\varepsilon^{m-1}\|_{\mathbb{L}^2}^2 \right)^{2p-1} \right] \leq C_p,$$

$$(iv) \mathbb{E} \left[\max_{1 \leq m \leq M} \|\Pi_\varepsilon^m\|_{L^2}^{2p} + \left(\sum_{m=1}^M \|\Pi_\varepsilon^m - \Pi_\varepsilon^{m-1}\|_{L^2}^2 \right)^{2p-1} \right] \leq \varepsilon^{-2p-1} C_p,$$

for some constant $C_p \geq 0$ depending only on $\|v_0\|_{L^{2p}(\Omega; \mathbb{L}^2)}$, $\|\pi_0\|_{L^{2p}(\Omega; L^2)}$, D , ν , $\|f\|_{L^{2p}(\Omega; L^2(0, T; \mathbb{H}^{-1}))}$, T , K_1 , p and K_2 , with $C_1 = C_{p=1}$.

Proof: Replace (φ_h, q_h) by $(V_\varepsilon^m, \Pi_\varepsilon^m)$ in Algorithm 1 and employ the identity $(a - b, a) = \frac{1}{2}(\|a\|_{\mathbb{L}^2}^2 - \|b\|_{\mathbb{L}^2}^2 - \|a - b\|_{\mathbb{L}^2}^2)$ together with Proposition 2.1-(ii), Cauchy-Schwarz, Poincaré and Young's inequalities:

$$\begin{aligned} & \frac{1}{2} \|V_\varepsilon^m\|_{\mathbb{L}^2}^2 - \frac{1}{2} \|V_\varepsilon^{m-1}\|_{\mathbb{L}^2}^2 + \frac{1}{4} \|V_\varepsilon^m - V_\varepsilon^{m-1}\|_{\mathbb{L}^2}^2 + \frac{\varepsilon}{2} \|\Pi_\varepsilon^m\|_{L^2}^2 - \frac{\varepsilon}{2} \|\Pi_\varepsilon^{m-1}\|_{L^2}^2 \\ & + \frac{\varepsilon}{2} \|\Pi_\varepsilon^m - \Pi_\varepsilon^{m-1}\|_{L^2}^2 + \frac{k\nu}{2} \|\nabla V_\varepsilon^m\|_{\mathbb{L}^2}^2 \leq \frac{C_D^2 k}{2\nu} \|f^m\|_{\mathbb{H}^{-1}}^2 + \|g(V_\varepsilon^{m-1}) \Delta_m W\|_{\mathbb{L}^2}^2 \\ & + (g(V_\varepsilon^{m-1}) \Delta_m W, V_\varepsilon^{m-1}). \end{aligned} \quad (4.4)$$

Summing equations (4.4) over m from 1 to $\ell \in \{1, \dots, M\}$, then applying the mathematical expectation, condition $\varepsilon \leq 1$, estimates (2.5) and (2.6) yield

$$\begin{aligned} & \mathbb{E} \left[\|V_\varepsilon^\ell\|_{\mathbb{L}^2}^2 + \varepsilon \|\Pi_\varepsilon^\ell\|_{L^2}^2 + k\nu \sum_{m=1}^{\ell} \|\nabla V_\varepsilon^m\|_{\mathbb{L}^2}^2 + \frac{1}{2} \sum_{m=1}^{\ell} \|V_\varepsilon^m - V_\varepsilon^{m-1}\|_{\mathbb{L}^2}^2 \right. \\ & \left. + \sum_{m=1}^{\ell} \varepsilon \|\Pi_\varepsilon^m - \Pi_\varepsilon^{m-1}\|_{L^2}^2 \right] \leq \mathbb{E} \left[\|v_0\|_{\mathbb{L}^2}^2 + \|\pi_0\|_{L^2}^2 + \frac{C_D^2}{\nu} \|f\|_{L^2(0, T; \mathbb{H}^{-1})}^2 \right. \\ & \left. + 2 \sum_{m=1}^{\ell} \|g(V_\varepsilon^{m-1}) \Delta_m W\|_{\mathbb{L}^2}^2 \right], \end{aligned} \quad (4.5)$$

where the mathematical expectation of last term in equation (4.4) vanishes due to the $\mathcal{F}_{t_{m-1}}$ -measurability of V_ε^{m-1} together with assumption (S_2) . On the other hand, the last term of inequality (4.5) can be handled through the Itô isometry and assumption (S_2) as follows:

$$\begin{aligned} & \mathbb{E} [\|g(V_\varepsilon^{m-1}) \Delta_m W\|_{\mathbb{L}^2}^2] = \mathbb{E} \left[\left\| \int_{t_{m-1}}^{t_m} g(V_\varepsilon^{m-1}) dW(t) \right\|_{\mathbb{L}^2}^2 \right] = k \mathbb{E} \left[\|g(V_\varepsilon^{m-1})\|_{\mathcal{L}_2(K, \mathbb{L}^2)}^2 \right] \\ & \leq 2kK_1^2 + 2kK_2^2 \mathbb{E} [\|V_\varepsilon^{m-1}\|_{\mathbb{L}^2}^2]. \end{aligned} \quad (4.6)$$

Thus, the discrete Grönwall inequality implies

$$\begin{aligned} & \max_{1 \leq m \leq M} \mathbb{E} [\|V_\varepsilon^m\|_{\mathbb{L}^2}^2 + \varepsilon \|\Pi_\varepsilon^m\|_{L^2}^2] + \sum_{m=1}^M \mathbb{E} \left[k\nu \|\nabla V_\varepsilon^m\|_{\mathbb{L}^2}^2 + \frac{1}{2} \|V_\varepsilon^m - V_\varepsilon^{m-1}\|_{\mathbb{L}^2}^2 \right] \\ & + \mathbb{E} \left[\sum_{m=1}^M \varepsilon \|\Pi_\varepsilon^m - \Pi_\varepsilon^{m-1}\|_{L^2}^2 \right] \leq C_1, \end{aligned} \quad (4.7)$$

where $C_1 > 0$ depends only on $\|v_0\|_{L^2(\Omega; \mathbb{L}^2)}$, $\|\pi_0\|_{L^2(\Omega; L^2)}$, D , ν , $\|f\|_{L^2(\Omega; L^2(0, T; \mathbb{H}^{-1}))}$, T , K_1 and K_2 . To terminate the proof of estimates (i) and (ii), it suffices to reconsider equation (4.4), sum it over m

from 1 to $\ell \in \{1, \dots, M\}$, take the maximum over ℓ , then apply the mathematical expectation to get

$$\begin{aligned} & \mathbb{E} \left[\max_{1 \leq \ell \leq M} (\|V_\varepsilon^\ell\|_{\mathbb{L}^2}^2 + \varepsilon \|\Pi_\varepsilon^\ell\|_{L^2}^2) \right] \leq \mathbb{E} \left[\|v_0\|_{\mathbb{L}^2}^2 + \|\pi_0\|_{L^2}^2 + C_D^2 \nu^{-1} \|f\|_{L^2(0,T;\mathbb{H}^{-1})}^2 \right. \\ & \left. + 2 \sum_{m=1}^M \|g(V_\varepsilon^{m-1}) \Delta_m W\|_{\mathbb{L}^2}^2 + 2 \max_{1 \leq \ell \leq M} \sum_{m=1}^{\ell} (g(V_\varepsilon^{m-1}) \Delta_m W, V_\varepsilon^{m-1}) \right]. \end{aligned} \quad (4.8)$$

The penultimate term is estimated in inequality (4.6). The last term is controlled by

$$\begin{aligned} & \lesssim 6 \mathbb{E} \left[\left(k \sum_{m=1}^M \|g(V_\varepsilon^{m-1})\|_{\mathcal{L}_2(K, \mathbb{L}^2)}^2 \|V_\varepsilon^{m-1}\|_{\mathbb{L}^2}^2 \right)^{\frac{1}{2}} \right] \\ & \leq \frac{3}{4} \mathbb{E} \left[\max_{1 \leq \ell \leq M} \|V_\varepsilon^\ell\|_{\mathbb{L}^2}^2 \right] + \mathbb{E} \left[\frac{3}{4} \|v_h^0\|_{\mathbb{L}^2}^2 + 3k \sum_{m=1}^M (K_1^2 + K_2^2 \|V_\varepsilon^{m-1}\|_{\mathbb{L}^2}^2) \right], \end{aligned}$$

where Young's inequality and assumption (S_2) are used together with the Davis inequality which is applicable since the integrand can be considered as a simple function with respect to time. Obviously, the first term on the right-hand side must be absorbed in the left side of equation (4.8) and the remaining terms can be readily controlled through estimates (2.5) and (4.7). This completes the proof of assertions (i) and (ii). Estimates (iii) and (iv) can be demonstrated as follows: let $p \geq 2$ be an integer. Summing equation (4.4) over m from 1 to $\ell \in \{1, \dots, M\}$, making use of estimates (2.5),(2.6), then raising both sides to the power 2^{p-1} yield

$$\begin{aligned} & \max_{1 \leq \ell \leq M} (\|V_\varepsilon^\ell\|_{\mathbb{L}^2}^{2p} + \varepsilon^{2p-1} \|\Pi_\varepsilon^\ell\|_{L^2}^{2p}) + \left(\sum_{m=1}^M (\|V_\varepsilon^m - V_\varepsilon^{m-1}\|_{\mathbb{L}^2}^2 + \varepsilon \|\Pi_\varepsilon^m - \Pi_\varepsilon^{m-1}\|_{L^2}^2) \right)^{2p-1} \\ & + \left(k\nu \sum_{m=1}^M \|\nabla V_\varepsilon^m\|_{\mathbb{L}^2}^2 \right)^{2p-1} \lesssim \|v_0\|_{\mathbb{L}^2}^{2p} + \|\pi_0\|_{L^2}^{2p} + C_D \nu^{-2p-1} \|f\|_{L^2(0,T;\mathbb{H}^{-1})}^{2p} \\ & + \left(\sum_{m=1}^M \|g(V_\varepsilon^{m-1}) \Delta_m W\|_{\mathbb{L}^2}^2 \right)^{2p-1} + \left(\max_{1 \leq \ell \leq M} \sum_{m=1}^{\ell} (g(V_\varepsilon^{m-1}) \Delta_m W, V_\varepsilon^{m-1}) \right)^{2p-1}. \end{aligned} \quad (4.9)$$

The mathematical expectation of the penultimate term is estimated through assumption (S_2) , inequality $|a + b|^q \leq 2^{q-1}(|a|^q + |b|^q)$, the Burkholder-Davis-Gundy and Young inequalities as follows:

$$\begin{aligned} & \mathbb{E} \left[\left(\sum_{m=1}^M \|g(V_\varepsilon^{m-1}) \Delta_m W\|_{\mathbb{L}^2}^2 \right)^{2p-1} \right] \leq M^{2p-1-1} \mathbb{E} \left[\sum_{m=1}^M \left\| \int_{t_{m-1}}^{t_m} g(V_\varepsilon^{m-1}) dW(t) \right\|_{\mathbb{L}^2}^{2p} \right] \\ & \lesssim M^{2p-1-1} \sum_{m=1}^M \mathbb{E} \left[\left(\int_{t_{m-1}}^{t_m} \|g(V_\varepsilon^{m-1})\|_{\mathcal{L}_2(K, \mathbb{L}^2)}^2 dt \right)^{2p-1} \right] \\ & \leq T^{2p-1-1} \mathbb{E} \left[k \sum_{m=1}^M \|g(V_\varepsilon^{m-1})\|_{\mathcal{L}_2(K, \mathbb{L}^2)}^{2p} \right] \\ & \lesssim T^{2p-1-1} K_1^{2p} + T^{2p-1} K_2^{2p} \mathbb{E} \left[k \sum_{m=1}^M \|V_\varepsilon^{m-1}\|_{\mathbb{L}^2}^{2p} \right]. \end{aligned} \quad (4.10)$$

The last term of equation (4.9) can be controlled through the Burkholder-Davis-Gundy inequality, assumption (S_2) , the Young and Hölder inequalities as follows

$$\begin{aligned}
& \mathbb{E} \left[\left(\max_{1 \leq \ell \leq M} \sum_{m=1}^{\ell} (g(V_\varepsilon^{m-1}) \Delta_m W, V_\varepsilon^{m-1}) \right)^{2p-1} \right] \lesssim \mathbb{E} \left[\left(k \sum_{m=1}^M \|g(V_\varepsilon^{m-1})\|_{\mathcal{L}_2(K, \mathbb{L}^2)}^2 \|V_\varepsilon^{m-1}\|_{\mathbb{L}^2}^2 \right)^{2p-2} \right] \\
& \leq \mathbb{E} \left[\max_{1 \leq m \leq M} \|V_\varepsilon^{m-1}\|_{\mathbb{L}^2}^{2p-1} k^{2p-2} M^{2p-2-1} \sum_{m=1}^M \|g(V_\varepsilon^{m-1})\|_{\mathcal{L}_2(K, \mathbb{L}^2)}^{2p-1} \right] \\
& \leq \frac{1}{4} \mathbb{E} \left[\max_{1 \leq m \leq M} \|V_\varepsilon^m\|_{\mathbb{L}^2}^{2p} \right] + \frac{1}{4} \mathbb{E} [\|v_0\|_{\mathbb{L}^2}^{2p}] + T^{2p-1-2} \mathbb{E} \left[\left(k \sum_{m=1}^M \|g(V_\varepsilon^{m-1})\|_{\mathcal{L}_2(K, \mathbb{L}^2)}^{2p-1} \right)^2 \right] \\
& \leq \frac{1}{4} \mathbb{E} \left[\max_{1 \leq m \leq M} \|V_\varepsilon^m\|_{\mathbb{L}^2}^{2p} \right] + \frac{1}{4} \mathbb{E} [\|v_0\|_{\mathbb{L}^2}^{2p}] + T^{2p-1-1} \mathbb{E} \left[k \sum_{m=1}^M (K_1 + K_2 \|V_\varepsilon^{m-1}\|_{\mathbb{L}^2})^{2p} \right] \\
& \lesssim \frac{1}{4} \mathbb{E} \left[\max_{1 \leq m \leq M} \|V_\varepsilon^m\|_{\mathbb{L}^2}^{2p} \right] + \frac{1}{4} \mathbb{E} [\|v_0\|_{\mathbb{L}^2}^{2p}] + T^{2p-1} K_1^{2p} + T^{2p-1-1} K_2^{2p} \mathbb{E} \left[k \sum_{m=1}^M \|V_\varepsilon^{m-1}\|_{\mathbb{L}^2}^{2p} \right].
\end{aligned}$$

Putting it all together and applying the discrete Gröwall inequality to equation (4.9) complete the proof. \square

4.2.2 Convergence

Stability properties derived in Lemma 4.3 will play a crucial role in this part, especially to offer convergence results to $\{(V_\varepsilon^m, \Pi_\varepsilon^m)\}_{m=1}^M$ as $\varepsilon, k, h \rightarrow 0$. For this purpose, a few new notations must be evoked along with one important lemma consisting of a monotonicity property that allows the convergence of Algorithm 1 toward equations (1.1) to occur. For all $m \in \{1, \dots, M\}$, the new notations read:

$$\begin{aligned}
(\mathcal{V}_{\varepsilon, k, h}^+(t), \Pi_{\varepsilon, k, h}^+(t)) &:= (V_\varepsilon^m, \Pi_\varepsilon^m), \quad \forall t \in (t_{m-1}, t_m], \\
(\mathcal{V}_{\varepsilon, k, h}^-(t), \Pi_{\varepsilon, k, h}^-(t)) &:= (V_\varepsilon^{m-1}, \Pi_\varepsilon^{m-1}), \quad \forall t \in [t_{m-1}, t_m).
\end{aligned}$$

There will also be similar notations in the upcoming part such as f^+ and r^- ; the reader may refer to section 2.2 for an adequate definition. Note that it is not mandatory for ε to be dependent on the discretization parameters k and h . If so, it suffices that $\varepsilon = \varepsilon(k, h) \rightarrow 0$ as $k, h \rightarrow 0$.

For instance, the penalty parameter ε may be linked to the time discretization step size k in a way that k/ε tends to 0 when both k, ε vanish. This idea is exposed in the below proposition, but will not be utilized for the convergence analysis of Algorithm 1.

Proposition 4.1 *Let $\{(V_\varepsilon^m, \Pi_\varepsilon^m)\}_{m=1}^M$ be the iterates of Algorithm 1. Then,*

$$\mathbb{E} \left[\sum_{m=1}^M \varepsilon \|\Pi_\varepsilon^m - \Pi_\varepsilon^{m-1}\|_{L^2}^2 \right] \leq 2C_1 \frac{k}{\varepsilon},$$

where C_1 appears in Lemma 4.3.

Proof: Let $q \in L^2(D) \setminus \{0\}$. By identity (2.4), it holds that $(\Pi_\varepsilon^m - \Pi_\varepsilon^{m-1}, q) = (\Pi_\varepsilon^m - \Pi_\varepsilon^{m-1}, \rho_h q)$. Therefore, using Algorithm 1, one obtains

$$\varepsilon (\Pi_\varepsilon^m - \Pi_\varepsilon^{m-1}, q) = -k(\operatorname{div}(V_\varepsilon^m), \rho_h q) \leq k\sqrt{2} \|\nabla V_\varepsilon^m\|_{\mathbb{L}^2} \|q\|_{L^2}, \quad (4.11)$$

thanks to the Cauchy-Schwarz inequality, the stability of ρ_h in $L^2(D)$ and the estimate $\|div \cdot\|_{L^2} \leq \sqrt{d} \|\nabla \cdot\|_{\mathbb{L}^2}$, with $d = 2$ is the dimension. Therefore,

$$\sup_{q \in L^2(D) \setminus \{0\}} \frac{\varepsilon (\Pi_\varepsilon^m - \Pi_\varepsilon^{m-1}, q)}{\|q\|_{L^2}} \leq k\sqrt{2} \|\nabla V_\varepsilon^m\|_{\mathbb{L}^2}.$$

Since $L^2(D)$ is the pivot space, the supremum in the above equation turns into $\varepsilon \|\Pi_\varepsilon^m - \Pi_\varepsilon^{m-1}\|_{L^2}$. Therefore, squaring both sides of the above equation, taking the sum over m from 1 to M , then applying the mathematical expectation return the following

$$\mathbb{E} \left[\sum_{m=1}^M \varepsilon \|\Pi_\varepsilon^m - \Pi_\varepsilon^{m-1}\|_{L^2}^2 \right] \leq \frac{2k}{\varepsilon} \mathbb{E} \left[k \sum_{m=1}^M \|\nabla V_\varepsilon^m\|_{\mathbb{L}^2}^2 \right].$$

Finally, a simple application of Lemma 4.3-(i) completes the proof. \square

Going back to the convergence demonstration of Algorithm 1, the following lemma states a monotonicity property of the operator $u \mapsto -\nu \Delta u + \hat{B}(u, u)$. This feature together with the Lipschitz-continuity of the diffusion coefficient g allow the avoidance of the Skorokhod theorem that forces the filtered probability space, which was defined in Section 2, to be exchanged with a new one.

Lemma 4.4 *Assume that $L_g \leq \sqrt{\frac{\nu}{2C_P^2}}$ where $C_P > 0$ is the Poincaré constant, and let $u, w \in \mathbb{H}_0^1$. For $z := u - w$, the following inequality holds true:*

$$\left\langle -\nu \Delta z + \hat{B}(u, u) - \hat{B}(w, w) + \frac{27}{2\nu^3} \|w\|_{\mathbb{L}^4}^4 z, z \right\rangle - \|g(u) - g(w)\|_{\mathcal{L}_2(K, \mathbb{L}^2)}^2 \geq 0.$$

Proof: From [17, Lemma 2.4], it holds that

$$\left\langle -\nu \Delta z + \hat{B}(u, u) - \hat{B}(w, w) + \frac{27}{2\nu^3} \|w\|_{\mathbb{L}^4}^4 z, z \right\rangle \geq \frac{\nu}{2} \|\nabla z\|_{\mathbb{L}^2}^2.$$

It suffices now to subtract from both sides the term $\|g(u) - g(w)\|_{\mathcal{L}_2(K, \mathbb{L}^2)}^2$, use assumption (S_2) , then employ the Poincaré inequality. \square

Besides Lemma 4.4, it is worth highlighting the strong convergence of $\{g(\mathcal{V}_{\varepsilon, k, h}^+) - g(\mathcal{V}_{\varepsilon, k, h}^-)\}_{k, h}$ in $L^2(\Omega; L^2(0, T; \mathcal{L}_2(K, \mathbb{L}^2)))$, which can be illustrated through assumption (S_2) and Lemma 4.3-(i) as follows

$$\begin{aligned} \mathbb{E} \left[\int_0^T \|g(\mathcal{V}_{\varepsilon, k, h}^+) - g(\mathcal{V}_{\varepsilon, k, h}^-)\|_{\mathcal{L}_2(K, \mathbb{L}^2)}^2 dt \right] &\leq L_g^2 k \mathbb{E} \left[\sum_{m=1}^M \|V_\varepsilon^m - V_\varepsilon^{m-1}\|_{\mathbb{L}^2}^2 \right] \\ &\leq L_g^2 C_1 k \rightarrow 0. \end{aligned} \quad (4.12)$$

The convergence demonstration down below is broken down into steps for clarity's sake.

Step1: Weak convergence and divergence-free

By virtue of Lemma 4.3, the sublinearity of g (see assumption (S_2)) and inequality (2.5), the sequences $\{\mathcal{V}_{\varepsilon, k, h}^+\}_{\varepsilon, k, h}$, $\{\sqrt{\varepsilon} \Pi_{\varepsilon, k, h}^+\}_{\varepsilon, k, h}$, $\{g(\mathcal{V}_{\varepsilon, k, h}^-)\}_{\varepsilon, k, h}$ are bounded in the Banach spaces $L^2(\Omega; L^\infty(0, T; \mathbb{L}^2) \cap L^2(0, T; \mathbb{H}_0^1))$, $L^2(\Omega; L^\infty(0, T; L^2(D)))$ and $L^2(\Omega; L^2(0, T; \mathcal{L}_2(K, \mathbb{L}^2)))$, respectively. Therefore, the Banach-Alaoglu theorem ensures the existence of the limiting functions $v \in L^2(\Omega; L^\infty(0, T; \mathbb{L}^2) \cap L^2(0, T; \mathbb{H}_0^1))$, $\chi \in L^2(\Omega; L^\infty(0, T; L^2(D)))$, $G_0 \in L^2(\Omega; L^2(0, T; \mathcal{L}_2(K, \mathbb{L}^2)))$ and two subsequences (still denoted as their original sequences) $\{\mathcal{V}_{\varepsilon, k, h}^+\}_{\varepsilon, k, h}$, $\{\sqrt{\varepsilon} \Pi_{\varepsilon, k, h}^+\}_{\varepsilon, k, h}$ such that

$$\mathcal{V}_{\varepsilon, k, h}^+ \overset{*}{\rightharpoonup} v \quad \text{in } L^2(\Omega; L^\infty(0, T; \mathbb{L}^2)), \quad (4.13)$$

$$\mathcal{V}_{\varepsilon,k,h}^+ \rightharpoonup v \quad \text{in } L^2(\Omega; L^2(0, T; \mathbb{H}_0^1)), \quad (4.14)$$

$$\sqrt{\varepsilon} \Pi_{\varepsilon,k,h}^+ \xrightarrow{*} \chi \quad \text{in } L^2(\Omega; L^\infty(0, T; L^2(D))), \quad (4.15)$$

$$g(\mathcal{V}_{\varepsilon,k,h}^-) \rightharpoonup G_0 \quad \text{in } L^2(\Omega; L^2(0, T; \mathcal{L}_2(K, \mathbb{L}^2))). \quad (4.16)$$

Besides convergence (4.16), it is also possible to acquire $g(\mathcal{V}_{\varepsilon,k,h}^+) \rightharpoonup G_0$ in $L^2(\Omega; L^2(0, T; \mathcal{L}_2(K, \mathbb{L}^2)))$ as follows: for all $\phi \in L^2(\Omega; L^2(0, T; \mathcal{L}_2(K, \mathbb{L}^2)))$,

$$\begin{aligned} \left(g(\mathcal{V}_{\varepsilon,k,h}^+) - G_0(t), \phi(t) \right)_{\mathcal{L}_2(K, \mathbb{L}^2)} &= \left(g(\mathcal{V}_{\varepsilon,k,h}^+) - g(\mathcal{V}_{\varepsilon,k,h}^-), \phi(t) \right)_{\mathcal{L}_2(K, \mathbb{L}^2)} \\ &+ \left(g(\mathcal{V}_{\varepsilon,k,h}^-) - G_0(t), \phi(t) \right)_{\mathcal{L}_2(K, \mathbb{L}^2)}. \end{aligned} \quad (4.17)$$

Now, integrate with respect to t , take the mathematical expectation, use results (4.12) and (4.16) to complete the proof.

The obtained function v is divergence-free. Indeed, let $q \in C_c^\infty(D)$ be a scalar function. From Algorithm 1, one has $\varepsilon (\Pi_\varepsilon^m - \Pi_\varepsilon^{m-1}, \rho_h q) = -k (\operatorname{div}(V_\varepsilon^m), \rho_h q)$. Summing both sides over m from 1 to M leads to $\int_0^T (\operatorname{div}(\mathcal{V}_{\varepsilon,k,h}^+), \rho_h q) dt = \varepsilon (\Pi_\varepsilon^0, \rho_h q) - \sqrt{\varepsilon} (\sqrt{\varepsilon} \Pi_{\varepsilon,k,h}^+(T), \rho_h q)$. The mathematical expectation of the right-hand side goes to 0 as $\varepsilon, k, h \rightarrow 0$ due to convergence (4.15) and estimate (2.5). Hence,

$$\mathbb{E} \left[\int_0^T (\operatorname{div}(\mathcal{V}_{\varepsilon,k,h}^+), q) dt \right] = \mathbb{E} \left[\int_0^T (\operatorname{div}(\mathcal{V}_{\varepsilon,k,h}^+), q - \rho_h q) dt \right] + \mathbb{E} \left[\int_0^T (\operatorname{div}(\mathcal{V}_{\varepsilon,k,h}^+), \rho_h q) dt \right]$$

converges to 0 as $\varepsilon, k, h \rightarrow 0$, thanks to estimate (2.3) and convergence $\operatorname{div}(\mathcal{V}_{\varepsilon,k,h}^+) \rightarrow \operatorname{div}(v)$ in $L^2(\Omega; L^2(0, T; L^2(D)))$ which follows straightforwardly from result (4.14). Subsequently, $\operatorname{div}(\mathcal{V}_{\varepsilon,k,h}^+) \rightharpoonup 0$ in $L^2(\Omega; L^2(0, T; L^2(D)))$ which implies $\operatorname{div}(v) = 0$ \mathbb{P} -a.s. and a.e. in $(0, T) \times D$.

Let $\mathcal{R}: \mathbb{H}_0^1 \rightarrow \mathbb{H}^{-1}$ be defined by $\mathcal{R}(u) := -\nu \Delta u + \hat{B}(u, u)$, for all $u \in \mathbb{H}_0^1$. From Algorithm 1, and for all $\varphi \in \mathcal{V}$ such that $\varphi_h := \Pi_h \varphi$, it follows

$$\begin{aligned} \int_0^T \langle \mathcal{R}(\mathcal{V}_{\varepsilon,k,h}^+) + \nabla \Pi_{\varepsilon,k,h}^+, \varphi_h \rangle dt &= - \left(\mathcal{V}_{\varepsilon,k,h}^+(T) - \mathcal{V}_{\varepsilon,k,h}^-(0), \varphi_h \right) + \int_0^T \langle f^+, \varphi_h \rangle dt \\ &+ \left(\int_0^T g(\mathcal{V}_{\varepsilon,k,h}^-) dW(t), \varphi_h \right). \end{aligned} \quad (4.18)$$

Owing to results (4.13) and (4.16) along with the strong convergence of f^+ in $L^2(\Omega; L^2(0, T; \mathbb{H}^{-1}))$ (see [25, Lemma III.4.9]), the mathematical expectation of the right-hand side of equation (4.18) is convergent. Therewith, define \mathcal{R}_0 by

$$\mathbb{E} \left[\int_0^T \langle \mathcal{R}_0(t), \varphi \rangle dt \right] = \lim_{\varepsilon, k, h \rightarrow 0} \mathbb{E} \left[\int_0^T \langle \mathcal{R}(\mathcal{V}_{\varepsilon,k,h}^+) + \nabla \Pi_{\varepsilon,k,h}^+, \Pi_h \varphi \rangle dt \right], \quad \forall \varphi \in \mathcal{V}.$$

Subsequently, the limiting function v satisfies \mathbb{P} -a.s. and for all $(t, \varphi) \in [0, T] \times \mathcal{V}$ the following:

$$(v(t) - v_0, \varphi) + \int_0^t \langle \mathcal{R}_0(s), \varphi \rangle ds = \int_0^t \langle f(s), \varphi \rangle ds + \left(\int_0^t G_0(s) dW(s), \varphi \right). \quad (4.19)$$

Step 2: Identification of \mathcal{R}_0 and G_0

For $\sigma \in C([0, T], \mathcal{V})$, define the finite element space of weakly divergence-free functions:

$$\mathbb{V}_h := \{ u_h \in \mathbb{H}_h \mid (\operatorname{div}(u_h), q_h) = 0, \forall q_h \in L_h \}.$$

Let $P_h: \mathbb{L}^2 \rightarrow \mathbb{V}_h$ be the projection operator from \mathbb{L}^2 onto \mathbb{V}_h such that for all $u \in \mathbb{L}^2$,

$$(P_h u, \varphi_h) = (u, \varphi_h), \quad \forall \varphi_h \in \mathbb{V}_h.$$

The space \mathbb{V}_h is required because the function σ that was introduced shortly before is divergence-free and shall be projected onto a finite element space that possesses a null divergence constraint. This will become clearer when the discrete pressure $\{\Pi_\varepsilon^m\}_{m=1}^M$ is dealt with in the sequel, especially because the limit of iterates $\{(V_\varepsilon^m, \Pi_\varepsilon^m)\}_{m=1}^M$ takes place in divergence-free spaces, as shown earlier for v . It is worth pointing out that the finite element space \mathbb{V}_h does not interact with $\{V_\varepsilon^m\}_{m=1}^M$ and $\{\Pi_\varepsilon^m\}_{m=1}^M$. In other words, the sequence $\{V_\varepsilon^m\}_{m=1}^M$ will remain non divergence-free and will never belong to \mathbb{V}_h . Now for all $m \in \{1, \dots, M\}$, denote $\sigma_h^+(t) := \sigma_h^m = P_h \sigma(t_m)$ and define $r^+(t) := r^m := \frac{27}{\nu^3} k \sum_{n=1}^m \|\sigma_h^n\|_{\mathbb{L}^4}^4$ for all $t \in (t_{m-1}, t_m]$, together with an exponential non-increasing function $\eta: [0, T] \rightarrow \mathbb{R}$ verifying $\eta(0) = 0$, and having the discrete forms $\eta^+(t) := \eta^m := e^{-r^+(t)}$ for all $t \in (t_{m-1}, t_m]$ and $\eta^-(t) := \eta^{m-1}$ for all $t \in [t_{m-1}, t_m)$. Setting $(\varphi_h, q_h) = (V_\varepsilon^m, \Pi_\varepsilon^m)$ in Algorithm 1, using Cauchy-Schwarz and Young's inequalities, identity $(a - b, a) = \frac{1}{2} \|a\|_{\mathbb{L}^2}^2 - \frac{1}{2} \|b\|_{\mathbb{L}^2}^2 + \frac{1}{2} \|a - b\|_{\mathbb{L}^2}^2$, and finally multiplying by η^{m-1} yield

$$\begin{aligned} & \eta^{m-1} (\|V_\varepsilon^m\|_{\mathbb{L}^2}^2 - \|V_\varepsilon^{m-1}\|_{\mathbb{L}^2}^2) + 2\eta^{m-1} k \langle \mathcal{R}(V_\varepsilon^m) + \nabla \Pi_\varepsilon^m, V_\varepsilon^m \rangle \leq 2\eta^{m-1} k \langle f^m, V_\varepsilon^m \rangle \\ & + \eta^{m-1} \|g(V_\varepsilon^{m-1}) \Delta_m W\|_{\mathbb{L}^2}^2 + 2\eta^{m-1} (g(V_\varepsilon^{m-1}) \Delta_m W, V_\varepsilon^{m-1}). \end{aligned} \quad (4.20)$$

Note that $\sum_{m=1}^M \eta^{m-1} (\|V_\varepsilon^m\|_{\mathbb{L}^2}^2 - \|V_\varepsilon^{m-1}\|_{\mathbb{L}^2}^2) = \int_0^T \eta^-(t) dt \|\mathcal{V}_{\varepsilon, k, h}^+\|_{\mathbb{L}^2}^2 dt$, and through equation (4.6), it holds that $\mathbb{E} [\|g(V_\varepsilon^{m-1}) \Delta_m W\|_{\mathbb{L}^2}^2] = k \mathbb{E} [\|g(V_\varepsilon^{m-1})\|_{\mathcal{L}_2(K, \mathbb{L}^2)}^2]$. Therefore, taking the sum over m from 1 to M , employing Proposition 2.2-(ii), then applying the mathematical expectation to equation (4.20) give

$$\begin{aligned} & \mathbb{E} \left[\eta^+(T) \|\mathcal{V}_{\varepsilon, k, h}^+(T)\|_{\mathbb{L}^2}^2 - \|\mathcal{V}_{\varepsilon, k, h}^-(0)\|_{\mathbb{L}^2}^2 \right] \leq \mathbb{E} \left[\int_0^T \|\mathcal{V}_{\varepsilon, k, h}^+\|_{\mathbb{L}^2}^2 dt \eta^+ dt \right] \\ & - \mathbb{E} \left[\int_0^T \eta^-(t) \langle 2\mathcal{R}(\mathcal{V}_{\varepsilon, k, h}^+) + 2\nabla \Pi_{\varepsilon, k, h}^+, \mathcal{V}_{\varepsilon, k, h}^+ \rangle dt \right] + \mathbb{E} \left[2 \int_0^T \eta^-(t) \langle f^+, \mathcal{V}_{\varepsilon, k, h}^+ \rangle dt \right] \\ & + \mathbb{E} \left[\int_0^T \eta^-(t) \|g(\mathcal{V}_{\varepsilon, k, h}^-)\|_{\mathcal{L}_2(K, \mathbb{L}^2)}^2 dt \right] =: I + II + III + IV, \end{aligned} \quad (4.21)$$

where the last term on the right-hand side of equation (4.20) vanishes after taking its expectation due to assumption (S_2) and the measurability of $\{V_\varepsilon^m\}_m$ (see Lemma 4.1). By virtue of Proposition 2.2-(iii), it follows that $d_t \eta^+ = -\frac{27}{\nu^3} \eta^- \|\sigma_h^+\|_{\mathbb{L}^4}^4 + \frac{27^2 k}{2\nu^6} e^{\delta(t)} \|\sigma_h^+\|_{\mathbb{L}^4}^8$, for some $\delta \in (-r^+, -r^-)$. Therefore,

$$I = -\mathbb{E} \left[\int_0^T \eta^-(t) \frac{27}{\nu^3} \|\sigma_h^+\|_{\mathbb{L}^4}^4 \|\mathcal{V}_{\varepsilon, k, h}^+\|_{\mathbb{L}^2}^2 dt \right] + \frac{27^2}{2\nu^6} k \mathbb{E} \left[\int_0^T \|\mathcal{V}_{\varepsilon, k, h}^+\|_{\mathbb{L}^2}^2 e^{\delta(t)} \|\sigma_h^+\|_{\mathbb{L}^4}^8 dt \right] =: I_1 + I_2.$$

Obviously, I_2 goes to 0 as $k, h, \varepsilon \rightarrow 0$ thanks to Lemma 4.3. I_1 can be rewritten as follows

$$\begin{aligned} I_1 &= -\frac{27}{\nu^3} \mathbb{E} \left[\int_0^T \eta^- \|\sigma_h^+\|_{\mathbb{L}^4}^4 \|\mathcal{V}_{\varepsilon, k, h}^+ - \sigma_h^+\|_{\mathbb{L}^2}^2 dt \right] \\ & - \frac{27}{\nu^3} \mathbb{E} \left[\int_0^T \eta^- \|\sigma_h^+\|_{\mathbb{L}^4}^4 \left\{ 2 \left(\mathcal{V}_{\varepsilon, k, h}^+, \sigma_h^+ \right) - \|\sigma_h^+\|_{\mathbb{L}^2}^2 \right\} dt \right] =: I_{1,1} + I_{1,2}. \end{aligned}$$

Making use of result (4.14) along with the strong convergence of $\{\sigma_h^m\}_m$ to σ in $C([0, T]; \mathbb{V})$, it can be easily shown that $II_{1,2} \rightarrow -\frac{27}{\nu^3} \mathbb{E} \left[\int_0^T \eta(t) \|\sigma(t)\|_{\mathbb{L}^4}^4 \left\{ 2 \langle v(t), \sigma(t) \rangle - \|\sigma(t)\|_{\mathbb{L}^2}^2 \right\} dt \right]$. On the other hand,

$$\begin{aligned} II &= -\mathbb{E} \left[\int_0^T \eta^- \langle 2\mathcal{R}(\mathcal{V}_{\varepsilon,k,h}^+) - 2\mathcal{R}(\sigma_h^+), \mathcal{V}_{\varepsilon,k,h}^+ - \sigma_h^+ \rangle dt \right] - \mathbb{E} \left[\int_0^T \eta^- \langle 2\nabla \Pi_{\varepsilon,k,h}^+, \mathcal{V}_{\varepsilon,k,h}^+ - \sigma_h^+ \rangle dt \right] \\ &\quad - \mathbb{E} \left[\int_0^T \eta^- \langle 2\mathcal{R}(\mathcal{V}_{\varepsilon,k,h}^+) + 2\nabla \Pi_{\varepsilon,k,h}^+ - 2\mathcal{R}(\sigma_h^+), \sigma_h^+ \rangle dt \right] - \mathbb{E} \left[\int_0^T \eta^- \langle 2\mathcal{R}(\sigma_h^+), \mathcal{V}_{\varepsilon,k,h}^+ \rangle dt \right] \\ &=: II_1 + II_2 + II_3 + II_4. \end{aligned}$$

By an integration by parts, II_2 can be rewritten as follows:

$$II_2 = 2\mathbb{E} \left[\int_0^T \eta^-(t) \left(\Pi_{\varepsilon,k,h}^+, \operatorname{div}(\mathcal{V}_{\varepsilon,k,h}^+) \right) dt \right] = 2\mathbb{E} \left[k \sum_{m=1}^M \eta^{m-1} \left(\Pi_{\varepsilon}^m, \operatorname{div}(\mathcal{V}_{\varepsilon}^m) \right) \right],$$

because $\sigma_h^+ \in \mathbb{V}_h$ i.e. $(\Pi_{\varepsilon,k,h}, \operatorname{div}(\sigma_h^+)) = 0$. Therefore, making use of Algorithm 1 yields

$$\begin{aligned} II_2 &= -2\varepsilon \mathbb{E} \left[\sum_{m=1}^M \eta^{m-1} \left(\Pi_{\varepsilon}^m - \Pi_{\varepsilon}^{m-1}, \Pi_{\varepsilon}^m \right) \right] \\ &= -\varepsilon \mathbb{E} \left[\sum_{m=1}^M \eta^{m-1} \left(\|\Pi_{\varepsilon}^m\|_{L^2}^2 - \|\Pi_{\varepsilon}^{m-1}\|_{L^2}^2 + \|\Pi_{\varepsilon}^m - \Pi_{\varepsilon}^{m-1}\|_{L^2}^2 \right) \right] \\ &= -\varepsilon \mathbb{E} \left[\int_0^T \eta^- dt \|\Pi_{\varepsilon,k,h}^+\|_{L^2}^2 \right] - \varepsilon \mathbb{E} \left[\sum_{m=1}^M \eta^{m-1} \|\Pi_{\varepsilon}^m - \Pi_{\varepsilon}^{m-1}\|_{L^2}^2 \right] =: II_{2,1} + II_{2,2}, \end{aligned}$$

thanks to the identity $(a - b, a) = \frac{1}{2} (\|a\|_{L^2}^2 - \|b\|_{L^2}^2 + \|a - b\|_{L^2}^2)$. Observe first that $II_{2,2} \leq 0$, which is true because $\{\eta^m\}_m$ is a nonnegative sequence. Moreover, by Proposition 2.2-(ii), it holds that

$$\begin{aligned} II_{2,1} &= \varepsilon \mathbb{E} \left[\int_0^T \|\Pi_{\varepsilon,k,h}^+\|_{L^2}^2 dt \eta^+(t) \right] - \varepsilon \mathbb{E} \left[\eta^+(T) \|\Pi_{\varepsilon,k,h}^+(T)\|_{L^2}^2 - \|\Pi_{\varepsilon,k,h}^-(0)\|_{L^2}^2 \right] \\ &\leq \varepsilon \mathbb{E} \left[\int_0^T \|\Pi_{\varepsilon,k,h}^+\|_{L^2}^2 dt \eta^+(t) \right] + \varepsilon \mathbb{E} \left[\|\Pi_{\varepsilon,k,h}^-(0)\|_{L^2}^2 \right] =: II_{2,1,1} + II_{2,1,2}. \end{aligned}$$

Proposition 2.2-(iii) and Lemma 4.3-(ii) imply

$$\begin{aligned} II_{2,1,1} &= \varepsilon \mathbb{E} \left[\int_0^T \|\Pi_{\varepsilon,k,h}^+\|_{L^2}^2 \left(-\frac{27}{\nu^3} \eta^- \|\sigma_h^+\|_{\mathbb{L}^4}^4 + \frac{27^2 k}{2\nu^6} e^{\delta(t)} \|\sigma_h^+\|_{\mathbb{L}^4}^8 \right) dt \right] \\ &\leq \frac{27^2}{2\nu^6} \|\sigma\|_{L^\infty(0,T;\mathbb{V})}^8 \mathbb{E} \left[\int_0^T \varepsilon \|\Pi_{\varepsilon,k,h}^+\|_{L^2}^2 \right] k \leq \frac{27^2 T}{2\nu^6} \|\sigma\|_{L^\infty(0,T;\mathbb{V})}^8 \mathbb{E} \left[\varepsilon \max_{1 \leq m \leq M} \|\Pi_{\varepsilon}^m\|_{L^2}^2 \right] k \\ &\leq \frac{27^2 T C_1}{2\nu^6} \|\sigma\|_{L^\infty(0,T;\mathbb{V})}^8 k \rightarrow 0 \text{ as } k \rightarrow 0, \end{aligned}$$

for some $\delta(t) \in (-r^+(t), -r^-(t))$. Furthermore, since the projector ρ_h is stable in $L^2(D)$, it follows that

$$II_{2,1,2} = \varepsilon \mathbb{E} \left[\|\rho_h^0\|_{L^2}^2 \right] \leq \varepsilon \mathbb{E} \left[\|\pi_0\|_{L^2}^2 \right] \rightarrow 0 \text{ as } \varepsilon \rightarrow 0.$$

Moreover, since $\{\sigma_h^m\}_m$ is strongly convergent toward σ in $C([0, T]; \mathbb{V})$, and by the definition of operator \mathcal{R}_0 , one obtains $II_3 \rightarrow -\mathbb{E} \left[\int_0^T \eta(t) \langle 2\mathcal{R}_0(t) - 2\mathcal{R}(\sigma(t)), \sigma(t) \rangle dt \right]$ as $k, h, \varepsilon \rightarrow 0$. Similarly, $II_4 \rightarrow -\mathbb{E} \left[\int_0^T \eta(t) \langle 2\mathcal{R}(\sigma(t)), v(t) \rangle dt \right]$, thanks to convergence (4.14). As mentioned in Step 1, $\{f^m\}_m$

converges strongly toward f in $L^2(\Omega; L^2(0, T; \mathbb{H}^{-1}))$. The latter together with convergence (4.14) imply that $III \rightarrow \mathbb{E} \left[2 \int_0^T \eta(t) \langle f(t), v(t) \rangle dt \right]$. Moving on to term IV , it can be reformulated as follows:

$$\begin{aligned} IV &= \mathbb{E} \left[\int_0^T \eta^-(t) \left\{ \|g(\mathcal{V}_{\varepsilon, k, h}^-) - g(\mathcal{V}_{\varepsilon, k, h}^+) \|_{\mathcal{L}_2(K, \mathbb{L}^2)}^2 + \|g(\mathcal{V}_{\varepsilon, k, h}^+) - g(\sigma_h^+) \|_{\mathcal{L}_2(K, \mathbb{L}^2)}^2 \right. \right. \\ &\quad - \|g(\sigma_h^+) \|_{\mathcal{L}_2(K, \mathbb{L}^2)}^2 + 2 \left(g(\mathcal{V}_{\varepsilon, k, h}^+), g(\sigma_h^+) \right)_{\mathcal{L}_2(K, \mathbb{L}^2)} \\ &\quad \left. \left. + 2 \left(g(\mathcal{V}_{\varepsilon, k, h}^-) - g(\mathcal{V}_{\varepsilon, k, h}^+), g(\mathcal{V}_{\varepsilon, k, h}^+) \right)_{\mathcal{L}_2(K, \mathbb{L}^2)} \right\} dt \right] := IV_1 + \dots + IV_5. \end{aligned}$$

From equation (4.12), it holds that $IV_1 \rightarrow 0$. Furthermore, Lemma 4.4 yields $I_{1,1} + II_1 + IV_2 \leq 0$, the strong convergence of $\{\sigma_h^m\}_m$ together with result (4.17) grant $IV_3 \rightarrow -\mathbb{E} \left[\int_0^T \eta(t) \|g(\sigma(t)) \|_{\mathcal{L}_2(K, \mathbb{L}^2)}^2 dt \right]$ and $IV_4 \rightarrow \mathbb{E} \left[2 \int_0^T \eta(t) (G_0(t), g(\sigma(t)))_{\mathcal{L}_2(K, \mathbb{L}^2)} dt \right]$. Finally, $IV_5 \rightarrow 0$ by virtue of convergences (4.12) and (4.17). Putting it all together, equation (4.21) becomes

$$\begin{aligned} \lim_{\varepsilon, k, h \rightarrow 0} \mathbb{E} \left[\eta^+(T) \| \mathcal{V}_{\varepsilon, k, h}^+(T) \|_{\mathbb{L}^2}^2 - \| \mathcal{V}_{\varepsilon, k, h}^-(0) \|_{\mathbb{L}^2}^2 \right] &\leq \mathbb{E} \left[\int_0^T \eta'(t) \left\{ 2(v, \sigma) - \|\sigma\|_{\mathbb{L}^2}^2 \right\} dt \right] \\ &- 2\mathbb{E} \left[\int_0^T \eta(t) \left\{ \langle \mathcal{R}_0 - \mathcal{R}(\sigma), \sigma \rangle + \langle \mathcal{R}(\sigma) - f, v \rangle + \frac{1}{2} \|g(\sigma)\|_{\mathcal{L}_2(K, \mathbb{L}^2)}^2 \right. \right. \\ &\quad \left. \left. - (G_0, g(\sigma))_{\mathcal{L}_2(K, \mathbb{L}^2)} \right\} dt \right], \end{aligned} \quad (4.22)$$

where $\eta(t) = \exp\left(-\frac{27}{\nu^3} \int_0^t \|\sigma(s)\|_{\mathbb{L}^4}^4 ds\right)$. Taking into account that $\mathbb{E} [\eta(T) \|v(T)\|_{\mathbb{L}^2}^2 - \|v_0\|_{\mathbb{L}^2}^2]$ is smaller than the left-hand side of equation (4.22) (thanks to result (4.13)), and applying Itô's formula to the process $(t, v) \mapsto \eta(t) \|v\|_{\mathbb{L}^2}^2$ (recall that v satisfies equation (4.19)) lead to

$$\begin{aligned} &\mathbb{E} \left[\int_0^T \eta'(t) \|v(t) - \sigma(t)\|_{\mathbb{L}^2}^2 dt \right] + \mathbb{E} \left[\int_0^T \eta(t) \|G_0(t) - g(\sigma(t))\|_{\mathcal{L}_2(K, \mathbb{L}^2)}^2 dt \right] \\ &\leq 2\mathbb{E} \left[\int_0^T \eta(t) \langle \mathcal{R}(\sigma(t)) - \mathcal{R}_0(t), \sigma(t) - v(t) \rangle dt \right], \quad \forall \sigma \in C([0, T]; \mathcal{V}). \end{aligned} \quad (4.23)$$

Arguing by density, it can be shown that inequality (4.23) holds for all $\sigma \in L^4(\Omega; L^\infty(0, T; \mathbb{H}) \cap L^2(\Omega; L^2(0, T; \mathbb{V}))$. Therefore, setting $\sigma = v$ yields $G_0 = g(v)$ \mathbb{P} -a.s. and a.e. in $[0, T] \times D$. With that said, the second term on the left-hand side of equation (4.23) cancels out. To identify \mathcal{R}_0 , it suffices to consider $\sigma = v + \mu u$ for $\mu > 0$ and $u \in L^4(\Omega; L^\infty(0, T; \mathbb{H})) \cap L^2(\Omega; L^2(0, T; \mathbb{V}))$. Subsequently,

$$\mu \mathbb{E} \left[\int_0^T \eta'(t) \|u(t)\|_{\mathbb{L}^2}^2 dt \right] \leq 2\mathbb{E} \left[\int_0^T \eta(t) \langle \mathcal{R}(v(t) + \mu u(t)) - \mathcal{R}_0(t), u(t) \rangle dt \right].$$

Letting $\mu \rightarrow 0$ and taking into consideration the hemicontinuity of the operator \mathcal{R} , one infers that $\mathbb{E} \left[\int_0^T \eta(t) \langle \mathcal{R}(v(t)) - \mathcal{R}_0(t), u(t) \rangle dt \right] \geq 0$, for all $u \in L^4(\Omega; L^\infty(0, T; \mathbb{H})) \cap L^2(\Omega; L^2(0, T; \mathbb{V}))$. Consequently, $\mathcal{R}_0 = \mathcal{R}(v)$ in $L^2(\Omega; L^2(0, T; \mathbb{H}^{-1}))$.

Step 3: Verification of v as NSE solution

The obtained function v is henceforth a solution to equations (1.1) in the sense of Definition 2.2. Indeed, the identifications in Step 2 turn equation (4.19) into

$$\begin{aligned} &(v(t), \varphi) + \nu \int_0^t (\nabla v(s), \nabla \varphi) ds + \int_0^t \langle \hat{B}(v(t), v(t)), \varphi \rangle ds \\ &= (v_0, \varphi) + \int_0^t \langle f(s), \varphi \rangle ds + \left(\int_0^t g(v(s)) dW(s), \varphi \right), \quad \forall \varphi \in \mathbb{V}. \end{aligned}$$

By definition, $\hat{B}(v, v) = ([v \cdot \nabla] + \frac{1}{2} \operatorname{div}(v))v = [v \cdot \nabla]v$, thanks to Step 2, where the null divergence of v was illustrated. Finally, $v \in L^2(\Omega; C([0, T]; \mathbb{L}^2))$ can be easily proven via equation (4.19) by using the standard approach in [20].

Step 4: Convergence of the whole sequence

Convergence results that were discovered within Step 1 are all up to a subsequence. However, due to the uniqueness of v (see [18, Proposition 3.2]), it follows that the whole sequence $\{\mathcal{V}_{\varepsilon, k, h}^+\}_{\varepsilon, k, h}$ is convergent toward v .

4.3 A linear version of Algorithm 1

In terms of simulations, a less time-consuming numerical scheme can be embodied through a linear Algorithm. This can be made up using a linearization of the trilinear term in Algorithm 1 as follows:

Algorithm 2 Starting from an initial datum $(v_h^0, p_h^0) \in \mathbb{H}_h \times L_h$, if $(V_\varepsilon^{m-1}, \Pi_\varepsilon^{m-1}) \in \mathbb{H}_h \times L_h$ is known for some $m \in \{1, \dots, M\}$, find $(V_\varepsilon^m, \Pi_\varepsilon^m) \in \mathbb{H}_h \times L_h$ that satisfies \mathbb{P} -a.s. the following:

$$\begin{cases} (V_\varepsilon^m - V_\varepsilon^{m-1}, \varphi_h) + k\nu(\nabla V_\varepsilon^m, \nabla \varphi) + k\hat{b}(V_\varepsilon^{m-1}, V_\varepsilon^m, \varphi_h) - (\Pi_\varepsilon^m, \operatorname{div}(\varphi_h)) \\ = k \langle f^m, \varphi_h \rangle + (g(V_\varepsilon^{m-1})\Delta_m W, \varphi_h), \quad \forall \varphi \in \mathbb{H}_h, \\ \frac{\varepsilon}{k} (\Pi_\varepsilon^m - \Pi_\varepsilon^{m-1}, q_h) + (\operatorname{div}(V_\varepsilon^m), q_h) = 0, \quad \forall q_h \in L_h, \end{cases}$$

where $f^m, \Delta_m W$ are defined in Algorithm 1 and $(V_\varepsilon^0, \Pi_\varepsilon^0) := (v_h^0, p_h^0)$.

Observe that $\hat{b}(V_\varepsilon^{m-1}, V_\varepsilon^m, V_\varepsilon^m) = 0$, thanks to Proposition 2.1-(ii). Therefore, iterates $\{(V_\varepsilon^m, \Pi_\varepsilon^m)\}_{m=1}^M$ of Algorithm 2 satisfy Lemmas 4.1, 4.3 and they fulfill better uniqueness properties than those of Algorithm 1, as demonstrated in Lemma 4.5. However, due to the infamous properties of \hat{b} , the initial datum v_h^0 should undergo a new assumption that consists of a uniform bound in h of $\|\nabla v_h^0\|_{\mathbb{L}^2}$, as explained beneath the proof of Lemma 4.5.

Lemma 4.5 Iterates $\{(V_\varepsilon^m, \Pi_\varepsilon^m)\}_{m=1}^M$ of Algorithm 2 are unique \mathbb{P} -a.s. in Ω and a.e. in $[0, T] \times D$.

Proof: Let $\{(V_\varepsilon^m, \Pi_\varepsilon^m)\}_{m=1}^M$ and $\{(U_\varepsilon^m, P_\varepsilon^m)\}_{m=1}^M$ be two solutions to Algorithm 2 such that $(V_\varepsilon^0, \Pi_\varepsilon^0) = (U_\varepsilon^0, P_\varepsilon^0) = (v_h^0, p_h^0)$. Denote $Z_\varepsilon^m := V_\varepsilon^m - U_\varepsilon^m$ and $Q_\varepsilon^m := \Pi_\varepsilon^m - P_\varepsilon^m$, for all $m \in \{0, 1, \dots, M\}$. The following equation is \mathbb{P} -a.s. satisfied by $\{(Z_\varepsilon^m, Q_\varepsilon^m)\}_{m=1}^M$:

$$\begin{cases} (Z_\varepsilon^m - Z_\varepsilon^{m-1}, \varphi_h) + k\nu(\nabla Z_\varepsilon^m, \nabla \varphi_h) + k \langle \hat{B}(V_\varepsilon^{m-1}, V_\varepsilon^m) - \hat{B}(U_\varepsilon^{m-1}, U_\varepsilon^m), \varphi_h \rangle \\ - k(Q_\varepsilon^m, \operatorname{div}(\varphi_h)) = ([g(V_\varepsilon^{m-1}) - g(U_\varepsilon^{m-1})]\Delta_m W, \varphi_h), \quad \forall \varphi \in \mathbb{H}_h, \\ \frac{\varepsilon}{k} (Q_\varepsilon^m - Q_\varepsilon^{m-1}, q_h) + (\operatorname{div}(Z_\varepsilon^m), q_h) = 0, \quad \forall q_h \in L_h. \end{cases} \quad (4.24)$$

For $m = 1$, it follows that $g(V_\varepsilon^0) - g(U_\varepsilon^0) = 0$ and $\hat{B}(V_\varepsilon^0, V_\varepsilon^1) - \hat{B}(U_\varepsilon^0, U_\varepsilon^1) = \hat{B}(V_\varepsilon^0 - U_\varepsilon^0, Z_\varepsilon^1) = 0$. Hence, setting $(\varphi_h, q_h) = (Z_\varepsilon^1, Q_\varepsilon^1)$ in equations (4.24) yields $\|Z_\varepsilon^1\|_{\mathbb{L}^2}^2 + \varepsilon\|Q_\varepsilon^1\|_{L^2}^2 + k\nu\|\nabla Z_\varepsilon^1\|_{\mathbb{L}^2}^2 = 0$ which implies $Z_\varepsilon^1 = Q_\varepsilon^1 = 0$ \mathbb{P} -a.s. and a.e. in $[0, T] \times D$. Arguing by induction completes the proof. \square

All steps that were conducted in section 4.2.2 are applicable to Algorithm 2, except for Lemma 4.4 which does not suit the associated bilinear operator \hat{B} since its variables are not identical. Therefore, a slight adjustment should take place, and it consists of the following:

In Step 1 of Section 4.2.2, $\mathcal{R}(\mathcal{V}_{\varepsilon, k, h}^+)$ shall be substituted by a new operator $\mathcal{S}(\mathcal{V}_{\varepsilon, k, h}^-, \mathcal{V}_{\varepsilon, k, h}^+) := -\nu\Delta\mathcal{V}_{\varepsilon, k, h}^+ + \hat{B}(\mathcal{V}_{\varepsilon, k, h}^-, \mathcal{V}_{\varepsilon, k, h}^+)$ and \mathcal{R}_0 by \mathcal{S}_0 which is defined by

$$\left[\int_0^T \langle \mathcal{S}_0(t), \varphi \rangle dt \right] = \lim_{\varepsilon, k, h \rightarrow 0} \mathbb{E} \left[\int_0^T \langle \mathcal{S}(\mathcal{V}_{\varepsilon, k, h}^-, \mathcal{V}_{\varepsilon, k, h}^+) + \nabla \Pi_{\varepsilon, k, h}^+, \Pi_h \varphi dt \rangle \right], \quad \forall \varphi \in \mathcal{V}.$$

Equation (4.21) remains unchanged because $\langle \mathcal{S}(\mathcal{V}_{\varepsilon,k,h}^-, \mathcal{V}_{\varepsilon,k,h}^+), \mathcal{V}_{\varepsilon,k,h}^+ \rangle = \langle \mathcal{R}(\mathcal{V}_{\varepsilon,k,h}^+), \mathcal{V}_{\varepsilon,k,h}^+ \rangle$, thanks to Proposition 2.1-(ii). However, when passing to the limit, term II_3 in Step 2 is not suitable for \mathcal{S}_0 , which is why it can be modified by employing Proposition 2.1-(ii) as follows:

$$\begin{aligned} II'_3 &= -2\mathbb{E} \left[\int_0^T \eta^- \left\langle \mathcal{S}(\mathcal{V}_{\varepsilon,k,h}^-, \mathcal{V}_{\varepsilon,k,h}^+) + \nabla II_{\varepsilon,k,h}^+ - \mathcal{S}(\sigma_h^-, \sigma_h^+), \sigma_h^+ \right\rangle dt \right] \\ &\quad - 2\mathbb{E} \left[\int_0^T \eta^- \left\langle \hat{B} \left(\mathcal{V}_{\varepsilon,k,h}^+ - \mathcal{V}_{\varepsilon,k,h}^-, \mathcal{V}_{\varepsilon,k,h}^+ \right), \sigma_h^+ \right\rangle dt \right] := II'_{3,1} + II'_{3,2}. \end{aligned}$$

$II'_{3,1}$ goes to $-2\mathbb{E} \left[\int_0^T \eta(t) \langle \mathcal{S}_0(t) - \mathcal{S}(\sigma(t), \sigma(t)), \sigma(t) \rangle dt \right]$ as $\varepsilon, k, h \rightarrow 0$. Consequently, the whole proof of section 4.2.2 becomes applicable to Algorithm 2, provided that $II'_{3,2}$ goes to 0. To this end, denote $Z_{\varepsilon,k,h} = \mathcal{V}_{\varepsilon,k,h}^+ - \mathcal{V}_{\varepsilon,k,h}^-$ and employ Proposition 2.1-(iii) to ensure:

$$\begin{aligned} \int_0^T \eta(t) \left\langle \hat{B} \left(Z_{\varepsilon,k,h}, \mathcal{V}_{\varepsilon,k,h}^+ \right), \sigma_h^+ \right\rangle dt &\lesssim \int_0^T \|Z_{\varepsilon,k,h}\|_{\mathbb{L}^2}^{\frac{1}{2}} \|\nabla Z_{\varepsilon,k,h}\|_{\mathbb{L}^2}^{\frac{1}{2}} \|\nabla \mathcal{V}_{\varepsilon,k,h}^+\|_{\mathbb{L}^2} \|\nabla \sigma_h^+\|_{\mathbb{L}^2} dt \\ &\lesssim k^{\frac{1}{4}} \left(\sum_{m=1}^M \|V_\varepsilon^m - V_\varepsilon^{m-1}\|_{\mathbb{L}^2}^2 \right)^{\frac{1}{4}} \left(k \sum_{m=1}^M \|\nabla(V_\varepsilon^m - V_\varepsilon^{m-1})\|_{\mathbb{L}^2}^2 \right)^{\frac{1}{4}} \left(k \sum_{m=1}^M \|\nabla V_\varepsilon^m\|_{\mathbb{L}^2}^2 \right)^{\frac{1}{2}}, \end{aligned}$$

thanks to the Hölder inequality and the high regularity of σ . Therewith,

$$II'_{3,2} \lesssim k^{\frac{1}{4}} \mathbb{E} \left[\sum_{m=1}^M \|V_\varepsilon^m - V_\varepsilon^{m-1}\|_{\mathbb{L}^2}^2 \right]^{\frac{1}{4}} \mathbb{E} \left[k \sum_{m=1}^M \|\nabla(V_\varepsilon^m - V_\varepsilon^{m-1})\|_{\mathbb{L}^2}^2 \right]^{\frac{1}{4}} \mathbb{E} \left[k \sum_{m=1}^M \|\nabla V_\varepsilon^m\|_{\mathbb{L}^2}^2 \right]^{\frac{1}{2}}.$$

The first and third expectations are bounded by virtue of Lemma 4.3-(i). Additionally, the second expectation, after undergoing a triangle inequality, can be controlled in a similar way provided that $\|\nabla v_h^0\|_{\mathbb{L}^2}$ is uniformly bounded in h . Consequently, $II'_{3,2} \lesssim k^{\frac{1}{4}} \rightarrow 0$.

One way of ensuring uniform boundedness in h of $\|\nabla v_h^0\|_{\mathbb{L}^2}$ is through the Ritz (also known as elliptic) operator $\mathcal{R}_h: \mathbb{H}_0^1 \rightarrow \mathbb{H}_h$, which is stable in \mathbb{H}_0^1 (see for instance [27]). In other words, setting $v_h^0 = \mathcal{R}_h v_0$ gets the job done, as long as $v_0 \in \mathbb{H}_0^1$. Another way is to use the already defined projection Π_h which can be an alternative for \mathcal{R}_h . This is true since the triangulation \mathcal{T}_h is quasi-uniform (see [6, Theorem 4]). With being said, an additional theorem can be given.

Theorem 4.1 *For $T > 0$, let $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{0 \leq t \leq T}, \mathbb{P})$ be a filtered probability space, $D \subset \mathbb{R}^2$ be a polygonal domain, assumptions (S_1) - (S_2) be satisfied, and $v_0 \in L^2(\Omega; \mathbb{H}_0^1)$. Given a positive integer M , define the discretization step size $k := \frac{T}{M}$ such that $k \in (0, 1)$ and I_k forms a uniform partition of the time interval $[0, T]$. Let $\varepsilon \in (0, 1)$ be a given penalty parameter, and $h \in (0, 1)$ be the space discretization step size such that the triangulation \mathcal{T}_h is quasi-uniform. Define the finite element triple $(\mathbb{H}_h, L_h, (v_h^0, p_h^0))$ such that the initial datum (v_h^0, p_h^0) belongs to (\mathbb{H}_h, L_h) , and $v_h^0 \in \{\mathcal{R}_h v_0, \Pi_h v_0\}$. Then, the following results hold:*

- For a given $(k, h, \varepsilon) \in (0, 1)^3$, there is a discrete solution $\{(V_\varepsilon^m, \Pi_\varepsilon^m)\}_{m=1}^M$ to Algorithm 2 satisfying Lemmas 4.1, 4.3, and 4.5.
- For a family $\{k, h, \varepsilon\}$ of parameters fulfilling $k, h, \varepsilon \rightarrow 0$ simultaneously, such that $v_h^0 \rightarrow v_0$ as $h \rightarrow 0$ in the space $L^2(\Omega; \mathbb{L}^2)$, the solution $\{(V_\varepsilon^m, \Pi_\varepsilon^m)\}_{m=1}^M$ of Algorithm 2 converges toward the unique strong solution of stochastic Navier-Stokes equations (1.1) in the sense of Definition 2.2.

4.4 How to properly choose ε regarding saddle point-based schemes?

For simplicity's sake and knowing that the Stokes problem establishes an insight into the Navier-Stokes equations, the foremost aim of this section will be to evaluate a Stokes version of Algorithm 1 against a specific numerical scheme of saddle point aspect of the following stochastic Stokes problem:

$$\begin{cases} \frac{\partial u}{\partial t} - \nu \Delta u + \nabla p = f + g(u) \frac{\partial W}{\partial t}, & \text{in } (0, T) \times D, \\ \operatorname{div}(u) = 0, & \text{in } (0, T) \times D, \\ u(0, \cdot) = v_0, & \text{in } D. \end{cases} \quad (4.25)$$

in order to choose the parameter ε effectively. The finite element spaces \mathbb{H}_h and L_h will be maintained throughout this section, and the discrete LBB (also known as inf-sup) condition

$$\sup_{\varphi_h \in \mathbb{H}_h} \frac{(\operatorname{div}(\varphi_h), q_h)}{\|\nabla \varphi_h\|_{\mathbb{L}^2}} \geq \beta \|q_h\|_{L^2}, \quad \forall q_h \in L_h, \quad (4.26)$$

will be required since numerical schemes of Stokes and Navier-Stokes problems dealing with saddle point techniques lack of velocity-pressure stability if such a condition was omitted. The constant $\beta > 0$ does not depend on the mesh size h . With that said, it is now meaningful to state the convective-free version of Algorithm 1:

$$\begin{cases} (U_\varepsilon^m - U_\varepsilon^{m-1}, \varphi_h) + k\nu (\nabla U_\varepsilon^m, \nabla \varphi_h) - k(p_\varepsilon^m, \operatorname{div}(\varphi_h)) \\ = k \langle f^m, \varphi_h \rangle + (g(U_\varepsilon^{m-1}) \Delta_m W, \varphi_h), \quad \forall \varphi_h \in \mathbb{H}_h, \\ \frac{\varepsilon}{k} (p_\varepsilon^m - p_\varepsilon^{m-1}, q_h) + (\operatorname{div}(U_\varepsilon^m), q_h) = 0, \quad \forall q_h \in L_h, \end{cases} \quad (4.27)$$

together with the following saddle point-based numerical scheme of the Stokes problem:

$$\begin{cases} (U^m - U^{m-1}, \varphi_h) + k\nu (\nabla U^m, \nabla \varphi_h) - k(p^m, \operatorname{div}(\varphi_h)) \\ = k \langle f^m, \varphi_h \rangle + (g(U^{m-1}) \Delta_m W, \varphi_h), \quad \forall \varphi_h \in \mathbb{H}_h, \\ (\operatorname{div}(U^m), q_h) = 0, \quad \forall q_h \in L_h. \end{cases} \quad (4.28)$$

Here, $\Delta_m W$ and f^m are identical to those of Algorithm 1, and the starting points $U_\varepsilon^0 = U^0 = \Pi_h v_0$. The convergence analysis of scheme (4.28) along with its convergence rate are provided in [11]. To come up with effective and adequate conditions upon the parameter ε , it suffices to investigate the quantity $\|U_\varepsilon^m - U^m\|$. This is logical because if u denotes the solution of Stokes equations (4.25), then $\|U_\varepsilon^m - u(t_m)\| \leq \|U_\varepsilon^m - U^m\| + \|U^m - u(t_m)\|$ grants the rate at which scheme (4.27) might converge. To this purpose, subtracting equations (4.27) and (4.28) yields for all $\varphi_h \in \mathbb{H}_h \setminus \{0\}$,

$$\begin{aligned} & (U_\varepsilon^m - U^m - (U_\varepsilon^{m-1} - U^{m-1}) - [g(U_\varepsilon^{m-1}) - g(U^{m-1})] \Delta_m W, \varphi_h) \\ & + k(p^m - p_\varepsilon^m, \operatorname{div}(\varphi_h)) = k\nu (\nabla(U^m - U_\varepsilon^m), \nabla \varphi_h) \leq k\nu \|\nabla(U^m - U_\varepsilon^m)\|_{\mathbb{L}^2} \|\nabla \varphi_h\|_{\mathbb{L}^2}. \end{aligned} \quad (4.29)$$

Dividing by $\|\nabla \varphi_h\|_{\mathbb{L}^2}$, taking the supremum over $\varphi_h \in \mathbb{H}_h \setminus \{0\}$ and employing the discrete LBB-condition (4.26) imply

$$\|p^m - p_\varepsilon^m\|_{L^2} \leq \frac{\nu}{\beta} \|\nabla(U^m - U_\varepsilon^m)\|_{\mathbb{L}^2}, \quad \forall m \in \{1, \dots, M\}. \quad (4.30)$$

Estimate (4.30) is true because $\omega \mapsto \sup_{\varphi_h \in \mathbb{H}_h \setminus \{0\}} \frac{(U_\varepsilon^m - U^m - (U_\varepsilon^{m-1} - U^{m-1}) - [g(U_\varepsilon^{m-1}) - g(U^{m-1})] \Delta_m W, \varphi_h)}{\|\nabla \varphi_h\|_{\mathbb{L}^2}}$ is

non-negative which results from the fact that \mathbb{H}_h is a vector space. In other words, this supremum can be roughly seen as the \mathbb{H}^{-1} -norm of $U_\varepsilon^m - U^m - (U_\varepsilon^{m-1} - U^{m-1}) - [g(U_\varepsilon^{m-1}) - g(U^{m-1})] \Delta_m W$. On the

other hand, setting $\varphi_h = U_\varepsilon^m - U^m$ in equation (4.29), using identity $2(a - b, a) = \|a\|_{\mathbb{L}^2}^2 - \|b\|_{\mathbb{L}^2}^2 + \|a - b\|_{\mathbb{L}^2}^2$, the Cauchy-Schwarz and Young inequalities return

$$\begin{aligned} & \frac{1}{2} \|U_\varepsilon^m - U^m\|_{\mathbb{L}^2}^2 - \frac{1}{2} \|U_\varepsilon^{m-1} - U^{m-1}\|_{\mathbb{L}^2}^2 + k\nu \|\nabla(U_\varepsilon^m - U^m)\|_{\mathbb{L}^2}^2 \leq k(p_\varepsilon^m - p^m, \operatorname{div}(U_\varepsilon^m)) \\ & + ([g(U_\varepsilon^{m-1}) - g(U^{m-1})] \Delta_m W, U_\varepsilon^{m-1} - U^{m-1}) + \frac{1}{2} \|[g(U_\varepsilon^{m-1}) - g(U^{m-1})] \Delta_m W\|_{\mathbb{L}^2}^2, \end{aligned} \quad (4.31)$$

where $(p_\varepsilon^m - p^m, \operatorname{div}(U^m)) = 0$, thanks to scheme (4.28). Summing the above equation over m from 1 to an arbitrary $\ell \in \{1, \dots, M\}$, taking its mathematical expectation, employing the Itô isometry to the last term on its right-hand side together with assumption (S_2) and making use of the identity $U_\varepsilon^0 = U^0$ yield

$$\begin{aligned} & \mathbb{E} \left[\frac{1}{2} \|U_\varepsilon^\ell - U^\ell\|_{\mathbb{L}^2}^2 + k\nu \sum_{m=1}^{\ell} \|\nabla(U_\varepsilon^m - U^m)\|_{\mathbb{L}^2}^2 \right] \leq \mathbb{E} \left[k \sum_{m=1}^{\ell} (p_\varepsilon^m - p^m, \operatorname{div}(U_\varepsilon^m)) \right] \\ & + \frac{L_g^2}{2} \mathbb{E} \left[k \sum_{m=1}^{\ell} \|U_\varepsilon^{m-1} - U^{m-1}\|_{\mathbb{L}^2}^2 \right], \end{aligned} \quad (4.32)$$

where the mathematical expectation of the penultimate term in equation (4.31) vanishes due to assumption (S_2) and the measurability of $\{U_\varepsilon^m\}_{m=1}^M$ and $\{U^m\}_{m=1}^M$. Attention will now turn toward the first term on the right-hand side of equation (4.32) which will eventually hand the upper-bound in terms of ε . Using equations (4.27), one obtains

$$\begin{aligned} J & := \mathbb{E} \left[k \sum_{m=1}^{\ell} (p_\varepsilon^m - p^m, \operatorname{div}(U_\varepsilon^m)) \right] = -\mathbb{E} \left[\varepsilon \sum_{m=1}^{\ell} (p_\varepsilon^m - p_\varepsilon^{m-1}, p_\varepsilon^m - p^m) \right] \\ & \leq \frac{\varepsilon\nu}{\beta} \mathbb{E} \left[\sum_{m=1}^{\ell} \|p_\varepsilon^m - p_\varepsilon^{m-1}\|_{L^2} \|\nabla(U_\varepsilon^m - U^m)\|_{\mathbb{L}^2} \right] \\ & \leq \frac{\sqrt{\varepsilon}\nu}{\beta} \mathbb{E} \left[\sum_{m=1}^M \varepsilon \|p_\varepsilon^m - p_\varepsilon^{m-1}\|_{L^2}^2 \right]^{1/2} \mathbb{E} \left[\sum_{m=1}^{\ell} \|\nabla(U_\varepsilon^m - U^m)\|_{\mathbb{L}^2}^2 \right]^{1/2} \\ & \leq \frac{\sqrt{\varepsilon}\nu}{4\beta^2} \mathbb{E} \left[\varepsilon \sum_{m=1}^M \|p_\varepsilon^m - p_\varepsilon^{m-1}\|_{L^2}^2 \right] + \sqrt{\varepsilon}\nu \mathbb{E} \left[\sum_{m=1}^{\ell} \|\nabla(U_\varepsilon^m - U^m)\|_{\mathbb{L}^2}^2 \right] \\ & \leq \frac{\sqrt{\varepsilon}\nu}{4\beta^2} C_1 + \sqrt{\varepsilon}\nu \mathbb{E} \left[\sum_{m=1}^{\ell} \|\nabla(U_\varepsilon^m - U^m)\|_{\mathbb{L}^2}^2 \right], \end{aligned} \quad (4.33)$$

thanks to the Cauchy-Schwarz and Young inequalities, estimate (4.30), and Lemma 4.3-(ii). In order to handle the last term on the right-hand side of equation (4.33), the penalty parameter ε shall undergo an assumption; that is, $\sqrt{\varepsilon} \leq k$. This way, it becomes absorbable in the left-hand side of equation (4.32). Finally, plug the result of estimate (4.33) in inequality (4.32) and make use of the discrete Grönwall inequality to achieve

$$\frac{1}{2} \max_{1 \leq m \leq M} \mathbb{E} \left[\|U_\varepsilon^m - U^m\|_{\mathbb{L}^2}^2 \right] + \mathbb{E} \left[(k - \sqrt{\varepsilon})\nu \sum_{m=1}^M \|\nabla(U_\varepsilon^m - U^m)\|_{\mathbb{L}^2}^2 \right] \leq \tilde{C}\sqrt{\varepsilon}, \quad (4.34)$$

for some constant $\tilde{C} > 0$ depending only on β, C_1, ν, L_g and T .

Estimate (4.34) seems to have the best upper-bound amongst the other possible ways of estimation. Besides, some calculation techniques may be inconsistent with the assumption $\sqrt{\varepsilon} \leq k$. For instance,

Proposition 4.1 could have been employed for the estimation of the term J , especially for the penultimate inequality in equation (4.33):

$$J \leq \frac{C_1 \nu}{2\beta^2} \frac{k}{\sqrt{\varepsilon}} + \sqrt{\varepsilon} \nu \mathbb{E} \left[\sum_{m=1}^M \|\nabla(U_\varepsilon^m - U^m)\|_{\mathbb{L}^2}^2 \right],$$

however, the second term on the right-hand side of the above inequality becomes non absorbable in the left-hand side of equation (4.32). Indeed, if the assumption $\sqrt{\varepsilon} \leq k$ is imposed, the obtained rate $\frac{k}{\sqrt{\varepsilon}}$ will no longer go to 0, which is senseless.

5 Numerical experiments and conclusion

The implementation within this section will be carried out through Algorithm 2 and a saddle point based-numerical scheme [4, Algorithm 3]:

Algorithm 3 Let $M \in \mathbb{N}$ and $V^0 = v_h^0 \in \mathbb{H}_h$ be given. For every $m \in \{1, \dots, M\}$, find an $\mathbb{H}_h \times L_h$ -valued (V^m, Π^m) such that

$$\begin{cases} (V^m - V^{m-1}, \varphi_h) + k\nu (\nabla V^m, \nabla \varphi_h) + k\hat{b}(V^{m-1}, V^m, \varphi_h) - k(\Pi^m, \operatorname{div}(\varphi_h)) \\ \quad = k\langle f^m, \varphi_h \rangle + (g(V^{m-1})\Delta_m W, \varphi_h), \quad \forall \varphi_h \in \mathbb{H}_h, \\ (\operatorname{div}(V^m), q_h) = 0, \quad \forall q_h \in L_h, \end{cases}$$

which will play the reference role with respect to the values of the parameter ε . The domain's meshing is carried out through the open source finite element mesh generator Gmsh [12], the implementation of the aforementioned algorithms is executed by the open source finite element software FEniCS [16], and the visualization is ensured via Paraview [1]. The simulation's configuration down below is set as follows: $T = 1$, $\nu = 1$, $h = 0.16$, $\varepsilon = 10^{-5}$, $k = 0.01$. For the sake of comparison, the space discretization will be conducted by the lower order Taylor-Hood (P_2/P_1) finite element for both algorithms 2 and 3. The initial data u_0 and π_0 are set to 0 which means that $v_h^0 = (0, 0)$ and $p_h^0 = 0$. The domain D is an L -shaped geometry whose figure and mesh are displayed in Figure 1.




Figure 1: The domain D and its mesh

The boundary condition

$$v(x, y) = \begin{cases} (1, 0) & \text{if } (x, y) \in \{0\} \times [0, 1], \\ (0, 0) & \text{elsewhere,} \end{cases}$$

is non-homogeneous, which is possible since a simple lifting technique can take the problem's boundary condition back to a homogeneous setting. The source term f takes on the value $(0, 0)$ and the diffusion coefficient $g = Id$ i.e. it is an additive noise. The Wiener increment $\Delta_m W$ is approximated as follows: let $J \in \mathbb{N}$ be non-zero, and W_1, W_2 be two independent $H_0^1(D)$ -valued Wiener processes such that $W = (W_1, W_2)$. Then,

$$\Delta_m W_\ell \approx \sqrt{k} \sum_{i,j=1}^J \xi_{i,j}^{\ell,m} e_{i,j}, \quad \ell \in \{1, 2\}.$$

The parameter J takes on the value 5, $\{(\xi_{i,j}^{1,m}, \xi_{i,j}^{2,m})\}_{i,j}^m$ is a family of independent identically distributed normal random variables, and $e_{i,j}(x, y) = \frac{2}{5} \sin(i\pi x/5) \sin(j\pi y/5)$ for all $i, j \in \mathbb{N}$. Although $\{e_{i,j}\}_{i,j}$ may not be the best choice for an L -shaped domain (because they represent the Laplace eigenfunctions on the square $(0, 5)^2$ with a Dirichlet boundary condition), they can be thought of herein as a restriction to D . The explicit formula of the Laplace eigenfunctions on an L -shaped domain is unknown as it is explained in [21]. With all that being said, it is now possible to exhibit the simulation results:




Figure 2: One realization of V^M (left) and V_ε^M (right) at time $T = 1$ for $\varepsilon = 10^{-5}$

As ε gets smaller, the difference between V^m and V_ε^m becomes indistinguishable. This fact is illustrated in an accurate way down below where the relationship between ε and the error $\mathbb{E} \left[\|V^M - V_\varepsilon^M\|_{\mathbb{L}^2}^2 \right]$ is exposed:




Figure 3: Error and error-variance in terms of ε

The computed error in figure 3 uses a Monte-Carlo method with 1000 realizations. The obtained curve was expected; it emphasizes the fact that ε should be taken as small as possible in order to guarantee accurate outcomes.

6 Conclusion

This paper provides a new approach to simulating the two-dimensional stochastic incompressible Navier-Stokes equations. The introduced technique can be thought of as a compromise between (strongly) divergence-free finite element methods and saddle point problems where a discrete LBB condition is required to prove the pressure's existence. No relationships were assumed between ε and the discretization parameters k and h , although it could have been possible. If so, ε must be solely linked to k on account of the penalizing term $\varepsilon\partial_t p^\varepsilon$ which offers a supplementary time regularity, meaning that h should not intervene. Whereas, in the case where the additional term was, for example, $\varepsilon\Delta\partial_t p^\varepsilon$; that is, the mass conservation equation of system (1.2) had the following form:

$$\operatorname{div}(v^\varepsilon) + \varepsilon\Delta\partial_t p^\varepsilon = 0,$$

then, the parameter ε may be expressed in terms of both discretization step sizes k and h because the Laplace operator offers a supplementary space-regularity to the pressure field p^ε , which needs to disappear when passing to the limit in order to recover a solution to the stochastic incompressible Navier-Stokes equations. In Section 4.4, ε was also linked to the time discretization step size k under a particular numerical scheme that involves a discrete LBB condition. This given relationship can dramatically deviate if another numerical method is selected to be compared with the proposed algorithm herein. After all, the most accurate assumption involving both k and ε can come to light during the convergence rate study of Algorithm 1 (or 2).

Declarations

Competing interests The author confirms the absence of any conflict of interest associated with this paper.

References

- [1] J. Ahrens, Berk Geveci, and Charles Law. “ParaView: An End-User Tool for Large Data Visualization”. In: *Vis. Handb.* (Jan. 2005).
- [2] Francesca Bonizzoni and Guido Kanschat. “H1 -conforming finite element cochain complexes and commuting quasi-interpolation operators on Cartesian meshes”. In: *Calcolo* 58.2 (2021), p. 18. ISSN: 0008-0624. DOI: 10.1007/s10092-021-00409-6.
- [3] Susanne Brenner and Ridgway Scott. *The mathematical theory of finite element methods*. Vol. 15. New York: Springer Sci. & Bus. Media, 2007, p. 400. DOI: 10.1007/978-0-387-75934-0.
- [4] Zdzislaw Brzeźniak, Erich Carelli, and Andreas Prohl. “Finite-element-based discretizations of the incompressible Navier–Stokes equations with multiplicative random forcing”. In: *IMAJNA*. 33.3 (Jan. 2013), pp. 771–824. ISSN: 0272-4979. DOI: 10.1093/imanum/drs032.

- [5] Erich Carelli and Andreas Prohl. “Rates of Convergence for Discretizations of the Stochastic Incompressible Navier-Stokes Equations”. In: *SIAM J. on Numer. Anal.* 50.5 (2012), pp. 2467–2496.
- [6] M. Crouzeix and V. Thomée. “The Stability in L_p and W_p^1 of the L_2 -Projection onto Finite Element Function Spaces”. In: *Mathematics of Computation* 48.178 (1987), pp. 521–532. ISSN: 00255718, 10886842.
- [7] Anne De Bouard and Arnaud Debussche. “A semi-discrete scheme for the stochastic nonlinear Schrödinger equation”. In: *Numerische Mathematik* 96.4 (2004), pp. 733–770.
- [8] Jim Douglas, Todd Dupont, and Lars Wahlbin. “The stability in L_q of the L_2 -projection into finite element function spaces”. In: *Numerische Mathematik* 23.3 (1974), pp. 193–197. DOI: 10.1007/BF01400302.
- [9] John A Evans and Thomas JR Hughes. “Isogeometric divergence-conforming B-splines for the unsteady Navier-Stokes equations”. In: *J. Comput. Phys.* 241 (2013), pp. 141–167.
- [10] Xiaobing Feng, Andreas Prohl, and Liet Vo. “Optimally convergent mixed finite element methods for the stochastic Stokes equations”. In: *IMA Journal of Numerical Analysis* 41.3 (2021), pp. 2280–2310.
- [11] Xiaobing Feng and Hailong Qiu. “Analysis of Fully Discrete Mixed Finite Element Methods for Time-dependent Stochastic Stokes Equations with Multiplicative Noise”. In: *J. of Sci. Comput.* 88.2 (2021), p. 31. DOI: 10.1007/s10915-021-01546-4.
- [12] Christophe Geuzaine and Jean-François Remacle. “Gmsh: A 3-D finite element mesh generator with built-in pre-and post-processing facilities”. In: *Int. j. numer. methods in eng.* 79.11 (2009), pp. 1309–1331.
- [13] Vivette Girault and Pierre-Arnaud Raviart. *Finite element methods for Navier-Stokes equations: theory and algorithms*. Vol. 5. Berlin Heidelberg: Springer Sci. & Bus. Media, 2012. DOI: <https://doi.org/10.1007/978-3-642-61623-5>.
- [14] O. A. Ladyzhenskaya et al. “The Mathematical Theory of Viscous Incompressible Flow”. In: *Phys. Today* 17.2 (1964), pp. 57–58. DOI: 10.1063/1.3051412.
- [15] JoséA. Langa, José Real, and Jacques Simon. “Existence and Regularity of the Pressure for the Stochastic Navier-Stokes Equations”. In: *Appl. Math. and Optim.* 48.3 (2003), pp. 195–210. DOI: 10.1007/s00245-003-0773-7.
- [16] Anders Logg, Kent-Andre Mardal, Garth N. Wells, et al. *Automated Solution of Differential Equations by the Finite Element Method*. Springer, 2012. ISBN: 978-3-642-23098-1. DOI: 10.1007/978-3-642-23099-8.
- [17] Utpal Manna, Jose-Luis Menaldi, and Sivaguru Sritharan. “Stochastic 2-D Navier-Stokes Equation with Artificial Compressibility”. In: *Commun. on Stoch Anal.* 1 (Mar. 2008). DOI: 10.31390/cosa.1.1.10.
- [18] Jose-Luis Menaldi and Sivaguru S Sritharan. “Stochastic 2-D Navier-Stokes Equation.” In: *Appl. Math. & Optim.* 46.1 (2002).
- [19] Michael Neilan and Duygu Sap. “Stokes elements on cubic meshes yielding divergence-free approximations”. In: *Calcolo* 53.3 (2016), pp. 263–283. DOI: 10.1007/s10092-015-0148-x.

- [20] E. Pardoux. “Equations aux dérivées partielles stochastiques non linéaires monotones”. PhD thesis. Université Paris XI, 1975.
- [21] J. K. Reid and J. E. Walsh. “An Elliptic Eigenvalue Problem for a Reentrant Region”. In: *J. Soc. Ind. Appl. Math.* 13.3 (1965), pp. 837–850. ISSN: 03684245.
- [22] Jie Shen. “On a new pseudocompressibility method for the incompressible Navier-Stokes equations”. In: *Appl. Numer. Math.* 21 (May 1996), pp. 71–90. DOI: 10.1016/0168-9274(95)00132-8.
- [23] Jie Shen. “On pressure stabilization method and projection method for unsteady Navier-Stokes equations”. In: *Adv. Comput. Methods Partial Differ. Equ.* Citeseer. 1992.
- [24] R. Témam. “Sur l’approximation de la solution des équations de Navier-Stokes par la méthode des pas fractionnaires (I)”. In: *Arch. for Ration. Mech. and Anal.* 32.2 (1969), pp. 135–153. DOI: 10.1007/BF00247678.
- [25] Roger Temam. *Navier-Stokes equations: theory and numerical analysis*. Vol. 343. Am. Math. Soc., 2001.
- [26] Roger Temam. “Une méthode d’approximation de la solution des équations de Navier-Stokes”. fr. In: *Bull. de la Soc. Math. de Fr.* 96 (1968), pp. 115–152. DOI: 10.24033/bsmf.1662.
- [27] Vidar Thomée. *Galerkin finite element methods for parabolic problems*. Vol. 25. Springer Science & Business Media, 2007.