



HAL
open science

Nash balanced assignment problem

Minh Hieu Nguyen, Mourad Baiou, Viet Hung Nguyen

► **To cite this version:**

Minh Hieu Nguyen, Mourad Baiou, Viet Hung Nguyen. Nash balanced assignment problem. 7th International Symposium on Combinatorial Optimization (ISCO), May 2022, Online Conference, France. hal-03656133v1

HAL Id: hal-03656133

<https://hal.science/hal-03656133v1>

Submitted on 1 May 2022 (v1), last revised 5 Oct 2023 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Nash balanced assignment problem

Minh Hieu Nguyen¹ , Mourad Baiou¹ , Viet Hung Nguyen^{1,2}

¹ INP Clermont Auvergne, Univ Clermont Auvergne, Mines Saint-Etienne, CNRS,
UMR 6158 LIMOS, 1 Rue de la Chebarde, Aubiere Cedex, France

² Corresponding author : viet.hung.nguyen@uca.fr
{minh.hieu.nguyen,mourad.baiou}@uca.fr

Abstract. In this paper, we consider a variant of the classic *Assignment Problem* (AP), called the *Balanced Assignment Problem* (BAP) [2]. The BAP seeks to find an assignment solution which has the smallest value of *max-min distance*: the difference between the maximum assignment cost and the minimum one. However, by minimizing only the max-min distance, the total cost of the BAP solution is neglected and it may lead to a very inefficient solution in terms of total cost. Hence, we propose a fair way based on Nash equilibrium [1] [3], [4] to inject the total cost into the objective function of the BAP for finding assignment solutions having a better trade-off between the two objectives: the first aims at minimizing the total cost and the second aims at minimizing the max-min distance. For this purpose, we introduce the concept of *Nash Fairness* (NF) solutions based on the definition of proportional-fair scheduling adapted in the context of the AP: a transfer of utilities between the total cost and the max-min distance is considered to be fair if the percentage increase in the total cost is smaller than the percentage decrease in the max-min distance and vice versa.

We first show the existence of a NF solution for the AP which is exactly the optimal solution minimizing the product of the total cost and the max-min distance. However, finding such a solution may be difficult as it requires to minimize a concave function. The main result of this paper is to show that finding all NF solutions can be done in polynomial time. For that, we propose a Newton-based iterative algorithm converging to NF solutions in polynomial time. It consists in optimizing a sequence of linear combinations of the two objective based on Weighted Sum Method [5]. Computational results on various instances of the AP are presented and commented.

Keywords: Combinatorial optimization · Balanced assignment problem · Proportional fairness · Proportional-fair scheduling · Weighted Sum Method

1 Introduction

The *Assignment Problem* (AP) is a fundamental combinatorial optimization problem. It can be formally defined as follows. Given a set n workers, a set of n jobs and a $n \times n$ cost matrix whose elements are positive representing the

assignment of any worker to any job, the AP aims at finding an one-to-one worker-job assignment (i.e a bipartite perfect matching) that minimizes certain objective functions.

In the classic AP, we seek to find an assignment solution minimizing the total cost. It is a well-known optimization problem that can be solved by Hungarian algorithm in $O(n^3)$ [7]. The *Balanced Assignment Problem* (BAP) is a variant of the classic AP where instead of minimizing the total cost, we minimize *the max-min distance* which is the difference between the maximum assignment cost and the minimum one in the assignment solution. In [2], the authors proposed an efficient threshold-based algorithm to solve the BAP in $O(n^4)$. However, by minimizing only the max-min distance, the total cost of the BAP solution is neglected and it may lead to a very inefficient solution in terms of total cost.

In this paper, to overcome the possible inefficiency of the solutions for the BAP, we propose a fair way based on Nash equilibrium to inject the total cost into the objective function of the BAP. Nash equilibrium is the most common optimality notion for sharing resources among users [1] [3],[4]. We are interested in assignment solutions for the AP achieving a Nash equilibrium between two players: the first aims at minimizing the total cost and the second aims at minimizing the max-min distance. For this purpose, we introduce the *Nash Fairness* (NF) solutions based on the definition of proportional-fair scheduling adapted in the context of the AP: a transfer of utilities between the total cost and the max-min distance is considered to be fair if the percentage increase in the total cost is smaller than the percentage decrease in the max-min distance and vice versa.

In fact, we have introduced the concept of NF solutions for the Balanced Traveling Salesman Problem (BTSP) in a recent paper [12]. In [12], we proposed an algorithm converging to particular NF solutions called *extreme NF solutions* having respectively smallest value of total cost and max-min distance. Similar to [12], in this current paper we also introduce the concept of NF solutions in the context of the AP. But, the main contribution of our work in this paper is a stronger result than in [12]: we provide an algorithm for finding all NF solutions in polynomial time.

The paper is organized as follows. In Section 2, we introduce a linear programming (LP) formulation for the BAP. The concept of NF solutions is presented in Section 3. In particular, we prove the existence of NF solutions for the AP and show that they are optimal solutions of a weighted sum objective problem. In Section 4, an algorithm for finding all NF solutions as well as computational results on various instances of the AP is given. Finally, conclusion and future works of this paper are discussed in Section 5.

2 LP formulation for BAP

We consider an AP with a $n \times n$ cost matrix whose elements $c_{i,j}$ are positive and they represent the cost assignments between worker i and job j . We first

present the linear programming (LP) formulation for the BAP as follows

$$\min Q = u - l \quad (1a)$$

$$\text{s.t. } \sum_{j \in [n]} x_{j,i} = \sum_{j \in [n]} x_{i,j} = 1 \quad \forall i \in [n] \quad (1b)$$

$$u \geq \sum_{j \in [n]} c_{i,j} x_{i,j} \quad \forall i \in [n] \quad (1c)$$

$$l \leq \sum_{j \in [n]} c_{i,j} x_{i,j} \quad \forall i \in [n] \quad (1d)$$

$$x_{i,j} \geq 0 \quad \forall i, j \in [n]. \quad (1e)$$

where $[n] = \{1, \dots, n\}$ and $x_{i,j}$ represents the assignment between worker i and job j corresponding to the cost assignment $c_{i,j}$. In order to calculate the max-min distance Q , we need to determine the maximum and the minimum assignment costs u and l in the assignment solution. Constraints (1c) obviously allow to bound u from below by the maximum assignment cost in the assignment solution. Similarly, constraints (1d) allow to bound l from above by the minimum assignment cost in the assignment solution. As $Q = u - l$ is minimized, u and l will respectively take the values of the maximum and the minimum assignment costs. We will show that this LP formulation has integral optimal solution corresponding to an assignment solution (i.e bipartite perfect matching).

Theorem 1. *This LP formulation always has an optimal solution where the variables take integer values.*

Proof. The objective function assures that u and l will be equal respectively to the maximum and the minimum assignment costs in the optimal solution. Consequently, the optimal solution of this LP is always integral because the constraints matrix of (1b) is totally unimodular (e.g see [6]) and the constraints (1c) and (1d) are simply bound constraints. \square

In the following we solve the classic AP as well as the BAP for several instances of the AP where we generate random uniform $c_{i,j}$ in $[1, 10^2]$. Optimal solutions of these instances are shown in Table 1 where *assignx* represents an instance of the AP with a cost matrix of dimension $x \times x$ and P, Q represent respectively the total cost and the max-min distance in the optimal solution. We use CPLEX 12.10 on a PC Intel Core i5-9500 3.00GHz with 6 cores and 6 threads for solving the classic AP and the BAP. We can see in each instance of the AP that the optimal solutions for the classic AP may be undesirable with respect to those for the BAP and vice versa: inefficient values of Q in the optimal solutions for the classic AP comparing with those in the optimal solutions for the BAP and inefficient values of P in the optimal solutions for the BAP comparing with those in the optimal solutions for the classic AP.

Hence, the purpose of this paper is using a fair way to inject the total cost into the objective function for finding assignment solutions having a better trade-off between the two objectives.

Table 1. Optimal solutions for the classic AP and the BAP

Instance	classic AP			BAP		
	P	Q	Time	P	Q	Time
assign3	100	6	0.01	200	3	0.01
assign6	114	15	0.01	173	10	0.01
assign17	68	10	0.01	242	3	2.43
assign25	189	27	0.03	2004	11	2.44
assign30	157	18	0.04	643	7	32.4

3 Nash fairness solutions for the AP

We have introduced the concept of Nash fairness (NF) solution for the Balanced Traveling Salesman Problem (BTSP) [12]. In this section, we restate the concept of NF solutions in the context of the AP and we put the proofs of theorems in Appendix.

3.1 Proportional fairness

The concept of NF solutions for the AP is closely related to the concept of proportional fairness for multiple players problem [1]. In the context of multiple players problem, let U be a set of possible *states of the world* or *alternatives* and let I be a finite set, representing a collection of individuals. For each $i \in I$, $u_i : U \rightarrow \mathbb{R}_+$ be a utility function, describing the amount of happiness an individual i derives from each possible state such that we prefer the alternative x to the alternative y if and only if $u_i(x) \geq u_i(y)$, $\forall i \in I$.

NF solutions for two-player problem [3] are defined by using the Nash standard of comparison: a transfer of utilities between the two players is considered to be fair if the percentage increase in the utility of one player is larger than the percentage decrease in utility of the other player [1].

Proportional fairness introduced by Bertsimas et al [1] is a generalized NF solution for multiple players. In that setting, the fair allocation should be such that, if compared to any other feasible allocation of utilities, the aggregate proportional change is less than or equal to 0 [3], [1], [4].

Definition 1. [1] $x^{NF} \in U$ is a NF solution for multiple players problem if and only if

$$\sum_{j=1}^n \frac{u_j(x) - u_j(x^{NF})}{u_j(x^{NF})} \leq 0, \forall x \in U, \quad (2)$$

where n is the number of players and $u_j(x) > 0$, $\forall j \in I, \forall x \in U$.

3.2 Characterization of NF solutions for the AP

Let P, Q now represent the total cost and the max-min distance in a feasible assignment solution for the AP. From the definitions of P and Q we have $P \geq Q \geq 0$. We first suppose that $Q > 0$. As P, Q now are two strictly positive utility functions, we have a two-player problem. In the usual definition of NF solutions [3], [1], an alternative assigned a greater value is preferred. However, in the context of the AP, we prefer the alternative assigned a smaller value for two utility functions P and Q . Thus, the aggregate proportional change should be greater than or equal to 0 in the definition of NF solutions for the AP. That is to say, the sum of relative gains when switching from NF solutions to another feasible solution is not negative in the context of the AP.

We denote the value solution for the total cost and the max-min distance corresponding to a feasible assignment solution by (P, Q) . Let (P^*, Q^*) be a NF solution for the AP, condition (2) can be translated into the context of the AP as follows

$$\frac{P - P^*}{P^*} + \frac{Q - Q^*}{Q^*} \geq 0, \forall (P, Q) \in \mathcal{S}, \quad (3)$$

which is equivalent to

$$PQ^* + QP^* \geq 2P^*Q^*, \forall (P, Q) \in \mathcal{S}, \quad (4)$$

where \mathcal{S} is the set of solutions (P, Q) corresponding to all feasible assignment solutions for the AP.

Note that in case $Q^* = 0$, the condition (4) is also satisfied. Hence, NF solution for the AP can be generally stated as follows

Lemma 1. [12] $(P^*, Q^*) \in \mathcal{S}$ is a NF solution for the AP if and only if $PQ^* + QP^* \geq 2P^*Q^*$, $\forall (P, Q) \in \mathcal{S}$.

Remark 1. An assignment solution having all equal assignment costs (i.e $Q = 0$) is a NF solution.

3.3 Existence of NF solutions

In this section, we first show the existence of NF solutions for the AP. Let us recall that in the multiple players problem mentioned in Section 3.1 where we prefer an alternative assigned a greater value, NF solutions can be obtained equivalently as the optimal solution of the problem

$$\max \sum_{j=1}^n \log u_j,$$

provided that U is convex. Notice that the above NF solution is equivalently the one maximizing the product of the utilities over U [1].

On the contrary in the AP, there exists NF solutions which can be obtained by minimizing instead of maximizing the product of the utilities.

Theorem 2. [12] $(P^*, Q^*) = \arg \min_{(P, Q) \in \mathcal{S}} PQ$ is a NF solution.

Proof. See Appendix. □

Theorem 2 proves the existence of NF solutions for the AP that minimize PQ , or equivalently minimize $(\log P + \log Q)$. We call such solutions *Product Nash Fairness* (PNF) solutions. However, finding PNF solutions may be difficult as it requires to minimize a concave function. In the following, we show that all NF solutions can be obtained by solving the following optimization problem

$$\mathcal{P}(\alpha) = \min \alpha P + Q \text{ s.t } (P, Q) \in \mathcal{S},$$

where $\alpha \in [0, 1]$ is a coefficient to be determined. For solving $\mathcal{P}(\alpha)$, we solve the LP formulation in Section 2 with $\alpha P + Q$ as the objective function instead of Q .

Let $\alpha \in \mathbb{R}_+$ and (P_α, Q_α) be an optimal solution of $\mathcal{P}(\alpha)$. Denote $T_\alpha := \alpha P_\alpha - Q_\alpha$ and $\mathcal{C}_0 := \{\alpha \in \mathbb{R}_+ | T_\alpha = 0\}$. Hence, if $\alpha \in \mathcal{C}_0$ (i.e $T_\alpha = 0$) then $\alpha \in [0, 1]$, otherwise $T_\alpha > P_\alpha - Q_\alpha \geq 0$.

Theorem 3. [12] $(P^*, Q^*) \in \mathcal{S}$ is a NF solution if and only if (P^*, Q^*) is an optimal solution of $\mathcal{P}(\alpha^*)$ where $\alpha^* = \frac{Q^*}{P^*}$.

Proof. See Appendix. □

Consequently, we state the following corollary.

Corollary 1. (P^*, Q^*) is NF solution if and only if $\alpha^* = \frac{Q^*}{P^*} \in \mathcal{C}_0$ and (P^*, Q^*) is an optimal solution of $\mathcal{P}(\alpha^*)$.

We call (P, Q) a Pareto-optimal solution for the AP if (P, Q) is an optimal solution of $\mathcal{P}(\alpha)$ where $\alpha \in [0, 1]$. By Theorem 3, a NF solution is necessarily a Pareto-optimal solution but not vice versa.

Proposition 1. *There may be more than one NF solution for the AP.*

Proof. See Appendix. □

The main question now is how to determine the coefficients of \mathcal{C}_0 corresponding one-to-one to all the NF solutions according to Theorem 3. In the next section, we present an algorithm for finding all NF solutions in polynomial time.

4 Finding all NF solutions for the AP

In [12], we proposed an algorithm converging to *extreme NF solutions* having respectively the smallest value of P and Q . In this section, we introduce another one to find all NF solutions in polynomial time. Obviously, they include the PNF solutions minimizing PQ .

4.1 Upper bound for the number of NF solutions

We call $(P, Q), (P', Q')$ two distinct solutions if $(P, Q) \not\equiv (P', Q')$. We will show that the number of NF solutions for the AP is at most $C_{n^2}^2 + n$ where $C_{n^2}^2 = \frac{n^2(n^2-1)}{2}$ by the following lemma and theorem.

Lemma 2. *If $(P, Q) \not\equiv (P', Q')$ are two distinct NF solutions having $Q, Q' > 0$ then $P \neq P'$ and $Q \neq Q'$.*

Proof. Suppose that $Q = Q' > 0$. Using the definition of NF solution we have

$$P'Q + Q'P \geq 2PQ \quad \text{and} \quad P'Q + Q'P \geq 2P'Q'.$$

which is equivalent to

$$P' + P \geq 2P \quad \text{and} \quad P' + P \geq 2P'.$$

Hence we obtain $P = P'$ which leads to a contradiction. By repeating the same argument for $P = P'$, we also have a contradiction. \square

Theorem 4. *The number of NF solutions for the AP is at most $C_{n^2}^2 + n$.*

Proof. If (P, Q) is a NF solution and $Q = 0$, the corresponding assignment solution has n equal assignment costs. For the AP with $n \times n$ cost matrix, we have n^2 assignments and consequently there are at most n distinct NF solutions having the same value $Q = 0$.

We consider now the NF solutions with $Q > 0$. We will show that the number of NF solutions having $Q > 0$ is at most $C_{n^2}^2$.

Let c_i^{max} and c_i^{min} be the maximum and the minimum assignment cost in the assignment solution corresponding to (P_i, Q_i) then $Q_i = c_i^{max} - c_i^{min}$. As shown in Lemma 2, for two distinct NF solutions $(P_i, Q_i), (P_j, Q_j)$ with both Q_i and Q_j strictly positive we obtain $Q_i \neq Q_j$ which is equivalent to $c_i^{max} - c_i^{min} \neq c_j^{max} - c_j^{min}$. We have then $(c_i^{max}, c_i^{min}) \not\equiv (c_j^{max}, c_j^{min})$. Thus, the assignment solutions corresponding to $(P_i, Q_i), (P_j, Q_j)$ have distinct pair of assignments representing the maximum and the minimum assignment cost. As we have at most n^2 distinct assignments, the number of distinct pairs of assignments is at most $C_{n^2}^2$. Consequently, the number of NF solutions having $Q > 0$ is at most $C_{n^2}^2$. Hence, the total number of NF solutions for the AP is at most $C_{n^2}^2 + n$. \square

By Theorem 4, the number of Pareto-optimal solutions having distinct values of Q is at most $C_{n^2}^2$.

4.2 Algorithm for finding all NF solutions

As shown in Theorem 3, each element $\alpha^* \in \mathcal{C}_0$ corresponds to a NF solution and vice versa. For finding all NF solutions, we aim at finding all elements of \mathcal{C}_0 . Our main idea is that from each $\alpha_0 \in [0, 1]$, we first use a procedure called *Find()* to find $\alpha_k \in \mathcal{C}_0$ satisfying α_k is the unique element $\in \mathcal{C}_0$ between α_0 and α_k .

Thus, let \mathcal{I} be the set containing the intervals $[\alpha_i, \alpha_j]$ corresponding to distinct Pareto-optimal solutions $(P_i, Q_i), (P_j, Q_j)$. We use another procedure called $Test()$ for verifying whether there exists a NF solution corresponding to $c_k \in [\alpha_i, \alpha_j]$ or not. Using these procedures, the algorithm for finding all NF solutions can be stated as follows.

Algorithm 1 Finding all NF solutions

Input: An AP with positive values in a $n \times n$ cost matrix.

Output: Set \mathcal{C}_0 whose elements correspond to all NF solutions for this AP.

```

1:  $c_0 \leftarrow Find(0)$ 
2:  $c_1 \leftarrow Find(1)$ 
3: if  $c_1 = c_0$  then
4:    $\mathcal{C}_0 = \{c_0\}$ 
5: else
6:    $\mathcal{I} = \{[c_0, c_1]\}, \mathcal{C}_0 = \{c_0, c_1\}$ 
7:   for  $[c_i, c_j] \in \mathcal{I}$  do
8:      $Test([c_i, c_j])$ 
9:   end for
10: end if
11: procedure FIND( $\alpha$ )
12:   repeat
13:     solve  $\mathcal{P}(\alpha)$  to obtain  $(P, Q)$ 
14:      $T \leftarrow \alpha P - Q$ 
15:      $\alpha \leftarrow Q/P$ 
16:   until  $T = 0$ 
17:   Return  $\alpha$ .
18: end procedure
19: procedure TEST( $[c_i, c_j]$ )
20:    $\alpha_k = \frac{Q_i - Q_j}{P_j - P_i}$ 
21:   if  $\mathcal{P}(\alpha_k)$  has an optimal solution different to  $(P_i, Q_i)$  and  $(P_j, Q_j)$  then
22:      $c_k \leftarrow Find(\alpha_k)$ 
23:     if  $c_i = c_k$  then
24:        $[c_i, c_j] \leftarrow [\alpha_k, c_j]$  ▷ Update the elements of  $\mathcal{I}$ 
25:        $Test([\alpha_k, c_j])$ 
26:     else if  $c_k = c_j$  then
27:        $[c_i, c_j] \leftarrow [c_i, \alpha_k]$  ▷ Update the elements of  $\mathcal{I}$ 
28:        $Test([c_i, \alpha_k])$ 
29:     else ▷  $c_k$  is a new element of  $\mathcal{C}_0$ 
30:        $[c_i, c_j] \leftarrow [c_i, c_k], [c_k, c_j]$  ▷ Update the elements of  $\mathcal{I}$ 
31:        $\mathcal{C}_0 \leftarrow \mathcal{C}_0 \cup c_k$ 
32:        $Test([c_i, c_k]), Test([c_k, c_j])$ 
33:     end if
34:   end if
35: end procedure

```

Let α_0 be the initial point, $T_i = \alpha_i P_i - Q_i$ and $\{\alpha_i\}$ denote the sequence constructed by Procedure $Find(\alpha_0)$. We show that Algorithm 1 explores all NF solutions in polynomial time by the following lemmas and theorem. Due to lack of space, we put some proofs in Appendix.

Lemma 3. [12] *Let $\alpha, \alpha' \in \mathbb{R}_+$ and $(P_\alpha, Q_\alpha), (P_{\alpha'}, Q_{\alpha'})$ be the optimal solutions of $\mathcal{P}(\alpha)$ and $\mathcal{P}(\alpha')$ respectively, if $\alpha \leq \alpha'$ then $P_\alpha \geq P_{\alpha'}$ and $Q_\alpha \leq Q_{\alpha'}$.*

Proof. See Appendix. \square

As a consequence of Lemma 3, if $(P_\alpha, Q_\alpha), (P_{\alpha'}, Q_{\alpha'})$ are optimal solutions of $\mathcal{P}(\alpha)$ and $\mathcal{P}(\alpha')$ and $P_\alpha < P_{\alpha'}$ (or $Q_\alpha > Q_{\alpha'}$) then we obtain $\alpha \geq \alpha'$.

Lemma 4. [12] *During the execution of Procedure $Find(\alpha_0)$ in Algorithm 1, $\alpha_i \in [0, 1], \forall i \geq 1$. Moreover, if $T_0 \geq 0$ then the sequence $\{\alpha_i\}$ is non-increasing and $T_i \geq 0, \forall i \geq 0$. Otherwise, if $T_0 \leq 0$ then the sequence $\{\alpha_i\}$ is non-decreasing and $T_i \leq 0, \forall i \geq 0$.*

Proof. See Appendix. \square

As a consequence of Lemma 4, if $T_k = 0$ then $\alpha_k = \alpha_{k+1}$, if $T_k > 0$ then $T_i > 0, \alpha_i > \alpha_{i+1}, \forall 0 \leq i \leq k$ and if $T_k < 0$ then $T_i < 0, \alpha_i < \alpha_{i+1}, \forall 0 \leq i \leq k$.

Lemma 5. [12] *From each $\alpha_0 \in [0, 1]$, Procedure $Find(\alpha_0)$ converges to a coefficient $\alpha_k \in C_0$ satisfying α_k is the unique element $\in C_0$ between α_0 and α_k .*

Proof. See Appendix. \square

Lemma 6. *Procedure $Find(\alpha_0)$ converges in polynomial time.*

Proof. Suppose that Procedure $Find(\alpha_0)$ converges to $\alpha_k \in C_0$ in $k+1$ iterations. We only consider the nontrivial case where $k > 0$ (i.e $T_0 \neq 0$).

If $T_0 > 0$ we have $T_i > 0, \forall 0 \leq i \leq k-1$.

By contradiction, we will show that if $T_{i+1} > 0$ then $P_i < P_{i+1}$ and $Q_i > Q_{i+1}$.

Let assume that $P_i \geq P_{i+1}$. According to Lemma 4, $\alpha_i \geq \alpha_{i+1}$ that implies $P_i \leq P_{i+1}$ as the result of Lemma 3. Thus, $P_i = P_{i+1}$. Moreover, $T_{i+1} = \alpha_{i+1} P_{i+1} - Q_{i+1} = \frac{Q_i P_{i+1} - Q_{i+1} P_i}{P_i} > 0$ leads to $Q_i P_{i+1} > Q_{i+1} P_i$. Since $P_i = P_{i+1} > 0$, we get $Q_i > Q_{i+1}$.

On the other hand, as (P_i, Q_i) is the optimal solution $\mathcal{P}(\alpha_i)$, it shows that $\alpha_i P_i + Q_i \leq \alpha_i P_{i+1} + Q_{i+1}$. Using $P_i = P_{i+1}$, we obtain $Q_i \leq Q_{i+1}$ which leads to a contradiction.

By repeating the same argument for $Q_i \leq Q_{i+1}$, we also have a contradiction.

Similarly, if $T_0 < 0$ we obtain the same conclusion.

Consequently, the execution of Procedure $Find(\alpha_0)$ explores k Pareto-optimal solutions having distinct values of Q . As the number of Pareto-optimal solutions having distinct value of Q is at most $C_{n^2}^2$, Procedure $Find(\alpha_0)$ converges after a polynomial number of iterations. Hence, Procedure $Find(\alpha_0)$ converges in polynomial time cause the LP formulation in Section 2 for $\mathcal{P}(\alpha)$ can be solved in polynomial time. \square

Now by using the following lemma, we show that Procedure $Test()$ can be used for verifying the existence of a Pareto-optimal solution (as well as NF solution) in each interval $[\alpha_i, \alpha_j]$.

Lemma 7. *Given an interval $[\alpha_i, \alpha_j]$ defined by $0 \leq \alpha_i < \alpha_j \leq 1$ corresponding to two distinct Pareto-optimal solutions (P_i, Q_i) and (P_j, Q_j) . Let $\alpha^* = \frac{Q_j - Q_i}{P_i - P_j}$, if $\mathcal{P}(\alpha^*)$ has no Pareto-optimal solution which is different to (P_i, Q_i) and (P_j, Q_j) , there does not have another one in $[\alpha_i, \alpha_j]$.*

Proof. Using Lemma 3 with $\alpha_i < \alpha_j$ and $(P_i, Q_i), (P_j, Q_j)$ be the two distinct Pareto-optimal solutions, we have $Q_i < Q_j, P_i > P_j$.

We first show that $\alpha^* \in [\alpha_i, \alpha_j]$.

Due to the optimality of (P_i, Q_i) and (P_j, Q_j) we obtain

$$\begin{aligned}\alpha_i P_i + Q_i &\leq \alpha_i P_j + Q_j, \\ \alpha_j P_j + Q_j &\leq \alpha_j P_i + Q_i,\end{aligned}$$

Hence, $\alpha_i \leq \frac{Q_j - Q_i}{P_i - P_j} \leq \alpha_j$ which leads to $\alpha_i \leq \alpha^* \leq \alpha_j$.

Now suppose that we do not obtain any Pareto-optimal solution by solving $\mathcal{P}(\alpha^*)$ which is different to (P_i, Q_i) and (P_j, Q_j) , we will show that there does not have another one in $[\alpha_i, \alpha_j]$.

Since $\alpha^* = \frac{Q_j - Q_i}{P_i - P_j}$, we have $\alpha^* P_i + Q_i = \alpha^* P_j + Q_j$. That means in this case (P_i, Q_i) and (P_j, Q_j) are two optimal solutions of $\mathcal{P}(\alpha^*)$. If there exists another Pareto-optimal solution (P, Q) of $\mathcal{P}(\alpha)$ where $\alpha \in [\alpha_i, \alpha_j]$, we have then $Q_i < Q < Q_j$ and $P_i > P > P_j$. Applying the consequence of Lemma 3 with $Q < Q_j$ we obtain $\alpha \leq \alpha^*$. Similarly, from $Q > Q_i$ we obtain $\alpha \geq \alpha^*$.

Hence, $\alpha = \alpha^*$ and then $\mathcal{P}(\alpha^*)$ has the Pareto-optimal solution (P, Q) which is different to (P_i, Q_i) and (P_j, Q_j) . It leads to a contradiction. \square

Theorem 5. *Algorithm 1 explores all NF solutions in polynomial time.*

Proof. As a consequence of Lemma 5, the interval $[c_0, c_1]$ contains all elements of \mathcal{C}_0 .

We known that the number of Pareto-optimal solutions having distinct values of Q is at most $C_{n^2}^2$. Consequently, $[c_0, c_1]$ can be separated by at most $C_{n^2}^2 - 1$ intervals $[c_i, c_j]$ such that $c_i < c_j$ correspond to two Pareto-optimal solutions having distinct values of Q and there intervals have no common points except the endpoints. By using Procedure $Test()$, each recursive call give us a Pareto-optimal solution or show that we have explored an interval having no Pareto-optimal solution and consequently no NF solution inside. As we use Procedure $Find(\alpha)$ in each recursive call, Procedure $Test()$ also terminates in polynomial time. Moreover, we obtain a NF solution from each Pareto-solution found with a corresponding coefficient $\in [c_0, c_1]$. Since Algorithm 1 terminated as the interval $[c_0, c_1]$ is totally explored, it found all NF solutions in polynomial time. Obviously, the PNF solutions minimizing PQ can be easily determined by comparing the products of all NF solutions. \square

4.3 Numerical results

Let us denote NFAP, PAF as the problems of finding all NF solutions and finding the NF solutions minimizing PQ for the AP. In this section, we conduct several experiments aiming at solving the NFAP with CPLEX 12.10 on some instances from the data sets of the AP [11] as well as on various instances that we generate random uniform their cost matrix. We also solve the classic AP and the BAP on the same instances. All the experiments are conducted on a PC Intel Core i5-9500 3.00GHz with 6 cores and 6 threads.

Table 2. Numerical results for the classic AP, BAP and NFBAP

Instance	classic AP			BAP			PNF				NFAP
	P	Q	Time	P	Q	Time	P	Q	Time	α	all NF solutions
assign3	100	6	0.01	200	3	0.05	140	4	0.11	0.028	(100,6), (140,4), (200,3)
assign4	70	9	0.01	196	3	0.25	196	3	1.20	0.015	(70,9), (80,8) (120,5), (196,3)
assign6	114	15	0.01	173	10	0.34	118	12	2.54	0.101	(118,12)
assign17	68	10	0.01	262	3	2.43	71	8	14.8	0.112	(71,8), (80,7), (130,4)
assign25	189	27	0.14	2004	11	7.12	189	27	41.2	0.142	(189,27), (452,14)
assign30	157	18	0.04	643	7	32.4	158	16	74.1	0.101	(158,16), (473,8)
assign45	6212	200	0.15	40937	54	85.0	6240	185	574	0.296	(6240,185), (7133,160) (9394,112), (12766,75)
assign75	8828	65	0.28	63860	36	122	9741	49	336	0.005	(9741,49)
assign100	305	6	0.58	661	3	34.3	310	3	85.6	0.009	(310,3)

Table 3. Sum of relatives gains when switching from PNF solutions to the optimal solutions of the classic AP and the BAP

Instance	Sum of relative gains	
	PNF vs classic AP	PNF vs BAP
assign3	0.214	0.178
assign4	1.357	0.000
assign6	0.216	0.299
assign17	0.207	2.065
assign25	0.000	9.010
assign30	0.118	2.507
assign45	0.076	4.852
assign75	0.232	5.290
assign100	0.983	1.132

Table 2 presents the numerical results in several instances with a range of dimension of cost matrix from 3×3 to 100×100 . We also provide the PNF

solution minimizing PQ and its corresponding value of α . We can see by the values of P and Q in this table that the PNF solution strikes a better trade-off between the total cost and the max-min distance comparing with those for the classic AP and the BAP. In particular, when the solutions for the classic AP and for the BAP are quite different: inefficient values of Q in the optimal solutions for the classic AP comparing with those in the optimal solutions for the BAP and inefficient values of P in the optimal solutions for the BAP comparing with those in the optimal solutions for the classic AP, the PNF solution offers almost a better alternative than the solution for the classic AP (respectively for the BAP) with a significant drop on the values of Q (respectively P) and a slight growth on the values of P (respectively Q). More precisely, Table 3 presents the sums of relative gains when switching from PNF solutions to the optimal solutions for the classic AP and the BAP. Note that their values are not negative as we mentioned in (3), Section 3.2 and values further from 0 are preferable for the PNF solutions because they have then a much better trade-off between P and Q . Table 2 also indicates that we only have several NF solutions and in most cases, the PNF solution is one of the extreme NF solutions having smallest value of P or Q . One important issue is the CPU time for solving the NFAP (approximately for finding the PNF solution) is quite huge comparing with the CPU time spent for solving the classic AP and the BAP. A deeper analysis on the iterations of Procedure $Find(\alpha)$ tells us that the CPU time spent for solving $P(\alpha)$ with small value of α occupies a very big part of the overall CPU time. Hence, a special-purpose algorithm for solving $P(\alpha)$ may be more interesting than simply optimizing a linear function over the LP given in Section 2.

5 Conclusion

In this paper, we have made use of Nash fairness equilibrium to achieve a trade-off between the efficiency estimated by the total cost and the balancedness estimated by the max-min distance in solutions for the *Assignment Problem* (AP). We have proven first the existence of *Nash Fairness* (NF) solutions for the AP. Second, we have designed an algorithm to find all NF solutions including the PNF solutions minimizing the product of total cost and max-min distance. Numerical results conducted on instances of the AP have shown that comparing with the optimal solutions for the BAP, the NF solutions found by our algorithm have almost much smaller total cost with a reasonable augmentation of the max-min distance and vice versa comparing with the optimal solutions for the classic AP. An important notice is that the results in this paper can be also applied to various balanced combinatorial optimization problems such as balanced traveling salesman problem [12], balanced spanning tree problem [10], etc. The future developments of our work are improving time complexity for Algorithm 1 by developing a special-purpose algorithm for solving $P(\alpha)$. Moreover, we are also interested in finding a better upper bound for the number of NF solutions as well as generating the concept of NF solutions for bi-objective optimization problems having positive objective functions.

References

1. D. Bertsimas, V.F. Farias, N. Trichakis, The Price of Fairness, *Operations Research*, January-February 2011, 59(1), pp 17-31.
2. S. Martello, W.R. Pulleyblank, P. Toth, D. de Werra, Balanced optimization problems, *Operations Research Letters* Volume 3, Number 5, August 1984.
3. FP Kelly, AK Maullo, DKH Tan, Rate control for communication networks : shadow prices, proportional fairness and stability, *Journal of the Operational Research Society* 49(3), November 1997.
4. W. Ogryczak, Hanan Luss, Michab Pioro, Dritan Nace, Artur Tomaszewski, Fair Optimization and Networks: A survey, *Journal of Applied Mathematics, Hindawi Publishing Coporation* (2014). pp 1-26.
5. R. Timothy Marler, Jasbir Singh Arora, The weighted sum method for multi-objective optimization: New insights, *Structural and Multidisciplinary Optimization* 41(6), June 2010. pp 853-862.
6. I. Heller, C.B. Tompkins, An Extension of a Theorem of Dantzig's, *Annals of Mathematics Studies, Princeton University Press* (1956). 247-254(38).
7. H. W. Kuhn, The Hungarian method for assignment problem, *Naval Research Logistics Quarterly* 2 (1-2): 83-97, March 1955.
8. W. Dinkelbach, On nonlinear fractional programming, *Management Sci.* (1967) 492-498(13).
9. J.R. Isbell, W.H. Marlow, Attrition games, *Naval Res. Logist. Quart* 3, (1956), pp 71-94.
10. Martello, Most and least uniform spanning trees, *Discrete Applied Mathematics* 15(2) 1986, pp 181-197.
11. J.E. Beasley, Linear programming on Clay supercomputer, *Journal of the Operational Research Society*, 41 (1990), pp 133-139.
12. M.H. Nguyen, Mourad Baiou, V.H. Nguyen and T.Q.T. Vo, Nash fairness solutions for balanced TSP, *International Network Optimization Conference, June 2022 (INOC2022)*.

APPENDIX

Proposition 1. *There may be more than one NF solution for the AP.*

Proof. Let us illustrate this by an instance of the AP having the following cost matrix :

$$A = \begin{bmatrix} 30 & 48 & 68 \\ 44 & 65 & 34 \\ 67 & 36 & 48 \end{bmatrix}$$

By verifying all feasible assignment solutions in this instance, we obtain easily three assignment solutions $(1-1, 2-3, 3-2)$, $(1-2, 2-1, 3-3)$ and $(1-3, 2-2, 3-1)$ corresponding to three NF solutions $(100, 6)$, $(140, 4)$ and $(200, 3)$. Note that $i-j$ where $1 \leq i, j \leq 3$ represents the assignment between worker i and job j in the solution of this instance. \square

We recall below the proofs of some recent results that we have published in [12]. They are needed to prove the new results presented in this paper. They are put in this appendix to facilitate the review of the paper.

Theorem 2. [12] $(P^*, Q^*) = \arg \min_{(P, Q) \in \mathcal{S}} PQ$ is a NF solution.

Proof. Obviously, there always exists a solution $(P^*, Q^*) \in \mathcal{S}$ such that

$$(P^*, Q^*) = \arg \min_{(P, Q) \in \mathcal{S}} PQ.$$

Now $\forall (P', Q') \in \mathcal{S}$ we have $P'Q' \geq P^*Q^*$. Then

$$P'Q^* + Q'P^* \geq 2\sqrt{P'Q'P^*Q^*} \geq 2P^*Q^*,$$

The first inequality holds by the Cauchy-Schwarz inequality.
Hence, (P^*, Q^*) is a NF solution. \square

Theorem 3. [12] $(P^*, Q^*) \in \mathcal{S}$ is a NF solution if and only if (P^*, Q^*) is an optimal solution of $\mathcal{P}(\alpha^*)$ where $\alpha^* = \frac{Q^*}{P^*}$.

Proof. Firstly, let (P^*, Q^*) be a NF solution and $\alpha^* = \frac{Q^*}{P^*}$. We will show that (P^*, Q^*) is an optimal solution of $\mathcal{P}(\alpha^*)$.

Since (P^*, Q^*) is a NF solution, we have

$$P'Q^* + Q'P^* \geq 2P^*Q^*, \quad \forall (P', Q') \in \mathcal{S}, \quad (6)$$

Since $\alpha^* = \frac{Q^*}{P^*}$, we have $\alpha^*P^* + Q^* = 2Q^*$.

Dividing two sides of (6) by $P^* > 0$ we obtain

$$2Q^* \leq \frac{Q^*}{P^*}P' + Q', \quad \forall (P', Q') \in \mathcal{S}, \quad (7)$$

So we deduce from (7)

$$\alpha^* P^* + Q^* \leq \alpha^* P' + Q', \quad \forall (P', Q') \in \mathcal{S},$$

Hence, (P^*, Q^*) is an optimal solution of $\mathcal{P}(\alpha^*)$.

Now suppose $\alpha^* = \frac{Q^*}{P^*}$ and (P^*, Q^*) is an optimal solution of $\mathcal{P}(\alpha^*)$, we show that (P^*, Q^*) is a NF solution.

If (P^*, Q^*) is not a NF solution, there exists a solution $(P', Q') \in \mathcal{S}$ such that

$$P'Q^* + Q'P^* < 2P^*Q^*,$$

We have then

$$\alpha P' + Q' = \frac{P'Q^* + Q'P^*}{P^*} < \frac{2P^*Q^*}{P^*} = \alpha^* P^* + Q^*,$$

which contradicts the optimality of (P^*, Q^*) . \square

Lemma 3. [12] *Let $\alpha, \alpha' \in \mathbb{R}_+$ and $(P_\alpha, Q_\alpha), (P_{\alpha'}, Q_{\alpha'})$ be the optimal solutions of $\mathcal{P}(\alpha)$ and $\mathcal{P}(\alpha')$ respectively, if $\alpha \leq \alpha'$ then $P_\alpha \geq P_{\alpha'}$ and $Q_\alpha \leq Q_{\alpha'}$.*

Proof. The optimality of (P_α, Q_α) and $(P_{\alpha'}, Q_{\alpha'})$ gives

$$\alpha P_\alpha + Q_\alpha \leq \alpha P_{\alpha'} + Q_{\alpha'}, \quad \text{and} \quad (8a)$$

$$\alpha' P_{\alpha'} + Q_{\alpha'} \leq \alpha' P_\alpha + Q_\alpha \quad (8b)$$

By adding both sides of (8a) and (8b), we obtain $(\alpha - \alpha')(P_\alpha - P_{\alpha'}) \leq 0$. Since $\alpha \leq \alpha'$, it follows that $P_\alpha \geq P_{\alpha'}$.

On the other hand, inequality (8a) implies $Q_{\alpha'} - Q_\alpha \geq \alpha(P_\alpha - P_{\alpha'}) \geq 0$ that leads to $Q_\alpha \leq Q_{\alpha'}$. \square

Lemma 4. [12] *During the execution of Procedure Find(α_0) in Algorithm 1, $\alpha_i \in [0, 1], \forall i \geq 1$. Moreover, if $T_0 \geq 0$ then the sequence $\{\alpha_i\}$ is non-increasing and $T_i \geq 0, \forall i \geq 0$. Otherwise, if $T_0 \leq 0$ then the sequence $\{\alpha_i\}$ is non-decreasing and $T_i \leq 0, \forall i \geq 0$.*

Proof. Since $P \geq Q \geq 0, \forall (P, Q) \in \mathcal{S}$, it follows that $\alpha_{i+1} = \frac{Q_i}{P_i} \in [0, 1], \forall i \geq 0$.

We first consider $T_0 \geq 0$. We proof $\alpha_i \geq \alpha_{i+1}, \forall i \geq 0$ by induction on i . For $i = 0$, we have $T_0 = \alpha_0 P_0 - Q_0 = P_0(\alpha_0 - \alpha_1) \geq 0$, it follows that $\alpha_0 \geq \alpha_1$. Suppose that our hypothesis is true until $i = k \geq 0$, we will prove that it is also true with $i = k + 1$.

Indeed, we have

$$\alpha_{k+1} - \alpha_{k+2} = \frac{Q_k}{P_k} - \frac{Q_{k+1}}{P_{k+1}} = \frac{Q_k P_{k+1} - P_k Q_{k+1}}{P_k P_{k+1}},$$

The inductive hypothesis gives $\alpha_k \geq \alpha_{k+1}$ that implies $P_{k+1} \geq P_k > 0$ and $Q_k \geq Q_{k+1} \geq 0$ according to Lemma 3. It leads to $Q_k P_{k+1} - P_k Q_{k+1} \geq 0$ and then $\alpha_{k+1} - \alpha_{k+2} \geq 0$.

Hence, we have $\alpha_i \geq \alpha_{i+1}$, $\forall i \geq 0$.

Consequently, $T_i = \alpha_i P_i - Q_i = P_i(\alpha_i - \alpha_{i+1}) \geq 0$, $\forall i \geq 0$.

Similarly, if $T_0 \leq 0$ we obtain that the sequence $\{\alpha_i\}$ is non-decreasing and $T_i \leq 0$, $\forall i \geq 0$. That concludes the proof. \square

Lemma 5. [12] *From each $\alpha_0 \in [0, 1]$, Procedure $Find(\alpha_0)$ converges to a coefficient $\alpha_k \in \mathcal{C}_0$ satisfying α_k is the unique element $\in \mathcal{C}_0$ between α_0 and α_k .*

Proof. As a consequence of Lemma 4, Procedure $Find(\alpha_0)$ converges to a coefficient $\alpha_k \in [0, 1]$, $\forall \alpha_0 \in [0, 1]$.

By the stopping criteria of Procedure $Find(\alpha_0)$, when $T_k = \alpha_k P_k - Q_k = 0$ we obtain $\alpha_k \in \mathcal{C}_0$ and (P_k, Q_k) is a NF solution. (Theorem 3)

If $T_0 = 0$ then obviously $\alpha_k = \alpha_0$. We consider $T_0 > 0$ and the sequence $\{\alpha_i\}$ is now non-negative, non-increasing. We will show that $[\alpha_k, \alpha_0] \cap \mathcal{C}_0 = \alpha_k$.

Suppose that we have $\alpha \in (\alpha_k, \alpha_0]$ and $\alpha \in \mathcal{C}_0$ corresponding to a NF solution (P, Q) . Then there exists $1 \leq i \leq k$ such that $\alpha \in (\alpha_i, \alpha_{i-1}]$. Since $\alpha \leq \alpha_{i-1}$, $P \geq P_{i-1}$ and $Q \leq Q_{i-1}$ due to Lemma 3. Thus, we get

$$\frac{Q}{P} \leq \frac{Q_{i-1}}{P_{i-1}} \quad (9)$$

By the definitions of α and α_i , inequality (9) is equivalent to $\alpha \leq \alpha_i$ which leads to a contradiction.

By repeating the same argument for $T_0 < 0$, we also have a contradiction. \square