



HAL
open science

Weakly and Strongly Aperiodic Subshifts of Finite Type on Baumslag-Solitar Groups

Solène Esnay, Etienne Moutot

► **To cite this version:**

Solène Esnay, Etienne Moutot. Weakly and Strongly Aperiodic Subshifts of Finite Type on Baumslag-Solitar Groups. *Theoretical Computer Science*, 2022, 917, pp.31-50. 10.1016/j.tcs.2022.03.010 . hal-03658172

HAL Id: hal-03658172

<https://hal.science/hal-03658172v1>

Submitted on 3 May 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Weakly and Strongly Aperiodic Subshifts of Finite Type on Baumslag-Solitar Groups

Solène J. Esnay^{1,2} and Etienne Moutot³
julien.esnay@ens-lyon.fr etienne.moutot@math.cnrs.fr

¹ Université de Lyon, CNRS, ENS de Lyon, UCBL, LIP, F-69342, Lyon Cedex 07, France

² Institut de Mathématiques de Toulouse, Université Paul Sabatier, 118 route de Narbonne, F-31062 Toulouse Cedex 9, France

³ CNRS, Aix Marseille Univ., I2M, Marseille, France

Abstract

We study the periodicity of subshifts of finite type (SFT) on Baumslag-Solitar groups. We show that for residually finite Baumslag-Solitar groups there exist both strongly and weakly-but-not-strongly aperiodic SFTs. In particular, this shows that unlike \mathbb{Z}^2 , but like \mathbb{Z}^3 , strong and weak aperiodic SFTs are different classes of SFTs in residually finite BS groups. More precisely, we prove that a weakly aperiodic SFT on $BS(m,n)$ due to Aubrun and Kari is, in fact, strongly aperiodic on $BS(1,n)$; and weakly but not strongly aperiodic on any other $BS(m,n)$. In addition, we exhibit an SFT which is weakly but not strongly aperiodic on $BS(1,n)$; and we show that there exists a strongly aperiodic SFT on $BS(n,n)$.

Introduction

The use of tilings as a computational model was initiated by Wang in the 60s [20] as a tool to study specific classes of logical formulas. His model consists in square tiles with colors on each side, that can be placed next to each other if the colors match. Wang studied the tilings of the discrete infinite plane (\mathbb{Z}^2) with these tiles, now called Wang tiles; a similar model exists to tile the infinite line (\mathbb{Z}) with two-sided dominoes. Wang realized that a key property of these tilings was the notion of periodicity. At the time, it was already known that if a set of dominoes tiled \mathbb{Z} , it was always possible to do it in a periodic fashion, and Wang suspected that it was the same for Wang tiles on \mathbb{Z}^2 . However, a few years later, one of his students, Berger, proved otherwise by providing a set of Wang tiles that tiled the plane but only aperiodically [5]. Numerous aperiodic sets of Wang tiles have been provided by many since then (see for example [13, 14, 18]).

The model of Wang tiles is actually equivalent to tilings using an arbitrary finite alphabet with adjacency rules, and it has become a part of symbolic dynamics, a more general way to encode a smooth dynamical system into symbolic states and trajectories. This approach by discretization (see [10] for a comprehensive historiography) is itself part of the field of discrete dynamical systems. In this broader context, it is interesting to study the set of all possible tilings for a given finite list of symbols and adjacency rules, called a Subshift of Finite Type (SFT). \mathbb{Z} -SFTs have been studied extensively, and many of their properties are known (see for example [15]). \mathbb{Z}^d -SFTs (for $d \geq 2$) seem to be much more complex, as many results from the unidimensional case cannot be directly transcribed to higher dimensions.

In recent years, the even wider class of SFTs built over Cayley graphs of groups has been attracting more and more attention [1, 6, 12].

In this broader setting, many questions about periodicity are still open [6]. For example there are many groups for which we do not know if there exists a non-empty SFT with only aperiodic configurations (called an aperiodic SFT). Another relevant question is the relation between two notions of aperiodicity, weak and strong aperiodicity: the first one requires all configurations to have infinitely many distinct translates; the second one that no configuration has any period whatsoever. They are equivalent for SFTs over \mathbb{Z} or \mathbb{Z}^2 , but this is not the case for other groups (notably \mathbb{Z}^3), and for many the relation is not even known.

Baumslag-Solitar groups of parameters m and n , commonly denoted by $BS(m, n)$, initially gathered interest in symbolic dynamics because of their simple description and rich properties. Notably, their domino problem is undecidable [3]; and Aubrun and Kari built a weakly aperiodic SFT in order to prove it. Their proof is essentially sketched in the general case, where it encodes piecewise affine maps, as it is done on \mathbb{Z}^2 in [14]. Only the $BS(2, 3)$ case is detailed in their paper; and an explicit period is provided for a given configuration, which shows that the resulting SFT, although weakly aperiodic, is not strongly aperiodic.

In the present paper, after a few definitions (Section 1); we thoroughly reintroduce Aubrun and Kari's construction in the general case, and provide a precise proof of its weak aperiodicity by encoding piecewise linear maps (Section 2) (although Aubrun and Kari sketched the proof for encoding piecewise affine maps, only piecewise linear ones are needed for the aperiodicity result). Then, we show that the resulting SFT is weakly but not strongly aperiodic on any $BS(m, n)$ (Section 2.2) (as sketched in [3]); except on $BS(1, n)$ where, with some extra work, we prove that it is strongly aperiodic (Section 2.4). Then, using a different technique based on substitutions on words, we exhibit a weakly but not strongly aperiodic SFT on $BS(1, n)$ (Section 3). Finally, by using tools from group theory and a theorem by Jeandel [12], we build a strongly aperiodic SFT on any $BS(n, n)$ (Section 4). These new results are summarized in the following table (in bold):

Group	Strongly aperiodic SFT	Weakly-not-strongly aperiodic SFT
\mathbb{Z}^2	Yes (Berger [5])	No (Folklore)
$BS(1, n)$	Yes, adapted from Aubrun-Kari (Section 2.4)	Yes, using substitutions (Section 3)
$BS(n, n)$	Yes, using a theorem by Jeandel (Section 4)	Yes (Section 2.2, Aubrun-Kari [3])
$BS(m, n)$?	Yes (Section 2.2, Aubrun-Kari [3])

The end result is that any residually finite Baumslag-Solitar group $BS(m, n)$ with $|m| \geq 2$ or $|n| \geq 2$ has a strongly aperiodic SFT and a weakly but not strongly aperiodic SFT. The remaining question is whether a general $BS(m, n)$ admits a strongly aperiodic SFT, when not in the $|m| = |n|$, $|m| = 1$ or $|n| = 1$ cases.

1 Definitions

1.1 Baumslag-Solitar groups

A group G is said to be *finitely generated* if it has a presentation $G = \langle S \mid R \rangle$ with S finite. Given a presentation $\langle S \mid R \rangle$ of a group G , its (right) *Cayley graph* is the graph $\Gamma_G = (G, E_\Gamma)$ whose vertices are the elements of G and the edges are of the form (g, gs) with $g \in G$ and $s \in S \cup S^{-1}$ a generator of G or its inverse.

A group G is said to be *residually finite* if for any $g \in G$ that is not the identity, there is a normal subgroup $N \triangleleft G$ of finite index such that $g \notin N$.

The groups we are interested in in this paper are the *Baumslag-Solitar groups*. They are defined, using two *nonzero* integers m, n as parameters, by the presentation

$$BS(m, n) = \langle a, b \mid ba^m b^{-1} = a^n \rangle.$$

Proposition 1.1 (Meskin [16]). *$BS(m, n)$ is residually finite $\Leftrightarrow |m| = 1$ or $|n| = 1$ or $|m| = |n|$.*

1.2 Subshifts and tilings

In a general sense, a *dynamical system* is a set S endowed with a topology U and on which a group acts; if \mathbb{Z} acts on it by the iteration of a function f , we may write (S, U, f) instead of (S, U, \mathbb{Z}) . Two dynamical systems (S, U, G) and (T, V, G) are *conjugate* if there exists a continuous bijection $\phi: S \rightarrow T$ so that for any $s \in S$ and any $g \in G$, $\phi(g \cdot s) = g \cdot \phi(s)$ (where G acts, respectively, on S and on T).

Let \mathcal{A} be a finite alphabet. Let G be a finitely generated group with presentation $\langle S \mid R \rangle$ and neutral element e . Let $x \in \mathcal{A}^G$ and $g, h \in G$: G naturally acts on the left on \mathcal{A}^G by $(g \cdot x)_h = x_{g^{-1}h}$. The set \mathcal{A}^G , when endowed with the product topology t and this action, forms a compact dynamical system $(\mathcal{A}^G, t, \cdot)$ called the *full shift* over G . We call $x \in \mathcal{A}^G$ a *configuration*.

A *pattern* p is a finite configuration $p \in \mathcal{A}^{P_p}$ where $P_p \subset G$ is finite. We say that a pattern $p \in \mathcal{A}^{P_p}$ *appears* in a configuration $x \in \mathcal{A}^G$ – or that x *contains* p – if there exists $g \in G$ such that for every $h \in P_p$, $(g \cdot x)_h = p_h$, in this case we write $p \sqsubset x$.

The *subshift* associated to a set of patterns \mathcal{F} , called set of *forbidden patterns*, is defined by

$$X_{\mathcal{F}} = \{x \in \mathcal{A}^G \mid \forall p \in \mathcal{F}, p \not\sqsubset x\}$$

that is, $X_{\mathcal{F}}$ is the set of all configurations that do not contain any pattern from \mathcal{F} . Note that there can be several sets of forbidden patterns defining the same subshift X . A subshift can equivalently be defined as a closed set under both the topology and the G -action.

If $X = X_{\mathcal{F}}$ with \mathcal{F} finite, then X is called a *Subshift of Finite Type*, SFT for short.

Let X be a subshift on a group G and $x \in X$. The *orbit* of x is $Orb_G(x) = \{g \cdot x \mid g \in G\}$ and its *stabilizer* $Stab_G(x) = \{g \in G \mid g \cdot x = x\}$. We say that x is a *strongly periodic configuration* if $|Orb_G(x)| < +\infty$, and that x is a *weakly periodic configuration* if $Stab_G(x) \neq \{e\}$. If no configuration in X is strongly periodic and the subshift is non-empty, then the subshift is said to be *weakly aperiodic*. If no configuration in X is weakly periodic and the subshift is non-empty, then the subshift is said to be *strongly aperiodic*.

A particular class of SFTs is obtained by considering Wang tiles over the Cayley graph of the group. This can be done for any finitely generated group, and it turns out that any SFT can be encoded into an equivalent set of Wang tiles. In order to make the definitions shorter, we limit ourselves to the definition of Wang tiles over $BS(m, n)$. A *Wang tileset* is a particular SFT where the alphabet is a set of *Wang tiles* τ , which are tuples of colors of the

form $s = (t_1^s, \dots, t_m^s, l^s, r^s, b_1^s, \dots, b_n^s)$. To make notations easier, we denote:

$$\begin{aligned} s(\text{top}_1) &= t_1^s \\ &\vdots \\ s(\text{top}_m) &= t_m^s \\ s(\text{left}) &= l^s \\ s(\text{right}) &= r^s \\ s(\text{bottom}_1) &= b_1^s \\ &\vdots \\ s(\text{bottom}_n) &= b_n^s \end{aligned}$$




Figure 1: A Wang tile of $BS(m, n)$

A *tiling* is then a configuration $T \in \tau^{BS(m, n)}$ over the group using the alphabet τ . We say that a tiling is *valid* if the colors of neighboring tiles match. That is, for any $g \in BS(m, n)$ and T_g the associated tile at position g , we must have:

$$\begin{aligned} T_g(\text{right}) &= T_{ga^m}(\text{left}) \\ T_g(\text{top}_k) &= T_{ga^{k-l}b}(\text{bottom}_l) \end{aligned}$$

for any $k \in \{1, \dots, m\}$ and $l \in \{1, \dots, n\}$. See Fig. 2 for an illustration of these rules.




Figure 2: Illustration of the neighbor rules for $BS(2, 2)$.

The set of all valid tilings for a tileset τ forms an SFT X^τ , since the tileset gives a finite number of local constraints based on a finite alphabet. In general, it is not necessarily simpler to consider Wang tilesets instead of local constraints on the Cayley graph; however, in [3]

the construction heavily uses the visual representation of tiles with numbers on the top and bottom that encode a multiplication by a real number, and this article will do the same.

1.3 Substitutions

Let \mathcal{A}^* be the set of (finite) words over \mathcal{A} . A *substitution* (or morphism) is a map $s: \mathcal{A} \rightarrow \mathcal{A}^*$. We say it is *uniform* of size $n \in \mathbb{N}$ if for every $a \in \mathcal{A}$, $|s(a)| = n$. The substitution s can be extended to \mathcal{A}^* by applying it to each letter of the word and concatenating the resulting words. We can also extend s to $\mathcal{A}^{\mathbb{N}_0}$ (resp. $\mathcal{A}^{\mathbb{N}}$) by doing the same, concatenating infinitely many words, the first letter of the first word being at position 0 (resp. 1). Finally, s can be extended to *pointed* biinfinite words (so on $\mathcal{A}^{\mathbb{Z}}$ in a certain way) as illustrated in Fig. 3.




Figure 3: How a substitution is applied to a biinfinite word: $y = s(x)$ with s a uniform substitution of size n

For $a \in \mathcal{A}$ with $s(a) = aw$ with $w \in \mathcal{A}^*$, we define the *positive infinite iteration* of s on a :

$$\overrightarrow{s}(a) = a w s(w) s^2(w) \cdots \in \mathcal{A}^{\mathbb{N}_0}.$$

In the same way, we define the *negative infinite iteration* of s on a :

$$\overleftarrow{s}(a) = \cdots s^2(w) s(w) w a \in \mathcal{A}^{\mathbb{Z}^-}.$$

For a word u (possibly biinfinite), we define its factor complexity

$$P_u(n) = |\{w \in \mathcal{A}^n \mid w \sqsubset u\}|$$

where $w \sqsubset u$ indicates that w is a subword of u . The factor complexity of a biinfinite word is bounded if and only if that word is periodic [7], that is, if it is made of the same finite word concatenated infinitely many times.

A *fixpoint* of a substitution s is a (possibly biinfinite) word u such that $s(u) = u$.

2 On a construction by Aubrun and Kari

In [3], Aubrun and Kari provide a weakly aperiodic Wang tileset on Baumslag-Solitar groups, with a proof focusing on the specific case of $BS(2, 3)$, for which they also present a period for one specific configuration, implying that the corresponding Wang tileset is not strongly aperiodic.

A more general version of the construction can be found in [4]. We repeat most of it here for the sake of completeness, since we study that construction in more details to obtain additional results: we will show that it yields a weakly but not strongly aperiodic SFT on any $BS(m, n)$ with $|m| \neq 1$ and $|n| \neq 1$, and a strongly aperiodic SFT on any $BS(1, n)$.

2.1 Aubrun and Kari's construction

Aubrun and Kari's construction works by encoding orbits of piecewise affine maps applied to real numbers. We will only apply their construction for piecewise linear maps, and begin this section with the necessary definitions.

2.1.1 Definitions

Definition 2.1 ((Representation by a sequence)). Let $i \in \mathbb{Z}$. We say that a binary biinfinite sequence $(x_k)_{k \in \mathbb{Z}} \in \{i, i+1\}^{\mathbb{Z}}$ represents a real number $x \in [i, i+1]$ if there exists a growing sequence of intervals $I_1 \subset I_2 \subset \dots \subseteq \mathbb{Z}$ of size at least $1, 2, \dots$ such that:

$$\lim_{k \rightarrow +\infty} \frac{\sum_{j \in I_k} x_j}{|I_k|} = x.$$

Note that if $(x_k)_{k \in \mathbb{Z}}$ is a representation of x , all the shifted sequences $(x_{l+k})_{k \in \mathbb{Z}}$ for every $l \in \mathbb{Z}$ are also representations of x . Note that a sequence $(x_k)_{k \in \mathbb{Z}}$ can *a priori* represent several distinct real numbers since different choices of interval sequences may make it converge to different points. A sequence always represents at least one real number by compactness of $[i, i+1]$.

Then, we define a generalization of piecewise linear maps: multiplicative systems. The main difference with piecewise linear maps is that points may have several images as definition intervals of different pieces might overlap.

Definition 2.2 (Multiplicative system). A *multiplicative system* is a finite set of non-zero linear maps

$$\mathcal{S} = \{f_1 : I_1 \rightarrow I'_1, \dots, f_k : I_k \rightarrow I'_k\}$$

with I_i and I'_i closed intervals of \mathbb{R} . Its *inverse* is defined to be

$$\mathcal{S}^{-1} = \{f_1^{-1} : I'_1 \rightarrow I_1, \dots, f_k^{-1} : I'_k \rightarrow I_k\}.$$

The *image* of $x \in \bigcup_i I_i$ is the set

$$\mathcal{S}(x) = \{y \in \bigcup_j I'_j \mid \exists i, f_i(x) = y\}.$$

The k -th iteration of \mathcal{S} on $x \in \bigcup_i I_i$ is then:

$$\begin{cases} \{y \in \mathbb{R} \mid \exists i_1, \dots, i_k, f_{i_k} \circ \dots \circ f_{i_1}(x) = y\} & \text{if } k > 0 \\ x & \text{if } k = 0 \\ \{y \in \mathbb{R} \mid \exists i_{-1}, \dots, i_{-k}, f_{i_{-k}}^{-1} \circ \dots \circ f_{i_{-1}}^{-1}(x) = y\} & \text{if } k < 0 \end{cases}$$

Note that if none of the intervals overlap, \mathcal{S} can be represented as a piecewise linear function and the definition of inverse and iteration coincide with the usual ones.

Definition 2.3 ((Immortal and periodic points)). Let $\mathcal{S} = \{f_1 : I_1 \rightarrow I'_1, \dots, f_k : I_1 \rightarrow I'_k\}$ be a multiplicative system. The real $x \in \mathbb{R}$ is *immortal* if for all $k \in \mathbb{Z}$,

$$\mathcal{S}^k(x) \cap \bigcup_i I_i \neq \emptyset$$

A *periodic point* for this system is a point $x \in \mathbb{R}$ such that there exists $k \in \mathbb{N}^*$ such that

$$x \in \mathcal{S}^k(x).$$

Definition 2.4 ((Level)). The *level* of $g \in BS(m, n)$ is the set $\mathcal{L}_g = \{ga^k \mid k \in \mathbb{Z}\}$.

When considering a tiling of $BS(m, n)$, given a line of tiles located between levels \mathcal{L}_g and $\mathcal{L}_{gb^{-1}}$, we talk about the upper side of the line to refer to level \mathcal{L}_g , and the lower side of the line to refer to level $\mathcal{L}_{gb^{-1}}$.

Definition 2.5 ((Height)). The *height* of $g \in BS(m, n)$ is, for any way of writing it as a word in $\{a, b, a^{-1}, b^{-1}\}^*$, its number of b 's minus its number of b^{-1} 's; it is denoted as $\|g\|_b$.

Since the only basic relation in $BS(m, n)$ uses one b and one b^{-1} , all writings of g as a word give the same height. Furthermore, it is actually the height of all elements in its level.

Definition 2.6 ((Multiplying tileset)). A set of tiles τ *multiplies* by $q \in \mathbb{Q}$ if for any tile $(t_1, \dots, t_m, l, r, b_1, \dots, b_n) \in \tau$ (see Fig. 1 for the notation),

$$q \frac{t_1 + \dots + t_m}{m} + l = \frac{b_1 + \dots + b_n}{n} + r. \quad (1)$$

Let τ be a tileset multiplying by $q \in \mathbb{Q}$. If we consider a line of N tiles of τ next to each other without tiling errors (as defined in Section 1.2), as left and right colors match, we can average Eq. (1):

$$qt + \frac{l}{N} = b + \frac{r}{N}. \quad (2)$$

where t is the average of the top labels of the line and b the average of the bottom ones. Therefore, if an infinite line has its upper side representing $x \in \mathbb{R}$ and its lower side representing $y \in \mathbb{R}$, taking the limit of Eq. (2) on a well chosen sequence of intervals gives:

$$qx = y.$$

Hence the name of *multiplying tileset* for τ .

2.1.2 A multiplying tileset

Let us define, in a fashion similar to [3], a couple of useful functions to build a multiplying tileset. Let $\alpha_{m,n} : \{a, b, a^{-1}, b^{-1}\}^* \rightarrow \mathbb{R}$ (or just α when m and n are clear) be defined by the recursion:

$$\begin{cases} \alpha(\varepsilon) = 0 \text{ where } \varepsilon \text{ is the empty word} \\ \alpha(wb) = \alpha(wb^{-1}) = \alpha(w) \\ \alpha(wa) = \alpha(w) + \left(\frac{n}{m}\right)^{\|w\|_b} \\ \alpha(wa^{-1}) = \alpha(w) - \left(\frac{n}{m}\right)^{\|w\|_b}. \end{cases}$$

The map α can be extended to elements of $BS(m, n)$, due to the fact that $\alpha(uba^m b^{-1}v) = \alpha(ua^n v)$ for any pair of words u and v in $\{a, b, a^{-1}, b^{-1}\}^*$: $\alpha(g)$ is then $\alpha(w)$ for any word representing g in the group.

Now, we define $\Phi : BS(m, n) \rightarrow \mathbb{R}^2$ as follows:

$$\Phi(g) = (\alpha(g), \|g\|_b).$$

The function Φ can be seen as a projection of every element of $BS(m, n)$ on the euclidean plane \mathbb{R}^2 .

Finally, let $\lambda : BS(m, n) \rightarrow \mathbb{R}$ be defined as

$$\lambda(g) = \frac{1}{m} \left(\frac{m}{n}\right)^{\|g\|_b} \alpha(g).$$


Let $q \in \mathbb{Q}$ and I an interval, let

$$\begin{aligned} t_j(g, x) &:= \lfloor (m\lambda(g) + j)x \rfloor - \lfloor (m\lambda(g) + (j-1))x \rfloor \text{ for } j = 1 \dots m \\ b_j(g, x) &:= \lfloor (n\lambda(g) + j)qx \rfloor - \lfloor (n\lambda(g) + (j-1))qx \rfloor \text{ for } j = 1 \dots n \\ l(g, x) &:= \frac{1}{m}q \lfloor m\lambda(g)x \rfloor - \frac{1}{n} \lfloor n\lambda(g)qx \rfloor \\ r(g, x) &:= \frac{1}{m}q \lfloor (m\lambda(g) + m)x \rfloor - \frac{1}{n} \lfloor (n\lambda(g) + n)qx \rfloor \end{aligned}$$

Then, we define the tileset $\tau_{q,I}$ as:

$$\tau_{q,I} = \{(t_1(g, x), \dots, t_m(g, x), l(g, x), r(g, x), b_1(g, x), \dots, b_n(g, x)) \mid g \in BS(m, n), x \in I\}.$$

One can show that Eq. (1) holds for these tiles.


 Figure 4: One tile from the tileset $\tau_{q,I}$.

Proposition 2.1 ([4, Proposition 6]). *Let $a \in \mathbb{Z}$, for any $I \subseteq [a, a + 1]$, $\tau_{q,I}$ is a tileset that multiplies by q .*

Let us define the *balanced representation* of x , which is the biinfinite sequence defined for any $z \in \mathbb{R}$ by

$$B_j(x, z) = \lfloor (z + j)x \rfloor - \lfloor (z + j - 1)x \rfloor.$$

Note that $B_j(x, z)$ does not depend on z , and can only take two values: $\lfloor x \rfloor$ or $\lfloor x \rfloor + 1$.

Proposition 2.2. *Let $q \in \mathbb{Q}$, $a \in \mathbb{Z}$ and $I \subseteq [a, a + 1]$. The upper side of any tile in $\tau_{q,I}$ is of the form*

$$B_j(x, m\lambda(g)), B_{j+1}(x, m\lambda(g)), \dots, B_{j+m}(x, m\lambda(g));$$

for $x \in I, j \in \mathbb{Z}, g \in BS(m, n)$. In particular, its labels are in $\{a, a + 1\}$. The lower side is of the form

$$B_j(qx, m\lambda(gb^{-1})), B_{j+1}(qx, m\lambda(gb^{-1})), \dots, B_{j+n}(qx, m\lambda(gb^{-1})).$$

Proof. Rewriting the top labels using balanced representation yields

$$B_j(x, m\lambda(g)), B_{j+1}(x, m\lambda(g)), \dots, B_{j+m}(x, m\lambda(g)).$$

Since each $B_j(x, m\lambda(g))$ is either $\lfloor x \rfloor$ or $\lfloor x \rfloor + 1$, and $x \in [a, a + 1]$, one obtains labels in $\{a, a + 1\}$. For the bottom side, note that $\lambda(gb^{-1}) = \frac{n}{m}\lambda(g)$, which gives labels

$$B_j(qx, m\lambda(gb^{-1})), B_{j+1}(qx, m\lambda(gb^{-1})), \dots, B_{j+n}(qx, m\lambda(gb^{-1})).$$

□

For our purpose we need to have a finite tileset, because subshifts use a finite alphabet.

Proposition 2.3. *Let $a \in \mathbb{Z}$, for any $I \subseteq [a, a + 1]$ the tileset $\tau_{q,I}$ is finite.*

Proof. Proposition 2.2 gives us that there are finitely many top and bottom labels. It remains to prove that there are also finitely many left and right labels.

First of all, one can check that $\lambda(ga^m) = \lambda(g) + 1$, and so $l(ga^m, x) = r(g, x)$. Consequently, we simply have to prove that l lies in a finite set. Let $q = \frac{q_1}{q_2}$ with $q_1, q_2 \in \mathbb{Z}$, and write

$$l(g, x) = \frac{nq \lfloor m\lambda(g)x \rfloor - m \lfloor n\lambda(g)qx \rfloor}{mn} = \frac{nq_1 \lfloor m\lambda(g)x \rfloor - mq_2 \lfloor n\lambda(g)qx \rfloor}{mnq_2}.$$

Since its numerator is an integer bounded by $-nq_1 =: k_1$ from below and $mq_2 =: k_2$ from above using usual inequalities on the floor function, we have that for any $g \in BS(m, n), x \in I$, $l(g, x)$ is in the finite set

$$\left\{ \frac{k_1}{mnq_2}, \frac{k_1 + 1}{mnq_2}, \dots, \frac{k_2}{mnq_2} \right\}.$$

□

Thanks to the multiplying property of $\tau_{q,[a,a+1]}$, we can use it to encode multiplicative systems in such a way that non-empty tilings corresponds to immortal points of the system. If we modify the tileset a little bit, we can encode multiplying systems defined on other intervals than $[a, a + 1]$.

Theorem 2.4. *Let $\mathcal{S} = \{f_1, \dots, f_N\}$ be a multiplicative system with*

$$\begin{aligned} f_i : I_i &\rightarrow \mathbb{R} \\ x &\mapsto q_i x, \end{aligned}$$

$q_i \in \mathbb{Q}$ and I_i interval with rational bounds included in some $[a_i, a_i + 1]$, $a_i \in \mathbb{Z}$. We can explicitly and algorithmically build an SFT $Y_{\mathcal{S}}$ with the following properties:

1. *any top of a line of tiles in a configuration $y \in Y_{\mathcal{S}}$ represents at least one real $x \in \bigcup_i I_i$.*
2. *if the top of a line of tiles represents a real $x \in \bigcup_i I_i$, then the bottom of that line represents a real in $\mathcal{S}(x)$;*
3. *$Y_{\mathcal{S}} \neq \emptyset$ if and only if \mathcal{S} has an immortal point;*

Proof. We build a tileset τ performing the computation by the linear functions $f_i : [a_i, a_i + 1] \rightarrow \mathbb{R}$; i.e. the linear maps with bigger intervals than the ones defining \mathcal{S} . In order to encode the multiplication correctly, one cannot simply take the union of all $\tau_i := \tau_{q_i, [a_i, a_i + 1]}$, because tiles coming from different f_i could be mixed on a single line. In order to "synchronize" the computations on every line, we create a product alphabet with the left and right colors of the tiles, and the number of the current function being used. This ensures that one line can have tiles from only one of the $\tau_{q_i, [a_i, a_i + 1]}$. Formally,

$$\tau = \left\{ (t_1, \dots, t_m, (l, i), (r, i), b_1, \dots, b_n) \mid (t_1, \dots, t_m, l, r, b_1, \dots, b_n) \in \tau_{q_i, [a_i, a_i + 1]} \right\}.$$

This way, we can interpret any line of a tiling by τ as being "of color" i for some $i \in \{1, \dots, N\}$.

Next, we restrict the intervals of reals that can be represented in each line of the SFT. Let us write $I_i = [a_i + \frac{d_1^i}{e_1^i}, a_i + 1 - \frac{d_2^i}{e_2^i}]$, for each i . $Y_{\mathcal{S}}$ will be X^τ with the following additional local constraints: on each line of color i , we force the upper side labels to respect

- every e_1^i consecutive labels must contain at least d_1^i labels $a_i + 1$; (3)

- every e_2^i consecutive labels must contain at least d_2^i labels a_i . (4)

Recall that the top labels of any line of color i only has labels a_i and $a_i + 1$ by definition of τ_i . From this, we can also deduce

- every e_1^i consecutive labels must contain at most $e_1^i - d_1^i$ labels a_i ;
- every e_2^i consecutive labels must contain at most $e_2^i - d_2^i$ labels $a_i + 1$.

Since these constraints are on e_1^i or e_2^i consecutive labels, and any other constraint from the τ_i 's is on neighboring tiles, $Y_{\mathcal{S}}$ is an SFT.

First, assume that $Y_{\mathcal{S}}$ is non-empty and contains some configuration y . Then, the sequence at the top of each level $\mathcal{L}_{b^k}, k \in \mathbb{Z}$ represents at least one real x_k , since it is a sequence made of at most two integers. Thanks to the multiplying property of each τ_i (Proposition 2.1), and the fact that the local rules of every line impose that any real represented belongs to an interval I_i , $(x_k)_{k \in \mathbb{Z}}$ is an infinite orbit of \mathcal{S} . Indeed, consider a real x represented by the upper side of a line of color i . We prove that $x \in I_i = [a + \frac{d_1}{e_1}, a + 1 - \frac{d_2}{e_2}]$ (we drop the i in this paragraph to make notations lighter). Let us write $(J_l)_{l \in \mathbb{N}}$ the intervals from Definition 2.1 (denoted as (I_i) there) on which we compute the mean for the represented real, and let r_l

denote the proportion of a 's over $(a + 1)$'s in the line representing x restricted to J_l . Thanks to the two conditions Eq. (3) and Eq. (4) (and the two we deduced), we have, for all l in \mathbb{N} ,

$$\frac{d_2}{e_2 - d_2} \leq r_l \leq \frac{e_1 - d_1}{d_1}. \quad (5)$$

Moreover, since x is the limit of the means computed on each J_l , one can show that

$$x = \lim_{l \rightarrow \infty} a + \frac{1}{1 + r_l}.$$

Using the left inequality of Eq. (5) gives that

$$\frac{1}{1 + r_l} \leq \frac{e_2 - d_2}{e_2} = 1 - \frac{d_2}{e_2}.$$

So

$$x \leq a + 1 - \frac{d_2}{e_2}.$$

Similarly, the right inequality of Eq. (5) gives

$$x \geq a + \frac{d_1}{e_1},$$

and so $x \in I_i$. This notably ensures Item 1 and, by the use of the τ_i 's, Item 2.

Now, assume that S has an immortal point x . Then there exists a sequence $(i_k)_{k \in \mathbb{Z}} \in \{1, \dots, N\}^{\mathbb{Z}}$ such that if we define

$$x_k = \begin{cases} x & \text{if } k = 0 \\ f_{i_{k-1}}(x_{k-1}) = q_{i_{k-1}} x_{k-1} & \text{if } k > 0, \\ f_{i_k}^{-1}(x_{k+1}) = \frac{1}{q_{i_k}} x_{k+1} & \text{if } k < 0 \end{cases},$$

then for all $k \in \mathbb{Z}$, $x_k \in \bigcup_i I_i$. For every $g \in BS(m, n)$, we place at g a tile from the tileset τ_{i_k} with $k = -\|g\|_b$, with colors:

$$\begin{aligned} t_j : t_j(g, x_k) &= \lfloor (m\lambda(g) + j) x_k \rfloor - \lfloor (m\lambda(g) + (j - 1)) x_k \rfloor \quad \text{for } j = 1 \dots m \\ b_j : b_j(g, x_k) &= \lfloor (n\lambda(g) + j) q_{i_k} x_k \rfloor - \lfloor (n\lambda(g) + (j - 1)) q_{i_k} x_k \rfloor \quad \text{for } j = 1 \dots n \\ l : l(g, x_k) &= \left(\frac{1}{m} q_{i_k} \lfloor m\lambda(g) x_k \rfloor - \frac{1}{n} \lfloor n\lambda(g) q_{i_k} x_k \rfloor, i_k \right) \\ r : r(g, x_k) &= \left(\frac{1}{m} q_{i_k} \lfloor (m\lambda(g) + m) x_k \rfloor - \frac{1}{n} \lfloor (n\lambda(g) + n) q_{i_k} x_k \rfloor, i_k \right) \end{aligned}$$

These tiles are obviously from the tileset τ . Recall that we have

$$\lambda(ga^m) = \lambda(g) + 1, \quad (6)$$

$$\lambda(gb) = \frac{m}{n} \lambda(g).$$

And therefore,

$$\begin{aligned} l(ga^m, x_k) &= \left(\frac{1}{m} q_{i_k} \lfloor m\lambda(ga^m) x_k \rfloor - \frac{1}{n} \lfloor n\lambda(ga^m) q_{i_k} x_k \rfloor, i_k \right) \\ &= \left(\frac{1}{m} q_{i_k} \lfloor m(\lambda(g) + 1) x_k \rfloor - \frac{1}{n} \lfloor n(\lambda(g) + 1) q_{i_k} x_k \rfloor, i_k \right) = r(g, x_k), \end{aligned}$$

and for any $j \in \{1, \dots, m\}, p \in \{1, \dots, n\}$,

$$\begin{aligned}
& b_p(ga^{j-p}b, x_{-||ga^{j-p}b||_b}) \\
&= \left[\left(n\lambda(ga^{j-p}b) + p \right) q_{i_{-||ga^{j-p}b||_b}} x_{-||ga^{j-p}b||_b} \right] - \left[\left(n\lambda(ga^{j-p}b) + (p-1) \right) q_{i_{-||ga^{j-p}b||_b}} x_{-||ga^{j-p}b||_b} \right] \\
&= \left[\left(n\frac{m}{n}\lambda(ga^{j-p}) + p \right) q_{i_{-||g||_{b-1}}} x_{-||g||_{b-1}} \right] - \left[\left(n\frac{m}{n}\lambda(ga^{j-p}) + (p-1) \right) q_{i_{-||g||_{b-1}}} x_{-||g||_{b-1}} \right] \\
&= \left[\left(m\left(\lambda(g) + \frac{j-p}{m}\right) + p \right) q_{i_{-||g||_{b-1}}} x_{-||g||_{b-1}} \right] - \left[\left(m\left(\lambda(g) + \frac{j-p}{m}\right) + (p-1) \right) q_{i_{-||g||_{b-1}}} x_{-||g||_{b-1}} \right] \\
&= \left[(m\lambda(g) + j) q_{i_{-||g||_{b-1}}} x_{-||g||_{b-1}} \right] - \left[(m\lambda(g) + (j-1)) q_{i_{-||g||_{b-1}}} x_{-||g||_{b-1}} \right] \\
&= \left[(m\lambda(g) + j) x_{-||g||_b} \right] - \left[(m\lambda(g) + (j-1)) x_{-||g||_b} \right] \\
&= t_j(g, x_{-||g||_b})
\end{aligned}$$

It remains to show that the labels at the top of every line follow the conditions (3) and (4). Fix some $g \in BS(m, n)$ and consider the level \mathcal{L}_g . Fix $k = -||g||_b$, denote $x := x_k, i := i_k$ and drop the i in the other variables to simplify notations. For $j \in \mathbb{Z}$, we write w_j for the label at position j of the considered level, with $t_1(g, x_k)$ being position 1. Remark that

$$w_j = \lfloor (m\lambda(g) + j)x \rfloor - \lfloor (m\lambda(g) + (j-1))x \rfloor$$

holds for all $j \in \mathbb{Z}$ thanks to Eq. (6). Assume that $x \geq a + \frac{d_1}{e_1}$. Let us write N_{j_0} the number of labels a that appear in the word $u = w_{j_0+1}w_{j_0+2} \cdots w_{j_0+e_1}$. If we sum all the labels of u , we have on the one hand

$$\sum_{j=j_0+1}^{j_0+e_1} w_j = N_{j_0}a + (e_1 - N_{j_0})(a+1) = e_1(a+1) - N_{j_0}.$$

And on the other hand,

$$\sum_{j=j_0+1}^{j_0+e_1} w_j = \lfloor (m\lambda(g) + j_0 + e_1)x \rfloor - \lfloor (m\lambda(g) + j_0)x \rfloor.$$

Therefore,

$$e_1(a+1) - N_{j_0} > (m\lambda(g) + j_0 + e_1)x - 1 - (m\lambda(g) + j_0)x = e_1x - 1 \geq e_1a + d_1 - 1,$$

which can be rearranged to

$$N_{j_0} \leq e_1 - d_1,$$

which implies (3). In the same way, if we assume $x \leq a + 1 - \frac{d_2}{e_2}$, we can show $N_{j_0} \geq d_2$, which is exactly (4). This shows Item 3. \square

Remark 2.5. Instead of considering a multiplicative system with rational bounds for the intervals, one can dilate them and encode a dilated system with integer bounds, which is equivalent to the original. For instance, instead of using a linear function f on $[a + \frac{d_1}{e_1}, a + 1 - \frac{d_2}{e_2}]$, one can consider that function f multiplied by e_1e_2 , on $[ae_1e_2 + d_1e_2, ae_1e_2 + e_1e_2 - d_2e_1]$ which has integer bounds. However, although this would yield a shorter proof, the results would only hold through some sort of conjugation, and we wanted to effectively build a Wang tileset for any multiplicative system.

Note that we only build a multiplying tileset in the current paper. Aubrun and Kari provided the details of the encoding of any finite set of affine maps into a tileset of $BS(m, n)$ in [4].

2.2 A weakly aperiodic SFT on $BS(m, n)$

The SFT $Y_{\mathcal{S}}$ previously defined on a given $BS(m, n)$ is also linked to the periodicity of \mathcal{S} .

Theorem 2.6. *If \mathcal{S} has no periodic point, then $Y_{\mathcal{S}}$ is a weakly aperiodic SFT.*

Proof. We prove the contrapositive.

Assume that $Y_{\mathcal{S}}$ has a strongly periodic configuration y , i.e. $|\text{Orb}_{BS(m,n)}(y)| = i$ with $i \in \mathbb{N}$. In particular, each set $\{ga^k g^{-1} \cdot y \mid k \in \mathbb{Z}\}$ is finite of cardinality lesser than i , and therefore for every $g \in BS(m, n)$, there exists $k_g \leq i$ such that $ga^{k_g} g^{-1} \cdot y = y$. If we define $p = i!$, we obtain that for all $g \in BS(m, n)$, $ga^p g^{-1} \cdot y = y$.

Let $g \in BS(m, n)$. Let $h \in \mathcal{L}_g$, i.e. there exists some $k \in \mathbb{Z}$ so that $h = ga^k$. Then

$$\begin{aligned} y_h &= (ga^p g^{-1} \cdot y)_h \\ &= y_{ga^{-p} g^{-1} h} \\ &= y_{ga^{k-p}} \\ &= y_{ha^{-p}} \end{aligned}$$

which means that the level \mathcal{L}_g is p -periodic.

Therefore any level is p -periodic, and consequently represents a unique rational number $\frac{c}{p}$. Since the alphabet of $Y_{\mathcal{S}}$ is finite, there are only finitely many different such rationals. Consequently, there exist two levels \mathcal{L}_{g^l} and \mathcal{L} with $l > 0$ that represent the same rational number x . By the multiplicative property of $Y_{\mathcal{S}}$,

$$x \in \mathcal{S}^l(x),$$

which means that x is a periodic point for \mathcal{S} . □

The consequence of this theorem is that, to obtain a weakly aperiodic SFT on $BS(m, n)$, we only need to explicitly build a multiplicative system with an immortal point but no periodic points. For the rest of this section, let

$$\begin{aligned} f_1 &: \left[\frac{1}{3}, 1\right] \rightarrow \left[\frac{2}{3}, 2\right] \\ & \quad x \mapsto 2x \end{aligned}$$

$$\begin{aligned} f_2 &: [1, 2] \rightarrow \left[\frac{1}{3}, \frac{2}{3}\right] \\ & \quad x \mapsto \frac{1}{3}x \end{aligned}$$

and

$$\mathcal{S}_0 = \{f_1, f_2\}.$$

It is easy to see that this system has immortal points (in fact, all points of $[\frac{1}{3}, 2]$ are immortal).

Proposition 2.7. *\mathcal{S}_0 has immortal points.*

Additionally, since 2 and 3 are relatively prime:

Proposition 2.8. *\mathcal{S}_0 has no periodic points.*

Corollary 2.9. *$Y_{\mathcal{S}_0}$ is non-empty and weakly aperiodic.*

However, this construction does not avoid weakly periodic configurations when m and n are not 1, as already remarked by Aubrun and Kari.

Proposition 2.10. *For any $m, n > 1$, $Y_{\mathcal{S}_0}$ on $BS(m, n)$ contains a weakly periodic tiling, with period $p = bab^{-1}aba^{-1}b^{-1}a^{-1}$.*

To prove it, we need the following lemma:

Lemma 2.11. *Let $p = bab^{-1}aba^{-1}b^{-1}a^{-1}$. For any $g \in BS(m, n)$, $\alpha(pg) = \alpha(g)$.*

Proof. Since $\|p\|_b = 0$, using the definition of α it is easy to show that $\alpha(pg) = \alpha(p) + \alpha(g)$ by recurrence on the length of g . Then, $\alpha(p) = \frac{n}{m} + 1 - \frac{n}{m} - 1 = 0$. \square

Proof of Proposition 2.10. We happen to have built a periodic configuration already: the one from the proof of Theorem 2.4.

Indeed, let x be any real in $[\frac{1}{3}, 2]$, since they are all immortal for \mathcal{S}_0 . Then there exists a sequence $(i_k)_{k \in \mathbb{Z}} \in \{1, 2\}^{\mathbb{Z}}$ such that if we define

$$x_k = \begin{cases} x & \text{if } k = 0 \\ f_{i_{k-1}}(x_{k-1}) = q_{i_{k-1}}x_{k-1} & \text{if } k > 0, \\ f_{i_k}^{-1}(x_{k+1}) = \frac{1}{q_{i_k}}x_{k+1} & \text{if } k < 0 \end{cases}$$

then for all $k \in \mathbb{Z}$, $x_k \in [\frac{1}{3}, 2]$. For every $g \in BS(m, n)$, we place at g a tile from the tileset τ_{i_k} with $k = -\|g\|_b$, with colors:

$$\begin{aligned} t_j : t_j(g, x_k) &= \lfloor (m\lambda(g) + j)x_k \rfloor - \lfloor (m\lambda(g) + (j-1))x_k \rfloor \quad \text{for } j = 1 \dots m \\ b_j : b_j(g, x_k) &= \lfloor (n\lambda(g) + j)q_{i_k}x_k \rfloor - \lfloor (n\lambda(g) + (j-1))q_{i_k}x_k \rfloor \quad \text{for } j = 1 \dots n \\ l : l(g, x_k) &= \left(\frac{1}{m}q_{i_k} \lfloor m\lambda(g)x_k \rfloor - \frac{1}{n} \lfloor n\lambda(g)q_{i_k}x_k \rfloor, i_k \right) \\ r : r(g, x_k) &= \left(\frac{1}{m}q_{i_k} \lfloor (m\lambda(g) + m)x_k \rfloor - \frac{1}{n} \lfloor (n\lambda(g) + n)q_{i_k}x_k \rfloor, i_k \right) \end{aligned}$$

We already checked that the resulting tiling y was in $Y_{\mathcal{S}_0}$, see the proof of Theorem 2.4. It remains to show that the tiles at g and at pg are the same for all $g \in BS(m, n)$, to conclude that $p^{-1} \cdot y = y$, or equivalently that p is a period of y . This is actually surprisingly easily, considering that for any $g \in BS(m, n)$, $\|pg\|_b = \|g\|_b$ so they use the same integer k and the same real x_k ; and $\lambda(g) = \lambda(pg)$, see Lemma 2.11. The tiles have the same labels as a consequence of this.

The y consequently defined is p -periodic. The only thing left to show is that p is a nontrivial element of $BS(m, n)$ as long as $m, n > 1$. Since it is freely reduced and does not contain ba^mb^{-1} or $b^{-1}a^nb$ as subwords, and since $BS(m, n)$ is a HNN extension $\mathbb{Z} *_\alpha$ with $\alpha: m\mathbb{Z} \rightarrow n\mathbb{Z}$, we can apply Britton's Lemma: p cannot be the neutral element.

Therefore $Y_{\mathcal{S}_0}$ contains a configuration with a nontrivial period, and consequently is not strongly aperiodic. \square

2.3 A deeper understanding of the configurations

We first present additional results on this tileset $Y_{\mathcal{S}_0}$ of $BS(m, n)$. Most of the ideas present in this section were already present in [11] in the context of tilings of the plane.

For a given line \mathcal{L}_g , we define the sequence $u_g := (y_{ga^i})_{i \in \mathbb{Z}}$ to be the sequence of digits on the line \mathcal{L}_g (its origin depending on g).

Let f be the following bijective continuous map:

$$f: \left[\frac{1}{3}, 2 \right] / \frac{1}{3} \sim 2 \rightarrow \left[\frac{1}{3}, 2 \right] / \frac{1}{3} \sim 2$$

$$f(x) = \begin{cases} f_1(x) = 2x & \text{if } x \in (\frac{1}{3}, 1) \\ f_2(x) = \frac{1}{3}x & \text{if } x \in (1, 2) \\ \frac{2}{3} & \text{if } x = 1 \\ \frac{2}{3} & \text{if } x = \frac{2}{3} \end{cases}$$

f is strongly related to \mathcal{S}_0 because for any $x \in [\frac{1}{3}, 2]$, for any $k \in \mathbb{Z}$,

$$\mathcal{S}_0^k(x) \Big/_{\frac{1}{3} \sim 2} = f^k(x). \quad (7)$$

due to the fact that for our particular \mathcal{S}_0 , $\mathcal{S}_0^k(x) \subseteq [\frac{1}{3}, 2]$.

An easy consequence is the following:

Lemma 2.12. *Let $y \in Y_{\mathcal{S}_0}$. Let $g \in BS(m, n)$. Let $x \in [\frac{1}{3}, 2]$ be a real represented by u_g ; then $u_{gb^{-1}}$ represents $f(x)$ (which means either $\frac{1}{3}$ or 2 if $f(x) = \bar{2}$).*

Proof. u_g represents at least one such x because of Item 1 of Theorem 2.4. The rest is due to Item 2 of Theorem 2.4 and Eq. (7). \square

It turns out that with our choice of multiplicative system \mathcal{S}_0 , any line can represent only one real number. We need several lemmas to prove this; all inspired by [11].

Lemma 2.13. *Let ϕ be defined as follows:*

$$\begin{aligned} \phi: [\tfrac{1}{3}, 2] \Big/_{\frac{1}{3} \sim 2} &\rightarrow [0, 1] \Big/_{0 \sim 1} \\ \phi(x) &= \frac{\log(x) + \log(3)}{\log(2) + \log(3)} \pmod{1} \end{aligned}$$

ϕ is a correctly defined mapping that conjugates the dynamical systems $([\frac{1}{3}, 2] \Big/_{\frac{1}{3} \sim 2}, t, f)$ and $([0, 1] \Big/_{0 \sim 1}, t', \phi \circ f \circ \phi^{-1})$, where t and t' are the usual topologies on the considered sets.

Proof. Since the action considered on $[0, 1] \Big/_{0 \sim 1}$ is $\phi \circ f \circ \phi^{-1}$, one only needs to check that ϕ is bijective and continuous to conclude that it yields a conjugation.

ϕ is clearly continuous everywhere except on $\bar{2}$; there, one can check that the left and right limits both lead to $\phi(\bar{2}) = 0$ which is correctly defined.

ϕ^{-1} is defined by $\Phi(y) = \frac{6^y}{3}$ except with collapsed images for 0 and 1. \square

Lemma 2.14. *The map $r := \phi \circ f \circ \phi^{-1}$ can be considered a rotation of irrational angle $\frac{\log(2)}{\log(2) + \log(3)}$ when identifying $[0, 1] \Big/_{0 \sim 1}$ and \mathbb{S}^1 .*

Proof. For every $\alpha \in \phi([\frac{1}{3}, 1])$,

$$\begin{aligned} \phi \circ f \circ \phi^{-1}(\alpha) &= \phi(2\phi^{-1}(\alpha)) \\ &= \frac{\log(2) + \log(\phi^{-1}(\alpha)) + \log(3)}{\log(2) + \log(3)} \pmod{1} \\ &= \alpha + \frac{\log(2)}{\log(2) + \log(3)} \pmod{1}. \end{aligned}$$

Similarly, for every $\alpha \in \phi((1, 2))$, one has

$$\begin{aligned} \phi \circ f \circ \phi^{-1}(\alpha) &= \phi(\tfrac{1}{3}\phi^{-1}(\alpha)) \\ &= \frac{\log(\phi^{-1}(\alpha))}{\log(2) + \log(3)} \pmod{1} \\ &= \alpha + \frac{\log(2)}{\log(2) + \log(3)} \pmod{1}. \end{aligned}$$

Finally, $\phi \circ f \circ \phi^{-1}(0) = \phi(f(\bar{2})) = \phi(\frac{2}{3}) = \frac{\log(2)}{\log(2) + \log(3)}$. \square

We now have the tools to prove the following key lemma:

Lemma 2.15. (*Uniqueness of representation*) *For any $y \in Y_{S_0}$, for any $g \in BS(m, n)$, the sequence $u_g = (y_{g^i})_{i \in \mathbb{Z}}$ represents a unique real number.*

Proof. Assume that u_g represents two distinct reals x and $z \in [\frac{1}{3}, 2]$. They cannot be $\frac{1}{3}$ and 2 because u_g uses only digits in $\{0, 1\}$ or in $\{1, 2\}$. Therefore they also are distinct reals in $[\frac{1}{3}, 2] / \frac{1}{3} \sim 2$.

For any $k \in \mathbb{Z}$, notice that $f^k(x) = \phi^{-1} \circ r^k \circ \phi(x)$ and same for z , from Lemma 2.13. We will study the behavior of $\phi(x)$ and $\phi(z)$ under iterations of r .

The angle $\frac{\log(2)}{\log(2)+\log(3)}$, of which r is a rotation by Lemma 2.14, is irrational. As a consequence, the sets $\{r^k \circ \phi(x) \mid k \in \mathbb{N}\}$ and $\{r^k \circ \phi(z) \mid k \in \mathbb{N}\}$ are both dense in \mathbb{S}^1 .

We introduce $d_{arc}(e^{2i\pi\theta}, e^{2i\pi\psi}) = m(\psi - \theta) \in [0, 1)$ for $\theta, \psi \in \mathbb{R}$, where $m(\psi - \theta)$ is the only real in $[0, 1)$ congruent to $\psi - \theta \pmod{1}$. We call d_{arc} the *oriented arc distance* (measured counterclockwise) between two elements of \mathbb{S}^1 . It is not a distance *per se* since it is not symmetric and has no triangular inequality, but its basic properties will suffice here. Since r is a rotation, it is easy to check that it preserves d_{arc} . Hence we have that $\forall k \in \mathbb{N}, d_{arc}(r^k \circ \phi(x), r^k \circ \phi(z))$ is constant equal to some $c \in [0, 1[$. Up to considering $d_{arc}(r^k \circ \phi(z), r^k \circ \phi(x))$ instead, and doing the following reasoning by swapping x and z , we can assume that $c \leq \frac{1}{2}$.

Let us split \mathbb{S}^1 between $A = \phi([\frac{1}{3}, 1))$, $B = \phi([1, 2))$, and $\{\phi(2)\} = \{\phi(\frac{1}{3})\} = \{0\}$. We want to show that there is some $l \in \mathbb{N}$ for which $r^l \circ \phi(x) \in B$ and $r^l \circ \phi(z) \in A$.

By density of $\{r^k \circ \phi(x) \mid k \in \mathbb{Z}\}$, there exists some $k_0 \in \mathbb{N}$ such that $d_{arc}(r^{k_0} \circ \phi(x), 0) < c$ and $r^{k_0} \circ \phi(x) \in B$. We cannot have $r^{k_0} \circ \phi(z) = 0$ without contradicting the previous inequality, hence it is either in A or in B . But if it was in B , then the arc from $r^{k_0} \circ \phi(x)$ to $r^{k_0} \circ \phi(z)$ would contain all of A . This is not possible because $|A|_{d_{arc}} > \frac{1}{2} \geq c$.

Hence there exists $l = k_0 \in \mathbb{N}$ such that $r^l \circ \phi(x) \in B$ and $r^l \circ \phi(z) \in A$.




Figure 5: Preservation of the oriented arc distance d_{arc} by r and intersection of the arc $(r^l \circ \phi(x), r^l \circ \phi(z))$ and the boundary between A and B .

Since $r^l = \phi \circ f^l \circ \phi^{-1}$, and considering the definitions of A and B , $f^l(z) \in (\frac{1}{3}, 1)$ and $f^l(x) \in [1, 2)$. This would cause $f^l(z)$ to be represented by a sequence of 0's and 1's (with an infinite number of 0's) and $f^l(x)$ by a sequence of 1's and 2's. However, the SFT Y_{S_0} is built such that a line contains only elements in $\{0, 1\}$ or $\{1, 2\}$, but not both (see proof of Theorem 2.4): this is a contradiction.

Therefore, x and z must be equal, hence the uniqueness of the real number represented by a given sequence. \square

Using previous results, we are now able to prove for $BS(m, n)$ that the real represented by the sequence u_g only depends on $\|g\|_b$, its “depth” in the Cayley graph.

Lemma 2.16. *Let $y \in Y_{S_0}$, and e the identity of $BS(m, n)$. Let x be the unique real represented by the sequence u_e . Then for every $g \in BS(m, n)$, u_g represents $f^{-\|g\|_b}(x)$ (with a choice between $\frac{1}{3}$ and 2, possibly different for different g 's, if the resulting value is $\bar{2}$).*

Proof. We prove the result by reasoning on words $w \in \{a, b, a^{-1}, b^{-1}\}^*$, by induction on their length. Note that we have no need of proving that different w 's representing the same g yield the same result, since this is guaranteed by Lemma 2.15.

The result is true for $g = e$.

Suppose the result is true for words of length $n \in \mathbb{N}_0$. Let w be a word of length n . Then:

- u_{wa} and $u_{wa^{-1}}$ represent the same real as u_w since they are the same sequence up to an index shift;
- $u_{wb^{-1}}$ represents $f(f^{-\|w\|_b}(x))$ due to Lemma 2.12 and the induction hypothesis, which is $f^{-\|wb^{-1}\|_b}(x)$;
- suppose u_{wb} represents y ; then u_w represents $f(y)$ due to Lemma 2.12. Then we have, by induction, $y = f^{-\|w\|_b^{-1}}(x) = f^{-\|wb\|_b}(x)$.

□

Remark 2.17. The previous proof heavily relies on the fact that f is a bijection on $[\frac{1}{3}, 2] / \frac{1}{3} \sim 2$, and that we do not have to differentiate between $\frac{1}{3}$ and 2 there.

2.4 A strongly aperiodic SFT on $BS(1, n)$

If m or n is equal to 1, then the previous weak period of Proposition 2.10 does not work anymore – it is a trivial element. In fact, we prove in this section that for $BS(1, n)$, Y_{S_0} is strongly aperiodic.

One key property of $BS(1, n)$ is that there is a simple quasi-normal form for all its elements.

Lemma 2.18. *(Quasi-normal form in $BS(1, n)$) For every $g \in BS(1, n)$, there are integers $k, m \in \mathbb{N}_0$ and $l \in \mathbb{Z}$ such that $g = b^{-k}a^l b^m$.*

Proof. From the definition of $BS(1, n)$, we have that $ba = a^nb$ (1), $ba^{-1} = a^{-n}b$ (2), $ab^{-1} = b^{-1}a^n$ (3) and $a^{-1}b^{-1} = b^{-1}a^{-n}$ (4). Consequently, taking an element of $BS(1, n)$ as a word w written with a and b , we can:

- Move each positive power of b to the right of the word using (1) and (2) repeatedly;
- Move each negative power of b to the left of the word using (3) and (4) repeatedly;

so that we finally get a form for the word w which is: $b^{-k}a^l b^m$ with $k, m \in \mathbb{N}_0$ and $l \in \mathbb{Z}$. □

Remark 2.19. A general normal form – the same, with k imposed to be minimal – can be obtained from Britton’s Lemma. The form obtained here is *not* unique ($a = b^{-1}a^nb$ for instance), but we use it because it admits a simple self-contained proof, and it is enough for what follows: the sum $m - k$ is constant for all writings of a given group element, hence we name it “quasi-normal”.

Proof. Indeed, suppose we have $b^{-k}a^l b^m = b^{-k'}a^{l'}b^{m'}$. Then

$$\begin{aligned} b^{-k}a^l &= b^{-k'}a^{l'}b^{-(m-m')} \\ &= b^{-k'-(m-m')}a^{l'}n^{m-m'} \end{aligned}$$

Hence we get $a^{l'n^{m-m'}-l} = b^{-k+k'+m-m'}$. Since it is clear that $a^i = b^j$ if and only if $i = j = 0$ in $BS(1, n)$, we obtain $m - k - (m' - k') = 0$ which is what we wanted. \square

This quasi-normal form is the only thing that missed to prove the following.

Theorem 2.20. *For every $n \geq 2$, the Baumslag-Solitar group $BS(1, n)$ admits a strongly aperiodic SFT.*

Proof. Let $y \in Y_{S_0}$, and $g \in \text{Stab}_{BS(1,n)}(y)$. Using Lemma 2.18, we can write $g = b^{-k}a^l b^m$ with $k, m \in \mathbb{N}_0, l \in \mathbb{Z}$.

Let x be the real represented by u_e . By Lemma 2.16, u_g represents $f^{k-m}(x)$. Since $g \in \text{Stab}_{BS(1,n)}(y)$, $u_g = u_e$ and so $f^{k-m}(x) = x$ by the uniqueness of the representation from Lemma 2.15. The aperiodicity of f then implies that $k = m$.

Let us assume $l \neq 0$. Then $g = b^{-k}a^l b^k$ and $g^n = b^{-k}(a^n)^l b^k$. We can reduce g^n to $b^{-k+1}a^l b^{k-1}$ using the relation $a^n = bab^{-1}$. More generally, we notice that for any positive integer i , iterating the process i times, we obtain that $g^{n^i} = b^{-k+i}a^l b^{k-i} \in \text{Stab}_{BS(1,n)}(y)$.

Since for all i , $g^{n^i} \in \text{Stab}_{BS(1,n)}(y)$, we can obtain a contradiction with an argument similar to Prop 6. of [3]. We have $b^j a^l b^{-j} \in \text{Stab}_{BS(1,n)}(y)$ for any $j \geq -k$. This means that $u_{bj} = u_{bj} a^l$ hence u_{bj} is a l -periodic sequence. We have a finite number of said sequences, since they can only use digits among $\{0, 1, 2\}$. Consequently, there are $j_1 \neq j_2$ such that the two levels $\mathcal{L}_{b^{j_1}}$ and $\mathcal{L}_{b^{j_2}}$ read the same sequence (up to index translation). These two levels represent respectively $f^{j_1}(x)$ and $f^{j_2}(x)$ due to Lemma 2.16, and since the two sequences on these levels are the same, $f^{j_1}(x) = f^{j_2}(x)$. This equality contradicts the fact that f has no periodic point, since we had $j_1 \neq j_2$.

As a consequence, any non-trivial $g \in BS(1, n)$ cannot be in $\text{Stab}_{BS(1,n)}(x)$, and we finally get that $\text{Stab}_{BS(1,n)}(x) = \{e\}$: Y_{S_0} is strongly aperiodic. \square

Following Theorem 2.20, a question remains: is the strong aperiodicity of Aubrun and Kari's SFT a property of the group $BS(1, n)$ itself, or does it only arise on carefully chosen SFTs, as Y_{S_0} ? Is this because $BS(1, n)$ behaves like \mathbb{Z}^2 and all its weakly aperiodic SFTs are also strongly aperiodic, or does Aubrun and Kari's construction happen to be "too much aperiodic"? It turns out that the latter is the correct answer, as we build in the following section an SFT on $BS(1, n)$ that is weakly but not strongly aperiodic.

3 A weakly but not strongly aperiodic SFT on $BS(1, n)$

Our weakly but not strongly aperiodic SFT will work by encoding specific substitutions into $BS(1, n)$. Indeed, the Cayley graph of $BS(1, n)$ is very similar to orbit graphs of uniform substitutions (see for example [2, 9] for a definition of orbit graphs and another example of a Cayley graph similar to an orbit graph). In this section, we find a set of substitutions that are easy to encode in $BS(1, n)$ (Section 3.1), and show how to do it (Section 3.2).

3.1 The substitutions σ_i

Let $\mathcal{A} = \{0, 1\}$. For $r \in \{0, \dots, n-1\}$, let $\sigma_r : \mathcal{A} \rightarrow \mathcal{A}^n$ be the following substitution:

$$\sigma_r : \begin{cases} 0 \mapsto 0^{n-r-1}10^r \\ 1 \mapsto 0^n \end{cases} .$$

We may also write $\sigma = \sigma_0$ and call the other ones the *shifts* of σ .

Note that, for $c \in \{0, 1\}$ and $i \in \{0, \dots, n-1\}$, $\sigma_r(c)_i = 0$ if and only if $c = 0$ and $i = n-r-1$ (starting to count from 0 the indices of the word $\sigma_r(c)$).

All $\sigma_r(0)$ are cyclic permutations of the same finite word. Denote ρ the *shift action* on a biinfinite word u , i.e. $\rho^j(u)_i = u_{i+j}$, as a way to write the action of \mathbb{Z} on $\{0, 1\}^{\mathbb{Z}}$.

Lemma 3.1. *For any biinfinite word $u \in \mathcal{A}^{\mathbb{Z}}$, any $i, r \in \{0, \dots, n-1\}$ and $j \in \mathbb{Z}$,*

$$(\sigma_r \circ \rho^j(u))_i = \sigma_r(u_j)_i = (\sigma_r(u))_{nj+i}.$$

Proof. For $i \in \{0, \dots, n-1\}$, $\sigma_r(\rho^j(u))_i$ depends on the letter of $\rho^j(u)$ at position 0 only, that is u_j (See Fig. 6), hence $\sigma_r(\rho^j(u))_i = \sigma_r(\rho^j(u))_0 = \sigma_r(u_j)_i$.

Similarly, the letter $(\sigma_r(u))_{nj+i}$ does not depend on the totality of u but only on u_j : it is the i th letter of $\sigma_r(u_j)$. \square




Figure 6: Illustration of Lemma 3.1.

Lemma 3.2. *For any $r \in \{0, \dots, n-1\}$,*

$$\sigma_r = \rho^r \circ \sigma.$$

Proof. Let $u \in \mathcal{A}^{\mathbb{Z}}$. Let $i, r \in \{0, \dots, n-1\}$ and $j \in \mathbb{Z}$.

$$\sigma(u)_{nj+i} = \begin{cases} 0 & \text{if } i \neq n-1 \\ \sigma_r(u)_{nj+i-r} & \text{if } i = n-1 \end{cases}$$

Considering that if $i \neq n-1$, $\sigma_r(u)_{nj+i-r} = 0$, we conclude that we always have $\sigma(u)_{nj+i} = \sigma_r(u)_{nj+i-r}$, and so $\sigma_r = \rho^r \circ \sigma$. \square

Lemma 3.3. *For $n \geq 3$, σ_1 has a unique fixpoint. For $n = 2$, σ_1 has no fixpoint but σ_1^2 has two fixpoints.*

Proof. Proposition 4 from [19] characterizes biinfinite fixpoints of substitutions. In the present case of σ_1 , [19] states that $w = \sigma_1(w)$ if and only if $w = y.x$ with $x = \overrightarrow{\sigma_1}(c)$ and $y = \overleftarrow{\sigma_1}(c')$ with $\sigma_1(c) = cv$ and $\sigma_1(c') = uc'$, $u, v \in \{0, 1\}^*$, $c, c' \in \{0, 1\}$. Notice that $\sigma_1(0) = 0^{n-2}10$ and $\sigma_1(1) = 0^n$, for $n \geq 3$, so the only choice for c and c' is $c = c' = 0$. Then σ_1 has a fixpoint that is $\overleftarrow{\sigma_1}(0).\overrightarrow{\sigma_1}(0)$ and which is unique.

For $n = 2$ the same reasoning concludes that σ_1 has no fixpoint. However, since $\sigma_1^2(0) = 0010$ and $\sigma_1^2(1) = 1010$, the same reasoning also yields that σ_1^2 has two fixpoints that are $\overleftarrow{\sigma_1^2}(0).\overrightarrow{\sigma_1^2}(0)$ and $\overleftarrow{\sigma_1^2}(0).\overrightarrow{\sigma_1^2}(1)$. \square

Lemma 3.4. *For every $k \in \mathbb{N}$ and every $i_1, \dots, i_k \in \{0, \dots, n-1\}$, the fixpoints of $s = \sigma_{i_k} \circ \dots \circ \sigma_{i_1}$ are aperiodic.*

Proof. To prove the aperiodicity of a fixpoint w of s (in the case where such a fixpoint exists), we follow a proof from [17], simplified for our specific case.

First, let us show that the two subwords 00 and 01 can be found in w .

- For 00, let us define $s' = \sigma_{i_{k-1}} \circ \dots \circ \sigma_{i_1}$. Then, by definition, $w = \sigma_{i_k}(s'(w))$ (by convention $s'(w) = w$ if $k = 1$). We are going to prove that $s'(w)$ always contains a 1. As a consequence, $w = \sigma_{i_k}(s'(w))$ contains 00 because $\sigma_{i_k}(1) = 0^n$. Suppose $s'(w) = \dots 000\dots$. If $k = 1$, it means that $w = \dots 000\dots$, but then $s(w) \neq w$ so this is impossible. If $k = 2$, then $s' = \sigma_{i_1}$ so the only way to have $s'(w) = \dots 000\dots$ is to have $w = \dots 111\dots$, but again $s(w) \neq w$. If $k \geq 3$, let us define $t = \sigma_{i_{k-3}} \circ \dots \circ \sigma_{i_1}$. With this notation, $w = \sigma_{i_k} \circ \sigma_{i_{k-1}} \circ \sigma_{i_{k-2}}(t(w))$. The assumption $s'(w) = \dots 000\dots$ causes $\sigma_{i_{k-2}}(t(w)) = \dots 111\dots$. However, this is impossible since $\dots 111\dots$ has no antecedent by $\sigma_{i_{k-2}}$. Therefore $s'(w)$ must contain a 1 and we can find 00 in w .
- For 01, the only way for w not to contain 01 is to be of the form $w = \dots 000\dots$, $w = \dots 111\dots$ or $w = \dots 1100\dots$. But it is clear that $s(\dots 000\dots) \neq \dots 000\dots$, $s(\dots 111\dots) \neq \dots 111\dots$ and $s(\dots 1100\dots) \neq \dots 1100\dots$ hence none of them can be fixpoints.

Hence $s(00)$ and $s(01)$ can also be found in w since $s(w) = w$. From this, we build by induction infinitely many words with two possible right extensions. We have $s(00) \neq s(01)$; consider the largest prefix on which they agree, call it u_2 , with $|u_2| > 1$. Then both u_20 and u_21 can be found in w . Hence $s(u_20)$ and $s(u_21)$ can also be found in w . We have $s(u_20) \neq s(u_21)$; consider the largest prefix on which they agree, call it u_3 , with $|u_3| > |u_2|$. Then both u_30 and u_31 can be found in w . Hence $s(u_30)$ and $s(u_31)$ can also be found in w .

By induction, we can build subwords of w as large as we want that have two choices for their last letter. Hence the factor complexity of w is unbounded, and so w is aperiodic (see Section 1.3). \square

3.2 Encoding substitutions in $BS(1, n)$

We now show how to encode such substitutions in SFTs of the group $BS(1, n)$ given by a tiling. We define the tiling τ_σ on $BS(1, n)$, $n \in \mathbb{N}$, $n \geq 2$, to be the set of tiles shown on Fig. 7 for all $c \in \{0, 1\}$ and $i \in \{0, \dots, n-1\}$. Remark that a tile is uniquely defined by the couple (c, i) .




Figure 7: Tiles of τ_σ : left and right colors are identical and equal to i , top color is c and bottom colors are equal to $\sigma_i(c)_0, \dots, \sigma_i(c)_{n-1}$.

This tiling will be the weakly but not strongly aperiodic tiling we are looking for. Lemmas 3.3 and 3.4 study the words that can appear on levels \mathcal{L}_g of the tiling, by looking at the fixpoints of σ_1 . They prove that no biinfinite word can be both a fixpoint for the σ_i 's and a periodic word, forbidding one direction of periodicity for any configuration we will encode with our tiling. This naturally leads to the following proposition:

Proposition 3.5. *No configuration of X^{τ_σ} can be a^k -periodic for any $k \in \mathbb{N}$.*

Proof. Suppose that there is a configuration x of X^{τ_σ} such that for any $g \in BS(1, n)$, $x_{a^k \cdot g} = x_g$ (a^k -periodicity). Call $w := (x_{aj})_{j \in \mathbb{Z}}$ the biinfinite word based on level \mathcal{L}_e . w

is k -periodic by a^k -periodicity of the configuration x . But w is also nk -periodic. Hence $(x_{ba^j})_{j \in \mathbb{Z}}$ is k -periodic. Indeed, by construction, when applying the correct substitution σ_i to x_{ba^j} and $x_{ba^{j+k}}$, one obtains the words $x_{a^{nj}} \dots x_{a^{nj+n-1}}$ and $x_{a^{nj+nk}} \dots x_{a^{nj+nk+n-1}}$ which are one and the same by nk -periodicity of w . Since there is only one preimage for a word by σ_i , $x_{ba^j} = x_{ba^{j+k}}$. By the same argument, one can show that for any integer $l > 0$, $(x_{b^l a^j})_{j \in \mathbb{Z}}$ must be k -periodic. However, these biinfinite sequences only use digits among $\{0, 1, 2\}$ so there is a finite number of such sequences. In particular, two of these sequences are the same. Since one is obtained from the other by applying the correct succession of σ_i 's, we get a periodic sequence that is a fixpoint of some $s = \sigma_{i_N} \circ \dots \circ \sigma_{i_1}$ for some $i_1, \dots, i_N \in \{0, \dots, n-1\}$. This contradicts Lemma 3.4. \square

Lemma 3.6. *There exists a weakly periodic configuration in $X^{\tau\sigma}$ for $n \geq 3$.*

Proof. We define w the unique fixpoint of σ_1 obtained thanks to Lemma 3.3.

Let $f(k) = \lfloor \frac{k}{n} \rfloor$ be the function that maps k to the quotient in the euclidean division of k by n and $r(k)$ its remainder. We also define $F(k) = f(k+1)$ and $R(k) = r(k+1)$. This means that $nF(k) + R(k) = k+1$, $F(ln) = l + \lfloor \frac{1}{n} \rfloor = l$, but also $F(k+n) = \lfloor \frac{k+1+n}{n} \rfloor = F(k) + 1$, and consequently $F^m(k+n^m) = F^m(k) + 1$.

$X^{\tau\sigma}$ is nonempty

We define a configuration x describing which tile (c_g, i_g) (a tile being uniquely defined by such a couple) is assigned to g , i.e. $x_g = (c_g, i_g)$, using the quasi-normal form $g = b^{-k} a^l b^m$. Then, we check that x does verify the adjacency rules. Define $x \in \tau_\sigma^{BS(1,n)}$ by

$$\begin{cases} x_{b^{-k} a^l} := (w_l, 1) \\ x_{b^{-k} a^l b^m} := (w_{F^m(l)}, R \circ F^{m-1}(l)) \text{ for } m > 0. \end{cases}$$

Remember that Lemma 2.18 states that any $g \in BS(1,n)$ can be written $b^{-k_1} a^{l_1} b^{m_1}$. Suppose it has a second form $b^{-k_2} a^{l_2} b^{m_2}$ with $k_2 > k_1$ up to exchanging the notations (were they equal, it is easy to prove the two forms would be the same). Then $b^{k_1-k_2} a^{l_2} b^{m_2-m_1} = a^{l_1}$, that is, $b^{k_1-k_2} a^{l_2-l_1} n^{k_2-k_1} b^{m_2-m_1} = e$. This means that $k_2 - k_1 = m_2 - m_1 > 0$ and $l_2 = l_1 n^{k_2-k_1}$. With that, we prove our x is well-defined. $k_2 > k_1$ causes $m_2 > 0$ in order to have $k_2 - k_1 = m_2 - m_1 > 0$. Consequently,

$$\begin{aligned} x_{b^{-k_2} a^{l_2} b^{m_2}} &= (w_{F^{m_2}(l_2)}, R \circ F^{m_2-1}(l_2)) \\ &= (w_{F^{m_1+k_2-k_1}(l_1 n^{k_2-k_1})}, R \circ F^{m_1+k_2-k_1-1}(l_1 n^{k_2-k_1})) \\ &= (w_{F^{m_1}(l_1)}, R \circ F^{m_1-1}(l_1)) \\ &= x_{b^{-k_1} a^{l_1} b^{m_1}} \end{aligned}$$

with a variation on the second to last line if $m_1 = 0$: we have $R \circ F^{k_2-k_1-1}(l_1 n^{k_2-k_1}) = R(l_1 n) = 1$.

Now, we prove that $x \in X^{\tau\sigma}$. Let $g = b^{-k} a^l b^m$.

- If $m > 0$, we have

$$\begin{aligned} x_{ga}(\text{left}) &= x_{b^{-k} a^{l+n^m} b^m}(\text{left}) \\ &= R \circ F^{m-1}(l + n^m) \\ &= R(F^{m-1}(l) + n) \\ &= R \circ F^{m-1}(l) \\ &= x_g(\text{right}). \end{aligned}$$

- If $m = 0$, we have

$$\begin{aligned} x_{ga}(\text{left}) &= x_{b^{-k}a^{l+1}}(\text{left}) \\ &= 1 \\ &= x_g(\text{right}). \end{aligned}$$

Let $j \in \{0, \dots, n-1\}$. We have

$$\begin{aligned} x_{ga^{-j}b}(\text{bottom}_{j+1}) &= x_{b^{-k}a^{l-jn^m}b^{m+1}}(\text{bottom}_{j+1}) && (g = b^{-k}a^l b^m) \\ &= \sigma_{R \circ F^m(l-jn^m)}(w_{F^{m+1}(l-jn^m)})_j && (\text{by definition of } x) \\ &= \sigma_{R \circ F^m(l-jn^m)}(w)_{nF^{m+1}(l-jn^m)+j} && (\text{Lemma 3.1}) \\ &= \sigma(w)_{nF^{m+1}(l-jn^m)+j+R \circ F^m(l-jn^m)} && (\text{Lemma 3.2}) \\ &= \sigma(w)_{F^m(l-jn^m)+j+1} && (\text{by definition of } F \text{ and } R) \\ &= \sigma_1(w)_{F^m(l-jn^m)+j} && (\text{Lemma 3.2}) \\ &= w_{F^m(l-jn^m)+j} && (\text{since } w \text{ is a fixpoint of } \sigma_1) \\ &= w_{F^m(l)} && (F^m(l-jn^m) = F^m(l) - j) \\ &= x_g(\text{top}) \end{aligned}$$

Consequently, x describes a valid configuration of X^{τ_σ} : all adjacency conditions are verified.

x is b -periodic

With the definition of x , it is easy to check that for any $g \in BS(1, n)$, $x_{bg} = x_g$. Hence it is a weakly periodic configuration. \square

We can now obtain our second main theorem:

Theorem 3.7. *The tileset τ_σ forms a weakly aperiodic but not strongly aperiodic SFT on $BS(1, n)$, $n \in \mathbb{N}$, $n \geq 2$.*

Proof. First, in the $n \geq 3$ case, there is a weakly periodic configuration in X^{τ_σ} , see Lemma 3.6. Hence it is not a strongly aperiodic SFT.

In the $n = 2$ case, we define u and v the two fixpoints of σ_1^2 (Lemma 3.3 again) and remark that $v = \sigma_1(u)$ and $u = \sigma_1(v)$. We define a configuration $x \in \tau_\sigma^{BS(1, n)}$ by:

$$\begin{aligned} x_{b^{-k}a^l} &:= \begin{cases} (u_l, 1) & \text{if } k+m \equiv 0 \pmod{2} \\ (v_l, 1) & \text{if } k+m \equiv 1 \pmod{2} \end{cases} \\ x_{b^{-k}a^l b^m} &:= \begin{cases} (u_{F^m(l)}, R \circ F^{m-1}(l)) & \text{for } m > 0 \text{ if } k+m \equiv 0 \pmod{2} \\ (v_{F^m(l)}, R \circ F^{m-1}(l)) & \text{for } m > 0 \text{ if } k+m \equiv 1 \pmod{2} \end{cases} \end{aligned}$$

and we use the same notations as in the proof of Lemma 3.6. The reasoning is also the same, except instead of using w an alternation appears between u and v in all the equations. As a consequence, the configuration is b^2 -periodic instead of b . Once again, X^{τ_σ} is consequently not strongly aperiodic.

Now, using Proposition 3.5, and since all powers of a are of infinite order in $BS(1, n)$, we get that for any valid configuration x of τ_σ , $|Orb_{BS(1, n)}(x)| = +\infty$, for any $n \geq 2$. Hence no configuration of τ_σ is strongly periodic, and so the SFT is weakly aperiodic. \square

4 A strongly aperiodic SFT on $BS(n, n)$

This section is a mere assembly of known results, that we think are worth gathering in the context of the current paper. It uses a theorem from [12] seen as an extension of the construction presented in [14]. The idea behind that theorem is that $G \times \mathbb{Z}$ admits a strongly aperiodic SFT as soon as G can encode piecewise affine functions. This is reflected by the PA' condition described in [12] and restated below.

Definition 4.1. Let $k \in \mathbb{N}$. Let $\mathcal{F} = \{f_i: P_i \rightarrow P'_i \mid i \in \{0, \dots, k\}\}$ be a finite set of piecewise affine rational homeomorphisms, where each P_i and P'_i is a finite union of bounded rational polytopes of \mathbb{R}^n . Let $D = \bigcap_{i=1}^k P_i \cap \bigcap_{i=1}^k P'_i$ be the common domain of all functions of \mathcal{F} and their inverses.

Let $S_{\mathcal{F}}$ be the closure of the set $\{f_i, f_i^{-1} \mid i \in \{1, \dots, k\}\}$ under composition. We define $G_{\mathcal{F}}$, the group $\{f|_D \mid f \in S_{\mathcal{F}}\}$.

A finitely generated group G is PA' -recognizable if there exists a finite set \mathcal{F} of piecewise affine rational homeomorphisms such that:

- (A) $G \cong G_{\mathcal{F}}$;
- (B) $\forall t \in D, \forall g \in \mathcal{F}, [\forall f \in \mathcal{F}, g(f(t)) = f(t)] \Rightarrow g = Id$.

Theorem 4.1 ([12], Th. 7). *If G is an infinite finitely generated PA' -recognizable group, then $\mathbb{Z} \times G$ admits a strongly aperiodic SFT.*

We need two additional propositions to obtain the desired result on $BS(n, n)$:

Proposition 4.2 ([6], Prop. 9 & 10). *If G is a finitely generated group and H is a finitely generated subgroup of G of finite index, then we have the following:*

$$\begin{aligned} H \text{ admits a weakly aperiodic SFT} &\Leftrightarrow G \text{ admits a weakly aperiodic SFT} \\ H \text{ admits a strongly aperiodic SFT} &\Leftrightarrow G \text{ admits a strongly aperiodic SFT.} \end{aligned}$$

The following proposition is known, but we include a self-contained proof.

Proposition 4.3. *$BS(n, n)$ admits $\mathbb{Z} \times \mathbb{F}_n$ as a subgroup of finite index, where \mathbb{F}_n is the free group of order n .*

Proof. Let H be the subgroup of $BS(n, n)$ generated by $\{a^n\} \cup \{a^i b a^{-i} \mid i \in \{0, \dots, n-1\}\}$. First, H is normal in $BS(n, n)$. We prove that $aHa^{-1} \subseteq H$ by verifying it on its generators: the only verification needed is $aa^{n-1}ba^{-(n-1)}a^{-1} = a^n b a^{-n} = b a^n a^{-n} = b$. Similarly, $a^{-1}Ha \subseteq H$; and finally, $bHb^{-1} \subseteq H$ (same for b^{-1}) since $b \in H$. Second, H is isomorphic to $\mathbb{Z} \times \mathbb{F}_n$ through the following isomorphism (denoting g_0, \dots, g_{n-1} the generators of \mathbb{F}_n and ε its identity):

$$\begin{aligned} \phi: \mathbb{Z} \times \mathbb{F}_n &\longrightarrow H \\ (1, \varepsilon) &\longmapsto a^n \\ (0, g_i) &\longmapsto a^i b a^{-i} \end{aligned}$$

It is a morphism by construction, which is correctly defined since the only basic relation of $\mathbb{Z} \times \mathbb{F}_n$, that is $(1, \varepsilon) \cdot (0, g_i) = (0, g_i) \cdot (1, \varepsilon)$, is preserved in H : $a^n a^i b a^{-i} = a^i a^n b a^{-i} = a^i b a^n a^{-i} = a^i b a^{-i} a^n$. Said morphism is surjective, because H is generated by a^n and $\{a^i b a^{-i} \mid i \in \{0, \dots, n-1\}\}$. Finally, it is also injective: let $g = (k, w) \in \mathbb{Z} \times \mathbb{F}_n$, with $w = (g_{i_1})^{e_1} \dots (g_{i_N})^{e_N}$ where the e_l are in $\{-1, +1\}$.

$$\phi(g) = e \Leftrightarrow a^{nk} a^{i_1} b^{e_1} a^{-i_1} a^{i_2} \dots a^{-i_{N-1}} a^{i_N} b^{e_N} a^{-i_N} = e$$

This form is a canonical form in H : any word in H can be uniquely written as such. Indeed, any word in H is a succession of generators of it, $a^{i_k} b^{e_k} a^{-i_k}$ and a^n . But a^n commutes

with all the other generators due to the relation of $BS(n, n)$, so such a form is always attainable. To prove it is unique, it is enough to prove it for e : suppose we have some $a^{nk}a^{i_1}b^{e_1}a^{-i_1}a^{i_2} \dots a^{-i_{N-1}}a^{i_N}b^{e_N}a^{-i_N} = e$. First, realize that no relation in $BS(n, n)$ allows to reduce the total power of a in a word, causing $k = 0$ necessarily. Then, consider the resulting word $a^{i_1}b^{e_1}a^{i_2-i_1} \dots a^{i_N-i_{N-1}}b^{e_N}a^{-i_N}$ in $BS(n, n)$: it cannot be reduced in $BS(n, n)$ since all powers of a between two b 's are of absolute value smaller than n .

As a consequence, the previous equality is true only when $k = 0$ and $w = \varepsilon$. Hence the injectivity of the map.

Moreover, any element of $BS(n, n)$ can be written in a form that much resembles the one mentioned above:

$$a^p a^{nk} a^{i_1} b^{e_1} a^{-i_1} \dots a^{i_N} b^{e_N} a^{-i_N}$$

with $p \in \{0, \dots, n-1\}$. To do so, first move all a^n 's in the rightmost power of a in the word, to the leftmost part of the word. Ensure that $-i_N$, the remaining power, is in $\{-n+1, \dots, -1, 0\}$. Then force a^{i_N} to appear on the left of the b itself to the left of a^{-i_N} , and call $-i_{N-1}$ the remaining power of a (it is in $\{-(n-1), \dots, -1, 0\}$ up to moving another a^n to the leftmost part of the word) before another b to the left. Repeat this operation until there is no b to the left of the power of a you consider, and split this final a^K into $a^p a^{nk} a^{i_1}$.

As a consequence, $BS(n, n)/H = \{\bar{e}, \bar{a}, \dots, \overline{a^{n-1}}\} \cong \mathbb{Z}/n\mathbb{Z}$. Hence H is of finite index in $BS(n, n)$. \square

Theorem 4.4. *For every $n \geq 2$, $BS(n, n)$ admits a strongly aperiodic SFT.*

Proof. First, finitely generated subgroups of compact groups of matrices on integers are PA' -recognizable (see [12], Proposition 5.12). \mathbb{F}_2 , the free group of order 2, is isomorphic to a subgroup of $SL_2(\mathbb{Z})$ (see [8, Lemma 2.3.2]), hence it is PA' -recognizable. It is also known (see [8, Corollary D.5.3]) that \mathbb{F}_n is a subgroup of \mathbb{F}_2 ; so it is isomorphic to a subgroup of $SL_2(\mathbb{Z})$ and PA' -recognizable too. Therefore by Theorem 4.1 $\mathbb{Z} \times \mathbb{F}_n$ admits a strongly aperiodic SFT. Using Proposition 4.2 and Proposition 4.3, we obtain that $BS(n, n)$ admits a strongly aperiodic SFT. \square

Conclusion

Baumslag-Solitar groups $BS(m, n)$ are residually finite if and only if $|m| = 1$, $|m| = |n|$ or $|m| = |n|$ (Proposition 1.1). Gathering results from Theorem 2.20, Theorem 3.7 and Theorem 4.4, and considering that $BS(-m, -n) \cong BS(-m, n) \cong BS(m, -n) \cong BS(m, n)$, we obtain the following:

Theorem 4.5. *Residually finite Baumslag-Solitar groups $BS(m, n)$ with $|m| \geq 2$ or $|n| \geq 2$ admit both strongly and weakly-not-strongly aperiodic SFTs.*

For non-residually finite Baumslag-Solitar groups, the existence of strongly aperiodic STF is still an open question.

In Section 3, we showed how to encode a particular set of substitutions into a tiling of $BS(1, n)$. An interesting question related to combinatorics on words would be to characterize the sets of substitutions that can be encoded using our technique. It is clear that the nature of $BS(1, n)$ impose some conditions on these substitutions, and it would be of independent interest to obtain a self-contained condition on the substitutions and study their properties.

Acknowledgments

The authors would like to thank Nathalie Aubrun for the interest she sparked about tilesets on the Baumslag-Solitar groups and the help she provided to understand her joint work with

Jarkko Kari. They also thank Silvère Gangloff for pointing them to [11] that provided the missing piece to prove Lemma 2.15; Jarkko Kari for his questions, that led to the writing of Section 3; and Pierre Guillon for his many remarks and useful suggestions that made the paper much more readable, even if it delayed a bit its publication.

The first author would like to thank Michael Schraudner and his PhD students Álvaro Bustos and Hugo Maturana, under whose supervision Section 4 was conceived as part of an internship, for their numerous insights in the proof of said theorem.

This publication was made possible through the support of the ID#61466 grant from the John Templeton Foundation, as part of the “The Quantum Information Structure of Spacetime (QISS)” Project (qiss.fr). The opinions expressed in this publication are those of the author(s) and do not necessarily reflect the views of the John Templeton Foundation

References

- [1] N. Aubrun, S. Barbieri, and E. Jeandel. About the domino problem for subshifts on groups. In *Trends in Mathematics*, pages 331–389. Springer International Publishing, 2018.
- [2] N. Aubrun, S. Barbieri, and E. Moutot. The Domino Problem is Undecidable on Surface Groups. In *44th International Symposium on Mathematical Foundations of Computer Science (MFCS 2019)*, volume 138 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 46:1–46:14, Dagstuhl, Germany, 2019.
- [3] N. Aubrun and J. Kari. Tiling Problems on Baumslag-Solitar groups. In *MCU’13*, pages 35–46, 2013.
- [4] N. Aubrun and J. Kari. Addendum to “Tilings problems on Baumslag-Solitar groups”. Jan 2021. arXiv: 2101.12470.
- [5] R. Berger. *The Undecidability of the Domino Problem*. PhD thesis, Harvard University, 1964.
- [6] D. Carroll and A. Penland. Periodic points on shifts of finite type and commensurability invariants of groups. *New York Journal of Mathematics*, 21, 2015.
- [7] J. Cassaigne and F. Nicolas. *Factor complexity*, pages 163–247. Encyclopedia of Mathematics and its Applications. Cambridge University Press, 2010.
- [8] T. Ceccherini-Silberstein and M. Coornaert. *Cellular Automata and Groups*. Springer Monographs in Mathematics. Springer Berlin Heidelberg, 2010.
- [9] D. Cohen and C. Goodman-Strauss. Strongly aperiodic subshifts on surface groups. *Groups, Geometry, and Dynamics*, 11(3):pp. 1041–1059, 2017.
- [10] E. Coven and Z. Nitecki. On the genesis of symbolic dynamics as we know it. *Colloquium Mathematicum*, 110, 12 2006.
- [11] B. Durand, G. Gamard, and A. Grandjean. Aperiodic tilings and entropy. In *Developments in Language Theory*, pages 166–177. Springer International Publishing, 2014.
- [12] E. Jeandel. Aperiodic Subshifts of Finite Type on Groups. *ArXiv e-prints*, Jan. 2015.
- [13] E. Jeandel and M. Rao. An aperiodic set of 11 Wang tiles. *Advances in Combinatorics*, Jan 2021.

- [14] J. Kari. On the undecidability of the tiling problem. In *Current Trends in Theory and Practice of Computer Science (SOFSEM)*, pages 74–82, 2008.
- [15] D. A. Lind and B. Marcus. *An Introduction to Symbolic Dynamics and Coding*. Cambridge University Press, New York, NY, USA, 1995.
- [16] S. Meskin. Nonresidually Finite One-Relator Groups. *Transactions of the American Mathematical Society*, 164:pp. 105–114, 1972.
- [17] J.-J. Pansiot. Decidability of periodicity for infinite words. *RAIRO - Theoretical Informatics and Applications*, 20(1):pp. 43–46, 1986.
- [18] R. M. Robinson. Undecidability and nonperiodicity for tilings of the plane. *Inventiones Mathematicae*, 12:pp. 177–209, 1971.
- [19] J. Shallit and M.-w. Wang. On two-sided infinite fixed points of morphisms. In G. Ciobanu and G. Păun, editors, *Fundamentals of Computation Theory*, pages 488–499, Berlin, Heidelberg, 1999. Springer Berlin Heidelberg.
- [20] H. Wang. Proving theorems by pattern recognition II. *Bell System Technical Journal*, 40(1-3):pp. 1–41, 1961.