



HAL
open science

Baseline Surface Radiation Network (BSRN): structure and data description (1992-2017)

Amelie Driemel, John Augustine, Klaus Behrens, Sergio Colle, Christopher Cox, Emilio Cuevas-Agulló, Fred M. Denn, Thierry Duprat, Masato Fukuda, Hannes Grobe, et al.

► To cite this version:

Amelie Driemel, John Augustine, Klaus Behrens, Sergio Colle, Christopher Cox, et al.. Baseline Surface Radiation Network (BSRN): structure and data description (1992-2017). *Earth System Science Data*, 2018, 10, pp.1491-1501. 10.5194/essd-10-1491-2018 . hal-03658664

HAL Id: hal-03658664

<https://hal.science/hal-03658664v1>

Submitted on 4 May 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Baseline Surface Radiation Network (BSRN): structure and data description (1992–2017)

Amelie Driemel¹, John Augustine², Klaus Behrens^{3,27}, Sergio Colle⁴, Christopher Cox⁵, Emilio Cuevas-Agulló⁶, Fred M. Denn⁷, Thierry Duprat⁸, Masato Fukuda⁹, Hannes Grobe¹, Martial Haeffelin¹⁰, Gary Hodges¹¹, Nicole Hyett¹², Osamu Ijima⁹, Ain Kallis¹³, Wouter Knap¹⁴, Vasilii Kustov¹⁵, Charles N. Long², David Longenecker^{2,*}, Angelo Lupi¹⁶, Marion Maturilli¹⁷, Mohamed Mimouni^{18,*}, Lucky Ntsangwane¹⁹, Hiroyuki Ogihara⁹, Xabier Olano²⁰, Marc Olefs²¹, Masao Omori⁹, Lance Passamani¹², Enio Bueno Pereira²², Holger Schmithüsen¹, Stefanie Schumacher¹, Rainer Sieger^{1,†}, Jonathan Tamlyn^{23,*}, Roland Vogt²⁴, Laurent Vuilleumier²⁵, Xiangao Xia²⁶, Atsumu Ohmura^{27,*}, and Gert König-Langlo^{1,*}

¹Alfred-Wegener-Institut Helmholtz-Zentrum für Polar- und Meeresforschung, Bremerhaven, Germany

²NOAA ESRL Global Monitoring Division, Boulder, CO, USA

³Deutscher Wetterdienst, Meteorologisches Observatorium Lindenberg – Richard-Aßmann-Observatorium, Tauche, Germany

⁴Universidade Federal de Santa Catarina, Florianópolis, Brasil

⁵CIRES/NOAA ESRL Physical Sciences Division, Boulder, CO, USA

⁶Izaña Atmospheric Research Center (AEMET), Tenerife, Spain

⁷Science Systems and Applications, Inc., Lanham, USA

⁸Meteo France, Carpentras, France

⁹Japan Meteorological Agency, Tokyo, Japan

¹⁰Laboratoire de Météorologie Dynamique, Paris, France

¹¹CIRES, University of Colorado Boulder, CO, USA

¹²Bureau of Meteorology, Melbourne, Australia

¹³Estonian Environment Agency, Tõravere, Estonia

¹⁴Royal Netherlands Meteorological Institute, De Bilt, the Netherlands

¹⁵Arctic and Antarctic Research Institute, St. Petersburg, Russia

¹⁶Institute of Atmospheric Sciences and Climate, National Research Council of Italy, Bologna, Italy

¹⁷Alfred-Wegener-Institut Helmholtz-Zentrum für Polar- und Meeresforschung, Potsdam, Germany

¹⁸Office National de la Météorologie, Algiers, Algeria

¹⁹South African Weather Service, Pretoria, South Africa

²⁰National Renewable Energy Centre, Sarriguren, Spain

²¹ZAMG – Zentralanstalt für Meteorologie und Geodynamik, Vienna, Austria

²²Instituto Nacional de Pesquisas Espaciais, São José dos Campos, Brasil

²³Met Office, Exeter, Devon, UK

²⁴Meteorology Climatology and Remote Sensing, Department Environmental Sciences, University of Basel, Basel, Switzerland

²⁵Federal Office of Meteorology and Climatology, MeteoSwiss, Payerne, Switzerland

²⁶LAGEO, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, China

²⁷Institute for Atmospheric and Climate Science, ETH Zurich, Zurich, Switzerland

* retired

† deceased

Correspondence: Amelie Driemel (amelie.driemel@awi.de)

Received: 25 January 2018 – Discussion started: 16 February 2018

Revised: 28 July 2018 – Accepted: 30 July 2018 – Published: 21 August 2018

Abstract. Small changes in the radiation budget at the earth’s surface can lead to large climatological responses when persistent over time. With the increasing debate on anthropogenic influences on climatic processes during the 1980s the need for accurate radiometric measurements with higher temporal resolution was identified, and it was determined that the existing measurement networks did not have the resolution or accuracy required to meet this need. In 1988 the WMO therefore proposed the establishment of a new international Baseline Surface Radiation Network (BSRN), which should collect and centrally archive high-quality ground-based radiation measurements in 1 min resolution. BSRN began its work in 1992 with 9 stations; currently (status 2018-01-01), the network comprises 59 stations (delivering data to the archive) and 9 candidates (stations recently accepted into the network with data forthcoming to the archive) distributed over all continents and oceanic environments. The BSRN database is the World Radiation Monitoring Center (WRMC). It is hosted at the Alfred Wegener Institute (AWI) in Bremerhaven, Germany, and now offers more than 10 300 months of data from the years 1992 to 2017. All data are available at <https://doi.org/10.1594/PANGAEA.880000> free of charge.

1 Introduction

The climate of our earth is decisively influenced by radiative processes. Even small changes in the radiation budget at the earth’s surface can lead to large climatological responses when persistent over time (e.g., Chylek et al., 2007; Kwok and Untersteiner, 2011). International data centers have long been archiving data that are relevant for studies aiming to detect and monitor long-term variability in the surface energy balance. For example, the Global Energy Balance Archive (GEBA) (<http://www.geba.ethz.ch/>¹) has archived monthly means of various energy flux components from around 2500 locations since 1985 (Wild et al., 2013, 2017). Similarly, the World Radiation Data Center (<http://wrdc.mgo.rssi.ru/>²), under the auspices of the World Meteorological Organization (WMO), has provided daily and monthly sums and monthly means of solar radiation fluxes since 1964 from more than 1000 stations. A small subset of these stations that are connected to the Global Atmosphere Watch (GAW) also provide hourly sums.

However, with the increasing debate on anthropogenic influences on climatic processes since the 1980s – e.g., anthropogenic aerosol emission (Charlson et al., 1992) – the need for accurate radiometric measurements with higher temporal resolution was identified, and it was determined that the existing measurement networks could not provide the necessary temporal resolution or accuracy. As a solution, in 1988 the WMO proposed the establishment of a new international Baseline Surface Radiation Network (BSRN) under the World Climate Research Programme (WCRP). The BSRN would provide high-temporal-resolution ground-based radiation measurements for the validation of satellite data, as well as the validation and improvement of radiative transfer calculations in climate models, and support the detection and monitoring of long-term changes in ground-level radiation fluxes. The requirements of BSRN stations that set the network apart from its predecessors include the follow-

ing (Ohmura et al., 1998; McArthur, 2005): a commitment for long-term involvement of the station under the lead of a radiation expert responsible for the quality of the data; measurements of direct, diffuse and global downwelling short-wave fluxes and the downwelling long-wave flux in 1 min resolution; and traceable calibrations of the radiation instruments to the World Radiometric Reference (WRR), which is maintained at the World Radiation Center (WRC) in Davos, Switzerland. Furthermore, stations should be representative of a relatively large surrounding area for use in satellite and climate model validation and should be manned at least during work days to allow for regular cleaning, maintenance and check of instruments. To guarantee high-quality measurements, ventilation of all radiation instruments, especially at sites with large variability in ambient temperature, should be installed.

A central data archive for BSRN was developed under the direction of Atsumu Ohmura at the Swiss Federal Institute of Technology (ETH) in Zurich (Gilgen et al., 1995; Hegner et al., 1998; Ohmura et al., 1998), called the World Radiation Monitoring Center (WRMC). Since 2008, the WRMC has been run by the Alfred Wegener Institute (AWI), Helmholtz Center for Polar and Marine Research in Bremerhaven, Germany (König-Langlo et al., 2013), and currently stores more than 10 300 months of radiometric data at 1 min temporal resolution. In this paper, we describe the structure of the BSRN and the WRMC, the data coverage and data quality procedures, and we provide information for potential users interested in accessing BSRN data from 1992 to 2017.

2 BSRN structure and measured values

BSRN began its work in 1992 with 9 stations distributed globally, including the Antarctic, Arctic, Atlantic and Pacific oceans, North America, Africa and Europe. Currently (status 2018-01-01), the network comprises 59 stations (delivering data to the archive) and 9 candidates (stations recently accepted into the network with data forthcoming to the archive), collectively representing all seven continents

¹last access: 11 June 2018

²last access: 11 June 2018

Table 1. Overview of the logical records (LR) contained in station-to-archive files, as well as the number of stations submitting these data to the WRMC (status December 2017). The last column states the standard title for BSRN datasets archived within PANGAEA*.

Logical record	Data type included in the LR	No. of stations	Dataset title in PANGAEA
LR0100	Global, diffuse, direct, long-wave down	59	Basic measurements of radiation
LR0300	Reflected, long-wave up	15	Basic and other measurements*
LR0500	Ultraviolet radiation	12	Ultra-violet measurements
LR1000	Synops	13	Meteorological synoptical observations
LR1100	Upper air soundings	30	Radiosonde measurements
LR1200	Total ozone	9	Ozone measurements
LR1300	Ceilometer data	3	Expanded measurements
LR3010	Radiation measurements from tower	13	Other measurements at 10 m

* In PANGAEA the LR0300 and LR0100 are archived as one dataset.

data (vertical profiles of temperature, humidity, wind and ozone) and data on the ozone column thickness. An up-to-date overview of all available measurements can be accessed here: <https://dataportal.pangaea.de/bsrn/>³; an explanation of the content of each logical record (LR) can be found in Table 1. These additional measurements are not required for BSRN stations (though they are recommended) and in this paper we focus on the core measurements of BSRN, the downward- and upward-directed long- and shortwave radiation fluxes from 1992 to 2017.

3 Data format and quality checks

3.1 Data format

The original submission format is the so-called station-to-archive file. The relatively complex, strictly defined format of the station-to-archive file is intended to encapsulate both the data and all relevant metadata within a single ASCII file. Metadata records include the contact information, calibration history, geographical characteristics of the station and the types of instruments. The actual data within station-to-archive files are divided into so-called logical records, which do not contain table headers (no parameter names and units); see König-Langlo et al. (2013) for a detailed explanation of the content of each logical record and Table 2 in the present paper for an overview of available data. All station-to-archive files are stored on, and are accessible via, a public ftp server. To increase usability and visibility of the files, the WRMC at the AWI also publishes the data within the Data Publisher for Earth & Environmental Science, PANGAEA (<http://bsrn.awi.de/data/data-retrieval-via-pangaea/>⁴) in a much more straightforward and intuitive format. BSRN datasets in PANGAEA contain a metadata header followed by the data table with parameter names and units; see, for example, <https://doi.org/10.1594/PANGAEA.874567> (König-Langlo, 2017) and Fig. 2 for a visualization of this data file.

³last access: 28 July 2018

⁴last access: 28 July 2018

3.2 Quality control

The methods for the initial stage of quality control are left up to the individual station scientists (some scientists are responsible for more than one station) and thus are not implemented uniformly across BSRN stations. However, BSRN does maintain recommendations (Long and Dutton, 2002; Long and Shi, 2008) as well as software tools (Schmithüsen et al., 2012). After this initial quality control, the data generated at BSRN stations are submitted in monthly granularity in the station-to-archive file format by the station scientist to the WRMC data curator. From 1992 to 2007, when the archive was maintained by the ETH in Zurich, the data were checked for physically possible limits within the WRMC and suspicious values were flagged, but not removed. Since being operated at the AWI, additional procedures have been enacted for data submitted to the WRMC. The data are now not only quality checked (see below), but suspicious data are reported back to the station scientist with an appeal to remove them and resubmit the file. The files that were archived using the old system (pre-2007) were adopted by AWI without further checks and without quality flags. However, many of the old files have since been exchanged with updated versions. Old files can be distinguished from new ones quite easily: if the data are from pre-2007 and the version number is 1 in the station to archive file (line 1, last number) or if it has 2007 as publication year in PANGAEA, it is an old file. Details on the quality checks performed at the AWI are given below.

Station scientists and users have the option of processing BSRN data using software developed for the BSRN community and WRMC called the “BSRN Toolbox” (Schmithüsen et al., 2012). The software provides a format check for the station-to-archive files (and for PANGAEA download files, see below) and can perform quality checks of the data as outlined in the BSRN Global Network recommended QC tests, V2.0 (Long and Dutton, 2002). The types of tests include those that identify data outside physically possible limits, as well as extremely rare limits. Additionally, a set of comparison metrics are available that flag data as suspect based on empirically established relationships relative




Figure 2. Example of a PanPlot2 (Sieger and Grobe, 2013) visualization of a monthly data file (January 2017) from station GVN (Neumayer, Antarctica); T2 is air temperature at 2 m height, RH is relative humidity and PoPoPoPo is air pressure.

to other variables (comparisons check, e.g., LWU to air temperature). A broader and more refined range of such tests is outlined by Long and Shi (2008) and their implementation is recommended for BSRN. The astronomical data necessary for many tests can be calculated using different algorithms (Iqbal, 1983; Solpos, Solar Position and Intensity, with refraction and Solpos without refraction; Michalsky; see, for example, Iqbal, 1983; Michalsky, 1988), most of them selectable within the BSRN Toolbox. At the AWI the data curator generally uses the Iqbal 1983 algorithm and the comparisons check. In all cases, an output file can be created, which includes the quality codes (checkbox “Quality codes”) and another file which plots the measured global radiation (SWD) vs. the calculated total horizontal radiation (SumSW), computed from the direct and diffuse radiation measurements ($\text{SumSW} = \text{DIF} + \text{DIR} \times \cos(z)$; see Fig. 3 for details). A comparison of SWD to SumSW is a good way of detecting instrument malfunctions⁵.

Since the BSRN Toolbox is subject to the GNU General Public License, it is freely available to everyone. With the help of the toolbox, BSRN station scientists as well as data users can easily download and convert station-to-archive files

⁵Although if the sun tracker for DIR and DIF is completely off then $\text{DIR} = 0$ and $\text{SumSW} = \text{SWD}$ (very good agreement), so there are situations where a perfect match does not indicate high-quality data. But this test is non-definitive in that failure does not indicate which of the three instruments might be producing the disagreement.

to user-friendly formats by clicking the option “Data”, which splits station-to-archive files up into logical records containing parameter names and units. In addition, quality checks can be applied on BSRN files (downloaded from both ftp and PANGAEA) using the option “Quality check”. All data can be visualized with the free software PanPlot2 (Sieger and Grobe, 2013).

Some quality issues cannot be easily solved. One prominent example is the case of zero-offsets: Thermophile radiometers are known to suffer a negative bias due to infrared loss to the sky (Dutton et al., 2001). Consequently, though physically impossible, negative radiation fluxes and values are often present in nighttime measurements. To reduce these so-called IR loss offsets, BSRN sensors are ventilated (Michalsky et al., 2017) and/or heated artificially. However, ventilation sometimes fails or is insufficient to compensate for large temperature variations. Depending on the location and instrument used, nighttime values more negative than -4 W m^{-2} can just be removed, or in other cases corrections can be made using algorithms that make use of collocated pyrogeometer data that are part of the BSRN instrument suite (Dutton et al., 2001; Long et al., 2001; Younkin and Long, 2004). However, if the negative bias also occurs – but is not evident – during the day, the BSRN recommends keeping the nighttime negatives in the archived files. Each individual BSRN customer can then decide how to treat the data depending on the scope and aim of their work. Ventilation is also used to help prevent the build-up of ice (e.g., rime




Figure 3. (a) Very good agreement of SumSW with SWD indicating high-quality measurements of SWD, DIR and DIF (please note that due to the thermal offset, combined with the cosine error of SWD the plot usually exhibits more scatter than shown here). (b) Three days of instrument failure shown by datapoints far below the 1 : 1 line. The toolbox flags values where $\text{SWD} / \text{SumSW}$ is not within $\pm 8\%$ of 1.0 for solar zenith angle (SZA) $< 75^\circ$, respectively, where $\text{SWD} / \text{SumSW}$ is not within $\pm 15\%$ of 1.0 for $93^\circ > \text{SZA} > 75^\circ$. For $\text{SumSW} < 50 \text{ W m}^{-2}$; however, the test is not effective.

or frost) on the sensor windows, though experience within the BSRN community indicates that ice remains an important vulnerability to measurement quality and data retention rates for stations in cold climates (BSRN, 2012). BSRN is working to formulate recommendations for the network.

A more thorough description of BSRN methods and quality control issues can be found in Lanconelli et al. (2011) and Matsui et al. (2012), where issues especially related to polar BSRN sites are elaborated on. Other interesting papers on the quality and possible biases of radiation measurements were published by Vuilleumier et al. (2014), who analyze the uncertainty of broadband shortwave radiation monitoring; Olefs et al. (2016), who expand on measurement and calibration standards within the Austrian radiation monitoring network; and Nyeki et al. (2017), who discuss pyrgeometer sensitivity on atmospheric integrated water vapor content.

4 Data availability

In <https://doi.org/10.1594/PANGAEA.880000> the downward and upward short- and long-wave data of all BSRN stations for the years 1992 to 2017 are archived. More than 10 300 monthly files (> 850 station years) of high-quality radiation data are thus available free of charge. The files are organized in zip files with one file containing all monthly data files for 1 year of downward and (if available) upward radiation data (LR0100 + 0300 or LR0100 and LR3010; see Table 1) from one station. The format of the files is tab delimited txt, and the structure is the same as if downloaded directly from PANGAEA (see Fig. 4). At the top of the files the metaheader gives details on the citation of the dataset; a link to the respective station-to-archive file on the ftp server;

and the location of the station, parameters, units, PI, methods etc., followed by the actual data table with abbreviated parameter names and units.

To be able to download the files, users should contact the WRMC staff (<http://bsrn.awi.de/contact-persons/>⁶) to request a BSRN account. By accepting this account, the data user agrees to comply with the guidelines defined at <http://bsrn.awi.de/data/conditions-of-data-release/>⁷. The reason for requiring that users register with BSRN is that the organization is a nonprofit voluntary network and it is important that station managers who commit resources to providing the data are able to receive appropriate acknowledgement, the correct citation of their data files in publications and are able to monitor the level of use of the station data. Therefore, the acknowledgement of the data release guidelines is an essential step in the data download process. There are plans to simplify the procedure by allowing the acceptance of the data release guidelines directly during the download of the data, but this feature is not operative yet. From 1 January to 31 December 2017, 140 requests from 35 countries were received and answered promptly (within 1 day). On average 800 visitors per month visit the BSRN website. Up to now, Web of Science reports a total of 100 publications related to BSRN, which have been cited more than 2500 times (accessed on 10 January 2018). Several hundred to several thousand BSRN files are downloaded each month. Unfortunately, to the best knowledge of the authors there is no data citation report yet available to check for BSRN dataset citations.

⁶last access: 28 July 2018

⁷last access: 28 July 2018


```

/* METADATA:
Citation: König-Langlo, Gert (2017): Basic and other measurements of radiation at Neumayer Station (2017-01). Alfred Wegener Institut Bremerhaven, PANGAEA, https://doi.org/10.1594/PANGAEA.874567
Other version: König-Langlo, Gert (2017): BSRN Station-to-archive file for Neumayer Station (2017-01). ftp://ftp.bsrn.awi.de/gvn/gvn0117.d
Project(s): Baseline Surface Radiation Network (BSRN) (URI: http://bsrn.awi.de)
Coverage: LATITUDE: -70.650000 * LONGITUDE: -8.250000
DATE/TIME START: 2017-01-01T00:00:00 * DATE/TIME END: 2017-01-31T23:59:00

Event(s): GVN (Georg von Neumayer) * LATITUDE: -70.650000 * LONGITUDE: -8.250000 * DATE/TIME: 1992-01-01T00:00:00 * ELEVAT
Parameter(s): DATE/TIME (date/time) * GEOCODE * PI: König-Langlo, Gert gert.koenig-langlo@awi.de
HEIGHT above ground [m] (height) * GEOCODE * PI: König-Langlo, Gert gert.koenig-langlo@awi.de
Short-wave downward (GLOBAL) radiation [W m-2] (SWD) * PI: König-Langlo, Gert gert.koenig-langlo@awi.de * METHOD
Direct radiation [W m-2] (DIR) * PI: König-Langlo, Gert gert.koenig-langlo@awi.de * METHOD: Pyrheliometer, Kipp & Zone
Diffuse radiation [W m-2] (DIF) * PI: König-Langlo, Gert gert.koenig-langlo@awi.de * METHOD: Pyranometer, Kipp & Zone
Long-wave downward radiation [W m-2] (LWD) * PI: König-Langlo, Gert gert.koenig-langlo@awi.de * METHOD: Pyrgeometer
Short-wave upward (REFLEX) radiation [W m-2] (SWU) * PI: König-Langlo, Gert gert.koenig-langlo@awi.de * METHOD: Py
Long-wave upward radiation [W m-2] (LWU) * PI: König-Langlo, Gert gert.koenig-langlo@awi.de * METHOD: Pyrgeometer
Air temperature at 2 m height [°C] (T2) * PI: König-Langlo, Gert gert.koenig-langlo@awi.de * METHOD: Thermometer
Humidity, relative [%] (RH) * PI: König-Langlo, Gert gert.koenig-langlo@awi.de * METHOD: Hygrometer
Station pressure [hPa] (PoPoPoPo) * PI: König-Langlo, Gert gert.koenig-langlo@awi.de * METHOD: Barometer
*/

```

Date/time	Height [m]	SWD [W m ⁻²]	DIR [W m ⁻²]	DIF [W m ⁻²]	LWD [W m ⁻²]	SWU [W m ⁻²]	LWU [W m ⁻²]
2017-01-01T00:00	2	19	0	19	243	18	303
2017-01-01T00:01	2	19	0	19	243	18	302
2017-01-01T00:02	2	20	0	19	245	18	302
2017-01-01T00:03	2	20	0	20	248	18	302
2017-01-01T00:04	2	20	0	19	246	18	302
2017-01-01T00:05	2	20	0	19	243	18	302
2017-01-01T00:06	2	20	0	19	244	18	302
2017-01-01T00:07	2	20	0	20	245	18	302
2017-01-01T00:08	2	21	0	20	250	19	302
2017-01-01T00:09	2	21	0	21	258	19	303

Figure 4. Example of the PANGAEA download format of a monthly data file (January 2017) from station GVN (Neumayer, Antarctica).

5 Some remarks for the best usage of BSRN data

The zip files which are publicly available at <https://doi.org/10.1594/PANGAEA.880000> contain all downward and (where available) upward radiation fluxes of one station and 1 year in monthly granularity. The “.tab” files can be opened in any text editor or spreadsheet program (e.g., Excel). They can be visualized individually (in the provided monthly granularity), for example, in PanPlot2 (see Sect. 3.2) by dragging the file onto the program icon. To obtain annual files one can, for example, drag all 12 months on the BSRN Toolbox and go to “Tools” ⇒ “Concatenate files”. One can then insert how many lines the program should skip (metadata header) and press OK. Be aware that the resulting annual file will be quite large.

PANGAEA itself offers comfortable ways to extract long periods of individual variables via the so-called data warehouse (<http://bsrn.awi.de/data/data-retrieval-via-pangaea/data-warehouse/>⁸). Go to <https://dataportals.pangaea.de/bsrn/>⁹, choose the station, logical record and time period and click on the respective value or “X” (for all datasets from that year and station), which will then lead to the respective search page on www.pangaea.de. Before the user can retrieve the selected files he/she has to log in at the very top right corner of this page (log in can be obtained from the WRMC director;

see <http://bsrn.awi.de/contact-persons/>¹⁰). After logging in, a green button labeled “Data Warehouse” appears on the top right corner of the page. A click on this button leads into the data warehouse, where parameter(s) can be chosen individually. Arithmetic averages can be calculated via “Method” but the results should be considered with care or should best not be used at all in cases where significant data gaps exist. In these cases the user should refer to more sophisticated averaging procedures as published in Roesch et al. (2011) or using a macro excel spreadsheet for hourly averages provided by Schild (2016). After clicking on “Start Data Warehouse Query” a window will pop up to download the output file. Depending on the amount of years, stations or parameters chosen the files can get quite large (easily > 100 MB).

A list of all radiation instruments used in BSRN is available at <https://www.pangaea.de/ddi?request=bsrn/BSRNMethods&format=html&title=BSRN+Methods>¹¹. To get all data from one instrument just copy–paste the instrument specifications into <https://www.pangaea.de/> and press search.

⁸last access: 28 July 2018

⁹last access: 28 July 2018

¹⁰last access: 28 July 2018

¹¹last access: 28 July 2018

6 Summary and outlook

In the 25 years of its existence, the Baseline Surface Radiation Network (BSRN) and its central archive have become an important component of global climate monitoring and climate research. Beginning as a contribution to the Global Energy and Water Cycle Experiment (GEWEX), the BSRN was later made part of the Global Climate Observing System (GCOS). Also, a formal cooperation agreement was established with the Network for the Detection of Atmospheric Composition Change (NDACC).

Validation of models (Wild, 2008; Müller Schmied et al., 2016) and satellite-based estimates of surface radiation (Pinker et al., 2005) remain the focus of users. However, the data have also been used for trend analyses (Long et al., 2009) and aerosol studies (Xia et al., 2007; Kim and Ramanathan, 2008). Over the past decade, BSRN data have increasingly been used by customers in the field of solar energy, who especially welcome the high temporal resolution of the data (e.g., Moshövel et al., 2015; Hofmann and Seckmeyer, 2017). Probably the first study that made full use of the BSRN network to estimate trends and constituted the solar brightening was Wild et al. (2005). More reviewed scientific papers referring to BSRN can be found at <http://bsrn.awi.de/other/publications/>¹².

Neither the BSRN stations nor the archive receive direct financial grants for their involvement in the BSRN. Despite this fact it has succeeded in developing a network that is setting standards for ground-level radiation measurements through volunteer efforts by a global community of scientists and technicians who have agreed to define and maintain a common set of standards. A total of 12 of the 59 stations have unfortunately closed, are no longer maintained, are inactive or seek funding. Thanks to the BSRN, the data collected by these stations remain available. However, BSRN is now in the process of adding nine new stations into the network (four from India, two from Taiwan, one from Russia, one from the Azores and one from Australia). One of the challenges the BSRN faces will therefore be the tremendous effort to continue its collection of high-quality, high-temporal-resolution measurements with as many representative stations as possible. As a fundamental environmental variable, the importance of permanently archived, freely accessible and high-quality radiation data cannot be valued highly enough.

Author contributions. All authors except for HG, SS, RS, AD and AO are responsible for station maintenance, data quality, technical aspects of the article; whereas HG, SS, RS, AD, GK-L, AO took care of the data curation and support within the WRMC. All authors except for RS gave input to manuscript, in which the main work has been done by AD, CL, GK-L.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. First and foremost, we would like to express our gratitude to all BSRN scientists and technicians working tediously to keep the high quality of BSRN standards up. Your dedicated work of calibrating, maintaining and quality checking sensors, instruments and data – often working extra hours to do so – is greatly appreciated. Thanks as well to all the institutions supporting the BSRN cause! We would also like to thank Wolfgang Cohrs for his excellent work in keeping the ftp server up and running, as well as his work for the BSRN homepage. We are grateful for the excellent and very useful comments of the two reviewers – your input definitely helped improve this paper. Last but not least we are indebted to the current (Skalde Lübberstedt) and former (Bonnie Raffel, Friedrich Richter) WRMC data curators – committed students with a knack for data visualization and the mind set on “finding the needle in the data haystack”. This article is dedicated to co-author Rainer Sieger, architect of the BSRN data archive in PANGAEA, responsible for maintaining the BSRN Toolbox and a good friend, who passed away unexpectedly in 2017; and also to Ells Dutton, the original and long-serving BSRN project manager for the first two decades, who was essential in establishing BSRN as a leader in surface radiation measurements and improvements. Ells Dutton passed away suddenly in 2012.

Edited by: David Carlson

Reviewed by: Martin Wild and one anonymous referee

References

- BSRN: Report of the 12th Baseline Surface Radiation Network (BSRN) Scientific Review and Workshop, Potsdam, Germany, 1–3 August 2012, 1–34, available at: <http://hdl.handle.net/10013/epic.09bd5b76-1cf5-4460-9be6-f523aea1ad34> (last access: 28 July 2018), 2012.
- Charlson, R. J., Schwartz, S. E., Hales, J. M., Cess, R. D., Coakley Jr., J. A., Hansen, J. E., and Hofmann, D. J.: Climate Forcing by Anthropogenic Aerosols, *Science*, 255, 423–430, <https://doi.org/10.1126/science.255.5043.423>, 1992.
- Chylek, P., Lohmann, U., Dubey, M., Mishchenko, M., Kahn, R., and Ohmura, A.: Limits on climate sensitivity derived from recent satellite and surface observations, *J. Geophys. Res.*, 112, D24S04, <https://doi.org/10.1029/2007JD008740>, 2007.
- Dutton, E. G., Michalsky, J. J., Stoffel, T., Forgan, B. W., Hickey, J., Nelson, D. W., Alberta, T. L., and Reda, I.: Measurement of broadband diffuse solar irradiance using current commercial instrumentation with a correction for thermal offset errors, *J. Atmos. Ocean. Tech.*, 18, 297–314, 2001.
- Gilgen H., Whitlock, C., Koch, F., Müller, G., Ohmura, A., Steiger, D., and Wheeler, R.: Technical plan for BSRN (Baseline Surface Radiation Network) data management, Version 2.1, WMO/TD-No. 443, WCRP/WMO, 1995.
- Hegner H., Müller, G., Nespor, V., Ohmura, A., Steigrad, R., and Gilgen, H.: Technical Plan for BSRN Data Management – 1998 Update, WMO/TD-No. 882, WCRP/WMO, available at:

¹²last access: 28 July 2018

- <http://hdl.handle.net/10013/epic.39581.d001> (last access: 28 July 2018), 1998.
- Hofmann, M. and Seckmeyer, G.: Influence of various irradiance models and their combination on simulation results of photovoltaic systems, *Energies*, 10, 1495, <https://doi.org/10.3390/en10101495>, 2017.
- Iqbal, M.: *Introduction to Solar Radiation*, Academic Press, New York, 1983.
- König-Langlo, G.: Basic and other measurements of radiation at Neumayer Station (2017-01), Alfred Wegener Institute, Helmholtz Center for Polar and Marine Research, Bremerhaven, <https://doi.org/10.1594/PANGAEA.874567>, 2017.
- König-Langlo, G., Sieger, R., Schmithüsen, H., Bäcker, A., Richter, F., and Dutton, E. G.: The Baseline Surface Radiation Network and its World Radiation Monitoring Centre at the Alfred Wegener Institute, GCOS Report 174: Update of the technical plan for BSRN data management, World Meteorological Organization (WMO), hdl:10013/epic.42596.d001, 2013.
- Kim, D. Y. and Ramanathan, V.: Solar radiation budget and radiative forcing due to aerosols and clouds, *J. Geophys. Res.*, 113, D02203, <https://doi.org/10.1029/2007JD008434>, 2008.
- Kwok, R. and Untersteiner, N.: The thinning of Arctic sea ice, *Physics Today*, 64, 36, <https://doi.org/10.1063/1.3580491>, 2011.
- Lanconelli, C., Busetto, M., Dutton, E. G., König-Langlo, G., Maturilli, M., Sieger, R., Vitale, V., and Yamanouchi, T.: Polar baseline surface radiation measurements during the International Polar Year 2007–2009, *Earth Syst. Sci. Data*, 3, 1–8, <https://doi.org/10.5194/essd-3-1-2011>, 2011.
- Long, C. N. and Dutton, E. G.: BSRN Global Network recommended QC tests, V2.0, BSRN Technical Report, hdl:10013/epic.38770.d001, 2002.
- Long, C. N. and Shi, Y.: An automated quality assessment and control algorithm for surface radiation measurements, *Open Atmos. Sci. J.*, 2, 23–37, <https://doi.org/10.2174/1874282300802010023>, 2008.
- Long, C. N., Younkin, K., and Powell, D. M.: Analysis of the Dutton et al. IR Loss Correction Technique Applied to ARM Diffuse SW Measurements, Eleventh ARM Science Team Meeting Proceedings, Atlanta, Georgia, 19–23 March 2001, available at: <http://hdl.handle.net/10013/epic.324172b4-606e-4a17-95f6-9337e26914da> (last access: 28 July 2018), 2001.
- Long, C. N., Dutton, E. G., Augustine, J. A., Wiscombe, W., Wild, M., McFarlane, S. A., and Flynn, C. J.: Significant Decadal Brightening of Downwelling Shortwave in the Continental US, *J. Geophys. Res.*, 114, D00D06, <https://doi.org/10.1029/2008JD011263>, 2009.
- Matsui, N., Long, C. N., Augustine, J., Halliwell, D., Uttal, T., Longenecker, D., Niebergall, O., Wendell, J., and Albee, R.: Evaluation of Arctic broadband surface radiation measurements, *Atmos. Meas. Tech.*, 5, 429–438, <https://doi.org/10.5194/amt-5-429-2012>, 2012.
- McArthur, L. J. B.: Baseline Surface Radiation Network (BSRN), WMO/TD-No. 879, WCRP/WMO, hdl:10013/epic.52032.d001, 2005.
- Michalsky, J.: The Astronomical Almanac's algorithm for approximate solar position (1950–2050), *Solar Energy*, 40, 227–235, 1988.
- Michalsky, J., Kutchenreiter, M., and Long, C. N.: Significant Improvements in Pyranometer Offsets Using Ventilation Strategies, *JAOTech*, 34, 1323–1332, <https://doi.org/10.1175/JTECH-D-16-0224.1>, 2017.
- Moshövel, J., Kairies, K.-P., Magnor, D., Leuthold, M., Bost, M., Gährs, S., Szczechowicz, E., Cramer, M., and Sauer, D. U.: Analysis of the maximal possible grid relief from PV-peak-power impacts by using storage systems for increased self-consumption, *Appl. Energ.*, 137, 567–575, <https://doi.org/10.1016/j.apenergy.2014.07.021>, 2015.
- Müller Schmied, H., Müller, R., Sanchez-Lorenzo, A., Ahrens, B., and Wild, M.: Evaluation of radiation components in a global freshwater model with station-based observations, *Water*, 8, 450, <https://doi.org/10.3390/w8100450>, 2016.
- Nyeki, S., Wacker, S., Gröbner, J., Finsterle, W., and Wild, M.: Revising shortwave and longwave radiation archives in view of possible revisions of the WSG and WISG reference scales: methods and implications, *Atmos. Meas. Tech.*, 10, 3057–3071, <https://doi.org/10.5194/amt-10-3057-2017>, 2017.
- Ohmura, A., Dutton, E. G., Forgan, B., Fröhlich, C., Gilgen, H., Hegner, H., Heimo, A., König-Langlo, G., McArthur, B., Müller, G., Philipona, R., Pinker, R., Whitlock, C. H., Dehne, K., and Wild, M.: Baseline surface radiation network (BSRN/WCRP): New precision radiometry for climate research, *B. Am. Meteorol. Soc.*, 79, 2115–2136, [https://doi.org/10.1175/1520-0477\(1998\)079<2115:BSRNBW>2.0.CO;2](https://doi.org/10.1175/1520-0477(1998)079<2115:BSRNBW>2.0.CO;2), 1998.
- Olefs, M., Baumgartner, D. J., Obleitner, F., Bichler, C., Foelsche, U., Pietsch, H., Rieder, H. E., Weihs, P., Geyer, F., Haiden, T., and Schöner, W.: The Austrian radiation monitoring network ARAD – best practice and added value, *Atmos. Meas. Tech.*, 9, 1513–1531, <https://doi.org/10.5194/amt-9-1513-2016>, 2016.
- Pinker, R. T., Zhang, B., and Dutton, E. G.: Do Satellites Detect Trends in Surface Solar Radiation?, *Science*, 308, 850–854, <https://doi.org/10.1126/science.1103159>, 2005.
- Roesch, A., Wild, M., Ohmura, A., Dutton, E. G., Long, C. N., and Zhang, T.: Assessment of BSRN radiation records for the computation of monthly means, *Atmos. Meas. Tech.*, 4, 339–354, <https://doi.org/10.5194/amt-4-339-2011>, 2011.
- Schild, P.: Macro-enabled Excel spreadsheet to calculate hourly-averages from BSRN .dat files, Bremerhaven, PANGAEA, hdl:10013/epic.48977.d002, 2016.
- Schmithüsen, H., Sieger, R., König-Langlo, G.: BSRN Toolbox V2.0 – a tool to create quality checked output files from BSRN datasets and station-to-archive files, Alfred Wegener Institute, Helmholtz Center for Polar and Marine Research, Bremerhaven, <https://doi.org/10.1594/PANGAEA.774827>, 2012.
- Sieger, R. and Grobe, H.: PanTool – a Swiss Army Knife for data conversion and recalculation, Alfred Wegener Institute, Helmholtz Center for Polar and Marine Research, Bremerhaven, PANGAEA, <https://doi.org/10.1594/PANGAEA.787549>, 2012.
- Sieger, R. and Grobe, H.: PanPlot 2 – software to visualize profiles and time series, Alfred Wegener Institute, Helmholtz Center for Polar and Marine Research, Bremerhaven, PANGAEA, <https://doi.org/10.1594/PANGAEA.816201>, 2013.
- Vuilleumier, L., Hauser, M., Félix, C., Vignola, F., Blanc, P., Kazantzidis, A., and Calpini, B.: Accuracy of ground surface broadband shortwave radiation monitoring, *J. Geophys. Res.-Atmos.*, 119, 13838–13860, <https://doi.org/10.1002/2014JD022335>, 2014.

- Wild, M.: Short-wave and long-wave surface radiation budgets in GCMs: a review based on the IPCC-AR4/CMIP3 models, *Tellus*, 60, 932–945, <https://doi.org/10.1111/j.1600-0870.2008.00342.x>, 2008.
- Wild, M., Gilgen, H., Roesch, A., Ohmura, A., Long, C., Dutton, E., Forgan, B., Kallis, A., Russak, V., and Tsvetkov, A.: From dimming to brightening: Decadal changes in solar radiation at the Earth's surface, *Science*, 308, 847–850, <https://doi.org/10.1126/science.1103215>, 2005.
- Wild, M., Folini, D., Schär, C., Loeb, N., Dutton, E. G., and König-Langlo, G.: The global energy balance from a surface perspective, *Clim. Dynam.*, 40, 3107–3134, <https://doi.org/10.1007/s00382-012-1569-8>, 2013.
- Wild, M., Ohmura, A., Schär, C., Müller, G., Folini, D., Schwarz, M., Hakuba, M. Z., and Sanchez-Lorenzo, A.: The Global Energy Balance Archive (GEBA) version 2017: a database for worldwide measured surface energy fluxes, *Earth Syst. Sci. Data*, 9, 601–613, <https://doi.org/10.5194/essd-9-601-2017>, 2017.
- Xia, X., Li, Z., Wang, P., Chen, H., and Cribb, M.: Estimation of aerosol effects on surface irradiance based on measurements and radiative transfer model simulations in northern China, *J. Geophys. Res.*, 112, D22S10, <https://doi.org/10.1029/2006JD008337>, 2007.
- Younkin, K. and Long, C. N.: Improved Correction of IR Loss in Diffuse Shortwave Measurements: An ARM Value Added Product, Atmospheric Radiation Measurement Program Technical Report, ARM TR-009, 50 pp., 2004.