



**HAL**  
open science

## A 2D model for a highly heterogeneous plate

Ahmed Boughammoura, Leila Rahmani, Ali Sili

► **To cite this version:**

Ahmed Boughammoura, Leila Rahmani, Ali Sili. A 2D model for a highly heterogeneous plate. *Ricerche di matematica*, In press, <10.1007/s11587-021-00671-4>. <hal-03662166>

**HAL Id: hal-03662166**

**<https://hal.science/hal-03662166v1>**

Submitted on 9 May 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

# A 2D model for a highly heterogeneous plate

Ahmed Boughammoura\*, Leila Rahmani<sup>†</sup> and Ali Sili<sup>‡</sup>

## Abstract

In this paper we investigate the  $2d$ -model for a thin plate  $\Omega_\varepsilon := \omega \times \varepsilon I$  of  $\mathbb{R}^3$  having two components: a circular stiff fiber  $F_\varepsilon$  and its complement the soft matrix  $M_\varepsilon$  with  $\frac{1}{\varepsilon^2}$  as a ratio between their respective elasticity coefficients. We prove that the limit model is associated to a nonlocal system involving Kirchoff-Love displacements in the fiber and we exhibit a corrector for the displacements in the initial cylindrical structure of  $\mathbb{R}^3$ .

**Keywords:** plate, thin structure, corrector, nonlocal.

**AMS Classification::** 35B25, 35B27, 35B40, 76M50, 74K10.

## 1 Introduction, notations and setting of the problem

The aim of this work is the study of the asymptotic behavior of the solutions of the linearized system of elasticity posed in a cylindrical domain  $\Omega_\varepsilon := \omega \times \varepsilon I$  of  $\mathbb{R}^3$  which is the configuration domain of a composite material. The material is made up of two components with high contrast: the first one  $F_\varepsilon$  representing the stiff part of the material has an elasticity tensor

---

\*Université de Monastir, Institut Supérieur d'Informatique et de Mathématiques de Monastir, Analyse et Contrôle des EDP, UR 13ES64, 5019, Monastir, Tunisia, email: ahmed.boughammoura@gmail.com

<sup>†</sup>Laboratoire de mathématiques pures et appliquées, Mouloud Mammeri University of Tizi-Ouzou, Algeria, email: rahmani-lei@yahoo.fr

<sup>‡</sup>Institut de Mathématiques de Marseille (I2M) UMR 7373, Aix-Marseille Université, CNRS, CMI, 39 rue F. Joliot-Curie, 13453 Marseille cedex 13, France, email: ali.sili@univ-amu.fr

with coefficients of order 1. The second component  $M_\varepsilon$  (the soft material) surrounds the first one and the coefficients of its elasticity tensor are of order  $\varepsilon^2$ .

Under an appropriate assumption on the volumes forces, we aim to approximate the behavior of the displacements and that of the associated tensors as the small parameter  $\varepsilon$  tends to zero. Hence the present work may be viewed as the  $3d - 2d$  version of the study addressed in [12] where the  $3d - 1d$  reduction of dimension problem was considered as well as the homogenization of a  $\varepsilon$ -periodic fibered medium inducing a local  $3d - 1d$  reduction of dimension. It was proved in [12] that the homogenized problem in a such setting is a copy of the one-dimensional problem obtained in the  $3d - 1d$  study. Although one can also consider here the  $\varepsilon$ -periodic homogenization problem of a medium containing  $\frac{1}{\varepsilon}$  cells which are the translates of  $\Omega_\varepsilon$  in such a way that the homogenization process leads to a local  $3d - 2d$  reduction of dimension, for the sake of brevity we restrict ourselves to the  $3d - 2d$  reduction of dimension problem arising in the single composite structure  $\Omega_\varepsilon$ . We consider the critical case where the ratio between the elasticity coefficients of the two components is equal to  $\frac{1}{\varepsilon^2}$  but other scalings may be considered as pointed out in [12], see also [9, 14].

On the other hand, we deal with general elasticity tensors including anisotropic materials, see also [3, 7, 11, 13]. Several studies on composite materials with high contrast between their components have been performed during the last years, see for instance [2, 4–6, 9–11].

It is known that for this kind of materials, the limit problem has in general a different structure than the starting problem. In particular, nonlocal phenomena can appear at the limit. We show here that the limit problem obtained after reduction of dimension  $3d - 2d$  is indeed a nonlocal problem. Theorem 3.2 below, which gives the limit problem, shows that the displacements in the circular fiber  $F$  are essentially of Kirchhoff-Love type and the associated equation may be obtained from system (29) by choosing  $\bar{z} = 0$ ; but to determine the limit displacements in the structure, another equation related to the matrix  $M$  (the outside of the fiber) is necessary. That equation is obtained by choosing  $\bar{u} = \bar{v} = 0$  in (29). This nonlocal phenomenon is emphasized in Theorem 3.4 which gives the limit of the average of the displacements  $\hat{u}^\varepsilon$  in the three-dimensional structure  $\Omega_\varepsilon$ . In particular convergences (33) and (34) show clearly that the limit displacements are determined after solving the two equations posed in  $F$  and in  $M$  respectively.

In terms of correctors, our result shows that the transversal displacements

$u_\alpha^\varepsilon$  behave as  $u_\alpha + \varepsilon v_\alpha + z_\alpha$  while the horizontal displacements  $u_3^\varepsilon$  behave as  $u_3 + \varepsilon^2 v_3 + \varepsilon z_3$ .

We now make more precise the notations we will use throughout the paper.

A vector  $x$  in  $\mathbb{R}^3$  is denoted by  $x = (x', x_3)$  where  $x_3$  denotes the vertical coordinate. Latin indices will usually range from 1 to 3, and Greek ones take values in  $\{1, 2\}$ ; the summation convention applies whenever indices are repeated. We write  $\partial_i := \frac{\partial}{\partial x_i}$  and  $\partial_{ij} := \frac{\partial^2}{\partial x_i \partial x_j}$ . The gradient with respect to  $x$  and  $x'$  are denoted by  $\nabla$  and  $\nabla'$  respectively. Given any  $\phi \in (\mathcal{D}'(\Omega))^3$ , we denote the strain of  $\phi$  by  $\mathbf{E}\phi := \text{sym}\nabla\phi = \frac{1}{2}(\nabla\phi + \nabla\phi^T)$ . We shall also use the following matrices notations

$$\nabla\phi = \left( \begin{array}{c|c} \partial_\alpha\phi_\beta & \partial_3\phi_\beta \\ \hline \partial_\alpha\phi_3 & \partial_3\phi_3 \end{array} \right),$$

$$\mathbf{E}\phi = \left( \begin{array}{c|c} (\mathbf{E}\phi)_{\alpha\beta} & (\mathbf{E}\phi)_{\alpha 3} \\ \hline (\mathbf{E}\phi)_{\alpha 3} & (\mathbf{E}\phi)_{33} \end{array} \right).$$

Let  $\omega \subset \mathbb{R}^2$  be an open, bounded, simply connected set with Lipschitz boundary  $\partial\omega$ . Let  $\mathbf{I} := (-\frac{1}{2}, \frac{1}{2})$ ,  $\mathbf{J} := (-\frac{r}{2}, \frac{r}{2})$ ,  $0 < r < \frac{1}{2}$ ,  $\Omega := \omega \times \mathbf{I}$ ,  $F := \omega \times \mathbf{J}$ ,  $M := \omega \times (\mathbf{I} \setminus \bar{\mathbf{J}})$ , where  $\bar{\mathbf{J}}$  denotes the closure of  $\mathbf{J}$ . According to Remark 1.1 below, physically we can think of  $\Omega$  as the reference configuration of a rescaled thin plate  $\Omega_\varepsilon$  reinforced by the fiber  $F_\varepsilon$  which is surrounded by the matrix  $M_\varepsilon$  occupied by a soft material. For every  $\varepsilon > 0$  we denote the diagonal matrix whose entries are 1, 1 and  $\varepsilon$  by  $R^\varepsilon := \text{diag}(1, 1, \varepsilon)$ , then the scaled gradient  $\nabla^\varepsilon\phi$  and the scaled strain  $\mathbf{E}^\varepsilon\phi$  are defined respectively by

$$\nabla^\varepsilon\phi := (R^\varepsilon)^{-1}\nabla\phi(R^\varepsilon)^{-1} = \left( \begin{array}{c|c} \partial_\alpha\phi_\beta & \frac{1}{\varepsilon}\partial_3\phi_\beta \\ \hline \frac{1}{\varepsilon}\partial_\alpha\phi_3 & \frac{1}{\varepsilon^2}\partial_3\phi_3 \end{array} \right), \quad (1)$$

and

$$\mathbf{E}^\varepsilon\phi := \text{sym}\nabla^\varepsilon\phi = \left( \begin{array}{c|c} (\mathbf{E}\phi)_{\alpha\beta} & \frac{1}{\varepsilon}(\mathbf{E}\phi)_{\alpha 3} \\ \hline \frac{1}{\varepsilon}(\mathbf{E}\phi)_{\alpha 3} & \frac{1}{\varepsilon^2}(\mathbf{E}\phi)_{33} \end{array} \right). \quad (2)$$

We are now in position to state the problem.

Let  $\mathbb{A} \in L^\infty(\Omega)$  be a symmetric fourth-order tensor field. We assume that  $\mathbb{A}$  fulfills the following assumptions:

$$\begin{cases} \mathbb{A}_{ijkl} = \mathbb{A}_{jikl} = \mathbb{A}_{klij}, \text{ a.e. in } \Omega, \\ \exists C > 0, \mathbb{A}_{ijkl}\xi_{kl}\xi_{ij} \geq C\xi_{ij}\xi_{ij}, \forall \xi \in \mathbb{R}^9 \text{ s.t. } \xi^T = \xi. \end{cases} \quad (3)$$

We shall assume that the plate is clamped at the lateral boundary of  $\Omega$  and subjected to body forces  $f \in L^2(\Omega; \mathbb{R}^3)$ , we thus set

$$H_L^1(\Omega) := \{u \in H^1(\Omega; \mathbb{R}^3) : u = 0 \text{ on } \partial\omega \times \mathbb{I}\}.$$

Consider the displacement field  $u^\varepsilon$  solution of the following system:

$$\begin{cases} u^\varepsilon \in H_L^1(\Omega), \forall \phi \in H_L^1(\Omega), \\ \int_{\Omega} (\chi_F + \varepsilon^2 \chi_M) \mathbb{A} \mathbf{E}^\varepsilon(u^\varepsilon) \cdot \mathbf{E}^\varepsilon(\phi) dx = \int_{\Omega} f \phi dx. \end{cases} \quad (4)$$

By virtue of the assumptions on the tensor  $\mathbb{A}$  and the body forces  $F$ , for every  $\varepsilon > 0$ , the problem (4) admits a unique solution by the Lax-Milgram Theorem.

**Remark 1.1** *As in [8], the homothety along the vertical axis defined by  $r^\varepsilon(x) := (x', \varepsilon x_3)$  transforms  $\Omega$ ,  $F$  and  $M$  respectively into  $\Omega_\varepsilon := \omega \times \varepsilon \mathbb{I}$ ,  $F_\varepsilon := \omega \times \varepsilon \mathbb{J}$  and  $M_\varepsilon := \omega \times \varepsilon (\mathbb{I} \setminus \bar{\mathbb{J}})$ . Then, the problem (4) is the variational version in the fixed reference configuration  $\Omega$  of the elasticity problem (6) below posed in the variable thin domain  $\Omega_\varepsilon$ . Indeed, for any  $v : \Omega \mapsto \mathbb{R}^3$  we define  $\hat{v} : \Omega_\varepsilon \mapsto \mathbb{R}^3$  by*

$$\hat{v}(x', \varepsilon x_3) := ((R^\varepsilon)^{-1}v \circ (r^\varepsilon)^{-1})(x', \varepsilon x_3) = (v_\alpha(x), \frac{1}{\varepsilon}v_3(x)). \quad (5)$$

In addition

$$\nabla \hat{v} = \mathbb{H}^\varepsilon v \circ (r^\varepsilon)^{-1}, \quad \mathbf{E} \hat{v} = \mathbf{E}^\varepsilon v \circ (r^\varepsilon)^{-1}.$$

With these new unknowns, the problem (4) may be rewritten as

$$\begin{cases} \hat{u}^\varepsilon \in H_L^1(\Omega_\varepsilon), \forall \phi \in H_L^1(\Omega_\varepsilon), \\ \int_{\Omega_\varepsilon} (\chi_{F_\varepsilon} + \varepsilon^2 \chi_{M_\varepsilon}) \mathbb{A}^\varepsilon \mathbf{E}(\hat{u}^\varepsilon) \cdot \mathbf{E}(\phi) dx = \int_{\Omega_\varepsilon} f^\varepsilon \phi dx \end{cases} \quad (6)$$

where  $H_L^1(\Omega_\varepsilon) := \{u \in H^1(\Omega_\varepsilon; \mathbb{R}^3) : u = 0 \text{ on } \partial\omega \times \varepsilon\mathbf{I}\}$ ,  $\mathbb{A}^\varepsilon := \mathbb{A} \circ (r^\varepsilon)^{-1}$ ,  $f^\varepsilon = R^\varepsilon f \circ (r^\varepsilon)^{-1}$ . Thus, the components of the loads are

$$f^\varepsilon = (f_\alpha \circ (r^\varepsilon)^{-1}, \varepsilon f_3 \circ (r^\varepsilon)^{-1}). \quad (7)$$

We study the behavior as  $\varepsilon \rightarrow 0$  of the sequence  $\{u^\varepsilon\}$ , solution of (4), through the forthcoming steps. That behavior will be described through Theorem 3.2 and then we will deduce the behavior of the sequence  $\hat{u}^\varepsilon$  solution of (6) in Theorem 3.4.

## 2 A priori estimates

First, we shall recall a rescaled Korn's inequality proved in [1]

$$\|\mathbf{E}\hat{v}\|_{L^2(F_\varepsilon)}^2 \geq C\varepsilon^2 \left( \|\hat{v}\|_{L^2(F_\varepsilon)}^2 + \|\nabla\hat{v}\|_{L^2(F_\varepsilon)}^2 \right), \quad \forall v \in H_L^1(F), \quad (8)$$

where

$$H_L^1(F) := \{u \in H^1(F; \mathbb{R}^3) : u = 0 \text{ on } \partial\omega \times \mathbf{J}\}$$

and  $\hat{v} = (R^\varepsilon)^{-1}v \circ (r^\varepsilon)^{-1}$ . As a consequence, we deduce

$$\|\mathbf{E}^\varepsilon v\|_{L^2(F)}^2 \geq C\varepsilon^2 \left( \|(R^\varepsilon)^{-1}v\|_{L^2(F)}^2 + \|\nabla^\varepsilon v\|_{L^2(F)}^2 \right), \quad \forall v \in H_L^1(F). \quad (9)$$

On the other hand, repeating the same arguments used in [12, Lemma 4.4], the following inequality holds true

$$\left( \|\nabla^\varepsilon v\|_{L^2(F)}^2 + \|\mathbf{E}^\varepsilon v\|_{L^2(M)}^2 \right) \geq C \|\nabla^\varepsilon v\|_{L^2(M)}^2, \quad \forall v \in H_L^1(\Omega). \quad (10)$$

For  $v \in H_L^1(\Omega)$  we set  $v' := (v_1, v_2)$ ,  $\nabla'v' := (\partial_\beta v_\alpha)_{\alpha, \beta=1,2}$  and

$$\mathcal{F}_\varepsilon(v) := \int_\Omega (\chi_F + \varepsilon^2 \chi_M) \mathbb{A} \mathbf{E}^\varepsilon v \cdot \mathbf{E}^\varepsilon v dx. \quad (11)$$

By using the ellipticity assumption (3) we obtain, for all  $v \in H_L^1(\Omega)$

$$\mathcal{F}_\varepsilon(v) \geq C \left( \|\mathbf{E}^\varepsilon v\|_{L^2(F)}^2 + \varepsilon^2 \|\mathbf{E}^\varepsilon v\|_{L^2(M)}^2 \right) \geq C \|\mathbf{E}v\|_{L^2(F)}^2 \geq C \|v\|_{H^1(F)}^2, \quad (12)$$

where the last inequality follows from the Korn's inequality. Furthermore, by virtue of (9) and (10) one has

$$\mathcal{F}_\varepsilon(v) \geq C\varepsilon^2 \left( \|\nabla^\varepsilon v\|_{L^2(F)}^2 + \|\mathbf{E}^\varepsilon v\|_{L^2(M)}^2 \right) \geq C\varepsilon^2 \|\nabla^\varepsilon v\|_{L^2(M)}^2. \quad (13)$$

From (13), one has for  $\varepsilon$  small enough,

$$\mathcal{F}_\varepsilon(v) \geq C\varepsilon^2 \left( \frac{1}{\varepsilon^2} \|\nabla' v_3\|_{L^2(M)}^2 + \frac{1}{\varepsilon^4} \|\partial_3 v_3\|_{L^2(M)}^2 \right) \geq C \|\nabla v_3\|_{L^2(M)}^2. \quad (14)$$

By means of the Poincaré's inequality, one has

$$\mathcal{F}_\varepsilon(v) \geq C \|v_3\|_{H^1(M)}^2. \quad (15)$$

Likewise, it holds that

$$\mathcal{F}_\varepsilon(v) \geq C \left( \varepsilon^2 \|\nabla' v'\|_{L^2(M)}^2 + \|\partial_3 v'\|_{L^2(M)}^2 \right) \geq C \|\partial_3 v'\|_{L^2(M)}^2. \quad (16)$$

From the following Poincaré's type inequality

$$\|v'\|_{L^2(M)}^2 \leq C \left( \|v'\|_{L^2(F)}^2 + \|\partial_3 v'\|_{L^2(M)}^2 \right),$$

(16) and (12), we get the following inequality

$$\mathcal{F}_\varepsilon(v) \geq C \|v'\|_{L^2(M)}^2. \quad (17)$$

It follows from (15) and (17) that

$$\mathcal{F}_\varepsilon(v) \geq C \left( \|v_3\|_{H^1(M)}^2 + \|v'\|_{L^2(\omega; H^1(\Gamma \setminus \bar{J}))}^2 \right). \quad (18)$$

As an immediate consequence of the previous inequalities, the following a priori estimates hold true.

**Theorem 2.1** *Let  $u_\varepsilon$  be the solution of problem (4). Then*

$$\sup_\varepsilon \left( \|u^\varepsilon\|_{H^1(F)} + \|\varepsilon \nabla^\varepsilon u^\varepsilon\|_{L^2(F)} + \|\mathbf{E}^\varepsilon u^\varepsilon\|_{L^2(F)} \right) \leq C, \quad (19)$$

$$\sup_\varepsilon \left( \|u_3^\varepsilon\|_{H^1(M)} + \|u_\alpha^\varepsilon\|_{L^2(\omega; H^1(\Gamma \setminus \bar{J}))} + \|\varepsilon \nabla^\varepsilon u^\varepsilon\|_{L^2(M)} \right) \leq C, \quad (20)$$

*in particular*

$$\sup_\varepsilon \left( \|u_3^\varepsilon\|_{H^1(\Omega)} + \|u_\alpha^\varepsilon\|_{L^2(\omega; H^1(\Gamma))} + \|\varepsilon \nabla^\varepsilon u^\varepsilon\|_{L^2(\Omega)} \right) \leq C, \quad \alpha = 1, 2. \quad (21)$$

**Proof.**

Taking  $\phi = u^\varepsilon$  in the problem (4), we derive

$$\mathcal{F}_\varepsilon(u^\varepsilon) \leq \|f\|_{L^2(\Omega)} \|u^\varepsilon\|_{L^2(\Omega)},$$

while from (12), (15) and (17) we find

$$\mathcal{F}_\varepsilon(u^\varepsilon) \geq C \|u^\varepsilon\|_{L^2(\Omega)}.$$

Thus

$$\sup_\varepsilon \mathcal{F}_\varepsilon(u^\varepsilon) \leq C.$$

Therefore, making use of (12)-(13) and then (13)-(18) we deduce (19) and (20) respectively.

### 3 Convergence results

We first define the following functional spaces:

$$H_{KL}(\Omega) := \{u \in H_L^1(\Omega) : (\mathbf{E}u)_{i3} = 0, i = 1, 2, 3\}$$

which is the space of Kirchhoff-Love displacements on  $\Omega$ ; it can be characterized also as

$$H_{KL}(\Omega) = \{u \in H_L^1(\Omega), \exists (g_\alpha, g_3) \in H_0^1(\omega) \times H_0^2(\omega) : u_\alpha = g_\alpha - x_3 \partial_\alpha g_3, u_3 = g_3\},$$

for brevity we set  $\mathcal{U} := H_{KL}(\Omega)$ ,

$$\mathcal{V} := \left\{ v \in L^2(\omega; H^1(\mathbb{J}))^3 : \int_{\mathbb{J}} v_i(x', t) dt = 0, \text{ a.e. } x' \in \omega, i = 1, 2, 3 \right\},$$

$$\mathcal{Z} := \left\{ z \in L^2(\omega; H_m^1(\mathbb{I}))^3 : z_i = 0, \text{ a.e. in } F, i = 1, 2, 3 \right\},$$

where  $H_m^1(\mathbb{I}) := \left\{ \psi \in H^1(\mathbb{I}) : \int_{\mathbb{I}} \psi(t) dt = 0 \right\}$ .

The following lemma is concerned with some convergence results.



**Lemma 3.1** *There exists  $(u, v, z) \in \mathcal{U} \times \mathcal{V} \times \mathcal{Z}$ , such that, up to a subsequence, we have*

$$\begin{aligned} u_\alpha^\varepsilon &\rightharpoonup u_\alpha + z_\alpha \text{ in } L^2(\omega; H^1(\mathbb{I})), \quad u_3^\varepsilon \rightharpoonup u_3 \text{ in } H^1(\Omega), \\ \mathbf{E}^\varepsilon u^\varepsilon \chi_F &\rightharpoonup \left( \begin{array}{c|c} (\mathbf{E}u)_{\alpha\beta} & \partial_3 v_\alpha \\ \partial_3 v_\alpha & \partial_3 v_3 \end{array} \right) \chi_F \text{ in } L^2(\Omega)^{3 \times 3}, \\ \varepsilon \mathbf{E}^\varepsilon u^\varepsilon \chi_M &\rightharpoonup \left( \begin{array}{c|c} 0 & \frac{1}{2} \partial_3 z_\alpha \\ \frac{1}{2} \partial_3 z_\alpha & \partial_3 z_3 \end{array} \right) \chi_M \text{ in } L^2(\Omega)^{3 \times 3}. \end{aligned}$$

**Proof.**

**Step 1: convergence of  $(u^\varepsilon)_\varepsilon$ .**

From estimate (19) which implies that  $u_\alpha^\varepsilon$  is bounded in  $H^1(F)$  and from estimate (20) which implies that  $u_\alpha^\varepsilon$  is bounded in  $L^2(\omega; H^1(I \setminus \bar{J}))$  for  $\alpha = 1, 2$ , we conclude that there exist  $u_\alpha \in H^1(\Omega)$  such that for a subsequence

$$u_\alpha^\varepsilon \rightharpoonup u_\alpha \text{ in } H^1(F), \quad (\mathbf{E}^\varepsilon u^\varepsilon)_{\alpha\beta} \chi_F \rightharpoonup (\mathbf{E}u)_{\alpha\beta} \chi_F \text{ in } L^2(F). \quad (22)$$

On the other hand, from (21) we get that up to a subsequence,

$$\begin{cases} u_\alpha^\varepsilon \rightharpoonup \tilde{u}_\alpha \text{ in } L^2(\omega; H^1(\mathbb{I})), \\ u_3^\varepsilon \rightharpoonup u_3 \text{ in } H^1(\Omega). \end{cases} \quad (23)$$

Moreover, due to the estimate  $\|\frac{1}{\varepsilon} \partial_3 u_3^\varepsilon\|_{L^2(\Omega)} \leq C$  which is a consequence of (21), we have

$$\partial_3 u_3^\varepsilon \rightarrow 0 \text{ in } L^2(\Omega)$$

in such a way that

$$\partial_3 u_3 = 0.$$

Thus, there exists  $g_3 \in H_0^1(\omega)$  such that

$$u_3(x) = g_3(x') \text{ for a.e. } x \in \Omega. \quad (24)$$

Comparing the first convergences in (22) and (23), we derive the equality

$$\tilde{u}_\alpha = u_\alpha \text{ in } F. \quad (25)$$

On the other hand, using (19) we infer

$$(\mathbf{E}u^\varepsilon)_{\alpha 3} \rightharpoonup (\mathbf{E}\tilde{u})_{\alpha 3} \text{ in } L^2(F).$$

Since

$$(\mathbf{E}u^\varepsilon)_{i3} \rightarrow 0, \text{ in } L^2(F)$$

we are led to

$$(\tilde{u}_\alpha, u_3) \in H_{KL}(F).$$

By definition of the space  $H_{KL}(F)$ , the component  $u_3$  belongs to  $H_0^2(\omega)$ , so that equality (24) implies that  $g_3 \in H_0^2(\omega)$ . Hence, there exist  $g_\alpha \in H_0^1(\omega)$  for  $\alpha = 1, 2$ , such that

$$\tilde{u}_\alpha(x) = g_\alpha(x') - x_3 \partial_\alpha g_3(x'), \quad u_3(x) = g_3(x') \text{ for a. e. } x \in F.$$

For a. e.  $x \in \Omega$ , we set

$$\begin{aligned} u_\alpha(x) &:= g_\alpha(x') - x_3 \partial_\alpha g_3(x') \in H_L^1(\Omega), \\ z_\alpha(x) &:= \tilde{u}_\alpha(x) - u_\alpha(x) \in L^2(\omega; H^1(\mathbb{I})), \end{aligned}$$

so that  $z_\alpha = 0$ , a. e. in  $F$  as seen from (25) and  $u := (u_\alpha, u_3) \in \mathcal{U}$  and

$$u_\alpha^\varepsilon \rightharpoonup u_\alpha + z_\alpha, \text{ in } L^2(\omega; H^1(\mathbb{I})).$$

## Step 2: convergence of $(\mathbf{E}^\varepsilon u^\varepsilon)_\varepsilon$ .

To identify the limit of the sequence  $(\mathbf{E}^\varepsilon u^\varepsilon)_{i3} \chi_F$ , we introduce the following sequences :

$$\hat{u}_\alpha^\varepsilon := \int_{-\frac{1}{2}}^{x_3} \frac{1}{\varepsilon} \partial_\alpha u_3^\varepsilon(x', t) dt, \quad (26)$$

$$\begin{aligned} v_\alpha^\varepsilon &:= \frac{1}{\varepsilon} u_\alpha^\varepsilon + \hat{u}_\alpha^\varepsilon - \int_{\mathbb{J}} \left( \frac{1}{\varepsilon} u_\alpha^\varepsilon + \hat{u}_\alpha^\varepsilon \right) dx_3, \\ v_3^\varepsilon &:= \frac{1}{\varepsilon^2} u_3^\varepsilon - \int_{\mathbb{J}} \frac{1}{\varepsilon^2} u_3^\varepsilon dx_3. \end{aligned} \quad (27)$$

Obviously, one has

$$\begin{aligned} \partial_3 v_\alpha^\varepsilon &= \frac{2}{\varepsilon} (\mathbf{E}u^\varepsilon)_{\alpha 3}, \\ \partial_3 v_3^\varepsilon &= \frac{1}{\varepsilon^2} (\mathbf{E}u^\varepsilon)_{33}. \end{aligned}$$

Since  $\frac{2}{\varepsilon}(\mathbf{E}u^\varepsilon)_{\alpha 3}$  and  $\frac{1}{\varepsilon^2}(\mathbf{E}u^\varepsilon)_{33}$  are bounded in  $L^2(F)$ , due to (19), and  $v^\varepsilon := (v_\alpha^\varepsilon, v_3^\varepsilon)$  has mean-value zero with respect to  $x_3$ , the sequence  $v^\varepsilon$  is bounded in  $L^2(\omega; H_m^1(\mathbf{J}))^3$ , where  $H_m^1(\mathbf{J}) := \left\{ \psi \in H^1(\mathbf{J}) : \int_{\mathbf{J}} \psi(t) dt = 0 \right\}$ .

Therefore there exists some  $v \in L^2(\omega; H_m^1(\mathbf{J}))^3$  such that

$$v^\varepsilon \rightharpoonup v \text{ in } L^2(\omega; H_m^1(\mathbf{J}))^3.$$

In particular, one has  $v \in \mathcal{V}$ ,

$$(\mathbf{E}^\varepsilon u^\varepsilon)_{\alpha 3} \chi_F \rightharpoonup \frac{1}{2} \partial_3 v_\alpha \chi_F \text{ in } L^2(\Omega),$$

and

$$(\mathbf{E}^\varepsilon u^\varepsilon)_{33} \chi_F \rightharpoonup \partial_3 v_3 \chi_F \text{ in } L^2(\Omega).$$

As a consequence, one obtains the following convergence

$$\mathbf{E}^\varepsilon u^\varepsilon \chi_F \rightharpoonup \left( \begin{array}{c|c} (\mathbf{E}u)_{\alpha\beta} & \frac{1}{2} \partial_3 v_\alpha \\ \hline \frac{1}{2} \partial_3 v_\alpha & \partial_3 v_3 \end{array} \right) \chi_F \text{ in } L^2(\Omega)^{3 \times 3},$$

where  $(u, v) \in \mathcal{U} \times \mathcal{V}$ .

We now seek for the limit of the sequence  $\varepsilon \mathbf{E}^\varepsilon u^\varepsilon \chi_M$ . To that aim, we consider the following sequence

$$\begin{aligned} z_\alpha^\varepsilon &:= u_\alpha^\varepsilon + \hat{u}_\alpha^\varepsilon - \int_{\mathbf{I}} (u_\alpha^\varepsilon + \hat{u}_\alpha^\varepsilon) dx_3, \\ z_3^\varepsilon &:= \frac{1}{\varepsilon} u_3^\varepsilon - \int_{\mathbf{I}} \frac{1}{\varepsilon} u_3^\varepsilon dx_3. \end{aligned} \tag{28}$$

where  $\hat{u}^\varepsilon$  is defined by (26). Since

$$\partial_3 z_\alpha^\varepsilon = 2(\mathbf{E}u^\varepsilon)_{\alpha 3}, \tag{29}$$

$$\partial_3 z_3^\varepsilon = \frac{1}{\varepsilon} (\mathbf{E}u^\varepsilon)_{33}. \tag{30}$$

Due to (21),  $2(\mathbf{E}u^\varepsilon)_{\alpha 3}$  and  $\frac{1}{\varepsilon} (\mathbf{E}u^\varepsilon)_{33}$  are bounded in  $L^2(\Omega)$  with mean-value zero with respect to  $x_3$ ; hence the sequence  $z^\varepsilon$  is bounded in  $L^2(\omega; H_m^1(\mathbf{I}))^3$ . Therefore there exists some  $\hat{z} \in L^2(\omega; H_m^1(\mathbf{I}))^3$ , such that

$$z^\varepsilon \rightharpoonup \hat{z} \text{ in } L^2(\omega; H_m^1(\mathbf{I}))^3.$$

We claim that  $\hat{z}_\alpha = z_\alpha$ . Indeed,

$$\partial_3 u_\alpha^\varepsilon \rightharpoonup \partial_3 u_\alpha + \partial_3 z_\alpha \text{ in } L^2(\Omega)$$

and

$$\partial_3 z_\alpha^\varepsilon \rightharpoonup \partial_3 \hat{z}_\alpha \text{ in } L^2(\Omega).$$

But

$$\partial_3 z_\alpha^\varepsilon = \partial_3 u_\alpha^\varepsilon + \partial_\alpha u_3^\varepsilon \rightharpoonup \partial_3 u_\alpha + \partial_3 z_\alpha + \partial_\alpha u_3 \text{ in } L^2(\Omega).$$

Thus

$$\partial_3 \hat{z}_\alpha = \partial_3 u_\alpha + \partial_3 z_\alpha + \partial_\alpha u_3.$$

Since  $u \in \mathcal{U}$ , we have

$$\partial_3 u_\alpha + \partial_\alpha u_3 = 0,$$

and therefore

$$\partial_3 \hat{z}_\alpha = \partial_3 z_\alpha \text{ i.e. } \hat{z}_\alpha(x) = z_\alpha(x) + c(x'), \text{ a.e. } x \in \Omega.$$

In particular

$$\hat{z}_\alpha(x) = c(x'), \text{ a.e. } x \in F$$

since

$$z_\alpha(x) = 0, \text{ a.e. } x \in F.$$

On the other hand one has by virtue of (20),

$$z^\varepsilon \rightarrow 0 \text{ in } L^2(F).$$

Hence

$$c(x') = 0, \text{ a.e. } x \in F$$

and therefore

$$\hat{z}_\alpha(x) = z_\alpha(x), \text{ a.e. } x \in \Omega.$$

This proves our claim.

Finally, from (29) and (30) we deduce that

$$\varepsilon(\mathbf{E}^\varepsilon u^\varepsilon)_{\alpha 3} \chi_M \rightharpoonup \frac{1}{2} \partial_3 z_\alpha \chi_M, \quad \varepsilon(\mathbf{E}^\varepsilon u^\varepsilon)_{33} \chi_M \rightharpoonup (\mathbf{E}z)_{33} \chi_M \text{ in } L^2(\Omega),$$

where  $z_3 := \hat{z}_3$ .

Moreover, since  $\varepsilon u_\alpha^\varepsilon \rightarrow 0$  in  $L^2(M)$  then  $\varepsilon(\mathbf{E}^\varepsilon u^\varepsilon)_{\alpha\beta} \chi_M \rightarrow 0$  in  $L^2(\Omega)$ . We conclude that

$$\varepsilon \mathbf{E}^\varepsilon u^\varepsilon \chi_M \rightharpoonup \left( \begin{array}{c|c} 0 & \frac{1}{2} \partial_3 z_\alpha \\ \hline \frac{1}{2} \partial_3 z_\alpha & \partial_3 z_3 \end{array} \right) \chi_M \text{ in } L^2(\Omega)^{3 \times 3}.$$

This ends the proof of Lemma 3.1.

In the following theorem, we state our main result.

**Theorem 3.2** *There exists  $(u, v, z) \in \mathcal{U} \times \mathcal{V} \times \mathcal{Z}$ , such that the sequence of solutions  $u^\varepsilon$  of the problem (4) fulfills the following strong convergences*

$$u_\alpha^\varepsilon \rightarrow u_\alpha + z_\alpha \text{ strongly in } L^2(\omega; H^1(\mathbb{I})), \quad u_3^\varepsilon \rightarrow u_3 \text{ strongly in } H^1(\Omega),$$

$$\mathbf{E}^\varepsilon u^\varepsilon \chi_F \rightarrow \left( \begin{array}{c|c} (\mathbf{E}u)_{\alpha\beta} & \frac{1}{2} \partial_3 v_\alpha \\ \hline \frac{1}{2} \partial_3 v_\alpha & \partial_3 v_3 \end{array} \right) \chi_F \text{ strongly in } L^2(\Omega)^{3 \times 3},$$

$$\varepsilon \mathbf{E}^\varepsilon u^\varepsilon \chi_M \rightarrow \left( \begin{array}{c|c} 0 & \frac{1}{2} \partial_3 z_\alpha \\ \hline \frac{1}{2} \partial_3 z_\alpha & \partial_3 z_3 \end{array} \right) \chi_M \text{ strongly in } L^2(\Omega)^{3 \times 3},$$

where the limit  $(u, v, z)$  is the unique solution of the problem

$$\left\{ \begin{array}{l} (u, v, z) \in \mathcal{U} \times \mathcal{V} \times \mathcal{Z}, \quad \forall (\bar{u}, \bar{v}, \bar{z}) \in \mathcal{U} \times \mathcal{V} \times \mathcal{Z}, \\ \int_{\Omega} \mathbb{A}(x) \left( \begin{array}{c|c} (\mathbf{E}u)_{\alpha\beta} & \frac{1}{2} \partial_3 v_\alpha \\ \hline \frac{1}{2} \partial_3 v_\alpha & \partial_3 v_3 \end{array} \right) \cdot \left( \begin{array}{c|c} (\mathbf{E}\bar{u})_{\alpha\beta} & \frac{1}{2} \partial_3 \bar{v}_\alpha \\ \hline \frac{1}{2} \partial_3 \bar{v}_\alpha & \partial_3 \bar{v}_3 \end{array} \right) \chi_F dx + \\ \int_{\Omega} \mathbb{A}(x) \left( \begin{array}{c|c} 0 & \frac{1}{2} \partial_3 z_\alpha \\ \hline \frac{1}{2} \partial_3 z_\alpha & \partial_3 z_3 \end{array} \right) \cdot \left( \begin{array}{c|c} 0 & \frac{1}{2} \partial_3 \bar{z}_\alpha \\ \hline \frac{1}{2} \partial_3 \bar{z}_\alpha & \partial_3 \bar{z}_3 \end{array} \right) \chi_M dx = \\ \int_{\Omega} (f_\alpha (\bar{u}_\alpha + \bar{z}_\alpha) + f_3 \bar{u}_3) dx. \end{array} \right. \quad (31)$$

**Proof.**

We first prove the convergence of the sequence of energies to the energy associated to the limit problem, assuming that the limit problem is the system (31). To that aim we will use the weak convergences proved in Lemma 3.1. We follow the argument already used in [12, Proof of Theorem 3.1].

Choosing  $(\bar{u}, \bar{v}, \bar{z}) = (u, v, z)$  in (31) and passing to the limit thanks to Lemma 3.1, we get

$$\begin{aligned} \lim_{\varepsilon \rightarrow 0} \int_{\Omega} (\chi_F + \varepsilon^2 \chi_M) \mathbb{A} \mathbf{E}^\varepsilon(u^\varepsilon) \cdot \mathbf{E}^\varepsilon(u^\varepsilon) dx &= \lim_{\varepsilon \rightarrow 0} \int_{\Omega} f u^\varepsilon dx \\ &= \int_{\Omega} \left( f_\alpha (u_\alpha + z_\alpha) + f_3 u_3 \right) dx \\ &= \int_{\Omega} (A E_F \cdot E_F \chi_F + A E_M \cdot E_M \chi_M) dx, \end{aligned}$$

where

$$E_F := \left( \begin{array}{c|c} (\mathbf{E}u)_{\alpha\beta} & \frac{1}{2} \partial_3 v_\alpha \\ \hline \frac{1}{2} \partial_3 v_\alpha & \partial_3 v_3 \end{array} \right), \quad E_M := \left( \begin{array}{c|c} 0 & \frac{1}{2} \partial_3 z_\alpha \\ \hline \frac{1}{2} \partial_3 z_\alpha & \partial_3 z_3 \end{array} \right).$$

Using the equivalence of the norms

$$L^2(\Omega) \ni E \longmapsto \int_{\Omega} A E \cdot E dx, \quad L^2(\Omega) \ni E \longmapsto \int_{\Omega} E \cdot E dx$$

we deduce that

$$\lim_{\varepsilon \rightarrow 0} \|(\chi_F + \varepsilon \chi_M) \mathbf{E}^\varepsilon(u^\varepsilon)\|_{L^2(\Omega)} = \|\chi_F E_F + \chi_M E_M\|_{L^2(\Omega)}.$$

Hence the strong convergence of the tensor sequences is proved. Now, by using the last strong convergence together with the following Korn's inequality

$$\|\mathbf{E}^\varepsilon(u^\varepsilon)\|_{L^2(F)} \geq \|\mathbf{E}(u^\varepsilon)\|_{L^2(F)} \geq C \|u^\varepsilon\|_{L^2(F)},$$

we deduce that  $u^\varepsilon$  is a Cauchy sequence in  $H^1(F)$ .

On the other hand, using (9) and (10) one has

$$\|\mathbf{E}^\varepsilon u^\varepsilon\|_{L^2(F)}^2 \geq C \varepsilon^2 \left( \|(R^\varepsilon)^{-1} u^\varepsilon\|_{L^2(F)}^2 + \|\nabla^\varepsilon u^\varepsilon\|_{L^2(F)}^2 \right)$$

and

$$\left( \|\nabla^\varepsilon u^\varepsilon\|_{L^2(F)}^2 + \|\mathbf{E}^\varepsilon u^\varepsilon\|_{L^2(M)}^2 \right) \geq C \|\nabla^\varepsilon u^\varepsilon\|_{L^2(M)}^2.$$

Hence,  $\varepsilon \nabla^\varepsilon u^\varepsilon$  is also a Cauchy sequence in  $L^2(\Omega)$ . It remains to prove that  $u_\alpha^\varepsilon$  and  $u_3^\varepsilon$  are two Cauchy sequences in  $L^2(\omega; H^1(\mathbb{I}))$  and  $H^1(M)$  respectively. To that aim, it suffices to use (18) to obtain

$$\|(\chi_F + \varepsilon \chi_M) \mathbf{E}^\varepsilon(u^\varepsilon)\|_{L^2(\Omega)}^2 = \mathcal{F}_\varepsilon(u^\varepsilon) \geq C \left( \|u_3^\varepsilon\|_{H^1(M)}^2 + \|u_\alpha^\varepsilon\|_{L^2(\omega; H^1(\mathbb{I}))}^2 \right)$$

then we can argue as previously to complete the proof.

We now prove that the limit problem is nothing but the system (31).

We choose a test function  $\phi^\varepsilon$  in (4) in the following form

$$\begin{cases} \phi_\alpha^\varepsilon = \bar{u}_\alpha + \varepsilon \bar{v}_\alpha + \bar{z}_\alpha \\ \phi_3^\varepsilon = \bar{u}_3 + \varepsilon^2 \bar{v}_3 + \varepsilon \bar{z}_3 \end{cases} \quad (32)$$

where  $(\bar{u}, \bar{v}, \bar{z}) \in \mathcal{U} \times \mathcal{D}(\omega; H_m^1(J))^3 \times \left\{ \mathcal{D}(\omega; H_m^1(I))^3 : z = 0, \text{ a.e. } x \in F \right\}$ . Since, from the definition of  $\mathcal{U}$ , one has  $(E\bar{u})_{i3} = 0$ , for all  $i = 1, 2, 3$ , an elementary calculation shows that

$$\begin{cases} (\mathbf{E}^\varepsilon \phi^\varepsilon)_{\alpha\beta} = (\mathbf{E}\bar{u})_{\alpha\beta} + \varepsilon(\mathbf{E}\bar{v})_{\alpha\beta} + (\mathbf{E}\bar{z})_{\alpha\beta}, \\ 2(\mathbf{E}^\varepsilon \phi^\varepsilon)_{\alpha 3} = \partial_3 \bar{v}_\alpha + \varepsilon \partial_\alpha \bar{v}_3 + \frac{1}{\varepsilon} \partial_3 \bar{z}_\alpha + \partial_\alpha \bar{z}_3, \\ (\mathbf{E}^\varepsilon \phi^\varepsilon)_{33} = (\mathbf{E}\bar{v})_{33} + \frac{1}{\varepsilon} (\mathbf{E}\bar{z})_{33}. \end{cases}$$

Using the following strong convergence

$$\begin{aligned} \mathbf{E}^\varepsilon \phi^\varepsilon \chi_F &\longrightarrow \left( \begin{array}{c|c} (\mathbf{E}\bar{u})_{\alpha\beta} & \frac{1}{2} \partial_3 \bar{v}_\alpha \\ \hline \frac{1}{2} \partial_3 \bar{v}_\alpha & \partial_3 \bar{v}_3 \end{array} \right) \chi_F \text{ in } L^2(\Omega)^{3 \times 3}, \\ \varepsilon \mathbf{E}^\varepsilon \phi^\varepsilon \chi_M &\longrightarrow \left( \begin{array}{c|c} 0 & \frac{1}{2} \partial_3 \bar{z}_\alpha \\ \hline \frac{1}{2} \partial_3 \bar{z}_\alpha & \partial_3 \bar{z}_3 \end{array} \right) \chi_M \text{ in } L^2(\Omega)^{3 \times 3}, \end{aligned}$$

we can pass to the limit in (4) as  $\varepsilon$  goes to zero to get (31) thanks to the density of  $\mathcal{D}(\omega; H_m^1(J))^3$  and  $\left\{ z \in \mathcal{D}(\omega; H_m^1(I))^3 : z_i = 0, \text{ a.e. } x \in F, i = 1, 2, 3 \right\}$  in  $\mathcal{V}$  and  $\mathcal{Z}$  respectively.

**Remark 3.3** *The space  $\mathcal{U} \times \mathcal{V} \times \mathcal{Z}$  is a Hilbert space for the following norm*

$$\|(u, v, z)\|^2 := \sum_{\alpha, \beta=1,2} \|(\mathbf{E}u)_{\alpha\beta}\|_{L^2(\Omega)}^2 + \sum_{i=1}^3 \left[ \|\partial_3 v_i\|_{L^2(\Omega)}^2 + \|\partial_3 z_i\|_{L^2(\Omega)}^2 \right].$$

*Therefore, using the Lax-Milgram Theorem we obtain the well-posedness of the limit problem (31) since the tensor  $A$  is coercive by virtue of (3) and  $f \in L^2(\Omega; \mathbb{R}^3)$ .*

We now turn back to the sequence of solutions of problem (6) to derive the following theorem from Theorem 3.2.

**Theorem 3.4** Let  $(u, v, z) \in \mathcal{U} \times \mathcal{V} \times \mathcal{Z}$  be the solution of the problem (31).

Let

$$\mathbf{E}_F := \left( \begin{array}{c|c} (\mathbf{E}u)_{\alpha\beta} & \frac{1}{2}\partial_3 v_\alpha \\ \hline \frac{1}{2}\partial_3 v_\alpha & \partial_3 v_3 \end{array} \right), \quad \mathbf{E}_M := \left( \begin{array}{c|c} 0 & \frac{1}{2}\partial_3 z_\alpha \\ \hline \frac{1}{2}\partial_3 z_\alpha & \partial_3 z_3 \end{array} \right).$$

Then the sequence of solutions  $\hat{u}^\varepsilon$  of the problem (6) fulfills the following convergences

$$\left\{ \begin{array}{l} \int_{\varepsilon J} \mathbf{E}\hat{u}^\varepsilon dx_3 \longrightarrow \int_J \mathbf{E}_F dx_3, \text{ in } L^2(\omega; \mathbb{R}^{3 \times 3}), \\ \int_{\varepsilon(I \setminus J)} \mathbf{E}\hat{u}^\varepsilon dx_3 \longrightarrow \int_{I \setminus J} \mathbf{E}_M dx_3 \text{ in } L^2(\omega; \mathbb{R}^{3 \times 3}), \\ \int_{\varepsilon I} \hat{u}_\alpha^\varepsilon dx_3 \longrightarrow U_\alpha(x') := \int_I (u_\alpha + z_\alpha) dx_3, \text{ in } L^2(\omega), \\ \int_{\varepsilon I} \varepsilon \hat{u}_3^\varepsilon dx_3 \longrightarrow u_3(x') \text{ in } L^2(\omega), \end{array} \right. \quad (33)$$

Moreover,  $U_\alpha$  may be written as

$$U_\alpha(x') = \int_I u_\alpha dx_3 + m_\alpha^0(x') \int_{I \setminus J} f_\alpha(x', x_3) dx_3 + m_\alpha^{00}(x'), \quad (34)$$

where  $m_\alpha^0$  and  $m_\alpha^{00}$  are given by

$$m_\alpha^0(x') := \int_{I \setminus J} z_\alpha^0(x', x_3) dx_3, \quad m_\alpha^{00}(x') := \int_{I \setminus J} z_\alpha^{00}(x', x_3) dx_3,$$

and  $z^0, z^{00}$  are respectively the solutions of the following problems

$$\left\{ \begin{array}{l} z^0 \in L^\infty(\omega, \mathcal{Z}^0), \mathcal{Z}^0 := \{z \in (H^1(I \setminus J))^3 : z = 0 \text{ on } \partial J\} \\ \int_{I \setminus J} A(x) \left( \begin{array}{c|c} 0 & \frac{1}{2}\partial_3 z_\alpha^0 \\ \hline \frac{1}{2}\partial_3 z_\alpha^0 & \partial_3 z_3^0 \end{array} \right) \left( \begin{array}{c|c} 0 & \frac{1}{2}\partial_3 \bar{z}_\alpha \\ \hline \frac{1}{2}\partial_3 \bar{z}_\alpha & \partial_3 \bar{z}_3 \end{array} \right) dx_3 = \int_{I \setminus J} \bar{z}_\alpha(x_3) dx_3, \\ \forall \bar{z} \in \mathcal{Z}^0, \text{ a.e. in } \omega, \end{array} \right. \quad (35)$$

$$\left\{ \begin{array}{l} z^{00} \in \mathcal{Z} \\ \int_M A(x) \left( \begin{array}{c|c} 0 & \frac{1}{2}\partial_3 z_\alpha^{00} \\ \hline \frac{1}{2}\partial_3 z_\alpha^{00} & \partial_3 z_3^{00} \end{array} \right) \left( \begin{array}{c|c} 0 & \frac{1}{2}\partial_3 \bar{z}_\alpha \\ \hline \frac{1}{2}\partial_3 \bar{z}_\alpha & \partial_3 \bar{z}_3 \end{array} \right) dx_3 = \\ \int_M \left( f_\alpha(x) - \int_{I \setminus J} f_\alpha(x) dx_3 \right) \bar{z}_\alpha(x) dx, \\ \forall \bar{z} \in \mathcal{Z}. \end{array} \right. \quad (36)$$



**Proof.**

The convergences (33) follow from the corresponding convergences stated in Theorem 3.2 by the change of variables (5).

Let us prove for instance the third convergence arising in (33) by the use of the first convergence arising in Theorem 3.2. Setting  $y_3 = \varepsilon x_3$  for  $x_3 \in I$  and bearing in mind that  $\hat{u}^\varepsilon$  is defined according to (5) by  $u_\alpha^\varepsilon(x', x_3) = \hat{u}^\varepsilon(x', \varepsilon x_3)$  for  $(x', x_3) \in \omega \times I$  and that the length of the interval  $I$  is equal to 1, we get thanks to the Cauchy-Schwarz inequality (with respect to  $x_3$ )

$$\left\{ \begin{array}{l} \int_\omega \left| \int_{\varepsilon I} \hat{u}_\alpha^\varepsilon(x', y_3) dy_3 - \int_I (u_\alpha(x', x_3) + z_\alpha(x', x_3)) dx_3 \right|^2 dx' = \\ \int_\omega \left| \int_I (u_\alpha^\varepsilon(x', x_3) - (u_\alpha(x', x_3) + z_\alpha(x', x_3))) dx_3 \right|^2 dx' \leq \\ \int_\omega \int_I |u_\alpha^\varepsilon(x', x_3) - (u_\alpha(x', x_3) + z_\alpha(x', x_3))|^2 dx_3 dx' \longrightarrow 0. \end{array} \right. \quad (37)$$

To prove (34), we first notice that one can derive the following equation satisfied by  $z_\alpha$  by choosing  $\bar{u} = \bar{v} = 0$  in (31),

$$\left\{ \begin{array}{l} z \in \mathcal{Z}, \quad \forall \bar{z} \in \mathcal{Z}, \\ \int_\Omega \mathbb{A}(x) \left( \begin{array}{c|c} 0 & \frac{1}{2} \partial_3 z_\alpha \\ \hline \frac{1}{2} \partial_3 z_\alpha & \partial_3 z_3 \end{array} \right) \cdot \left( \begin{array}{c|c} 0 & \frac{1}{2} \partial_3 \bar{z}_\alpha \\ \hline \frac{1}{2} \partial_3 \bar{z}_\alpha & \partial_3 \bar{z}_3 \end{array} \right) \chi_M dx = \\ \int_\Omega f_\alpha \bar{z}_\alpha dx. \end{array} \right. \quad (38)$$

Taking advantage from the linearity of (38), one can check easily by a uniqueness argument that  $z_\alpha$  may be identified as  $z_\alpha = z_\alpha^1 + z_\alpha^2$  where  $z_\alpha^1$  and  $z_\alpha^2$  are solutions of (38) but with right hand sides defined respectively by

$$\int_{I \setminus J} f_\alpha(x) dx_3 \quad \text{and} \quad \left( f_\alpha(x) - \int_{I \setminus J} f_\alpha(x) dx_3 \right).$$

By linearity and using once again a uniqueness argument, we conclude that

$$z_\alpha^1 = \left( \int_{I \setminus J} f_\alpha(x) dx_3 \right) z_\alpha^0 \quad \text{and} \quad z_\alpha^2 = z_\alpha^{00}$$

where  $z^0$  and  $z^{00}$  are the solutions of (35) and (36) respectively. Hence,  $U_\alpha$  defined in the third convergence (33) may be written as (34). This completes the proof of Theorem 3.4.

## References

- [1] G. ANZELLOTTI, S. BALDO, D. PERCIVALE, Dimension reduction in variational problems, asymptotic development in  $\Gamma$ -convergence and thin structures in elasticity, *Asymptotic Anal.*, **9** (1994), 61–100.
- [2] M. BELLIEUD & G. BOUCHITTÉ, Homogenization of a soft elastic material reinforced by fibers, *Asymptot. Anal.*, **32** (2002), no. 2, 153–183.
- [3] A. BOUGHAMMOURA, Homogenization and correctors for composite media with coated and highly anisotropic fibers, *Elect. J. Differential Equations*, (2012), no. 6, 1–27.
- [4] A. BRAIDES, M. BRIANE, & J. CASADO-DIAZ, Homogenization of non-uniformly bounded periodic diffusion energies in dimension two, *Nonlinearity*, **22** (2009), 1459–1480.
- [5] A. BRAIDES, V.-C. PIAT, & A. PIATNITSKI, A variational approach to double-porosity problems, *Asympt. Analysis*, **39** (2004), 281–308.
- [6] A. BRILLARD & M. EL JARROUDI, Asymptotic behaviour of a cylindrical elastic structure periodically reinforced along identical fibers, *IMA J. Appl. Math.*, **66** (2001), 567–590.
- [7] K. D. CHEREDNICHENKO, V. P. SMYSHLYAEV & V. V. ZHIKOV, Nonlocal limits for composite media with highly anisotropic periodic fibers, *Proceed. Royal. Soc. Edinburgh Sect. A*, **136** (2006), 87–144.
- [8] P.G. CIARLET & P. DESTUYNDER, A justification of the two-dimensional linear plate model, *J. Mécanique*, **18** (1979), 315–344.
- [9] A. GAUDIELLO & A. SILI, Homogenization of highly oscillating boundaries with strongly contrasting diffusivity, *SIAM J. Math. Anal.*, **47** (2015), no. 3, 1671–1692.

- [10] M. MABROUK & A. BOUGHAMMOURA, Homogénéisation d'un milieu élastique fortement hétérogène, *Comptes Rendus Mécanique*, **330**, 8 (2002), 543–548.
- [11] F. MURAT & A. SILI, Problèmes monotones dans des cylindres de faible diamètre formés de matériaux hétérogènes, *C.R. Acad. Sci. Paris Sér. I Math.*, **320** (1995), 1199–1204.
- [12] R. PARONI & A. SILI, Non-local effects by homogenization or 3D-1D dimension reduction in elastic materials reinforced by stiff fibers, *Journal of Differential Equations*, **260**, 3 (2016), 2026–2059.
- [13] A. SILI, Homogénéisation dans des cylindres minces, *C.R. Acad. Sci. Paris Sér. I Math.*, **332** (2001), 777–782.
- [14] A. SILI, Diffusion through a composite structure with a high contrasting diffusivity, *Asymptot. Anal.*, **89** (2014), no. 1-2, 173–187.