



HAL
open science

The subjective experience of recollection and familiarity in Alzheimer's disease

Dimitrios Kapogiannis, Mohamad El Haj

► **To cite this version:**

Dimitrios Kapogiannis, Mohamad El Haj. The subjective experience of recollection and familiarity in Alzheimer's disease. Behavioral and Brain Sciences, 2020, 42, pp.e290. 10.1017/S0140525X19001766 . hal-03662240

HAL Id: hal-03662240

<https://hal.science/hal-03662240>

Submitted on 19 May 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NoDerivatives 4.0 International License



Published in final edited form as:

Behav Brain Sci. ; 42: e290. doi:10.1017/S0140525X19001766.

The subjective experience of recollection and familiarity in Alzheimer's Disease.

Dimitrios KAPOGIANNIS^a, Mohamad EL HAJ^{b,c,d}

^aLaboratory of Neurosciences, National Institute on Aging, Baltimore, MD, USA

^bUniv Nantes, Laboratoire de psychologie des Pays de la Loire, LPPL, EA 4638, F-44000 Nantes, France

^cUnité de Gériatrie, Centre Hospitalier de Tourcoing, Tourcoing, France

^dInstitut Universitaire de France, Paris, France

Abstract

While the Integrative Memory model for Alzheimer's disease is interesting, it may benefit from incorporating the subjective experience of recollection. We therefore offer complementary lines of interpretation to explain how recollection and familiarity in Alzheimer's disease can be dissociated based not only on accounts of their neural correlates but, critically, on the subjective experience of memory in patients.

The Integrative Memory model, as proposed by Bastin et al., is theoretically and clinically relevant as it provides a framework for the chronological pattern of recollection and familiarity processes in Alzheimer's Disease depending on the brain regions most affected at each stage of the disease. That being said, the framework can be extended to include the subjective experience of both recollection and familiarity and their abnormalities in patients with Alzheimer's Disease. Recollection and familiarity have been associated with distinct subjective experiences; recollection is typically associated with a subjective experience of "mental time travel", in which we experience the conscious sensation of travelling back in time to relive the original event and to see it in our mind's eye, whereas familiarity is typically associated with a feeling of knowing characterized by a vague and unspecific experience of remembering (Tulving, 2002). Regarding Alzheimer's disease, patients typically demonstrate a shift from the ability to mentally relive past events (i.e., a shift from recollection) to a general sense of familiarity that may be expressed by the patients as a sense of "having experienced this before" (El Haj, Antoine, Nandrino, & Kapogiannis, 2015).

The decline of recollection in Alzheimer's disease can be attributed to decline in specific processes of the recollective experience. This decline has been pointed out by research demonstrating decline in the recollective experience during retrieval of past personal events in patients with Alzheimer's disease, including decline in subjective processes, such as

reliving, travel in time, remembering, realness, rehearsal, and visual imagery (El Haj, Kapogiannis, & Antoine, 2016). Among these subjective processes, decline of visual imagery seems to play a key role in the decline of recollective experience in patients with Alzheimer's disease. The decline of visual imagery in Alzheimer's disease seems to deprive patients from the ability to retrieve and manipulate mental images during retrieval, and also deprive them from visual cues that mediate and/or accelerate their search through memory stores (El Haj, Gallouj, & Antoine, 2019; El Haj, Moustafa, Gallouj, & Robin, 2019).

The relationship between decline of visual imagery and decline of the recollective experience in Alzheimer's disease can be understood by highlighting research using the Field/Observer paradigm, which has been widely used to assess the subjective experience of recollection and familiarity in general populations (Nigro & Neisser, 1983; Rice & Rubin, 2011). In one study in Alzheimer's disease, patients were invited to retrieve past personal events and, after a delay, provide a "Field" response, if they can visualize the event through their own eyes, or an "Observer" response if they can visualize themselves in the scene as a spectator would (El Haj, Moustafa, et al., 2019). Results demonstrated increased "Field" and decreased "Observer" responses in patients, suggesting a diminished ability of patients with Alzheimer's disease to construct vivid images when recollecting the past.

In summary, the decline of the recollective experience during retrieval can be associated with declines in several components of the subjective experience, such as reliving, travel in time, remembering, realness, and rehearsal. Critically, decline of visual imagery seems to deprive patients from the ability to construct mental images when recollecting the past. We propose that these processes should be taken into account by The Integrative Memory model, as proposed by Bastin et al., to better account for recollection deficits in patients with Alzheimer's disease. We believe that a decline in the subjective experience during retrieval in patients with Alzheimer's disease leads to a decline of recollection and, consequently, to the emergence of a general sense of familiarity that is typically associated with a sense of "having experienced this before". As pointed out by Bastin et al., the Integrative Memory model is not comprehensive and should evolve to incorporate other mechanisms. Therefore, we propose that the Integrative Memory model by Bastin et al is sufficiently flexible to include these subjective processes and may be enriched by this inclusion.

Acknowledgments

This research was supported in part (DK) by the Intramural Research Program of the National Institute on Aging, NIH.

References

- El Haj M, Antoine P, Nandrino JL, & Kapogiannis D (2015). Autobiographical memory decline in Alzheimer's disease, a theoretical and clinical overview. *Ageing Res Rev*, 23(Pt B), 183–192. doi: 10.1016/j.arr.2015.07.001 [PubMed: 26169474]
- El Haj M, Gallouj K, & Antoine P (2019). Mental imagery and autobiographical memory in Alzheimer's disease. *Neuropsychology*.
- El Haj M, Kapogiannis D, & Antoine P (2016). Phenomenological Reliving and Visual Imagery During Autobiographical Recall in Alzheimer's Disease. *Journal of Alzheimer's Disease*, 52(2), 421–431. doi: 10.3233/JAD-151122

- El Haj M, Moustafa AA, Gallouj K, & Robin F (2019). Visual imagery: The past and future as seen by patients with Alzheimer's disease. *Consciousness and Cognition*, 68, 12–22. doi: 10.1016/j.concog.2018.12.003 [PubMed: 30593998]
- Nigro G, & Neisser U (1983). Point of view in personal memories. *Cognitive Psychology*, 15(4), 467–482. doi: 10.1016/0010-0285(83)90016-6
- Rice HJ, & Rubin DC (2011). Remembering from any angle: the flexibility of visual perspective during retrieval. *Consciousness and Cognition*, 20(3), 568–577. doi: 10.1016/j.concog.2010.10.013 [PubMed: 21109466]
- Tulving E (2002). Episodic memory: from mind to brain. *Annual Review of Psychology*, 53, 1–25. doi: 10.1146/annurev.psych.53.100901.135114