



**HAL**  
open science

# Semiclassical Weyl Formula for Elliptic Operators with Non-smooth Coefficients

Lech Zielinski

► **To cite this version:**

Lech Zielinski. Semiclassical Weyl Formula for Elliptic Operators with Non-smooth Coefficients. 2022.  
hal-03662996

**HAL Id: hal-03662996**

**<https://hal.science/hal-03662996v1>**

Preprint submitted on 9 May 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Semiclassical Weyl Formula for Elliptic Operators with Non-smooth Coefficients

by  
Lech Zielinski

LMPA, Centre Mi-Voix, Université du Littoral,  
B.P. 699, 62228 Calais Cedex, France  
e-mail: Lech.Zielinski@lmpa.univ-littoral.fr

**Abstract.** We consider the Weyl formula for the asymptotic number of eigenvalues of self-adjoint elliptic differential operators with coefficients which have Hölder continuous first order derivatives. Our aim is to prove that the Weyl formula holds with a remainder usually considered in the case of operators with smooth coefficients.

**Mathematics Subject Classification (2000):** 35P20.

**Key words:** spectral asymptotics, semiclassical approximation, elliptic operator, Weyl formula.

## 1 Introduction

This paper presents a generalization of our earlier results described in [14], [16], [17] and motivated by the well-known result of L. Hörmander [5] on spectral asymptotics of a self-adjoint elliptic operator  $A$  with smooth coefficients on a smooth (boundaryless) manifold  $M$ . If  $M$  is a compact manifold of dimension  $d$  and  $2m$  is the order of  $A$ , then the associated counting function  $\mathcal{N}(A, \lambda)$ , defined as the number of eigenvalues (counted with their multiplicities) smaller than  $\lambda$ , satisfies the Weyl formula

$$\mathcal{N}(A, \lambda) = c\lambda^{\frac{d}{2m}} + O(\lambda^{\frac{d-\mu}{2m}}), \quad (1.1)$$

where  $c = c_{A,M}$  is a constant and  $\mu = 1$ .

It is natural to ask if (1.1) still holds for the operators with non-smooth coefficients. In [16] we proved that for coefficients which are Hölder continuous of exponent  $r \in ]0; 1]$ , the formula (1.1) holds if  $\mu < r$ . In [17] we proved (1.1) with  $\mu = 1$  assuming that the first order derivatives of coefficients are Lipschitz continuous and in this paper we show a way of replacing the hypothesis of Lipschitz continuity by Hölder continuity. Moreover we use the semiclassical framework, which allows us to obtain the classical Weyl formula (1.1) as a corollary.

Let  $\mathcal{B}^r$  denote the set of bounded, Hölder continuous functions on  $\mathbb{R}^d$ , i.e.  $a \in \mathcal{B}^r$  means that  $a \in L^\infty(\mathbb{R}^d)$  and there is  $C > 0$  such that

$$|a(x) - a(y)| \leq C|x - y|^r \quad (x, y \in \mathbb{R}^d).$$

Further on  $m \in \mathbb{N} \setminus \{0\}$  and we fix  $r \in ]0; 1]$ . For  $\nu, \bar{\nu} \in \mathbb{N}^d$ ,  $|\nu|, |\bar{\nu}| \leq m$ , let  $a_{\nu, \bar{\nu}}$  be real valued coefficients satisfying  $a_{\nu, \bar{\nu}} = a_{\bar{\nu}, \nu}$  and

$$\partial_{x_j} a_{\nu, \bar{\nu}} \in \mathcal{B}^r \quad (j = 1, \dots, d). \quad (1.2)$$

We assume moreover the existence of a constant  $c > 0$  such that

$$\sum_{|\nu|=|\bar{\nu}|=m} a_{\nu, \bar{\nu}}(x) \xi^{\nu+\bar{\nu}} \geq c|\xi|^{2m} \quad (x, \xi \in \mathbb{R}^d) \quad (1.3)$$

and for  $h > 0$  we introduce the quadratic form  $\mathcal{A}_h$  defined for  $\varphi, \psi \in C_0^m(\mathbb{R}^d)$  by the formula

$$\mathcal{A}_h[\varphi, \psi] = \sum_{|\nu|, |\bar{\nu}| \leq m} (a_{\nu, \bar{\nu}}(hD)^\nu \varphi, (hD)^{\bar{\nu}} \psi), \quad (1.4)$$

where  $(\cdot, \cdot)$  is the scalar product of  $L^2(\mathbb{R}^d)$  and  $(hD)^\nu = (-ih)^{|\nu|} \partial^\nu / \partial x^\nu$ .

The ellipticity hypothesis (1.3) ensures the fact that  $\mathcal{A}_h$  is bounded from below and its closure defines a self-adjoint operator  $A_h$ . We introduce

$$a(x, \xi) = \sum_{|\nu|, |\bar{\nu}| \leq m} a_{\nu, \bar{\nu}}(x) \xi^{\nu+\bar{\nu}} \quad (1.5)$$

and for  $E \in \mathbb{R}$  we denote

$$\Gamma_E = a^{-1}(] - \infty; E]) = \{(x, \xi) \in \mathbb{R}^{2d} : a(x, \xi) < E\}. \quad (1.6)$$

We will present a proof of

**Theorem 1.1** *Let  $E_0, E \in \mathbb{R}$  be such that  $E_0 < E$  and  $\Gamma_E$  is bounded. Then the operators  $A_h$  described above have discrete spectrum in  $] -\infty; E_0]$  for  $h \in ]0; h_0]$ , if  $h_0 > 0$  is fixed small enough. We assume moreover*

$$a(x, \xi) = E_0 \Rightarrow \nabla_\xi a(x, \xi) \neq 0. \quad (1.7)$$

*Then, for  $h \in ]0; h_0]$ , the counting function  $\mathcal{N}(A_h, E_0)$  satisfies the asymptotic formula*

$$\mathcal{N}(A_h, E_0) = |\Gamma_{E_0}|(2\pi h)^{-d} + O(h^{1-d}), \quad (1.8)$$

*where  $|\Gamma_{E_0}|$  denotes the Lebesgue measure of  $\Gamma_{E_0}$ .*

## 2 Regularization of coefficients

Let  $\gamma_1 \in C_0^\infty(\mathbb{R}^d)$  be such that  $\gamma_1(-x) = \gamma_1(x)$ ,  $\int \gamma_1(x) dx = 1$  and for  $\varepsilon > 0$  set  $\gamma_\varepsilon(x) = \varepsilon^{-d} \gamma_1(x/\varepsilon)$ . We fix  $\delta \in ]0; 1[$  and introduce

$$a_{\nu, \bar{\nu}, h}(x) = (a_{\nu, \bar{\nu}} * \gamma_{h^\delta})(x) = \int a_{\nu, \bar{\nu}}(y) \gamma_1(h^{-\delta}(x - y)) h^{-\delta d} dy. \quad (2.1)$$

It is easy to see (cf. Section 7(A) ) that (1.2) ensures

$$|a_{\nu, \bar{\nu}}(x) - a_{\nu, \bar{\nu}, h}(x)| \leq Ch^{\delta(1+r)}, \quad (2.2)$$

$$|\partial_x^\alpha a_{\nu, \bar{\nu}, h}(x)| \leq C_\alpha h^{-\delta(|\alpha| - 1 - r)_+}, \quad (2.3)$$

where  $s_+ = (s + |s|)/2$  is the positive part of  $s \in \mathbb{R}$ . We define

$$p_h(x, \xi) = \sum_{|\nu|, |\bar{\nu}| \leq m} a_{\nu, \bar{\nu}, h}(x) \xi^{\nu + \bar{\nu}} \quad (2.4)$$

and assume further on  $(1 + r)\delta \geq 1$ , hence (2.2) ensures

$$|\partial_\xi^\alpha (a - p_h)(x, \xi)| \leq C_\alpha h \langle \xi \rangle^{2m - |\alpha|}, \quad (2.5)$$

where  $\langle \xi \rangle = (1 + |\xi|^2)^{1/2}$ . Moreover the operator

$$P_h = \sum_{|\nu|, |\bar{\nu}| \leq m} (hD)^\nu a_{\nu, \bar{\nu}, h}(x) (hD)^{\bar{\nu}} \quad (2.6)$$

is an approximation of  $A_h$  satisfying

$$\pm(A_h - P_h) \leq Ch\langle hD \rangle^{2m} = Ch(I - h^2\Delta)^m$$

and  $P_h^- \leq A_h \leq P_h^+$  holds (in the sens of quadratic forms) with

$$P_h^\pm = P_h \pm Ch\langle hD \rangle^{2m}. \quad (2.7)$$

Our principal aim is to show

**Theorem 2.1** *The assertion of Theorem 1.1 holds if  $A_h$  is replaced by  $P_h^\pm$ .*

First of all we remark that the discreteness of the spectrum of  $P_h^\pm$  is a standard result and analogical reasoning can be applied to  $A_h$  (cf. Section 7(B) for details). Then the asymptotic formula (1.8) follows from the estimates of the counting functions for  $P_h^\pm$  due to the min-max principle (cf. [12]), which ensures

$$P_h^- \leq A_h \leq P_h^+ \Rightarrow \mathcal{N}(P_h^+, E) \leq \mathcal{N}(A_h, E) \leq \mathcal{N}(P_h^-, E). \quad (2.8)$$

In the remaining part of this section we describe a version of Theorem 2.1 formulated suitably for the proof that will follow.

Let  $\mathbf{1}_Z : \mathbb{R} \rightarrow \{0, 1\}$  be the characteristic function of  $Z \subset \mathbb{R}$  and set

$$\mathcal{N}_h(p_h, E) = \int_{p_h(x, \xi) \leq E} \frac{dx d\xi}{(2\pi h)^d} = \int \frac{dx d\xi}{(2\pi h)^d} \mathbf{1}_{]-\infty; E]} \circ p_h. \quad (2.9)$$

Further on we also write  $\mathbf{1}_Z(P_h^\pm)$  to denote the spectral projector of  $P_h^\pm$  on a Borel set  $Z \subset \mathbb{R}$  and due to (2.8) and (2.11), to obtain the asymptotic formula (1.8) it suffices to show

$$\mathrm{tr} \mathbf{1}_{]-\infty; E_0]}(P_h^\pm) = \mathcal{N}(P_h^\pm, E_0) = \mathcal{N}_h(p_h, E_0) + O(h^{1-d}). \quad (2.15)$$

To show (2.15) we introduce

$$\begin{cases} g \in C_0^\infty(]E_0 - 2c_0; E_0 + 2c_0[), & g = 1 \text{ on } [E_0 - c_0; E_0 + c_0], \\ \tilde{g} \in C_0^\infty(]-\infty; E_0 - c_0]) \end{cases} \quad (2.16)$$

and assume  $\tilde{g} + g^2 = 1$  on  $[-E'_0; E_0 + c_0]$  where  $E'_0 > 0$  is chosen sufficiently large to ensure  $-E'_0 \leq \inf p_h^\pm$  and  $-E'_0 \leq \inf \sigma(P_h^\pm)$  for  $h \in ]0; h_0]$ .

Then for  $E \in [E_0 - c_0; E_0 + c_0]$  we can write the decompositions

$$\mathcal{N}_h(p_h, E) = \int \frac{dx d\xi}{(2\pi h)^d} (\tilde{g} + g^2 \mathbf{1}_Z)(p_h(x, \xi)),$$

$$\mathbf{1}_{]-\infty; E]}(P_h^\pm) = (\tilde{g} + g^2 \mathbf{1}_Z)(P_h^\pm),$$

where  $Z = [E_0 - 2c_0; E]$ . Now it is clear that (2.15) is a consequence of

**Theorem 2.2** *One has*

$$\mathrm{tr} \tilde{g}(P_h^\pm) = \int \frac{dx d\xi}{(2\pi h)^d} \tilde{g} \circ p_h + O(h^{1-d}), \quad (2.17)$$

$$\mathrm{tr} (g^2 \mathbf{I}_Z)(P_h^\pm) = \int \frac{dx d\xi}{(2\pi h)^d} (g^2 \mathbf{I}_Z) \circ p_h + O(h^{1-d}). \quad (2.18)$$

Further on  $\bar{c} > 0$  is small enough and  $f_0 \in C_0^\infty([\bar{c}/2; \bar{c}/2])$ . Then we define  $f_1 = f_0 * f_0 \in C_0^\infty([\bar{c}; \bar{c}])$  and

$$\tilde{f}_h(\lambda) = \mathcal{F}_h^{-1} f_1(\lambda) = \int_{-\infty}^{\infty} e^{i\lambda t/h} f_1(t) \frac{dt}{2\pi h}. \quad (2.19)$$

We note that  $\tilde{f}_h(\lambda) = \tilde{f}_1(\lambda/h)/h$ ,  $\tilde{f}_1$  is rapidly decreasing and  $\tilde{f}_1 > 0$ . Moreover we can choose  $f_0$  such that  $\int_{-\infty}^{\infty} \tilde{f}_h = f_1(0) = 1$ . Then we use

$$\tilde{f}_h^Z(\lambda) = (\mathbf{I}_Z * \tilde{f}_h)(\lambda) = \int_Z d\lambda' \tilde{f}_h(\lambda - \lambda') \quad (2.20)$$

as an approximation of  $\mathbf{I}_Z$ . In Section 7(C) we recall a proof of

**Proposition 2.3** *The estimate (2.18) follows if*

$$\mathrm{tr} (g^2 \tilde{f}_h^Z)(P_h^\pm) = \int \frac{dx d\xi}{(2\pi h)^d} (g^2 \tilde{f}_h^Z) \circ p_h + O(h^{1-d}) \quad (2.21)$$

holds uniformly with respect to  $Z = [E'; E]$ .

### 3 Pseudodifferential approximation

In this section we show the estimate (2.17). We use pseudodifferential operators in a reasoning which will be developed in Section 4 to treat (2.18). We denote by  $\|\cdot\|$  the norm of bounded operators  $B(L^2(\mathbb{R}^d))$  and  $\|B\|_{\mathrm{tr}} = \mathrm{tr} (B^* B)^{1/2}$  is the trace class norm. If  $b_h \in C_0^\infty(\mathbb{R}^{2d})$  then we denote by  $b_h(x, hD)$  the pseudodifferential operator acting on  $\varphi \in \mathcal{S}(\mathbb{R}^d)$  according to the formula

$$(b_h(x, hD)\varphi)(x) = \int \frac{dy d\xi}{(2\pi h)^d} e^{i(x-y)\xi/h} b_h(x, \xi) \varphi(y). \quad (3.1)$$

If  $b_h, l_h \in C_0^\infty(\mathbb{R}^{2d})$ , then  $b_h(x, hD)l_h(x, hD)^*$  has the integral kernel

$$\mathcal{K}_h(x, y) = \int \frac{d\xi}{(2\pi h)^d} e^{i(x-y)\xi/h} b_h(x, \xi) \overline{l_h(y, \xi)},$$

hence

$$\mathrm{tr} b_h(x, hD) l_h(x, hD)^* = \int dx \mathcal{K}_h(x, x) = \int \frac{dx d\xi}{(2\pi h)^d} b_h \bar{l}_h. \quad (3.2)$$

In this section we consider  $\tilde{g} \in C_0^\infty(\cdot - \infty; E_1)$  with  $E_1$  such that  $\Gamma_{E_1+2c_1}$  is bounded for a certain  $c_1 > 0$ . Then using (2.5) we can find  $h_0 > 0$  small enough to ensure

$$0 < h < h_0 \Rightarrow \mathrm{supp}(\tilde{g} \circ p_h) \subset \Gamma_{E_1+c_1}. \quad (3.3)$$

Next we fix an auxiliary  $h$ -independent cut-off function  $l \in C_0^\infty(\mathbb{R}^{2d})$  such that  $l = 1$  on a neighbourhood of  $\overline{\Gamma_{E_1+c_1}}$  and  $l \geq 0$ . We denote  $L_h = l(x, hD)$  and recall standard estimates of pseudodifferential operators:

$$\|L_h\| = O(1), \quad \|L_h\|_{\mathrm{tr}} = O(h^{-d}). \quad (3.4)$$

Moreover (3.2) allows us to write

$$\int \frac{dy d\xi}{(2\pi h)^d} \tilde{g} \circ p_h = \int \frac{dy d\xi}{(2\pi h)^d} l(\tilde{g} \circ p_h) = \mathrm{tr}(\tilde{g} \circ p_h)(x, hD) L_h^* \quad (3.5)$$

and since  $|\mathrm{tr} B| \leq \|B\|_{\mathrm{tr}}$ , it is clear that (2.17) follows from

$$\|\tilde{g}(P_h^\pm) - L_h^*(\tilde{g} \circ p_h)(x, hD)\|_{\mathrm{tr}} = O(h^{1-d}). \quad (3.6)$$

Our proof of (3.6) is based on an auxiliary lemma (cf. Section 7(D)):

**Lemma 3.1** *If  $\tilde{g}$  and  $L_h$  are as above, then for every  $N \in \mathbb{N}$  one has*

$$\|\tilde{g}(P_h^\pm)(I - L_h)\|_{\mathrm{tr}} = O(h^N). \quad (3.7)$$

Since  $\|\tilde{g}(P_h^\pm)L_h\|_{\mathrm{tr}} \leq \|\tilde{g}(P_h^\pm)\| \|L_h\|_{\mathrm{tr}} = O(h^{-d})$  due to (3.4), Lemma 3.1 ensures  $\|\tilde{g}(P_h^\pm)\|_{\mathrm{tr}} = O(h^{-d})$ . Another immediate corollary of (3.7) is

$$\|\tilde{g}(P_h^\pm) - L_h^* \tilde{g}(P_h^\pm) L_h\|_{\mathrm{tr}} = O(h^N)$$

and we can conclude that instead of (3.6) it suffices to show

$$\|L_h^* \tilde{g}(P_h^\pm) L_h - L_h^*(\tilde{g} \circ p_h)(x, hD)\|_{\mathrm{tr}} = O(h^{1-d}). \quad (3.6')$$

Further on  $g_1 \in \mathcal{S}(\mathbb{R})$  denotes the Fourier transform of  $\tilde{g}$ , i.e. we have  $\tilde{g}(\lambda) = \int_{-\infty}^{\infty} \frac{dt}{2\pi} g_1(t) e^{it\lambda}$  and

$$(\tilde{g} \circ p_h)(x, \xi) = \tilde{g}(p_h(x, \xi)) l(x, \xi) = \int_{-\infty}^{\infty} \frac{dt}{2\pi} g_1(t) (e^{itp_h} l)(x, \xi), \quad (3.8)$$

$$\tilde{g}(P_h^\pm) = \int_{-\infty}^{\infty} \frac{dt}{2\pi} g_1(t) e^{itP_h^\pm}. \quad (3.9)$$

Therefore we can write

$$\tilde{g}(P_h^\pm)L_h - (\tilde{g} \circ p_h)(x, hD) = \int_{-\infty}^{\infty} \frac{dt}{2\pi} g_1(t) R_{h,t}$$

with

$$R_{h,t} = e^{itP_h^\pm} L_h - (e^{itp_h l})(x, hD) \quad (3.10)$$

and the left hand side of (3.6') can be estimated by

$$\int_{-\infty}^{\infty} \frac{dt}{2\pi} |g_1(t)| \|L_h^* R_{h,t}\|_{\text{tr}}. \quad (3.11)$$

However  $\|L_h^* R_{h,t}\|_{\text{tr}} \leq \|L_h^*\|_{\text{tr}} \|R_{h,t}\| \leq Ch^{-d} \|R_{h,t}\|$  and  $g_1$  is rapidly decaying, hence in order to estimate the quantity (3.11) by  $O(h^{1-d})$  it suffices to show

**Lemma 3.2** *There is  $C > 0$  such that  $\|R_{h,t}\| \leq Ch\langle t \rangle^C$ .*

Before starting the proof of Lemma 3.2 it is useful to introduce some notations. We write the differential operators  $P_h^\pm$  in the standard form

$$P_h^\pm = \sum_{|\alpha| \leq 2m} a_{\nu, \bar{\nu}, h}^\pm(x) (hD)^\alpha = p_h^\pm(x, hD), \quad (3.12)$$

$$p_h^\pm(x, \xi) = \sum_{|\alpha| \leq 2m} a_{\nu, \bar{\nu}, h}^\pm(x) \xi^\alpha \quad (3.13)$$

and it is easy to check that (2.5) still holds with  $p_h^\pm$  instead of  $p_h$ .

Let  $\tilde{m} \in \mathbb{R}$  and  $\delta_1 \geq \delta_0 \geq 0$  be such that  $\delta_0 + \delta_1 < 1$ . If  $b = \{b_h\}_{0 < h \leq h_0}$  is a family of smooth functions satisfying the estimates

$$|\partial_\xi^\alpha \partial_x^\beta b_h(x, \xi)| \leq C_{n, \alpha, \beta} h^{-\tilde{m} - |\alpha|\delta_0 - |\beta|\delta_1} \langle \xi \rangle^{-n} \quad (3.14)$$

(for every  $\alpha, \beta \in \mathbb{N}^d$  and  $n \in \mathbb{N}$ ), then we write  $b \in S_{\delta_0, \delta_1}^{\tilde{m}}$ .

If  $l \in C_0^\infty(\mathbb{R}^{2d})$  is  $h$ -independent, then the families  $lp_h$ ,  $l\tilde{p}_h^\pm$ ,  $le^{itp_h}$  belong to  $S_{0, \delta}^0$  and well-known  $L^2$ -estimates of pseudodifferential operators (cf. [2],[6]) give

$$b \in S_{0, \delta}^{\tilde{m}} \Rightarrow \exists C > 0, \quad \|(e^{itp_h} lb_h)(x, hD)\| \leq Ch^{-\tilde{m}} \langle t \rangle^C. \quad (3.15)$$



If  $s \in \mathbb{R}$  then we write  $B_h = L_h + O(h^s)$  if and only if there is a constant  $C > 0$  such that  $\|B_h - L_h\| \leq Ch^s$  holds for all  $h \in ]0; h_0]$ . Moreover further on we adopt the convention to drop the index  $h$  whenever there is no confusion. Instead of  $P_h^\pm, L_h, R_{h,t}, p_h, p_h^\pm$ , we are going to write simply  $P, L, R_t, p, p^\pm$ .

**Proof of Lemma 3.2** We are going to obtain suitable estimates of

$$\tilde{R}_t = (i\partial_t + P)(e^{itp}l)(x, hD) \quad (3.16)$$

and the assertion of Lemma 3.2 will follow from

$$R_t = -[e^{i(t-\tau)P}(e^{i\tau p}l)(x, hD)]_{\tau=0}^{\tau=t} = i \int_0^t d\tau e^{i(t-\tau)P} \tilde{R}_\tau. \quad (3.17)$$

The standard formula of the composition with a differential operator gives

$$p^\pm(x, hD)(e^{itp}l)(x, hD) = \tilde{q}_t(x, hD), \quad (3.18)$$

with

$$\tilde{q}_t = \sum_{|\alpha| \leq 2m} \frac{(-ih)^{|\alpha|}}{\alpha!} \partial_x^\alpha (e^{itp}l) \partial_\xi^\alpha p^\pm = e^{itp} \sum_{k=0}^{2m} t^k q_k, \quad (3.19)$$

$$q_k = \sum_{k \leq |\alpha| \leq 2m} \sum_{\substack{\alpha_0 + \dots + \alpha_k = \alpha \\ \alpha_j \neq 0 \text{ if } j \neq 0}} c_{\alpha_0, \dots, \alpha_k} \partial_x^{\alpha_0} l \partial_x^{\alpha_1} p \dots \partial_x^{\alpha_k} p \partial_\xi^\alpha p^\pm \quad (3.20)$$

and  $q_k \in S_{0,\delta}^{-k}$  follows from  $-|\alpha| + \sum_{j=1}^k \delta(|\alpha_j| - 1) \leq -k - (1 - \delta)(|\alpha| - k)$ . However (3.15) ensures

$$q_k \in S_{0,\delta}^{-k} \Rightarrow (e^{itp}q_k)(x, hD) = O(h^k \langle t \rangle^{C_k}),$$

hence

$$P(e^{itp}l)(x, hD) = (e^{itp}q_0)(x, hD) + O(h \langle t \rangle^C).$$

To complete the proof we note that  $l(p^\pm - p) \in S_{0,\delta}^{-1}$  and

$$q_0 - lp^\pm = \sum_{1 \leq |\alpha| \leq 2m} \frac{(-ih)^{|\alpha|}}{\alpha!} \partial_x^\alpha l \partial_\xi^\alpha p^\pm \in S_{0,\delta}^{-1} \quad (3.21)$$

ensure  $q_0 - lp \in S_{0,\delta}^{-1}$ , hence

$$i\partial_t(e^{itp}l)(x, hD) + (e^{itp}q_0)(x, hD) = (e^{itp}(q_0 - lp))(x, hD) = O(h \langle t \rangle^C).$$

## 4 Refinement

Due to Proposition 2.3, to complete the proof of Theorem 2.2 it suffices to show (2.21). We will use a reasoning of Section 3 in a refined form. We keep notations introduced before and in particular we drop the index  $h$ .

To begin we remark that replacing  $\tilde{g}$  by  $g \in C_0^\infty(]E_0 - c_0; E_0 + c_0[)$  in (3.6), we can introduce  $l_0 = g \circ p$ ,  $L_0 = l_0(x, hD)$  and write

$$\mathrm{tr}(g^2 \tilde{f}_h^Z)(P) = \mathrm{tr} L_0^* \tilde{f}_h^Z(P) L_0 + O(h^{1-d}). \quad (4.1)$$

As before  $Z = [E'; ]$  and we use the Fourier transform

$$\begin{aligned} f_h^Z(t) &:= \mathcal{F}_h \tilde{f}_h^Z(t) = \int_{-\infty}^{\infty} d\lambda e^{-it\lambda/h} \tilde{f}_h^Z(\lambda) \\ &= f_1(t) \int_{E'}^E d\lambda e^{-it\lambda/h} = f_1(t) \frac{e^{-it(E+E')/(2h)}}{t/(2h)} \sin \frac{(E'-E)t}{2h} \end{aligned}$$

to write

$$\mathrm{tr} L_0^* \tilde{f}_h^Z(P) L_0 = \int_{-\infty}^{\infty} \frac{dt}{2\pi h} f_h^Z(t) \mathrm{tr} L_0^* e^{itP/h} L_0, \quad (4.2)$$

$$\int \frac{dx d\xi}{(2\pi h)^d} l_0^2(\tilde{f}_h^Z \circ p) = \int \frac{dx d\xi}{(2\pi h)^d} \int_{-\infty}^{\infty} \frac{dt}{2\pi h} f_h^Z(t) l_0^2 e^{itp/h}. \quad (4.3)$$

However (3.2) allows us to express (4.3) as

$$\int_{-\infty}^{\infty} \frac{dt}{2\pi h} f_h^Z(t) \mathrm{tr} L_0^*(l_0 e^{itp/h})(x, hD) \quad (4.3')$$

and introducing

$$R_{t/h}^0 = e^{itP/h} L_0 - (e^{itp/h} l_0)(x, hD) \quad (4.4)$$

we find that (2.21) is equivalent to the estimate

$$\int_{-\infty}^{\infty} \frac{dt}{2\pi h} f_h^Z(t) \mathrm{tr} L_0^* R_{t/h}^0 = O(h^{1-d}). \quad (4.5)$$

Next we observe that (2.5) and (2.10) ensure

$$E_0 - 2c_0 \leq p(x, \xi) \leq E_0 + 2c_0 \Rightarrow |\nabla_\xi p(x, \xi)| \geq c_0 \quad (4.6)$$

for  $h \in ]0; h_0[$  if  $h_0 > 0$  is fixed small enough. The condition (4.6) ensures

**Lemma 4.1** *Assume that  $b \in S_{0,\delta}^0$ . Then for every  $n \in \mathbb{N}$  one can find  $b_n \in S_{0,\delta}^0$  such that*

$$t^n \int d\xi b e^{itp/h} = h^n \int d\xi b_n e^{itp/h}. \quad (4.7(n))$$

**Proof.** Using (4.6) in a standard reasoning we obtain

$$\forall b \in S_{0,\delta}^0 \quad \exists \tilde{b}_j \in S_{0,\delta}^0, \quad b = \sum_{j=1}^d \tilde{b}_j \partial_{\xi_j} p. \quad (4.8)$$

Therefore performing the integration by parts

$$t \int d\xi \sum_{j=1}^d b_j \partial_{\xi_j} p e^{itp/h} = ih \int d\xi \sum_{j=1}^d \partial_{\xi_j} \tilde{b}_j e^{itp/h}$$

we obtain the assertion of Lemma 4.1 for  $n = 1$  with  $b_1 = \sum_{j=1}^d i \partial_{\xi_j} \tilde{b}_j$ . Obviously the general case follows by induction with respect to  $n$ .  $\triangle$ .

We will use Lemma 4.1 to replace  $R_{t/h}^0$  in (4.5) by

$$R_{t/h} = e^{itP/h} L_0 - (e^{itp/h} l_t)(x, hD), \quad (4.4')$$

where  $l_t = l_0 + tl'_0 + \frac{1}{2}t^2 l''_0$  with certain  $l'_0 \in S_{0,\delta}^0$  and  $l''_0 \in S_{0,\delta}^1$ . In fact  $\text{tr} L_0^*(R_{t/h} - R_{t/h}^0)$  can be expressed as

$$\int \frac{dx d\xi}{(2\pi h)^d} l_0 (l_t - l_0) e^{itp/h} = h^{1-d} \int dx d\xi \tilde{l} e^{itp/h},$$

with a certain  $\tilde{l} \in S_{0,\delta}^0$  (due to Lemma 4.1 with  $n = 1$  and 2). Thus

$$\int_{-\infty}^{\infty} \frac{dt}{2\pi h} f_h^Z(t) \text{tr} L_0^*(R_{t/h} - R_{t/h}^0) = h^{1-d} \int dx d\xi \tilde{l} (f_h^Z \circ p)$$

is clearly  $O(h^{1-d})$ , hence (4.5) is equivalent to

$$\int_{-\infty}^{\infty} \frac{dt}{2\pi h} f_h^Z(t) \text{tr} L_0^* R_{t/h} = O(h^{1-d}). \quad (4.5')$$

obtain (4.5') it suffices to show that

$$\sup_{|t| \leq \bar{c}} |\text{tr} L_0^* R_{t/h}| \leq Ch^{1-d+\varepsilon} \quad (4.9)$$

holds if  $\bar{c}, \varepsilon > 0$  are small enough. The main result of this section is

## 5 Construction of the approximation

$$[\text{Op}_t(b), x_j/h] = t/overh\mu\text{Op}_t(\partial_{\xi_j}pb) + \text{Op}_t(i\partial_{\xi_j}b),$$

$$[\text{Op}_t(b), D_j] = t/overh\mu\text{Op}_t(\partial_{x_j}pb) + \text{Op}_t(i(\partial_{x_j} + \partial_{y_j})b),$$

In this section  $\bar{N} \in \mathbf{N}$  is fixed and we describe the construction of

$$Q_{\bar{N},n}(t) = \left( e^{-itp} q_{\bar{N},n}(t) \right) (x, D) \quad (4.1)$$

such that (3.15) holds. The proof of (3.14) will be given in Sections 5-7.

For  $N = 0, \dots, \bar{N}$  we are going to consider

$$q_{N,n}(t)(x, \xi) = \sum_{0 \leq k \leq N} t^k q_{k,n}^0(x, \xi) \quad (4.2(N))$$

where  $q_{N,n}(0) = q_{0,n}^0 = l_n$  and for  $k \geq 1$  we have

$$q_{k,n}^0 \in S(hp(h^{\sigma/2}\tilde{p})^{k-1}, g), \quad \text{supp } q_{k,n}^0 \subset \mathcal{B}_n(\bar{c}/2). \quad (4.3(k))$$

We construct  $q_{N,n}(t)$  by induction with respect to  $N$  and its final step corresponding to  $N = \bar{N}$  gives  $q_{\bar{N},n}(t)$  to be used in (4.1).

For a smooth function  $(t, x, \xi) \rightarrow b(t, x, \xi) \in \mathbf{C}$  we denote

$$\mathcal{P}_{\bar{N}}b(t) = \sum_{|\alpha| \leq \bar{N}} (-i)^{|\alpha|} \partial_{\xi}^{\alpha} \left( b(t) \overline{\partial_x^{\alpha} p_0} \right) / \alpha!,$$

where  $p_0$  is given by (3.24).

**Proposition 5.1** *Let  $N = 0, \dots, \bar{N}$ . Then we can find  $q_{N,n}(t)$  satisfying (4.2(N)) with  $q_{N,n}(0) = l_n$ ,  $q_{k,n}^0$  satisfying (4.3(k)) for  $k = 1, \dots, N$  and*

$$(\partial_t + i\mathcal{P}_{\bar{N}}) \left( e^{-itp} q_{N,n}(t) \right) = e^{-itp} \tilde{q}_{N,n}^0(t) \quad (4.4(N))$$

holds with

$$\tilde{q}_{N,n}^0(t)(x, \xi) = \sum_{N \leq k \leq N + \bar{N}} t^k \tilde{q}_{N,k,n}^0(x, \xi), \quad (4.5(N))$$

$$\tilde{q}_{N,k,n}^0 \in S(hp(h^{\sigma/2}\tilde{p})^k, g) \text{ for } k = N, \dots, N + \bar{N} \quad (4.6(N))$$

and  $\text{supp } \tilde{q}_{N,k,n}^0 \subset \mathcal{B}_n(\bar{c}/2)$ .

**Proof.** We introduce the notation

$$\tilde{\mathcal{P}}_{\bar{N}}q(t) = e^{itp}(\partial_t + i\mathcal{P}_{\bar{N}})(e^{-itp}q(t)).$$

If  $q(t, x, \xi) = q_0(x, \xi)$  is independent of  $t$ , then

$$\tilde{\mathcal{P}}_{\bar{N}}q(t) = \sum_{0 \leq k \leq \bar{N}} t^k \tilde{q}_k^0 \quad (4.7)$$

with

$$\begin{aligned} \tilde{q}_0^0 &= i(\bar{p}_0 - p)q_0 - \sum_{1 \leq |\alpha| \leq \bar{N}} (-i)^{|\alpha|+1} \partial_\xi^\alpha (q_0 \overline{\partial_x^\alpha p_0}) / \alpha!, \\ \tilde{q}_k^0 &= \sum_{\substack{|\alpha_0 + \dots + \alpha_k| \leq \bar{N} \\ \alpha_j \neq 0 \text{ if } j \neq 0}} c_{\alpha_0, \dots, \alpha_k} \partial_\xi^{\alpha_0} (q_0 \overline{\partial_x^{\alpha_0 + \dots + \alpha_k} p_0}) \partial_\xi^{\alpha_1} p \dots \partial_\xi^{\alpha_k} p \quad (k = 1, \dots, \bar{N}). \end{aligned}$$

Using the fact that the estimates (2.8) still hold with  $p_0$  instead of  $p$  we find

$$q_0 \in S(m, g) \implies \partial_\xi^{\alpha_0} (q_0 \partial_x^\alpha \bar{p}_0) \in S(mph^{1+\sigma|\alpha|} \langle \xi \rangle^{|\alpha| - |\alpha_0|}, g) \quad (4.8)$$

if  $\alpha \neq 0$ . Moreover the estimates (2.10') imply

$$\partial_\xi^{\alpha_1} p \dots \partial_\xi^{\alpha_k} p \in S(\tilde{p}^k \langle \xi \rangle^{-|\alpha_1| - \dots - |\alpha_k|}, g) \quad (4.9)$$

if  $\alpha_1, \dots, \alpha_k \neq 0$ . Combining (4.8), (4.9) with  $\alpha = \alpha_0 + \dots + \alpha_k \implies |\alpha| \geq \max\{k, 1\}$  in the expression of  $\tilde{q}_k^0$  we find

$$q_0 \in S(m, g) \implies \tilde{q}_k^0 \in S(mph^{1+\sigma \max\{k, 1\}} \tilde{p}^k, g) \quad (4.10)$$

(where in the case  $k = 0$  we use moreover (3.24)).

Therefore in the case  $N = 0$  when we take  $q(t) = l_n \in S(1, g)$  in (4.7), we obtain (4.4(0)) with  $\tilde{q}_{0,n}^0$  of the form (4.5(0)) and (4.10) with  $m = 1$  imply (4.6(0)), i.e. Proposition 4.1 holds for  $N = 0$ .

Further on we assume that the statement of Proposition 4.1 holds for a given  $N \leq \bar{N} - 1$  and we prove that it still holds for  $N + 1$  instead of  $N$ .

Using the induction hypothesis to express  $\tilde{\mathcal{P}}_{\bar{N}}q_{N,n}(t)$  we find

$$\begin{aligned} \tilde{\mathcal{P}}_{\bar{N}}q_{N+1,n}(t) &= \tilde{\mathcal{P}}_{\bar{N}}(t^{N+1}q_{N+1,n}^0) + \tilde{\mathcal{P}}_{\bar{N}}q_{N,n}(t) = \\ &t^N \left( (N+1)q_{N+1,n}^0 + \tilde{q}_{N,N,n}^0 \right) + t^{N+1} \tilde{\mathcal{P}}_{\bar{N}}q_{N,n+1}^0 + \sum_{N+1 \leq k \leq N+\bar{N}} t^k \tilde{q}_{N,k,n}^0. \end{aligned}$$

In order to obtain (4.5( $N + 1$ )) it suffices to cancel the term with  $t^N$  taking

$$q_{N+1,n}^0 = -\tilde{q}_{N,N,n}^0/(N + 1) \quad (4.11)$$

and we have  $\tilde{q}_{N,N,n}^0 \in S(hp(h^{\sigma/2}\tilde{p})^N, g)$  by the induction hypothesis. Let us introduce the following notation :

$$a_n(t) \in \sum_{k \in \mathcal{K}} t^k S(m(k), g) \iff a_n(t) = \sum_{k \in \mathcal{K}} t^k a_{k,n}^0 \text{ with } a_{k,n}^0 \in S(m(k), g).$$

Then using  $ph \leq \tilde{p}$  and  $\max\{k, 1\} \geq (k + 1)/2$  we can write (4.7), (4.10) in the following form

$$q(t) = q_0 \in S(m, g) \implies \tilde{\mathcal{P}}_{\bar{N}} q(t) \in \sum_{0 \leq k \leq \bar{N}} t^k S(m(h^{\sigma/2}\tilde{p})^{k+1}, g). \quad (4.12)$$

Since (4.11) gives (4.3( $N + 1$ ))), applying (4.12) with  $m = ph(h^{\sigma/2}\tilde{p})^N$  we find

$$t^{N+1}\tilde{\mathcal{P}}_{\bar{N}} q_{N,n+1}^0 \in \sum_{0 \leq k \leq \bar{N}} t^{N+1+k} S(ph(h^{\sigma/2}\tilde{p})^{N+k+1}, g)$$

and (4.6( $N + 1$ )) follows.  $\triangle$

**Proposition 5.2** *Let  $Q_{\bar{N},n}(t)$  be defined by (4.1) with  $q_{\bar{N},n}$  given by (4.2( $\bar{N}$ )) where  $q_{n,0}^0 = l_n$  and  $q_{k,n}^0$  satisfy (4.3( $k$ )) for  $k = 1, \dots, \bar{N}$ . Then (3.15) holds.*

**Proof.** For  $k \in \mathbf{N}$ ,  $n \in \mathbf{N}^*$ ,  $q \in C_0^\infty(\mathbf{R}^d \times \mathbf{R}^d)$  and  $\lambda \in \mathbf{R}$  we denote

$$\mathcal{N}_{k,n}(q, \lambda) = \int_{-\infty}^{\infty} dt \check{f}_{n,\lambda}^\pm(t) t^k J_t(q), \quad (4.13)$$

where

$$J_t(q) = (2\pi)^{-d} \int e^{-itp(x,\xi)} q(x, \xi) dx d\xi. \quad (4.14)$$

Since  $t^k \check{f}_{n,\lambda}^\pm(t)$  is the Fourier inverse of  $i^k (f_{n,\lambda}^\pm)^{(k)}$ , changing the order of integrals (4.13) and (4.14) we find

$$\mathcal{N}_{k,n}(q, \lambda) = (2\pi)^{-d} \int i^k (f_{n,\lambda}^\pm)^{(k)}(p(x, \xi)) q(x, \xi) dx d\xi. \quad (4.15)$$

Since  $(e^{-itp}q)(x, D)L_n^*$  has the integral kernel

$$(x, y) \rightarrow (2\pi)^{-d} \int e^{-itp(x,\xi)} q(x, \xi) l_n(y, \xi) d\xi,$$

we have  $\text{tr } L_n^*(e^{-itp}q)(x, D) = J_t(ql_n)$  and

$$\int_{-\infty}^{\infty} dt \check{f}_{n,\lambda}^{\pm}(t) \text{tr } Q_{\bar{N},n}(t)L_n^* = \sum_{0 \leq k \leq \bar{N}} \mathcal{N}_{k,n}(q_{k,n}^0 l_n, \lambda). \quad (4.16)$$

For  $k = 0$  we have  $q_{0,n}^0 = l_n$ , hence (4.15) and (3.8) give

$$\begin{aligned} & \left| \mathcal{N}_{0,n}(q_{0,n}^0 l_n, \lambda) - (2\pi)^{-d} \int_{p < \lambda} l_n(x, \xi)^2 dx d\xi \right| \\ &= \left| (2\pi)^{-d} \int (f_{n,\lambda}^{\pm} - \mathbf{1}_{]-\infty, \lambda[}) (p(x, \xi)) l_n(x, \xi)^2 dx d\xi \right| \\ &\leq \int_{\lambda - h_n p_n < p < \lambda + h_n p_n} \mathbf{1}_{\mathcal{B}_n(\bar{c})}(x, \xi) dx d\xi. \end{aligned}$$

It remains to show that the estimate

$$|\mathcal{N}_{k,n}(q_n, \lambda)| \leq \bar{C} \int_{\lambda - h_n p_n < p < \lambda + h_n p_n} \mathbf{1}_{\mathcal{B}_n(\bar{c})}(x, \xi) dx d\xi \quad (4.17(k))$$

holds with  $q_n = q_{k,n}^0 l_n$ ,  $k = 1, \dots, \bar{N}$ .

The expression (4.15) allows to write the obvious inequality

$$|\mathcal{N}_{k,n}(q_n, \lambda)| \leq \int |(f_{n,\lambda}^{\pm})^{(k)}(p(x, \xi))| |q_n(x, \xi)| dx d\xi. \quad (4.18(k))$$

In the case  $k = 1$  the above inequality leads to the following :

**Corollary 5.3** *If  $q_n \in S(ph, g)$  are such that  $\text{supp } q_n \subset \mathcal{B}_n(\bar{c})$ , then*

*$|q_n| \leq Ch_n p_n$  and using (3.9(1)) in (4.18(1)) we obtain (4.17(1)).*

*In particular (4.17(1)) holds with  $q_n = q_{1,n}^0 l_n$ .*

In order to prove (4.17(k)) for  $q_n = q_{k,n}^0 l_n$ ,  $k \geq 2$ , let  $\chi \in C_0^\infty(]-2; 2[)$  be such that  $\chi = 1$  on  $[-1; 1]$  and for  $s > 0$  let

$$\chi_s(x, \xi) = \chi(m(x, \xi)/s^2) \quad \text{with } m = \langle \xi \rangle^2 |\nabla_\xi p|^2 h^{-2} p^{-2}. \quad (4.19)$$

We consider also  $\tilde{\chi}_s := 1 - \chi_s$  and remark that using  $1 + m \in S(1 + m, g)$  it is easy to check that  $\chi_s, \tilde{\chi}_s \in S(1, g)$ . Moreover

$$(x, \xi) \in \text{supp } \chi_s \implies \tilde{p}(x, \xi) = ((1 + m^{1/2})hp)(x, \xi) \leq (1 + 2s)(hp)(x, \xi). \quad (4.20)$$

Therefore  $|q_{k,n}^0 l_n \chi_s| \leq C_k h_n^k p_n^k$  and using (3.9(k)) in (4.18(k)) we find that (4.17(k)) holds with  $q_n = q_{k,n}^0 l_n \chi_s$ ,  $k \geq 2$ .

Thus it remains to show that (4.17(k)) holds with  $q_n = q_{k,n}^0 l_n \tilde{\chi}_s$ ,  $k \geq 2$ .

To obtain this result it suffices to show

**Lemma 5.4** *If  $q_n \in S(m, g)$  then we can find  $\tilde{q}_n \in S(m/\tilde{p}, g)$  such that  $\text{supp } \tilde{q}_n \subset \text{supp } q_n$  and*

$$t J_t(q_n \tilde{\chi}_s) = J_t(\tilde{q}_n \tilde{\chi}_{s/2}). \quad (4.21)$$

Indeed, iterating the assertion of Lemma 4.4 we can write

$$t^k J_t(q_{k,n}^0 l_n \tilde{\chi}_s) = t^{k-1} J_t(q_{1,k,n} \tilde{\chi}_{s/2}) = \dots = t^{k-\bar{k}} J_t(q_{\bar{k},k,n} \tilde{\chi}_{s/2^{\bar{k}}}). \quad (4.22)$$

for some  $q_{\bar{k},k,n} \in S(ph\tilde{p}^{k-1-\bar{k}}, g)$  with  $\text{supp } q_{\bar{k},k,n} \subset \mathcal{B}_n(\bar{c}/2)$ . Thus using (4.22) with  $\bar{k} = k - 1$  we obtain

$$\mathcal{N}_{k,n}(q_{k,n}^0 l_n \tilde{\chi}_s, \lambda) = \mathcal{N}_{1,n}(q_n, \lambda)$$

with  $q_n \in S(hp, g)$ ,  $\text{supp } q_n \subset \mathcal{B}_n(\bar{c}/2)$ , and (4.17(1)) holds due to Corollary 4.3.

**Proof of Lemma 4.4.** Using  $\langle \xi \rangle^2 |\nabla_\xi p|^2 \in S(\tilde{p}^2, g)$  and

$$s^2 \tilde{p}(x, \xi)^2 \leq 2s^2 (hp)(x, \xi)^2 + 2\langle \xi \rangle^2 |\nabla_\xi p(x, \xi)|^2 \leq 2(s^2 + 1) \langle \xi \rangle^2 |\nabla_\xi p(x, \xi)|^2$$

for  $(x, \xi) \in \text{supp } \tilde{\chi}_s$ , we obtain  $\tilde{\chi}_s \langle \xi \rangle^{-2} |\nabla_\xi p|^{-2} \in S(\tilde{p}^{-2}, g)$  due to (3.20) and combining with  $\langle \xi \rangle^2 \partial_{\xi_j} p \in S(\langle \xi \rangle \tilde{p}, g)$  we can define

$$\tilde{\chi}_{j,s} := \tilde{\chi}_s \partial_{\xi_j} p |\nabla_\xi p|^{-2} \in S(\langle \xi \rangle / \tilde{p}, g). \quad (4.23)$$

Writing  $q_n \tilde{\chi}_s = \sum_{j=1}^d q_n \tilde{\chi}_{j,s} \partial_{\xi_j} p$  and integrating by parts we find

$$\begin{aligned} t J_t(q_n \tilde{\chi}_{j,s} \partial_{\xi_j} p) &= (2\pi)^{-d} \int i \partial_{\xi_j} \left( e^{-itp(x,\xi)} \right) (q_n \tilde{\chi}_{j,s})(x, \xi) dx d\xi \\ &= -i J_t(\partial_{\xi_j} (q_n \tilde{\chi}_{j,s})), \end{aligned}$$

which completes the proof due to  $\partial_{\xi_j} (q_n \tilde{\chi}_{j,s}) \in S(m/\tilde{p}, g)$  and  $\text{supp } \tilde{\chi}_{j,s} \cap \text{supp } \chi_{s/2} = \emptyset$ .  $\triangle$

## 6 Preliminary remarks about the approximation error

In this section we begin a study of  $\text{tr } \tilde{Q}_{\bar{N},n}(t - \tau) e^{-i\tau P} L_n^*$  with the purpose to establish (3.18), which implies (3.14).

To abbreviate notations we denote the elements of  $\mathcal{V}$  by the letter  $\nu$ , adopting the following convention : if the index  $\nu$  appears in a formula simultaneously



with a letter  $n$ ,  $t$  or  $\tau$ , then  $\nu = (n, t, \tau)$ . Moreover the notation  $s_\nu = \tilde{s}_\nu + O(m_{n,t})$  means that  $|s_{n,t,\tau} - \tilde{s}_{n,t,\tau}| \leq C m_{n,t}$  holds with a constant  $C > 0$  independent of  $\nu = (n, t, \tau) \in \mathcal{V}$ .

Assume that  $Q_\nu = (e^{-itp} q_\nu)(x, D)$  with  $q_\nu \in S(m, g)$ ,  $\text{supp } q_\nu \subset \mathcal{B}_n(\bar{c})$  and  $Y_\nu \in B(L^2(\mathbf{R}^d))$  for  $\nu \in \mathcal{V}$ . Then Lemma 9.1 allows to estimate

$$\begin{aligned} |t^k \text{tr } Q_\nu e^{-i\tau P} Y_\nu| &\leq |t|^k \|Q_\nu\|_{\text{tr}} \|Y_\nu\| \leq \\ &\leq C m_n h_n^{-C_0} \langle h_n p_n t \rangle^{k+C_0} \|Y_\nu\|, \end{aligned} \quad (5.1)$$

where  $m_n$  are as in (3.6).

The family  $\{Y_\nu\}_{\nu \in \mathcal{V}} \subset B(L^2(\mathbf{R}^d))$  will be called negligible if for every  $N \in \mathbf{N}$  we can find  $C(N) > 0$  such that

$$\|Y_\nu\| = O\left(h_n^N \langle h_n p_n t \rangle^{C(N)}\right). \quad (5.2)$$

If  $N', k \in \mathbf{N}$  and  $\{Y_\nu\}_{\nu \in \mathcal{V}}$  is negligible, then (5.1) allows to find  $C(k, N') > 0$  such that

$$t^k \text{tr } Q_\nu e^{-i\tau P} Y_\nu = O\left(h_n^{N'} \langle h_n p_n t \rangle^{C(k, N')}\right). \quad (5.3)$$

**Proposition 6.1** *Let  $c > \bar{c}/2$  and consider  $b_\nu \in S(m, g)$  satisfying the condition  $\text{supp } b_\nu \cap \mathcal{B}_n(c) = \emptyset$ . Then the family*

$$R_\nu = b_\nu^{\mathcal{W}}(x, D) e^{-i\tau P} L_n^* \quad (5.4)$$

*is negligible.*

Before giving the proof of this result we describe its consequences.

We assume  $\bar{c} > c > c' > \bar{c}/2$  and note that the method of ch. 18.4 [16] (cf. the beginning of Section 3) allows to find  $\tilde{\theta}_k^- \in C_0^\infty(\mathcal{B}_{\bar{x}(k)}^-(c))$  such that  $\tilde{\theta}_k^- = 1$  on  $\mathcal{B}_{\bar{x}(k)}^-(c')$  and

$$|\partial^\alpha \tilde{\theta}_k^-(x)| \leq C_\alpha v(x)^{-\rho|\alpha|}.$$

If  $\tilde{\Theta}_k^-$  denotes the operator of multiplication by  $\tilde{\theta}_k^-$ , then Proposition 5.1 ensures that  $(I - \tilde{\Theta}_{k_n}^-) e^{-i\tau P} L_n^*$  and  $P(I - \tilde{\Theta}_{k_n}^-) e^{-i\tau P} L_n^*$  are negligible, hence for every  $N \in \mathbf{N}$  we can find  $C(N) > 0$  such that

$$\begin{aligned} \text{tr } \tilde{Q}_{\bar{N}, n}(t - \tau) e^{-i\tau P} L_n^* = \\ \text{tr } \tilde{Q}_{\bar{N}, n}(t - \tau) \tilde{\Theta}_{k_n}^- e^{-i\tau P} L_n^* + O\left(h_n^N \langle h_n p_n t \rangle^{C(N)}\right). \end{aligned} \quad (5.5)$$

Further on we prefer using the operators  $\tilde{Q}_{\bar{N},n}(t-\tau)\tilde{\Theta}_{k_n^-}^-$  instead of  $\tilde{Q}_{\bar{N},n}(t-\tau)$  and for this purpose we introduce new notations.

**Symbols depending on**  $(x, \xi, y) \in \mathbf{R}^d \times \mathbf{R}^d \times \mathbf{R}^d$ .

If  $q_\nu \in C_0^\infty(\mathbf{R}^d \times \mathbf{R}^d \times \mathbf{R}^d)$ , then  $\text{Op}(e^{i(t-\tau)p}q_\nu)$  will denote the integral operator with the kernel

$$(x, y) \rightarrow \mathcal{K}_\nu(q, x, y) = (2\pi)^{-d} \int e^{i(x-y)\xi - i(t-\tau)p(x,\xi)} q_\nu(x, \xi, y) d\xi. \quad (5.6)$$

For  $c > 0$  and  $n \in \mathbf{N}^*$  we will denote

$$\tilde{\mathcal{B}}_n(c) := \mathcal{B}_{\bar{x}(k_n^-)}^-(c) \times \mathcal{B}_{\bar{\xi}(k_n^+)}^+ \times \mathcal{B}_{\bar{x}(k_n^-)}^-(c)$$

and for  $m \in \mathcal{M}_{\bar{c}}(\tilde{g})$  we will write  $q_\nu \in \tilde{S}(m)$  if and only if there is  $c < \bar{c}$  such that  $q_\nu \in C_0^\infty(\tilde{\mathcal{B}}_n(c))$  and

$$|\partial_{x,y}^\beta \partial_\xi^\alpha q_\nu(x, \xi, y)| \leq C_{\alpha,\beta} m_n h_n^{|\sigma|} \langle \xi \rangle^{|\beta| - |\alpha|}$$

holds for every  $\alpha \in \mathbf{N}^d$ ,  $\beta \in \mathbf{N}^{2d}$ , where  $m_n$  are as in (3.6).

These notations will be used below to express the approximation error in a particular form, similar to (4.13), (4.14), (4.16).

**Expressions of**  $\text{tr} \tilde{Q}_{\bar{N},n}(t-\tau)\tilde{\Theta}_{k_n^-}^- e^{-i\tau P} L_n^*$ .

We have  $p^{\mathcal{W}}(x, D) = p_0(x, D) = p_0(x, D)^*$  and

$$Q_{\bar{N},n}(t)p_0(x, D)^* = \text{Op}\left(e^{-itp}\tilde{q}_{\bar{N},n,t}\right)$$

with

$$\tilde{q}_{\bar{N},n,t}(x, \xi, y) = q_{\bar{N},n}(t)(x, \xi) \overline{p_0(y, \xi)}.$$

Writing the Taylor's development of  $\tilde{q}_{\bar{N},n,t}(x, \xi, \cdot)$  in  $x$  and applying standard integrations by parts in the integrals of the form (5.6) based on the equality  $(x-y)^\alpha e^{i(x-y)\xi} = i^{|\alpha|} \partial_{\xi_j}^\alpha (e^{i(x-y)\xi})$ , we find

$$\tilde{Q}_{\bar{N},n}(t) = \text{Op}\left(e^{-itp}(\tilde{q}_{\bar{N},n}^0 + \tilde{q}_{\bar{N},n}^1)(t)\right),$$

where  $\tilde{q}_{\bar{N},n}^0(t)$  are as in Proposition 4.1 and the remainder term of the Taylor's development of order  $\bar{N}$  gives

$$\tilde{q}_{\bar{N},n}^1(t)(x, \xi, y) = e^{itp(x,\xi)} (\bar{N} + 1) \int_0^1 ds (1-s)^{\bar{N}} \tilde{q}_{\bar{N},n}(t, s)(x, \xi, y) \quad (5.7)$$

with

$$\begin{aligned} \tilde{q}_{\bar{N},n}(t,s)(x,\xi,y) = \\ \sum_{|\alpha|=\bar{N}+1} (-i)^{|\alpha|} \left( \partial_{\xi}^{\alpha} \left( q_{\bar{N},n}(t) e^{-itp} \right) (x,\xi) \overline{\partial_x^{\alpha} p_0(x+s(y-x),\xi)} \right) / \alpha!. \end{aligned}$$

We can write

$$\begin{aligned} \tilde{q}_{\bar{N},n}^0(t-\tau)(x,\xi) \tilde{\theta}_{k_n}^-(y) &= \sum_{\bar{N} \leq k \leq 2\bar{N}} (t-\tau)^k \tilde{q}_{\bar{N},k,n}^0(x,\xi) \tilde{\theta}_{k_n}^-(y) \\ &= \sum_{\bar{N} \leq k \leq 2\bar{N}} t^k \tilde{q}_{\bar{N},k,\nu}^0(x,\xi,y), \end{aligned}$$

where  $\nu = (n, t, \tau)$  according to our convention and

$$\tilde{q}_{\bar{N},k,\nu}^0(x,\xi,y) = \left(1 - \frac{\tau}{t}\right)^k \tilde{q}_{\bar{N},k,n}^0(x,\xi) \tilde{\theta}_{k_n}^-(y).$$

It is clear that  $\tilde{q}_{\bar{N},k,\nu}^0 \in \tilde{S}(ph^{\bar{N}\sigma/2}\tilde{p}^k)$  and using the form of  $q_{\bar{N},n}^0(t)$  in (5.7) we find a similar expression

$$\tilde{q}_{\bar{N},n}^1(t-\tau)(x,\xi,y) \tilde{\theta}_{k_n}^-(y) = \sum_{0 \leq k \leq 2\bar{N}} t^k \tilde{q}_{\bar{N},k,\nu}^1(x,\xi,y)$$

with  $\tilde{q}_{\bar{N},k,\nu}^1 \in \tilde{S}(ph^{\bar{N}\sigma/2}\tilde{p}^k)$ .

If  $Y_{\nu} \in B(L^2(\mathbf{R}^d))$  and  $q_{\nu} \in \tilde{S}(m)$  then the quantity

$$J_{\nu}(q, Y) := \text{tr} \left( \text{Op}(e^{i(\tau-t)p} q_{\nu}) e^{-i\tau P} Y_{\nu} \right), \quad (5.8)$$

is well defined due to Lemma 9.1 and analogically to (5.1) we have

$$|t^k J_{\nu}(q, Y)| \leq C m_n h_n^{-C_0} \langle h_n p_n t \rangle^{k+C_0} \|Y_{\nu}\|, \quad (5.9)$$

where  $m_n$  are as in (3.6).

Using the above notation we can state the following conclusion

**Corollary 6.2** *There exist  $\tilde{q}_{\bar{N},k,\nu} \in \tilde{S}(ph^{\bar{N}\sigma/2}\tilde{p}^k)$  such that*

$$\text{tr} \tilde{Q}_{\bar{N},n}(t-\tau) \tilde{\Theta}_{k_n}^- e^{-i\tau P} L_n^* = \sum_{0 \leq k \leq 2\bar{N}} t^k J_{\nu}(\tilde{q}_{\bar{N},k}, Y) \quad (5.10)$$

holds with  $Y_{\nu} = L_n^*$ .

We will complete the proof of Theorem 2.1 by showing

**Proposition 6.3** *Assume  $N_0, k \in \mathbf{N}^*$ ,  $Y_\nu = L_n^*$  and  $q_\nu \in \tilde{S}(m)$ . Then one can find  $k_0 \in \mathbf{N}^*$  and  $C(N_0) > 0$  such that*

$$t^k J_\nu(q, Y) = \sum_{1 \leq \bar{k} \leq k_0} J_\nu(q_{\bar{k}}, Y_{\bar{k}}) + O\left(h_n^{N_0} \langle h_n p_n t \rangle^{C(N_0)}\right). \quad (5.11)$$

holds for certain families of symbols  $\{q_{\bar{k}, \nu}\}_{\nu \in \mathcal{V}}$  and operators  $\{Y_{\bar{k}, \nu}\}_{\nu \in \mathcal{V}}$  satisfying

$$q_{\bar{k}, \nu} \in \tilde{S}(m \tilde{p}^{-k}), \quad \|Y_{\bar{k}, \nu}\| = O\left(\langle h_n p_n t \rangle^{C(N_0)}\right) \quad (5.12)$$

for  $\bar{k} = 1, \dots, k_0$ .

Indeed, using (5.10) and Proposition 5.3 with  $q_\nu = \tilde{q}_{\bar{N}, k, \nu}$ ,  $m = p h^{\bar{N}\sigma/2} \tilde{p}^k$ , we can write

$$\text{tr } \tilde{Q}_{\bar{N}, n}(t-\tau) \tilde{\Theta}_{k_n^-}^- e^{-i\tau P} L_n^* = \sum_{1 \leq \bar{k} \leq k(\bar{N})} J_\nu(q_{\bar{N}, \bar{k}}, Y_{\bar{N}, \bar{k}}) + O\left(h_n^{N_0} \langle h_n p_n t \rangle^{C(\bar{N}, N_0)}\right),$$

with

$$q_{\bar{N}, \bar{k}, \nu} \in \tilde{S}(p h^{\bar{N}\sigma/2}), \quad \|Y_{\bar{N}, \bar{k}, \nu}\| = O\left(\langle h_n p_n t \rangle^{C(\bar{N}, N_0)}\right),$$

hence choosing  $\bar{N} = \bar{N}(N_0)$  large enough we can ensure

$$J_\nu(q_{\bar{N}, \bar{k}}, Y_{\bar{N}, \bar{k}}) = O\left(h_n^{N_0} \langle h_n p_n t \rangle^{\bar{C}(\bar{N}, N_0)}\right)$$

due to (5.9). This proves (3.18) completing the proof of Theorem 3.2 and 2.1.

**Proof of Proposition 5.1.** The method of ch. 18.4 [16] allows to find symbols  $l_n^0(x, \xi) = \theta_{k_n^-}^{0-}(x) \theta_{k_n^+}^{0+}(\xi)$  satisfying

$$\begin{aligned} l_n^0 \in S(1, g), \quad \partial_{x_j} l_n^0 \in S(v(x)^{-\rho}, g), \quad \text{supp } l_n^0 \subset \mathcal{B}_n(c), \\ l_n^0 = 1 \text{ on } \mathcal{B}_n(c') \text{ with } c > c' > \bar{c}/2. \end{aligned} \quad (5.13)$$

We set  $L_n^0 = l_n^0(x, D)$  and note that  $\|b_\nu^\mathcal{W}(x, D) L_n^0\| = O(h_n^N)$  for every  $N \in \mathbf{N}$  holds due to  $\text{supp } b_\nu \cap \text{supp } l_n^0 = \emptyset$  (cf. Theorem 18.5.4 and 18.6.3 [16]). Therefore it suffices to show that

$$\|(I - L_n^0) e^{-i\tau P} L_n^*\| = O\left(h_n^{k\sigma} \langle h_n p_n \tau \rangle^k\right) \quad (5.14(k))$$

holds for every  $k \in \mathbf{N}$  if  $l_n^0$  satisfies (5.13). Obviously (5.14( $k$ )) holds for  $k = 0$  and we will prove the general statement by induction with respect to  $k \in \mathbf{N}$ .

Further on we assume that  $\sigma > 0$  is small enough to ensure  $\rho'(1 + \sigma) \leq \rho$  and  $r'(1 + \sigma) \leq 1$ . This assumption implies  $v(x)^{-\rho} \langle \xi \rangle^{-1} \leq h(x, \xi)^{1+\sigma}$  and  $\partial_{x_j} l_n^0 \partial_{\xi_j} p, \partial_{x_j} p \partial_{\xi_j} l_n^0 \in S(ph^{1+\sigma}, g)$  if  $l_n^0$  satisfies (5.13). Thus (3.23) ensures

$$[I - L_n^0, P] = -[L_n^0, P] = \tilde{l}_n^{\mathcal{W}}(x, D) \quad \text{with } \tilde{l}_n \in S(ph^{1+\sigma}, g). \quad (5.15)$$

Moreover we can find

$$\begin{aligned} l_n^1 &\in S(1, g), \quad \partial_{x_j} l_n^1 \in S(v(x)^{-\rho}, g), \quad \text{supp } l_n^1 \subset \mathcal{B}_n(c'), \\ l_n^1 &= 1 \quad \text{on } \mathcal{B}_n(c'') \quad \text{with } c' > c'' > \bar{c}/2 \end{aligned} \quad (5.16)$$

and setting  $L_n^1 = l_n^1(x, D)$  we have  $\text{supp } \tilde{l}_n \cap \text{supp } l_n^1 = \emptyset$ , hence

$$\|\tilde{l}_n^{\mathcal{W}}(x, D)L_n^1\| = O(h_n^N) \quad \text{for every } N \in \mathbf{N}. \quad (5.17)$$

Since  $\|(I - L_n^0)L_n^*\| = O(h_n^N)$  for every  $N \in \mathbf{N}$  and

$$(I - L_n^0)e^{-i\tau P}L_n^* = e^{-i\tau P}(I - L_n^0)L_n^* + [I - L_n^0, e^{-i\tau P}]L_n^*, \quad (5.18)$$

it remains to estimate the norm of

$$[I - L_n^0, e^{-i\tau P}]L_n^* = \int_0^1 ds \tau e^{i(s-1)\tau P} [I - L_n^0, P] e^{-is\tau P} L_n^*. \quad (5.19)$$

and due to (5.17) it suffices to estimate

$$|\tau| \|\tilde{l}_n^{\mathcal{W}}(x, D)(I - L_n^1)e^{-is\tau P}L_n^*\|. \quad (5.20)$$

However we have  $\|\tilde{l}_n^{\mathcal{W}}(x, D)\| \leq Cp_n h_n^{1+\sigma}$  (cf. Theorem 18.6.3 [16]) and using the induction hypothesis we can assume that (5.14( $k$ )) holds with  $L_n^1$  instead of  $L_n^0$  and  $c''$  instead of  $c'$ . Thus the quantity (5.20) can be estimated by

$$C|\tau| p_n h_n^{1+\sigma} h_n^{k\sigma} \langle h_n p_n \tau \rangle^k \leq Ch_n^{(k+1)\sigma} \langle h_n p_n \tau \rangle^{k+1},$$

completing the proof of (5.14( $k + 1$ )).  $\triangle$ .

## 7 Auxiliary commutator formulas

**Notations.** We will write  $b_\nu \in S_n(m, g)$  if and only if there exist  $c_0 < \bar{c}$ ,  $l_n^0 \in S(1, g)$  satisfying  $\text{supp } l_n^0 \subset \mathcal{B}_n(c_0)$  and  $(1 - l_n^0)b_\nu \in S(h^N, g)$  for every  $N \in \mathbf{N}$ .

Then Theorem 18.5.4 [16] ensures  $b\tilde{b}, b \odot \tilde{b} \in S_n(m\tilde{m}, g)$  if  $b_\nu \in S_n(m, g)$ ,  $\tilde{b}_\nu \in S(\tilde{m}, g)$  and (3.23) still holds with  $b, \tilde{b}, S(m', g)$  replaced by  $b_\nu, \tilde{b}_\nu, S_n(m', g)$ . Moreover  $b_\nu \in S_n(m, g)$  implies

$$|\partial_x^\beta \partial_\xi^\alpha b_\nu(x, \xi)| \leq C_{\alpha, \beta} m_n h_n^{|\sigma|} |\langle \xi \rangle^{|\beta| - |\alpha|} \quad (6.1)$$

where  $m_n$  are as in (3.6) and Theorem 8.6.3 [16] ensures

$$\|b_\nu^{\mathcal{W}}(x, D)\| \leq C m_n, \quad \|(I - l_n^0)^{\mathcal{W}}(x, D) b_\nu^{\mathcal{W}}(x, D)\| = O(h_n^N) \quad (6.2)$$

for every  $N \in \mathbf{N}$ .

We introduce the following formal notation

$$Y(\tau, B) := e^{-i\tau P} B e^{i\tau P}. \quad (6.3)$$

Let  $\{Y_\nu\}_{\nu \in \mathcal{V}}$  be a family of bounded operators and let  $m \in \mathcal{M}_{\bar{c}}(\tilde{g})$ . We write  $Y_\nu \in \mathcal{Y}(m)$  if and only if there exist  $N \in \mathbf{N}$ ,  $C_0 > 0$ , the weights  $m(k, k') \in \mathcal{M}_{\bar{c}}(\tilde{g})$ , the symbols  $b_{k, k', \nu} \in S_n(m(k, k'), g)$  and functions  $s_{k, k'} : [0; 1]^N \rightarrow \mathbf{R}$ ,  $s_{k, \nu} : [0; 1]^N \rightarrow \mathbf{C}$ , satisfying

$$\prod_{k'=1}^N m(k, k') \leq m, \quad |s_{k, k'}(w)| \leq C_0, \quad |s_{k, \nu}(w)| \leq C_0 \langle h_n p_n t \rangle^{C_0},$$

for  $k, k' = 1, \dots, N$  and

$$Y_\nu = \sum_{k=1}^N \int_{[0; 1]^N} dw s_{k, \nu}(w) Y(s_{k, 1}(w)\tau, B_{k, 1, \nu}) \dots Y(s_{k, N}(w)\tau, B_{k, N, \nu}) + R_\nu \quad (6.4)$$

where  $B_{k, k', \nu} = b_{k, k', \nu}^{\mathcal{W}}(x, D)$  and the family  $\{R_\nu\}_{\nu \in \mathcal{V}}$  is negligible.

Taking  $s_{k, \nu}(w) = s_{k, k'}(w) = 1$  we can forget the integration with respect to  $w$ , hence  $Y(\tau, B_{k, 1, \nu}) \dots Y(\tau, B_{k, N, \nu}) \in \mathcal{Y}(m)$  and more generally

$$Y_\nu \in \mathcal{Y}(m), \quad \tilde{Y}_\nu \in \mathcal{Y}(\tilde{m}) \implies Y_\nu \tilde{Y}_\nu \in \mathcal{Y}(m\tilde{m}), \quad (6.5)$$

$$Y_\nu \in \mathcal{Y}(m) \implies \|Y_\nu\| \leq C m_n \langle h_n p_n t \rangle^C. \quad (6.6)$$

**Reformulation of Proposition 5.3.** In Section 7 we will prove

**Proposition 7.1** *Assume  $q_{0,\nu} \in \tilde{S}(m)$ ,  $Y_{0,\nu} \in \mathcal{Y}(1)$  and  $N_0 \in \mathbf{N}$ . Then one can find  $k_0 \in \mathbf{N}^*$  and  $C(N_0) > 0$  such that*

$$tJ_\nu(q_0, Y_0) = \sum_{1 \leq \bar{k} \leq k_0} (J_\nu(q_{\bar{k}}, Y_{\bar{k}}) + tJ_\nu(q_{-\bar{k}}, Y_{-\bar{k}})) + O\left(h_n^{N_0} \langle h_n p_n t \rangle^{C(N_0)}\right) \quad (6.7)$$

holds with certain symbols  $q_{\pm\bar{k},\nu}$  and operators  $Y_{\pm\bar{k},\nu}$  satisfying

$$q_{\bar{k},\nu} \in \tilde{S}(m/\tilde{p}), \quad q_{-\bar{k},\nu} \in \tilde{S}(mh^\sigma), \quad Y_{\pm\bar{k},\nu} \in \mathcal{Y}(1) \quad (6.8)$$

for  $\bar{k} = 1, \dots, k_0$ .

It is easy to see that Proposition 6.1 implies Proposition 5.3. Indeed, first of all we note that the assertion of Proposition 6.1 can be applied to express  $tJ_\nu(q_{-\bar{k}}, Y_{-\bar{k}})$ ,  $\bar{k} = 1, \dots, k_0$  and iterating this procedure  $N$  times we find the expression of  $tJ_\nu(q_0, Y_0)$  in the form (6.7) with new symbols  $q_{-\bar{k},\nu} \in \tilde{S}(mh^{N\sigma})$ ,  $\bar{k} = 1, \dots, k_N$ . Thus for  $N = N(N_0)$  large enough all terms  $tJ_\nu(q_{-\bar{k}}, Y_{-\bar{k}})$ ,  $\bar{k} = 1, \dots, k_N$ , become  $O\left(h_n^{N_0} \langle h_n p_n t \rangle^{C(N_0)}\right)$ , i.e. the assertion of Proposition 6.1 holds with  $q_{-\bar{k}} = 0$  for  $\bar{k} \geq 1$ . This proves Proposition 5.3 in the case  $k = 1$  and it is clear that the general case follows after  $k$  iterations.

In the remaining part of this section we describe the properties of  $Y_\nu \in \mathcal{Y}(m)$  needed in the proof of Proposition 6.1. More precisely we consider the commutator of  $Y_\nu$  with the operator of multiplication by the  $j$ -th coordinate, denoted by  $x_j$ .

**Lemma 7.2** *Assume  $Y_\nu \in \mathcal{Y}(m)$ . Then there exist*

$$Y_\nu^+ \in \mathcal{Y}(\langle \xi \rangle^{-1}m), \quad Y_\nu^- \in \mathcal{Y}(\langle \xi \rangle^{-1}\tilde{p}h^\sigma m) \quad (6.9)$$

such that  $[Y_\nu, x_j] = Y_\nu^+ + \tau Y_\nu^-$ .

**Proof.** Let  $B_{k,k',\nu} = b_{k,k',\nu}^{\mathcal{W}}(x, D)$  with  $b_{k,k',\nu} \in S_n(m(k, k'), g)$ . If we know that

$$[Y(\tau, B_{k,k',\nu}), x_j] = Y_{k,k',\nu}^+ + \tau Y_{k,k',\nu}^- \quad (6.10)$$

holds with

$$Y_{k,k',\nu}^+ \in \mathcal{Y}(\langle \xi \rangle^{-1}m(k, k')), \quad Y_{k,k',\nu}^- \in \mathcal{Y}(\langle \xi \rangle^{-1}\tilde{p}h^\sigma m(k, k')),$$

then commuting succesively  $x_j$  with  $Y(s_{k,k'}(w)\tau, B_{k,k',\nu})$ ,  $k' = 1, \dots, N$ , we obtain easily the general statement of Lemma 6.2.

To begin we write

$$[Y(\tau, B_{k,k',\nu}), x_j] = e^{-i\tau P} [B_{k,k',\nu}, Y(-\tau, x_j)] e^{i\tau P} \quad (6.11)$$

and denote

$$P_j := [iP, x_j] = \partial_{\xi_j} p^{\mathcal{W}}(x, D) = \partial_{\xi_j} p_0(x, D). \quad (6.12)$$

Then we can write the Taylor formula

$$\begin{aligned} Y(-\tau, x_j) &= x_j - \tau \partial_{\tau} Y(0, x_j) + \tau^2 \int_0^1 ds (1-s) \partial_{\tau}^2 Y(-s\tau, x_j) \\ &= x_j + \tau P_j + \tau e^{-i\tau P} \bar{Y}(\tau, [i\tau P, P_j]) e^{i\tau P}, \end{aligned} \quad (6.13)$$

where

$$\bar{Y}(\tau, B) = \tau \int_0^1 ds (1-s) Y((-s-1)\tau, B).$$

Using (6.13) we can express the commutator (6.11) in the form

$$Y(\tau, [B_{k,k',\nu}, x_j]) + \tau Y(\tau, [B_{k,k',\nu}, P_j]) + \tau [Y(\tau, B_{k,k',\nu}), \bar{Y}(\tau, [i\tau P, P_j])] \quad (6.14)$$

and since  $[B_{k,k',\nu}, x_j] = \partial_{\xi_j} b_{k,k',\nu}^{\mathcal{W}}(x, D)$ , it is clear that the first term of (6.13) is in  $\mathcal{Y}(\langle \xi \rangle^{-1} m(k, k'))$ . Next we check that

$$\partial_{x_j} b_{k,k',\nu} \partial_{\xi_j} \partial_{\xi_j} p \in S_n(\langle \xi \rangle^{-1} \tilde{p} h^{\sigma} m(k, k'), g)$$

$$\partial_{\xi_j} b_{k,k',\nu} \partial_{x_j} \partial_{\xi_j} p S_n(\langle \xi \rangle^{-1} p h^{1+\sigma} m(k, k'), g) \subset S_n(\langle \xi \rangle^{-1} \tilde{p} h^{\sigma} m(k, k'), g)$$

due to  $hp \leq \tilde{p}$ , hence (3.23) implies

$$b_{k,k',\nu} \odot \partial_{\xi_j} p - \partial_{\xi_j} p \odot b_{k,k',\nu} \in S_n(\langle \xi \rangle^{-1} \tilde{p} h^{\sigma} m(k, k'), g),$$

and consequently  $Y(\tau, [B_{k,k',\nu}, P_j]) \in \mathcal{Y}(\langle \xi \rangle^{-1} \tilde{p} h^{\sigma} m(k, k'))$ .

Moreover we have

$$\partial_{x_j} p \partial_{\xi_j} \partial_{\xi_j} p, \partial_{\xi_j} p \partial_{x_j} \partial_{\xi_j} p \in S(\langle \xi \rangle^{-1} \tilde{p} h^{1+\sigma} p, g),$$

hence (3.23) ensures

$$p \odot \partial_{\xi_j} p - \partial_{\xi_j} p \odot p \in S(\langle \xi \rangle^{-1} \tilde{p} h^{1+\sigma} p, g). \quad (6.15)$$



Introducing  $l_n^0 \in S(1, g)$  such that  $(I - L_n^0)B_{k, k', \nu}$  is negligible with  $L_n^0 = l_n^0(x, D)$  and  $\text{supp } l_n^0 \subset \mathcal{B}_n(c_0)$  with  $c_0 < \bar{c}$ , we can write

$$Y(\tau, B_{k, k', \nu})\bar{Y}(\tau, [P_j, i\tau P]) = Y(\tau, B_{k, k', \nu})\bar{Y}(\tau, L_n^0[P_j, i\tau P]) + R_{k, k', \nu} \quad (6.16)$$

with  $\{R_{k, k', \nu}\}_{\nu \in \mathcal{V}}$  negligible. However using

$$\langle h_n p_n \tau \rangle^{-1} | \tau | l_n^0 \in S_n(h^{-1} p^{-1}, g)$$

and (6.15) we find  $p_{j, \nu} \in S_n(\langle \xi \rangle^{-1} \tilde{p} h^\sigma, g)$  such that

$$\langle h_n p_n \tau \rangle^{-1} \tau L_n^0[P_j, i\tau P] = p_{j, \nu}^{\mathcal{W}}(x, D),$$

hence  $\bar{Y}(\tau, L_n^0[P_j, i\tau P]) \in \mathcal{Y}(\langle \xi \rangle^{-1} \tilde{p} h^\sigma)$ . Therefore the right hand side of (6.16) belongs to  $\mathcal{Y}(\langle \xi \rangle^{-1} \tilde{p} h^\sigma m(k, k'))$  and  $\bar{Y}(\tau, [P_j, i\tau P])Y(\tau, B_{k, k', \nu})$  belongs to the same class, i.e. (6.14) gives the desired decomposition (6.10).  $\triangle$

**Corollary 7.3** *Let  $Y_{0, \nu} \in \mathcal{Y}(1)$ . If  $P_j = [iP, x_j]$ , then one has*

$$\tau P_j e^{-i\tau P} Y_{0, \nu} = [e^{-i\tau P} Y_{0, \nu}, x_j] + e^{-i\tau P} (Y_{0, \nu}^+ + \tau Y_{0, \nu}^-) \quad (6.17)$$

with some  $Y_{0, \nu}^+ \in \mathcal{Y}(\langle \xi \rangle^{-1})$  and  $Y_{0, \nu}^- \in \mathcal{Y}(\langle \xi \rangle^{-1} \tilde{p} h^\sigma)$ .

Indeed, using (6.13) to express  $[e^{-i\tau P}, x_j]$  and applying Lemma 6.2 with  $Y_\nu = Y_{0, \nu}$  we can write

$$\begin{aligned} [e^{-i\tau P} Y_{0, \nu}, x_j] &= [e^{-i\tau P}, x_j] Y_{0, \nu} + e^{-i\tau P} [Y_{0, \nu}, x_j] \\ &= -\tau P_j e^{-i\tau P} Y_{0, \nu} - \tau e^{-i\tau P} \bar{Y}(\tau, [P_j, i\tau P]) Y_{0, \nu} + e^{-i\tau P} (Y_\nu^+ + \tau Y_\nu^-). \end{aligned}$$

It remains to remark that the reasoning of the proof of Lemma 6.2 ensures the fact that  $Y_{0, \nu}^+ = Y_\nu^+$  and  $Y_{0, \nu}^- = -\bar{Y}(\tau, [P_j, i\tau P]) Y_{0, \nu} + Y_\nu^-$  belong to the indicated classes.  $\triangle$

## 8 End of the proof of Theorem 2.1

Throughout this section we use the following notation

$$Q_\nu = \text{Op}(e^{i(\tau-t)p} q_\nu) \quad \text{with} \quad q_\nu \in \tilde{S}(\langle \xi \rangle m / \tilde{p}). \quad (7.1)$$

We adopt the convention that the symbol  $(x, \xi) \rightarrow p(x, \xi)$  can be considered as  $(x, \xi, y) \rightarrow p(x, \xi)$ , allowing to define  $q_\nu \partial_{\xi_j} p \in \tilde{S}(m)$  by the formula

$$(q_\nu \partial_{\xi_j} p)(x, \xi, y) = q_\nu(x, \xi, y) \partial_{\xi_j} p(x, \xi). \quad (7.2)$$

**Lemma 8.1** *If  $Q_\nu$  and  $q_\nu \partial_{\xi_j} p$  are as in (7.1), (7.2), then*

$$[Q_\nu, x_j] = (t - \tau) \text{Op}(e^{i(\tau-t)p} q_\nu \partial_{\xi_j} p) + \text{Op}(e^{i(\tau-t)p} i \partial_{\xi_j} q_\nu). \quad (7.3)$$

**Proof.** Since the integral kernel of  $[Q_\nu, x_j]$  is

$$(x, y) \rightarrow (2\pi)^{-d} \int (y_j - x_j) e^{i(x-y)\xi + i(\tau-t)p(x,\xi)} q_\nu(x, \xi, y) d\xi \quad (7.4)$$

and  $(y_j - x_j) e^{i(x-y)\xi} = -i \partial_{\xi_j} e^{i(x-y)\xi}$ , the integration by parts allows to write (7.4) in the form

$$(x, y) \rightarrow (2\pi)^{-d} \int e^{i(x-y)\xi + i(\tau-t)p(x,\xi)} ((t - \tau) \partial_{\xi_j} p q_\nu + i \partial_{\xi_j} q_\nu)(x, \xi, y) d\xi,$$

which gives (7.3).  $\triangle$

The computation of the composition kernel gives

$$\text{Op}(e^{i(\tau-t)p} q_\nu) b_\nu(x, D)^* = \text{Op}(e^{i(\tau-t)p} (q_\nu \bullet b_\nu)) \quad (7.5)$$

with

$$(q_\nu \bullet b_\nu)(x, \xi, y) = (2\pi)^{-d} \int e^{i(\tilde{y}-y)(\xi-\tilde{\xi})} q_\nu(x, \xi, \tilde{y}) \overline{b_\nu(y, \tilde{\xi})}$$

and the usual Taylor development with integrations by parts give

$$q_\nu \bullet b_\nu = q_\nu \bullet_{N'} b_\nu + r_{N', \nu}, \quad (7.6)$$

with

$$(q_\nu \bullet_{N'} b_\nu)(x, \xi, y) = \sum_{|\alpha| < N'} (-i)^{|\alpha|} \partial_y^\alpha q_\nu(x, \xi, y) \overline{\partial_\xi^\alpha b_\nu(y, \xi)} / \alpha!, \quad (7.7)$$

$$r_{N', \nu} = \sum_{|\alpha| = N'} (q_{\alpha, \nu}(z) \bullet \partial_\xi^\alpha b_\nu)|_{z=y},$$

$$q_{\alpha, \nu}(z)(x, \xi, y) = \int_0^1 \partial_y^\alpha q_\nu(x, \xi, z + s(y-z)) \frac{i^{N'} N' ds}{\alpha!}.$$

Then for an arbitrary  $N \in \mathbf{N}$  we can find  $N' = N'(N)$  such that the family of operators  $R_{N', \nu} = \text{Op}(e^{i(\tau-t)p} r_{N', \nu})$  satisfies

$$\|R_{N', \nu}\| \leq C_N h_n^N \langle h_n p_n t \rangle^{C(N)}. \quad (7.8)$$

Denoting  $(q_\nu \bar{b}_\nu^\#)(x, \xi, y) = q_\nu(x, \xi, y) \overline{b_\nu(y, \xi)}$  we can write

$$q_\nu \in \tilde{S}(\tilde{m}), b_\nu \in S(m', g) \Rightarrow q_\nu \bullet_{N'} b_\nu \in \tilde{S}(\tilde{m}m'), q_\nu \bullet_{N'} b_\nu - q_\nu \bar{b}_\nu^\# \in \tilde{S}(\tilde{m}m'h^\sigma).$$

If in particular  $q_\nu \in \tilde{S}(\langle \xi \rangle m / \tilde{p})$ ,  $\partial_{\xi_j} p_0 \in S(\langle \xi \rangle^{-1} \tilde{p}, g)$ , then

$$Q_\nu P_j = \text{Op}(e^{i(\tau-t)p}(q_\nu \bullet \partial_{\xi_j} p_0)), \quad q_\nu \bullet_{N'} \partial_{\xi_j} p_0 \in \tilde{S}(m),$$

$$q_\nu \bullet_{N'} \partial_{\xi_j} p_0 - q_\nu \overline{\partial_{\xi_j} p_0}^\# \in \tilde{S}(h^\sigma m).$$

Since (3.20) ensures  $\partial_{\xi_j} p - \overline{\partial_{\xi_j} p_0} \in S(\langle \xi \rangle^{-1} h^{1+\sigma} p, g) \subset S(\langle \xi \rangle^{-1} h^\sigma \tilde{p}, g)$ ,

$$q_{N', \nu}^- := q_\nu \bullet_{N'} \partial_{\xi_j} p_0 - q_\nu \overline{\partial_{\xi_j} p}^\# \in \tilde{S}(h^\sigma m). \quad (7.9)$$

For  $j = 1, \dots, d$  we introduce

$$p_{(j), \nu}(x, \xi, y) := \left(1 - \frac{\tau}{t}\right) \partial_{\xi_j} p(x, \xi) + \frac{\tau}{t} \partial_{\xi_j} p(y, \xi). \quad (7.10)$$

**Proposition 8.2** *Let  $q_\nu \in \tilde{S}(\langle \xi \rangle m / \tilde{p})$ ,  $Y_{0, \nu} \in \mathcal{Y}(1)$  and  $N_0 \in \mathbf{N}$ . Then we can find  $C(N_0) > 0$  such that*

$$t J_\nu(q p_{(j)}, Y_0) = \sum_{1 \leq k \leq 2} (J_\nu(q_k, Y_k) + t J_\nu(q_{-k}, Y_{-k})) + O\left(h_n^{N_0} \langle h_n p_{nt} \rangle^{C(N_0)}\right) \quad (7.11)$$

where for  $k = 1$  and  $2$  we have

$$q_{k, \nu} \in \tilde{S}(m / \tilde{p}), \quad q_{-k, \nu} \in \tilde{S}(mh^\sigma), \quad Y_{k, \nu} \in \mathcal{Y}(1), \quad Y_{-k, \nu} \in \mathcal{Y}(1). \quad (7.12)$$

**Proof.** By definition of  $p_{(j), \nu}$  we have

$$t \text{Op}(e^{i(\tau-t)p} q_\nu p_{(j), \nu}) = (t - \tau) \text{Op}(e^{i(\tau-t)p} q_\nu \partial_{\xi_j} p) + \tau \text{Op}(e^{i(\tau-t)p} q_\nu \partial_{\xi_j} p^\#) \quad (7.13)$$

and applying (7.3), (7.9), we can write the above expression as

$$[Q_\nu, x_j] + Q_\nu \tau P_j + \text{Op}(e^{i(\tau-t)p} (\tau q_{N', \nu}^- - i \partial_{\xi_j} q_\nu)) + \tau R_{N', \nu}$$

with  $R_{N', \nu}$  satisfying (7.8). Thus  $t J_\nu(q p_{(j)}, Y_0)$  can be written as

$$\begin{aligned} & \text{tr} [Q_\nu, x_j] e^{-i\tau P} Y_{0, \nu} + \text{tr} Q_\nu \tau P_j e^{-i\tau P} Y_{0, \nu} + \\ & J_\nu(-i \partial_{\xi_j} q, Y_0) + \tau J_\nu(q_{N', \nu}^-, Y_0) + O(h_n^{N_0} \langle h_n p_{nt} \rangle^{C(N_0)}). \end{aligned} \quad (7.14)$$

Due to Corollary 6.3, the sum of two first terms in (7.14) equals to

$$\begin{aligned}
& \operatorname{tr} [Q_\nu, x_j] e^{-i\tau P} Y_{0,\nu} + \operatorname{tr} Q_\nu [e^{-i\tau P} Y_{0,\nu}, x_j] + J_\nu(q, Y_0^+) + \tau J_\nu(q, Y_0^-) \\
&= \operatorname{tr} [Q_\nu e^{-i\tau P} Y_{0,\nu}, x_j] + J_\nu(q, Y_0^+) + \tau J_\nu(q, Y_0^-) \\
&= J_\nu(q, Y_0^+) + \tau J_\nu(q, Y_0^-).
\end{aligned} \tag{7.15}$$

Thus we obtain (7.11) with

$$\begin{aligned}
q_{1,\nu} &= i\partial_{\xi_j} q_\nu, & q_{-1,\nu} &= \frac{\tau}{t} q_{N',\nu}^-, & Y_{1,\nu} &= Y_{-1,\nu} = Y_{0,\nu}, \\
q_{2,\nu} &= \langle \bar{\xi}(k_n^+) \rangle^{-1} q_\nu, & Y_{2,\nu} &= \langle \bar{\xi}(k_n^+) \rangle Y_{0,\nu}^+, \\
q_{-2,\nu} &= \langle \bar{\xi}(k_n^+) \rangle^{-1} \tilde{p}_n h_n^\sigma q_\nu, & Y_{-2,\nu} &= \langle \bar{\xi}(k_n^+) \rangle \tilde{p}_n^{-1} h_n^{-\sigma} Y_{0,\nu}^-,
\end{aligned}$$

where according to our convention  $\tilde{p}_n = \tilde{p}_n(\bar{x}(k_n^-), \bar{\xi}(k_n^+))$ .  $\triangle$ .

**Proof of Proposition 6.1.** Let  $\chi_s, \tilde{\chi}_s$  be as in the proof of Proposition 4.2. Then  $q_{0,\nu} \in \tilde{S}(m, g) \implies \tilde{q}_\nu := q_{0,\nu} \chi_s h_n^{-1} p_n^{-1} \in \tilde{S}(m/\tilde{p}, g)$  and

$$tJ_\nu(q_0 \chi_s, Y_0) = J_\nu(\tilde{q}, \tilde{Y}) \text{ with } \tilde{Y}_\nu = h_n p_n t Y_{0,\nu} \in \mathcal{Y}(1).$$

Therefore it suffices to prove the statement of Proposition 6.1 with  $q_{0,\nu} \tilde{\chi}_s$  instead of  $q_{0,\nu}$ .

Further on we assume  $s = 1$  and note that for  $(x, \xi) \in \operatorname{supp} \tilde{\chi}_1$  we have

$$\langle \xi \rangle^{-1} \tilde{p}(x, \xi)/2 \leq |\nabla_\xi p(x, \xi)| \leq 2 \langle \xi \rangle^{-1} \tilde{p}(x, \xi) \tag{7.16}$$

and due to (2.10), for  $(x, \xi, y) \in \operatorname{supp} q_{0,\nu} \tilde{\chi}_1$  we have

$$\begin{aligned}
|\nabla_\xi p(x, \xi)| - \left( \sum_{j=1}^d p_{(j),\nu}^2 \right)^{1/2} &\leq |\nabla_\xi p(x, \xi) - \nabla_\xi p(y, \xi)| \\
&\leq \langle \xi \rangle^{-1} \tilde{p}(x, \xi)/4 \leq |\nabla_\xi p(x, \xi)|/2,
\end{aligned}$$

which implies

$$\left( \sum_{j=1}^d p_{(j),\nu}^2 \right)^{1/2} \geq |\nabla_\xi p(x, \xi)|/2 \geq \langle \xi \rangle^{-1} \tilde{p}(x, \xi)/4. \tag{7.17}$$

Using (7.16), (7.17), we can write  $q_{0,\nu} \tilde{\chi}_1 = \sum_{j=1}^d q_{(j),\nu} p_{(j),\nu}$  with

$$q_{(j),\nu} = q_{0,\nu} \tilde{\chi}_1 p_{(j),\nu} \left( \sum_{j=1}^d p_{(j),\nu}^2 \right)^{-1} \in \tilde{S}(\langle \xi \rangle m/\tilde{p}, g)$$

and we complete the proof applying Proposition 7.2 with  $q_\nu = q_{(j),\nu}$ .

## References

- [1] Chazarain, J.: Spectre d'un hamiltonien quantique et mécanique classique, *Comm. Partial Diff. Eq.* (1980), 595-644.
- [2] Dimassi, M., Sjöstrand, J.: *Spectral asymptotics in the semiclassical limit*, London mathematical Society Lecture Note Series 268, Cambridge University Press, 1999.
- [3] Helffer, B., Robert D., Comportement semi-classique du spectre des hamiltoniens quantiques elliptiques, *Ann. Inst. Fourier, Grenoble* **31** (3) (1981), 169-223.
- [4] Helffer, B., Robert D., Calcul fonctionnel par la transformation de Mellin et opérateurs admissibles, *J. Funct. Anal.* **53** (3) (1983), 246-268.
- [5] Hörmander, L.: The spectral function of an elliptic operator, *Acta Math.* **121** (1968), 173-218.
- [6] Hörmander, L.: *The analysis of linear partial differential operators*, Springer-Verlag, vol. 1, 2, 3, 4, Berlin-Heidelberg-New York-Tokyo, 1983, 1985.
- [7] Ivrii, V.: *Microlocal analysis and precise spectral asymptotics*, Springer-Verlag, Berlin, 1998.
- [8] Ivrii, V.: Sharp spectral asymptotics for operators with irregular coefficients, *Internat. Math. Res. Notices* (2000), 1155–1166.
- [9] Métivier, G.: Valeurs propres des problèmes aux limites irréguliers, *Bull. Soc. Math. France Mem.* **51-52** (1977), 125–219.
- [10] Miyazaki, Y.: A sharp asymptotic remainder estimate for the eigenvalues of operators associated with strongly elliptic sesquilinear forms, *Japan. J. Math.* **15** (1989), 65–97.
- [11] Miyazaki, Y.: The eigenvalue distribution of elliptic operators with Hölder continuous coefficients, *Osaka J. Math.* **28** (1991), 935-973; Part 2, *Osaka J. Math.* **30** (1993), 267–302.
- [12] M. Reed, M. and Simon, B.: *Methods of modern mathematical physics*, vol. I-IV, Academic Press, New York-San Francisco-London, 1972, 1975, 1979.

- [13] Robert, D.: *Autour de l'approximation semi-classique*, Birkhäuser, Boston 1987.
- [14] Zielinski, L.: Asymptotic behaviour of eigenvalues of differential operators with irregular coefficients on a compact manifold, *C. R. Acad. Sci. Paris Sér. I Math.* **310**, (1990), 563–568.
- [15] Zielinski, L.: Asymptotic distribution of eigenvalues for elliptic boundary value problems, *Asymptot. Anal.* **16** (1998), 181–201.
- [16] Zielinski, L.: Asymptotic distribution of eigenvalues of some elliptic operators with intermediate remainder estimates, *Asymptot. Anal.* **17** (1998), 93–120.
- [17] Zielinski, L.: Sharp spectral asymptotics and Weyl formula for elliptic operators with non-smooth coefficients, *Math. Phys. Anal. Geom.* **2** (1999), 291–321; Part 2: *Colloq. Math.* **92** (2002), 1–18.