



Convergence of a finite-volume scheme for a heat equation with a multiplicative Lipschitz noise

Caroline Bauzet, Flore Nabet, Kerstin Schmitz, Aleksandra Zimmermann

► To cite this version:

Caroline Bauzet, Flore Nabet, Kerstin Schmitz, Aleksandra Zimmermann. Convergence of a finite-volume scheme for a heat equation with a multiplicative Lipschitz noise. 2022. hal-03663571v1

HAL Id: hal-03663571

<https://hal.science/hal-03663571v1>

Preprint submitted on 10 May 2022 (v1), last revised 27 Oct 2022 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Convergence of a finite-volume scheme for a heat equation with a multiplicative Lipschitz noise

Caroline Bauzet*, Flore Nabet[†], Kerstin Schmitz[‡],
Aleksandra Zimmermann[‡]

Abstract

We study here the approximation by a finite-volume scheme of a heat equation forced by a Lipschitz continuous multiplicative noise in the sense of Itô. More precisely, we consider a discretization which is semi-implicit in time and a two-point flux approximation scheme (TPFA) in space. Since the nonlinearity in the stochastic integral is not compatible with the weak convergence obtained by the *a priori* estimates, we adapt the method based on the theorem of Prokhorov and on Skorokhod's representation theorem in order to show stochastically strong convergence of the scheme towards the unique variational solution of our parabolic problem.

Keywords: Stochastic heat equation • Multiplicative Lipschitz noise • Finite-volume method • Stochastic compactness method • Variational approach • Convergence analysis.

Mathematics Subject Classification (2020): 60H15 • 35K05 • 65M08.

1 Introduction

Let $\Lambda \subset \mathbb{R}^2$ be a bounded, open, connected and polygonal set. Moreover let $(\Omega, \mathcal{A}, \mathbb{P})$ be a probability space endowed with a right-continuous, complete filtration $(\mathcal{F}_t)_{t \geq 0}$ and let $(W(t))_{t \geq 0}$ be a standard one-dimensional Brownian motion with respect to $(\mathcal{F}_t)_{t \geq 0}$ on $(\Omega, \mathcal{A}, \mathbb{P})$.

For $T > 0$, we consider a nonlinear stochastic heat equation under Neumann boundary

¹Aix Marseille Univ, CNRS, Centrale Marseille, LMA UMR 7031, Marseille, France, caroline.bauzet@univ-amu.fr, +33 4 84 52 56 31

²CMAP, CNRS, Ecole Polytechnique, Institut Polytechnique de Paris, 91128 Palaiseau, France, flore.nabet@polytechnique.edu, +33 1 69 33 45 65

³Universität Duisburg-Essen, Fakultät für Mathematik, Essen, Germany, kerstin.schmitz@uni-due.de, +49 201 183 6130, aleksandra.zimmermann@uni-due.de, +49 201 183 7292

conditions:

$$\begin{aligned} du - \Delta u \, dt &= g(u) \, dW(t), & \text{in } \Omega \times (0, T) \times \Lambda; \\ u(0, \cdot) &= u_0, & \text{in } \Omega \times \Lambda; \\ \nabla u \cdot \mathbf{n} &= 0, & \text{on } \Omega \times (0, T) \times \partial\Lambda; \end{aligned} \tag{1.1}$$

where \mathbf{n} denotes the unit normal vector to $\partial\Lambda$ outward to Λ . We assume the following hypotheses on the data:

H_1 : $u_0 \in L^2(\Omega; H^1(\Lambda))$ is \mathcal{F}_0 -measurable.

H_2 : $g : \mathbb{R} \rightarrow \mathbb{R}$ is a Lipschitz continuous function with Lipschitz constant $L \geq 0$.

H_3 : $g(0) = 0$.

Remark 1.1. Hypothesis H_3 has been made in order to avoid additional technicalities in the proofs. Since H_2 implies $|g(r)| \leq C_L(1 + |r|)$ for all $r \in \mathbb{R}$ and a constant $C_L \geq 0$ only depending on the Lipschitz constant $L \geq 0$ of g and $|g(0)|$, our scheme may be applied without assuming H_3 and also for square integrable, additive noise with appropriate measurability assumptions.

Remark 1.2. We further remark that our results do not depend on the space dimension and may be generalized to polyhedral domains of \mathbb{R}^3 with straightforward calculations.

1.1 Concept of solution and main result

The theoretical framework associated with Problem (1.1) is well established in the literature. Indeed, we can find many existence and uniqueness results for various concepts of solutions associated with this problem such as mild solutions, variational solutions, pathwise solutions and weak solutions. In the present paper we will be interested in the concept of solution as defined below, which we will call a variational solution:

Definition 1.3. *A variational solution to Problem (1.1) is an $(\mathcal{F}_t)_{t \geq 0}$ -adapted stochastic process*

$$u \in L^2(\Omega; C([0, T]; L^2(\Lambda))) \cap L^2(\Omega; L^2(0, T; H^1(\Lambda)))$$

such that for all $t \in [0, T]$,

$$u(t) - u_0 - \int_0^t \Delta u(s) \, ds = \int_0^t g(u(s)) \, dW(s)$$

in $L^2(\Lambda)$ and a.s. in Ω .

The main result of this paper is to propose a finite-volume scheme for the approximation of such a variational solution and to show its stochastically strong convergence by passing to the limit with respect to the time and space discretization parameters. This is stated in the following convergence result:

Theorem 1.4. *Assume that hypotheses H_1 to H_3 hold. Let $(\mathcal{T}_m)_{m \in \mathbb{N}}$ be a sequence of admissible finite-volume meshes of Λ in the sense of Definition 2.1 such that the mesh size h_m tends to 0 and $(N_m)_{m \in \mathbb{N}} \in \mathbb{N}^*$ a sequence of positive numbers which tends to infinity. For a fixed $m \in \mathbb{N}$, let u_{h_m, N_m}^r and u_{h_m, N_m}^l be the finite-volume approximations defined by (2.1), (2.3)-(2.4) with $\mathcal{T} = \mathcal{T}_m$ and $N = N_m$. Then $(u_{h_m, N_m}^r)_{m \in \mathbb{N}}$ and $(u_{h_m, N_m}^l)_{m \in \mathbb{N}}$ converge in $L^2(\Omega; L^2(0, T; L^2(\Lambda)))$ to the variational solution of Problem (1.1) in the sense of Definition 1.3.*

1.2 State of the art

The study of numerical schemes for stochastic partial differential equations (SPDEs) has attracted a lot of attention in the last decades and there exists an extensive literature on this topic. A list of references for the numerical analysis of SPDEs and an overview of the state of the art is given in [16], [2] and [32].

Concerning the theoretical and numerical study of stochastic heat equations, semigroup techniques may be used to construct mild solutions (see, e.g., [14]). However, from the point of view of applications and mathematical modeling, it is often interesting to consider first-order perturbations of the stochastic heat equation and more complicated, nonlinear second order operators, such as the p -Laplacian or the porous medium operator. For these nonlinear SPDEs, the semigroup approach is not available and variational techniques have been developed in [33], [29] and [30].

In the numerical analysis of variational solutions to parabolic SPDEs, spatial discretizations of finite-element type have been frequently used (see, e.g., [8], [11] and the references therein). On the other hand, for stochastic scalar conservation laws, finite-volume schemes have been studied in [5], [4], [6], [26], [31], [19], [3] and [20]. To the best of our knowledge, there are only a few results on finite-volume schemes for parabolic SPDEs. Let us mention the work of [7] where the authors proposed a convergence result of a finite-volume scheme for the approximation of a stochastic heat equation with linear multiplicative noise.

1.3 Aim of the study

In this contribution, we want to extend the finite-volume approximation results in the hyperbolic case to the stochastic heat equation with Lipschitz continuous multiplicative noise. Having applications to nonlinear operators and also to degenerate parabolic-hyperbolic problem with stochastic force in mind for the future, we propose a method for the convergence of the scheme which does not rely on mild solutions and results from semigroup theory. The main technical challenge is then a lack of compactness since we consider a nonlinear multiplicative noise. Indeed, from the *a priori* estimates, we get up to subsequences weak convergence results in several functional spaces for our finite-volume approximations and this mode of convergence is not enough to indentify the weak limit of the nonlinear term in the stochastic integral. Therefore, we first show the convergence towards a martingale solution by adapting the stochastic compactness method based on Skorokhod's representation theorem. Then, using a famous argument of pathwise

uniqueness (see, e.g., [28]), we obtain the stochastically strong convergence result stated in Theorem 1.4.

1.4 Outline of the paper

The paper is organized as follows. The next section concerns the introduction of the finite-volume framework: the definition of an admissible finite-volume mesh on Λ will be stated and the associated notations of discrete unknowns will be given. Then the notions of discrete gradient and discrete H^1 -seminorm will be introduced. In a last subsection, we will introduce our finite-volume scheme together with the associated finite-volume approximations. The remainder of the paper is then devoted to the proof of the convergence of this approximations towards the variational solution of (1.1). To do so, we will prove in Section 3 several stability estimates satisfied by these approximations, but also a boundedness result on the approximation of the stochastic integral. These estimates will allow us to pass the limit in the numerical scheme in Section 4. More precisely, we apply the classical stochastic compactness argument (see, e.g., [10]). By the theorem of Prokhorov, we will get convergence in law (up to subsequences) of our finite-volume approximations. At the cost of a change of probability space, the Skorokhod representation theorem will allow us to obtain almost sure convergence of the proposed finite-volume scheme. Then, a martingale identification argument will help us in order to recover at the limit the desired stochastic integral. In this way, we have shown that our finite-volume scheme converges to a martingale solution of (1.1), i.e., the stochastic basis is not fixed, but enters an unknown in the equation. Next, we show pathwise uniqueness of solutions to (1.1). This, together with a classical argument of Gyöngy and Krylov (see [28]) allows us to deduce convergence in probability of the scheme with respect to the initial stochastic basis.

2 The finite-volume framework

2.1 Admissible finite-volume meshes and notations

In order to perform a finite-volume approximation of the variational solution of Problem (1.1) on $[0, T] \times \Lambda$ we need first of all to set a choice for the temporal and spatial discretization. For the time discretization, let $N \in \mathbb{N}^*$ be given. We define the fixed time step $\Delta t = \frac{T}{N}$ and divide the interval $[0, T]$ in $0 = t_0 < t_1 < \dots < t_N = T$ equidistantly with $t_n = n\Delta t$ for all $n \in \{0, \dots, N-1\}$. For the space discretization, we consider finite-volume admissible meshes in the sense of the following definition.

Definition 2.1. (*Admissible finite-volume mesh*) An admissible finite-volume mesh \mathcal{T} of Λ (see Fig. 1) is given by a family of open polygonal and convex subsets K , called control volumes of \mathcal{T} , satisfying the following properties:

- $\bar{\Lambda} = \bigcup_{K \in \mathcal{T}} \bar{K}$.

- If $K, L \in \mathcal{T}$ with $K \neq L$ then $\text{int } K \cap \text{int } L = \emptyset$.
- If $K, L \in \mathcal{T}$, with $K \neq L$ then either the 1-dimensional Lebesgue measure of $\overline{K} \cap \overline{L}$ is 0 or $\overline{K} \cap \overline{L}$ is the edge of the mesh denoted $\sigma = K|L$ separating the control volumes K and L .
- To each control volume $K \in \mathcal{T}$, we associate a point $x_K \in \overline{K}$ (called the center of K) such that: If $K, L \in \mathcal{T}$ are two neighbouring control volumes the straight line between the centers x_K and x_L is orthogonal to the edge $\sigma = K|L$.




Figure 1: Notations of the mesh \mathcal{T} associated with Λ

Once an admissible finite-volume mesh \mathcal{T} of Λ is fixed, we will use the following notations.

Notations.

- $h = \text{size}(\mathcal{T}) = \sup\{\text{diam}(K) : K \in \mathcal{T}\}$, the mesh size.
- $d_h \in \mathbb{N}$ the number of control volumes $K \in \mathcal{T}$ with $h = \text{size}(\mathcal{T})$.
- λ_2 denotes the two-dimensional Lebesgue measure.
- \mathcal{E} is the set of the edges of the mesh \mathcal{T} and we define $\mathcal{E}_{\text{int}} := \{\sigma \in \mathcal{E} : \sigma \not\subseteq \partial\Lambda\}$, $\mathcal{E}_{\text{ext}} := \{\sigma \in \mathcal{E} : \sigma \subseteq \partial\Lambda\}$.
- For $K \in \mathcal{T}$, \mathcal{E}_K is the set of edges of K and $m_K := \lambda_2(K)$.
- Let $K, L \in \mathcal{T}$ be two neighbouring control volumes. For $\sigma = K|L \in \mathcal{E}_{\text{int}}$, let m_σ be the length of σ and $d_{K|L}$ the distance between x_K and x_L .
- For neighbouring control volumes $K, L \in \mathcal{T}$, we denote the unit vector on the edge $\sigma = K|L$ pointing from K to L by \mathbf{n}_{KL} .
- For $\sigma = K|L \in \mathcal{E}_{\text{int}}$, the diamond D_σ (see Fig. 2) is the open quadrangle whose diagonals are the edge σ and the segment $[x_K, x_L]$. For $\sigma \in \mathcal{E}_{\text{ext}} \cap \mathcal{E}_K$, we define $D_\sigma := K$. Then, $\overline{\Lambda} = \bigcup_{\sigma \in \mathcal{E}} \overline{D_\sigma}$.




Figure 2: Notations on a diamond cell D_σ for $\sigma \in \mathcal{E}_{\text{int}}$

- $m_{D_\sigma} := \lambda_2(D_\sigma)$ is the two-dimensional Lebesgue measure of the diamond D_σ . Note that for $\sigma \in \mathcal{E}_{\text{int}}$, we have $m_{D_\sigma} = \frac{m_\sigma d_{K|L}}{2}$.

Using these notations, we introduce a positive number

$$\text{reg}(\mathcal{T}) = \max \left(\mathcal{N}, \max_{\substack{K \in \mathcal{T} \\ \sigma \in \mathcal{E}_K}} \frac{\text{diam}(K)}{d(x_K, \sigma)} \right)$$

(where \mathcal{N} is the maximum of edges incident to any vertex) that measures the regularity of a given mesh and is useful to perform the convergence analysis of finite-volume schemes. This number should be uniformly bounded when the mesh size tends to 0 for the convergence results to hold.

2.2 Discrete unknowns and piecewise constant functions

From now on and unless otherwise specified, we consider $N \in \mathbb{N}^*$, $\Delta t = \frac{T}{N}$ and \mathcal{T} an admissible finite-volume mesh of Λ in the sense of Definition 2.1 with a mesh size h . For $n \in \{0, \dots, N-1\}$ given, the idea of a finite-volume scheme for the approximation of Problem (1.1) is to associate to each control volume $K \in \mathcal{T}$ and time t_n a discrete unknown value denoted $u_K^n \in \mathbb{R}$, expected to be an approximation of $u(t_n, x_K)$, where u is the variational solution of (1.1). Before presenting the numerical scheme satisfied by the discrete unknowns $\{u_K^n, K \in \mathcal{T}, n \in \{0, \dots, N-1\}\}$, let us introduce some general notations.

For any arbitrary vector $(w_K^n)_{K \in \mathcal{T}} \in \mathbb{R}^{d_h}$ we can define the piecewise constant function $w_h^n : \Lambda \rightarrow \mathbb{R}$ by

$$w_h^n(x) := \sum_{K \in \mathcal{T}} w_K^n \mathbb{1}_K(x), \quad \forall x \in \Lambda.$$

Note that since the mesh \mathcal{T} is fixed, by the continuous embedding

$$(w_K^n)_{K \in \mathcal{T}} \mapsto \sum_{K \in \mathcal{T}} \mathbb{1}_K w_K^n$$

the space \mathbb{R}^{d_h} can be considered as a finite-dimensional subspace of $L^2(\Lambda)$ and we may naturally identify the function and the vector

$$w_h^n \equiv (w_K^n)_{K \in \mathcal{T}} \in \mathbb{R}^{d_h}.$$

Then, knowing for all $n \in \{0, \dots, N\}$ the function w_h^n , we can define the following piecewise constant functions in time and space $w_{h,N}^r, w_{h,N}^l : [0, T] \times \Lambda \rightarrow \mathbb{R}$ by

$$\begin{aligned} w_{h,N}^r(t, x) &:= \sum_{n=0}^{N-1} w_h^{n+1}(x) \mathbb{1}_{[t_n, t_{n+1})}(t) \text{ if } t \in [0, T) \text{ and } w_{h,N}^r(T, x) := w_h^N(x), \\ w_{h,N}^l(t, x) &:= \sum_{n=0}^{N-1} w_h^n(x) \mathbb{1}_{[t_n, t_{n+1})}(t) \text{ if } t \in (0, T] \text{ and } w_{h,N}^l(0, x) := w_h^0(x). \end{aligned} \quad (2.1)$$

As for the piecewise constant function in space, since \mathcal{T} and N are fixed, by the continuous embedding

$$(w_K^n)_{n \in \{0, \dots, N-1\}} \mapsto \sum_{\substack{K \in \mathcal{T} \\ n \in \{0, \dots, N-1\}}} \mathbb{1}_K \mathbb{1}_{(t_n, t_{n+1}]} w_K^n$$

the space $\mathbb{R}^{d_h \times N}$ can be considered as a finite-dimensional subspace of $L^2(0, T; L^2(\Lambda))$ and we may naturally identify

$$\begin{aligned} w_{h,N}^l &\equiv (w_K^n)_{\substack{K \in \mathcal{T} \\ n \in \{0, \dots, N-1\}}} \in \mathbb{R}^{d_h \times N}, \\ w_{h,N}^r &\equiv (w_K^{n+1})_{\substack{K \in \mathcal{T} \\ n \in \{0, \dots, N-1\}}} \in \mathbb{R}^{d_h \times N}. \end{aligned}$$

We can also define the piecewise affine in time and piecewise constant in space reconstruction $\widehat{w}_{h,N} : [0, T] \times \Lambda \rightarrow \mathbb{R}$ by

$$\widehat{w}_{h,N}(t, x) := \sum_{n=0}^{N-1} \mathbb{1}_{[t_n, t_{n+1})}(t) \left(\frac{w_h^{n+1}(x) - w_h^n(x)}{\Delta t} (t - t_n) + w_h^n(x) \right). \quad (2.2)$$

Remark 2.2. Note that in the rest of the paper, when we will consider a time and space function $\alpha : [0, T] \times \Lambda \rightarrow \mathbb{R}$ on all the space Λ (respectively the time interval $[0, T]$) at a fixed time $t \in [0, T]$ (respectively at a fixed $x \in \Lambda$) we will omit the space (respectively time) variable in the notations and write $\alpha(t)$ (respectively $\alpha(x)$) instead of $\alpha(t, \cdot)$ (respectively $\alpha(\cdot, x)$).

2.3 Discrete norms and discrete gradient

Fix $n \in \{0, \dots, N-1\}$ and consider for the remainder of this subsection an arbitrary vector $(w_K^n)_{K \in \mathcal{T}} \in \mathbb{R}^{d_h}$ and use its natural identification with the piecewise constant function in space $w_h^n \equiv (w_K^n)_{K \in \mathcal{T}}$. We introduce in what follows the notions of discrete gradient and discrete norms for such a function w_h^n .

Definition 2.3 (Discrete L^2 -norm). We define the L^2 -norm of $w_h^n \in \mathbb{R}^{d_h}$ as follows

$$\|w_h^n\|_{L^2(\Lambda)} = \left(\sum_{K \in \mathcal{T}} m_K |w_K^n|^2 \right)^{\frac{1}{2}}.$$

Definition 2.4 (Discrete gradient). We define the gradient operator ∇^h that maps scalar fields $w_h^n \in \mathbb{R}^{d_h}$ into vector fields of $(\mathbb{R}^2)^{e_h}$ (where e_h is the number of edges in the mesh \mathcal{T}), we set $\nabla^h w_h^n = (\nabla_\sigma^h w_h^n)_{\sigma \in \mathcal{E}}$ with

$$\nabla_\sigma^h w_h^n := \begin{cases} 2 \frac{w_L^n - w_K^n}{d_{K|L}} \mathbf{n}_{KL}, & \text{if } \sigma = K|L \in \mathcal{E}_{\text{int}}; \\ 0, & \text{if } \sigma \in \mathcal{E}_{\text{ext}}. \end{cases}$$

Definition 2.5 (Discrete H^1 -seminorm). We define the H^1 -seminorm of $w_h^n \in \mathbb{R}^{d_h}$ as follows

$$|w_h^n|_{1,h} := \left(\sum_{\sigma \in \mathcal{E}_{\text{int}}} \frac{m_\sigma}{d_{K|L}} |w_K^n - w_L^n|^2 \right)^{\frac{1}{2}}.$$

Notation. If not marked otherwise, for an edge $\sigma \in \mathcal{E}_{\text{int}}$ we denote the neighbouring control volumes by K and L , i.e., $\sigma = K|L$. In particular we use this notation in sums.

Remark 2.6. Note that in particular,

$$\|\nabla^h w_h^n\|_{(L^2(\Lambda))^2}^2 = \sum_{\sigma \in \mathcal{E}_{\text{int}}} m_{D_\sigma} |\nabla_\sigma^h w_h^n|^2 = 2 \sum_{\sigma \in \mathcal{E}_{\text{int}}} \frac{m_\sigma}{d_{K|L}} |w_K^n - w_L^n|^2 = 2 |w_h^n|_{1,h}^2.$$

Remark 2.7. If we consider another arbitrary vector $\tilde{w}_h^n \equiv (\tilde{w}_K^n)_{K \in \mathcal{T}} \in \mathbb{R}^{d_h}$, then by rearranging the sum we get the following rule of discrete partial integration

$$\sum_{K \in \mathcal{T}} \sum_{\sigma \in \mathcal{E}_K \cap \mathcal{E}_{\text{int}}} \frac{m_\sigma}{d_{K|L}} (w_K^n - w_L^n) \tilde{w}_K^n = \sum_{\sigma \in \mathcal{E}_{\text{int}}} \frac{m_\sigma}{d_{K|L}} (w_K^n - w_L^n) (\tilde{w}_K^n - \tilde{w}_L^n).$$

We have now all the necessary definitions and notations to present the finite-volume scheme studied in this paper. This is the aim of the next subsection.

2.4 The finite-volume scheme

Firstly, we define the vector $u_h^0 \equiv (u_K^0)_{K \in \mathcal{T}} \in \mathbb{R}^{d_h}$ by the discretization of the initial condition u_0 of Problem (1.1) over each control volume:

$$u_K^0 := \frac{1}{m_K} \int_K u_0(x) dx, \quad \forall K \in \mathcal{T}. \quad (2.3)$$

The finite-volume scheme we propose reads, for this given initial \mathcal{F}_0 -measurable random vector $u_h^0 \in \mathbb{R}^{d_h}$:

For any $n \in \{0, \dots, N-1\}$, knowing $u_h^n \equiv (u_K^n)_{K \in \mathcal{T}} \in \mathbb{R}^{d_h}$ we search for $u_h^{n+1} \equiv (u_K^{n+1})_{K \in \mathcal{T}} \in \mathbb{R}^{d_h}$ such that, for almost every $\omega \in \Omega$, the vector u_h^{n+1} is solution to the following random equations

$$\frac{m_K}{\Delta t}(u_K^{n+1} - u_K^n) + \sum_{\sigma \in \mathcal{E}_{\text{int}} \cap \mathcal{E}_K} \frac{m_\sigma}{d_{K|L}}(u_K^{n+1} - u_L^{n+1}) = \frac{m_K}{\Delta t}g(u_K^n)\Delta_{n+1}W, \quad \forall K \in \mathcal{T}, \quad (2.4)$$

where $\Delta_{n+1}W$ denotes the increments of the Brownian motion between t_{n+1} and t_n :

$$\Delta_{n+1}W := W(t_{n+1}) - W(t_n) \text{ for } n \in \{0, \dots, N-1\}.$$

Proposition 2.8 (Existence of a discrete solution). *Assume that hypotheses H_1 to H_3 hold. Let \mathcal{T} be an admissible finite-volume mesh of Λ in the sense of Definition 2.1 with a mesh size h and $N \in \mathbb{N}^*$. Then, there exists a unique solution $(u_h^n)_{1 \leq n \leq N} \in (\mathbb{R}^{d_h})^N$ to Problem (2.4) associated with the initial vector u_h^0 defined by (2.3). Additionally, for any $n \in \{0, \dots, N\}$, u_h^n is a \mathcal{F}_{t_n} -measurable random vector.*

The solution $(u_h^n)_{1 \leq n \leq N} \in (\mathbb{R}^{d_h})^N$ of the scheme (2.3)-(2.4) is then used to build the right and left finite-volume approximations $u_{h,N}^r$ and $u_{h,N}^l$ defined by (2.1) for the variational solution u of Problem (1.1).

Proof. (Proposition 2.8). Set $n \in \{0, \dots, N-1\}$. For $K \in \mathcal{T}$ and a.s. in Ω , note that (2.4) can be rewritten in the following way:

$$m_K(u_K^{n+1} - f_K^n) + \Delta t \sum_{\sigma \in \mathcal{E}_{\text{int}} \cap \mathcal{E}_K} \frac{m_\sigma}{d_{K|L}}(u_K^{n+1} - u_L^{n+1}) = 0, \quad (2.5)$$

where $f_K^n := g(u_K^n)\Delta_{n+1}W + u_K^n$. For $f_h^n \equiv (f_K^n)_{K \in \mathcal{T}} \in \mathbb{R}^{d_h}$ and a.e. $\omega \in \Omega$, we define the functional $J_h^n : \mathbb{R}^{d_h} \rightarrow \mathbb{R}$ by

$$\begin{aligned} J_h^n(w_h) &= \frac{1}{2} \left(\int_{\Lambda} (w_h)^2 dx + \Delta t |w_h|_{1,h}^2 \right) - \int_{\Lambda} w_h f_h^n dx \\ &= \frac{1}{2} \sum_{K \in \mathcal{T}} m_K (w_K)^2 + \frac{\Delta t}{2} \sum_{\sigma \in \mathcal{E}_{\text{int}}} \frac{m_\sigma}{d_{K|L}} (w_K - w_L)^2 - \sum_{K \in \mathcal{T}} m_K w_K f_K^n \end{aligned}$$

for any $w_h \equiv (w_K)_{K \in \mathcal{T}} \in \mathbb{R}^{d_h}$. From a straightforward calculation it is easy to see that J_h^n is strictly convex and coercive. Therefore J_h^n admits a unique minimizer $u_h^{n+1} \in \mathbb{R}^{d_h}$ and the associated sequence $(u_h^n)_{1 \leq n \leq N} \in (\mathbb{R}^{d_h})^N$ is the unique solution of (2.5) a.s. in Ω . If we assume that u_h^n is \mathcal{F}_{t_n} -measurable, then by the properties of the Brownian motion, f_h^n is $\mathcal{F}_{t_{n+1}}$ -measurable and consequently the random variable $\omega \mapsto J_h^n(w_h)(\omega)$ is $\mathcal{F}_{t_{n+1}}$ -measurable for any $w_h \in \mathbb{R}^{d_h}$. In particular, $\omega \mapsto u_h^{n+1}(\omega) = \min_{w_h \in \mathbb{R}^{d_h}} J_h^n(w_h)(\omega)$ is $\mathcal{F}_{t_{n+1}}$ -measurable. Then, it follows by iteration, that for a given, \mathcal{F}_0 -measurable random variable $u_h^0 \in \mathbb{R}^{d_h}$, for any $n \in \{0, \dots, N-1\}$ there exists a $\mathcal{F}_{t_{n+1}}$ -measurable function $u_h^{n+1} \in \mathbb{R}^{d_h}$ such that $(u_h^n)_{1 \leq n \leq N} \in (\mathbb{R}^{d_h})^N$ is solution to Problem (2.4) associated with the initial vector u_h^0 which concludes the proof. \square

3 Stability estimates

We will derive in this section several stability estimates satisfied by the discrete solution $(u_h^n)_{1 \leq n \leq N} \in (\mathbb{R}^{d_h})^N$ of the scheme (2.3)-(2.4) given by Proposition 2.8, and also by the associated right and left finite-volume approximations $u_{h,N}^r$ and $u_{h,N}^l$ defined by (2.1).

3.1 Bounds on the finite-volume approximations

We start by giving a bound on the discrete initial data.

Lemma 3.1. *Let u_0 be a given function satisfying assumption H_1 . Then, the associated discrete initial data $u_h^0 \in \mathbb{R}^{d_h}$ defined by (2.3) satisfies \mathbb{P} -a.s. in Ω ,*

$$\|u_h^0\|_{L^2(\Lambda)} \leq \|u_0\|_{L^2(\Lambda)}.$$

The proof is a direct consequence of the definition of u_h^0 and the Cauchy-Schwarz inequality.

We can now give the bounds on the discrete solutions which is one of the key points of the proof of the convergence theorem.

Proposition 3.2 (Bounds on the discrete solutions). *There exist constants $C_1, C_2 > 0$, not depending on the discretization parameters $N \in \mathbb{N}^*$ and h , such that*

$$\begin{aligned} \sup_{n \in \{1, \dots, N\}} \mathbb{E} \left[\|u_h^n\|_{L^2(\Lambda)}^2 \right] &\leq C_1, \\ \mathbb{E} \left[\|u_h^n\|_{L^2(\Lambda)}^2 \right] + \mathbb{E} \left[\sum_{k=0}^{n-1} \|u_h^{k+1} - u_h^k\|_{L^2(\Lambda)}^2 \right] + 2\Delta t \sum_{k=0}^{n-1} \mathbb{E} [|u_h^{k+1}|_{1,h}^2] &\leq C_2, \quad \forall n \in \{1, \dots, N\}. \end{aligned}$$

Proof. We fix $n \in \{1, \dots, N\}$. For any $k \in \{0, \dots, n-1\}$ we multiply the numerical scheme (2.4) with u_K^{k+1} and sum over $K \in \mathcal{T}$ to obtain

$$\begin{aligned} &\sum_{K \in \mathcal{T}} \frac{m_K}{\Delta t} (u_K^{k+1} - u_K^k) u_K^{k+1} + \sum_{K \in \mathcal{T}} \sum_{\sigma \in \mathcal{E}_{\text{int}} \cap \mathcal{E}_K} \frac{m_\sigma}{d_{K|L}} (u_K^{k+1} - u_L^{k+1}) u_K^{k+1} \\ &= \sum_{K \in \mathcal{T}} \frac{m_K}{\Delta t} g(u_K^k) u_K^k \Delta_{k+1} W + \sum_{K \in \mathcal{T}} \frac{m_K}{\Delta t} g(u_K^k) (u_K^{k+1} - u_K^k) \Delta_{k+1} W. \end{aligned} \tag{3.1}$$

We consider the terms separately: For the first term on the left-hand side we find

$$\sum_{K \in \mathcal{T}} \frac{m_K}{\Delta t} (u_K^{k+1} - u_K^k) u_K^{k+1} = \frac{1}{2} \sum_{K \in \mathcal{T}} \frac{m_K}{\Delta t} (|u_K^{k+1}|^2 - |u_K^k|^2 + |u_K^{k+1} - u_K^k|^2).$$

For the second term on the left-hand side of (3.1), we obtain by using Remark 2.7

$$\sum_{K \in \mathcal{T}} \sum_{\sigma \in \mathcal{E}_{\text{int}} \cap \mathcal{E}_K} \frac{m_\sigma}{d_{K|L}} (u_K^{k+1} - u_L^{k+1}) u_K^{k+1} = \sum_{\sigma \in \mathcal{E}_{\text{int}}} \frac{m_\sigma}{d_{K|L}} |u_K^{k+1} - u_L^{k+1}|^2.$$

Taking expectation in (3.1), the first expression on the right-hand side of (3.1) vanishes, since

$$\mathbb{E} [g(u_K^k) u_K^k \Delta_{k+1} W] = 0.$$

In the second term we apply Young's inequality in order to keep all necessary terms. Then, taking expectation and using the Itô isometry we obtain

$$\begin{aligned} \mathbb{E} [g(u_K^k)(u_K^{k+1} - u_K^k) \Delta_{k+1} W] &\leq \mathbb{E} [|g(u_K^k) \Delta_{k+1} W|^2] + \frac{1}{4} \mathbb{E} [|u_K^{k+1} - u_K^k|^2] \\ &\leq \Delta t \mathbb{E} [|g(u_K^k)|^2] + \frac{1}{4} \mathbb{E} [|u_K^{k+1} - u_K^k|^2] \end{aligned}$$

for any $K \in \mathcal{T}$. Altogether we find

$$\begin{aligned} &\frac{1}{2\Delta t} \int_{\Lambda} \mathbb{E} [|u_h^{k+1}|^2 - |u_h^k|^2] dx + \frac{1}{4\Delta t} \int_{\Lambda} \mathbb{E} [|u_h^{k+1} - u_h^k|^2] dx + \mathbb{E} [|u_h^{k+1}|_{1,h}^2] \\ &\leq \int_{\Lambda} \mathbb{E} [|g(u_h^k)|^2] dx. \end{aligned}$$

Summing over $k \in \{0, \dots, n-1\}$ and multiplying with $2\Delta t$ we obtain

$$\begin{aligned} &\mathbb{E} [\|u_h^n\|_{L^2(\Lambda)}^2 - \|u_h^0\|_{L^2(\Lambda)}^2] + \frac{1}{2} \sum_{k=0}^{n-1} \mathbb{E} [\|u_h^{k+1} - u_h^k\|_{L^2(\Lambda)}^2] + 2\Delta t \sum_{k=0}^{n-1} \mathbb{E} [|u_h^{k+1}|_{1,h}^2] \\ &\leq 2\Delta t \sum_{k=0}^{n-1} \mathbb{E} [\|g(u_h^k)\|_{L^2(\Lambda)}^2]. \end{aligned} \tag{3.2}$$

Since g is Lipschitz continuous, $g(0) = 0$ and the second and third term in (3.2) are nonnegative, there arises

$$\mathbb{E} [\|u_h^n\|_{L^2(\Lambda)}^2] \leq \mathbb{E} [\|u_h^0\|_{L^2(\Lambda)}^2] + 2L^2 \Delta t \sum_{k=0}^{n-1} \mathbb{E} [\|u_h^k\|_{L^2(\Lambda)}^2].$$

Applying the discrete Gronwall lemma yields

$$\mathbb{E} [\|u_h^n\|_{L^2(\Lambda)}^2] \leq (1 + 2L^2 T) \mathbb{E} [\|u_h^0\|_{L^2(\Lambda)}^2] e^{2L^2 T}. \tag{3.3}$$

From (3.3) and Lemma 3.1 we may conclude that there exists a constant $C_1 > 0$ such that

$$\sup_{n \in \{1, \dots, N\}} \mathbb{E} [\|u_h^n\|_{L^2(\Lambda)}^2] \leq C_1$$

and since g is Lipschitz continuous with $g(0) = 0$ also

$$2\Delta t \sum_{k=0}^{n-1} \mathbb{E} [\|g(u_h^k)\|_{L^2(\Lambda)}^2] \leq 2L^2 \Delta t \sum_{k=0}^{n-1} \mathbb{E} [\|u_h^k\|_{L^2(\Lambda)}^2] \leq 2L^2 T C_1 \tag{3.4}$$

for all $n \in \{1, \dots, N\}$. From (3.2), Lemma 3.1 and (3.4) it now follows that

$$\begin{aligned} & \mathbb{E} \left[\|u_h^n\|_{L^2(\Lambda)}^2 \right] + \frac{1}{2} \sum_{k=0}^{n-1} \mathbb{E} \left[\|u_h^{k+1} - u_h^k\|_{L^2(\Lambda)}^2 \right] + 2\Delta t \sum_{k=0}^{n-1} \mathbb{E} [|u_h^{k+1}|_{1,h}^2] \\ & \leq \mathbb{E} \left[\|u_0\|_{L^2(\Lambda)}^2 \right] + 2L^2TC_1 := C_2 \end{aligned}$$

for all $n \in \{1, \dots, N\}$. \square

We are now interested in the bounds on the right and left finite-volume approximations defined by (2.1). As a direct consequence of Proposition 3.2 we get a $L^2(\Omega; L^2(0, T; L^2(\Lambda)))$ -bound on these approximations.

Lemma 3.3. *The sequences $(u_{h,N}^r)_{h,N}$ and $(u_{h,N}^l)_{h,N}$ are bounded independently of the discretization parameters $N \in \mathbb{N}^*$ and h in $L^2(\Omega; L^2(0, T; L^2(\Lambda)))$.*

Thanks to Proposition 3.2 we can also obtain a $L^2(\Omega; L^2(0, T; L^2(\Lambda)))$ -bound on the discrete gradients of the finite-volume approximations.

Lemma 3.4. *There exist constants $K_1, K_2 \geq 0$ not depending on the discretization parameters $N \in \mathbb{N}^*$ and h such that*

$$\int_0^T \mathbb{E} [|u_{h,N}^r|_{1,h}^2] dt \leq K_1 \quad (3.5)$$

and

$$\int_0^T \mathbb{E} [|u_{h,N}^l|_{1,h}^2] dt \leq K_2. \quad (3.6)$$

Proof.

$$\int_0^T \mathbb{E} [|u_{h,N}^r|_{1,h}^2] dt = \Delta t \sum_{k=0}^{N-1} \mathbb{E} [|u_h^{k+1}|_{1,h}^2]$$

and therefore (3.5) follows directly from Proposition 3.2. Using the definition of $u_{h,N}^l$ and (3.5), we get

$$\int_0^T \mathbb{E} [|u_{h,N}^l|_{1,h}^2] dt \leq \Delta t \mathbb{E} [|u_h^0|_{1,h}^2] + \Delta t \sum_{k=0}^{N-1} \mathbb{E} [|u_h^{k+1}|_{1,h}^2] \leq \Delta t \mathbb{E} [|u_h^0|_{1,h}^2] + K_1.$$

Since u_0 is assumed to be in $L^2(\Omega; H^1(\Lambda))$, from [24, Lemma 9.4], it follows that there exists $C_\Lambda \geq 0$ depending on the mesh regularity $\text{reg}(\mathcal{T})$ such that,

$$\mathbb{E} [|u_h^0|_{1,h}^2] \leq C_\Lambda \mathbb{E} [\|\nabla u_0\|_{L^2(\Lambda)}^2]$$

and therefore (3.6) follows. \square

We end this section by a bound on the discrete solution which will be useful for obtaining the time translate estimate and the bounds on the Gagliardo seminorm.

Lemma 3.5. *There exists a constant $K_3 \geq 0$ independent of the discretization parameters $N \in \mathbb{N}^*$ and h , such that*

$$\mathbb{E} \left[\max_{n \in \{0, \dots, N\}} \|u_h^n\|_{L^2(\Lambda)}^2 \right] \leq K_3.$$

Proof. For $N \in \mathbb{N}$, we choose an arbitrary $k \in \{0, \dots, N-1\}$ and an arbitrary $K \in \mathcal{T}$. Testing the implicit scheme (2.4) with u_K^{k+1} yields after taking the sum over $K \in \mathcal{T}$ and applying Remark 2.7

$$\sum_{K \in \mathcal{T}} \frac{m_K}{\Delta t} (u_K^{k+1} - u_K^k) u_K^{k+1} + \sum_{\sigma \in \mathcal{E}_{\text{int}}} \frac{m_\sigma}{d_{K|L}} |u_K^{k+1} - u_L^{k+1}|^2 = \sum_{K \in \mathcal{T}} \frac{m_K}{\Delta t} g(u_K^k) u_K^{k+1} \Delta_{k+1} W.$$

This provides by Cauchy-Schwarz and Young's inequalities

$$\begin{aligned} & \frac{1}{2} \left(\|u_h^{k+1}\|_{L^2(\Lambda)}^2 - \|u_h^k\|_{L^2(\Lambda)}^2 + \|u_h^{k+1} - u_h^k\|_{L^2(\Lambda)}^2 \right) \\ & \leq \left(\int_{t_k}^{t_{k+1}} g(u_h^k) dW(s), u_h^{k+1} - u_h^k \right)_{L^2(\Lambda)} + \left(\int_{t_k}^{t_{k+1}} g(u_h^k) dW(s), u_h^k \right)_{L^2(\Lambda)} \\ & \leq \frac{1}{2} \left\| \int_{t_k}^{t_{k+1}} g(u_h^k) dW(s) \right\|_{L^2(\Lambda)}^2 + \frac{1}{2} \|u_h^{k+1} - u_h^k\|_{L^2(\Lambda)}^2 + \left(\int_{t_k}^{t_{k+1}} g(u_h^k) dW(s), u_h^k \right)_{L^2(\Lambda)}. \end{aligned}$$

We obtain

$$\|u_h^{k+1}\|_{L^2(\Lambda)}^2 - \|u_h^k\|_{L^2(\Lambda)}^2 \leq \left\| \int_{t_k}^{t_{k+1}} g(u_h^k) dW(s) \right\|_{L^2(\Lambda)}^2 + 2 \left(\int_{t_k}^{t_{k+1}} g(u_h^k) dW(s), u_h^k \right)_{L^2(\Lambda)}.$$

For $n \in \{1, \dots, N\}$ fixed, we sum over $k = \{0, \dots, n-1\}$ to obtain

$$\|u_h^n\|_{L^2(\Lambda)}^2 \leq \sum_{k=0}^{n-1} \left\| \int_{t_k}^{t_{k+1}} g(u_h^k) dW(s) \right\|_{L^2(\Lambda)}^2 + 2 \sum_{k=0}^{n-1} \left(\int_{t_k}^{t_{k+1}} g(u_h^k) dW(s), u_h^k \right)_{L^2(\Lambda)} + \|u_h^0\|_{L^2(\Lambda)}^2.$$

From this, taking the maximum over $n \in \{1, \dots, N\}$ first and then the expectation there

arises by applying Itô isometry since u_h^k is \mathcal{F}_{t_k} -measurable

$$\begin{aligned}
& \mathbb{E} \left[\max_{n=1, \dots, N} \|u_h^n\|_{L^2(\Lambda)}^2 \right] \\
& \leq \mathbb{E} \left[\sum_{k=0}^{N-1} \left\| \int_{t_k}^{t_{k+1}} g(u_h^k) dW(s) \right\|_{L^2(\Lambda)}^2 \right] \\
& \quad + 2 \mathbb{E} \left[\max_{n=1, \dots, N} \sum_{k=0}^{n-1} \left(\int_{t_k}^{t_{k+1}} g(u_h^k) dW(s), u_h^k \right)_{L^2(\Lambda)} \right] + \mathbb{E} \left[\|u_h^0\|_{L^2(\Lambda)}^2 \right] \quad (3.7) \\
& \leq \sum_{k=0}^{N-1} \int_{t_k}^{t_{k+1}} \mathbb{E} \left[\|g(u_h^k)\|_{L^2(\Lambda)}^2 \right] ds \\
& \quad + 2 \mathbb{E} \left[\max_{n=1, \dots, N} \sum_{k=0}^{n-1} \int_{t_k}^{t_{k+1}} (g(u_h^k), u_h^k)_{L^2(\Lambda)} dW(s) \right] + \mathbb{E} \left[\|u_h^0\|_{L^2(\Lambda)}^2 \right].
\end{aligned}$$

We can estimate the second term by the Burkholder-Davis-Gundy inequality

$$\begin{aligned}
& 2 \mathbb{E} \left[\max_{n=1, \dots, N} \sum_{k=0}^{n-1} \int_{t_k}^{t_{k+1}} (g(u_h^k), u_h^k)_{L^2(\Lambda)} dW(s) \right] \\
& \leq 2 \mathbb{E} \left[\sup_{t \in [0, T]} \left| \int_0^t (g(u_{h,N}^l), u_{h,N}^l)_{L^2(\Lambda)} dW(s) \right| \right] \\
& \leq 2C_B \mathbb{E} \left[\left(\int_0^T |(g(u_{h,N}^l), u_{h,N}^l)_{L^2(\Lambda)}|^2 ds \right)^{\frac{1}{2}} \right].
\end{aligned}$$

Now we apply Cauchy-Schwarz and Young's inequalities (with $\alpha > 0$)

$$\begin{aligned}
& 2C_B \mathbb{E} \left[\left(\int_0^T |(g(u_{h,N}^l), u_{h,N}^l)_{L^2(\Lambda)}|^2 ds \right)^{\frac{1}{2}} \right] \\
& \leq 2C_B \mathbb{E} \left[\left(\sup_{t \in [0, T]} \|u_{h,N}^l(t)\|_{L^2(\Lambda)}^2 \int_0^T \|g(u_{h,N}^l)\|_{L^2(\Lambda)}^2 ds \right)^{\frac{1}{2}} \right] \\
& \leq 2C_B \mathbb{E} \left[\frac{\alpha}{2} \sup_{t \in [0, T]} \|u_{h,N}^l(t)\|_{L^2(\Lambda)}^2 + \frac{1}{2\alpha} \int_0^T \|g(u_{h,N}^l)\|_{L^2(\Lambda)}^2 ds \right] \\
& \leq C_B \alpha \mathbb{E} \left[\max_{n=1, \dots, N} \|u_h^n\|_{L^2(\Lambda)}^2 \right] + \frac{C_B L^2}{\alpha} \mathbb{E} \left[\int_0^T \|u_{h,N}^l\|_{L^2(\Lambda)}^2 ds \right].
\end{aligned}$$

Plugging the above estimate in (3.7), we arrive at

$$\begin{aligned}
\mathbb{E} \left[\max_{n=1, \dots, N} \|u_h^n\|_{L^2(\Lambda)}^2 \right] & \leq \left(L^2 + \frac{C_B L^2}{\alpha} \right) \int_0^T \mathbb{E} \left[\|u_{h,N}^l\|_{L^2(\Lambda)}^2 \right] ds \\
& \quad + C_B \alpha \mathbb{E} \left[\max_{n=1, \dots, N} \|u_h^n\|_{L^2(\Lambda)}^2 \right] + \mathbb{E} \left[\|u_h^0\|_{L^2(\Lambda)}^2 \right].
\end{aligned}$$

Choosing $\alpha > 0$ such that $1 - C_B\alpha > 0$, we find a constant $C(\alpha, L) > 0$ such that

$$\mathbb{E} \left[\max_{n=1, \dots, N} \|u_h^n\|_{L^2(\Lambda)}^2 \right] \leq C(\alpha, L) \left(\int_0^T \mathbb{E} \left[\|u_{h,N}^l\|_{L^2(\Lambda)}^2 \right] ds + \mathbb{E} \left[\|u_h^0\|_{L^2(\Lambda)}^2 \right] \right).$$

Now, the assertion follows by Lemmas 3.1 and 3.3. \square

3.2 Time and space translate estimates

For the stochastic compactness argument in Subsection 4.2, we need a uniform bound on $(u_{h,N}^l)_{h,N}$ in the spaces $L^2(\Omega; L^2(0, T; W^{\alpha,2}(\Lambda)))$ and $L^2(\Omega; W^{\alpha,2}(0, T; L^2(\Lambda)))$ for $\alpha \in (0, \frac{1}{2})$. In order to prove the bound in $L^2(\Omega; L^2(0, T; W^{\alpha,2}(\Lambda)))$, we establish a uniform estimate on the space translates of $(u_{h,N}^l)_{h,N}$ in Lemma 3.8. The proof of the bound in $L^2(\Omega; W^{\alpha,2}(0, T; L^2(\Lambda)))$ is more complicated. Lemma 3.6 is a technical result for the proof of Lemma 3.7, where we show a uniform estimate on time translates of $(u_{h,N}^l - M_{h,N})_{h,N}$. Thanks to Lemma 3.7, we may conclude a uniform bound on $(u_{h,N}^l - M_{h,N})_{h,N}$ in $L^2(\Omega; W^{\alpha,2}(0, T; L^2(\Lambda)))$ in Lemma 3.10. Then the desired bound on $(u_{h,N}^l)_{h,N}$ is obtained in Lemma 3.11 by using the additional information that $(M_{h,N})_{h,N}$ is bounded in $L^2(\Omega; W^{\alpha,2}(0, T; L^2(\Lambda)))$.

Lemma 3.6. *There exists a constant $K_4 > 0$, independent of the discretization parameters $N \in \mathbb{N}^*$ and h , such that for all $\tau \in (0, T)$ there holds*

$$\mathbb{E} \left[\int_0^{T-\tau} \|u_{h,N}^l(t+\tau) - M_{h,N}^l(t+\tau) - (u_{h,N}^l(t) - M_{h,N}^l(t))\|_{L^2(\Lambda)}^2 dt \right] \leq K_4\tau, \quad (3.8)$$

where $M_{h,N}^l$ is defined for any $(t, x) \in [0, T] \times \Lambda$ by

$$M_{h,N}^l(t, x) := \int_0^{t_n} g(u_{h,N}^l(s, x)) dW(s) \text{ if } t \in [t_n, t_{n+1}) \text{ with } n \in \{0, \dots, N-1\}.$$

Proof. Let $\tau \in (0, T)$ be fixed. In the following, we set $\varphi_h^N(t, x) := u_{h,N}^l(t, x) - M_{h,N}^l(t, x)$ for $t \in [0, T], x \in \Lambda$. Note that by definition $M_K^n := M_{h,N}^l(t_n, x_K)$ and $\varphi_K^n := u_K^n - M_K^n$ for $n \in \{0, \dots, N\}, K \in \mathcal{T}$. For $t \in (0, T - \tau)$ let $n_0(t), n_1(t) \in \{0, \dots, N-1\}$ be the unique nonnegative integer satisfying

$$n_0(t)\Delta t \leq t < (n_0(t) + 1)\Delta t \quad \text{and} \quad n_1(t)\Delta t \leq t + \tau < (n_1(t) + 1)\Delta t.$$

There holds \mathbb{P} -a.s in Ω

$$\begin{aligned} \int_0^{T-\tau} \|u_{h,N}^l(t+\tau) - M_{h,N}^l(t+\tau) - (u_{h,N}^l(t) - M_{h,N}^l(t))\|_{L^2(\Lambda)}^2 dt \\ = \int_0^{T-\tau} \sum_{K \in \mathcal{T}} m_K |\varphi_K^{n_1(t)} - \varphi_K^{n_0(t)}|^2 dt =: \int_0^{T-\tau} A(t) dt. \end{aligned}$$

If $n_0(t) = n_1(t)$ holds, we have $A(t) = 0$. So we only consider $t \in (0, T - \tau)$ with $n_1(t) > n_0(t)$. We get

$$\begin{aligned}
A(t) &= \sum_{K \in \mathcal{T}} m_K(\varphi_K^{n_1(t)} - \varphi_K^{n_0(t)})(\varphi_K^{n_1(t)} - \varphi_K^{n_0(t)}) \\
&= \sum_{K \in \mathcal{T}} m_K(\varphi_K^{n_1(t)} - \varphi_K^{n_0(t)}) \sum_{n=n_0(t)}^{n_1(t)-1} (\varphi_K^{n+1} - \varphi_K^n) \\
&= \sum_{K \in \mathcal{T}} m_K(\varphi_K^{n_1(t)} - \varphi_K^{n_0(t)}) \sum_{n=0}^{N-1} \chi_{n+1}(t, t + \tau)(\varphi_K^{n+1} - \varphi_K^n) \\
&= \sum_{n=0}^{N-1} \chi_{n+1}(t, t + \tau) \sum_{K \in \mathcal{T}} (\varphi_K^{n_1(t)} - \varphi_K^{n_0(t)}) m_K(\varphi_K^{n+1} - \varphi_K^n),
\end{aligned}$$

where $\chi_n(t, t + \tau) = 1$ if $n\Delta t \in [t, t + \tau)$ and $\chi_n(t, t + \tau) = 0$ otherwise. Using [\(2.4\)](#) yields, we obtain

$$A(t) = -\Delta t \sum_{n=0}^{N-1} \chi_{n+1}(t, t + \tau) \sum_{K \in \mathcal{T}} (\varphi_K^{n_1(t)} - \varphi_K^{n_0(t)}) \sum_{\sigma \in \mathcal{E}_{\text{int}} \cap \mathcal{E}_K} \frac{m_\sigma}{d_{K|L}} (u_K^{n+1} - u_L^{n+1}).$$

Rearranging the sum in the same way as for discrete partial integration (see [Remark 2.7](#)), using the definition of φ_h^N and the notation $u_K^{N,l} := u_{h,N}^l(x_K)$ for $K \in \mathcal{T}$ there arises

$$\begin{aligned}
A(t) &= -\Delta t \sum_{n=0}^{N-1} \chi_{n+1}(t, t + \tau) \sum_{\sigma \in \mathcal{E}_{\text{int}}} \frac{m_\sigma}{d_{K|L}} (u_K^{n+1} - u_L^{n+1}) \left(\varphi_K^{n_1(t)} - \varphi_L^{n_1(t)} - (\varphi_K^{n_0(t)} - \varphi_L^{n_0(t)}) \right) \\
&= -\Delta t \sum_{n=0}^{N-1} \chi_{n+1}(t, t + \tau) \sum_{\sigma \in \mathcal{E}_{\text{int}}} \frac{m_\sigma}{d_{K|L}} (u_K^{n+1} - u_L^{n+1}) [u_K^{n_1(t)} - u_L^{n_1(t)} - u_K^{n_0(t)} + u_L^{n_0(t)}] \\
&\quad + \Delta t \sum_{n=0}^{N-1} \chi_{n+1}(t, t + \tau) \sum_{\sigma \in \mathcal{E}_{\text{int}}} \frac{m_\sigma}{d_{K|L}} (u_K^{n+1} - u_L^{n+1}) \int_{n_0(t)\Delta t}^{n_1(t)\Delta t} \left(g(u_K^{N,l}) - g(u_L^{N,l}) \right) dW(s) \\
&=: A_1(t) + A_2(t),
\end{aligned}$$

By Cauchy-Schwarz and Young inequalities we get

$$\begin{aligned}
A_1(t) &\leq \frac{\Delta t}{2} \sum_{n=0}^{N-1} \chi_{n+1}(t, t + \tau) |u_h^{n+1}|_{1,h}^2 + \frac{\Delta t}{2} \sum_{n=0}^{N-1} \chi_{n+1}(t, t + \tau) |u_h^{n_1(t)} - u_h^{n_0(t)}|_{1,h}^2 \\
&\leq \frac{\Delta t}{2} \sum_{n=0}^{N-1} \chi_{n+1}(t, t + \tau) |u_h^{n+1}|_{1,h}^2 + \Delta t \sum_{n=0}^{N-1} \chi_{n+1}(t, t + \tau) (|u_h^{n_1(t)}|_{1,h}^2 + |u_h^{n_0(t)}|_{1,h}^2).
\end{aligned}$$

Consequently,

$$\mathbb{E} \left[\int_0^{T-\tau} A_1(t) dt \right] \leq I_1 + I_2$$

where

$$I_1 = \frac{1}{2} \int_0^{T-\tau} \sum_{n=0}^{N-1} \chi_{n+1}(t, t+\tau) \mathbb{E} [\Delta t |u_h^{n+1}|_{1,h}^2] dt$$

and

$$I_2 = \int_0^{T-\tau} \sum_{n=0}^{N-1} \chi_{n+1}(t, t+\tau) \Delta t \mathbb{E} [|u_h^{n_0(t)}|_{1,h}^2 + |u_h^{n_1(t)}|_{1,h}^2] dt.$$

Since

$$\begin{aligned} \chi_{n+1}(t, t+\tau) = 1 &\Leftrightarrow (n+1)\Delta t \in [t, t+\tau) \\ &\Leftrightarrow t - \tau \leq (n+1)\Delta t - \tau < t \leq (n+1)\Delta t, \end{aligned}$$

there arises

$$\int_0^{T-\tau} \chi_{n+1}(t, t+\tau) dt = \int_{(n+1)\Delta t - \tau}^{(n+1)\Delta t} 1 dt = \tau. \quad (3.9)$$

Using this and (3.5), we have

$$I_1 = \frac{\tau}{2} \mathbb{E} \left[\int_0^T |u_{h,N}^r|_{1,h}^2 ds \right] \leq \frac{K_1 \tau}{2}.$$

Now we write $I_2 = I_{2,1} + I_{2,2}$ where,

$$\begin{aligned} I_{2,1} &= \int_0^{T-\tau} \sum_{n=0}^{N-1} \chi_{n+1}(t, t+\tau) \Delta t \mathbb{E} [|u_h^{n_0(t)}|_{1,h}^2] dt, \\ I_{2,2} &= \int_0^{T-\tau} \sum_{n=0}^{N-1} \chi_{n+1}(t, t+\tau) \Delta t \mathbb{E} [|u_h^{n_1(t)}|_{1,h}^2] dt. \end{aligned}$$

We note that, for any $m \in \{0, \dots, N-1\}$, if $t \in [t_m, t_{m+1})$ then the definition of n_0 implies $n_0(t) = m$ and therefore

$$I_{2,1} \leq \sum_{m=0}^{N-1} \left(\int_{t_m}^{t_{m+1}} \sum_{n=0}^{N-1} \chi_{n+1}(t, t+\tau) dt \right) \Delta t \mathbb{E} [|u_h^m|_{1,h}^2]$$

From [27, Lemma 6.2] it follows that

$$\int_{t_m}^{t_{m+1}} \sum_{n=0}^{N-1} \chi_{n+1}(t, t+\tau) dt \leq \tau, \quad \forall m \in \{0, \dots, N-1\}.$$

Therefore, thanks to Lemma 3.4 one has

$$I_{2,1} \leq \tau \Delta t \sum_{m=0}^{N-1} \mathbb{E} [|u_h^m|_{1,h}^2] = \tau \int_0^T \mathbb{E} [|u_{h,N}^l|_{1,h}^2] dt \leq K_2 \tau.$$

Analogously, for any $m \in \{0, \dots, N-1\}$, if $t \in [t_m - \tau, t_{m+1} - \tau)$, then the definition of n_1 implies $n_1(t) = m$ and therefore

$$I_{2,2} \leq \sum_{m=0}^{N-1} \Delta t \mathbb{E} [|u_h^m|_{1,h}^2] \int_{t_m - \tau}^{t_{m+1} - \tau} \sum_{n=0}^{N-1} \chi_{n+1}(t, t + \tau) dt \leq K_2 \tau$$

by [27, Lemma 6.2] and Lemma 3.4, where $\chi_{n+1}(t, t + \tau) = 0$ for $t < 0$. Combining the previous estimates we arrive at

$$\mathbb{E} \left[\int_0^{T-\tau} A_1(t) dt \right] \leq \left(\frac{K_1}{2} + 2K_2 \right) \tau. \quad (3.10)$$

Now we consider A_2 . Applying Young's inequality we find

$$\begin{aligned} A_2(t) &\leq \frac{\Delta t}{2} \sum_{n=0}^{N-1} \chi_{n+1}(t, t + \tau) \sum_{\sigma \in \mathcal{E}_{\text{int}}} \frac{m_\sigma}{d_{K|L}} (u_K^{n+1} - u_L^{n+1})^2 \\ &\quad + \frac{\Delta t}{2} \sum_{n=0}^{N-1} \chi_{n+1}(t, t + \tau) \sum_{\sigma \in \mathcal{E}_{\text{int}}} \frac{m_\sigma}{d_{K|L}} \left| \int_{n_0(t)\Delta t}^{n_1(t)\Delta t} \left(g(u_K^{N,l}) - g(u_L^{N,l}) \right) dW(s) \right|^2 \\ &=: A_{2,1}(t) + A_{2,2}(t). \end{aligned}$$

We have

$$\mathbb{E} \left[\int_0^{T-\tau} A_{2,1}(t) dt \right] = \frac{\Delta t}{2} \mathbb{E} \left[\sum_{n=0}^{N-1} |u_h^{n+1}|_{1,h}^2 \int_0^{T-\tau} \chi_{n+1}(t, t + \tau) dt \right].$$

By (3.9) and Lemma 3.4 we may conclude

$$\mathbb{E} \left[\int_0^{T-\tau} A_{2,1}(t) dt \right] = \frac{\tau}{2} \mathbb{E} \left[\int_0^T |u_{h,N}^r|_{1,h}^2 ds \right] \leq \frac{K_1 \tau}{2}. \quad (3.11)$$

For the study of the term $A_{2,2}$ we recall the notation $u_K^{N,l} := u_{h,N}^l(x_K)$ for $K \in \mathcal{T}$. From the Burkholder-Davis-Gundy inequality [10, Proposition 2.3.8] it follows that there exists a constant $C_B \geq 0$ such that, for any $t \in (0, T - \tau)$ with $n_0(t) < n_1(t)$,

$$\begin{aligned} \mathbb{E} \left[\left| \int_{n_0(t)\Delta t}^{n_1(t)\Delta t} \left(g(u_K^{N,l}) - g(u_L^{N,l}) \right) dW(s) \right|^2 \right] &\leq 4 \mathbb{E} \left[\sup_{r \in [0, T]} \left| \int_0^r g(u_K^{N,l}) - g(u_L^{N,l}) dW(s) \right|^2 \right] \\ &\leq 4C_B \mathbb{E} \left[\int_0^T |g(u_K^{N,l}) - g(u_L^{N,l})|^2 ds \right]. \end{aligned}$$

Therefore,

$$\begin{aligned}
& \mathbb{E} \left[\int_0^{T-\tau} A_{2,2}(t) dt \right] \\
&= \frac{\Delta t}{2} \int_0^{T-\tau} \sum_{n=0}^{N-1} \chi_{n+1}(t, t+\tau) \sum_{\sigma \in \mathcal{E}_{\text{int}}} \frac{m_\sigma}{d_{K|L}} \mathbb{E} \left[\left| \int_{n_0(t)\Delta t}^{n_1(t)\Delta t} \left(g(u_K^{N,l}) - g(u_L^{N,l}) \right) dW(s) \right|^2 \right] dt \\
&\leq 4C_B \frac{\Delta t}{2} \int_0^{T-\tau} \sum_{n=0}^{N-1} \chi_{n+1}(t, t+\tau) \sum_{\sigma \in \mathcal{E}_{\text{int}}} \frac{m_\sigma}{d_{K|L}} \mathbb{E} \left[\int_0^T |g(u_K^{N,l}) - g(u_L^{N,l})|^2 ds \right] dt \\
&\leq 2C_B L^2 \Delta t \int_0^{T-\tau} \sum_{n=0}^{N-1} \chi_{n+1}(t, t+\tau) \sum_{\sigma \in \mathcal{E}_{\text{int}}} \frac{m_\sigma}{d_{K|L}} \mathbb{E} \left[\int_0^T |u_K^{N,l} - u_L^{N,l}|^2 ds \right] dt \\
&= 2C_B L^2 \Delta t \int_0^{T-\tau} \sum_{n=0}^{N-1} \chi_{n+1}(t, t+\tau) dt \int_0^T \mathbb{E} [|u_{h,N}^l|_{1,h}^2] ds.
\end{aligned}$$

Because of (3.9) there holds

$$\int_0^{T-\tau} \sum_{n=0}^{N-1} \chi_{n+1}(t, t+\tau) dt = \sum_{n=0}^{N-1} \tau = N\tau.$$

Therefore from (3.6) it follows

$$\mathbb{E} \left[\int_0^{T-\tau} A_{2,2}(t) dt \right] \leq 2C_B L^2 T \tau K_2. \quad (3.12)$$

Finally (3.8) holds from (3.10), (3.11) and (3.12). \square

Lemma 3.7. *There exists a constant $K_5 > 0$, independent of the discretization parameters $N \in \mathbb{N}^*$ and h , such that for all $\tau \in (0, T)$ there holds*

$$\mathbb{E} \left[\int_0^{T-\tau} \|u_{h,N}^l(t+\tau) - M_{h,N}(t+\tau) - (u_{h,N}^l(t) - M_{h,N}(t))\|_{L^2(\Lambda)}^2 dt \right] \leq K_5 \tau, \quad (3.13)$$

where for any $(t, x) \in [0, T] \times \Lambda$

$$M_{h,N}(t, x) := \int_0^t g(u_{h,N}^l(s, x)) dW(s). \quad (3.14)$$

Proof. Let $0 < \tau < T$. We can write using the fact that for any $a, b, c \in \mathbb{R}$, $|a + b + c|^2 \leq$

$$3(|a|^2 + |b|^2 + |c|^2)$$

$$\begin{aligned} & \mathbb{E} \left[\int_0^{T-\tau} \|u_{h,N}^l(t+\tau) - M_{h,N}(t+\tau) - (u_{h,N}^l(t) - M_{h,N}(t))\|_{L^2(\Lambda)}^2 dt \right] \\ & \leq 3\mathbb{E} \left[\int_0^{T-\tau} \|u_{h,N}^l(t+\tau) - M_{h,N}^l(t+\tau) - (u_{h,N}^l(t) - M_{h,N}^l(t))\|_{L^2(\Lambda)}^2 dt \right] \\ & \quad + 3\mathbb{E} \left[\int_0^{T-\tau} \|M_{h,N}(t+\tau) - M_{h,N}(t)\|_{L^2(\Lambda)}^2 dt \right] \\ & \quad + 3\mathbb{E} \left[\int_0^{T-\tau} \|M_{h,N}^l(t+\tau) - M_{h,N}^l(t)\|_{L^2(\Lambda)}^2 dt \right] \\ & =: 3(I_1 + I_2 + I_3). \end{aligned}$$

From Lemma 3.6 we know that $I_1 \leq K_4\tau$.

By the Itô isometry and Lemma 3.5 we get

$$\begin{aligned} I_2 &= \int_0^{T-\tau} \int_t^{t+\tau} \mathbb{E} \left[\|g(u_{h,N}^l)\|_{L^2(\Lambda)}^2 \right] ds dt \\ &\leq L^2 \int_0^{T-\tau} \int_t^{t+\tau} \mathbb{E} \left[\sup_{t \in [0, T]} \|u_{h,N}^l\|_{L^2(\Lambda)}^2 \right] ds dt \leq L^2 K_3 T \tau. \end{aligned}$$

For $t \in [0, T]$, let $n_0(t), n_1(t) \in \{0, \dots, N-1\}$ be defined by

$$n_0(t)\Delta t \leq t < (n_0(t) + 1)\Delta t \quad \text{and} \quad n_1(t)\Delta t \leq t + \tau < (n_1(t) + 1)\Delta t.$$

From the Itô isometry and Lemma 3.5 there arises

$$I_3 = \int_0^{T-\tau} \int_{n_0(t)\Delta t}^{n_1(t)\Delta t} \mathbb{E} \left[\|g(u_{h,N}^l)\|_{L^2(\Lambda)}^2 \right] ds dt \leq \int_0^{T-\tau} L^2 K_3 \int_{n_0(t)\Delta t}^{n_1(t)\Delta t} 1 ds dt.$$

Similarly to the proof of Lemma 3.6, let $\chi_n(t, t+\tau) = 1$ for $n \in \mathbb{N}$ if $n\Delta t \in (t, t+\tau]$ and 0 otherwise. Taking (3.9) into account, we can continue the above estimate by

$$\begin{aligned} I_3 &= \int_0^{T-\tau} L^2 K_3 \sum_{n=0}^{N-1} \chi_{n+1}(t, t+\tau) \int_{t_n}^{t_{n+1}} 1 ds dt \\ &= L^2 K_3 \Delta t \sum_{n=0}^{N-1} \int_0^{T-\tau} \chi_{n+1}(t, t+\tau) dt \leq L^2 K_3 T \tau, \end{aligned}$$

and the assertion follows. \square

We conclude by giving the estimation of space translate. We do not give the proof here as it is similar to the one given in [24, Theorem 10.3].

Lemma 3.8. *Let $\bar{u}_{h,N}^l$ be $d\mathbb{P} \otimes dt \otimes dx$ -a.s. defined by $\bar{u}_{h,N}^l = u_{h,N}^l$ on $\Omega \times (0, T) \times \Lambda$ and $\bar{u}_{h,N}^l = 0$ on $\Omega \times (\mathbb{R}^3 \setminus ((0, T) \times \Lambda))$. Then there exists a constant $C \geq 0$, only depending on Λ , such that for all $\eta \in \mathbb{R}^2$ and almost every $t \in [0, T]$ and \mathbb{P} -a.s in Ω*

$$\int_{\mathbb{R}^2} |\bar{u}_{h,N}^l(t, x+\eta) - \bar{u}_{h,N}^l(t, x)|^2 dx \leq C|\eta| \left(|u_{h,N}^l(t)|_{1,h}^2 + \|u_{h,N}^l(t)\|_{L^2(\Lambda)}^2 \right).$$

3.3 Bound on the Gagliardo seminorm

Our target is to apply [25, Theorem 2.1] which claims, that for any $\alpha \in (0, \frac{1}{2})$

$$W^{\alpha,2}(0, T; L^2(\Lambda)) \cap L^2(0, T; W^{\alpha,2}(\Lambda))$$

is compactly embedded in

$$L^2(0, T; L^2(\Lambda)).$$

We denote by $[\cdot]_{W^{\alpha,2}(\Lambda)}$ the Gagliardo seminorm such that for any function $w : \Lambda \rightarrow \mathbb{R}$ one has,

$$[w]_{W^{\alpha,2}(\Lambda)} = \left(\int_{\Lambda} \int_{\Lambda} \frac{|w(x) - w(y)|^2}{|x - y|^{2+2\alpha}} dx dy \right)^{\frac{1}{2}}.$$

Lemma 3.9. *For any fixed $\alpha \in (0, \frac{1}{2})$, the sequences $(u_{h,N}^l)_{h,N}$ and $(u_{h,N}^r)_{h,N}$ are bounded in $L^2(\Omega; L^2(0, T; W^{\alpha,2}(\Lambda)))$ independently of the discretization parameters $N \in \mathbb{N}^*$ and h .*

Proof. We fix $0 < \alpha < \frac{1}{2}$ and $R > 0$. For almost every $t \in (0, T)$, thanks to Lemma 3.8

$$\begin{aligned} & \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{|\bar{u}_{h,N}^l(t, x) - \bar{u}_{h,N}^l(t, y)|^2}{|x - y|^{2+2\alpha}} dx dy \\ &= \int_{|\eta| > R} \int_{\mathbb{R}^2} \frac{|\bar{u}_{h,N}^l(t, x) - \bar{u}_{h,N}^l(t, x + \eta)|^2}{|\eta|^{2(1+\alpha)}} dx d\eta \\ & \quad + \int_{|\eta| < R} \int_{\mathbb{R}^2} \frac{|\bar{u}_{h,N}^l(t, x) - \bar{u}_{h,N}^l(t, x + \eta)|^2}{|\eta|^{2(1+\alpha)}} dx d\eta \\ &\leq 4 \|\bar{u}_{h,N}^l(t)\|_{L^2(\mathbb{R}^2)}^2 \int_{|\eta| > R} |\eta|^{-2(1+\alpha)} d\eta \\ & \quad + C \left(|u_{h,N}^l(t)|_{1,h}^2 + \|u_{h,N}^l(t)\|_{L^2(\Lambda)}^2 \right) \int_{|\eta| < R} |\eta|^{-2(1+\alpha)+1} d\eta \\ &= 4 \|u_{h,N}^l(t)\|_{L^2(\Lambda)}^2 \int_0^{2\pi} \int_R^\infty r^{-2(1+\alpha)} r dr d\varphi \\ & \quad + C \left(|u_{h,N}^l(t)|_{1,h}^2 + \|u_{h,N}^l(t)\|_{L^2(\Lambda)}^2 \right) \int_0^{2\pi} \int_0^R r^{-2\alpha-1} r dr d\varphi \\ &=: \tilde{C}_1 \|u_{h,N}^l(t)\|_{L^2(\Lambda)}^2 + \tilde{C}_2 |u_{h,N}^l(t)|_{1,h}^2 \end{aligned}$$

where the constants $\tilde{C}_1, \tilde{C}_2 \geq 0$ only depend on Λ and $R > 0$. Consequently we have

$$\begin{aligned} \int_0^T [u_{h,N}^l(t)]_{W^{\alpha,2}(\Lambda)}^2 dt &\leq \int_0^T \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \frac{|\bar{u}_{h,N}^l(t, x) - \bar{u}_{h,N}^l(t, y)|^2}{|x - y|^{2+2\alpha}} dx dy dt \\ &\leq \int_0^T \left(\tilde{C}_1 \|u_{h,N}^l(t)\|_{L^2(\Lambda)}^2 + \tilde{C}_2 |u_{h,N}^l(t)|_{1,h}^2 \right) dt. \end{aligned}$$

Therefore thanks to Lemmas 3.4 and 3.5 we get

$$\begin{aligned}\mathbb{E} [\|u_{h,N}^l\|_{L^2(0,T;W^{\alpha,2}(\Lambda))}] &= \mathbb{E} \left[\int_0^T (\|u_{h,N}^l(t)\|_{L^2(\Lambda)} + [u_{h,N}^l(t)]_{W^{\alpha,2}(\Lambda)})^2 dt \right] \\ &\leq 2(1 + \tilde{C}_1) \int_0^T \mathbb{E} [\|u_{h,N}^l(t)\|_{L^2(\Lambda)}^2] dt + 2\tilde{C}_2 \int_0^T \mathbb{E} [|u_{h,N}^l(t)|_{1,h}^2] dt \\ &\leq 2TK_3(1 + \tilde{C}_1) + 2\tilde{C}_2K_2.\end{aligned}$$

So $(u_{h,N}^l)_{h,N}$ is bounded in $L^2(\Omega; L^2(0, T; W^{\alpha,2}(\Lambda)))$ and also $(u_{h,N}^r)_{h,N}$ with similar arguments. \square

In order to establish the $L^2(\Omega; W^{\alpha,2}(0, T; L^2(\Lambda)))$ -bound on the discrete solutions we give the following auxiliary result:

Lemma 3.10. *For any fixed $\alpha \in (0, \frac{1}{2})$, the sequence $(u_{h,N}^l - M_{h,N})_{h,N}$ defined by (3.14) is bounded in $L^2(\Omega; W^{\alpha,2}(0, T; L^2(\Lambda)))$ independently of the discretization parameters $N \in \mathbb{N}^*$ and h .*

Proof. For any $x \in \Lambda$, let $\bar{\varphi}_{h,N}(t, x) := u_{h,N}^l(t, x) - M_{h,N}(t, x)$ for $t \in [0, T]$ and $\bar{\varphi}_{h,N}(t, x) = 0$ for $t \in \mathbb{R} \setminus [0, T]$. We get by applying Fubini's theorem

$$\begin{aligned}\mathbb{E} \left[\int_0^T \int_0^T \frac{\|\bar{\varphi}_{h,N}(s) - \bar{\varphi}_{h,N}(t)\|_{L^2(\Lambda)}^2}{|t - s|^{1+2\alpha}} ds dt \right] \\ = \mathbb{E} \left[\int_0^T \int_{-t}^{T-t} \frac{\|\bar{\varphi}_{h,N}(t + \tau) - \bar{\varphi}_{h,N}(t)\|_{L^2(\Lambda)}^2}{|\tau|^{1+2\alpha}} d\tau dt \right] \\ = \mathbb{E} \left[\int_0^T \int_{-\tau}^{T-\tau} \frac{\|\bar{\varphi}_{h,N}(t + \tau) - \bar{\varphi}_{h,N}(t)\|_{L^2(\Lambda)}^2}{|\tau|^{1+2\alpha}} dt d\tau \right].\end{aligned}$$

Note, that for any $x \in \Lambda$, $\bar{\varphi}_{h,N}(t, x) = 0$ for $t \in (-\infty, 0)$, so we get by (3.13)

$$\begin{aligned}\mathbb{E} \left[\int_0^T \int_{-\tau}^{T-\tau} \frac{\|\bar{\varphi}_{h,N}(t + \tau) - \bar{\varphi}_{h,N}(t)\|_{L^2(\Lambda)}^2}{|\tau|^{1+2\alpha}} dt d\tau \right] \\ = \int_0^T |\tau|^{-1-2\alpha} \int_{-\tau}^0 \mathbb{E} [\|\bar{\varphi}_{h,N}(t + \tau)\|_{L^2(\Lambda)}^2] dt d\tau \\ + \int_0^T |\tau|^{-1-2\alpha} \int_0^{T-\tau} \mathbb{E} [\|\bar{\varphi}_{h,N}(t + \tau) - \bar{\varphi}_{h,N}(t)\|_{L^2(\Lambda)}^2] dt d\tau \\ \leq \int_0^T |\tau|^{-1-2\alpha} \int_0^\tau \mathbb{E} [\|\bar{\varphi}_{h,N}(s)\|_{L^2(\Lambda)}^2] ds d\tau + K_5 \int_0^T |\tau|^{-2\alpha} d\tau.\end{aligned}\tag{3.15}$$

Let $s \in (0, \tau)$ be arbitrary. By Itô isometry there holds

$$\mathbb{E} [\|\bar{\varphi}_{h,N}(s)\|_{L^2(\Lambda)}^2] \leq 2\mathbb{E} [\|u_{h,N}^l(s)\|_{L^2(\Lambda)}^2] + 2\mathbb{E} \left[\int_0^s \|g(u_{h,N}^l(t))\|_{L^2(\Lambda)}^2 dt \right].$$

Using Lemma 3.5, we obtain

$$\begin{aligned} \sup_{s \in [0, T]} \mathbb{E} \left[\|\bar{\varphi}_{h,N}(s)\|_{L^2(\Lambda)}^2 \right] &\leq 2 \max_{n=0, \dots, N-1} \mathbb{E} \left[\|u_h^n\|_{L^2(\Lambda)}^2 \right] + 2L^2 \mathbb{E} \left[\int_0^T \|u_{h,N}^l(t)\|_{L^2(\Lambda)}^2 dt \right] \\ &\leq 2(1 + L^2 T) K_3 =: C. \end{aligned}$$

Using this estimate in (3.15) yields

$$\begin{aligned} \mathbb{E} \left[\int_0^T \int_0^T \frac{\|\bar{\varphi}_{h,N}(s) - \bar{\varphi}_{h,N}(t)\|_{L^2(\Lambda)}^2}{|t - s|^{1+2\alpha}} ds dt \right] \\ \leq \int_0^T \tau^{-2\alpha-1} \tau C d\tau + K_5 \int_0^T \tau^{-2\alpha} d\tau \leq \tilde{C} \int_0^T \tau^{-2\alpha} d\tau \end{aligned}$$

and the above integral is finite for $\alpha \in (0, \frac{1}{2})$. \square

Lemma 3.11. *For any fixed $\alpha \in (0, \frac{1}{2})$, the sequence $(u_{h,N}^l)_{h,N}$ is bounded in $L^2(\Omega; W^{\alpha,2}(0, T; L^2(\Lambda)))$ independently of the discretization parameters $N \in \mathbb{N}^*$ and h .*

Proof. From Lemma 3.10 we know that $(u_{h,N}^l - M_{h,N})_{h,N}$ is bounded in $L^2(\Omega; W^{\alpha,2}(0, T; L^2(\Lambda)))$. Since

$$\int_0^T \mathbb{E} \left[\|u_{h,N}^l\|_{L^2(\Lambda)}^2 \right] dt \leq T \mathbb{E} \left[\sup_{t \in [0, T]} \|u_{h,N}^l\|_{L^2(\Lambda)}^2 \right] \leq T K_3$$

by Lemma 3.5, we get by applying [25, Lemma 2.1] that $(M_{h,N})_{h,N}$ is bounded in $L^2(\Omega; W^{\alpha,2}(0, T; L^2(\Lambda)))$. \square

4 Convergence of the finite-volume scheme

We now have all the necessary material to pass to the limit in the numerical scheme.

In the sequel, for $m \in \mathbb{N}^*$, let $(\mathcal{T}_m)_m$ be a sequence of admissible meshes of Λ in the sense of Definition 2.1 such that the mesh size h_m tends to 0 when m tends to $+\infty$ and let $(N_m)_m \subset \mathbb{N}$ be a sequence with $\lim_{m \rightarrow +\infty} N_m = +\infty$ and $\Delta t_m := \frac{T}{N_m}$.

For the sake of simplicity we shall use the notations $\mathcal{T} = \mathcal{T}_m$, $h = \text{size}(\mathcal{T}_m)$, $\Delta t = \Delta t_m$ and $N = N_m$ when the m -dependency is not useful for the understanding of the reader.

4.1 Weak convergence of finite-volume approximations

First, thanks to the bounds on the discrete solutions, we obtain the following weak convergences.

Lemma 4.1. *There exist not relabeled subsequences of $(u_{h,N}^r)_m$ and of $(u_{h,N}^l)_m$ respectively and a function $u \in L^2(\Omega; L^2(0, T; H^1(\Lambda)))$ such that*

$$u_{h,N}^l \rightharpoonup u \text{ and } u_{h,N}^r \rightharpoonup u$$

for $m \rightarrow +\infty$ weakly in $L^2(\Omega; L^2(0, T; L^2(\Lambda)))$.

Proof. From Lemma 3.3 it follows that the sequences $(u_{h,N}^r)_m$, $(u_{h,N}^l)_m$ respectively are bounded in $L^2(\Omega; L^2(0, T; L^2(\Lambda)))$, thus, up to a not relabeled subsequence, weakly convergent in $L^2(\Omega; L^2(0, T; L^2(\Lambda)))$ towards possibly distinct elements u , \tilde{u} respectively. Moreover, from Lemma 3.4 and Remark 2.6, it follows that

$$\|\nabla^h u_{h,N}^r\|_{L^2(\Omega \times (0,T) \times \Lambda)}^2 \leq 2K_1.$$

Consequently, there exists $\chi \in L^2(\Omega; L^2(0, T; L^2(\Lambda)))$ such that, passing to a not relabeled subsequence if necessary, $\nabla^h u_{h,N}^r \rightharpoonup \chi$ weakly in $L^2(\Omega; L^2(0, T; L^2(\Lambda)))$ for $m \rightarrow +\infty$. With similar arguments as in [23, Lemma 2] and [24, Theorem 14.3] we get the additional regularity $u \in L^2(\Omega; L^2(0, T; H^1(\Lambda)))$ and $\chi = \nabla u$. Since, by Proposition 3.2,

$$\mathbb{E} \left[\|u_{h,N}^r - u_{h,N}^l\|_{L^2(0,T;L^2(\Lambda))}^2 \right] = \Delta t \mathbb{E} \left[\sum_{n=0}^{N-1} \|u_h^{n+1} - u_h^n\|_{L^2(\Lambda)}^2 \right] \leq C_2 \Delta t, \quad (4.1)$$

it follows that $(u_{h,N}^r - u_{h,N}^l)_m$ converges to 0 strongly in $L^2(\Omega; L^2(0, T; L^2(\Lambda)))$ when $m \rightarrow +\infty$, hence also weakly and therefore $u = \tilde{u}$. \square

Our aim is to show that u is the unique solution to (1.1). But weak convergence is not enough to pass to the limit in the nonlinear diffusion term of our finite-volume scheme. Therefore we will apply the method of stochastic compactness.

4.2 The stochastic compactness argument

From Lemmas 3.9 and 3.11 we get immediately the following bounds.

Lemma 4.2. *For any fixed $\alpha \in (0, \frac{1}{2})$, $(u_{h,N}^l)_m$ is bounded in $L^2(\Omega; L^2(0, T; W^{\alpha,2}(\Lambda))) \cap L^2(\Omega; W^{\alpha,2}(0, T; L^2(\Lambda)))$ independently of the parameter $m \in \mathbb{N}^*$.*

In the following, for a random variable X defined on a probability space $(\Omega, \mathcal{A}, \mathbb{P})$ the law of X will be denoted by $\mathbb{P} \circ X^{-1}$.

Lemma 4.3. *The sequence of laws $(\mathbb{P} \circ (u_{h,N}^l)^{-1})_m$ on $L^2(0, T; L^2(\Lambda))$ is tight.*

Proof. For better readability we define $V := L^2(0, T; L^2(\Lambda))$ and $\mathcal{W} := W^{\alpha,2}(0, T; L^2(\Lambda)) \cap L^2(0, T; W^{\alpha,2}(\Lambda))$. By [25, Theorem 2.1] we know that \mathcal{W} is compactly embedded in V . Let $\varepsilon > 0$ be arbitrary. For any $R > 0$ the ball $B_{\mathcal{W}}(0, R) := \{v \in \mathcal{W} : \|v\|_{\mathcal{W}} \leq R\}$ is compact in V . There holds

$$[\mathbb{P} \circ (u_{h,N}^l)^{-1}](B_{\mathcal{W}}(0, R)) = 1 - [\mathbb{P} \circ (u_{h,N}^l)^{-1}](B_{\mathcal{W}}(0, R)^c) = 1 - \int_{\{\|u_{h,N}^l\|_{\mathcal{W}} > R\}} 1 \, d\mathbb{P},$$

where we can estimate further

$$\int_{\{\|u_{h,N}^l\|_{\mathcal{W}} > R\}} 1 \, d\mathbb{P} \leq \frac{1}{R^2} \int_{\{\|u_{h,N}^l\|_{\mathcal{W}} > R\}} \|u_{h,N}^l\|_{\mathcal{W}}^2 \, d\mathbb{P} \leq \frac{1}{R^2} \int_{\Omega} \|u_{h,N}^l\|_{\mathcal{W}}^2 \, d\mathbb{P} \leq \frac{C}{R^2}$$

for a constant $C > 0$. In the last inequality we used that $(u_{h,N}^l)_{h,N}$ is, thanks to Lemma 4.2, bounded in $L^2(\Omega; \mathcal{W})$. It follows

$$[\mathbb{P} \circ (u_{h,N}^l)^{-1}](B_{\mathcal{W}}(0, R)) \geq 1 - \frac{C}{R^2}.$$

If we choose an appropriate R , the assertion follows. \square

For the next lemmas, we recall that the initial value of Problem (1.1) denoted u_0 is \mathcal{F}_0 -measurable and belongs to $L^2(\Omega; H^1(\Lambda))$. Moreover, its spatial discretization denoted u_h^0 is defined by (2.3). In the following, we will write $(W(t))_{t \geq 0} =: W$, whenever the t -dependence is not relevant for the argumentation.

In order to apply Skorokhod theorem and to prove the almost surely convergence, we begin by proving the convergence in law.

Lemma 4.4. *For $m \in \mathbb{N}^*$ we consider the sequence of random vectors*

$$Y_m = ((u_{h_m, N_m}^l, u_{h_m, N_m}^r - u_{h_m, N_m}^l, W, u_{h_m}^0)$$

with values in

$$\mathcal{X} := L^2(0, T; L^2(\Lambda)) \times L^2(0, T; L^2(\Lambda)) \times C([0, T]) \times L^2(\Lambda).$$

There exists a not relabeled subsequence of $(Y_m)_m$ converging in law, i.e., there exists a probability measure $\mu_\infty = (\mu_\infty^1, \delta_0, \mathbb{P} \circ W^{-1}, \mathbb{P} \circ (u_0)^{-1})$ on \mathcal{X} such that

$$\mathbb{E}[f(Y_m)] \xrightarrow{m \rightarrow +\infty} \int_{\mathcal{X}} f d\mu_\infty$$

for all bounded, continuous functions $f : \mathcal{X} \rightarrow \mathbb{R}$.

Proof. We recall that a subsequence of $(Y_m)_m$ converges in law if and only if all its components converge in law. The tightness of laws of $(u_{h_m, N_m}^l)_m$ was shown in Lemma 4.3. Then, from the Theorem of Prokhorov (see [9, Theorem 5.1]) it follows that, passing to a not relabeled subsequence if necessary, $(u_{h_m, N_m}^l)_m$ converges in law towards a probability measure μ_∞^1 defined on $L^2(0, T; L^2(\Lambda))$.

Clearly, as a constant sequence, the Brownian motion W converges in law towards $\mathbb{P} \circ W^{-1}$. Since $(u_{h_m}^0)_m$ converges to u_0 in $L^2(\Lambda)$ for $m \rightarrow +\infty$ a.s. in Ω (see [1, Proposition 3.5]), it follows that $(u_{h_m}^0)_m$ converges in law towards $\mathbb{P} \circ (u_0)^{-1}$. From (4.1) it follows that $(u_{h_m, N_m}^r - u_{h_m, N_m}^l)_m$ converges to 0 for $m \rightarrow +\infty$ in $L^2(\Omega; L^2(0, T; L^2(\Lambda)))$ and this convergence implies for all bounded, continuous functions $f : L^2(0, T; L^2(\Lambda)) \rightarrow \mathbb{R}$

$$\int_{L^2(0, T; L^2(\Lambda))} f d(\mathbb{P} \circ (u_{h_m, N_m}^r - u_{h_m, N_m}^l)^{-1}) = \mathbb{E}[f(u_{h_m, N_m}^r - u_{h_m, N_m}^l)] \xrightarrow{m \rightarrow +\infty} \mathbb{E}[f(0)],$$

hence the convergence in law of $(u_{h_m, N_m}^r - u_{h_m, N_m}^l)_m$ towards δ_0 . \square

Thanks to Lemma 4.4 we may apply the Skorokhod representation Theorem (see [9, Theorem 6.7]): There exists a probability space $(\Omega', \mathcal{A}', \mathbb{P}')$ and random variables

$$Y'_m = (v_m, z_m, B_m, v_m^0), \quad u_\infty, W_\infty, v_0$$

with

$$\begin{aligned} \mathbb{P}' \circ (Y'_m)^{-1} &= \mathbb{P} \circ (Y_m)^{-1} \text{ for all } m \in \mathbb{N}, \\ \mathbb{P}' \circ (v_0)^{-1} &= \mathbb{P} \circ (u_0)^{-1}, \\ \mathbb{P}' \circ (u_\infty)^{-1} &= \mu_\infty^1, \\ \mathbb{P}' \circ (W_\infty)^{-1} &= \mathbb{P} \circ W^{-1} \end{aligned}$$

and such that

$$\begin{aligned} v_m &\xrightarrow{m \rightarrow +\infty} u_\infty && \text{in } L^2(0, T; L^2(\Lambda)), \mathbb{P}'\text{-a.s. in } \Omega' \\ z_m &\xrightarrow{m \rightarrow +\infty} 0 && \text{in } L^2(0, T; L^2(\Lambda)), \mathbb{P}'\text{-a.s. in } \Omega' \\ B_m &\xrightarrow{m \rightarrow +\infty} W_\infty && \text{in } C([0, T]), \mathbb{P}'\text{-a.s. in } \Omega' \\ v_m^0 &\xrightarrow{m \rightarrow +\infty} v_0 && \text{in } L^2(\Lambda), \mathbb{P}'\text{-a.s. in } \Omega'. \end{aligned} \tag{4.2}$$

In Lemmas 4.5 and 4.6 we will show that, thanks to equality in law, v_m and z_m are in fact finite-volume functions with the same piecewise constant structure as u_{h_m, N_m}^l and $u_{h_m, N_m}^r - u_{h_m, N_m}^l$, respectively.

Lemma 4.5. *For $m \in \mathbb{N}^*$ fixed, v_m is a step function with respect to time and space in the sense that there exists $v_{h_m, N_m}^l \in \mathbb{R}^{d_{h_m} \times N_m}$ such that \mathbb{P}' -a.s. in Ω' $v_m = v_{h_m, N_m}^l$. Moreover, $v_{h_m, N_m}^l(0, x) := v_{h_m}^0(x) = v_m^0(x)$ for all $x \in \Lambda$ and, in particular $v_m^0 = v_{h_m}^0$ is a spatial step function.*

Proof. By [35, Lemma A3] with $E = L^2(0, T; L^2(\Lambda))$ and $F = \mathbb{R}^{d_{h_m} \times N_m}$ it follows that there exists $(v_K^n)_{\substack{K \in \mathcal{T}_m \\ n \in \{0, \dots, N_m-1\}}}$ in $\mathbb{R}^{d_{h_m} \times N_m}$ such that

$$v_m \equiv (v_K^n)_{\substack{K \in \mathcal{T}_m \\ n \in \{0, \dots, N_m-1\}}} \quad \mathbb{P}'\text{-a.s. in } \Omega'.$$

In the same manner with $E = L^2(\Lambda)$ and $F = \mathbb{R}^{d_{h_m}}$ it follows that there exists $(\tilde{v}_K^0)_{K \in \mathcal{T}_m}$ in $\mathbb{R}^{d_{h_m}}$ such that

$$v_m^0 \equiv (\tilde{v}_K^0)_{K \in \mathcal{T}_m} \quad \mathbb{P}'\text{-a.s. in } \Omega'.$$

We recall the notation of Subsection 2.2 and in particular that

$$u_{h_m, N_m}^l \equiv (u_K^n)_{\substack{K \in \mathcal{T}_m \\ n \in \{0, \dots, N_m-1\}}} \quad \mathbb{P}\text{-a.s. in } \Omega.$$

For any $K \in \mathcal{T}_m$ we consider the non-negative, Borel measurable mapping

$$\begin{aligned} \xi_K^0 : \mathbb{R}^{d_{hm}} \times \mathbb{R}^{d_{hm} \times N_m} &\rightarrow \mathbb{R} \\ ((a_M)_M, (b_M^k)_{M,k}) &\mapsto |a_K - b_K^0|. \end{aligned}$$

Since

$$\mathbb{P} \circ ((u_M^0)_M, (u_M^k)_{M,k})^{-1} = \mathbb{P}' \circ ((\tilde{v}_M^0)_M, (v_M^k)_{M,k})^{-1},$$

it follows that

$$0 = \mathbb{E} [\xi_K^0((u_M^0)_M, (u_M^k)_{M,k})] = \mathbb{E}' [\xi_K^0((\tilde{v}_M^0)_M, (v_M^k)_{M,k})] = \mathbb{E}' [| \tilde{v}_K^0 - v_K^0 |]$$

and therefore, for all $x \in K$ and all $K \in \mathcal{T}_m$, $v_m(0, x) = v_K^0 = \tilde{v}_K^0 = v_m^0(x)$ \mathbb{P}' -a.s. in Ω' . \square

Lemma 4.6. For $m \in \mathbb{N}^*$ fixed, $z_m(t, x) = v_K^{n+1} - v_K^n$ for all $(t, x) \in (t_n, t_{n+1}] \times K$ and \mathbb{P}' -a.s. in Ω' for any $K \in \mathcal{T}_m$ and $n \in \{0, \dots, N_m - 1\}$, where $(v_K^n)_{\substack{K \in \mathcal{T}_m \\ n \in \{0, \dots, N_m - 1\}}}$ is defined as in the proof of Lemma 4.5.

Proof. With similar arguments as in Lemma 4.5 it follows that there exists $(z_K^n)_{\substack{K \in \mathcal{T}_m \\ n \in \{0, \dots, N_m - 1\}}} \in \mathbb{R}^{d_{hm} \times N_m}$ such that

$$z_m \equiv (z_K^n)_{\substack{K \in \mathcal{T}_m \\ n \in \{0, \dots, N_m - 1\}}} \quad \mathbb{P}'\text{-a.s. in } \Omega'.$$

For any fixed $K \in \mathcal{T}_m$, $n \in \{0, \dots, N_m - 1\}$, the mapping

$$\Phi_K^n : \mathbb{R}^{d_{hm} \times N_m} \times \mathbb{R}^{d_{hm} \times N_m} \rightarrow \mathbb{R}, ((a_M^k)_{M,k}, (b_M^k)_{M,k}) \mapsto |a_K^{n+1} - a_K^n - b_K^n|$$

is non-negative and Borel measurable. Since

$$\mathbb{P} \circ ((u_M^k)_{M,k}, (u_M^{k+1} - u_M^k)_{M,k})^{-1} = \mathbb{P}' \circ ((v_M^k)_{M,k}, (z_M^k)_{M,k})^{-1},$$

it follows that for any $K \in \mathcal{T}_m$ and all $n \in \{0, \dots, N_m - 1\}$

$$\begin{aligned} 0 &= \mathbb{E} [\Phi_K^n((u_M^k)_{M,k}, (u_M^{k+1} - u_M^k)_{M,k})] = \mathbb{E}' [\Phi_K^n((v_M^k)_{M,k}, (z_M^k)_{M,k})] \\ &= \mathbb{E}' [|v_K^{n+1} - v_K^n - z_K^n|]. \end{aligned}$$

Therefore, for all $K \in \mathcal{T}_m$ and all $n \in \{0, \dots, N_m - 1\}$ there holds $z_K^n = v_K^{n+1} - v_K^n$ \mathbb{P}' -a.s. in Ω' . \square

Next we prove that the finite-volume function $(v_h^n)_{1 \leq n \leq N}$ we have just constructed verifies the following numerical scheme.

Lemma 4.7. For $m \in \mathbb{N}^*$ fixed, any $n \in \{0, \dots, N_m - 1\}$ and any $K \in \mathcal{T}_m$, v_K^{n+1} satisfies the semi-implicit equation

$$\frac{m_K}{\Delta t}(v_K^{n+1} - v_K^n) + \sum_{\sigma \in \mathcal{E}_{\text{int}} \cap \mathcal{E}_K} \frac{m_\sigma}{d_{K|L}}(v_K^{n+1} - v_L^{n+1}) - \frac{m_K}{\Delta t}g(v_K^n)\Delta_{n+1}B_m = 0 \quad (\text{NIE})$$

\mathbb{P}' -a.s. in Ω' , where $\Delta_{n+1}B_m := B_m(t_{n+1}) - B_m(t_n)$.

Proof. From Lemma 4.6 it follows that $z_K^n = v_K^{n+1} - v_K^n$, \mathbb{P}' -a.s. in Ω' for all $K \in \mathcal{T}_m$ and all $n \in \{0, \dots, N_m - 1\}$. Then, for arbitrary $K \in \mathcal{T}_m$, $n \in \{0, \dots, N_m - 1\}$ the mapping

$$\begin{aligned} \Psi_K^n : \mathbb{R}^{d_{hm} \times N_m} \times \mathbb{R}^{d_{hm} \times N_m} \times C([0, T]) &\rightarrow \mathbb{R} \\ ((a_M^k)_{M,k}, (b_M^k)_{M,k}, f) &\mapsto \left| \frac{m_K}{\Delta t}b_K^n + \sum_{\sigma \in \mathcal{E}_{\text{int}} \cap \mathcal{E}_K} \frac{m_\sigma}{d_{K|L}}(b_K^n + a_K^n) - (b_L^n + a_L^n) \right. \\ &\quad \left. - \frac{m_K}{\Delta t}g(a_K^n)(f(t_{n+1}) - f(t_n)) \right| \end{aligned}$$

is non-negative and Borel measurable. Since

$$\mathbb{P} \circ ((u_M^k)_{M,k}, (u_M^{k+1} - u_M^k)_{M,k}, W)^{-1} = \mathbb{P}' \circ ((v_M^k)_{M,k}, (z_M^k)_{M,k}, B_m)^{-1},$$

from Proposition 2.8 it follows that

$$\begin{aligned} 0 &= \mathbb{E} [\Psi_K^n((u_M^k)_{M,k}, (u_M^{k+1} - u_M^k)_{M,k}, W)] = \mathbb{E}' [\Psi_K^n((v_M^k)_{M,k}, (z_M^k)_{M,k}, B_m)] \\ &= \mathbb{E}' \left[\frac{m_K}{\Delta t}(v_K^{n+1} - v_K^n) + \sum_{\sigma \in \mathcal{E}_{\text{int}} \cap \mathcal{E}_K} \frac{m_\sigma}{d_{K|L}}(v_K^{n+1} - v_L^{n+1}) - \frac{m_K}{\Delta t}g(v_K^n)\Delta_{n+1}B_m \right]. \end{aligned}$$

Thus, for all $K \in \mathcal{T}_m$, $n \in \{0, \dots, N_m - 1\}$ and \mathbb{P}' -a.s. in Ω'

$$0 = \frac{m_K}{\Delta t}(v_K^{n+1} - v_K^n) + \sum_{\sigma \in \mathcal{E}_{\text{int}} \cap \mathcal{E}_K} \frac{m_\sigma}{d_{K|L}}(v_K^{n+1} - v_L^{n+1}) - \frac{m_K}{\Delta t}g(v_K^n)\Delta_{n+1}B_m.$$

□

4.3 The martingale identification argument

In this subsection, we show that each B_m is a Brownian motion with respect to the filtration given in Definition 4.8. With this result at hand, we may show that $(W_\infty(t))_{t \geq 0}$ is a Brownian motion with respect to the filtration given in Definition 4.11. In Lemma 4.13 we prove that u_∞ is admissible for the stochastic Itô integral with respect to $(W_\infty(t))_{t \geq 0}$. Finally, in Lemma 4.14 we provide an approximation result for the stochastic Itô integrals.

Definition 4.8. For $t \in [0, T]$ we define \mathcal{F}_t^m to be the smallest sub- σ -field of \mathcal{A}' generated by v_m^0 and $B_m(s)$ for $0 \leq s \leq t$. The right-continuous, \mathbb{P}' -augmented filtration of $(\mathcal{F}_t^m)_{t \in [0, T]}$ denoted by $(\mathfrak{F}_t^m)_{t \in [0, T]}$ is defined by

$$\mathfrak{F}_t^m := \bigcap_{s > t} \sigma[\mathcal{F}_s^m \cup \{\mathcal{N} \in \mathcal{A}' : \mathbb{P}'(\mathcal{N}) = 0\}]$$

for any $t \in [0, T]$.

Remark 4.9. We recall that for the augmented filtration and for given processes $(X_t)_{t \geq 0}$, $(Y_t)_{t \geq 0}$ such that $(X_t)_{t \geq 0}$ is adapted and $Y_t = X_t$ holds a.s. for all t , it holds true that $(Y_t)_{t \geq 0}$ is also adapted.

Lemma 4.10. $(v_m)_m$ is adapted to $(\mathfrak{F}_t^m)_{t \in [0, T]}$ and $(B_m(t))_{t \in [0, T]}$ is a Brownian motion with respect to $(\mathfrak{F}_t^m)_{t \in [0, T]}$.

Proof. Since $(\mathfrak{F}_t^m)_{t \in [0, T]}$ is a filtration induced by v_m^0 and B_m , in particular v_m^0 is \mathfrak{F}_0^m -measurable. Thus, applying the same arguments as in the proof of Proposition 2.8, from (NIE) it follows that v_m is adapted to $(\mathfrak{F}_t^m)_{t \in [0, T]}$. Since $\mathbb{P}' \circ (B_m)^{-1} = \mathbb{P} \circ W^{-1}$, we get the following results:

- $\mathbb{E}'[|B_m(0)|] = \mathbb{E}[|W(0)|] = 0$, hence $B_m(0) = 0$ \mathbb{P}' -a.s. in Ω' .
- By Burkholder-Davis-Gundy inequality there exists a constant $C_B > 0$ such that

$$\mathbb{E}' \left[\sup_{t \in [0, T]} |B_m(t)|^2 \right] = \mathbb{E} \left[\sup_{t \in [0, T]} |W(t)|^2 \right] \leq C_B T^{\frac{1}{2}} < \infty. \quad (4.3)$$

- For all $0 \leq s \leq t \leq T$ and all bounded, continuous functions $\psi : C_b(L^2(\Lambda) \times C([0, s]) \rightarrow \mathbb{R}$

$$0 = \mathbb{E}[(W(t) - W(s))\psi(u_{h_m}^0, W|_{[0, s]})] = \mathbb{E}'[(B_m(t) - B_m(s))\psi(v_m^0, B_m|_{[0, s]})], \quad (4.4)$$

and

$$\begin{aligned} 0 &= \mathbb{E}[(W^2(t) - W^2(s) - (t - s))\psi(u_{h_m}^0, W|_{[0, s]})] \\ &= \mathbb{E}'[(B_m^2(t) - B_m^2(s) - (t - s))\psi(v_m^0, B_m|_{[0, s]})] \end{aligned} \quad (4.5)$$

Recalling Definition 4.8, $\mathcal{F}_t^m = \sigma_t(v_m^0, B_m)$ for $t \in [0, T]$. The real-valued random variable

$$\Omega' \ni \omega' \mapsto \psi(v_m^0(\omega'), B_m|_{[0, s]}(\omega'))$$

is \mathcal{F}_s^m -measurable. Using the properties of conditional expectation from (4.4) it follows that

$$\begin{aligned} 0 &= \mathbb{E}'[(B_m(t) - B_m(s))\psi(v_m^0, B_m|_{[0, s]})] \\ &= \mathbb{E}'[\mathbb{E}'((B_m(t) - B_m(s))\psi(v_m^0, B_m|_{[0, s]})|\mathcal{F}_s^m)] \\ &= \mathbb{E}'[\psi(v_m^0, B_m|_{[0, s]})\mathbb{E}'(B_m(t) - B_m(s)|\mathcal{F}_s^m)]. \end{aligned} \quad (4.6)$$

Since (4.6) applies to every bounded and continuous function $\psi : C_b(L^2(\Lambda) \times C([0, s]) \rightarrow \mathbb{R}$, from the Lemma of Doob-Dynkin it follows that

$$0 = \mathbb{E}'[\mathbb{1}_A \mathbb{E}'(B_m(t) - B_m(s)|\mathcal{F}_s^m)]$$

for all \mathcal{F}_s^m -measurable subsets $A \in \mathcal{A}'$ and for all $0 \leq s \leq t \leq T$. From the above equation it now follows that $\mathbb{E}'(B_m(t) - B_m(s) | \mathcal{F}_s^m) = 0$ \mathbb{P}' -a.s. in Ω' for all $0 \leq s \leq t \leq T$ and therefore $(B_m(t))_{t \in [0, T]}$ is a martingale with respect to $(\mathcal{F}_t^m)_{t \in [0, T]}$. Using [17, p.75] we may conclude that $(B_m(t))_{t \in [0, T]}$ is also a martingale with respect to the augmented filtration $(\mathfrak{F}_t^m)_{t \in [0, T]}$. With similar arguments from (4.5) it follows that $((B_m(t))^2 - t)_{t \in [0, T]}$ is a martingale with respect to $(\mathfrak{F}_t^m)_{t \in [0, T]}$ and consequently the quadratic variation process $\langle\langle B_m \rangle\rangle_t$ of $(B_m(t))_{t \in [0, T]}$ is given by t for all $t \in [0, T]$ (for the Definition of the quadratic variation of a stochastic process see [8, Definition 2.19]). Summarizing the above results, $(B_m(t))_{t \in [0, T]}$ is a square integrable martingale with respect to $(\mathfrak{F}_t^m)_{t \in [0, T]}$ starting in 0 with almost surely continuous paths and quadratic variation $\langle\langle B_m \rangle\rangle_t = t$. From [14, Theorem 3.11] $(B_m(t))_{t \in [0, T]}$ is a Brownian motion with respect to $(\mathfrak{F}_t^m)_{t \in [0, T]}$. \square

In the following, we want to show firstly that the stochastic process $(W_\infty(t))_{t \in [0, T]} := W_\infty$ is a Brownian motion and secondly that a filtration may be chosen in order to have compatibility of u_∞ with stochastic integration in the sense of Itô with respect to W_∞ . Since u_∞ is a random variable taking values in $L^2(0, T; L^2(\Lambda))$, $u_\infty(t, \cdot)$ is only defined for a.e. $t \in [0, T]$ and the construction of an appropriate filtration induced by u_∞ becomes delicate.

Definition 4.11. For $t \in [0, T]$ let \mathcal{F}_t^∞ be the smallest sub- σ -field of \mathcal{A}' generated by v_0 , $W_\infty(s)$ and $\int_0^s u_\infty(r) dr$ for $0 \leq s \leq t$. The right-continuous, \mathbb{P}' -augmented filtration of $(\mathcal{F}_t^\infty)_{t \in [0, T]}$ denoted by $(\mathfrak{F}_t^\infty)_{t \in [0, T]}$ is defined by

$$\mathfrak{F}_t^\infty := \bigcap_{s > t} \sigma[\mathcal{F}_s^\infty \cup \{\mathcal{N} \in \mathcal{A}' : \mathbb{P}'(\mathcal{N}) = 0\}]$$

for $t \in [0, T]$.

In the following, we will show that W_∞ is a Brownian motion with respect to $(\mathfrak{F}_t^\infty)_{t \in [0, T]}$ and u_∞ admits a $(\mathfrak{F}_t^\infty)_{t \in [0, T]}$ -predictable representative.

Lemma 4.12. There holds $B_m \xrightarrow{m \rightarrow +\infty} W_\infty$ in $L^2(\Omega'; C([0, T]))$ and $(W_\infty(t))_{t \in [0, T]}$ is a Brownian motion with respect to $(\mathfrak{F}_t^\infty)_{t \in [0, T]}$.

Proof. Combining (4.3) with $\mathbb{P}' \circ (W_\infty)^{-1} = \mathbb{P} \circ W^{-1}$ it follows that

$$\mathbb{E}' \left[\sup_{t \in [0, T]} |W_\infty(t)|^2 \right] = \mathbb{E} \left[\sup_{t \in [0, T]} |W(t)|^2 \right] < \infty$$

and consequently $W_\infty \in L^2(\Omega'; C([0, T]))$. Moreover, since $\mathbb{P}' \circ B_m^{-1} = \mathbb{P} \circ W^{-1}$, it follows that

$$\lim_{m \rightarrow \infty} \mathbb{E}' \left[\sup_{t \in [0, T]} |B_m(t)|^2 \right] = \mathbb{E}' \left[\sup_{t \in [0, T]} |W_\infty(t)|^2 \right].$$

We already know, that, for $m \rightarrow +\infty$, B_m converges to W_∞ in $C([0, T])$ a.s. in Ω' . Therefore, a version of the Lemma of Brézis and Lieb (see [35, Lemma A2]) provides the desired convergence result in $L^2(\Omega'; C([0, T]))$. From

$$\mathbb{P}' \circ (v_m^0, B_m, v_m)^{-1} = \mathbb{P} \circ (u_{h_m}^0, W, u_{h_m, N_m}^l)^{-1}$$

it follows that for any $0 \leq s \leq t \leq T$ and every bounded and continuous function $\psi : L^2(\Lambda) \times C([0, s]) \times C([0, s]; L^2(\Lambda)) \rightarrow \mathbb{R}$, we have

$$\begin{aligned} & \mathbb{E}' \left[(B_m(t) - B_m(s)) \psi \left(v_m^0, B_m|_{[0, s]}, \int_0^\cdot v_m(r) dr|_{[0, s]} \right) \right] \\ &= \mathbb{E} \left[(W(t) - W(s)) \psi \left(u_{h_m}^0, W|_{[0, s]}, \int_0^\cdot u_{h_m, N_m}^l(r) dr|_{[0, s]} \right) \right]. \end{aligned} \quad (4.7)$$

Now, using the fact that $u_{h_m}^0$ is \mathcal{F}_0 -measurable, by construction $\int_0^s u_{h_m, N_m}^l(r) dr$ is \mathcal{F}_s -measurable for all $m \in \mathbb{N}$ and that $(W(t))_{t \geq 0}$ is a martingale with respect to $(\mathcal{F}_t)_{t \geq 0}$ one gets that

$$\mathbb{E} \left[(W(t) - W(s)) \psi \left(u_{h_m}^0, W|_{[0, s]}, \int_0^\cdot u_{h_m, N_m}^l(r) dr|_{[0, s]} \right) \right] = 0. \quad (4.8)$$

We recall that, \mathbb{P}' -a.s. in Ω' , $v_m^0 \rightarrow v_0$ in $L^2(\Lambda)$ and $B_m \rightarrow W_\infty$ in $C([0, T])$, hence also in $C([0, s])$ for all $0 \leq s \leq t \leq T$. Moreover, by Cauchy-Schwarz inequality,

$$\begin{aligned} & \left\| \int_0^\cdot v_m(r) dr - \int_0^\cdot u_\infty(r) dr \right\|_{C([0, s]; L^2(\Lambda))}^2 = \sup_{z \in [0, s]} \left\| \int_0^z (v_m(r) - u_\infty(r)) dr \right\|_{L^2(\Lambda)}^2 \\ & \leq \sup_{z \in [0, s]} \left(\int_0^z \|v_m(r) - u_\infty(r)\|_{L^2(\Lambda)}^2 dr \right) \leq T \int_0^T \|v_m(r) - u_\infty(r)\|_{L^2(\Lambda)}^2 dr. \end{aligned}$$

Since $v_m \xrightarrow{m \rightarrow +\infty} u_\infty$ in $L^2(0, T; L^2(\Lambda))$ \mathbb{P}' -a.s. in Ω' , it follows that

$$\int_0^\cdot v_m(r) dr \xrightarrow{m \rightarrow +\infty} \int_0^\cdot u_\infty(r) dr$$

in $C([0, s]; L^2(\Lambda))$, \mathbb{P}' -a.s. in Ω' . Using the upgraded convergence of B_m towards W_∞ in $L^2(\Omega'; C([0, T]))$, the convergence results from above and Lebesgue's dominated convergence theorem it follows that

$$\begin{aligned} & \lim_{m \rightarrow \infty} \mathbb{E}' \left[(B_m(t) - B_m(s)) \psi \left(v_m^0, B_m|_{[0, s]}, \int_0^\cdot v_m(r) dr|_{[0, s]} \right) \right] \\ &= \mathbb{E}' \left[(W_\infty(t) - W_\infty(s)) \psi \left(v_0, W_\infty|_{[0, s]}, \int_0^\cdot u_\infty(r) dr|_{[0, s]} \right) \right]. \end{aligned} \quad (4.9)$$

Now, combining (4.7), (4.8) and (4.9) it follows that

$$\mathbb{E}' \left[(W_\infty(t) - W_\infty(s)) \psi \left(v_0, W_\infty|_{[0, s]}, \int_0^\cdot u_\infty(r) dr|_{[0, s]} \right) \right] = 0. \quad (4.10)$$

From (4.10) it follows that $\mathbb{E}'(W_\infty(t) - W_\infty(s) | \mathcal{F}_s^\infty) = 0$ \mathbb{P}' -a.s. in Ω' for all $0 \leq s \leq t \leq T$ and therefore $(W_\infty(t))_{t \in [0, T]}$ is a martingale with respect to $(\mathcal{F}_t^\infty)_{t \in [0, T]}$. Using [17, p.75] we may conclude that $(W_\infty(t))_{t \in [0, T]}$ is also a martingale with respect to the augmented filtration $(\mathfrak{F}_t^\infty)_{t \in [0, T]}$. With similar arguments it follows that

$$\mathbb{E}' \left[(W_\infty^2(t) - W_\infty^2(s) - (t-s)) \psi \left(v_0, W_\infty|_{[0, s]}, \int_0^\cdot u_\infty(r) dr|_{[0, s]} \right) \right] = 0,$$

therefore $((W_\infty(t))^2 - t)_{t \in [0, T]}$ is a martingale with respect to $(\mathfrak{F}_t^\infty)_{t \in [0, T]}$ and consequently the quadratic variation process $\langle\langle W_\infty \rangle\rangle_t$ of $(W_\infty(t))_{t \in [0, T]}$ is given by t for all $t \in [0, T]$. Summarizing the above results, $(W_\infty(t))_{t \in [0, T]}$ is a square integrable martingale with respect to $(\mathfrak{F}_t^\infty)_{t \in [0, T]}$ starting in 0 with almost surely continuous paths for all $t \in [0, T]$ and quadratic variation $\langle\langle W_\infty \rangle\rangle_t = t$. From [14, Theorem 3.11] it follows that $(W_\infty(t))_{t \in [0, T]}$ is a Brownian motion with respect to $(\mathfrak{F}_t^\infty)_{t \in [0, T]}$. \square

By [10, Theorem 2.6.3] it is always possible to choose $(\Omega', \mathcal{A}', \mathbb{P}') = ([0, 1], \mathcal{B}([0, 1]), \lambda)$, where $\mathcal{B}([0, 1])$ denotes the Borel sets on $[0, 1]$ and λ denotes the Lebesgue measure on $[0, 1]$. We will need this particular choice of the new probability space in the proof of the following Lemma.

We recall that, for a filtered probability space $(\Omega, \mathcal{A}, \mathbb{P})$ with $\mathcal{F} = (\mathcal{F}_t)_{t \geq 0}$ and $T > 0$, the predictable σ -field on $\Omega \times [0, T]$ is the σ -field generated by the sets

$$(s, t] \times F_s, \quad 0 \leq s < t \leq T, \quad F_s \in \mathcal{F}_s \text{ and } \{0\} \times F_0, \quad F_0 \in \mathcal{F}_0.$$

For more details on stochastic integration, we refer to [14].

Lemma 4.13. *There exists a $(\mathfrak{F}_t^\infty)_{t \in [0, T]}$ -predictable, $d\mathbb{P}' \otimes dt$ -representative of u_∞ .*

Proof. For $\delta > 0$ we define $u_\infty^\delta : \Omega' \times [0, T] \rightarrow L^2(\Lambda)$ by

$$u_\infty^\delta(t) := \frac{1}{\delta} \int_{(t-\delta)^+}^t u_\infty(s) ds = \frac{1}{\delta} \left(\int_0^t u_\infty(s) ds - \int_0^{(t-\delta)^+} u_\infty(s) ds \right)$$

where the integrals on the right-hand side are understood as Bochner integrals with values in $L^2(\Lambda)$. Since u_∞^δ is an $(\mathfrak{F}_t^\infty)_{t \in [0, T]}$ -adapted stochastic process with a.s. continuous paths, it is predictable with respect to $(\mathfrak{F}_t^\infty)_{t \in [0, T]}$. For fixed $k \in \mathbb{N}$ the cut-off function $T_k : \mathbb{R} \rightarrow [-k, k]$ defined by $T_k(r) := r$ if $|r| < k$ and $T_k(r) := \text{sign}(r)k$ if $|r| \geq k$ induces a continuous operator $L^2(\Lambda) \ni v \mapsto T_k(v) \in L^2(\Lambda)$. Hence the stochastic process

$$\Omega' \times [0, T] \ni (\omega', t) \mapsto T_k(u_\infty^\delta(\omega', t)) \in L^2(\Lambda)$$

is $(\mathfrak{F}_t^\infty)_{t \in [0, T]}$ -predictable. Again we recall that, \mathbb{P}' -a.s. in Ω' , $v_m \xrightarrow{m \rightarrow +\infty} u_\infty$ in $L^2(0, T; L^2(\Lambda))$, hence also in $L^1(0, T; L^2(\Lambda))$ and therefore

$$\lim_{m \rightarrow +\infty} \int_0^T \|v_m(t)\|_{L^2(\Lambda)} dt = \int_0^T \|u_\infty(t)\|_{L^2(\Lambda)} dt$$

\mathbb{P}' -a.s. in Ω' . Using Fatou's lemma, $\mathbb{P}' \circ (v_m)^{-1} = \mathbb{P} \circ (u_{h_m, N_m}^l)^{-1}$, and the Cauchy-Schwarz inequality we obtain

$$\begin{aligned} \mathbb{E}' \left[\int_0^T \|u_\infty(t)\|_{L^2(\Lambda)} dt \right] &\leq \liminf_{m \rightarrow \infty} \mathbb{E}' \left[\int_0^T \|v_m(t)\|_{L^2(\Lambda)} dt \right] \\ &= \liminf_{m \rightarrow \infty} \mathbb{E} \left[\int_0^T \|u_{h_m, N_m}^l(t)\|_{L^2(\Lambda)} dt \right] \leq \sqrt{T} \liminf_{m \rightarrow \infty} \|u_{h_m, N_m}^l\|_{L^2(\Omega; L^2(0, T; L^2(\Lambda)))}^2. \end{aligned}$$

From Lemma 3.3 it follows that the right-hand side of the equation is uniformly bounded and consequently, $u_\infty \in L^1(\Omega'; L^1(0, T; L^2(\Lambda)))$. In particular, $u_\infty \in L^1(\Omega'; L^1(0, T; L^1(\Lambda)))$. Since $(\Omega', \mathcal{A}', \mathbb{P}') = ([0, 1], \mathcal{B}([0, 1]); \lambda)$ according to [21, Remark after Proposition 1.8.1] we have

$$L^1(\Omega'; L^1(0, T; L^1(\Lambda))) \cong L^1(\Omega' \times (0, T); L^1(\Lambda)) \cong L^1(0, T; L^1(\Omega'; L^1(\Lambda))).$$

For almost every $t \in (0, T)$ and $0 < \delta < t$ we have

$$\begin{aligned} \|u_\infty^\delta(t) - u_\infty(t)\|_{L^1(\Omega'; L^1(\Lambda))} &= \left\| \frac{1}{\delta} \int_{(t-\delta)^+}^t (u_\infty(s) - u_\infty(t)) ds \right\|_{L^1(\Omega'; L^1(\Lambda))} \\ &\leq \frac{1}{\delta} \int_{(t-\delta)^+}^t \|u_\infty(s) - u_\infty(t)\|_{L^1(\Omega'; L^1(\Lambda))} ds. \end{aligned}$$

By the generalisation of Lebesgue differentiation theorem for vector-valued functions (see, e.g. [18, Theorem 9, Chapter II]) the right-hand side of the above inequality goes to 0 for $\delta \rightarrow 0^+$ for almost every $t \in (0, T)$, hence $u_\infty^\delta(t) \rightarrow u_\infty(t)$ a.e in $L^1(\Omega'; L^1(\Lambda))$ for $\delta \rightarrow 0^+$. Then, Lebesgue's dominated convergence theorem provides

$$\lim_{\delta \rightarrow 0^+} T_k(u_\infty^\delta) = T_k(u_\infty)$$

in $L^1(0, T; L^1(\Omega'; L^1(\Lambda)))$, thus also in $L^1(\Omega' \times (0, T); L^1(\Lambda))$.

Thus, passing to a not relabeled subsequence if necessary, $(T_k(u_\infty^\delta(\omega', t)))_{\delta > 0}$ converges for almost every (ω', t) in $\Omega' \times (0, T)$ to $T_k(u_\infty(\omega', t))$ in $L^1(\Lambda)$ as $\delta \rightarrow 0^+$ and therefore $T_k(u_\infty(\omega', t))$ has a $d\mathbb{P}' \otimes dt$ representative that is $(\mathfrak{F}_t^\infty)_{t \in [0, T]}$ -predictable for every $k \in \mathbb{N}$. Obviously there holds

$$u_\infty(\omega', t) = \sup_{k \in \mathbb{N}} T_k(u_\infty(\omega', t)) \text{ in } L^1(\Lambda) \text{ for a.e. } (\omega', t) \text{ in } \Omega' \times (0, T),$$

where the set of measure zero can be chosen independently of $k \in \mathbb{N}$. This provides the existence of a $d\mathbb{P}' \otimes dt$ representative of u_∞ that is $(\mathfrak{F}_t^\infty)_{t \in [0, T]}$ -predictable. \square

Lemma 4.14. *Passing to a not relabeled subsequence if necessary,*

$$\mathcal{M}_{h_m, N_m} \xrightarrow{m \rightarrow +\infty} \mathcal{M}_\infty \text{ in } L^2(0, T; L^2(\Lambda)) \text{ } \mathbb{P}'\text{-a.s. in } \Omega', \quad (4.11)$$

where for $t \in [0, T]$, $x \in \Lambda$, \mathbb{P}' -a.s. in Ω' ,

$$\begin{aligned}\mathcal{M}_{h_m, N_m}(t, x) &:= \int_0^t g(v_{h_m, N_m}^l(s, x)) dB_m(s) \\ \mathcal{M}_\infty(t, x) &:= \int_0^t g(u_\infty(s, x)) dW_\infty(s).\end{aligned}$$

Proof. From Lemma 4.12, we know that $(B_m)_m$ converges in $L^2(\Omega'; C([0, T]))$ towards W_∞ which is a Brownian motion with respect to $(\mathfrak{F}_t^\infty)_{t \in [0, T]}$. Particularly, this convergence result also holds in probability in $C([0, T])$. Moreover, from the convergence (4.2) and Lemma 4.5, we know that $(v_{h_m, N_m}^l)_m$ converges towards u_∞ in $L^2(0, T; L^2(\Lambda))$, \mathbb{P}' -a.s. in Ω' , thus up to a subsequence denoted in the same way, using the Lipschitz property of g , it follows that $(g(v_{h_m, N_m}^l))_m$ converges towards $g(u_\infty)$ in probability in $L^2(0, T; L^2(\Lambda))$. Now, we can apply Lemma 2.1 in [15] and conclude that the convergence in (4.11) holds true in probability in $L^2(0, T; L^2(\Lambda))$ and therefore, passing to a subsequence if necessary, the assertion follows. \square

4.4 Convergence towards a martingale solution

For the sake of simplicity we use the notations $\mathcal{T} = \mathcal{T}_m$, $h = h_m$, $\Delta t = \Delta t_m$ and $N = N_m$. For any $n \in \{0, \dots, N\}$ and $K \in \mathcal{T}$, setting $\mathcal{M}_K^n = \mathcal{M}_{h, N}(t_n, x_K)$ we can define $\widehat{\mathcal{M}}_{h, N}$ using Definition (2.2) and we obtain the following strong convergence result in $L^p(\Omega'; L^2(0, T; L^2(\Lambda)))$.

Lemma 4.15. *Passing to a not relabeled subsequence if necessary, we have the following convergence results for any $1 \leq p < 2$:*

$$\begin{aligned}v_{h, N}^l, v_{h, N}^r \text{ and } \widehat{v}_{h, N} &\xrightarrow{m \rightarrow +\infty} u_\infty && \text{in } L^p(\Omega'; L^2(0, T; L^2(\Lambda))), \\ \mathcal{M}_{h, N} \text{ and } \widehat{\mathcal{M}}_{h, N} &\xrightarrow{m \rightarrow +\infty} \mathcal{M}_\infty && \text{in } L^p(\Omega'; L^2(0, T; L^2(\Lambda)))\end{aligned}$$

and

$$v_h^0 \xrightarrow{m \rightarrow +\infty} v_0 \quad \text{in } L^p(\Omega'; L^2(\Lambda)).$$

Moreover, $u_\infty \in L^2(\Omega'; L^2(0, T; H^1(\Lambda)))$ and $\mathcal{M}_\infty \in L^2(\Omega'; C([0, T]; L^2(\Lambda)))$.

Proof. We recall that thanks to convergence (4.2), $(v_{h, N}^l)_m$ converges to u_∞ for $m \rightarrow \infty$ in $L^2(0, T; L^2(\Lambda))$ \mathbb{P}' -a.s. in Ω' . Since $\mathbb{P}' \circ (v_{h, N}^l)^{-1} = \mathbb{P} \circ (u_{h, N}^l)^{-1}$, from Lemma 3.3 it follows that there exists a constant $C \geq 0$ such that

$$\mathbb{E}' \left[\|v_{h, N}^l\|_{L^2(0, T; L^2(\Lambda))}^2 \right] \leq C \tag{4.12}$$

for all $m \in \mathbb{N}$ and from Fatou's lemma we obtain $u_\infty \in L^2(\Omega'; L^2(0, T; L^2(\Lambda)))$. The convergence of $(v_{h, N}^l)_m$ towards u_∞ in $L^p(\Omega'; L^2(0, T; L^2(\Lambda)))$ is a consequence of (4.12) and of the theorem of Vitali (see, e.g., [21, Corollaire 1.3.3]). Now, using (4.1) we get

$$\mathbb{E}' \left[\|v_{h, N}^r - v_{h, N}^l\|_{L^2(0, T; L^2(\Lambda))}^2 \right] = \mathbb{E} \left[\|u_{h, N}^r - u_{h, N}^l\|_{L^2(0, T; L^2(\Lambda))}^2 \right] \xrightarrow{m \rightarrow +\infty} 0,$$

hence $(v_{h,N}^r - v_{h,N}^l) \rightarrow 0$ in $L^2(\Omega'; L^2(0, T; L^2(\Lambda)))$ as $m \rightarrow +\infty$ and, thanks to the continuous embedding of $L^2(\Omega'; L^2(0, T; L^2(\Lambda)))$ into $L^p(\Omega'; L^2(0, T; L^2(\Lambda)))$, also in $L^p(\Omega'; L^2(0, T; L^2(\Lambda)))$ for all $1 \leq p < 2$. Therefore, $v_{h,N}^r \rightarrow u_\infty$ as $m \rightarrow +\infty$ in $L^p(\Omega'; L^2(0, T; L^2(\Lambda)))$ for all $1 \leq p < 2$. Now we apply a similar argumentation to $(\widehat{v}_{h,N})_m$. We have

$$\begin{aligned} \mathbb{E}' \left[\|v_{h,N}^l - \widehat{v}_{h,N}\|_{L^2(0,T;L^2(\Lambda))}^2 \right] &= \mathbb{E}' \left[\sum_{n=0}^{N-1} \|v_h^{n+1} - v_h^n\|_{L^2(\Lambda)}^2 \int_{t_n}^{t_{n+1}} \left(\frac{t - t_n}{\Delta t} \right)^2 dt \right] \\ &= \frac{\Delta t}{3} \mathbb{E}' \left[\sum_{n=0}^{N-1} \|v_h^{n+1} - v_h^n\|_{L^2(\Lambda)}^2 \right]. \end{aligned} \quad (4.13)$$

Repeating the arguments of Proposition 3.2 on (NIE) it follows that there exists a constant $C'_2 \geq 0$ such that

$$\mathbb{E}' \left[\int_0^T |v_{h,N}^r|_{1,h}^2 dt \right] + \mathbb{E}' \left[\sum_{n=0}^{N-1} \|v_h^{n+1} - v_h^n\|_{L^2(\Lambda)}^2 \right] \leq C'_2. \quad (4.14)$$

Combining (4.13) with (4.14) it follows that $(v_{h,N}^l - \widehat{v}_{h,N}) \rightarrow 0$ in $L^2(\Omega'; L^2(0, T; L^2(\Lambda)))$ and we may conclude that $\widehat{v}_{h,N} \rightarrow u_\infty$ in $L^p(\Omega'; L^2(0, T; L^2(\Lambda)))$ for all $1 \leq p < 2$. Using (4.14) and the same arguments as in the proof of Lemma 4.1 on the discrete gradient $\nabla^h v_{h,N}^r$ of $v_{h,N}^r$ it follows that, passing to a not relabeled subsequence if necessary, $(\nabla^h v_{h,N}^r)_m$ converges weakly in $L^2(\Omega'; L^2(0, T; L^2(\Lambda)^2))$ towards ∇u_∞ , hence $u_\infty \in L^2(\Omega'; L^2(0, T; H^1(\Lambda)))$. We recall that, according to Lemma 4.14, $\mathcal{M}_{h,N} \rightarrow \mathcal{M}_\infty$ for $m \rightarrow +\infty$ in $L^2(0, T; L^2(\Lambda))$ \mathbb{P}' -a.s. in Ω' . Using the Lipschitz continuity of g , the Burkholder-Davis-Gundy inequality and Lemma 3.3 it follows that there exist constants $C_B \geq 0$, $C \geq 0$ such that

$$\begin{aligned} \mathbb{E}' \left[\sup_{t \in [0, T]} \|\mathcal{M}_{h,N}\|_{L^2(\Lambda)}^2 \right] &\leq C_B L^2 \mathbb{E}' \left[\int_0^T \|v_{h,N}^l\|_{L^2(\Lambda)}^2 dt \right] \\ &= C_B L^2 \mathbb{E} \left[\int_0^T \|u_{h,N}^l\|_{L^2(\Lambda)}^2 dt \right] \leq C_B L^2 C. \end{aligned} \quad (4.15)$$

Now, the convergence of $(\mathcal{M}_{h,N})_m$ towards \mathcal{M}_∞ in $L^p(\Omega'; L^2(0, T; L^2(\Lambda)))$ for all $1 \leq p < 2$ follows from (4.15) and the theorem of Vitali (see, e.g., [21, Corollaire 1.3.3]). Using the Itô isometry, the Lipschitz continuity of g and Lemma 3.3 it follows that there exists

$C_3 \geq 0$ such that

$$\begin{aligned}
& \mathbb{E}' \left[\int_0^T \|\mathcal{M}_{h,N} - \widehat{\mathcal{M}}_{h,N}\|_{L^2(\Lambda)}^2 dt \right] \\
&= \mathbb{E}' \left[\sum_{n=0}^{N-1} \int_{t_n}^{t_{n+1}} \left\| \int_{t_n}^t g(v_{h,N}^l) dB_m - \frac{t-t_n}{\Delta t} \int_{t_n}^{t_{n+1}} g(v_{h,N}^l) dB_m \right\|_{L^2(\Lambda)}^2 dt \right] \\
&\leq 2\mathbb{E}' \left[\sum_{n=0}^{N-1} \int_{t_n}^{t_{n+1}} \left\| \int_{t_n}^t g(v_{h,N}^l) dB_m \right\|_{L^2(\Lambda)}^2 + \left(\frac{t-t_n}{\Delta t} \right)^2 \left\| \int_{t_n}^{t_{n+1}} g(v_{h,N}^l) dB_m \right\|_{L^2(\Lambda)}^2 dt \right] \\
&= 2\mathbb{E}' \left[\sum_{n=0}^{N-1} \int_{t_n}^{t_{n+1}} \left(\int_{t_n}^t \|g(v_{h,N}^l)\|_{L^2(\Lambda)}^2 ds + \left(\frac{t-t_n}{\Delta t} \right)^2 \int_{t_n}^{t_{n+1}} \|g(v_{h,N}^l)\|_{L^2(\Lambda)}^2 ds \right) dt \right] \\
&\leq \frac{8\Delta t}{3} L^2 \mathbb{E}' \left[\int_0^T \|v_{h,N}^l\|_{L^2(\Lambda)}^2 dt \right] \leq C_3 \Delta t \xrightarrow{m \rightarrow +\infty} 0.
\end{aligned}$$

Hence, $\mathcal{M}_{h,N} - \widehat{\mathcal{M}}_{h,N} \rightarrow 0$ for $m \rightarrow +\infty$ in $L^2(\Omega'; L^2(0, T; L^2(\Lambda)))$ and therefore $\widehat{\mathcal{M}}_{h,N} \rightarrow \mathcal{M}_\infty$ for $m \rightarrow +\infty$ in $L^p(\Omega'; L^2(0, T; L^2(\Lambda)))$ for all $1 \leq p < 2$. Recalling that \mathcal{M}_∞ is a stochastic Itô integral with respect to the Brownian motion $(W_\infty(t))_{t \geq 0}$, we may conclude that \mathcal{M}_∞ has \mathbb{P}' -a.s. continuous paths in $L^2(\Lambda)$. From (4.15) and Fatou's lemma it now follows that $\mathcal{M}_\infty \in L^2(\Omega'; C([0, T]; L^2(\Lambda)))$. Since $\mathbb{P}' \circ (v_h^0)^{-1} = \mathbb{P} \circ (u_h^0)^{-1}$, from Lemma 3.1 it follows that

$$\mathbb{E}' \left[\|v_h^0\|_{L^2(\Lambda)}^2 \right] = \mathbb{E} \left[\|u_h^0\|_{L^2(\Lambda)}^2 \right] \leq \mathbb{E} \left[\|u_0\|_{L^2(\Lambda)}^2 \right].$$

Together with the \mathbb{P}' -a.s. convergence of $v_h^0 \xrightarrow{m \rightarrow +\infty} v_0$ in $L^2(\Lambda)$ from the convergence result (4.2), the last assertion follows again from the theorem of Vitali (see, e.g., [21, Corollaire 1.3.3]). \square

We now have all the necessary tools to pass to the limit in the scheme.

Proposition 4.16. *There exists a subsequence of $(\widehat{v}_{h,N})_m$, still denoted by $(\widehat{v}_{h,N})_m$, converging for $m \rightarrow +\infty$ towards a $(\mathfrak{F}_t^\infty)_{t \in [0, T]}$ -adapted stochastic process u_∞ with values in $L^2(\Lambda)$ and \mathbb{P}' -a.s. continuous paths. Moreover, $u_\infty \in L^2(\Omega'; L^2(0, T; H^1(\Lambda)))$ satisfies for all $t \in [0, T]$,*

$$u_\infty(t) - v_0 - \int_0^t \Delta u_\infty ds = \int_0^t g(u_\infty) dW_\infty \quad \text{in } L^2(\Lambda) \text{ and } \mathbb{P}'\text{-a.s. in } \Omega'.$$

Proof. Let $A \in \mathcal{A}'$, $\xi \in \mathcal{D}(\mathbb{R})$ with $\xi(T) = 0$ and $\varphi \in \mathcal{D}(\mathbb{R}^2)$ with $\nabla \varphi \cdot \mathbf{n} = 0$ on $\partial \Lambda$. Moreover we define the piecewise constant function $\varphi_h(x) := \varphi(x_K)$ for $x \in K$, $K \in \mathcal{T}$. For $K \in \mathcal{T}$, $n \in \{0, \dots, N-1\}$ and $t \in [t_n, t_{n+1})$ we multiply (NIE) with $\mathbb{1}_A \xi(t) \varphi(x_K)$ to obtain

$$\mathbb{1}_A \xi(t) \frac{m_K}{\Delta t} [v_K^{n+1} - v_K^n - g(v_K^n) \Delta_{n+1} B_m] \varphi(x_K) + \mathbb{1}_A \xi(t) \sum_{\sigma \in \mathcal{E}_{\text{int}} \cap \mathcal{E}_K} \frac{m_\sigma}{d_{K|L}} (v_K^{n+1} - v_L^{n+1}) \varphi(x_K) = 0.$$

Summing over $K \in \mathcal{T}$, integrating over $[t_n, t_{n+1}]$, summing over $n = 0, \dots, N-1$ and integrating over Ω' yields

$$\begin{aligned}
0 &= \int_{\Omega'} \sum_{n=0}^{N-1} \int_{t_n}^{t_{n+1}} \sum_{K \in \mathcal{T}} m_K \mathbb{1}_A \xi(t) \frac{1}{\Delta t} [v_K^{n+1} - v_K^n - g(v_K^n) \Delta_{n+1} B_m] \varphi(x_K) dt d\mathbb{P}' \\
&\quad + \int_{\Omega'} \sum_{n=0}^{N-1} \int_{t_n}^{t_{n+1}} \mathbb{1}_A \xi(t) \sum_{K \in \mathcal{T}} \sum_{\sigma \in \mathcal{E}_{\text{int}} \cap \mathcal{E}_K} \frac{m_\sigma}{d_{K|L}} (v_K^{n+1} - v_L^{n+1}) \varphi(x_K) dt d\mathbb{P}' \\
&=: T_{1,m} + T_{2,m}.
\end{aligned} \tag{4.16}$$

In the following, we will pass to the limit with $m \rightarrow +\infty$ on the right-hand side of (4.16). Using partial integration we obtain

$$\begin{aligned}
T_{1,m} &= \int_A \int_0^T \int_\Lambda \partial_t [\widehat{v}_{h,N} - \widehat{\mathcal{M}}_{h,N}] \xi(t) \varphi_h(x) dx dt d\mathbb{P}' \\
&= - \int_A \int_0^T \int_\Lambda [\widehat{v}_{h,N} - \widehat{\mathcal{M}}_{h,N}] \xi'(t) \varphi_h(x) dx dt d\mathbb{P}' - \int_A \int_\Lambda v_h^0 \xi(0) \varphi_h(x) dx d\mathbb{P}'.
\end{aligned}$$

Thanks to the convergence results of Lemma 4.15, passing to a not relabeled subsequence if necessary, we can pass to the limit and obtain

$$\begin{aligned}
&- \int_A \int_0^T \int_\Lambda [\widehat{v}_{h,N} - \widehat{\mathcal{M}}_{h,N}] \xi'(t) \varphi_h(x) dx dt d\mathbb{P}' - \int_A \int_\Lambda v_h^0 \xi(0) \varphi_h(x) dx d\mathbb{P}' \\
&\xrightarrow{m \rightarrow +\infty} - \int_A \int_0^T \int_\Lambda [u_\infty - \mathcal{M}_\infty] \xi'(t) \varphi(x) dx dt d\mathbb{P}' - \int_A \int_\Lambda v_0 \xi(0) \varphi(x) dx d\mathbb{P}'.
\end{aligned}$$

Now our aim is to show the following convergence result:

$$T_{2,m} \xrightarrow{m \rightarrow +\infty} - \int_A \int_0^T \int_\Lambda \xi(t) \Delta \varphi(x) u_\infty dx dt d\mathbb{P}'.$$

First, we can note that the term $T_{2,m}$ can be written as

$$T_{2,m} = \int_{\Omega'} \sum_{n=0}^{N-1} \int_{t_n}^{t_{n+1}} \mathbb{1}_A \xi(t) \sum_{K \in \mathcal{T}} v_K^{n+1} \sum_{\sigma \in \mathcal{E}_{\text{int}} \cap \mathcal{E}_K} m_\sigma \left(\frac{\varphi(x_K) - \varphi(x_L)}{d_{K|L}} \right) dt d\mathbb{P}'.$$

Then, since $\nabla \varphi \cdot \mathbf{n} = 0$ on $\partial \Lambda$, thanks to the Stokes formula one has,

$$\int_K \Delta \varphi(x) dx = \int_{\partial K} \nabla \varphi(x) \cdot \mathbf{n} d\sigma(x) = \sum_{\sigma \in \mathcal{E}_{\text{int}} \cap \mathcal{E}_K} \int_\sigma \nabla \varphi(x) \cdot \mathbf{n}_{KL} d\sigma(x).$$

Thus, we have

$$\begin{aligned}
T_{2,m} &= \\
&- \int_{\Omega'} \sum_{n=0}^{N-1} \int_{t_n}^{t_{n+1}} \mathbb{1}_A \xi(t) \sum_{K \in \mathcal{T}} v_K^{n+1} \left(\int_K \Delta \varphi(x) dx - \sum_{\sigma \in \mathcal{E}_{\text{int}} \cap \mathcal{E}_K} \int_{\sigma} \nabla \varphi(x) \cdot \mathbf{n}_{KL} d\sigma(x) \right) dt d\mathbb{P}' \\
&+ \int_{\Omega'} \sum_{n=0}^{N-1} \int_{t_n}^{t_{n+1}} \mathbb{1}_A \xi(t) \sum_{K \in \mathcal{T}} v_K^{n+1} \sum_{\sigma \in \mathcal{E}_{\text{int}} \cap \mathcal{E}_K} m_{\sigma} \left(\frac{\varphi(x_K) - \varphi(x_L)}{d_{K|L}} \right) dt d\mathbb{P}' \\
&= - \int_{\Omega'} \int_0^T \mathbb{1}_A \xi(t) \int_{\Lambda} v_{h,N}^r \Delta \varphi(x) dx dt d\mathbb{P}' \\
&+ \int_{\Omega'} \sum_{n=0}^{N-1} \int_{t_n}^{t_{n+1}} \mathbb{1}_A \xi(t) \sum_{\sigma \in \mathcal{E}_{\text{int}}} m_{\sigma} (v_K^{n+1} - v_L^{n+1}) R_{\sigma}^{\varphi} dt d\mathbb{P}',
\end{aligned}$$

with

$$R_{\sigma}^{\varphi} = \frac{1}{m_{\sigma}} \int_{\sigma} \nabla \varphi(x) \cdot \mathbf{n}_{KL} d\sigma(x) - \frac{\varphi(x_L) - \varphi(x_K)}{d_{K|L}}.$$

Using Lemma 4.15 and passing to a not relabeled subsequence if necessary, one gets

$$- \int_{\Omega'} \int_0^T \mathbb{1}_A \xi(t) \int_{\Lambda} v_{h,N}^r \Delta \varphi(x) dx dt d\mathbb{P}' \xrightarrow{m \rightarrow +\infty} - \int_A \int_0^T \int_{\Lambda} \xi(t) u_{\infty} \Delta \varphi(x) dx dt d\mathbb{P}'.$$

Concerning the second term in $T_{2,m}$, for any $\sigma = K|L \in \mathcal{E}_{\text{int}}$, the orthogonality condition implies $x_L - x_K = d_{K|L} \mathbf{n}_{KL}$, thus thanks to the Taylor formula for any $x \in \sigma$ one has,

$$\nabla \varphi(x) \cdot \mathbf{n}_{KL} = \frac{\varphi(x_L) - \varphi(x_K)}{d_{K|L}} + \mathcal{O}(h),$$

that gives,

$$R_{\sigma}^{\varphi} \leq C_{\varphi} h.$$

Therefore, thanks to the Cauchy-Schwarz inequality and inequality (4.14) the second term in $T_{2,m}$ satisfies,

$$\begin{aligned}
&\left| \int_{\Omega'} \sum_{n=0}^{N-1} \int_{t_n}^{t_{n+1}} \mathbb{1}_A \xi(t) \sum_{\sigma \in \mathcal{E}_{\text{int}}} m_{\sigma} (v_K^{n+1} - v_L^{n+1}) R_{\sigma}^{\varphi} dt d\mathbb{P}' \right| \\
&\leq C_{\varphi} h \int_{\Omega'} \sum_{n=0}^{N-1} \int_{t_n}^{t_{n+1}} \mathbb{1}_A |\xi(t)| \left(\sum_{\sigma \in \mathcal{E}_{\text{int}}} m_{\sigma} d_{K|L} \right)^{\frac{1}{2}} \left(\sum_{\sigma \in \mathcal{E}_{\text{int}}} m_{\sigma} \frac{|v_K^{n+1} - v_L^{n+1}|^2}{d_{K|L}} \right)^{\frac{1}{2}} \\
&\leq \sqrt{2} C_{\varphi} |\Lambda|^{\frac{1}{2}} h \int_{\Omega'} \int_0^T \mathbb{1}_A |\xi(t)| |v_{h,N}^r|_{1,h} \\
&\leq \sqrt{2} C_{\varphi} |\Lambda|^{\frac{1}{2}} h \|\xi \mathbb{1}_A\|_{L^2(\Omega' \times (0,T))} \left(\mathbb{E}' \left[\int_0^T |v_{h,N}^r|_{1,h}^2 dt \right] \right)^{\frac{1}{2}} \xrightarrow{m \rightarrow +\infty} 0.
\end{aligned}$$

Thus, we have shown that

$$\begin{aligned}
& - \int_0^T \int_{\Lambda} \left(u_{\infty} - \int_0^t g(u_{\infty}) dW_{\infty} \right) \xi'(t) \varphi(x) dx dt - \int_{\Lambda} v_0 \xi(0) \varphi(x) dx \\
& = \int_0^T \int_{\Lambda} u_{\infty} \Delta \varphi(x) \xi(t) dx dt = - \int_0^T \int_{\Lambda} \nabla u_{\infty} \cdot \nabla \varphi(x) \xi(t) dx dt
\end{aligned} \tag{4.17}$$

\mathbb{P}' -a.s. in Ω' for all $\xi \in \mathcal{D}(\mathbb{R})$ with $\xi(T) = 0$ and all $\varphi \in \mathcal{D}(\mathbb{R}^2)$ such that $\nabla \varphi \cdot \mathbf{n} = 0$ on $\partial\Lambda$. By [22, Theorem 1.1] the set $\{\varphi \in \mathcal{D}(\mathbb{R}^2) \mid \nabla \varphi \cdot \mathbf{n} = 0 \text{ on } \partial\Lambda\}$ is dense in $H^1(\Lambda)$ and therefore (4.17) applies to all $\varphi \in H^1(\Lambda)$.

In the following, we denote the dual space of $H^1(\Lambda)$ by $H^1(\Lambda)^*$, recall that

$$H^1(\Lambda) \hookrightarrow L^2(\Lambda) \hookrightarrow H^1(\Lambda)^*$$

with continuous and dense embeddings and we will denote the $H^1(\Lambda)$ - $H^1(\Lambda)^*$ duality bracket by $\langle \cdot, \cdot \rangle$ and the $L^2(\Lambda)$ scalar product by (\cdot, \cdot) . With the additional information

$$u_{\infty} \in L^2(\Omega'; L^2(0, T; H^1(\Lambda)))$$

from Lemma 4.15, it follows that

$$\Delta u_{\infty} \in L^2(\Omega'; L^2(0, T; H^1(\Lambda)^*))$$

and

$$- \int_{\Lambda} \int_0^T \nabla u_{\infty} \cdot \nabla \varphi(x) \xi(t) dx dt = \int_0^T \langle \Delta u_{\infty}, \varphi \rangle \xi(t) dt \tag{4.18}$$

\mathbb{P}' -a.s. in Ω' , for all $\xi \in \mathcal{D}(\mathbb{R})$ such that $\xi(T) = 0$ and all $\varphi \in H^1(\Lambda)$. Combining (4.17) with (4.18) and with the identity

$$- \int_{\Lambda} v_0 \varphi(x) \xi(0) dx = \int_0^T \int_{\Lambda} v_0 \varphi(x) \xi'(t) dx dt \tag{4.19}$$

(see, [34, Lemma 7.3]), from Fubini's theorem it follows that

$$\left\langle - \int_0^T \left(u_{\infty}(t) - \int_0^t g(u_{\infty}) dW_{\infty} - v_0 \right) \xi'(t) dt, \varphi \right\rangle = \left\langle \int_0^T \Delta u_{\infty} \xi(t) dt, \varphi \right\rangle$$

\mathbb{P}' -a.s. in Ω' for all $\xi \in \mathcal{D}(\mathbb{R})$ such that $\xi(T) = 0$ and all $\varphi \in H^1(\Lambda)$ and therefore

$$- \int_0^T \left(u_{\infty}(t) - \int_0^t g(u_{\infty}) dW_{\infty} - v_0 \right) \xi'(t) dt = \int_0^T \Delta u_{\infty} \xi(t) dt$$

in $H^1(\Lambda)^*$, for all $\xi \in \mathcal{D}(\mathbb{R})$ such that $\xi(T) = 0$, \mathbb{P}' -a.s. in Ω' and by a separability argument the exceptional set in Ω' may be chosen independently of φ and ξ . Consequently, (see, e.g. [12, Proposition A6])

$$u_{\infty} - \int_0^t g(u_{\infty}) dW_{\infty} - v_0 \in W^{1,2}(0, T; H^1(\Lambda)^*) \quad \mathbb{P}'\text{-a.s. in } \Omega'$$

and

$$\frac{d}{dt} \left(u_\infty - \int_0^t g(u_\infty) dW_\infty - v_0 \right) = \Delta u_\infty \quad \text{in } L^2(\Omega'; L^2(0, T; H^1(\Lambda)^*)). \quad (4.20)$$

Since g is Lipschitz continuous, from the chain rule for Sobolev functions it follows that $g(u_\infty) \in L^2(\Omega'; L^2(0, T; H^1(\Lambda)))$ and

$$\nabla \left(\int_0^t g(u_\infty) dW_\infty \right) = \int_0^t g'(u_\infty) \nabla u_\infty dW_\infty,$$

hence $u_\infty - \int_0^t g(u_\infty) dW_\infty \in L^2(\Omega'; L^2(0, T; H^1(\Lambda)))$. From [34, Lemma 7.3] we obtain $u_\infty \in L^2(\Omega'; C([0, T]; L^2(\Lambda)))$ and together with (4.20) the following rule of partial integration for all $0 \leq t \leq T$, \mathbb{P}' -a.s. in Ω' :

$$\begin{aligned} & \left(u_\infty(t) - \int_0^t g(u_\infty) dW_\infty - v_0, \zeta(t) \right) - (u_\infty(0) - v_0, \zeta(0)) \\ &= \int_0^t \langle \Delta u_\infty(s), \zeta(s) \rangle ds + \int_0^t \left\langle \zeta'(s), u_\infty(s) - \int_0^s g(u_\infty) dW_\infty - v_0 \right\rangle ds \end{aligned} \quad (4.21)$$

for all $\zeta \in L^2(0, T; H^1(\Lambda))$ with $\zeta' \in L^2(0, T; H^1(\Lambda)^*)$. Choosing $\zeta(t, x) = \xi(t)\varphi(x)$ with $\varphi \in H^1(\Lambda)$, $\xi \in \mathcal{D}(\mathbb{R})$ with $\xi(T) = 0$ in (4.21), we get

$$\begin{aligned} & \left(u_\infty(t) - \int_0^t g(u_\infty) dW_\infty - v_0, \varphi \right) \xi(t) - (u_\infty(0) - v_0, \varphi) \xi(0) \\ &= \int_0^t \xi(s) \langle \Delta u_\infty(s), \varphi \rangle ds + \int_0^t \xi'(s) \left(u_\infty(s) - \int_0^s g(u_\infty) dW_\infty - v_0, \varphi \right) ds \end{aligned} \quad (4.22)$$

\mathbb{P}' -a.s. in Ω' . The particular choice of $t = T$ and $\xi \in \mathcal{D}(\mathbb{R})$ with $\xi(T) = 0$ and $\xi(0) = 1$ in (4.22) combined with (4.17), (4.18) and (4.19) yields

$$(u_\infty(0) - v_0, \varphi) = 0 \quad \text{for all } \varphi \in H^1(\Lambda), \mathbb{P}'\text{-a.s. in } \Omega'$$

and therefore $u_\infty(0) = v_0$ \mathbb{P}' -a.s. in Ω' .

Now, we fix $t \in [0, T]$ and choose $\xi \in \mathcal{D}(\mathbb{R})$ with $\xi(T) = 0$ and $\xi(s) = 1$ for all $s \in [0, t]$. With this choice, from (4.22) we obtain

$$\left(u_\infty(t) - \int_0^t g(u_\infty) dW_\infty - u_\infty(0), \varphi \right) = \int_0^t \langle \Delta u_\infty(s), \varphi \rangle ds \quad (4.23)$$

\mathbb{P}' -a.s. in Ω' for all $\varphi \in H^1(\Lambda)$. Since, for fixed $\varphi \in H^1(\Lambda)$,

$$t \mapsto \left(u_\infty(t) - \int_0^t g(u_\infty) dW_\infty - u_\infty(0), \varphi \right) \quad \text{and} \quad t \mapsto \int_0^t \langle \Delta u_\infty(s), \varphi \rangle ds$$

are continuous in $[0, T]$, \mathbb{P}' -a.s. in Ω' , the exceptional set in Ω' in (4.23) may be chosen idenpendently of $t \in [0, T]$ and (4.23) also holds for $t = T$. This yields

$$u_\infty(t) - u_\infty(0) - \int_0^t g(u_\infty) dW_\infty = \int_0^t \Delta u_\infty(s) ds \quad \text{in } H^1(\Lambda)^* \text{ and } \mathbb{P}'\text{-a.s. in } \Omega'$$

and, since the left-hand side of the above equation is in $L^2(\Lambda)$, also in $L^2(\Lambda)$. \square

Remark 4.17. Applying the chain rule in (4.21) for $t = T$ and $\zeta = \Psi \in \mathcal{D}(\mathbb{R} \times \mathbb{R}^2)$ such that $\Psi(T, \cdot) = 0$ we immediately get that u_∞ is a weak solution, i.e.,

$$\begin{aligned} & \int_0^T \int_\Lambda u_\infty(t, x) \partial_t \Psi(t, x) dx dt - \int_0^T \int_\Lambda \nabla u_\infty(t, x) \cdot \nabla \Psi(t, x) dx dt + \int_\Lambda u_0(x) \Psi(0, x) dx \\ &= \int_0^T \int_\Lambda \int_0^t g(u_\infty(s, x)) dW_\infty(s) \partial_t \Psi(t, x) dx dt \end{aligned}$$

\mathbb{P}' -a.s. in Ω' .

4.5 Strong convergence of finite-volume approximations

In the previous subsections, we have shown that our finite-volume approximations converge towards a *martingale solution* of (1.1), i.e., the stochastic basis

$$(\Omega', \mathcal{A}', \mathbb{P}', (\mathfrak{F}_t^\infty)_{t \in [0, T]}, (W_\infty(t))_{t \in [0, T]})$$

is not a-priori given, but part of the solution. In this subsection, we want to show convergence of our finite-volume approximations with respect to the initially given stochastic basis

$$(\Omega, \mathcal{A}, \mathbb{P}, (\mathcal{F}_t)_{t \geq 0}, (W(t))_{t \geq 0}).$$

To do so, we will proceed in several steps. First, *pathwise uniqueness* of the heat equation with multiplicative Lipschitz noise is a consequence of Proposition 4.18: Roughly speaking, martingale solutions of (1.1) on a joint stochastic basis and with respect to the same initial datum coincide. In the proof of Proposition 4.20, we construct two convergent finite-volume approximations with respect to a joint stochastic basis, namely $(v_{\nu_k}^l)$ and $(v_{\rho_k}^l)$, from the function $(u_{h,N}^l)$ of our original finite-volume scheme using the theorems of Prokhorov and Skorokhod. Then, as a consequence of pathwise uniqueness, the limits coincide and we may apply [28, Lemma 1.1] in order to obtain the convergence in probability of $(u_{h,N}^l)$. Thanks to our previous result we can improve the convergence and pass to the limit in the originally given finite-volume scheme (see Lemma 4.21).

Proposition 4.18. *Let $(\Omega, \mathcal{A}, \mathbb{P}, (\mathcal{F}_t)_{t \geq 0}, (W(t))_{t \geq 0})$ be a stochastic basis and u_1, u_2 be solutions to (1.1) with respect to the \mathcal{F}_0 -measurable initial values u_0^1 and u_0^2 in $L^2(\Omega; L^2(\Lambda))$ respectively on $(\Omega, \mathcal{A}, \mathbb{P}, (\mathcal{F}_t)_{t \geq 0}, (W(t))_{t \geq 0})$. Then, there exists a constant $C \geq 0$ such that*

$$\mathbb{E} \left[\|u_1(t) - u_2(t)\|_{L^2(\Lambda)}^2 \right] \leq C \mathbb{E} \left[\|u_0^1 - u_0^2\|_{L^2(\Lambda)}^2 \right]$$

for all $t \in [0, T]$.

Proof. We apply the Itô formula ([30, Theorem 4.2.5]) to the process $u_1 - u_2$, discard the nonnegative term on the left-hand side of the resulting equation and take expectation. Then, the assertion is a straightforward consequence of Gronwall's inequality. \square

Remark 4.19. If u_1, u_2 are solutions to (1.1) on $(\Omega, \mathcal{A}, \mathbb{P}, (\mathcal{F}_t)_{t \geq 0}, (W(t))_{t \geq 0})$ with respect to the same initial value u_0 , from Proposition 4.18 it follows that $u_1(t) = u_2(t)$ in $L^2(\Lambda)$ for all $t \in [0, T]$, \mathbb{P} -a.s. in Ω . Since u_1, u_2 have continuous paths in $L^2(\Lambda)$, the exceptional set in Ω may be chosen independently of $t \in [0, T]$ and it follows that $u_1 = u_2$ in $L^2(0, T; L^2(\Lambda))$ \mathbb{P} -a.s. in Ω .

Proposition 4.20. *Let $(u_{h,N}^l)_m$ be given by Proposition 2.8. Then, there exists a subsequence of $(u_{h,N}^l)_m$, still denoted by $(u_{h,N}^l)_m$, converging for $m \rightarrow +\infty$ in $L^2(0, T; L^2(\Lambda))$ \mathbb{P} -a.s. in Ω towards a stochastic process u with values in $L^2(\Lambda)$, having \mathbb{P} -a.s. continuous paths and belonging to $L^2(\Omega; L^2(0, T; H^1(\Lambda)))$.*

Proof. For the sake of simplicity, we will write $u_m^l := u_{h_m, N_m}^l$ and $u_m^r := u_{h_m, N_m}^r$ in the following. We consider an arbitrary pair of subsequences $(u_\nu^l)_\nu, (u_\rho^l)_\rho$ of $(u_m^l)_m$. Our aim is to apply [28, Lemma 1.1], therefore we show that there exists a joint subsequence $(u_{\nu_k}^l, u_{\rho_k}^l)_k$ converging in law to a probability measure η on $L^2(0, T; L^2(\Lambda))^2$ such that

$$\eta(\{(x, y) \in L^2(0, T; L^2(\Lambda))^2 \mid x = y\}) = 1.$$

We define the random vector-valued sequence $(Y_{\nu, \rho})_{\nu, \rho}$ by

$$Y_{\nu, \rho} := (u_\nu^l, u_\rho^l, (u_\nu^r - u_\nu^l), (u_\rho^r - u_\rho^l), W, u_\nu^0, u_\rho^0, (u_\nu^0 - u_\rho^0))$$

for any $\nu, \rho \in \mathbb{N}^*$ and extract a joint subsequence

$$Y_k := (u_{\nu_k}^l, u_{\rho_k}^l, (u_{\nu_k}^r - u_{\nu_k}^l), (u_{\rho_k}^r - u_{\rho_k}^l), W, u_{\nu_k}^0, u_{\rho_k}^0, (u_{\nu_k}^0 - u_{\rho_k}^0))$$

for any $k \in \mathbb{N}$ that converges in law towards a probability measure

$$\eta_\infty := (\eta_\infty^1, \eta_\infty^2, \delta_0, \delta_0, \mathbb{P} \circ W^{-1}, \mathbb{P} \circ (u_0)^{-1}, \mathbb{P} \circ (u_0)^{-1}, \delta_0)$$

where we include the difference of the random initial data into the vector to ensure, that $u_{\nu_k}^0$ and $u_{\rho_k}^0$ converge to the same limit. With straightforward modifications of the arguments of Subsections 4.2-4.4, we can find random elements $u_\infty^1, u_\infty^2, v_0$,

$$Y_k' = (v_{\nu_k}^l, v_{\rho_k}^l, z_{\nu_k}, z_{\rho_k}, W_k, v_{\nu_k}^0, v_{\rho_k}^0, (v_{\nu_k}^0 - v_{\rho_k}^0))$$

such that

$$\begin{aligned} \mathbb{P}' \circ (Y_k')^{-1} &= \mathbb{P} \circ (Y_k)^{-1} \text{ for all } k \in \mathbb{N}, \\ \mathbb{P}' \circ (v_0)^{-1} &= \mathbb{P} \circ (u_0)^{-1}, \\ \mathbb{P}' \circ (u_\infty^1)^{-1} &= \eta_\infty^1, \\ \mathbb{P}' \circ (u_\infty^2)^{-1} &= \eta_\infty^2 \end{aligned}$$

and a joint stochastic basis $(\Omega', \mathcal{A}', \mathbb{P}', (\mathfrak{F}_t^\infty)_{t \in [0, T]}, (W_\infty(t))_{t \in [0, T]})$ such that u_∞^1 and u_∞^2 are both solutions of (1.1) with initial value v_0 on $(\Omega', \mathcal{A}', \mathbb{P}', (\mathfrak{F}_t^\infty)_{t \in [0, T]}, (W_\infty(t))_{t \in [0, T]})$. Thus, from Proposition 4.18 and Remark 4.19 it follows for $\eta = (\eta_\infty^1, \eta_\infty^2)$

$$\begin{aligned} 1 = \mathbb{P}'(\{u_\infty^1 = u_\infty^2\}) &= \mathbb{P}' \circ (u_\infty^1, u_\infty^2)^{-1}(\{(x, y) \in L^2(0, T; L^2(\Lambda))^2 \mid x = y\}) \\ &= \eta(\{(x, y) \in L^2(0, T; L^2(\Lambda))^2 \mid x = y\}). \end{aligned}$$

From [28, Lemma 1.1] we get the convergence of u_m^l in probability to a random element \tilde{u} in $L^2(0, T; L^2(\Lambda))$. Obviously by Lemma 4.1 $\tilde{u} = u$. We can extract a not relabeled subsequence of $(u_m^l)_m$ that converges a.s. in $L^2(0, T; L^2(\Lambda))$ and in particular

$$\|u_m^l\|_{L^2(0, T; L^2(\Lambda))} \xrightarrow{m \rightarrow \infty} \|u\|_{L^2(0, T; L^2(\Lambda))} \quad \mathbb{P}\text{-a.s. in } \Omega.$$

From Lemma 3.4 and Fatou's Lemma we obtain $u \in L^2(\Omega; L^2(0, T; L^2(\Lambda)))$. For all $f : L^2(0, T; L^2(\Lambda)) \rightarrow \mathbb{R}$ continuous and bounded it follows from Lebesgue dominated convergence theorem that

$$\mathbb{E} [f(u_m^l)] \xrightarrow{m \rightarrow \infty} \mathbb{E} [f(u)],$$

so $\mathbb{P} \circ (u_m^l)^{-1} \rightarrow \eta_\infty^1$ for $m \rightarrow \infty$ and by Lemma 4.4 $\eta_\infty^1 = \mu_\infty^1$. Hence there holds in particular

$$\mathbb{E} [\|u_m^l\|_{L^2(0, T; L^2(\Lambda))}^2] \xrightarrow{m \rightarrow \infty} \mathbb{E} [\|u\|_{L^2(0, T; L^2(\Lambda))}^2] < \infty.$$

From Vitali's theorem we get $u_m^l \xrightarrow{m \rightarrow \infty} u$ in $L^p(\Omega; L^2(0, T; L^2(\Lambda)))$ for $1 < p < 2$. But from Lemma 4.1 we know $u_m^l \rightharpoonup u$ for $m \rightarrow +\infty$ weakly in $L^q(\Omega; L^2(0, T; L^2(\Lambda)))$ for $1 \leq q \leq 2$ with $u \in L^2(\Omega; L^2(0, T; H^1(\Lambda)))$. Now the Lemma of Brézis and Lieb (see [13]) yields $u_m^l \xrightarrow{m \rightarrow \infty} u$ in $L^2(\Omega; L^2(0, T; L^2(\Lambda)))$. \square

Lemma 4.21. *The semi-implicit finite-volume scheme (2.4) converges towards u given in Proposition 4.20 which is a solution to (1.1) in the sense of Definition 1.3.*

Proof. With similar arguments as in Proposition 4.20 it follows that $u_{h,N}^r$ and $\hat{u}_{h,N}$ converge to u in $L^2(\Omega; L^2(0, T; L^2(\Lambda)))$ for $m \rightarrow \infty$. Moreover there holds $g(u_{h,N}^l) \xrightarrow{m \rightarrow \infty} g(u)$ in $L^2(\Omega; L^2(0, T; L^2(\Lambda)))$. Therefore

$$M_{h,N} = \int_0^\cdot g(u_{h,N}^l) dW \xrightarrow{m \rightarrow \infty} \int_0^\cdot g(u) dW \quad \text{in } L^2(\Omega; C([0, T]; L^2(\Lambda))).$$

As shown in Lemma 4.15 it follows $\widehat{M}_{h,N} \xrightarrow{m \rightarrow \infty} \int_0^\cdot g(u) dW$ in $L^2(\Omega; L^2(0, T; L^2(\Lambda)))$. Now we consider the semi-implicit finite-volume scheme (2.4), multiply it with $\mathbb{1}_A \xi \varphi$, where $A \in \mathcal{A}$, $\xi \in \mathcal{D}(\mathbb{R})$ with $\xi(T) = 0$ and $\varphi \in \mathcal{D}(\mathbb{R}^2)$ with $\nabla \varphi \cdot \mathbf{n} = 0$, then we sum over $K \in \mathcal{T}$, integrate over $[t_n, t_{n+1})$ and sum over $n = 0, \dots, N-1$ to get

$$\begin{aligned} 0 &= \int_\Omega \sum_{n=0}^{N-1} \int_{t_n}^{t_{n+1}} \sum_{K \in \mathcal{T}} m_K \mathbb{1}_A \xi(t) \frac{1}{\Delta t} [u_K^{n+1} - u_K^n - g(u_K^n) \Delta_{n+1} W] \varphi(x_K) dt d\mathbb{P} \\ &\quad + \int_\Omega \sum_{n=0}^{N-1} \int_{t_n}^{t_{n+1}} \sum_{K \in \mathcal{T}} \mathbb{1}_A \xi(t) \sum_{K \in \mathcal{T}} \sum_{\sigma \in \mathcal{E}_{\text{int}} \cap \mathcal{E}_K} \frac{m_\sigma}{d_{K|L}} (u_K^{n+1} - u_L^{n+1}) \varphi(x_K) dt d\mathbb{P} \\ &=: T_{1,m} + T_{2,m}. \end{aligned}$$

If we define $\varphi_h(x) := \varphi(x_K)$ for $x \in K$, $K \in \mathcal{T}$, there holds

$$\begin{aligned} T_{1,m} &= \int_A \int_0^T \int_\Lambda \partial_t [\widehat{u}_{h,N} - \widehat{M}_{h,N}] \xi(t) \varphi_h(x) dx dt d\mathbb{P} \\ &= - \int_A \int_0^T \int_\Lambda (\widehat{u}_{h,N} - \widehat{M}_{h,N}) \xi'(t) \varphi_h(x) dx dt d\mathbb{P} - \int_A \int_\Lambda u_h^0 \varphi_h(x) dx d\mathbb{P}. \end{aligned}$$

From [1, Proposition 3.5] we know that $u_h^0 \xrightarrow{m \rightarrow +\infty} u_0$ in $L^2(\Lambda)$, \mathbb{P} -a.s. in Ω , and thanks to Lemma 3.1 we can apply Lebesgue's dominated convergence theorem. The passage to the limit is analogous to that on Ω' . \square

Acknowledgments This work has been supported by the SIMALIN project (ANR-19-CE40-0016) of the French National Research Agency, by the German Research Foundation project ZI 1542/3-1 and the Procope programme for Project-Related Personal Exchange (France-Germany).

References

- [1] B. Andreianov, F. Boyer, and F. Hubert. Discrete duality finite volume schemes for Leray-Lions-type elliptic problems on general 2D meshes. *Numer. Methods Partial Differential Equations*, 23(1):145–195, 2007.
- [2] R. Anton, D. Cohen, and L. Quer-Sardanyons. A fully discrete approximation of the one-dimensional stochastic heat equation. *IMA J. Numer. Anal.*, 40(1):247–284, 2020.
- [3] C. Bauzet, V. Castel, and J. Charrier. Existence and uniqueness result for an hyperbolic scalar conservation law with a stochastic force using a finite volume approximation. *J. Hyperbolic Differ. Equ.*, 17(2):213–294, 2020.
- [4] C. Bauzet, J. Charrier, and T. Gallouët. Convergence of flux-splitting finite volume schemes for hyperbolic scalar conservation laws with a multiplicative stochastic perturbation. *Math. Comp.*, 85(302):2777–2813, 2016.
- [5] C. Bauzet, J. Charrier, and T. Gallouët. Convergence of monotone finite volume schemes for hyperbolic scalar conservation laws with multiplicative noise. *Stoch. Partial Differ. Equ. Anal. Comput.*, 4(1):150–223, 2016.
- [6] C. Bauzet, J. Charrier, and T. Gallouët. Numerical approximation of stochastic conservation laws on bounded domains. *ESAIM Math. Model. Numer. Anal.*, 51(1):225–278, 2017.

- [7] C. Bauzet and F. Nabet. Convergence of a finite-volume scheme for a heat equation with a multiplicative stochastic force. In *Finite volumes for complex applications IX—methods, theoretical aspects, examples—FVCA 9, Bergen, Norway, June 2020*, volume 323 of *Springer Proc. Math. Stat.*, pages 275–283. Springer, Cham, 2020.
- [8] Ł. Bañas, Z. Brzeźniak, M. Neklyudov, and A. Prohl. *Stochastic ferromagnetism*, volume 58 of *De Gruyter Studies in Mathematics*. De Gruyter, Berlin, 2014.
- [9] P. Billingsley. *Convergence of probability measures*. Wiley Series in Probability and Statistics: Probability and Statistics. John Wiley & Sons, Inc., New York, second edition, 1999.
- [10] D. Breit, E. Feireisl, and M. Hofmanová. *Stochastically forced compressible fluid flows*, volume 3 of *De Gruyter Series in Applied and Numerical Mathematics*. De Gruyter, Berlin, 2018.
- [11] D. Breit, M. Hofmanová, and S. Loisel. Space-time approximation of stochastic p -Laplace-type systems. *SIAM J. Numer. Anal.*, (4):2218–2236, 2021.
- [12] H. Brézis. *Opérateurs maximaux monotones et semi-groupes de contractions dans les espaces de Hilbert*. North-Holland Mathematics Studies, No. 5. North-Holland Publishing Co., Amsterdam-London; American Elsevier Publishing Co., Inc., New York, 1973.
- [13] H. Brézis and E. Lieb. A relation between pointwise convergence of functions and convergence of functionals. *Proc. Amer. Math. Soc.*, 88(3):486–490, 1983.
- [14] G. Da Prato and J. Zabczyk. *Stochastic equations in infinite dimensions*, volume 152 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, second edition, 2014.
- [15] A. Debussche, N. Glatt-Holtz, and R. Temam. Local martingale and pathwise solutions for an abstract fluids model. *Physica D: Nonlinear Phenomena*, 240(14):1123–1144, 2011.
- [16] A. Debussche and J. Printems. Weak order for the discretization of the stochastic heat equation. *Math. Comp.*, 78(266):845–863, 2009.
- [17] Claude Dellacherie and Paul-André Meyer. *Probabilités et potentiel. Chapitres V à VIII*. Actualités Scientifiques et Industrielles [Current Scientific and Industrial Topics], No. 1385. Hermann, Paris, revised edition, 1980.
- [18] J. Diestel and J. J. Uhl, Jr. *Vector measures*. Mathematical Surveys, No. 15. American Mathematical Society, Providence, R.I., 1977.
- [19] S. Dotti and J. Vovelle. Convergence of approximations to stochastic scalar conservations laws. *Archive for Rational Mechanics and Analysis*, 1(1):1–53, 2018.

- [20] S. Dotti and J. Vovelle. Convergence of the finite volume method for scalar conservation laws with multiplicative noise: an approach by kinetic formulation. *Stochastic and Partial Differential Equations : Analysis and Computations*, 8(4):265–310, 2020.
- [21] J. Droniou. Intégration et Espaces de Sobolev à Valeurs Vectorielles. <https://hal.archives-ouvertes.fr/hal-01382368/document>, 2001.
- [22] J. Droniou. A density result in Sobolev spaces. *J. Math. Pures App.*, 81(7):697–714, 2002.
- [23] R. Eymard and T. Gallouët. H-convergence and numerical scheme for elliptic problems. *SIAM J. Numer. Anal.*, 41(2):539–562, 2003.
- [24] R. Eymard, T. Gallouët, and R. Herbin. Finite volume methods. In *Handbook of numerical analysis, Vol. VII*, Handb. Numer. Anal., VII, pages 713–1020. North-Holland, Amsterdam, 2000.
- [25] F. Flandoli and D. Gatarek. Martingale and stationary solutions for stochastic Navier-Stokes equations. *Prob. Th. Rel. Fields*, 102:367–391, 1995.
- [26] T. Funaki, Y. Gao, and D. Hilhorst. Convergence of a finite volume scheme for a stochastic conservation law involving a Q -Brownian motion. *Discrete Contin. Dyn. Syst. Ser. B*, 23(4):1459–1502, 2018.
- [27] T. Gallouët, R. Herbin, A. Larcher, and J.-C. Latché. Analysis of a fractional-step scheme for the \mathbf{P}_1 radiative diffusion model. *Comput. Appl. Math.*, 35(1):135–151, 2016.
- [28] I. Gyöngy and N. Krylov. Existence of strong solutions for Itô’s stochastic equations via approximations. *Probab. Th. Rel. Fields*, 105:143–158, 1996.
- [29] N. V. Krylov and B. L. Rozovskii. Stochastic evolution equations. *Journal of Soviet Mathematics*, 16(4):1233–1277, 1981.
- [30] W. Liu and M. Röckner. *Stochastic partial differential equations: an introduction*. Universitext. Springer, Cham, 2015.
- [31] A. Majee. Convergence of a flux-splitting finite volume scheme for conservation laws driven by Lévy noise. *Applied Mathematics and Computation*, 338(1):676–697, 2018.
- [32] M. Ondrejat, A. Prohl, and N. Walkington. Numerical Approximation of Nonlinear SPDE’s, arXiv:2003.06001.
- [33] É. Pardoux. *Équations aux dérivées partielles stochastiques non linéaires monotones*. Ph.d. thesis, University Paris Sud, 1975.

- [34] Tomáš Roubíček. *Nonlinear partial differential equations with applications*, volume 153 of *International Series of Numerical Mathematics*. Birkhäuser/Springer Basel AG, Basel, second edition, 2013.
- [35] G. Vallet and A. Zimmermann. Well-posedness for nonlinear SPDEs with strongly continuous perturbation. *Proc. Roy. Soc. Edinburgh Sect. A*, 151(1):265–295, 2021.