



**HAL**  
open science

# A note on the pp conjecture for sheaves of spaces of orderings

Pawel Gladki

► **To cite this version:**

Pawel Gladki. A note on the pp conjecture for sheaves of spaces of orderings. Communications in Mathematics, 2016, Volume 24 (2016), Issue 1 (1), pp.1 - 5. 10.1515/cm-2016-0001 . hal-03664921

**HAL Id: hal-03664921**

**<https://hal.science/hal-03664921>**

Submitted on 11 May 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

## A note on the pp conjecture for sheaves of spaces of orderings

*Paweł Gładki*

**Abstract.** In this note we provide a direct and simple proof of a result previously obtained by Astier stating that the class of spaces of orderings for which the pp conjecture holds true is closed under sheaves over Boolean spaces.

The theory of abstract spaces of orderings was developed by Murray Marshall in a series of papers from the late 1970s, and provides an abstract framework for studying orderings of fields and the reduced theory of quadratic forms in general. We will refer to the monograph [5] as far as background, notation, and main results are concerned. A space of orderings is a pair  $(X, G)$  such that  $X$  is a nonempty set,  $G$  is a subgroup of  $\{1, -1\}^X$ , which contains the constant function  $-1$ , separates points of  $X$ , and satisfies some additional axioms (see [5] for details).

A space of orderings  $(X, G)$  has a natural topology introduced by the family of subbasic clopen Harrison sets:

$$H(a) = \{x \in X : a(x) = 1\},$$

for a given  $a \in G$ , which makes  $X$  into a Boolean space [5, Theorem 2.1.5].

For any multiplicative group  $G$  of exponent 2 with distinguished element  $-1$ , we set  $X = \{x \in \chi(G) : x(-1) = -1\}$  and call the pair  $(X, G)$  a fan. A fan is also a space of orderings [5, Theorem 3.1.1]. We can also consider fans within a bigger space of orderings, and for this we need the notion of a subspace of a space  $(X, G)$  – a subset  $Y \subseteq X$  is called a subspace of  $(X, G)$ , if  $Y$  is expressible in the form  $\bigcap_{a \in S} H(a)$ , for some subset  $S \subseteq G$ . For any subspace  $Y$  we will denote by  $G|_Y$  the group of all restrictions  $a|_Y$ ,  $a \in G$ . The pair  $(Y, G|_Y)$  is a space of orderings itself [5, Theorem 2.4.3]. Finally, if  $(X, G)$  is a space of orderings, by a fan in  $(X, G)$  we understand a subspace  $V$  such that the space  $(V, G|_V)$  is a fan.

---

2010 MSC: Primary 11E10; Secondary 12D15

Key words: spaces of orderings, pp conjecture

DOI: 10.1515/cm-2016-0001

Sheaves of spaces of orderings had been defined in [4, Chapter 8], where the original results are phrased in terms of reduced Witt rings, and recently studied in [1] and [3]. The definition of the sheaf that we use here is exactly the one taken from [3]. Assume  $(X_i, G_i)$  is a space of orderings for each  $i \in I$ , where  $I$  is a Boolean space. Assume further that  $X = \bigcup_{i \in I} X_i$ , the disjoint union of  $X_i$ 's, is equipped with a topology such that

- (1)  $X$  is a Boolean space,
- (2) the inclusion map  $\iota_i : X_i \hookrightarrow X$  is continuous, for each  $i \in I$ ,
- (3) the projection map  $\pi : X \rightarrow I$  is continuous, and
- (4) if  $(i_\lambda)_{\lambda \in D}$  is any net in  $I$  converging to  $i \in I$  and if  $\sigma_1^\lambda, \sigma_2^\lambda, \sigma_3^\lambda, \sigma_4^\lambda$  is a 4-element fan in  $X_{i_\lambda}$  such that  $\sigma_j^\lambda$  converges to  $\sigma_j \in X_i$  for each  $j = 1, 2, 3, 4$ , then  $\sigma_1 \sigma_2 \sigma_3 \sigma_4 = 1$ .

Furthermore, let

$$G := \{\phi \in \text{Cont}(X, \{\pm 1\}) : \phi|_{X_i} \in G_i \forall i \in I\}$$

where  $G_i$  is identified with its image in  $\text{Cont}(X_i, \{\pm 1\})$ ,  $i \in I$ , under the natural embedding. Then  $(X, G)$  is a space of orderings called the sheaf of spaces  $(X_i, G_i)$  over the Boolean space  $I$ .

The pp conjecture has been posed in [6], and, as proven in [2], has a negative solution in general, although it is still interesting to investigate numerous examples of spaces where it is valid. Recall that, for a space of orderings  $(X, G)$ , a positive primitive (pp for short) formula  $P(\underline{a})$  with  $n$  quantifiers and  $k$  parameters in  $G$  is of the form

$$P(\underline{a}) = \exists \underline{t} \bigwedge_{j=1}^m p_j(\underline{t}, \underline{a}) \in D_X(1, q_j(\underline{t}, \underline{a})),$$

where  $\underline{t} = (t_1, \dots, t_n)$ ,  $\underline{a} = (a_1, \dots, a_k)$ , for  $a_1, \dots, a_k \in G$ , and  $p_j(\underline{t}, \underline{a})$ ,  $q_j(\underline{t}, \underline{a})$  are  $\pm$  products of some of the  $t_s$ 's and  $a_r$ 's,  $s \in \{1, \dots, n\}$ ,  $r \in \{1, \dots, k\}$ , for  $j \in \{1, \dots, m\}$ . The pp conjecture then asks whether it is true that if a pp formula holds in every finite subspace of a space of orderings, then it also holds in the whole space.

Our main goal in this note is the following:

**Theorem 1.** *If the pp conjecture holds in  $(X_i, G_i)$ , for all  $i \in I$ , then it also holds in the sheaf  $(X, G)$  of  $(X_i, G_i)$  over the Boolean space  $I$ .*

This is essentially Theorem 2.2 in [1], where the author uses the language and methods from model theory. Here we provide the proof that uses only some simple topology and set theory along with the standard notation used in the theory of spaces of orderings. We begin with a series of lemmas that come from [4], where they are stated for abstract Witt rings, which are basically the same things as abstract spaces of orderings; nonetheless, we shall provide proofs here for the sake of completeness of presentation, as well as the language used.

**Lemma 1.** *For all  $i \in I$ , and for every  $f \in G_i$ , there exists  $\phi \in \text{Cont}(X, \{\pm 1\})$  such that  $\phi|_{X_i} = f$ .*

*Proof.* Fix  $i \in I$  and  $f \in G_i$ . Consider the subsets  $A := \iota_i(H(f))$  and  $B := \iota_i(H(-f))$  of  $X$ . Since  $H(f)$  and  $H(-f)$  are clopen subsets of a compact space, they are compact, and since  $\iota_i$  is continuous,  $A$  and  $B$  are compact. Moreover, since  $X$  is Hausdorff, for every  $x \in A$  and  $y \in B$  there exist clopen neighborhoods  $U_x$  of  $x$  and  $V_{x,y}$  of  $y$  such that  $U_x \cap V_{x,y} = \emptyset$ . Thus  $V_x = \bigcup_{y \in B} V_{x,y}$  forms an open cover of  $B$  such that  $U_x \cap V_x = \emptyset$ . Since  $B$  is compact, there exists a finite subcover of  $V_x$ , and thus  $V_x$  is clopen. Further,  $C = \bigcup_{x \in A} U_x$  forms an open cover of  $A$  that, again, can be assumed finite, so that  $C$  is clopen. Let  $D = \bigcap_{x \in A} V_x$ . Then  $C \cap D = \emptyset$  and  $D$  is clearly closed. Since  $X$  is normal, it follows that there are disjoint open sets  $U$  and  $V$  such that  $C \subset U$ ,  $D \subset V$  and  $U \cap V = \emptyset$ .

$U$  and  $V$  as open sets are unions of clopen sets covering  $A$  and  $B$ , respectively. Using compactness one more time we may assume  $U$  and  $V$  to be clopen. The function  $\phi : X \rightarrow \{\pm 1\}$  given by

$$\phi(x) = \begin{cases} 1 & \text{if } x \in U \\ -1 & \text{if } x \in X \setminus U \end{cases}$$

is clearly continuous and, as  $V \subset X \setminus U$ ,  $\phi|_{X_i} = f$ . □

**Lemma 2.** *(cf. [4, Lemma 8.6]) For all  $i \in I$ , and for every  $\phi \in \text{Cont}(X, \{\pm 1\})$  such that  $\phi|_{X_i} \in G_i$  there exists a clopen neighborhood  $J_i$  of  $i$  such that  $\phi|_{X_j} \in G_j$ , for  $j \in J_i$ .*

*Proof.* Fix  $i \in I$  and  $\phi \in \text{Cont}(X, \{\pm 1\})$ . Suppose, a *contrario*, that for every clopen neighborhood  $J$  of  $i$  there exists  $j_J \in J$  such that  $\phi|_{X_{j_J}} \notin G_{j_J}$ . By [5, Corollary 3.2.4], for each  $j_J$  there exists a 4-element fan  $\sigma_1^{j_J}, \sigma_2^{j_J}, \sigma_3^{j_J}, \sigma_4^{j_J}$  in  $(X_{j_J}, G_{j_J})$  such that

$$\phi(\sigma_1^{j_J}) \cdot \phi(\sigma_2^{j_J}) \cdot \phi(\sigma_3^{j_J}) \cdot \phi(\sigma_4^{j_J}) = -1.$$

Direct the family  $\mathcal{J}$  of all clopen neighborhoods  $J$  of  $i$  by inclusion. The net  $(i_J)_{J \in \mathcal{J}}$  is then convergent to  $i$ . The net  $(\sigma_1^{i_J})_{J \in \mathcal{J}}$  is then a net in the compact space  $X$  and thus has a cluster point  $\sigma_1$ . Let  $(\sigma_1^{i_J})_{J \in \mathcal{J}_1}$  be a cofinal subnet of  $(\sigma_1^{i_J})_{J \in \mathcal{J}}$  that converges to  $\sigma_1$ ,  $\mathcal{J}_1 \subset \mathcal{J}$ . Similarly, the net  $(\sigma_2^{i_J})_{J \in \mathcal{J}_1}$  is then a net in the compact space  $X$  and thus has a cluster point  $\sigma_2$ . Let  $(\sigma_2^{i_J})_{J \in \mathcal{J}_2}$  be a cofinal subnet of  $(\sigma_2^{i_J})_{J \in \mathcal{J}_1}$  that converges to  $\sigma_2$ ,  $\mathcal{J}_2 \subset \mathcal{J}_1$ . Repeating the argument two more times we eventually get nets  $(\sigma_3^{i_J})_{J \in \mathcal{J}_3}$  convergent to  $\sigma_3$ ,  $\mathcal{J}_3 \subset \mathcal{J}_2$ , and  $(\sigma_4^{i_J})_{J \in \mathcal{J}_4}$  convergent to  $\sigma_4$ ,  $\mathcal{J}_4 \subset \mathcal{J}_3$ . Then, as each constructed net is a subnet of the previous one,  $(\sigma_1, \sigma_2, \sigma_3, \sigma_4)$  is the limit of the net

$$(\sigma_1^{j_J}, \sigma_2^{j_J}, \sigma_3^{j_J}, \sigma_4^{j_J})_{J \in \mathcal{J}_4}.$$

But then

$$\phi(\sigma_1^{j_J}) \cdot \phi(\sigma_2^{j_J}) \cdot \phi(\sigma_3^{j_J}) \cdot \phi(\sigma_4^{j_J}) = -1,$$

for  $J \in \mathcal{J}_4$ , and by continuity of  $\phi$  also

$$\phi(\sigma_1) \cdot \phi(\sigma_2) \cdot \phi(\sigma_3) \cdot \phi(\sigma_4) = -1,$$

which contradicts axiom (4) of a sheaf. □

**Corollary 1.** (cf. [4, Lemma 8.7]) For all  $i \in I$  the restriction map  $G \rightarrow G_i$  given by  $f \mapsto f|_{X_i}$  is surjective.

*Proof.* This follows immediately from Lemmas 1 and 2 □

**Lemma 3.** (cf. [4, Lemma 8.8]) For all  $i \in I$ , and for every open neighborhood  $U$  of  $X_i$  there exist a clopen neighborhood  $K_i$  of  $i$  such that  $X_i \subset \pi^{-1}(K_i) \subset U$ .

*Proof.* This follows directly from the fact that  $X \setminus U$  is compact. □

We now proceed to the proof of Theorem 1

*Proof.* As in the discussion preceding the proof, let

$$P(\underline{a}) = \exists \underline{t} \bigwedge_{j=1}^m p_j(\underline{t}, \underline{a}) \in D_X(1, q_j(\underline{t}, \underline{a}))$$

be a pp formula that holds on every finite subspace of  $(X, G)$ . Here  $\underline{t} = (t_1, \dots, t_n)$ ,  $\underline{a} = (a_1, \dots, a_k)$ , for  $a_1, \dots, a_k \in G$ , and  $p_j(\underline{t}, \underline{a})$ ,  $q_j(\underline{t}, \underline{a})$  are  $\pm$  products of some of the  $t_s$ 's and  $a_r$ 's,  $s \in \{1, \dots, n\}$ ,  $r \in \{1, \dots, k\}$ , for  $j \in \{1, \dots, m\}$ . Our goal is to show that  $P(\underline{a})$  holds in  $(X, G)$ . Let  $a_r^i := a_r|_{X_i}$ , for  $r \in \{1, \dots, k\}$  and  $i \in I$  and define

$$P^i(\underline{a}^i) = \exists \underline{t}^i \bigwedge_{j=1}^m p_j(\underline{t}^i, \underline{a}^i) \in D_{X_i}(1, q_j(\underline{t}^i, \underline{a}^i)),$$

where  $\underline{t}^i = (t_1^i, \dots, t_n^i)$ ,  $\underline{a}^i = (a_1^i, \dots, a_k^i)$ .

$P^i(\underline{a}^i)$  holds on every space  $(X_i, G_i)$ ,  $i \in I$ . Indeed, fix  $i \in I$  and fix a finite subspace  $\{y_1, \dots, y_l\}$  of  $X_i$ . Let  $Y$  be the finite subspace of  $(X, G)$  generated by  $y_1, \dots, y_l$  in the sense of [5, p. 33]. Then  $P(\underline{a})$  holds on  $Y$  with some  $u_1, \dots, u_n \in G$  verifying it, so let  $u_s^i := u_s|_{X_i}$ ,  $s \in \{1, \dots, n\}$ . Then  $P^i(\underline{a}^i)$  holds on  $\{y_1, \dots, y_l\}$  with  $u_1^i, \dots, u_n^i \in G_i$  verifying it. Since  $\{y_1, \dots, y_l\}$  was chosen arbitrarily and since the pp conjecture holds true for the space  $(X_i, G_i)$ ,  $P^i(\underline{a}^i)$  holds on  $(X_i, G_i)$  with some  $t_1^i, \dots, t_n^i \in G_i$  verifying it.

By Corollary 1 we may assume that  $t_1^i, \dots, t_n^i \in G$ , for  $i \in I$ . Let

$$U_i = \bigcap_{j=1}^m [H(p_j(\underline{t}^i, \underline{a})) \cup H(-q_j(\underline{t}^i, \underline{a}))], \text{ for } i \in I.$$

Then  $X_i \subset U_i$  and  $P(\underline{a})$  holds true on the open set  $U_i$  with  $t_1^i, \dots, t_n^i$  verifying it. By Lemma 3, let  $K_i$  be a clopen neighborhood of  $i$  such that  $X_i \subset \pi^{-1}(K_i) \subset U_i$ , for  $i \in I$ . By Lemma 2, let  $J_s^i$  be a clopen neighborhood of  $i$  such that  $t_s^i|_{X_j} \in G_j$ , for  $j \in J_s^i$ ,  $s \in \{1, \dots, n\}$ ,  $i \in I$ . Let  $J_i = J_1^i \cap \dots \cap J_n^i$ ,  $i \in I$ . Define the clopen neighborhoods  $I_i = J_i \cap K_i$ ,  $i \in I$ .

Now  $\bigcup_{i \in I} I_i$  is an open cover of the compact space  $I$ , so we may choose  $I_1, \dots, I_\ell$  among  $I_i$ ,  $i \in I$ , so that  $I = I_1 \cup \dots \cup I_\ell$ . Replacing  $I_1, \dots, I_\ell$  by

$$I_1, I_2 \setminus I_1, I_3 \setminus (I_1 \cup I_2), \dots, I_\ell \setminus (I_1 \cup \dots \cup I_{\ell-1}),$$

if necessary, we might as well assume  $I_1, \dots, I_\ell$  are pairwise disjoint. Define the functions  $t_s : X \rightarrow \{\pm 1\}$  by

$$t_s(x) = \begin{cases} t_s^1(x) & \text{if } x \in X_i \text{ for } i \in I_1, \\ t_s^2(x) & \text{if } x \in X_i \text{ for } i \in I_2, \\ \vdots & \\ t_s^\ell(x) & \text{if } x \in X_i \text{ for } i \in I_\ell, \end{cases}$$

for  $s \in \{1, \dots, n\}$ . Clearly  $t_s$  is continuous, and  $t_s|_{X_i} \in G_i$ , for  $i \in I$ , so that  $t_s \in G$ ,  $s \in \{1, \dots, n\}$ . Moreover,  $P(\underline{a})$  holds on  $(X, G)$  with  $t_1, \dots, t_n$  verifying it, which concludes the proof.  $\square$

## Acknowledgement

The author would like to express his thanks to the referee for his careful reading and useful improvements that made the presentation more comprehensible.

## References

- [1] V. Astier: On some sheaves of reduced special groups. *Arch. for Math. Logic* 46 (5) (2007) 481–488.
- [2] P. Gładki, M. Marshall: The pp conjecture for spaces of orderings of rational conics. *J. Algebra Appl.* 6 (2) (2007) 245–257.
- [3] P. Gładki, M. Marshall: Quotients of index two and general quotients in a space of orderings. *Fund. Math.* 229 (2015) 255–275.
- [4] M. Marshall: *Abstract Witt rings*. Queen's University, Kingston, Ontario (1980).
- [5] M. Marshall: Spaces of orderings and abstract real spectra. *Lecture Notes in Mathematics* 1636 (1996).
- [6] M. Marshall: Open questions in the theory of spaces of orderings. *J. Symbolic Logic* 67 (1) (2002) 341–352.

*E-mail:* pawel.gladki@us.edu.pl

*Author's addresses:*

INSTITUTE OF MATHEMATICS, UNIVERSITY OF SILESIA, UL. BANKOWA 14, KATOWICE,  
POLAND, 40-007

DEPARTMENT OF COMPUTER SCIENCE, AGH UNIVERSITY OF SCIENCE AND TECHNOLOGY,  
AL. MICKIEWICZA 30, KRAKÓW, POLAND, 30-059

*Received:* 24 August, 2015

*Accepted for publication:* 2 May, 2016

*Communicated by:* Karl Dilcher