



HAL
open science

A Note on Transcendental Power Series Mapping the Set of Rational Numbers into Itself

Diego Marques, Elaine Silva

► **To cite this version:**

Diego Marques, Elaine Silva. A Note on Transcendental Power Series Mapping the Set of Rational Numbers into Itself. *Communications in Mathematics*, 2017, Volume 25 (2017), Issue 1 (1), pp.1 - 4. 10.1515/cm-2017-0001 . hal-03664934

HAL Id: hal-03664934

<https://hal.science/hal-03664934v1>

Submitted on 11 May 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

A Note on Transcendental Power Series Mapping the Set of Rational Numbers into Itself

Diego Marques, Elaine Silva

Abstract. In this note, we prove that there is no transcendental entire function $f(z) \in \mathbb{Q}[[z]]$ such that $f(\mathbb{Q}) \subseteq \mathbb{Q}$ and $\text{den } f(p/q) = F(q)$, for all sufficiently large q , where $F(z) \in \mathbb{Z}[z]$.

1 Introduction

A real number ξ is called a *Liouville number*, if for any real number $\omega > 0$ there exists a rational number p/q , with $q > 1$, such that

$$0 < \left| \xi - \frac{p}{q} \right| < q^{-\omega}.$$

In his pioneer book, Maillet [3, Chapter III] proved that the set of the Liouville numbers is preserved under rational functions with rational coefficients. Based on this result, in 1984, Mahler [2] posed the following question

Question 1. *Are there transcendental entire functions $f(z)$ such that if ξ is any Liouville number, then so is $f(\xi)$?*

Recently, some authors (see [4], [5], [7]) constructed classes of Liouville numbers which are mapped into Liouville numbers by transcendental entire functions. For example, to prove this, Marques and Moreira [4] showed the existence of transcendental entire functions f , such that $f(\mathbb{Q}) \subseteq \mathbb{Q}$ and $\text{den } f(p/q) < q^{8q^2}$, for all $p/q \in \mathbb{Q}$, with $q > 1$ (where $\text{den } z$ denotes the denominator of the rational number z). Moreover, their proof implies that the Mahler's question has an affirmative answer if the answer to the below question is also 'yes' (see also [7, Theorem 2.1]).

2010 MSC: 11J81

Key words: Liouville numbers, Mahler's question, power series

DOI: 10.1515/cm-2017-0001

Question 2. Are there transcendental entire functions $f(z)$ such that $f(\mathbb{Q}) \subseteq \mathbb{Q}$ and

$$\text{den } f(p/q) \leq F(q),$$

for some fixed polynomial $F(z) \in \mathbb{Z}[z]$ and for all sufficiently large q ?

In 2015, Marques, Ramirez and Silva [6] proved that the answer for the previous question is ‘no’ for lacunary power series in $\mathbb{Q}[[z]]$ (see [1] for the definition of lacunary power series as well as some results related to their arithmetic properties). Moreover, their proof also implies that there is no transcendental entire function $f(z) \in \mathbb{Q}[[z]]$ such that $f(\mathbb{Q}) \subseteq \mathbb{Q}$ and

$$\text{den } f(p/q) = o(q).$$

In an attempt of answering the previous question, a natural question arises: Could $\text{den } f(p/q)$ be a polynomial in q for all sufficiently large q ?

In this paper, we shall answer this previous question by proving that

Theorem 1. There is no transcendental entire function $f(z) \in \mathbb{Q}[[z]]$ such that $f(\mathbb{Q}) \subseteq \mathbb{Q}$ and

$$\text{den } f(p/q) = F(q),$$

for all sufficiently large q , where $F(z) \in \mathbb{Z}[z]$.

2 The proof

Suppose, towards a contradiction, that for some $F(z) \in \mathbb{Z}[z]$ with degree $m \geq 1$ (the case $m = 0$ was solved in Remark 2.1 of [6]), there exists such a function, say $f(z) = \sum_{k \geq 0} a_k z^k \in \mathbb{Q}[[z]]$.

Thus, for all sufficiently large q , we have that $f(1/q) = n(q)/F(q)$, where $n(q)$ is an integer. Then

$$f\left(\frac{1}{q}\right) - \left(a_0 + \frac{a_1}{q} + \cdots + \frac{a_{m-1}}{q^{m-1}}\right) = \sum_{k \geq m} \frac{a_k}{q^k}.$$

By setting $A = \prod_{i=0}^{m-1} \text{den}(a_i)$, we have

$$A \frac{n(q)}{F(q)} - \left(b_0 + \frac{b_1}{q} + \cdots + \frac{b_{m-1}}{q^{m-1}}\right) = \frac{Aa_m}{q^m} + A \sum_{k \geq m+1} \frac{a_k}{q^k},$$

where $b_i = Aa_i \in \mathbb{Z}$. Therefore,

$$A \frac{n(q)}{F(q)} - \frac{C(q)}{q^{m-1}} = \frac{Aa_m}{q^m} + A \sum_{k \geq m+1} \frac{a_k}{q^k},$$

where $C(z) \in \mathbb{Z}[z]$ has degree $\leq m - 1$. After multiplying by $F(q)$, we obtain

$$An(q) - \frac{D(q)}{q^{m-1}} = \frac{Aa_m F(q)}{q^m} + A \sum_{k \geq m+1} \frac{a_k F(q)}{q^k},$$

where $D(z) \in \mathbb{Z}[z]$ has degree $\leq 2m - 1$. However, we can write $D(q)/q^{m-1} = E(q)/q^{m-1} + G(q)$, where $E, G \in \mathbb{Z}[z]$, $\deg E \leq m - 2$ and $\deg G \leq m$. Now, write

$$An(q) - G(q) = \frac{Aa_m F(q)}{q^m} + A \sum_{k \geq m+1} \frac{a_k F(q)}{q^k} + \frac{E(q)}{q^{m-1}}. \quad (1)$$

Now, we want to evaluate the limit in the right-hand side above when $q \rightarrow \infty$. Let ϵ be the leading coefficient of $F(z)$ (which we can assume to be ≥ 1). Note that $\lim_{q \rightarrow \infty} F(q)/q^m = \epsilon$ and $\lim_{q \rightarrow \infty} E(q)/q^{m-1} = 0$. Now, we need to calculate the limit of the summatory. For that, take a real number δ such that $0 < \delta < 1/\epsilon \leq 1$. Then $\delta^k < 1/\epsilon$, for all $k \geq m + 1$. Thus,

$$\frac{q^k}{\delta^k} \geq \frac{q^{m+1}}{\delta^k} > qF(q)$$

for all sufficiently large q (since the degree of $zF(z)$ is $m + 1$ and its leading coefficient is ϵ). Hence,

$$\frac{|F(q)|}{q^k} = \frac{F(q)}{q^k} < \frac{1}{q\delta^k}$$

(for all sufficiently large q and for all $k \geq m + 1$) and so

$$\left| \sum_{k \geq m+1} \frac{a_k F(q)}{q^k} \right| \leq \frac{1}{q} \sum_{k \geq m+1} \frac{|a_k|}{\delta^k}.$$

Since $\sum_{k \geq m+1} |a_k|/\delta^k < \infty$ (by the absolute convergence of $\sum_{k \geq 0} a_k z^k$ in \mathbb{C}), then

$$\lim_{q \rightarrow \infty} \sum_{k \geq m+1} \frac{a_k F(q)}{q^k} = 0.$$

In conclusion, the right-hand side of (1) tends to $Aa_m \epsilon$ as $q \rightarrow \infty$. Therefore, for all sufficiently large q , it holds that

$$0 \leq |An(q) - G(q)| \leq Aa_m \epsilon + 1.$$

Since $An(q) - G(q)$ is an integer, then there exist $t \in \mathbb{Z}$ and an infinite set $S \subseteq \mathbb{N}$ such that $An(q) - G(q) = t$ for all $q \in S$. Thus

$$f\left(\frac{1}{q}\right) = \frac{n(q)}{F(q)} = \frac{G(q) + t}{AF(q)} = \frac{e_0 + e_1 q + \dots + e_m q^m}{d_0 + d_1 q + \dots + d_m q^m} = \frac{P(1/q)}{Q(1/q)}, \quad (2)$$

where $P(z) = \sum_{i=0}^m e_i z^{m-i}$ and $Q(z) = \sum_{i=0}^m d_i z^{m-i}$.

Let r be a positive real number such that $r < \min\{|z| : Q(z) = 0\}$ (observe that $Q(0) = d_m = A\epsilon \neq 0$). Then the function $h(z)$ given by

$$h(z) = \frac{P(z)}{Q(z)}$$

is analytic on the interval $(-r, r)$. Moreover, by (2), we have that the analytic functions $f(z)$ and $h(z)$ coincide on the set $\{1/q : q \in S \cap (1/r, \infty)\} \subseteq (-r, r)$ which has a limit point in $(-r, r)$. Thus, by the identity principle for analytic functions, we have that $f(z) = h(z)$ on $(-r, r)$. In particular, the entire functions $Q(z)f(z)$ and $P(z)$ coincide on $(-r, r)$ yielding, by the same principle, that they have equal values for all $z \in \mathbb{C}$. Hence the function $f(z)$ satisfies $P(z, f(z)) = 0$, for all $z \in \mathbb{C}$, where $P(x, y) = Q(x)y - P(x)$ (which is a nonzero polynomial). However, this contradicts the transcendence of f . The proof is then complete. \square

Acknowledgement

Diego Marques was supported in part by CNPq and FAP/DF, Brazil. Elaine Silva was supported in part by Capes-Brazil.

References

- [1] K. Mahler: Arithmetic properties of lacunary power series with integral coefficients. *J. Austral. Math. Soc.* 5 (1965) 56–64.
- [2] K. Mahler: Some suggestions for further research. *Bull. Austral. Math. Soc.* 29 (1984) 101–108.
- [3] E. Maillet: *Introduction à la Théorie des Nombres Transcendants et des Propriétés Arithmétiques des Fonctions*. Gauthier-Villars, Paris (1906).
- [4] D. Marques, C.G. Moreira: A variant of a question proposed by K. Mahler concerning Liouville numbers. *Bull. Austral. Math. Soc.* 91 (2015) 29–33.
- [5] D. Marques, J. Ramirez: On transcendental analytic functions mapping an uncountable class of U -numbers into Liouville numbers. *Proc. Japan Acad. Ser. A Math. Sci.* 91 (2015) 25–28.
- [6] D. Marques, J. Ramirez, E. Silva: A note on lacunary power series with rational coefficients. *Bull. Austral. Math. Soc.* 93 (2015) 1–3.
- [7] D. Marques, J. Schleisitz: On a problem posed by Mahler. *J. Austral. Math. Soc.* 100 (2016) 86–107.

Authors' address

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE DE BRASÍLIA, BRASÍLIA, 70910-900,
BRAZIL

E-mail: diego@mat.unb.br, elainecris@mat.unb.br

Received: 27 November, 2015

Accepted for publication: 14 July, 2016

Communicated by: Karl Dilcher