



**HAL**  
open science

# Erratum to: Variations by generalized symmetries of local Noether strong currents equivalent to global canonical Noether currents

Marcella Palese

► **To cite this version:**

Marcella Palese. Erratum to: Variations by generalized symmetries of local Noether strong currents equivalent to global canonical Noether currents. *Communications in Mathematics*, 2017, Volume 25 (2017), Issue 1 (1), pp.71 - 72. 10.1515/cm-2017-0007 . hal-03664939

**HAL Id: hal-03664939**

**<https://hal.science/hal-03664939v1>**

Submitted on 11 May 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

## Erratum to: Variations by generalized symmetries of local Noether strong currents equivalent to global canonical Noether currents

Marcella Palese

**Abstract.** We correct misprints in a formula in the last sentence at the end of page 129; the first paragraph of subsection 4.1; misprints at the end of page 132 and in Proposition 1 at page 133 of the paper ‘Variations by generalized symmetries of local Noether strong currents equivalent to global canonical Noether currents’, Communications in Mathematics 24 (2016), 125–135. DOI: 10.1515/cm-2016-0009

### 1 Errata corrige

- The formulae in the last sentence at the end of page 129 should read as follows.

$$\begin{aligned}\mathcal{E}_{n+1}(\eta) &= 0 & \mathfrak{d}\eta &= 0 \\ \delta_n(\eta) \neq 0 &\implies \mathfrak{d}\lambda_i \neq 0,\end{aligned}$$

- The first paragraph of subsection 4.1 should read as follows.

Suppose that  $\mathfrak{d}\mathcal{L}_{j^r\Xi}\lambda_i = 0$ . We are here particularly concerned with the case when  $\mathcal{L}_{j^r\Xi}\lambda_i \neq 0$  i.e.  $j^r\Xi$  is a generalized symmetry, that means  $\mathcal{L}_{j^r\Xi}\lambda_i =_{\text{loc.}} d_H\beta_i$ . In fact, since  $\mathcal{E}_n(\mathcal{L}_{j^r\Xi}\lambda_i) = 0$  then  $\mathcal{L}_{j^r\Xi}\lambda_i$  defines a cohomology class and we have  $\delta_{n-1}(\mathcal{L}_{j^r\Xi}\lambda_i) \neq 0$  which implies  $\mathfrak{d}\beta_i \neq 0$ . Notice that if  $\mathfrak{d}\mathcal{L}_{j^r\Xi}\lambda_i = 0$  then  $0 = \mathfrak{d}d_H\beta_i = d_H\mathfrak{d}\beta_i$ .

- the last line at page 132 should read:

[...] always satisfied for vertical generalized symmetries of global dynamical forms.

- Proposition 1 at page 133 should read as follows.

**Proposition 1.** *Let  $j^r\Xi$  be a vertical generalized symmetry. The coboundary of the strong Noether currents is locally exact, i.e.  $d_H(\mathfrak{d}(\nu_i + \epsilon_i)) = 0$ .*

*Proof.* To get the assertion it is enough to prove that  $\mathfrak{d}\mathcal{L}_{j^r\Xi}\lambda_i = 0$ . To prove this we note that we have, identically,

$$\delta_n \mathcal{E}_n(\mathfrak{d}\mathcal{L}_{j^r\Xi}\lambda_i) = 0.$$

Indeed, by linearity

$$\delta_n \mathcal{E}_n(\mathfrak{d}\mathcal{L}_{j^r\Xi}\lambda_i) = \delta_n \mathfrak{d}\mathcal{E}_n(\mathcal{L}_{j^r\Xi}\lambda_i) = \delta_n \mathfrak{d}\mathcal{L}_{j^r\Xi}\mathcal{E}_n(\lambda_i),$$

and being  $j^r\Xi$  a generalized symmetry  $\mathcal{L}_{j^r\Xi}\mathcal{E}_n(\lambda_i) = 0$ , thus we get immediately the result, because this implies  $\mathfrak{d}(\mathfrak{d}\mathcal{L}_{j^r\Xi}(\lambda_i) - i(d_H\bar{\beta}_i)) = 0$ , i.e.  $d_H\bar{\beta}_i = \mathfrak{d}d_H\bar{\psi}_i$ ; for vertical generalized symmetries we have  $\mathfrak{d}\bar{\psi}_i = d_H\hat{\psi}_{ij}$  which gives us the results.

Note that the result holds true for any generalized symmetry (not necessarily vertical) of  $\eta_{\lambda_i}$  generating Noether conserved currents for Lagrangians of the type  $\lambda_i - d_H\mu_i$ , with  $\mu_i$  satisfying  $d_H\beta_i = d_H\mathcal{L}_{j^r\Xi}\mu_i$  and  $\mathcal{L}_{j^r\Xi}\mu_i = d_H\psi_i + \mathfrak{d}\bar{\psi}_i$ . A necessary condition for any generalized symmetry to generate Noether strong currents with closed coboundary is  $\mathcal{L}_{j^r\Xi}\mathcal{L}_{j^r\Xi}\mathfrak{d}\lambda_i = 0$ .  $\square$

*Author's address:*

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORINO, VIA C. ALBERTO 10, I-10123  
TORINO, ITALY

*E-mail:* marcella.palese@unito.it

*Received:* 10 April, 2017