



HAL
open science

Geometry of Mus-Sasaki metric

Abderrahim Zagane, Mustapha Djaa

► **To cite this version:**

Abderrahim Zagane, Mustapha Djaa. Geometry of Mus-Sasaki metric. Communications in Mathematics, 2018, Volume 26 (2018), Issue 2 (2), pp.113 - 126. 10.2478/cm-2018-0008 . hal-03664954

HAL Id: hal-03664954

<https://hal.science/hal-03664954v1>

Submitted on 11 May 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

Geometry of Mus-Sasaki metric

Abderrahim Zagane, Mustapha Djaa

Abstract. In this paper, we introduce the Mus-Sasaki metric on the tangent bundle TM as a new natural metric non-rigid on TM . First we investigate the geometry of the Mus-Sasakian metrics and we characterize the sectional curvature and the scalar curvature.

1 Introduction

The geometry of the tangent bundle TM equipped with Sasaki metric has been studied by many authors Sasaki [16], K. Yano and S. Ishihara [18], P. Dombrowski [7], A. Salimov, A. Gezer and N. Cengiz (see [2], [8], [12], [13], [14]) etc. The rigidity of Sasaki metric has incited some geometers to construct and study other metrics on TM . J. Cheeger and D. Gromoll has introduced the notion of Cheeger-Gromoll metric [3], this metric has been studied also by many authors (see [1], [4], [5], [6], [9], [10], [11], [15], [17]).

In this paper, we introduce the Mus-Sasaki metric on the tangent bundle TM as a new natural metric non-rigid on TM . First we investigate the geometry of the Mus-Sasakian metrics and we characterize the sectional curvature (Theorem 4 and Theorem 5) and the scalar curvature (Theorem 6 and Theorem 7).

1.1 Basic Notions and Definition on TM

Let (M, g) be an m -dimensional Riemannian manifold and (TM, π, M) be its tangent bundle. A local chart $(U, x^i)_{i=1, \dots, n}$ on M induces a local chart

$$(\pi^{-1}(U), x^i, y^j)_{i=1, \dots, n}$$

2010 MSC: 53A45, 53C20, 58E20

Key words: Horizontal lift, vertical lift, Mus-Sasaki metric, scalar curvature.

Affiliation:

Abderrahim Zagane – Department of Mathematics, University Ahmed BenBella
 Oran 1

GMFAMI Laboratory, Relizane University, Algeria

E-mail: Zagane2016@gmail.com

Mustapha Djaa – Laboratory of Geometry Analysis and Applications
 University of Saida, Algeria

on TM . Denote by Γ_{ij}^k the Christoffel symbols of g and by ∇ the Levi-Civita connection of g .

We have two complementary distributions on TM , the vertical distribution \mathcal{V} and the horizontal distribution \mathcal{H} , defined by:

$$\begin{aligned}\mathcal{V}_{(x,u)} &= \text{Ker}(d\pi_{(x,u)}) = \left\{ a^i \frac{\partial}{\partial y^i} \Big|_{(x,u)} : a^i \in \mathbb{R} \right\} \\ \mathcal{H}_{(x,u)} &= \left\{ a^i \frac{\partial}{\partial x^i} \Big|_{(x,u)} - a^i u^j \Gamma_{ij}^k \frac{\partial}{\partial y^k} \Big|_{(x,u)} : a^i \in \mathbb{R} \right\},\end{aligned}$$

where $(x, u) \in TM$, such that $T_{(x,u)}TM = \mathcal{H}_{(x,u)} \oplus \mathcal{V}_{(x,u)}$.

Let $X = X^i \frac{\partial}{\partial x^i}$ be a local vector field on M . The vertical and the horizontal lifts of X are defined by

$$X^V = X^i \frac{\partial}{\partial y^i} \tag{1}$$

$$X^H = X^i \frac{\delta}{\delta x^i} = X^i \left\{ \frac{\partial}{\partial x^i} - y^j \Gamma_{ij}^k \frac{\partial}{\partial y^k} \right\} \tag{2}$$

For consequences, we have $(\frac{\partial}{\partial x^i})^H = \frac{\delta}{\delta x^i}$ and $(\frac{\partial}{\partial x^i})^V = \frac{\partial}{\partial y^i}$, then $(\frac{\delta}{\delta x^i}, \frac{\partial}{\partial y^i})_{i=1, \dots, n}$ is a local adapted frame on TTM .

Proposition 1. (see [18]) *Let (M, g) be a Riemannian manifold and R its tensor curvature, then for all vector fields $X, Y \in \Gamma(TM)$ and $p \in T^2M$ we have:*

$$\begin{aligned}[X^H, Y^H]_p &= [X, Y]_p^H - (R_x(X, Y)u)^V, \\ [X^H, Y^V]_p &= (\nabla_X Y)_p^V, \\ [X^V, Y^V]_p &= 0,\end{aligned}$$

where $p = (x, u)$.

Definition 1. Let (M, g) be a Riemannian manifold and $K: TM \rightarrow TM$ be a smooth bundle endomorphism of TM . Then the vertical and horizontal vector fields VK and HK are defined on TM by

$$\begin{aligned}VK: TM &\rightarrow TTM \\ (x, u) &\mapsto (K(u))^V, \\ HK: TM &\rightarrow TTM \\ (x, u) &\mapsto (K(u))^H,\end{aligned}$$

locally we have

$$VK = y^i K_i^j \frac{\partial}{\partial y^j} = y^i \left(K \left(\frac{\partial}{\partial x^i} \right) \right)^V \tag{3}$$

$$HK = y^i K_i^j \frac{\partial}{\partial x^j} - y^i y^k K_i^j \Gamma_{jk}^s \frac{\partial}{\partial y^s} = y^i \left(K \left(\frac{\partial}{\partial x^i} \right) \right)^H. \tag{4}$$

2 Mus-Sasaki metric.

Definition 2. Let (M, g) be a Riemannian manifold and $f: M \times \mathbb{R} \rightarrow (0, +\infty)$. On the tangent bundle TM , we define a Mus-Sasaki metric noted g_f^S by

$$\begin{aligned} g_f^S(X^H, Y^H)_{(x,u)} &= g_x(X, Y) \\ g_f^S(X^H, Y^V)_{(x,u)} &= 0 \\ g_f^S(X^V, Y^V)_{(x,u)} &= f(x, r)g_x(X, Y) \end{aligned}$$

where $X, Y \in \Gamma(TM)$, $(x, u) \in TM$ and $r = g(u, u)$. The function f is called twisting function.

Note that, if $f = 1$ then g_f^S is the Sasaki metric [18].

Lemma 1. Let (M, g) be a Riemannian manifold, then for all $x \in M$ and $u = u^i \frac{\partial}{\partial x^i} \in T_x M$, we have the following

$$\begin{aligned} X^H(g(u, u))_{(x,u)} &= 0 \\ X^H(g(Y, u))_{(x,u)} &= g(\nabla_X Y, u)_x \\ X^V(g(u, u))_{(x,u)} &= 2g(X, u)_x \\ X^V(g(Y, u))_{(x,u)} &= g(X, Y)_x \end{aligned}$$

Proof. Locally, Lemma 1 follows from formulas (1) and (2). □

From Lemma 2 we obtain

Lemma 2. Let (M, g) be a Riemannian manifold, $F: \mathbb{R}^2 \rightarrow [0, +\infty]$, $\alpha: M \rightarrow (0, +\infty)$ and $\beta: \mathbb{R} \rightarrow (0, +\infty)$ be smooth functions. If $f(x, r) = F(\alpha(x), \beta(r))$, then we have the following

$$\begin{aligned} X^V(f)_{(x,u)} &= 2\beta'(r)g_x(X, u) \frac{\partial F}{\partial t}(\alpha(x), \beta(r)) \\ X^H(f)_{(x,u)} &= g_x(\text{grad}_M \alpha, X) \frac{\partial F}{\partial s}(\alpha(x), \beta(r)) = X(\alpha) \frac{\partial F}{\partial s}(\alpha(x), \beta(r)) \end{aligned}$$

where $(x, u) \in TM$ and $r = g_x(u, u)$.

In the following, we consider $f(x, r) = F(\alpha(x), \beta(r))$, where $F: \mathbb{R}^2 \rightarrow [0, +\infty]$, $\alpha: M \rightarrow (0, +\infty)$ and $\beta: \mathbb{R} \rightarrow (0, +\infty)$ are smooth functions.

Lemma 3. Let (M, g) be a Riemannian manifold. If $f(x, r) = F(\alpha(x), \beta(r))$ and ∇ (resp. $\tilde{\nabla}$) denote the Levi-Civita connection of (M, g) (resp. (TM, g_f^S)), then we have:

$$\begin{aligned} g_f^S(\tilde{\nabla}_{X^H} Y^H, Z^H) &= g_f^S((\nabla_X Y)^H, Z^H) \\ g_f^S(\tilde{\nabla}_{X^H} Y^H, Z^V) &= -\frac{1}{2}g_f^S((R(X, Y)u)^V, Z^V) \end{aligned}$$

$$\begin{aligned}
g_f^S(\tilde{\nabla}_{X^H} Y^V, Z^H) &= g_f^S\left(\frac{f}{2}(R(u, Y)X)^H, Z^H\right) \\
g_f^S(\tilde{\nabla}_{X^H} Y^V, Z^V) &= g_f^S\left(\frac{1}{2}X(\alpha)\frac{\partial \ln F}{\partial s}Y^V + (\nabla_X Y)^V, Z^V\right) \\
g_f^S(\tilde{\nabla}_{X^V} Y^H, Z^H) &= g_f^S\left(\left(\frac{f}{2}R(u, X)Y\right)^H, Z^H\right) \\
g_f^S(\tilde{\nabla}_{X^V} Y^H, Z^V) &= g_f^S\left(\frac{1}{2}Y(\alpha)\frac{\partial \ln F}{\partial s}X^V, Z^V\right) \\
g_f^S(\tilde{\nabla}_{X^V} Y^V, Z^H) &= g_f^S\left(-\frac{1}{2}g(X, Y)\frac{\partial F}{\partial s}(\text{grad}(\alpha))^H, Z^H\right) \\
g_f^S(\tilde{\nabla}_{X^V} Y^V, Z^V) &= \beta'\frac{\partial \ln F}{\partial t}g_f^S(g(X, u)Y^V + g(Y, u)X^V - g(X, Y)U^V, Z^V)
\end{aligned}$$

The proof of Lemma 3 follows directly from Kozul formula, Definition 2, Lemma 1 and Lemma 2.

As a direct consequence of Lemma 3, we get the following theorem.

Theorem 1. *Let (M, g) be a Riemannian manifold. If $f(x, r) = F(\alpha(x), \beta(r))$ and ∇ (resp. $\tilde{\nabla}$) denote the Levi-Civita connection of (M, g) (resp. (TM, g_f^S)), then we have:*

$$\begin{aligned}
(\tilde{\nabla}_{X^H} Y^H)_p &= (\nabla_X Y)_p^H - \frac{1}{2}(R_x(X, Y)u)^V \\
(\tilde{\nabla}_{X^H} Y^V)_p &= (\nabla_X Y)_p^V + \frac{f(x, r)}{2}(R_x(u, Y)X)^H + \frac{1}{2}X(\alpha)\frac{\partial \ln F}{\partial s}(\alpha(x), \beta(r))Y_p^V \\
(\tilde{\nabla}_{X^V} Y^H)_p &= \frac{f(x, r)}{2}(R_x(u, X)Y)^H + \frac{1}{2}Y(\alpha)\frac{\partial \ln F}{\partial s}(\alpha(x), \beta(r))X_p^V \\
(\tilde{\nabla}_{X^V} Y^V)_p &= \beta'(r)\frac{\partial \ln F}{\partial t}(\alpha(x), \beta(r))[g_x(Y, U)X_p^V + g_x(X, U)Y_p^V - g_x(X, Y)U_p^V] \\
&\quad - \frac{1}{2}g_x(X, Y)\frac{\partial F}{\partial s}(\alpha(x), \beta(r))(\text{grad}_M \alpha)_p^H
\end{aligned}$$

for all vector fields $X, Y \in \Gamma(TM)$ and $p = (x, u) \in TM$, where R denotes the curvature tensor of (M, g) .

From Definition 1 and Theorem 1 we have:

Proposition 2. *Let (M, g) be a Riemannian manifold and $\tilde{\nabla}$ be the Levi-Civita connection of the tangent bundle (TM, g_f^S) equipped with the Mus-Sasaki metric. If K is a tensor field of type $(1, 1)$ on M , then:*

$$\begin{aligned}
(\tilde{\nabla}_{X^H} H K)_{(x, u)} &= H(\nabla_X K)(x, u) - \frac{1}{2}\left(R_x(X_x, K_x(u))u\right)^V \\
(\tilde{\nabla}_{X^H} V K)_{(x, u)} &= V(\nabla_X K)(x, u) + \frac{f(x, r)}{2}\left(R_x(u, K_x(u))X_x\right)^H \\
&\quad + \frac{1}{2}X(\alpha)\frac{\partial \ln F}{\partial s}(\alpha(x), \beta(r))(V K)_p
\end{aligned}$$

$$\begin{aligned}
 (\tilde{\nabla}_{X^V} HK)_{(x,u)} &= (K(X))_{(x,u)}^H + \frac{f(x,r)}{2} \left(R_x(u, X_x) K_x(u) \right)^H \\
 &\quad + \frac{1}{2} g_x(\text{grad}_M \alpha, K_x(u)) \frac{\partial \ln F}{\partial s}(\alpha(x), \beta(r)) X_p^V \\
 (\tilde{\nabla}_{X^V} VK)_{(x,u)} &= (K(X))_{(x,u)}^V - \frac{1}{2} g_x(X, K_x(u)) \frac{\partial F}{\partial s}(\alpha(x), \beta(r)) (\text{grad}_M \alpha)_p^H \\
 &\quad + \beta'(r) \frac{\partial \ln F}{\partial t}(\alpha(x), \beta(r)) \left[g_x(K_x(u), u) X_p^V \right. \\
 &\quad \left. + g_x(X, U)(VK)_p - g_x(X, K_x(u)) U_p^V \right]
 \end{aligned}$$

where $(x, u) \in TM$ and $X \in \Gamma(TM)$.

3 Curvatures of Mus-Sasaki metric.

Theorem 2. Let (M, g) be a Riemannian manifold and (TM, g_f^S) its tangent bundle equipped with the Mus-Sasaki metric. If R (resp. \tilde{R}) denote the Riemann curvature tensor of M (resp. TM), then we have the following formulae.

$$\begin{aligned}
 \tilde{R}_{(x,u)}(X^H, Y^H)Z^H &= \left\{ \frac{1}{2}(\nabla_Z R)(X, Y)u + \frac{1}{2}Z(\alpha) \frac{\partial F}{\partial s}(\alpha(x), \beta(r))R(X, Y) \right. \\
 &\quad \left. - \frac{1}{4}X(\alpha) \frac{\partial \ln F}{\partial s}(R(Y, Z)u + \frac{1}{4}Y(\alpha) \frac{\partial \ln F}{\partial s}R(X, Z)u) \right\}_x^V \\
 &\quad + \left\{ R(X, Y)Z + \frac{f(x,r)}{4}R(u, R(Z, Y)u)X \right. \\
 &\quad \left. + \frac{f(x,r)}{4}R(u, R(X, Z)u)Y + \frac{f(x,r)}{2}R(u, R(X, Y)u)Z \right\}_x^H \quad (5)
 \end{aligned}$$

$$\begin{aligned}
 \tilde{R}(X^H, Y^H)Z^V &= \left\{ R(X, Y)Z - \frac{f(x,r)}{4}R(X, R(u, Z)Y)u \right. \\
 &\quad \left. + \frac{f(x,r)}{4}R(Y, R(u, Z)X)u \right. \\
 &\quad \left. + \beta'(r) \frac{\partial \ln F}{\partial t} [g(Z, u)R(X, Y)u - g(Z, R(X, Y)u)u] \right\}^V \quad (6) \\
 &\quad + \left\{ \frac{f(x,r)}{2} [(\nabla_X R)(u, Z)Y - (\nabla_Y R)(u, Z)X \right. \\
 &\quad \left. + \frac{1}{2}X(\alpha) \frac{\partial \ln F}{\partial s}R(u, Z)Y - \frac{1}{2}Y(\alpha) \frac{\partial \ln F}{\partial s}R(u, Z)X] \right. \\
 &\quad \left. - \frac{1}{2}g(R(X, Y)u, Z) \frac{\partial \ln F}{\partial s} \text{grad}_M(\alpha) \right\}^H
 \end{aligned}$$

$$\begin{aligned}
\tilde{R}(X^H, Y^V)Z^V &= \left\{ \beta' X(\alpha) \frac{\partial^2 \ln F}{\partial s \partial t} \left[g(Z, U)Y + g(Y, U)Z - g(Y, Z)U \right] \right. \\
&\quad - \beta' \frac{\partial \ln F}{\partial t} g_x(Y, Z)(\nabla_X U) - \frac{f}{4} g(\text{grad}_M \alpha, R(u, Z)X) \frac{\partial \ln F}{\partial s} Y \\
&\quad - \left. \beta' X(\alpha) g(Y, u) \frac{\partial^2 \ln F}{\partial t \partial s} Z + \frac{1}{4} g(Y, Z) \frac{\partial F}{\partial s} (R(X, \text{grad}_M \alpha)u) \right\}^V \\
&\quad + \left\{ \frac{f}{2} \beta' \frac{\partial \ln F}{\partial t} \left[g(Z, u)(R(u, Y)X) + g(Y, u)(R(u, Z)X) \right] g \right. \\
&\quad - \frac{1}{2} X(\alpha) g_x(Y, Z) \frac{\partial^2 F}{\partial s^2} (\text{grad}_M \alpha) \\
&\quad - \frac{1}{2} g(Y, Z) \frac{\partial F}{\partial s} (\nabla_X \text{grad}_M \alpha) - \beta' g(Y, u) \frac{\partial F}{\partial t} (R(u, Z)X) \\
&\quad - \frac{f}{2} (R(Y, Z)X)^H - \frac{f^2}{4} (R(u, Y)R(u, Z)X) \\
&\quad \left. + \frac{f}{4} X(\alpha) g(Y, Z) \left(\frac{\partial \ln F}{\partial s} \right)^2 (\text{grad}_M \alpha) \right\}^H
\end{aligned} \tag{7}$$

$$\begin{aligned}
\tilde{R}(X^H, Y^V)Z^H &= \left\{ \frac{1}{2} R(X, Z)Y - \frac{f(x, r)}{4} R(X, R(u, Y)Z)u + \frac{1}{2} X(Z(\alpha)) \frac{\partial \ln F}{\partial s} Y \right. \\
&\quad + \frac{1}{2} X(\alpha) Z(\alpha) \left(\frac{\partial^2 \ln F}{\partial s^2} + \frac{1}{2} \left(\frac{\partial \ln F}{\partial s} \right)^2 \right) Y - \frac{1}{2} (\nabla_X Z)(\alpha) \frac{\partial \ln F}{\partial s} Y \\
&\quad + \left. \frac{\beta'(r)}{2} \frac{\partial \ln F}{\partial t} \left[g(Y, u)R(X, Z)u - g(Y, R(X, Z)u)U \right] \right\}^V \\
&\quad + \left\{ \frac{1}{2} X(\alpha) \frac{\partial F}{\partial s} R(u, Y)Z + \frac{f(x, r)}{4} Z(\alpha) \frac{\partial \ln F}{\partial s} R(u, Y)X \right. \\
&\quad \left. + \frac{f(x, r)}{2} (\nabla_X R)(u, Y)Z - \frac{1}{4} g(Y, R(X, Z)u) \frac{\partial F}{\partial s} \text{grad}_M(\alpha) \right\}^H
\end{aligned} \tag{8}$$

$$\begin{aligned}
\tilde{R}_{(x, u)}(X^V, Y^V)Z^H &= \left\{ \beta' \frac{\partial F}{\partial t} \left[g(X, U)R(u, Y)Z - g(Y, U)R(u, X)Z \right] + fR(X, Y)Z \right. \\
&\quad + \left. \frac{f^2}{4} \left[R(u, X)R(u, Y)Z - R(u, Y)R(u, X)Z \right] \right\}_x^H \\
&\quad + \left\{ \beta' Z(\alpha) \frac{\partial^2 \ln F}{\partial s \partial t} \left[g(X, u)Y - g(Y, u)X \right] \right. \\
&\quad + \frac{f}{4} \frac{\partial \ln F}{\partial s} \left[g(\text{grad}_M \alpha, R(u, Y)Z)X \right. \\
&\quad \left. - g(\text{grad}_M \alpha, R(u, X)Z)Y \right] \Big\}_x^V
\end{aligned} \tag{9}$$

$$\begin{aligned}
 \tilde{R}_{(x,u)}(X^V, Y^V)Z^V &= \left\{ 2 \left[\beta' \frac{\partial \ln F}{\partial t} + (\beta')^2 \frac{\partial^2 \ln F}{\partial t^2} \right] \left[g(X, u)g(Z, U)Y \right. \right. \\
 &\quad - g(X, u)g(Y, Z)U - g(Y, u)g(Z, U)X + g(Y, u)g(X, Z)U \left. \right] \\
 &\quad + \beta' \frac{\partial \ln F}{\partial t} \left[g(X, Z)Y - g(Y, Z)X \right] \\
 &\quad + (\beta')^2 \left(\frac{\partial \ln F}{\partial t} \right)^2 \left[g(Y, u)g(Z, U)X - g(Y, U)g(X, Z)U \right. \\
 &\quad - g(X, u)g(Z, U)Y + g(X, u)g(Y, Z)U \\
 &\quad \left. - g(Y, Z)\|u\|^2 X + g(X, Z)\|u\|^2 Y \right] \\
 &\quad + \frac{f}{4} \left(\frac{\partial \ln F}{\partial s} \right)^2 \|\text{grad}_M \alpha\|^2 \left[g(X, Z)Y - g(Y, Z)X \right] \Big\}_x^V \\
 &\quad + \left\{ \beta' \frac{\partial \ln F}{\partial t} \frac{\partial F}{\partial s} \left[g(Y, Z)g(X, u) - g(X, Z)g(Y, u) \right] \text{grad}_M \alpha \right. \\
 &\quad + \frac{f}{4} \frac{\partial F}{\partial s} \left[g(X, Z)R(u, Y) \text{grad}_M \alpha - g(Y, Z)R(u, X) \text{grad}_M \alpha \right] \\
 &\quad \left. + \beta' \frac{\partial^2 \ln F}{\partial s \partial t} \left[g(Y, u)g(X, Z) - g(X, u)g(Y, Z) \right] \text{grad}_M \alpha \right\}_x^H
 \end{aligned} \tag{10}$$

for all $(x, u) \in TM$ and $X, Y, Z \in \Gamma(TM)$.

Proof. Let

$$\begin{aligned}
 K_1: TM &\rightarrow TM \\
 u &\mapsto R(Y, Z)u.
 \end{aligned}$$

Then we have:

$$\begin{aligned}
 \tilde{\nabla}_{X^H} \tilde{\nabla}_{Y^H} Z^H &= \tilde{\nabla}_{X^H} (\nabla_Y Z)^H - \frac{1}{2} \tilde{\nabla}_{X^H} V K_1 \\
 &= (\nabla_X \nabla_Y Z)^H - \frac{1}{2} (R(X, \nabla_Y Z)u)^V - \frac{1}{2} (\nabla_X R(Y, Z)u)^V \\
 &\quad - \frac{f}{4} (R_x(u, R_x(Y, Z)u)X)^H - \frac{1}{4} X(\alpha) \frac{\partial \ln F}{\partial s} V K_1
 \end{aligned}$$

$$\begin{aligned}
 \tilde{\nabla}_{X^H} \tilde{\nabla}_{Y^H} Z^H &= (\nabla_X \nabla_Y Z)^H - \frac{1}{2} (R(X, \nabla_Y Z)u)^V - \frac{1}{2} [(\nabla_X R)(Y, Z)u]^V \\
 &\quad - \frac{1}{2} [R(\nabla_X Y, Z)u + R(Y, \nabla_X Z)u]^V \\
 &\quad - \frac{f}{4} (R_x(u, R_x(Y, Z)u)X)^H - \frac{1}{4} X(\alpha) \frac{\partial \ln F}{\partial s} (R(Y, Z)u)^V
 \end{aligned} \tag{11}$$

$$\begin{aligned}
\tilde{\nabla}_{Y^H} \tilde{\nabla}_{X^H} Z^H &= (\nabla_Y \nabla_X Z)^H - \frac{1}{2} (R(Y, \nabla_X Z)u)^V - \frac{1}{2} [(\nabla_Y R)(X, Z)u]^V \\
&\quad - \frac{1}{2} [R(\nabla_Y X, Z)u + R(X, \nabla_Y Z)u]^V \\
&\quad - \frac{f}{4} (R_x(u, R_x(X, Z)u)Y)^H - \frac{1}{4} Y(\alpha) \frac{\partial \ln F}{\partial s} (R(X, Z)u)^V
\end{aligned} \tag{12}$$

$$\begin{aligned}
\tilde{\nabla}_{[X, Y]^H} Z^H &= (\nabla_{[X, Y]} Z)^H - \frac{1}{2} [R([X, Y], Z)u]^V - \frac{f(x, r)}{2} (R(u, R(X, Y)u)Z)^H \\
&\quad - \frac{1}{2} Z(\alpha) \frac{\partial \ln F}{\partial s} (R(X, Y)u)^V.
\end{aligned} \tag{13}$$

From (11), (12), (13) and the second Bianchi identity

$$(\nabla_X R)(Y, Z) + (\nabla_Y R)(Z, X) + (\nabla_Z R)(X, Y) = 0$$

we obtain the formula (5).

The other formulae are obtained by a similar calculation. \square

From Definition 2 and Theorem 2 we get

Theorem 3. *Let (M, g) be a Riemannian manifold and (TM, g_f^S) its tangent bundle equipped with the Mus-Sasaki metric. If R (resp. \tilde{R}) denote the Riemann curvature tensor of M (resp. TM), then we have*

$$\begin{aligned}
g_f^S(\tilde{R}(X^H, Y^H)Y^H, X^H) &= g(R(X, Y)Y, X) - \frac{3f}{4} \|R(Y, X)u\|^2 \\
g_f^S(\tilde{R}(X^H, Y^V)Y^V, X^H) &= \frac{f}{4} |X(\alpha)|^2 \|Y\|^2 \left(\frac{\partial \ln F}{\partial s} \right)^2 - \frac{1}{2} |X(\alpha)|^2 \|Y\|^2 \frac{\partial^2 F}{\partial s^2} \\
&\quad - \frac{1}{2} \|Y\|^2 \frac{\partial F}{\partial s} g(\nabla_X \text{grad}_M \alpha, X) + \frac{f^2}{4} \|R(u, Y)X\|^2 \\
g_f^S(\tilde{R}(X^V, Y^V)Y^V, X^V) &= 2f \left[\beta'' \frac{\partial \ln F}{\partial t} + (\beta')^2 \frac{\partial^2 \ln F}{\partial t^2} \right] \left[2g(X, u)g(Y, u)g(X, Y) \right. \\
&\quad \left. - \|Y\|^2 |g(X, u)|^2 - \|X\|^2 |g(Y, u)|^2 \right] \\
&\quad + f\beta' \frac{\partial \ln F}{\partial t} \left[|g(X, Y)|^2 - \|X\|^2 \|Y\|^2 \right] \\
&\quad + f(\beta')^2 \left(\frac{\partial \ln F}{\partial t} \right)^2 \left[\|X\|^2 |g(Y, u)|^2 - 2g(X, u)g(Y, u)g(X, Y) \right. \\
&\quad \left. + \|Y\|^2 |g(X, u)|^2 - \|X\|^2 \|Y\|^2 \|u\|^2 + |g(X, Y)|^2 \|u\|^2 \right] \\
&\quad + \frac{f^2}{4} \left(\frac{\partial \ln F}{\partial s} \right)^2 \|\text{grad}_M \alpha\|^2 \left[|g(X, Y)|^2 - \|X\|^2 \|Y\|^2 \right].
\end{aligned}$$

for any $X, Y \in \Gamma(TM)$.

By Theorem 3 and definition of sectional curvature tensor we get

Theorem 4. *Let (M, g) be a Riemannian manifold and (TM, g_f^S) its tangent bundle equipped with the Mus-Sasaki metric. If K (resp. \tilde{K}) denote the sectional curvature tensor of M (resp. TM), then for any orthonormal vector fields $X, Y \in \Gamma(TM)$, we have*

$$\begin{aligned} \tilde{K}(X^H, Y^H) &= K(X, Y) - \frac{3f}{4} \|R(X, Y)u\|^2. \\ \tilde{K}(X^H, Y^V) &= \frac{f}{4} \|R(u, Y)X\|^2 + \frac{1}{4} |X(\alpha)|^2 \left(\frac{\partial \ln F}{\partial s} \right)^2 - \frac{1}{2f} |X(\alpha)|^2 \frac{\partial^2 F}{\partial s^2} \\ &\quad - \frac{1}{2f} \frac{\partial F}{\partial s} g(\nabla_X \text{grad}_M \alpha, X) \\ \tilde{K}(X^V, Y^V) &= -\frac{2}{f} \beta'' \frac{\partial \ln F}{\partial t} \left[|g(X, u)|^2 + |g(Y, u)|^2 \right] - \frac{1}{4f^2} \left(\frac{\partial F}{\partial s} \right)^2 \|\text{grad}_M \alpha\|^2 \\ &\quad - \frac{(\beta')^2}{f} \left(\frac{\partial \ln F}{\partial t} \right)^2 \left[\|u\|^2 - |g(X, u)|^2 - |g(Y, u)|^2 \right] - \frac{\beta'}{f} \frac{\partial \ln F}{\partial t} \end{aligned}$$

Theorem 5. *Let (M, g) be a Riemannian manifold of constant sectional curvature λ and (TM, g_f^S) its tangent bundle equipped with the Mus-Sasaki metric. If \tilde{K} denote the sectional curvature tensor of TM , then for any orthonormal vector fields $X, Y \in \Gamma(TM)$, we have*

$$\begin{aligned} \tilde{K}(X^H, Y^H) &= \lambda - \frac{3f\lambda^2}{4} \left[g(X, u)^2 + g(Y, u)^2 \right] \\ \tilde{K}(X^H, Y^V) &= \frac{f\lambda^2}{4} g(u, X)^2 + \frac{1}{4} |X(\alpha)|^2 \left(\frac{\partial \ln F}{\partial s} \right)^2 - \frac{1}{2f} |X(\alpha)|^2 \frac{\partial^2 F}{\partial s^2} \\ &\quad - \frac{1}{2f} \frac{\partial F}{\partial s} g(\nabla_X \text{grad}_M \alpha, X) \\ \tilde{K}(X^V, Y^V) &= -\frac{2}{f} \left[\beta'' \frac{\partial \ln F}{\partial t} + (\beta')^2 \frac{\partial^2 \ln F}{\partial t^2} \right] \left[|g(X, u)|^2 + |g(Y, u)|^2 \right] - \frac{\beta'}{f} \frac{\partial \ln F}{\partial t} \\ &\quad - \frac{(\beta')^2}{f} \left(\frac{\partial \ln F}{\partial t} \right)^2 \left[\|u\|^2 - |g(X, u)|^2 - |g(Y, u)|^2 \right] \\ &\quad - \frac{1}{4} \left(\frac{\partial \ln F}{\partial s} \right)^2 \|\text{grad}_M \alpha\|^2 \end{aligned}$$

The proof of Theorem 5 is deduced from the Theorem 4 and the following equations

$$\begin{aligned} R(X, Y)Z &= \lambda(g(X, Z)Y - g(Y, Z)X) \\ \|R(X, Y)u\|^2 &= \lambda^2 [g(X, u)^2 + g(Y, u)^2] \\ \|R(u, Y)X\|^2 &= \lambda^2 g(u, X)^2 \end{aligned}$$

for all $(x, u) \in TM$ and $X, Y, Z \in \Gamma(TM)$

Theorem 6. Let (M, g) be a Riemannian manifold and (TM, g_f^S) its tangent bundle equipped with the Mus-Sasaki metric. If σ (resp. $\tilde{\sigma}$) denote the scalar curvature of M (resp. TM), then for any local orthonormal frame (E_1, \dots, E_m) on M , we have

$$\begin{aligned} \tilde{\sigma} = & \sigma + \frac{2-3f}{4} \sum_{i,j=1}^m \|R(E_i, E_j)u\|^2 - \frac{m}{f^2} \frac{\partial F}{\partial s} \text{trace}[g(\nabla_{\text{grad}_M} \alpha, *)] \\ & + \frac{m}{f^2} \|\text{grad}(\alpha)\|^2 \left[\frac{1}{2f} \left(\frac{\partial F}{\partial s} \right)^2 - \frac{\partial^2 F}{\partial s^2} - \frac{(m-1)}{4f^2} \left(\frac{\partial F}{\partial s} \right)^2 \right] \\ & - \frac{4(m-1)}{f^3} \left[\beta'' \frac{\partial \ln F}{\partial t} + (\beta')^2 \frac{\partial^2 \ln F}{\partial t^2} + \frac{(m-2)}{4} (\beta')^2 \left(\frac{\partial \ln F}{\partial t} \right)^2 \right] \|u\|^2 \\ & - \frac{m(m-1)}{f^3} \beta' \frac{\partial \ln F}{\partial t} \end{aligned}$$

Proof. Let (E_1, \dots, E_m) be local orthonormal frame on M , then

$$\left(E_1^H, \dots, E_m^H, \frac{1}{\sqrt{f}} E_1^V, \dots, \frac{1}{\sqrt{f}} E_m^V \right)$$

are a local orthonormal frame on TM . Locally we obtain

$$\begin{aligned} \sum_{i,j=1}^m \|R(u, E_i)E_j\|^2 &= \sum_{i,j=1}^m g(R(u, E_i)E_j, R(u, E_i)E_j) \\ &= \sum_{i,j,k,l,s=1}^m u_k u_l g(R(E_k, E_i)E_j, E_s) g(R(E_l, E_i)E_j, E_s) \\ &= \sum_{i,j,k,l,s=1}^m u_k u_l g(R(E_j, E_s)E_k, E_i) g(R(E_j, E_s)E_l, E_i) \\ &= \sum_{i,j,s=1}^m g(R(E_j, E_s)u, g(R(E_j, E_s)u, E_i)E_i) \\ &= \sum_{i,j=1}^m \|R(E_i, E_j)u\|^2 \end{aligned} \tag{14}$$

From Theorem 4 and definition of scalar curvature, we have

$$\tilde{\sigma} = \sum_{i,j=1}^m \tilde{K}(E_i^H, E_j^H) + \frac{2}{f} \sum_{i,j=1}^m \tilde{K}(E_i^H, E_j^V) + \frac{1}{f^2} \sum_{i \neq j=1}^m \tilde{K}(E_i^V, E_j^V)$$

$$\begin{aligned}
 \tilde{\sigma} = & \sigma - \sum_{i,j=1}^m \frac{3f}{4} \|R(E_i, E_j)u\|^2 + \frac{2}{f} \sum_{i,j=1}^m \left[\frac{f}{4} \|R(u, E_j)E_i\|^2 + \frac{1}{4} |E_i(\alpha)|^2 \left(\frac{\partial \ln F}{\partial s} \right)^2 \right. \\
 & - \frac{1}{2f} |E_i(\alpha)|^2 \frac{\partial^2 F}{\partial s^2} - \frac{1}{2f} \frac{\partial F}{\partial s} g(\nabla_{E_i} \text{grad}_M \alpha, E_i) \left. \right] \\
 & - \frac{1}{f^2} \sum_{i \neq j=1}^m \left\{ \frac{2}{f} \left[\beta'' \frac{\partial \ln F}{\partial t} + (\beta')^2 \frac{\partial^2 \ln F}{\partial t^2} \right] \left[|g(E_i, u)|^2 + |g(E_j, u)|^2 \right] \right. \\
 & + \frac{\beta'}{f} \frac{\partial \ln F}{\partial t} + \frac{(\beta')^2}{f} \left(\frac{\partial \ln F}{\partial t} \right)^2 \left[\|u\|^2 - |g(E_i, u)|^2 - |g(E_j, u)|^2 \right] \\
 & \left. + \frac{1}{4} \left(\frac{\partial \ln F}{\partial s} \right)^2 \|\text{grad}_M \alpha\|^2 \right\} \tag{15}
 \end{aligned}$$

$$\begin{aligned}
 \tilde{\sigma} = & \sigma - \sum_{i,j=1}^m \frac{3f}{4} \|R(E_i, E_j)u\|^2 + \frac{1}{2} \sum_{i,j=1}^m \|R(u, E_j)E_i\|^2 + \frac{m}{2f} \|\text{grad}(\alpha)\|^2 \left(\frac{\partial \ln F}{\partial s} \right)^2 \\
 & - \frac{m}{f^2} \|\text{grad}(\alpha)\|^2 \frac{\partial^2 F}{\partial s^2} - \frac{m}{f^2} \frac{\partial F}{\partial s} \text{trace}[g(\nabla_{\text{grad}_M} \alpha, *)] \\
 & - \frac{1}{f^2} \left\{ \frac{4(m-1)}{f} \left[\beta'' \frac{\partial \ln F}{\partial t} + (\beta')^2 \frac{\partial^2 \ln F}{\partial t^2} \right] \|u\|^2 + m(m-1) \frac{\beta'}{f} \frac{\partial \ln F}{\partial t} \right. \\
 & \left. + \frac{(\beta')^2}{f} \left(\frac{\partial \ln F}{\partial t} \right)^2 (m-1)(m-2) \|u\|^2 + \frac{m(m-1)}{4} \left(\frac{\partial \ln F}{\partial s} \right)^2 \|\text{grad}_M \alpha\|^2 \right\}
 \end{aligned}$$

$$\begin{aligned}
 \tilde{\sigma} = & \sigma - \sum_{i,j=1}^m \frac{3f}{4} \|R(E_i, E_j)u\|^2 + \frac{1}{2} \sum_{i,j=1}^m \|R(u, E_j)E_i\|^2 + \frac{m}{2f} \|\text{grad}(\alpha)\|^2 \left(\frac{\partial \ln F}{\partial s} \right)^2 \\
 & - \frac{m}{f^2} \|\text{grad}(\alpha)\|^2 \frac{\partial^2 F}{\partial s^2} - \frac{m}{f^2} \frac{\partial F}{\partial s} \text{trace}[g(\nabla_{\text{grad}_M} \alpha, *)] \\
 & - \frac{4(m-1)}{f^3} \left[\beta'' \frac{\partial \ln F}{\partial t} + (\beta')^2 \frac{\partial^2 \ln F}{\partial t^2} \right] \|u\|^2 - \frac{m(m-1)}{f^3} \beta' \frac{\partial \ln F}{\partial t} \\
 & - \frac{(m-1)(m-2)}{f^3} (\beta')^2 \left(\frac{\partial \ln F}{\partial t} \right)^2 \|u\|^2 - \frac{m(m-1)}{4f^2} \left(\frac{\partial \ln F}{\partial s} \right)^2 \|\text{grad}_M \alpha\|^2
 \end{aligned}$$

Using formula (14) we deduce

$$\begin{aligned}
 \tilde{\sigma} = & \sigma + \frac{2-3f}{4} \sum_{i,j=1}^m \|R(E_i, E_j)u\|^2 + \frac{m}{2f} \|\text{grad}(\alpha)\|^2 \left(\frac{\partial \ln F}{\partial s} \right)^2 \\
 & - \frac{m}{f^2} \|\text{grad}(\alpha)\|^2 \frac{\partial^2 F}{\partial s^2} - \frac{m}{f^2} \frac{\partial F}{\partial s} \text{trace}[g(\nabla_{\text{grad}_M} \alpha, *)] \\
 & - \frac{4(m-1)}{f^3} \left[\beta'' \frac{\partial \ln F}{\partial t} + (\beta')^2 \frac{\partial^2 \ln F}{\partial t^2} \right] \|u\|^2 - \frac{m(m-1)}{f^3} \beta' \frac{\partial \ln F}{\partial t} \\
 & - \frac{(m-1)(m-2)}{f^3} (\beta')^2 \left(\frac{\partial \ln F}{\partial t} \right)^2 \|u\|^2 - \frac{m(m-1)}{4f^2} \left(\frac{\partial \ln F}{\partial s} \right)^2 \|\text{grad}_M \alpha\|^2
 \end{aligned}$$

$$\begin{aligned}
\tilde{\sigma} &= \sigma + \frac{2-3f}{4} \sum_{i,j=1}^m \|R(E_i, E_j)u\|^2 - \frac{m}{f^2} \frac{\partial F}{\partial s} \text{trace}[g(\nabla_{\text{grad}_M} \alpha, *)] \\
&\quad + \frac{m}{f^2} \|\text{grad}(\alpha)\|^2 \left[\frac{1}{2f} \left(\frac{\partial F}{\partial s} \right)^2 - \frac{\partial^2 F}{\partial s^2} - \frac{(m-1)}{4f^2} \left(\frac{\partial F}{\partial s} \right)^2 \right] \\
&\quad - \frac{4(m-1)}{f^3} \left[\beta'' \frac{\partial \ln F}{\partial t} + (\beta')^2 \frac{\partial^2 \ln F}{\partial t^2} + \frac{(m-2)}{4} (\beta')^2 \left(\frac{\partial \ln F}{\partial t} \right)^2 \right] \|u\|^2 \\
&\quad - \frac{m(m-1)}{f^3} \beta' \frac{\partial \ln F}{\partial t} \quad \square
\end{aligned}$$

Corollary 1. *Let (M, g) be a Riemannian manifold of constant sectional curvature λ and (TM, g_f^S) its tangent bundle equipped with the Mus-Sasaki metric. If $\tilde{\sigma}$ denote the scalar curvature of TM , then for any local orthonormal frame (E_1, \dots, E_m) on M , we have*

$$\begin{aligned}
\tilde{\sigma} &= (m-1)\lambda \left[m + \frac{(2-3f)}{2} \lambda \|u\|^2 \right] - \frac{m}{f^2} \frac{\partial F}{\partial s} \text{trace}[g(\nabla_{\text{grad}_M} \alpha, *)] \\
&\quad + \frac{m}{f^2} \|\text{grad}(\alpha)\|^2 \left[\frac{1}{2f} \left(\frac{\partial F}{\partial s} \right)^2 - \frac{\partial^2 F}{\partial s^2} - \frac{(m-1)}{4f^2} \left(\frac{\partial F}{\partial s} \right)^2 \right] \\
&\quad - \frac{4(m-1)}{f^3} \left[\beta'' \frac{\partial \ln F}{\partial t} + (\beta')^2 \frac{\partial^2 \ln F}{\partial t^2} + \frac{(m-2)}{4} (\beta')^2 \left(\frac{\partial \ln F}{\partial t} \right)^2 \right] \|u\|^2 \\
&\quad - \frac{m(m-1)}{f^3} \beta' \frac{\partial \ln F}{\partial t} \quad (16)
\end{aligned}$$

Proof. Taking account that $\sigma = m(m-1)\lambda$ and for any vector fields $X, Y, Z \in TM$

$$R(X, Y)Z = \lambda(g(Y, Z)Y - g(X, Z)Y)$$

then we obtain

$$\begin{aligned}
\sum_{i,j=1}^m \|R(E_i, E_j)u\|^2 &= \lambda^2 \sum_{\substack{i \neq j \\ i,j=1}}^m \|g(E_j, u)E_i - g(E_i, u)E_j\|^2 \\
&= \lambda^2 \sum_{\substack{i \neq j \\ i,j=1}}^m (\|g(E_j, u)\|^2 + \|g(E_i, u)\|^2) \\
&= 2(m-1)\lambda^2 \|u\|^2 \quad (17)
\end{aligned}$$

From Theorem 6 and formula (17) we deduce

$$\begin{aligned}
\tilde{\sigma} &= (m-1)\lambda \left[m + \frac{(2-3f)}{2} \lambda \|u\|^2 \right] - \frac{m}{f^2} \frac{\partial F}{\partial s} \text{trace}[g(\nabla_{\text{grad}_M} \alpha, *)] \\
&\quad + \frac{m}{f^2} \|\text{grad}(\alpha)\|^2 \left[\frac{1}{2f} \left(\frac{\partial F}{\partial s} \right)^2 - \frac{\partial^2 F}{\partial s^2} - \frac{(m-1)}{4f^2} \left(\frac{\partial F}{\partial s} \right)^2 \right] \\
&\quad - \frac{4(m-1)}{f^3} \left[\beta'' \frac{\partial \ln F}{\partial t} + (\beta')^2 \frac{\partial^2 \ln F}{\partial t^2} + \frac{(m-2)}{4} (\beta')^2 \left(\frac{\partial \ln F}{\partial t} \right)^2 \right] \|u\|^2 \\
&\quad - \frac{m(m-1)}{f^3} \beta' \frac{\partial \ln F}{\partial t} \quad (18)
\end{aligned}$$

□

Theorem 7. Let (M, g) be a Riemannian manifold and (TM, g_f^S) its tangent bundle equipped with the Mus-Sasaki metric. If $f = \text{const}$ then (TM, g_f^S) has a constant scalar curvature if and only if $f = \frac{3}{2}$ and (M, g) has a constant scalar curvature, or (M, g) is flat.

Proof. From Theorem 6 we have

$$\tilde{\sigma} = \sigma + \frac{(2-3f)}{4} \sum_{i,j=1}^m \|R(E_i, E_j)u\|^2$$

then $\tilde{\sigma}$ is constant if and only if $\sigma = \text{const}$ and $f = \frac{3}{2}$ or $R = 0$. □

Corollary 2. Let (M, g) be a Riemannian manifold and (TM, g_f^S) its tangent bundle equipped with the Mus-Sasaki metric. If $f = \frac{3}{2}$ then $\tilde{\sigma} = \sigma$.

References

- [1] M.T.K. Abbassi, M. Sarih: On Natural Metrics on Tangent Bundles of Riemannian Manifolds. *Archivum Mathematicum* 41 (2005) 71–92.
- [2] N. Cengiz, A.A. Salimov: Diagonal lift in the tensor bundle and its applications. *Appl. Math. Comput.* 142 (2–3) (2003) 309–319.
- [3] J. Cheeger, D. Gromoll: On the structure of complete manifolds of nonnegative curvature. *Ann. of Math.* 96 (2) (1972) 413–443.
- [4] M. Djaa, N.E.H. Djaa, R. Nasri: Natural Metrics on T^2M and Harmonicity. *International Electronic Journal of Geometry* 6 (1) (2013) 100–111.
- [5] M. Djaa, J. Gancarzewicz: The geometry of tangent bundles of order r. *Boletín Academia, Gaceta de Ciencias* 4 (1985) 147–165.
- [6] N.E.H. Djaa, S. Ouakkas, M. Djaa: Harmonic sections on the tangent bundle of order two. *Annales Mathematicae et Informaticae* 38 (2011) 15–25.
- [7] P. Dombrowski: On the Geometry of the Tangent Bundle. *J. Reine Angew. Math.* 210 (1962) 73–88.
- [8] A. Gezer: On the Tangent Bundle With Deformed Sasaki Metric. *International Electronic Journal of Geometry* 6 (2) (2013) 19–31.
- [9] S. Gudmundsson, E. Kappos: On the Geometry of the Tangent Bundle with the Cheeger-Gromoll Metric. *Tokyo J. Math.* 25 (1) (2002) 75–83.
- [10] O. Kowalski and M. Sekizawa: On Riemannian Geometry Of Tangent Sphere Bundles With Arbitrary Constant Radius. *Archivum Mathematicum* 44 (2008) 391–401.
- [11] E. Musso, F. Tricerri: Riemannian Metrics on Tangent Bundles. *Ann. Mat. Pura Appl.* 150 (4) (1988) 1–19.
- [12] A.A. Salimov, F. Agca: Some Properties of Sasakian Metrics in Cotangent Bundles. *Mediterranean Journal of Mathematics* 8 (2) (2011) 243–255.
- [13] A.A. Salimov, A. Gezer: On the geometry of the $(1,1)$ -tensor bundle with Sasaki type metric. *Chinese Annals of Mathematics* 32 (3) (2011) 369–386.

- [14] A.A. Salimov, A. Gezer, K. Akbulut: Geodesics of Sasakian metrics on tensor bundles. *Mediterr. J. Math* 6 (2) (2009) 135–147.
- [15] A.A. Salimov, S. Kazimova: Geodesics of the Cheeger-Gromoll Metric. *Turk. J. Math.* 33 (2009) 99–105.
- [16] S. Sasaki: On the differential geometry of tangent bundles of Riemannian manifolds II. *Tohoku Math. J.* 14 (1962) 146–155.
- [17] M. Sekizawa: Curvatures of Tangent Bundles with Cheeger-Gromoll Metric. *Tokyo J. Math.* 14 (2) (1991) 407–417.
- [18] K. Yano, S. Ishihara: *Tangent and Cotangent Bundles*. Marcel Dekker. INC., New York (1973).

Received: 04 April, 2017

Accepted for publication: 15 May, 2018

Communicated by: Haizhong Li