



HAL
open science

A Study on Φ -recurrence τ -curvature tensor in (k, μ) -contact metric manifolds

Gurupadavva Ingalahalli, C.S. Bagewadi

► **To cite this version:**

Gurupadavva Ingalahalli, C.S. Bagewadi. A Study on Φ -recurrence τ -curvature tensor in (k, μ) -contact metric manifolds. Communications in Mathematics, 2018, Volume 26 (2018), Issue 2 (2), pp.1 - 10. 10.2478/cm-2018-0009 . hal-03664955

HAL Id: hal-03664955

<https://hal.science/hal-03664955>

Submitted on 11 May 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

A Study on ϕ -recurrence τ -curvature tensor in (k, μ) -contact metric manifolds

Gurupadavva Ingalahalli, C.S. Bagewadi

Abstract. In this paper we study ϕ -recurrence τ -curvature tensor in (k, μ) -contact metric manifolds.

1 Introduction

In [11], S. Tanno introduced the notion of k -nullity distribution of a contact metric manifold as a distribution such that the characteristic vector field ξ of the contact metric manifold belongs to the distribution. The contact metric manifold with ξ belonging to the k -nullity distribution is called $N(k)$ -contact metric manifold and such a manifold is also studied by various authors. Generalizing this notion in 1995, D.E. Blair, T. Koufogiorgos and B.J. Papantoniou [2] introduced the notion of a contact metric manifold with ξ belonging to the (k, μ) -nullity distribution, where k and μ are real constants. In particular, if $\mu = 0$ then the notion of (k, μ) -nullity distribution reduces to the notion of k -nullity distribution.

In [13], M.M. Tripathi and et al. introduced the τ -curvature tensor which consists of known curvatures like conformal, concircular, projective, M -projective, W_i -curvature tensor ($i = 0, \dots, 9$) and W_j^* -curvature tensor ($j = 0, 1$). Further, in [14] and [15] M.M. Tripathi and et al. studied τ -curvature tensor in K -contact, Sasakian and semi-Riemannian manifolds. Later in [6] the authors studied some properties of τ -curvature tensor and they obtained some interesting results.

2010 MSC: 53C15, 53C25, 53D15.

Key words: Contact metric manifold, curvature tensor, Ricci tensor, Ricci operator.

Affiliation:

Gurupadavva Ingalahalli – Department of Mathematics, Kuvempu University,
Shankaraghatta, 577 451, Shimoga, Karnataka, India.

E-mail: gurupadavva@gmail.com

C.S. Bagewadi – Department of Mathematics, Kuvempu University,
Shankaraghatta, 577 451, Shimoga, Karnataka, India.

E-mail: prof_bagewadi@yahoo.co.in

The notion of local symmetry of a Riemannian manifold has been weakened by many authors in several ways to a different extent. In the context of contact geometry the notion of ϕ -symmetry is introduced and studied by E. Boeckx, P. Buecken and L. Vanhecke [3] with several examples. As a weaker version of local symmetry, T. Takahashi [12] introduced the notion of locally ϕ -symmetry on a Sasakian manifold. Generalizing the notion of ϕ -symmetry De et al. [5] introduced the notion of ϕ -recurrent Sasakian manifold. In [4], the authors studied ϕ -recurrent $N(k)$ -contact metric manifolds. Motivated by all these work in this paper we study the ϕ -recurrent τ -curvature tensor in (k, μ) -contact metric manifold.

2 Preliminaries

A $(2n + 1)$ -dimensional differential manifold M is said to have an almost contact structure (ϕ, ξ, η) if it carries a tensor field ϕ of type $(1, 1)$, a vector field ξ and 1-form η satisfying

$$\phi^2 = -I + \eta \otimes \xi, \quad \eta(\xi) = 1, \quad \eta \circ \phi = 0, \quad \phi\xi = 0. \quad (1)$$

Let g be a compatible Riemannian metric with almost contact structure (ϕ, ξ, η) such that,

$$g(\phi X, \phi Y) = g(X, Y) - \eta(X)\eta(Y), \quad (2)$$

$$g(\phi X, Y) = -g(X, \phi Y) \quad g(X, \xi) = \eta(X), \quad (3)$$

where X, Y are vector fields defined on M . Then the structure (ϕ, ξ, η, g) on M is said to have an almost contact metric structure and the manifold M equipped with this structure is called an almost contact metric manifold.

An almost contact metric structure (ϕ, ξ, η, g) becomes a contact metric structure if

$$d\eta(X, Y) = g(X, \phi Y),$$

for all vector fields X, Y on M .

Given a contact metric manifold $M(\phi, \xi, \eta, g)$, we define a $(1, 1)$ tensor field h by $h = \frac{1}{2}\mathfrak{L}\phi$, where \mathfrak{L} denotes the Lie differentiation. Then h is symmetric and satisfies $h\phi = -\phi h$. Also we have $\text{tr}(h) = \text{tr}(\phi h) = 0$ and $h\xi = 0$. Moreover, if ∇ denotes the Riemannian connection on M , then the following relation holds:

$$\nabla_X \xi = -\phi X - \phi hX. \quad (4)$$

In contact metric manifold $M(\phi, \xi, \eta, g)$, the (k, μ) -nullity distribution is

$$p \rightarrow N_p(k, \mu) = \left\{ Z \in T_p M : R(X, Y)Z = k[g(Y, Z)X - g(X, Z)Y] + \mu[g(Y, Z)hX - g(X, Z)hY] \right\}, \quad (5)$$

for all vector fields $X, Y \in T_p M$ and k, μ are real numbers and R is the curvature tensor. Hence, if the characteristic vector field ξ belongs to the (k, μ) -nullity distribution, then we have

$$R(X, Y)\xi = k[\eta(Y)X - \eta(X)Y] + \mu[\eta(Y)hX - \eta(X)hY]. \quad (6)$$

Thus a contact metric manifold satisfying (6) is called a (k, μ) -contact metric manifold. In particular, if $\mu = 0$, then the notion of (k, μ) -nullity distribution reduces to the notion of k -nullity distribution introduced by S. Tanno [11]. In a (k, μ) -contact metric manifold the following relations hold [2], [9]:

$$h^2 = (k - 1)\phi^2, \quad k \leq 1, \quad (7)$$

$$(\nabla_X \phi)Y = g(X + hX, Y)\xi - \eta(Y)[X + hX],$$

$$\begin{aligned} (\nabla_X h)Y &= [(1 - k)g(X, \phi Y) + g(X, h\phi Y)]\xi \\ &\quad + \eta(Y)[h(\phi X + \phi hX)] - \mu\eta(X)\phi hY, \end{aligned}$$

$$R(\xi, X)Y = k[g(X, Y)\xi - \eta(Y)X] + \mu[g(hX, Y)\xi - \eta(Y)hX],$$

$$\begin{aligned} \eta(R(X, Y)Z) &= k[g(Y, Z)\eta(X) - g(X, Z)\eta(Y)] \\ &\quad + \mu[g(hY, Z)\eta(X) - g(hX, Z)\eta(Y)], \end{aligned}$$

$$S(X, \xi) = 2nk\eta(X), \quad (8)$$

$$\begin{aligned} S(X, Y) &= [2(n - 1) - n\mu]g(X, Y) + [2(n - 1) + \mu]g(hX, Y) \\ &\quad + [2(1 - n) + n(2k + \mu)]\eta(X)\eta(Y), \quad n \geq 1, \end{aligned} \quad (9)$$

$$\begin{aligned} QX &= [2(n - 1) - n\mu]X + [2(n - 1) + \mu]hX \\ &\quad + [2(1 - n) + n(2k + \mu)]\eta(X)\xi, \quad n \geq 1, \end{aligned} \quad (10)$$

$$r = 2n[2n - 2 + k - n\mu],$$

$$S(\phi X, \phi Y) = S(X, Y) - 2nk\eta(X)\eta(Y) - 2(2n - 2 + \mu)g(hX, Y), \quad (11)$$

where S is the Ricci tensor of type $(0, 2)$, Q is the Ricci operator, that is, $S(X, Y) = g(QX, Y)$ and r is the scalar curvature of the manifold. From (3), it follows that

$$(\nabla_X \eta)Y = g(X + hX, \phi Y).$$

Definition 1. A (k, μ) -contact metric manifold M is said to be locally ϕ -symmetric if

$$\phi^2((\nabla_W R)(X, Y)Z) = 0,$$

for all vector fields X, Y, Z, W orthogonal to ξ . This notion was introduced by T. Takahashi [12] for Sasakian manifolds.

A field that is at every point and for every direction proportional to its covariant differential is called recurrent. Based on this concept we define the following definition:

Definition 2. A (k, μ) -contact metric manifold M is said to be ϕ -recurrent if and only if there exists a non zero 1-form A such that

$$\phi^2((\nabla_W R)(X, Y)Z) = A(W)R(X, Y)Z,$$

for all arbitrary vector fields X, Y, Z, W which are not necessarily orthogonal to ξ .

If the 1-form A vanishes identically, then the manifold is said to be a locally ϕ -symmetric manifold.

Definition 3. A (k, μ) -contact metric manifold M is said to be ϕ - τ -recurrent if and only if there exists a non zero 1-form A such that

$$\phi^2((\nabla_W \tau)(X, Y)Z) = A(W)\tau(X, Y)Z,$$

for all arbitrary vector fields X, Y, Z, W which are not necessarily orthogonal to ξ .

The τ -curvature tensor [13] is given by

$$\begin{aligned} \tau(X, Y)Z &= a_0R(X, Y)Z + a_1S(Y, Z)X + a_2S(X, Z)Y + a_3S(X, Y)Z \\ &\quad + a_4g(Y, Z)QX + a_5g(X, Z)QY + a_6g(X, Y)QZ \\ &\quad + a_7r[g(Y, Z)X - g(X, Z)Y], \end{aligned} \quad (12)$$

where a_0, \dots, a_7 are all constants on M . For different values of a_0, \dots, a_7 the τ -curvature tensor reduces to the curvature tensor R , quasi-conformal curvature tensor, conformal curvature tensor, conharmonic curvature tensor, concircular curvature tensor, pseudo-projective curvature tensor, projective curvature tensor, M -projective curvature tensor, W_i -curvature tensors ($i = 0, \dots, 9$), W_j^* -curvature tensors ($j = 0, 1$).

3 ϕ - τ -recurrent (k, μ) -contact metric manifold

In this section, we define ϕ - τ -recurrent (k, μ) -contact metric manifold

$$\phi^2((\nabla_W \tau)(X, Y)Z) = A(W)\tau(X, Y)Z, \quad (13)$$

for all vector fields X, Y, Z, W . By using (9) and (10) in (12), we get

$$\begin{aligned} \tau(X, Y)Z &= a_0R(X, Y)Z + a_1[\alpha g(Y, Z)X + \beta g(hY, Z)X + \gamma \eta(Y)\eta(Z)X] \\ &\quad + a_2[\alpha g(X, Z)Y + \beta g(hX, Z)Y + \gamma \eta(X)\eta(Z)Y] \\ &\quad + a_3[\alpha g(X, Y)Z + \beta g(hX, Y)Z + \gamma \eta(X)\eta(Y)Z] \\ &\quad + a_4g(Y, Z)[\alpha X + \beta hX + \gamma \eta(X)\xi] \\ &\quad + a_5g(X, Z)[\alpha Y + \beta hY + \gamma \eta(Y)\xi] \\ &\quad + a_6g(X, Y)[\alpha Z + \beta hZ + \gamma \eta(Z)\xi] \\ &\quad + a_7r[g(Y, Z)X - g(X, Z)Y], \end{aligned} \quad (14)$$

where $\alpha = [2(n-1) - n\mu]$, $\beta = [2(n-1) + \mu]$ and $\gamma = [2(1-n) + n(2k + \mu)]$.

Differentiating (14) with respect to W , we obtain

$$\begin{aligned}
(\nabla_W \tau)(X, Y)Z &= a_0(\nabla_W R)(X, Y)Z \\
&+ a_1 \left[\beta g((\nabla_W h)Y, Z)X + \gamma \{ (\nabla_W \eta)Y \eta(Z)X + (\nabla_W \eta)Z \eta(Y)X \} \right] \\
&+ a_2 \left[\beta g((\nabla_W h)X, Z)Y + \gamma \{ (\nabla_W \eta)X \eta(Z)Y + (\nabla_W \eta)Z \eta(X)Y \} \right] \\
&+ a_3 \left[\beta g((\nabla_W h)X, Y)Z + \gamma \{ (\nabla_W \eta)X \eta(Y)Z + (\nabla_W \eta)Y \eta(X)Z \} \right] \\
&+ a_4 g(Y, Z) \left[\beta (\nabla_W h)X + \gamma \{ (\nabla_W \eta)(X)\xi + \eta(X)\nabla_W \xi \} \right] \\
&+ a_5 g(X, Z) \left[\beta (\nabla_W h)Y + \gamma \{ (\nabla_W \eta)(Y)\xi + \eta(Y)\nabla_W \xi \} \right] \\
&+ a_6 g(X, Y) \left[\beta (\nabla_W h)Z + \gamma \{ (\nabla_W \eta)(Z)\xi + \eta(Z)\nabla_W \xi \} \right] \\
&+ a_7 (\nabla_W r)[g(Y, Z)X - g(X, Z)Y].
\end{aligned} \tag{15}$$

By virtue of (1), (13), we have

$$-(\nabla_W \tau)(X, Y)Z + \eta((\nabla_W \tau)(X, Y)Z)\xi = A(W)\tau(X, Y)Z. \tag{16}$$

By taking an inner product with U , we obtain

$$-g((\nabla_W \tau)(X, Y)Z, U) + \eta((\nabla_W \tau)(X, Y)Z)g(\xi, U) = A(W)g(\tau(X, Y)Z, U). \tag{17}$$

Let $\{e_i : i = 1, 2, \dots, 2n + 1, \}$ be an orthonormal basis of the tangent space at any point of the manifold. Putting $X = U = e_i$ in (17) and taking summation over i , we get

$$-g((\nabla_W \tau)(e_i, Y)Z, e_i) + \eta((\nabla_W \tau)(e_i, Y)Z)g(\xi, e_i) = A(W)g(\tau(e_i, Y)Z, e_i). \tag{18}$$

By using (15) in (18), we obtain

$$\begin{aligned}
&-a_0(\nabla_W S)(Y, Z) - [2na_1 + a_2 + a_5] \left[\beta \left\{ [(1-k)g(W, \phi Y) + g(W, h\phi Y)]\eta(Z) \right. \right. \\
&+ \eta(Y)g(h(\phi W + \phi hW), Z) - \mu\eta(W)g(\phi hY, Z) \left. \left. \right\} + \gamma \left\{ g(W + hW, \phi Y)\eta(Z) \right. \right. \\
&+ \eta(Y)g(W + hW, \phi Z) \left. \left. \right\} \right] - (a_3 + a_6) \left[\beta \left\{ [(1-k)g(W, \phi Z) + g(W, h\phi Z)]\eta(Y) \right. \right. \\
&+ \eta(Z)g(h(\phi W + \phi hW), Y) - \mu\eta(W)g(\phi hZ, Y) \left. \left. \right\} + \gamma \left\{ g(W + hW, \phi Z)\eta(Y) \right. \right. \\
&+ \eta(Z)g(W + hW, \phi Y) \left. \left. \right\} \right] - a_7(\nabla_W r)[2ng(Y, Z)] + a_0\eta((\nabla_W R)(\xi, Y)Z) \\
&+ a_2[\beta g(h(\phi W + \phi hW), Z)\eta(Y) + \gamma g(W + hW, \phi Z)\eta(Y)] \\
&+ a_3[\beta g(h(\phi W + \phi hW), Y)\eta(Z) + \gamma g(W + hW, \phi Y)\eta(Z)] \\
&+ a_5[\beta \{(1-k)g(W, \phi Y) + g(W, h\phi Y)\}\eta(Z) + \gamma\eta(Z)g(W + hW, \phi Y)] \\
&+ a_6[\beta \{(1-k)g(W, \phi Z) + g(W, h\phi Z)\}\eta(Y) + \gamma\eta(Y)g(W + hW, \phi Z)] \\
&+ a_7(\nabla_W r)[g(Y, Z) - \eta(Y)\eta(Z)] \\
&= A(W)[a_4 + 2na_7]rg(Y, Z) + A(W)[a_0 + (2n + 1)a_1 + a_2 + a_3 + a_5 + a_6]S(Y, Z).
\end{aligned} \tag{19}$$

Putting $Z = \xi$ in (19) and on simplification, we get

$$\begin{aligned} & -a_0(\nabla_W S)(Y, \xi) - (2na_1 + a_2 + a_6)[\beta\{(1-k)g(W, \phi Y) + g(W, h\phi Y)\} \\ & + \gamma g(W + hW, \phi Y)] - 2na_7(\nabla_W r)\eta(Y) \\ & = A(W)\eta(Y)[[a_0 + (2n+1)a_1 + a_2 + a_3 + a_5 + a_6]2nk + [a_4 + 2na_7]r]. \end{aligned} \quad (20)$$

We know that

$$(\nabla_W S)(Y, \xi) = \nabla_W S(Y, \xi) - S(\nabla_W Y, \xi) - S(Y, \nabla_W \xi). \quad (21)$$

By using (4), (8) in (21), we obtain

$$(\nabla_W S)(Y, \xi) = S(Y, \phi W) + S(Y, \phi hW) - 2nkg(Y, \phi W) - 2nkg(Y, \phi hW). \quad (22)$$

Substituting (22) in (20), we get

$$\begin{aligned} & -a_0\{S(Y, \phi W) + S(Y, \phi hW) - 2nkg(Y, \phi W) - 2nkg(Y, \phi hW)\} \\ & - (2na_1 + a_2 + a_6)[\beta\{(1-k)g(W, \phi Y) + g(W, h\phi Y)\} + \gamma g(W + hW, \phi Y)] \\ & - 2na_7(\nabla_W r)\eta(Y) \\ & = A(W)\eta(Y)[[a_0 + (2n+1)a_1 + a_2 + a_3 + a_5 + a_6]2nk + [a_4 + 2na_7]r]. \end{aligned} \quad (23)$$

Replacing Y by ϕY in (23) and simplifying, we have

$$\begin{aligned} & -a_0S(Y, W) - a_0S(Y, hW) + [2a_0\beta + 2nka_0 + (2na_1 + a_2 + a_6)(\beta + \gamma)]g(hW, Y) \\ & + [2nka_0 + (2na_1 + a_2)\{\beta(1-k) + \gamma\} + a_6\gamma - (2a_0 + a_6)\beta(k-1)]g(W, Y) \\ & + [(2a_0 + a_6)\beta(k-1) - (2na_1 + a_2)\{\beta(1-k) + \gamma\} - a_6\gamma]\eta(W)\eta(Y) = 0. \end{aligned} \quad (24)$$

Replacing W by hW in (24) and by virtue of (7), (9) and on simplification, we get

$$\begin{aligned} & -a_0S(Y, hW) + a_0(k-1)S(Y, W) \\ & + (k-1)[2a_0\beta + (2na_1 + a_2 + a_6)(\beta + \gamma)]\eta(W)\eta(Y) \\ & + [2nka_0 + (2na_1 + a_2)\{\beta(1-k) + \gamma\} + a_6\gamma - (2a_0 + a_6)\beta(k-1)]g(hW, Y) \\ & - (k-1)[2a_0\beta + 2nka_0 + (2na_1 + a_2 + a_6)(\beta + \gamma)]g(W, Y) = 0. \end{aligned} \quad (25)$$

Subtracting (24) and (25) and by virtue of (9), we obtain

$$g(Y, hW) = \frac{E}{F}g(Y, W) + \frac{G}{F}\eta(Y)\eta(W), \quad (26)$$

where

$$\begin{aligned} E &= [a_0(\beta - 2nk) + (2na_1 + a_2 + a_6)\gamma] \\ F &= [a_0(2\beta - \alpha) + \beta(2na_1 + a_2)] \end{aligned}$$

and

$$G = \gamma(-a_0 + 2na_1 + a_2 + a_6).$$

By substituting (26) in (24), we get

$$S(Y, W) = \left[\frac{NE}{a_0 F} + \frac{P}{a_0} \right] g(Y, W) + \left[\frac{GN}{F a_0} + \frac{Q}{a_0} \right] \eta(Y)\eta(W),$$

where

$$N = [2nk - \alpha + 2\beta]a_0 + (2na_1 + a_2 + a_6)(\beta + \gamma),$$

$$P = [2nka_0 - \beta(k - 1)[a_0 + 2na_1 + a_2 + a_6] + \gamma[2na_1 + a_2 + a_6]]$$

and

$$Q = [\beta(k - 1)[a_0 + 2na_1 + a_2 + a_6] - \gamma[2na_1 + a_2 + a_6]].$$

Hence, we state the following:

Theorem 1. *A ϕ - τ -recurrent (k, μ) -contact metric manifold is an η -Einstein manifold with $a_0 \neq 0$.*

4 η - τ -Ricci-recurrent (k, μ) -contact metric manifold

Definition 4. A (k, μ) -contact metric manifold M is said to be η - τ -Ricci-recurrent if it satisfies the condition

$$(\nabla_X S_\tau)(\phi Y, \phi Z) = A(X)S_\tau(\phi Y, \phi Z), \quad (27)$$

for all vector fields X, Y, Z on M

From (12), we have

$$S_\tau(Y, Z) = [a_0 + (2n + 1)a_1 + a_2 + a_3 + a_5 + a_6]S(Y, Z) + r[a_4 + 2na_7]g(Y, Z). \quad (28)$$

Replacing $Y = \phi Y$ and $Z = \phi Z$ in (28), we obtain

$$S_\tau(\phi Y, \phi Z) = [a_0 + (2n + 1)a_1 + a_2 + a_3 + a_5 + a_6]S(\phi Y, \phi Z) + [a_4 + 2na_7]rg(\phi Y, \phi Z). \quad (29)$$

Differentiating (29) with respect to X , we get

$$(\nabla_X S_\tau)(\phi Y, \phi Z) = [a_0 + (2n + 1)a_1 + a_2 + a_3 + a_5 + a_6](\nabla_X S)(\phi Y, \phi Z) + [a_4 + 2na_7](\nabla_X r)g(\phi Y, \phi Z). \quad (30)$$

By using (30) and (29) in (27), we have

$$L(\nabla_X S)(\phi Y, \phi Z) + M(\nabla_X r)g(\phi Y, \phi Z) = A(X)\{LS(\phi Y, \phi Z) + Mrg(\phi Y, \phi Z)\}, \quad (31)$$

where $L = [a_0 + (2n + 1)a_1 + a_2 + a_3 + a_5 + a_6]$ and $M = [a_4 + 2na_7]$.

Now, differentiating (11), we have

$$\begin{aligned} (\nabla_X S)(\phi Y, \phi Z) &= (\nabla_X S)(Y, Z) - 2nk[-\eta(Z)g(Y, \phi X) - \eta(Z)g(Y, \phi hX) \\ &\quad - \eta(Y)g(Z, \phi X) - \eta(Y)g(Z, \phi hX)] \\ &\quad - 2[2n - 2 + \mu][(1 - k)g(X, \phi Y)\eta(Z) \\ &\quad + g(X, h\phi Y)\eta(Z) + \eta(Y)g(h(\phi X + \phi hX), Z) \\ &\quad - \mu\eta(X)g(\phi hY, Z)] \\ &\quad + \eta(Y)S(X + hX, \phi Z) + \eta(Z)S(\phi Y, X + hX). \end{aligned} \quad (32)$$

Substituting (32) in (31) and on simplification, we obtain

$$\begin{aligned}
(\nabla_X S)(Y, Z) &= -2nk[\eta(Z)g(Y, \phi X) + \eta(Z)g(Y, \phi hX) + \eta(Y)g(Z, \phi X) \\
&\quad + \eta(Y)g(Z, \phi hX)] + 2[2n - 2 + \mu][(1 - k)g(X, \phi Y)\eta(Z) \\
&\quad + g(X, h\phi Y)\eta(Z) + \eta(Y)g(h(\phi X + \phi hX), Z) - \mu\eta(X)g(\phi hY, Z)] \\
&\quad - \eta(Y)S(X + hX, \phi Z) - \eta(Z)S(\phi Y, X + hX) \\
&\quad - \frac{M}{L}(\nabla_X r)g(\phi Y, \phi Z) + A(X)\{S(Y, Z) - 2nk\eta(Y)\eta(Z) \\
&\quad - 2[2n - 2 + \mu]g(hY, Z) + \frac{Mr}{L}g(\phi Y, \phi Z)\}.
\end{aligned} \tag{33}$$

Let $\{e_i : i = 1, 2, \dots, 2n + 1\}$ be an orthonormal frame field at any point of the manifold. Then contracting Y and Z in (33), we have

$$dr(X) = A(X) \left[r - \frac{2nkL}{(L + 2nM)} \right]. \tag{34}$$

Again, contracting over X and Z in (33), we get

$$\begin{aligned}
\frac{1}{2}dr(Y) &= \left[-2nk \operatorname{tr}(\phi) + 2(2n - 2 + \mu) \operatorname{tr}(h\phi h) - \operatorname{tr}(Q\phi) - \operatorname{tr}(Qh\phi) + \frac{(\xi r)M}{L} \right] \eta(Y) \\
&\quad - \left[2nk + \frac{rM}{L} \right] A(\xi)\eta(Y) - \frac{M}{L}dr(Y) + S(Y, \rho) - 2(2n - 2 + \mu)A(hY) \\
&\quad + \frac{rM}{L}A(Y).
\end{aligned} \tag{35}$$

By using (34) in (35) and on simplification, we get

$$\begin{aligned}
A(Y) \left[\frac{r}{2} - \frac{nk(L + 2M)}{(L + 2nM)} \right] &= \left[-2nk \operatorname{tr}(\phi) + 2(2n - 2 + \mu) \operatorname{tr}(h\phi h) \right. \\
&\quad \left. - \operatorname{tr}(Q\phi) - \operatorname{tr}(Qh\phi) - \frac{2nk(L + M + 2nM)}{(L + 2nM)} A(\xi) \right] \eta(Y) \\
&\quad + S(Y, \rho) - 2(2n - 2 + \mu)A(hY). \tag{36}
\end{aligned}$$

Replacing $Y = hY$ in (36) and by virtue of (9), we obtain

$$A(hY) = \frac{2(L + 2nM)}{[(r - 2\alpha)(L + 2nM) - 2nk(L + 2M)]} [\beta(k - 1)\{A(Y) - A(\xi)\eta(Y)\}], \tag{37}$$

where $\alpha = [2(n - 1) - n\mu]$ and $\beta = [2(n - 1) + \mu]$.

Substituting (37) in (36), we have

$$\begin{aligned} A(Y) & \left[\frac{r}{2} - \frac{nk(L+2M)}{(L+2nM)} + \frac{4\beta^2(L+2nM)(k-1)}{[(r-2\alpha)(L+2nM) - 2nk(L+2M)]} \right] \\ & = [-2nk \operatorname{tr}(\phi) + 2\beta \operatorname{tr}(h\phi h) - \operatorname{tr}(Q\phi) - \operatorname{tr}(Qh\phi)]\eta(Y) \\ & + \left[\frac{4\beta^2(L+2nM)(k-1)}{[(r-2\alpha)(L+2nM) - 2nk(L+2M)]} - \frac{2nk(L+M+2nM)}{(L+2nM)} \right] A(\xi)\eta(Y) \\ & \quad + S(Y, \rho). \end{aligned} \quad (38)$$

Putting $Y = \xi$ in (38), we get

$$A(\xi) \left[\frac{r}{2} - \frac{nkL}{(L+2nM)} \right] = [-2nk \operatorname{tr}(\phi) + 2\beta \operatorname{tr}(h\phi h) - \operatorname{tr}(Q\phi) - \operatorname{tr}(Qh\phi)]. \quad (39)$$

From (39) and (38), we have

$$\begin{aligned} S(Y, \rho) & = \left[\frac{r}{2} - \frac{nk(L+2M)}{(L+2nM)} + \frac{4\beta^2(L+2nM)(k-1)}{[(r-2\alpha)(L+2nM) - 2nk(L+2M)]} \right] g(Y, \rho) \\ & + \left[-\frac{r}{2} + \frac{nkL}{(L+2nM)} - \frac{4\beta^2(L+2nM)(k-1)}{[(r-2\alpha)(L+2nM) - 2nk(L+2M)]} \right. \\ & \quad \left. + \frac{2nk(L+M+2nM)}{(L+2nM)} \right] \eta(Y)\eta(\rho). \end{aligned} \quad (40)$$

From (40), we have

$$\begin{aligned} QY & = \left[\frac{r}{2} - \frac{nk(L+2M)}{(L+2nM)} + \frac{4\beta^2(L+2nM)(k-1)}{[(r-2\alpha)(L+2nM) - 2nk(L+2M)]} \right] Y \\ & + \left[-\frac{r}{2} + \frac{nkL}{(L+2nM)} - \frac{4\beta^2(L+2nM)(k-1)}{[(r-2\alpha)(L+2nM) - 2nk(L+2M)]} \right. \\ & \quad \left. + \frac{2nk(L+M+2nM)}{(L+2nM)} \right] \eta(Y)\xi. \end{aligned} \quad (41)$$

Hence, we state the following:

Theorem 2. *If the Ricci tensor of a (k, μ) -contact metric manifold is η - τ -Ricci-recurrent then its Ricci tensor along the associated vector field of the 1-form is given by (40) and the eigen value of the Ricci tensor with respect to the characteristic vector ξ is given by (41).*

Acknowledgement

The authors are very grateful to the referees for their valuable suggestions and opinions.

References

- [1] D.E. Blair: *Contact metric manifolds in Riemannian geometry*. Springer-Verlag, Berlin-New-York (1976). Lecture Notes in Mathematics 509
- [2] D.E. Blair, T. Koufogiorgos, B.J. Papantoniou: Contact metric manifolds satisfying a nullity condition. *Israel J. Math.* 91 (1995) 189–214.
- [3] E. Boeckx, P. Buecken, L. Vanhecke: ϕ -symmetric contact metric spaces. *Glasgow Math. J.* 41 (1999) 409–416.
- [4] U.C. De, A.K. Gazi: On ϕ -recurrent $N(k)$ -contact metric manifolds. *Math. J. Okayama Univ.* 50 (2008) 101–112.
- [5] U.C. De, A.A. Shaikh, S. Biswas: On ϕ -recurrent Sasakian manifolds. *Novi Sad J. Math.* 33 (2003) 13–48.
- [6] H.G. Nagaraja, G. Somashekhara: τ -curvature tensor in (k, μ) -contact metric manifolds. *Mathematica Aeterna* 2 (6) (2012) 523–532.
- [7] B.J. Papantoniou: Contact Riemannian manifolds satisfying $R(\xi, X) \cdot R = 0$ and $\xi \in (k, \mu)$ -nullity distribution. *Yokohama Math. J.* 40 (2) (1993) 149–161.
- [8] C.R. Premalatha, H.G. Nagaraja: On Generalized (k, μ) -space forms. *Journal of Tensor Society* 7 (2013) 29–38.
- [9] A.A. Shaikh, K.K. Baishya: On (k, μ) -contact metric manifolds. *Differential Geometry – Dynamical Systems* 8 (2006) 253–261.
- [10] R. Sharma, D.E. Blair: Conformal motion of contact manifolds with characteristic vector field in the k -nullity distribution. *Illinois J. Math.* 42 (1998) 673–677.
- [11] S. Tanno: Ricci curvatures of contact Riemannian manifolds. *Tohoku Math. J.* 40 (1988) 441–448.
- [12] T. Takahashi: Sasakian ϕ -symmetric spaces. *Tohoku Math. J.* 29 (1977) 91–113.
- [13] M.M. Tripathi, P. Gupta: τ -curvature tensor on a semi-Riemannian manifold. *J. Adv. Math. Stud.* 4 (1) (2011) 117–129.
- [14] M.M. Tripathi, P. Gupta: On τ -curvature tensor in K-contact and Sasakian manifolds. *International Electronic Journal of Geometry* 4 (2011) 32–47.
- [15] M.M. Tripathi, P. Gupta: $(N(k), \xi)$ -semi-Riemannian manifolds: Semisymmetries. *arXiv:1202.6138v[math.DG]* 28 (2012).

Received: 24 April, 2017

Accepted for publication: 15 May, 2018

Communicated by: Haizhong Li