



**HAL**  
open science

# Hilbert series of the Grassmannian and k-Narayana numbers

Lukas Braun

► **To cite this version:**

Lukas Braun. Hilbert series of the Grassmannian and k-Narayana numbers. Communications in Mathematics, 2019, Volume 27 (2019), Issue 1 (1), pp.27 - 41. 10.2478/cm-2019-0003 . hal-03664962

**HAL Id: hal-03664962**

**<https://hal.science/hal-03664962v1>**

Submitted on 11 May 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

## Hilbert series of the Grassmannian and $k$ -Narayana numbers

Lukas Braun

**Abstract.** We compute the Hilbert series of the complex Grassmannian using invariant theoretic methods. This is made possible by showing that the denominator of the  $q$ -Hilbert series is a Vandermonde-like determinant. We show that the  $h$ -polynomial of the Grassmannian coincides with the  $k$ -Narayana polynomial. A simplified formula for the  $h$ -polynomial of Schubert varieties is given. Finally, we use a generalized hypergeometric Euler transform to find simplified formulae for the  $k$ -Narayana numbers, i.e. the  $h$ -polynomial of the Grassmannian.

### Introduction

Consider the Grassmannian  $X = \text{Gr}(k, n)$  of  $k$ -dimensional vector subspaces of a given  $n$ -dimensional complex vector space and its homogeneous coordinate ring  $R = \bigoplus_j R_j$  defined by the Plücker embedding. Recall that the associated Hilbert series is

$$H(X) = \sum_{j \geq 0} \dim(R_j) t^j.$$

The Hilbert function  $j \mapsto \dim(R_j)$  is, up to finitely many values, a polynomial in  $j$ , the Hilbert polynomial of  $\text{Gr}(k, n)$ . Moreover, the Hilbert series is represented as a rational function in  $j$  with a denominator polynomial of degree  $k(k(n-k)+1)$ . The numerator is then called the  $h$ -polynomial of  $\text{Gr}(k, n)$ . Various approaches leading to explicit formulae for the Hilbert polynomial of  $\text{Gr}(k, n)$  can be found for example in [2], [5], [6], [7], [8], [10], [15], [18], [20].

Mukai used an invariant theoretic approach in [15] to compute the Hilbert polynomial of the Grassmannian in the special case  $k = 2$ . The aim of the present

---

2010 MSC: 14M15, 13D40, 33C90

Key words: Hilbert series of the Grassmannian, Narayana numbers, Euler's hypergeometric transform

Affiliation:

Mathematisches Institut, Universität Tübingen, Auf der Morgenstelle 10, 72076

Tübingen, Germany

E-mail: braun@math.uni-tuebingen.de

paper is to generalize this invariant theoretic method to arbitrary  $k$ . In fact, Mukai's Hilbert polynomial for  $\text{Gr}(2, n)$  is the special case  $k = 2$ ,  $r = n - 1$  of the order polynomial

$$\mathfrak{N}_k(r, j) := \prod_{i=0}^{k-1} \binom{r+i+j-1}{i+j} \binom{r+i-1}{i}^{-1},$$

which is a polynomial in  $j$ , appearing in Sulanke's papers [21], [22]. In these papers, he generalized the (two-dimensional) *Narayana numbers* to the  $k$ -*Narayana numbers*

$$N_k(r, j) := \sum_{l=0}^j (-1)^{j-l} \binom{kr+1}{j-l} \mathfrak{N}_k(r+1, l)$$

for arbitrary dimension  $k$ . We call the order polynomial  $\mathfrak{N}_k(r, j)$  the *multiset  $k$ -Narayana numbers* in the following. For combinatorial interpretations of these numbers, we refer to Section 1. The generating series

$$N_{k,r} := \sum_{j=0}^{\binom{k-1}{r-1}} N_k(r, j) t^j, \quad \mathfrak{N}_{k,r} := \sum_{j \geq 0} \mathfrak{N}_k(r, j) t^j,$$

of which the first is a polynomial in  $t$ , are called the  $k$ -*Narayana polynomial* and the  $k$ -*Narayana series* respectively. Computing the Hilbert series of the Grassmannian using the invariant theoretic method involves a Vandermonde-like determinant as a crucial ingredient, and we arrive at our first main theorem:

**Theorem 1.** *The  $h$ -polynomial of the Grassmannian  $\text{Gr}(k, n)$  is the  $k$ -Narayana polynomial  $N_{k, n-k}$  and its Hilbert series is the  $k$ -Narayana series  $\mathfrak{N}_{k, n-k+1}$ .*

Finally, we express the  $k$ -Narayana series as a hypergeometric function  ${}_kF_{k-1}$ . This leads to the observation that the simplified formula  $\frac{1}{r} \binom{r}{j} \binom{r}{j+1}$  for the 2-Narayana numbers is a direct consequence of Euler's hypergeometric transformation

$${}_2F_1 \left( \begin{matrix} a, b \\ c \end{matrix}; t \right) = (1-t)^{c-a-b} {}_2F_1 \left( \begin{matrix} c-a, c-b \\ c \end{matrix}; t \right).$$

This transformation has been generalized in [14] – see also [12], [13] – to the generalized hypergeometric function  ${}_kF_{k-1}$ . Applying this generalized Euler transformation to the  $k$ -Narayana series, we express the  $k$ -Narayana polynomial as a hypergeometric function, i.e. we find a new (multiplicative) formula for the  $k$ -Narayana numbers:

**Theorem 3.** *For the  $k$ -Narayana numbers  $N_k(r, j)$ , we have the product formula*

$$N_k(r, j) = \frac{1}{j+1} \binom{(k-1)(r-1)}{j} \binom{(k-1)(r-1)+1}{j} \prod_{i=1}^{\binom{k-2}{r-2}} \frac{\eta_i + j}{\eta_i},$$

where the  $\eta_i$  are the zeros of a certain polynomial of degree  $(k-2)(r-2)$ .

The paper is divided in three sections. In the first, we study different generalizations of Narayana numbers, relations among them and express their generating functions as hypergeometric functions. The second section is devoted to the computation of Hilbert series of the Grassmannian using invariant theoretic methods. In addition, a simplified formula – compared to that of [16] – for the  $h$ -polynomial of Schubert varieties is derived. The third and last section uses the generalized hypergeometric Euler transformation of [14] to find a new multiplicative formula for the  $k$ -Narayana numbers.

The author wishes to thank Jürgen Hausen as well as the anonymous referee for many helpful comments.

## 1 The $k$ -Narayana numbers

The *Narayana numbers*

$$N(r, j) := \frac{1}{r} \binom{r}{j} \binom{r}{j+1} = \frac{1}{j+1} \binom{r-1}{j} \binom{r}{j}$$

count the number of Dyck paths in the plane from  $(0, 0)$  to  $(r, r)$  with exactly  $j$  ascents. They have been generalised in numerous ways. We exhibit some of these generalizations. The  $k$ -Narayana numbers  $N_k(r, j)$  count the number of paths along the lattice  $\mathbf{Z}^k$  from the origin to  $(r, \dots, r)$ , staying in the region  $\{0 \leq x_1 \leq \dots \leq x_k\}$  and having  $j$  ascents. An ascent here is a pair of successive steps so that the second step is an increase in a coordinate with higher index than the first one. They have been observed by Sulanke in the papers [22] and [21] and are given by the formula

$$N_k(r, j) := \sum_{l=0}^j (-1)^{j-l} \binom{kr+1}{j-l} \prod_{i=0}^{k-1} \binom{r+i+l}{r} \binom{r+i}{r}^{-1}.$$

In his initial paper [17], Narayana introduced the Narayana numbers in another context than that of the Dyck paths that coincides for  $k = 2$ . Consider the following setting: paths with  $j$  steps from the origin to a point  $(a_1, \dots, a_k)$  in  $\mathbf{Z}^k$  with  $a_1 \geq \dots \geq a_k \geq j$ . The steps comply with the following rules:

1. in each step, each coordinate increases at least by one.
2. if  $a_{il}$  is the  $i$ -th coordinate after the  $l$ -th step, then  $a_{1l} \geq \dots \geq a_{kl} \geq l$  holds.

The number  $(a_1, \dots, a_k)_j$  of such paths according to Theorem 1 of [17] is given by

$$(a_1, \dots, a_k)_j := \left| \begin{array}{ccc} \binom{a_1-1}{j} & \cdots & \binom{a_k-1}{j+k-1} \\ \vdots & \ddots & \vdots \\ \binom{a_1-1}{j-k+1} & \cdots & \binom{a_k-1}{j} \end{array} \right| = \det \left( \binom{a_l-1}{j+l-i} \right)_{1 \leq i, l \leq k}.$$

We see that by setting  $k = 2$ ,  $a_1 = a_2 = r$  one gets the ordinary Narayana numbers, while for  $k = 1$ , one gets the binomial coefficients. We now consider two small modifications, namely the numbers

$$\begin{aligned}
[a_1, \dots, a_k]_j &:= \begin{vmatrix} \binom{a_1+j-1}{j} & \dots & \binom{a_k+j-1}{j+k-1} \\ \vdots & \ddots & \vdots \\ \binom{a_1+j-1}{j-k+1} & \dots & \binom{a_k+j-1}{j} \end{vmatrix} = \det \left( \binom{a_l + j - 1}{j + l - i} \right)_{1 \leq i, l \leq k}, \\
\{a_1, \dots, a_k\}_j &:= \begin{vmatrix} \binom{a_k+j-1}{j} & \dots & \binom{a_1+j+k-2}{j+k-1} \\ \vdots & \ddots & \vdots \\ \binom{a_k+j-k}{j-k+1} & \dots & \binom{a_1+j-1}{j} \end{vmatrix} = \det \left( \binom{a_{k-l+1} + j + l - i - 1}{j + l - i} \right)_{1 \leq i, l \leq k} \\
&= \begin{vmatrix} \binom{a_1+j-1}{j} & \dots & \binom{a_1+j-1}{j+k-1} \\ \vdots & \ddots & \vdots \\ \binom{a_k+j-k}{j-k+1} & \dots & \binom{a_k+j-k}{j} \end{vmatrix} = \det \left( \binom{a_i + j - i}{j + l - i} \right)_{1 \leq i, l \leq k}.
\end{aligned}$$

The numbers  $[a_1, \dots, a_k]_j$  turn up in enumerative combinatorics, see for example [1], while the numbers  $\{a_1, \dots, a_k\}_j$  give the Hilbert polynomial of Schubert varieties due to the formula of Hodge and Pedoe, see Theorem III on page 387 of [8] and also [4], [16]. The equality of the two different formulae for  $\{a_1, \dots, a_k\}_j$  is given by Lemma 7 of [4]. In all cases, setting  $k = 1$  gives the multiset coefficients

$$\binom{\binom{a}{b}}{\binom{a}{b}} = \binom{a + b - 1}{b}.$$

If all  $a_i$  are equal, which has been considered for the numbers  $[a_1, \dots, a_k]_j$  in [1], [11], [19], then we have the following identity:

**Lemma 1.** *Let  $r, k - 1 \in \mathbf{Z}_{\geq 0}$ . It holds*

$$\underbrace{\{r + k - 1, \dots, r + k - 1\}}_k = \underbrace{\{r, \dots, r\}}_k = \underbrace{\{j, \dots, j\}}_k r.$$

*Proof.* The second equality follows directly by transposing the matrix  $M_k(r) := \binom{r+j-1}{j+l-i}_{1 \leq i, l \leq k}$  and applying the binomial identity  $\binom{a}{b} = \binom{a}{a-b}$ . We prove the first one. We see that the last rows of  $M_k(r)$  and of  $M'_k(r+k-1) := \binom{r+k+j-i-1}{j+l-i}_{1 \leq i, l \leq k}$  are the same. Moreover, the lower arguments of the binomial coefficients in the  $(i, l)$ -th entry of  $M_k(r)$  and of  $M'_k(r+k-1)$  are the same. The upper arguments in  $M'_k(r+k-1)$  decrease by one if  $i$  increases by one. Now recall the binomial identity

$$\binom{a}{b} - \binom{a-1}{b-1} = \binom{a-1}{b}.$$

By the elementary row operations of subtracting the second from the first, the third from the second and so on till we subtract the  $k$ -th from the  $(k-1)$ -th row and applying the above binomial identity, we decrease the upper arguments of the binomial coefficients in the first  $k-1$  rows of  $M'_k(r+k-1)$  each by one without changing the determinant. In particular, now the last two rows of  $M_k(r)$  and our new  $M'_k(r+k-1)$  are the same. We do the same again, now only with the first  $k-1$  rows and achieve that the last three rows of the two matrices coincide. After

doing this  $k-1$  times, we have transferred  $M'_k(r+k-1)$  to  $M_k(r)$  without changing the determinant and the assertion is proven.  $\square$

**Definition 1.** Let  $k, r, j \in \mathbf{Z}_{\geq 1}$ . Then we call the numbers

$$\mathcal{N}_k(r, j) := \underbrace{(r, \dots, r)}_k, \quad \mathfrak{N}_k(r, j) := \underbrace{[r, \dots, r]}_k = \underbrace{[j, \dots, j]}_k$$

the *simple  $k$ -Narayana numbers* and the *multiset  $k$ -Narayana numbers* respectively and furthermore

$$\mathcal{N}_{k,r}(t) := \sum_{j=0}^r \mathcal{N}_k(r, j)t^j, \quad \mathfrak{N}_{k,r}(t) := \sum_{j=0}^{\infty} \mathfrak{N}_k(r, j)t^j, \quad N_{k,r}(t) := \sum_{j=0}^{(r-1)(k-1)} N_k(r, j)t^j$$

the *simple  $k$ -Narayana polynomial*, the  *$k$ -Narayana series* and the  *$k$ -Narayana polynomial* respectively.

Apart from [1], they turn up in [11] and [19]. In [1] as well as in [19], closed formulae for  $\mathfrak{N}_k(r, j)$  are given, so that we get the following:

**Proposition 1.** Let  $k, r, j \in \mathbf{Z}_{\geq 0}$  or  $k \geq 2, r > j \geq 0$  respectively. We have the identities

$$\begin{aligned} \mathfrak{N}_k(r, j) &= \prod_{i=1}^k \binom{j+i}{r-1} \binom{i}{r-1}^{-1} = \prod_{i=1}^k \binom{r-1+i}{j} \binom{i}{j}^{-1}, \\ \mathcal{N}_k(r, j) &= \prod_{i=1}^k \binom{r-1+i}{j} \binom{i}{j}^{-1} = \prod_{i=0}^{k-1} \binom{r+i}{j} \binom{j+i}{j}^{-1}. \end{aligned}$$

*Proof.* The first identity is Theorem 3.3 in [1]. The second follows by interchanging  $j+1$  and  $r$ . The third by setting  $r' := r+j-1$  in the second one and the last is a simple index shift.  $\square$

**Corollary 1.** We have the following relations between  $k$ -Narayana numbers and polynomials:

$$\begin{aligned} \mathfrak{N}_k(r, j) &= \mathcal{N}_k(r+j-1, j) \\ &= \sum_{l \geq 0} \binom{k(r-1)+j-l}{k(r-1)} N_k(r-1, l), \\ \mathcal{N}_k(r, j) &= \sum_{l=0}^j (-1)^{j-l} \binom{kr+1}{j-l} \mathcal{N}_k(r+l, l) \\ &= \sum_{l=0}^j (-1)^{j-l} \binom{kr+1}{j-l} \mathfrak{N}_k(r+1, l) \\ \mathfrak{N}_{k,r}(t) &= \frac{N_{k,r-1}(t)}{(1-t)^{k(r-1)+1}} \end{aligned}$$

*Proof.* The second equality is Proposition 4 of [21], the last one is stated on page 4 of [21], the others follow directly from the definitions and Proposition 1.  $\square$

Now we express the  $k$ -Narayana polynomials and series in terms of hypergeometric functions in order to find simplified formulae using a generalized Euler transform.

**Proposition 2.** *Let  $\mathcal{N}_{k,r}(t)$  be the simple  $k$ -Narayana polynomial and  $\mathfrak{N}_{k,r}(t)$  the  $k$ -Narayana series. Let further  ${}_pF_q \left( \begin{smallmatrix} a_1, \dots, a_p \\ b_1, \dots, b_q \end{smallmatrix}; t \right)$  be the generalized hypergeometric function. Then we have*

$$\begin{aligned}\mathcal{N}_{k,r}(t) &= {}_kF_{k-1} \left( \begin{matrix} -r, \dots, -r-k+1 \\ 2, \dots, k \end{matrix}; (-1)^k t \right), \\ \mathfrak{N}_{k,r}(t) &= {}_kF_{k-1} \left( \begin{matrix} r, \dots, r+k-1 \\ 2, \dots, k \end{matrix}; t \right).\end{aligned}$$

*Proof.* We have

$$\begin{aligned}\mathfrak{N}_{k,r}(t) &= \sum_{j \geq 0} \prod_{i=1}^k \binom{j+i}{r-1} \binom{i}{r-1}^{-1} t^j \\ &= \sum_{j \geq 0} \frac{(j+r-1)! \cdots (j+r+k-2)!}{(r-1)!^k j! \cdots (j+k-1)!} \frac{(r-1)!^k 1! \cdots k!}{(r-1)! \cdots (r+k-2)!} t^j \\ &= \sum_{j \geq 0} \frac{(r)_j \cdots (r+k-1)_j}{(2)_j \cdots (k)_j} \frac{t^j}{j!} = {}_kF_{k-1} \left( \begin{matrix} r, \dots, r+k-1 \\ 2, \dots, k \end{matrix}; t \right), \\ \mathcal{N}_{k,r}(t) &= \sum_{j=0}^r \prod_{i=0}^{k-1} \binom{r+i}{j} \binom{j+i}{j}^{-1} t^j \\ &= \sum_{j=0}^r \frac{r! \cdots (r+k-1)!}{j!^k (r-j)! \cdots (r+k-1-j)!} \frac{j!^k 1! \cdots (k-1)!}{j! \cdots (j+k-1)!} t^j \\ &= \sum_{j=0}^r \frac{(-r)_j \cdots (-r-k+1)_j}{(2)_j \cdots (k)_j} \frac{(-1)^{kj} t^j}{j!} \\ &= {}_kF_{k-1} \left( \begin{matrix} -r, \dots, -r-k+1 \\ 2, \dots, k \end{matrix}; (-1)^k t \right).\end{aligned}$$

$\square$

## 2 Hilbert series of the Grassmannian

Denote by  $\text{Gr}(k, n)$  the complex Grassmannian of  $k$ -dimensional vector subspaces in  $n$ -dimensional vector space. The Hilbert series of its homogeneous coordinate ring under the Plücker embedding has been investigated for example in [2], [5], [6], [7], [8], [15], [18], [20]. Mukai gives a closed formula for the Hilbert polynomial in terms of binomial coefficients in [15] while [5] gives a closed formula for the generating rational function, both in the case  $\text{Gr}(2, n)$ . In [6], Hodge conjectured a

closed formula for the Hilbert polynomial in the general case  $\text{Gr}(k, n)$ , which was proved by Littlewood in [10]. This formula is the following:

$$d_{k,n}(j) = \frac{(n+j)! \cdots (n+j-k)!}{j! \cdots (k+j)!} \frac{1! \cdots k!}{(n-k)! \cdots n!}.$$

The homogeneous coordinate ring of  $\text{Gr}(k, n)$  equals  $\mathbf{C}[V^n]^{\text{SL}_k}$ , where the action of  $\text{SL}_k$  on  $V^n$  is induced by multiplication from the left on the  $k$ -dimensional vector space  $V$ , see for example [9], Theorem 9.3.6. So the Hilbert series of  $\text{Gr}(k, n)$  and  $\mathbf{C}[V^n]^{\text{SL}_k}$  coincide.

The following theorem gives a formula for the Hilbert series in terms of binomial coefficients and an expression of the Hilbert series as a rational function.

**Theorem 1.** *The  $h$ -polynomial of the complex Grassmannian  $\text{Gr}(k, n)$  is the  $k$ -Narayana polynomial  $N_{k,n-k}$  and its Hilbert series is the  $k$ -Narayana series  $\mathfrak{N}_{k,n-k+1}$ . That is to say:*

$$H(\text{Gr}(k, n)) = \mathfrak{N}_{k,n-k+1}(t^k) = \frac{N_{k,n-k}(t)}{(1-t^k)^{k(n-k)+1}}.$$

**Lemma 2.** *For  $k \geq 2$ , the following equality holds in  $\mathbf{C}(z_1, \dots, z_k)$ :*

$$\prod_{1 \leq i < j \leq k} \left(1 - \frac{z_i}{z_j}\right) = \left| \left( z_i^{j-i} \right)_{1 \leq i, j \leq k} \right|$$

*Proof.* The Vandermonde matrix

$$\left( z_i^{j-1} \right)_{1 \leq i, j \leq k}$$

has the determinant

$$\left| \left( z_i^{j-1} \right)_{1 \leq i, j \leq k} \right| = \prod_{1 \leq i < j \leq k} (z_j - z_i).$$

So we have

$$\begin{aligned} \left| \left( z_i^{j-i} \right)_{1 \leq i, j \leq k} \right| &= \left( \prod_{i=1}^k z_i^{1-i} \right) \left| \left( z_i^{j-1} \right)_{1 \leq i, j \leq k} \right| \\ &= \left( \prod_{i=1}^k z_i^{1-i} \right) \left( \prod_{1 \leq i < j \leq k} (z_j - z_i) \right) \\ &= \prod_{1 \leq i < j \leq k} \left( 1 - \frac{z_i}{z_j} \right). \quad \square \end{aligned}$$

*Proof of Theorem 1.* Let  $n \geq k$ . We begin with what Mukai [15] calls the  $q$ -Hilbert series of the action of  $SL_k$  on  $V^n$  induced by multiplication from the left on the  $k$ -dimensional vector space  $V$ .



Let  $T = SL_k \cap \text{Diag}(k) \cong (\mathbf{C}^*)^{k-1}$  be the standard maximal torus of  $SL_k$  and  $X(T) \cong \mathbf{Z}^{k-1}$  its character lattice with standard basis  $e_1, \dots, e_{k-1}$ . Let

$$\lambda_i := \sum_{j=1}^i e_j \in X(T), \quad i = 1, \dots, k-1$$

be the fundamental weights and

$$\begin{aligned} \alpha_1 &:= e_1 - e_2 = -\lambda_2 + 2\lambda_1 && \in X(T) \\ \alpha_i &:= e_i - e_{i+1} = -\lambda_{i+1} + 2\lambda_i - \lambda_{i-1} && \in X(T), \quad i = 2, \dots, k-2 \\ \alpha_{k-1} &:= 2e_{k-1} + \sum_{j=1}^{k-2} e_j = 2\lambda_{k-1} - \lambda_{k-2} && \in X(T) \end{aligned}$$

the simple roots. Define  $q_i := \chi_{\lambda_i}$  for  $i = 1, \dots, k-1$ . So the set of positive roots is

$$\Phi^+ = \left\{ \sum_{i=a}^b \alpha_i \mid 1 \leq a \leq b \leq k-1 \right\} = \left\{ \frac{q_a q_b}{q_{a-1} q_{b+1}} \mid 1 \leq a \leq b \leq k-1, q_0 = q_k = 1 \right\}.$$

On  $V$ , the action of  $T$  is given by the matrix

$$\text{Diag}(q_1, q_2 q_1^{-1}, q_3 q_2^{-1}, \dots, q_{k-1} q_{k-2}^{-1}, q_{k-1}^{-1}).$$

Now with [3], Remark 4.6.10, the Hilbert series of  $\mathbf{C}[V^n]^{\text{SL}_k}$  is the coefficient of  $q_1^0 \cdots q_{k-1}^0$  when we set  $q_0 = q_k = 1$  in the  $q$ -Hilbert series:

$$\begin{aligned} H_q(\mathbf{C}[V^n]^{\text{SL}_k}) &= \frac{\prod_{1 \leq a \leq b \leq k-1} \left(1 - \frac{q_a q_b}{q_{a-1} q_{b+1}}\right)}{\prod_{j=1}^k \left(1 - t \frac{q_j}{q_{j-1}}\right)^n} \\ &= \frac{\left| \left( \left( \frac{q_i}{q_{i-1}} \right)^{j-i} \right)_{1 \leq i, j \leq k} \right|}{\prod_{j=1}^k \left(1 - t \frac{q_j}{q_{j-1}}\right)^n} \\ &= \left| \left( \left( \frac{q_i}{q_{i-1}} \right)^{j-i} \right)_{1 \leq i, j \leq k} \right| \sum_{i_1, \dots, i_k=0}^{\infty} \prod_{j=1}^k \binom{n}{i_j} \left( t \frac{q_j}{q_{j-1}} \right)^{i_j} \\ &= \sum_{\sigma \in S_k} \text{sgn}(\sigma) \sum_{i_1, \dots, i_k=0}^{\infty} \prod_{j=1}^k \binom{n}{i_j} t^{i_j} \left( \frac{q_j}{q_{j-1}} \right)^{\sigma(j) + i_j - j} \end{aligned}$$

For the second equality, we used Lemma 2 with  $z_i = q_i/q_{i-1}$ . Now for the coefficient of  $q_1^0 \cdots q_{k-1}^0$ , for each  $\sigma \in S_k$ , all the exponents  $\sigma(j) + i_j - j$  must be the same, so

that we get:

$$\begin{aligned}
H(\mathbf{C}[V^n]^{SL_k}) &= \sum_{\sigma \in S_k} \operatorname{sgn}(\sigma) \sum_{\substack{i_1, \dots, i_k \geq 0 \\ \sigma(1) + i_1 - 1 \\ \dots \\ \sigma(k) + i_k - k}} \prod_{j=1}^k \binom{n}{i_j} t^{i_j} \\
&= \sum_{\sigma \in S_k} \operatorname{sgn}(\sigma) \sum_{\substack{i_1 \geq \sigma(1) - 1 \\ \dots \\ i_k \geq \sigma(k) - k \\ i_1 = \dots = i_k}} \prod_{j=1}^k \binom{n}{i_j - \sigma(j) + j} t^{i_j - \sigma(j) + j} \\
&= \sum_{\sigma \in S_k} \operatorname{sgn}(\sigma) \sum_{l=0}^k \prod_{j=1}^k \binom{n}{l - \sigma(j) + j} t^{l - \sigma(j) + j} \\
&= \sum_{l=0}^{\infty} t^{kl} \sum_{\sigma \in S_k} \operatorname{sgn}(\sigma) \binom{n}{l - \sigma(j) + j} = \sum_{l=0}^{\infty} \left| \binom{n}{l - i + j} \right|_{1 \leq i, j \leq k} t^{kl} \\
&= \sum_{l=0}^{\infty} \{n, \dots, n\}_l t^{kl} = \sum_{l=0}^{\infty} [n - k + 1, \dots, n - k + 1]_l t^{kl} \\
&= \mathfrak{N}_{k, n-k+1}(t^k) = \frac{N_{k, n-k}(t)}{(1 - t^k)^{k(n-k)+1}}.
\end{aligned}$$

The second and third equalities hold because for each  $\sigma$ , if there is a  $1 \leq j \leq k$  with  $\sigma(j) - j > 0$ , there must be a  $j'$  with  $\sigma(j') - j' > 0$ . Furthermore, since  $k \leq n$ , for all  $1 \leq j \leq k$  we have  $-n < \sigma(j) - j$ . Thus the additional as well as the removed summands all are zero and the second and third equality hold. The last identity directly follows from Corollary 1.  $\square$

**Remark 1.** Of course the quotient of  $V^n$  by  $SL_k$  is also defined for  $n < k$ , but the only invariant functions in these cases are the constant ones, so that  $H(\mathbf{C}[V^n]^{SL_k}) = 1$  holds here.

The equality of our formula for the Hilbert polynomial and the one of Hodge is immediate. There is another way to compute it using the Borel-Weil-Bott Theorem, see Section 5.3 of [2]. For the connection to the approach [20] of Sturmfels, see also the paper [18].

As we already stated in Section 1, the numbers  $\{a_1, \dots, a_k\}_j$  with  $1 \leq a_1 < \dots < a_k \leq n$  give the Hilbert polynomials of Schubert varieties

$$X(a_1, \dots, a_k) := \{W \in \operatorname{Gr}(k, n) \mid \dim(W \cap \langle e_1, \dots, e_{a_i} \rangle) \geq i, i = 1, \dots, k\}.$$

From this, Nanduri in [16] deduced a closed form for the  $h$ -polynomial of  $X(a_1, \dots, a_k)$ . We give a slightly simpler formula in the following, which reduces to the formula of Sulanke for the  $k$ -Narayana numbers when we set  $a_1 = \dots = a_k = n$ .

**Proposition 3.** *The  $i$ -th coefficient  $h_i$  of the  $h$ -polynomial of the Schubert variety*

$X(a_1, \dots, a_k)$  of dimension  $d$  is given by

$$h_i = \sum_{l=0}^i (-1)^l \{a_1, \dots, a_k\}_{i-l} \binom{d}{l}.$$

*Proof.* According to the proof of Proposition 2.9 of [16], the coefficient  $h_i$  is given by

$$h_i = \sum_{l=0}^i (-1)^l \{a_1, \dots, a_k\}_{i-l} \binom{d-1}{l} - \sum_{l=0}^{i-1} (-1)^l \{a_1, \dots, a_k\}_{i-l-1} \binom{d-1}{l}.$$

We have

$$\begin{aligned} h_i &= \sum_{l=0}^i (-1)^l \{a_1, \dots, a_k\}_{i-l} \binom{d-1}{l} - \sum_{l=1}^{i-1} (-1)^{l-1} \{a_1, \dots, a_k\}_{i-l} \binom{d-1}{l-1} \\ &= \sum_{l=0}^i (-1)^l \{a_1, \dots, a_k\}_{i-l} \binom{d-1}{l} + \sum_{l=0}^{i-1} (-1)^l \{a_1, \dots, a_k\}_{i-l} \binom{d-1}{l-1} \\ &= \sum_{l=0}^i (-1)^l \{a_1, \dots, a_k\}_{i-l} \binom{d}{l}. \end{aligned} \quad \square$$

### 3 Generalized hypergeometric Euler transform

For integers  $p, q \geq 0$  and  $a_1, \dots, a_p, b_1, \dots, b_q \in \mathbf{C}$ , where no  $b_i$  is an integer smaller than one, recall the generalized hypergeometric function

$${}_pF_q \left( \begin{matrix} a_1, \dots, a_p \\ b_1, \dots, b_q \end{matrix}; t \right) := \sum_{k=0}^{\infty} \frac{(a_1)_k \cdots (a_p)_k}{(b_1)_k \cdots (b_q)_k} \frac{t^k}{k!},$$

where  $(a)_k$  is the Pochhammer symbol standing for the rising factorial

$$a(a+1) \cdots (a+k-1)$$

with  $(a)_0 = 1$ . Consider the Euler transformation for the ordinary hypergeometric function  ${}_2F_1$ :

$${}_2F_1 \left( \begin{matrix} a, b \\ c \end{matrix}; t \right) = (1-t)^{c-a-b} {}_2F_1 \left( \begin{matrix} c-a, c-b \\ c \end{matrix}; t \right).$$

This identity in particular shows that the 2-Narayana polynomial is a hypergeometric function, i.e. the identity

$$\sum_{l=0}^j (-1)^{j-l} \binom{2r+1}{j-l} \binom{r+l}{r} \binom{r}{r}^{-1} = \frac{1}{r} \binom{r}{j} \binom{r}{j+1}.$$

The hypergeometric Euler transform has been generalized to  ${}_pF_q$  in different ways, see [12, Theorem 2.1] and [13, Theorem 3] as well as [14, Theorem 4]. We will

use Theorem 4 of [14] to express  $k$ -Narayana numbers as products for  $k \geq 3$  as well. This leads to a representation of the  $h$ -polynomials of Grassmannians as hypergeometric functions for  $k \geq 3$ .

In order to do this, we first state the generalized hypergeometric Euler transform. Denote by  $\left\{ \begin{smallmatrix} x \\ y \end{smallmatrix} \right\}$  the Stirling numbers of the second kind. Consider for  $r \in \mathbf{Z}_{\geq 1}$  numbers  $m_1, \dots, m_r \in \mathbf{Z}_{\geq 1}$  and set  $m = \sum m_r$ . Furthermore let  $a, b, c, f_1, \dots, f_r \in \mathbf{C}$  meet the requirements

$$(c - a - m)_m, (c - b - m)_m, (1 + a + b - c)_m \neq 0.$$

Now for  $j, k = 0, \dots, m$  denote by  $\sigma_j$  the coefficient of  $x^j$  in the polynomial  $(f_1 + x)_{m_1} \cdots (f_r + x)_{m_r}$  and set

$$\begin{aligned} A_k &:= \sum_{j=k}^m \left\{ \begin{matrix} j \\ k \end{matrix} \right\} \sigma_j, \\ G_k(t) &:= {}_3F_2 \left( \begin{matrix} k - m, k - t, 1 - c - t \\ 1 + b + k - c - t, 1 + a + k - c - t \end{matrix}; 1 \right), \\ Q(t) &:= \sum_{k=0}^m (-1)^k A_k (a)_k (b)_k (t)_k (c - a - m - t)_{m-k} (c - b - m - t)_{m-k} G_k(-t). \end{aligned}$$

With these definitions, we get the following:

**Theorem 2 (Generalized hypergeometric Euler transform, [14], Th. 4).** *Let  $\eta_1, \dots, \eta_m$  be the nonvanishing zeros of the polynomial  $Q(t)$ . Then for  $|t| < 1$ , the following equality holds true:*

$$\begin{aligned} & {}_{r+2}F_{r+1} \left( \begin{matrix} a, b, f_1 + m_1, \dots, f_r + m_r \\ c, f_1, \dots, f_r \end{matrix}; t \right) \\ &= (1 - t)^{c-a-b-m} {}_{m+2}F_{m+1} \left( \begin{matrix} c - a - m, c - b - m, \eta_1 + 1, \dots, \eta_m + 1 \\ c, \eta_1, \dots, \eta_m \end{matrix}; t \right). \end{aligned}$$

Now consider the case of the  $k$ -Narayana series

$$\mathfrak{N}_{k,r}(t) = {}_kF_{k-1} \left( \begin{matrix} r, \dots, r + k - 1 \\ 2, \dots, k \end{matrix}; t \right)$$

with  $k \geq 3$ . Letting  $c := 2, a := r + k - 2, b := r + k - 1$  and for  $i = 3, \dots, k$  furthermore  $f_i := i, m_i := r - 3$ , so that  $m = \sum m_i = (k - 2)(r - 3)$ , we have that none of

$$\begin{aligned} & (4 - r - k - (k - 2)(r - 3))_{(k-2)(r-3)}, \\ & (3 - r - k - (k - 2)(r - 3))_{(k-2)(r-3)}, \\ & (2(r + k) - 4)_{(k-2)(r-3)} \end{aligned}$$

equals zero. We thus get:

**Theorem 3.** *The  $k$ -Narayana polynomial can be expressed as the hypergeometric function*

$$N_{k,r} = {}_{m+2}F_{m+1} \left( \begin{matrix} (k-1)(1-r), (k-1)(1-r)-1, \eta_1+1, \dots, \eta_m+1 \\ 2, \eta_1, \dots, \eta_m \end{matrix}; t \right)$$

with  $m = (k-2)(r-2)$ . Moreover, for the  $k$ -Narayana numbers  $N_k(r, j)$ , we have the product formula

$$N_k(r, j) = \frac{1}{j+1} \binom{(k-1)(r-1)}{j} \binom{(k-1)(r-1)+1}{j} \prod_{i=1}^{(k-2)(r-2)} \frac{\eta_i + j}{\eta_i},$$

where the  $\eta_i$  are the zeros in  $t$  of the polynomial

$$\sum_{l=0}^{(k-2)(r-2)} (-1)^l A_l (-k+1)r-3k-4-t \binom{(k-2)(r-2)-l}{(k-2)(r-2)-l} (-k+1)r-3k-3-t \binom{(k-2)(r-2)-l}{(k-2)(r-2)-l} \\ \times (r+k-1)_l (r+k)_l (t)_l {}_3F_2 \left( \begin{matrix} l-(k-2)(r-2), l+t, t-1 \\ r+k-1+l+t, r+k-2+l+t \end{matrix}; 1 \right).$$

*Proof.* Letting  $c = 2$ ,  $a = r+k-2$ ,  $b = r+k-1$  and for  $i = 3, \dots, k$  furthermore  $f_i = i$ ,  $m_i = r-3$  as stated above, the requirements of Theorem 2 are fulfilled and applying it we get

$$\mathfrak{N}_{k,r}(t) = {}_kF_{k-1} \left( \begin{matrix} r+k-2, r+k-1, r, \dots, r+k-3 \\ 2, 3, \dots, k \end{matrix}; t \right) \\ = \frac{{}_{m+2}F_{m+1} \left( \begin{matrix} r+2k-kr-2, r+2k-kr-3, \eta_1+1, \dots, \eta_m+1 \\ 2, \eta_1, \dots, \eta_m \end{matrix}; t \right)}{(1-t)^{(r-1)k+1}}$$

with  $m = (k-2)(r-3)$ . So with Corollary 1, we have

$$N_{k,r-1} = {}_{m+2}F_{m+1} \left( \begin{matrix} r(1-k)+2k-2, r(1-k)+2k-3, \eta_1+1, \dots, \eta_m+1 \\ 2, \eta_1, \dots, \eta_m \end{matrix}; t \right)$$

and thus with  $m' = (k-2)(r-2)$

$$N_k(r, j) = \frac{((k-1)(1-r))_j ((k-1)(1-r)-1)_j (\eta_1+1)_j \cdots (\eta_{m'}+1)_j}{(2)_j (\eta_1)_j \cdots (\eta_{m'})_j j!} \\ = \frac{1}{j+1} \binom{(k-1)(r-1)}{j} \binom{(k-1)(r-1)+1}{j} \prod_{i=1}^{(k-2)(r-2)} \frac{\eta_i + j}{\eta_i}.$$

□

**Corollary 2.** *The  $h$ -polynomial of the Grassmannian  $\text{Gr}(k, n)$  is given by*

$${}_{m+2}F_{m+1} \left( \begin{matrix} (k-1)(1-n+k), (k-1)(1-n+k)-1, \eta_1+1, \dots, \eta_m+1 \\ 2, \eta_1, \dots, \eta_m \end{matrix}; t \right)$$

with  $m = (n - k - 2)(k - 2)$ .

**Remark 2.** Consider the case  $k > r$ . Here we have the reduction

$$\begin{aligned} \mathfrak{N}_{k,r}(t) &= {}_kF_{k-1} \left( \begin{matrix} r, \dots, k, k+1, \dots, r+k-1 \\ 2, \dots, r-1, r, \dots, k \end{matrix}; t \right) \\ &= {}_{r-1}F_{r-2} \left( \begin{matrix} k+1, \dots, (k+1) + (r-1) - 1 \\ 2, \dots, r-1 \end{matrix}; t \right) = \mathfrak{N}_{r-1, k+1}(t), \end{aligned}$$

from what follows  $N_{k,r}(t) = N_{r,k}(t)$ . In this case, it makes sense to first use this reduction and afterwards apply Theorem 3 to get a simpler formula.

Moreover, the polynomial  $Q(t)$ , of which the zeros have to be computed, looks complicated. It is nevertheless no problem to compute these zeros with the help of computer algebra systems, but we propose another method that also provides some comparison between the computational complexity of the two different formulae: We know that the zeros  $\eta_i$  exist and that there are  $(k-2)(r-2)$  of them. So instead of computing them from the polynomial  $Q(t)$ , one can leave them as indeterminates, call the resulting function  $N_k(r, j)[\eta_1, \dots, \eta_{(k-2)(r-2)}]$ , compute  $N_k(r, j_l)$  for different  $j_1, \dots, j_{(k-2)(r-2)}$  and solve the system of equations

$$N_k(r, j_l)[\eta_1, \dots, \eta_{(k-2)(r-2)}] = N_k(r, j_l), \quad l = 1, \dots, (k-2)(r-2).$$

This is equivalent to a system of multilinear equations in the  $\eta_i$ , providing the same set of zeros as  $Q(t)$ . One should of course choose the values of  $j$  for which  $N_k(r, j)$  is easiest to compute with the old formula.

We can now roughly estimate the difference in complexity of the two formulae: we know that the  $N_k(r, j)$  are symmetric, but the binomial coefficients in the new formula are not, i.e.

$$N_k(r, j) = N_k(r, (r-1)(k-1) - j),$$

$$N_k(r, j)[\eta_1, \dots, \eta_{(k-2)(r-2)}] \neq N_k(r, j)[\eta_1, \dots, \eta_{(k-2)(r-2)}].$$

Moreover, computations suggest that there are always exactly  $2 \lfloor (k-2)(r-2)/2 \rfloor$  nonvanishing zeroes of  $Q(t)$ , which means we have to compute  $\lfloor (k-2)(r-2)/2 \rfloor$  values of  $N_k(r, j_l)$  in order to compute the  $\eta_i$ . We implemented the classical formula as well as the new one using both  $Q(t)$  and the systems of equations in Maple, available as ancillary worksheet file. Examples suggest that computing the  $\eta_i$  using  $Q(t)$  is faster than using the system of equations, but we can not exclude that this is due to our implementation. Nevertheless, once the  $\eta_i$  are known for fixed  $k$  and  $r$ , they can be stored and used to compute the  $N_k(r, j)$  more efficiently.

**Example 1.** We give some formulae from our computations in Maple for  $k = 3$  and low values of  $r$  in the following.

$r$	$N_k(r, j)$
4	$\frac{1}{105(j+1)} \binom{6}{j} \binom{7}{j} (3 + \sqrt{114} - j) (-3 + \sqrt{114} + j)$
5	$\frac{1}{63(j+1)} \binom{8}{j} \binom{9}{j} (4 + \sqrt{79} - j) (-4 + \sqrt{79} + j)$
6	$\frac{1}{27720(j+1)} \binom{10}{j} \binom{11}{j} (4 + j) (14 - j) (-5 + 2\sqrt{130} + j) (5 + 2\sqrt{130} - j)$
7	$\frac{1}{123552000(j+1)} \binom{12}{j} \binom{13}{j} (-60 + \sqrt{17450 - 10\sqrt{682705}} + 10j) (60 + \sqrt{17450 - 10\sqrt{682705}} - 10j) \times (-60 + \sqrt{17450 + 10\sqrt{682705}} + 10j) (60 + \sqrt{17450 + 10\sqrt{682705}} - 10j)$

## References

- [1] J. Berman, P. Köhler: Cardinalities of finite distributive lattices. *Mitt. Math. Sem. Giessen* 121 (1976) 103–124.
- [2] J.V. Chipalkatti: Notes on Grassmannians and Schubert varieties. *Queen’s Papers in Pure and Applied Mathematics* 13 (119) (2000).
- [3] H. Derksen, G. Kemper: *Computational invariant theory*. Springer (2015).
- [4] S.R. Ghorpade: Note on Hodge’s postulation formula for Schubert varieties. In: *Lecture Notes in Pure and Applied Mathematics*. Marcel Dekker, New York (2001) 211–220.
- [5] B.H. Gross, N.R. Wallach: On the Hilbert polynomials and Hilbert series of homogeneous projective varieties. *Arithmetic geometry and automorphic forms* 19 (2011) 253–263.
- [6] W.V.D. Hodge: A note on  $k$ -connexes. In: *Mathematical Proceedings of the Cambridge Philosophical Society*. Cambridge University Press (1942) 129–143.
- [7] W.V.D Hodge: Some enumerative results in the theory of forms. In: *Mathematical Proceedings of the Cambridge Philosophical Society*. Cambridge University Press (1943) 22–30.
- [8] W.V.D. Hodge, D. Pedoe: *Methods of algebraic geometry, Volume 2*. Cambridge University Press (1952).
- [9] V. Lakshmbai, J. Brown: *The Grassmannian Variety: Geometric and Representation-theoretic Aspects*. Springer (2015).
- [10] D.E. Littlewood: On the number of terms in a simple algebraic form. In: *Mathematical Proceedings of the Cambridge Philosophical Society*. Cambridge University Press (1942) 394–396.
- [11] P.A. MacMahon: *Combinatory analysis*. Cambridge University Press (1916).
- [12] R. Maier: A generalization of Euler’s hypergeometric transformation. *Transactions of the American Mathematical Society* 358 (1) (2006) 39–57.
- [13] A.R. Miller, R.B. Paris: Euler-type transformations for the generalized hypergeometric function  ${}_{r+2}F_{r+1}(x)$ . *Zeitschrift für angewandte Mathematik und Physik* 62 (1) (2011) 31–45.

- [14] A.R. Miller, R.B. Paris: Transformation formulas for the generalized hypergeometric function with integral parameter differences. *Rocky Mountain Journal of Mathematics* 43 (1) (2013) 291–327.
- [15] S. Mukai: *An introduction to invariants and moduli*. Cambridge University Press (2003). Cambridge Studies in Advanced Mathematics 81
- [16] R. Nanduri: Hilbert coefficients of Schubert varieties in Grassmannians. *Journal of Algebra and Its Applications* 14 (3) (2015) 1550036.
- [17] T.V. Narayana: A partial order and its applications to probability theory. *Sankhya: The Indian Journal of Statistics* (1959) 91–98.
- [18] F. Santos, C. Stump, V. Welker: Noncrossing sets and a Graßmann associahedron. In: *Forum of Mathematics, Sigma*. Cambridge University Press (2017) 49p.
- [19] R.P. Stanley: Theory and application of plane partitions. Part 2. *Studies in Applied Mathematics* 50 (3) (1971) 259–279.
- [20] B. Sturmfels: *Gröbner bases and convex polytopes*. American Mathematical Society (1996). University Lecture Series Vol. 8.
- [21] R.A. Sulanke: Generalizing Narayana and Schröder numbers to higher dimensions. *The Electronic Journal of Combinatorics* 11 (1) (2004) 54p.
- [22] R.A. Sulanke: Three dimensional Narayana and Schröder numbers. *Theoretical computer science* 346 (2–3) (2005) 455–468.

Received: 4 June, 2018

Accepted for publication: 18 December, 2018

Communicated by: Eric Swartz