



HAL
open science

A simple construction of basic polynomials invariant under the Weyl group of the simple finite-dimensional complex Lie algebra

Askold M. Perelomov

► **To cite this version:**

Askold M. Perelomov. A simple construction of basic polynomials invariant under the Weyl group of the simple finite-dimensional complex Lie algebra. *Communications in Mathematics*, 2021, Volume 28 (2020), Issue 3 (3), pp.301 - 305. 10.2478/cm-2020-0024 . hal-03664999

HAL Id: hal-03664999

<https://hal.science/hal-03664999>

Submitted on 11 May 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

A simple construction of basic polynomials invariant under the Weyl group of the simple finite-dimensional complex Lie algebra

Askold M. Perelomov

Abstract. For every simple finite-dimensional complex Lie algebra, I give a simple construction of all (except for the Pfaffian) basic polynomials invariant under the Weyl group. The answer is given in terms of the two basic polynomials of smallest degree.

1 Basics

For necessary information on Lie algebras and Lie groups, see [5]. Recall the notation: let \mathfrak{g} be a simple finite-dimensional complex Lie algebra of rank l , let R_+ (resp. R_-) be the set of its positive (resp. negative) roots, and $\{\alpha_1, \dots, \alpha_l\}$ the set of simple roots. Let the Weyl group $W_{\mathfrak{g}}$ of the root system R act in the space $V = \mathbb{R}^l$, let $(-, -)$ be the non-degenerate $W_{\mathfrak{g}}$ -invariant bilinear form on V , such that $(x, y) = \sum_{j=1}^l x_j y_j$ for any $x, y \in V$; let h be the Coxeter number; $\delta = \sum_{j=1}^l b_j \alpha_j$ the highest root; $b = \max b_j$.

As is known (by abuse of notation we denote the Lie algebra $\mathfrak{sl}(n+1)$ by the symbol of its root system A_n in Cartan's notation, and similarly for the other

2020 MSC: 11F22

Key words: Weyl group, invariant polynomial

Affiliation:

Institute for Theoretical and Experimental Physics, B. Cheremushkinskaya ul. 25,
 117259, Moscow, Russia
 E-mail: apereलोmo@gmail.com

simple Lie algebras),

$$b = \begin{cases} 1 & \text{for } A_l, \\ 2 & \text{for } B_l, C_l \text{ and } D_l, \\ 3 & \text{for } G_2 \text{ and } E_6, \\ 4 & \text{for } F_4 \text{ and } E_7, \\ 6 & \text{for } E_8; \end{cases} \quad h = \begin{cases} l + 1 & \text{for } A_l, \\ 2l & \text{for } B_l, \text{ and } C_l, \\ 2l - 2 & \text{for } D_l, \\ 12 & \text{for } E_6 \text{ and } F_4, \\ 18 & \text{for } E_7, \\ 30 & \text{for } E_8, \\ 6 & \text{for } G_2. \end{cases} \quad (1)$$

If $\alpha = \sum_{j=1}^l n_j \alpha_j \in R_+$, then we define the *height* of α to be $ht(\alpha) = \sum_{j=1}^l n_j$.

As it was shown in [1], the algebra of invariant polynomials is generated by l basic homogeneous polynomials of degrees d_j for $j = 1, \dots, l$ such that $d_1 = 2$, $d_j \leq d_{j+1}$, $d_l = h$.

Note that

$$d_j = \begin{cases} j + 1 & \text{for } A_l \\ 2j & \text{for } B_l \text{ and } C_l, \end{cases} \quad (2)$$

$$\{d_j \mid j = 1, \dots, l - 1, l\} = \{2, 4, \dots, 2(l - 1), l\}, \text{ as ordered sets, for } D_l. \quad (3)$$

For the exceptional simple Lie algebras, the d_j were first found in [6] using results of [7]:

$$\{d_1, \dots, d_l\} = \begin{cases} \{2, 6\} & \text{for } G_2; \\ \{2, 6, 8, 12\} & \text{for } F_4; \\ \{2, 5, 6, 8, 9, 12\} & \text{for } E_6; \\ \{2, 6, 8, 10, 12, 14, 18\} & \text{for } E_7; \\ \{2, 8, 12, 14, 18, 20, 24, 30\} & \text{for } E_8. \end{cases} \quad (4)$$

Note that the degrees satisfy duality relations

$$d_j + d_{l+1-j} = h + 2. \quad (5)$$

Denote by I_j the invariant polynomial of degree d_j in eq. (2), (3), and (4).

2 Degrees and exponents

Let us remind a characterization of *exponents*, i.e., numbers $m_j := d_j - 1$, given in [2]. Let n_k be the number of roots of height k . Then $n_k - n_{k+1}$ is the number of times k occurs as an exponent of \mathfrak{g} .

Note that $d_1=2$ for all simple Lie algebras. Recall the values of b , see (1). The values of d_2, d_3 and d_4 are given by

Theorem 1. *We have*

$$\begin{aligned} d_2 &= b + 2 && \text{for all } \mathfrak{g}, \text{ except } G_2, \\ d_3 &= 2b && \text{for all exceptional } \mathfrak{g}, \text{ except } G_2, \\ d_4 &= 2b + 2 && \text{for } E_6, E_7 \text{ and } E_8. \end{aligned}$$

The other quantities d_j can be obtained from duality.

The proof follows easily from Table (4) and duality eq. (5) borrowed from any sufficiently thick book (e.g., [5]) and papers [6], [7]. Below we give an independent proof.

Proof. We give the proof only for the most complicated case $\mathfrak{g} = E_8$. For all other cases the proof is analogous.

We enumerate the simple roots of E_8 first along the Dynkin diagram, starting from the longest end of the branch, as the simple roots of A_7 , and set

$$\begin{aligned} (\alpha_8, \alpha_5) &= -1, \\ (\alpha_8, \alpha_k) &= 0 \quad \text{for } k \neq 5, 8, \\ (\alpha_j, \alpha_j) &= 2 \quad \text{for } j = 1, \dots, 8. \end{aligned}$$

Then, as it is well-known [5], the highest root δ satisfies the conditions:

$$\begin{aligned} (\alpha_1, \delta) &= 1, \\ (\alpha_j, \delta) &= 0 \quad \text{for } j = 2, \dots, 8, \end{aligned}$$

and hence it has the form

$$\delta = 2\alpha_1 + 3\alpha_2 + 4\alpha_3 + 5\alpha_4 + 6\alpha_5 + 4\alpha_6 + 2\alpha_7 + 3\alpha_8, \quad ht(\delta) = 29 = h - 1.$$

Let us consider the set of positive roots decreasing in height. We denote

$$\beta_j := \alpha_1 + \alpha_2 + \dots + \alpha_j \quad \text{for } j = 1, 2, \dots, 8.$$

Then we see that

(1) One root for the each height in the interval $[h - 1, h - b] = [29, 24]$, namely,

$$\delta, \delta - \beta_1, \delta - \beta_2, \delta - \beta_3, \delta - \beta_4, \text{ and } \delta - \beta_5.$$

(2) Two roots for the each height in the interval $[h - b - 1, h - 2b + 2] = [23, 20]$, namely,

$$\begin{array}{ll} \delta - \beta_6, & \delta - (\beta_5 + \alpha_8); \\ \delta - \beta_7, & \delta - (\beta_6 + \alpha_8); \\ \delta - \beta_8, & \delta - (\beta_6 + \alpha_5 + \alpha_8); \\ \delta - (\beta_8 + \alpha_5), & \delta - (\beta_6 + \alpha_4 + \alpha_5 + \alpha_8). \end{array}$$

Note that this is due to the fact that node 5 in the Dynkin diagram for E_8 is the branch node, and $b = 6$.

(3) Three roots for height $h - 2b + 1 = 19$, namely,

$$\delta - (\beta_8 + \alpha_5 + \alpha_6), \quad \delta - (\beta_8 + \alpha_4 + \alpha_5), \quad \delta - (\beta_6 + \alpha_3 + \alpha_4 + \alpha_5 + \alpha_8).$$

Hence, from Kostant's characterization of exponents [2] we have

$$m_7 = h - 1 - b = 23, \quad m_6 = h - 2b + 1 = 19,$$

and from the duality we have

$$m_2 = b + 1 = 7, \quad m_3 = 2b - 1 = 11.$$

Note that $m_1 = 1$ and $m_8 = 29 = h - 1$. So,

$$d_2 = m_2 + 1 = b + 2 = 8, \quad d_3 = m_3 + 1 = 2b = 12.$$

We analogously obtain $d_4 = 2b + 2 = 14$. □

3 Main Theorem

For an explicit construction of $W_{\mathfrak{g}}$ -invariant polynomials, see the papers by Mehta [4] and by Macdonald [3].

Let $\Delta = \sum \frac{\partial^2}{\partial x_j^2}$, and $AI = (\nabla I_2, \nabla I)$ for any polynomial I , where

$$\nabla := \left(\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_l} \right);$$

clearly, $\deg AI = \deg I + b$.

Theorem 2. *All basic $W_{\mathfrak{g}}$ -invariant polynomials for all algebras \mathfrak{g} , except for D_l and G_2 , can be obtained by applying A^k for some k to polynomials I_1 and I_3 , where $I_3 = \Delta(AI_2)$.*

More precisely, for the root systems A_l, B_l, C_l , and D_l , we have

$$I_{j+1} = \begin{cases} A^j I_1 & \text{for } j = 0, \dots, l - 1 & \text{for the root systems } A_l, B_l, \text{ and } C_l, \\ A^j I_1 & \text{for } j = 0, \dots, l - 2 & \text{for the root system } D_l, \end{cases}$$

and we obtain all basic polynomials, except for the Pfaffian for D_l .

For the root systems F_4, E_6, E_7 and E_8 the basic invariant polynomials have the form: $A^j I_1$ and $A^k I_3$, where

$$\begin{array}{lll} j = 0, 1 & \text{and } k = 0, 1 & \text{for } F_4; \\ j = 0, 1, 2 & \text{and } k = 0, 1, 2 & \text{for } E_6; \\ j = 0, 1, 2, 3, 4 & \text{and } k = 0, 1 & \text{for } E_7; \\ j = 0, 1, 2, 3 & \text{and } k = 0, 1, 2, 3 & \text{for } E_8. \end{array}$$

Proof. Consider the set of polynomials $A^j I_1$ and $A^k I_3$ for fixed Lie algebra \mathfrak{g} of rank l from the list of Theorem 2. We obtain all l first $W_{\mathfrak{g}}$ -invariant polynomials I_l , except for the Pfaffian for D_l . The explicit formulas in the paper [4] imply that these l polynomials are algebraically independent. So we may take these polynomials, and the Pfaffian for a basis of $W_{\mathfrak{g}}$ -invariant polynomials. □

4 Acknowledgments

I am thankful to D. Leites for useful remarks; he also improved my English in this paper.

References

- [1] C. Chevalley: Invariants of finite groups generated by reflections. *American Journal of Mathematics* 77 (4) (1955) 778–782.
- [2] B. Kostant: The principal three-dimensional subgroup and the Betti numbers of a complex simple Lie group. *American Journal of Mathematics* 81 (1959) 973–1032.
- [3] I.G. Macdonald: Orthogonal polynomials associated with root systems. *Séminaire Lotharingien de Combinatoire* 45 (2000) B45a.
- [4] M.L. Mehta: Basic sets of invariant polynomials for finite reflection groups. *Communications in Algebra* 16 (5) (1988) 1083–1098.
- [5] A.L. Onishchik, E.B. Vinberg: *Lie groups and algebraic groups*. Springer (1990).
- [6] G. Racah: Sulla caratterizzazione delle rappresentazioni irriducibili dei gruppi semisemplici di Lie. *Lincei-Rend. Sc. fis. mat. e nat* 8 (1950) 108–112.
- [7] E. Witt: Spiegelungsgruppen und aufzählung halbeinfacher liescher Ringe. *Abhandlungen aus dem mathematischen Seminar der Universität Hamburg* 14 (1) (1941) 289–322.

Received: 22 June 2019

Accepted for publication: 2 July 2019

Communicated by: Pasha Zusmanovich