



HAL
open science

Which hydrological activity in weathered bedrock during flash floods?

Audrey Douinot, H el ene Roux, Pierre-Andr e Garambois, K evin Larnier, Denis Dartus

► To cite this version:

Audrey Douinot, H el ene Roux, Pierre-Andr e Garambois, K evin Larnier, Denis Dartus. Which hydrological activity in weathered bedrock during flash floods?. 8th HyMeX Workshop - Hydrological cycle in Mediterranean Experiment - HyMeX 2014, Sep 2014, Valetta, Malta. pp. 1-1. <hal-03665191>

HAL Id: hal-03665191

<https://hal.science/hal-03665191v1>

Submitted on 11 May 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin ee au d ep ot et  a la diffusion de documents scientifiques de niveau recherche, publi es ou non,  emanant des  tablissements d'enseignement et de recherche fran ais ou  trangers, des laboratoires publics ou priv es.



HAL Authorization



Open Archive TOULOUSE Archive Ouverte (OATAO)

OATAO is an open access repository that collects the work of Toulouse researchers and makes it freely available over the web where possible.

This is an author-deposited version published in : <http://oatao.univ-toulouse.fr/>
Eprints ID : 13870

To link to this article :

URL : <http://www.hymex.org/?page=public/workshops/8/programme>

To cite this version : Douinot, Audrey and Roux, H  l  ne and Garambois, Pierre-Andr   and Larnier, Kevin and Dartus, Denis *Which hydrological activity in weathered bedrock during flash floods? The case of Mediterranean catchments*. (2014) In: 8th HyMeX Workshop - Hydrological cycle in Mediterranean Experiment - HyMeX 2014, 15 September 2014 - 18 September 2014 (Valetta, Malta). (Unpublished)

Any correspondance concerning this service should be sent to the repository administrator: staff-oatao@listes-diff.inp-toulouse.fr


Which hydrological activity in weathered bedrock during flash floods? The case of Mediterranean catchments

Audrey Douinot, Hélène Roux, Pierre-André Garambois, Kévin Larnier, Denis Dartus


Université de Toulouse ; INPT, UPS ; IMFT (Institut de Mécanique des Fluides de Toulouse) ; F-31400 Toulouse, France
CNRS ; IMFT ; F-31400 Toulouse, France

Objectives

- Assessing hydrological dynamic into weathered rock layer with:
 - hydrological modeling: MARINE (Roux et al., 2011)
 - baseflow separation (Eckhardt, 2005)
- Case study: 8 catchments of French mediterranean area



MARINE model description




A physical process oriented model with 5 parameters to calibrate (τ , soil depth: C_z)

Estimation of flow and storage into weathered rock layers through MARINE modeling


2 hydrological model processing :

- MARINE forcing with soil depth from BD-sol: subsurface flow estimations
- MARINE forcing with likelihood calibration: flow into weathered rock layers (WR F.) estimations




$C_z \times$
Correction coefficient of soil depth

- Significant hydrological dynamic into deep layers during flash flood
- Groundwater behavior specific to rock layer



% of flow through deep horizons: estimation on several flash flood events (* number of events)

Hydrograph decomposition (Eckhardt, 2005)



Using 2 parameters recursive filter to assess baseflow

$$b_t = \frac{(1 - BFI) \cdot a \cdot b_{t-1} + (1 - a) \cdot BFI \cdot y_t}{1 - a \cdot BFI}$$

b_t : baseflow at time t BFI : baseflow index
 y_t : streamflow at time t a : recession constant

Recession constants are computed making recession analysis. The BFI parameter is fixed to 0.8 as (Eckhardt, 2005) suggests.

- Significant baseflow during flash flood : it represents between 10 % and 46 % of the flash flood streamflow volumes (statistic achieved with 14 events)
- There isn't any correlation between baseflow contribution and catchment geology, because of the fixed BFI parameter to apply the decomposition method

Conclusion & Perspectives

Flows through deep horizons were estimated to represent up to 29 % and 46% of the total hydrograph according to MARINE hydrological modelling and hydrograph decomposition respectively. These results are consistent with previous studies (Vannier et al, 2013) that underline hydrological activity into weathered bedrock layers with significant contribution to flash flood processing.

Similar results are provided by both independent hydrological modeling and hydrograph decomposition methods, that give them consistency.

For more accurate results the hydrological model should detail subsurface flows and flows into deeper horizons with appropriate physical modeling. Hydrograph decomposition could help to find different flow contributions. The method has to be improved, by calibrating BFI parameter and computing the recession constant by focusing on flash flood events.

References

- Roux, H., Labat, D., Garambois, P. A., Maubourguet, M.-M., Chorda, J. & Dartus, D. 2011. A physically-based parsimonious hydrological model for flash floods in Mediterranean catchment *Nat. Hazards Earth Syst. Sci. J1 - NHESS*, 161, 2567–2582.
- Eckhardt, K. 2005. How to construct recursive digital filters for baseflow separation. *Hydrological Processes*, 19, 507-515.
- Vannier, O., Braud, I., Anquetin, S. 2013. Regional estimation of catchment-scale soil properties by means of streamflow recession analysis for use in distributed hydrological models. *Hydrological Processes*.