

Collembola dispersion, selection, and biological interactions in urban ecosystems: a review

Sophie S. Joimel, Antonin Jules, L. Vieublé-Gonod

▶ To cite this version:

Sophie S. Joimel, Antonin Jules, L. Vieublé-Gonod. Collembola dispersion, selection, and biological interactions in urban ecosystems: a review. Environmental Chemistry Letters, 2022, 20 (3), pp.2123-2133. 10.1007/s10311-022-01406-z . hal-03665441

HAL Id: hal-03665441 https://hal.science/hal-03665441v1

Submitted on 11 May 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Collembola dispersion, selection, and biological interactions in urban ecosystems: a review 2 3 Sophie Joimel^a, Antonin Jules^a, Laure Vieublé Gonod^a 4 5 ^a Université Paris-Saclay, INRAE, AgroParisTech, UMR EcoSys, 78850 Thiverval-Grignon 6 7 * Corresponding author: Sophie Joimel Université Paris-Saclay, INRAE, AgroParisTech, UMR EcoSys, 78850 Thiverval-Grignon 8 9 Phone: + 33 1 30 81 59 58 Fax: + 33 1 30 81 55 63 10 E-mail address: sophie.boulanger-joimel@agroparistech.fr 11 12 13 Abstract Most of the ecological predictions that are intended to be general have been made on the basis 14 of observations of aerial organisms, ignoring subterranean organisms, which are subject to 15 other - and complex - constraints compared from those present on the surface. Collembola, a 16 17 soil fauna group, are good biological indicators of soil quality and could be used as model

17 soil rauna group, are good biological indicators of soil quality and could be used as model 18 organisms in order to understand assembly rules of soil organisms in cities. Here we review 19 available literature on soil Collembola conducted on 75 publications dealing with urban 20 context.

The major points are the following: (1) Soil properties and landscape characteristics influenced soil communities. The importance of landscape is also highlighted by the need of connectivity between urban green spaces in order to allow dispersion of Collembola. However, studies of Collembola communities in urban context are recent, offering little hindsight to understand their assembly. (2) A gap of references values, complexities by a lack of methods standardisation, is observable. Especially, knowledge on dispersion and biotic
interactions are still too rare. (3) Moreover, studies on temporal aspect in order to understand
better the colonisation processes are almost absent of the corpus.

29

30

31 Keywords (4-6): Urban soil, soil biodiversity, assembly rules, dispersal, habitat, biotic
32 interactions

33

34

35 **1. Introduction**

36 Protecting biodiversity in the coming decades is a major challenge as the global assessment 37 report on biodiversity and ecosystem services has estimated that around 1 million animal and 38 plant species are threatened by extinction (IPBES, 2019).

39 Representing only 3% of the Earth's surface (MEA 2005), urban environments have long been neglected by ecological and biological sciences. Until 1995, only approximately one hundred 40 41 publications in ecology focused on the urban environment. Today, urban ecology papers 42 account for 14% of all ecology papers (Barot et al., 2019). This interest is due in particular to the increase in societal issues related to the city due to (i) the city is the living environment for 43 the majority of the Earth's inhabitants. Since 1950, the urbanization process has been 44 undergoing a change that accelerated in the 1990s with the increase in urban population; (ii) 45 46 urban ecosystems must be multifunctional. Cities, which are very "mineral", must evolve to become multifunctional and resilient in order to meet the new needs of ever-increasing 47 48 numbers of city dwellers; and (iii) the urban environment condenses multiple environmental problems such as greenhouse gas emissions, air, water and soil pollution and urban waste 49 50 management (Liu et al., 2014). The emergence of these societal issues is reflected in the

promotion of a more present nature in cities, which must provide the multiple services 51 necessary for the proper functioning of cities. In a context of biodiversity erosion, we can 52 wonder to what extent the multiplication and diversification of urban green spaces can impact 53 54 biodiversity. Preliminary results showed that the effects of urbanization on biodiversity vary according to taxonomic groups, environmental conditions and socio-economic contexts (Liu 55 et al., 2014). However, some authors seem to agree on several findings: urbanization modifies 56 the biodiversity of organisms present in cities. In terms of taxonomic diversity, an increase as 57 58 well as a decrease along an urbanization gradient can be observed between groups (McKinney, 2008). In terms of composition, urban biodiversity is made up of native and 59 exotic species, often generalist and synanthropic, and remnants of less tolerant species that 60 may disappear (Shochat et al., 2006). These conclusions were drawn as a result of studies on 61 birds, plants and more recently epigean arthropods (e.g. Marzluff, 2019; Sukopp, 2008). 62

However, urban soil biodiversity is largely under-studied. To neglect soil biodiversity - or subterranean biodiversity - is to neglect a quarter of global taxonomic diversity (Decaëns et al., 2006). This compartment is often neglected because of its lack of visibility and the difficulties inherent in its study, which often gives it the title of third biotic frontier - after the canopies of tropical forests and abysses - or black box (André et al., 1994).

Among soil organisms, Collembola, that belong to the mesofauna (0.1 at 2 mm), are known to perform major functions in soils: organic matter decomposition, micro-aggregation and population regulation (Hopkin, 1997; Rusek, 1998). In addition, their great diversity and abundance in soils makes it possible to evaluate effect of environmental factors on their community in different urban habitats. Finally, contrary to larger soil organisms such as earthworms, they are present even in soils of low depth such as soils of green roofs.

In this context, the objective of this study was to draw a state of Collembola communities incities in order to draw conclusions on the part of biodiversity of urban soils and to guide

future research. In order to understand the provision of biodiversity service and the 76 functioning of ecosystems, it is necessary to understand the determinisms of biodiversity 77 distribution at different scales. Also, the synthesis is divided in three parts articulated within 78 the conceptual framework of filter theory (Lortie et al., 2004). We therefore selected three 79 questions: (i) how do Collembola arrive in the city (capacities and means of dispersal)?, (ii) 80 what are the links between the biodiversity of Collembola and habitat characteristics (e.g. 81 reference to ecological niches)? and (iii) What are the relationships between Collembola and 82 83 other organisms (interactions, co-occurences)?

84

86

85 2. Bibliometric methods and analysis

The research focused on urban Collembola was conducted in April 2021 in ISI web of 87 Knowledge among all database offered. Key words selected for topic were: "(collembol* or 88 89 springtail*) and (urban* or city or cities or technosol* or anthroposol*). As showed in Fig.1, 90 the first studies concerning urban Collembola communities appeared between the 60's and the 80's, and were more regular only from the 90's. This concurs with the development of urban 91 92 ecology (Liu et al., 2014; Young and Wolf, 2006) and other ecological urban soil studies (Guilland et al., 2018). Nowadays, the proportion of publications dealing with urban 93 Collembola represents around 4-5% of the total number of publications dealing with 94 Collembola. These figures are of the same order of magnitude as those highlighted by 95 Guilland et al. (2018) for the biology of urban soils, confirming that urban soil organisms are 96 97 still poorly studied and unevenly between taxa.




Figure 1 : Number of publications dealing with Collembola since 1956; light grey bars: number of publications dealing with Collembola; black bars: number of publications dealing with Collembola in a urban context and dark grey line: proportion of publications in a urban context

99

98

100 172 articles came out of the query. A first selection was applied on the base of titles 101 and abstracts to keep papers that mention soil Collembola communities in an urban habitat, 102 their dispersal capacity towards or within cities or interactions with other organisms in urban 103 soils. Thus, papers describing a new species or solely evaluating physiological responses were 104 excluded. Finally, 75 publications were gathered as a corpus for analysis (S1, S2). From each 105 article, information concerning study location, land-use, sampling methods as well as 106 abundance, density, richness and species composition was collected. The access was restricted 107 to the abstract for 22 articles: 17 were not published in English and 5 were not accessible, 108 limiting the available information. Thus the in-depth analysis of the results was carried out 109 only on 53 articles.

A disparity was observed in the geographic origin of the studies. Indeed, 51% of them were conducted in Europe and 26% in Asia (n=75). Rare works were recorded in Africa, South America and Oceania while they host distinctive and probably endemic species (Culik and Filho, 2003; Hopkin, 1997; Janion-Scheepers et al., 2015). As the vast majority of the unavailable papers were from Asia and more precisely from China (17 out of 22), Europe
represents 66% of the publications of the reduced corpus (n=53).

116 Besides, aims and methods differed between the studies. A minor part of them, mainly from 117 China, was focused on describing the composition of urban soil fauna communities, giving 118 little insights about Collembola assemblages according to the three filters. The others tested 119 the effect of environmental parameters on the soil communities, providing results that can be helpful in the understanding of Collembola responses to urban pressures. Among them, the 120 121 most tested parameters were soil characteristics, land-use and metal contaminations. Differences in taxonomic levels were also noted in the restricted corpus (n=53) for which 122 50% of the publications dealt exclusively with Collembola species. The others were multi-123 organisms studies and often used a wider taxonomic level such as arthropods orders 124 (microarhtropods, invertebrates). Furthermore, several studies do not indicate mean species 125 126 richness, relative abundance per habitat or site or species composition but only a total number 127 of species or organisms in the study (e.g. Biffi et al., 2019; Rochefort et al., 2013). When 128 abundance values (number of individuals per mass of soil) were provided, they were converted into density values (number of individuals per square meter), but a bias lies in the 129 130 variable depth of soil cores used that is not included in the conversion. Moreover, there was 131 no standardized method for sampling. Volume or surface of soil cores, number of replicates, 132 collection tools (pitfall traps or soil cores), extraction units, and seasons often vary between 133 studies, whereas they influence results (Lang, 2000; Querner and Bruckner, 2010; Santorufo 134 et al., 2014b).

Each publication was carefully read to determine if the study concerned habitat, dispersal or biotic interactions. Whenever a habitat or environmental characteristics were specified, the habitat was considered, which was the case in 89% of the papers (n=53). Biotic interactions and dispersal ways were described much more rarely, 52% and 28% respectively 139 (n=53). The former was considered whenever Collembola were studied simultaneously with
140 other taxa, even though only nine studies really investigated interactions between organisms.
141 The latter was taken into account for each article that mentioned a way for Collembola to
142 move from a place to another.

143 There is no official or consensual classification to characterise urban green spaces (de 144 Groot et al., 2002) or urban soils which are often gathered on the term-of SUITMA (soils of 145 urban, industrial, traffic, mining and military areas (Joimel et al., 2016; Morel et al., 2014). 146 For the purpose of this study, we chose to address this classification through their function 147 within the city. We adapted the PPG17 guidelines of United Kingdom, which listed 17 148 distinctive types of green spaces predominantly focused on land uses. We added green roofs, 149 and removed habitats related to water land uses. In total 15 distinctive types of green spaces were defined : (i) Parks & gardens: country, urban and local parks, public gardens, (ii) street 150 151 sides, rails, roads, (iii) Institutional grounds, (iv) Built-up areas residential land, (v) Industrial 152 and commercial sites, (vi) educational establishments, (vii) Outdoor sports facilities : 153 golfcourses, tennis, sport fields, (viii) Churchyards, cemeteries, burial grounds, (ix) 154 Brownfield land: quarries, wastelands, landfills, vacant and derelict land, (x) Community green spaces: allotments, community gardens, urban farms, (xi) Horticultural establishments, 155 156 (xii) Natural & semi-natural green spaces: woodlands, forests, reserves, heathlands, grassland, 157 meadow, conservation land, (xiii) Avenue trees, (xiv) Agricultural land, farms, ranches, (xv) 158 green roofs and vertical forms.

- 159
- 160 161

3. Dispersion and colonisation in urban environment

Dispersion refers to the mobility or transport of organisms from a source location to a urban
habitat. Colonisation refers to the spatio-temporal dynamics in one habitat. The distinction
between dispersion and colonisation in the studied corpus is not always simple. We estimated

that only 10 articles dealt with Collembola dispersion and few articles treated it as a main
topic (e.g. Joimel et al., 2018; Rumble et al., 2018), illustrating the lack of knowledge in the
connectivity of the belowground communities in urban habitats. Concerning colonisation,
only three publications studied colonisation (Burrow et al., 2018; Rumble and Gange, 2013;
Santorufo et al., 2021).

170

Dispersion is one of the major drivers that rules assemblages patterns at a local or landscape 171 172 scale (Lortie et al., 2004; Zobel, 1997). Collembola are known to disperse in different ways 173 (Dunger et al., 2002). Dispersion may be the result of active (organism controlling the 174 locomotion leading to dispersal process) and passive (movement depending on external forces 175 such as wind, water currents or animals and materials used for the construction of Technosols) 176 mechanisms (Clobert et al., 2012), and concerns both airborne and soil organisms (Dunger, 177 1998). Although, their dispersion abilities are not well understood (Auclerc et al., 2009), 178 Collembola are supposed to actively disperse, probably in a range of few hundreds of meters 179 (Ponge, 2013). However, in urban context, experiments and insights about active dispersion of 180 Collembola is still absent from the published literature.

On the other hand several studies offer food to thought about how passive dispersion takes 181 182 place in urbanised landscape, especially the five articles concerning green roofs (Joimel et al., 2018; Rumble et al., 2018; Rumble and Gange, 2013; Schindler et al., 2019; Schrader and 183 184 Böning, 2006). Indeed, in studying mechanisms behind dispersion, it has been shown that 185 Collembola are able to passively disperse those newly formed urban soils as they were present in all the roofs substrates. Nevertheless Collembola diversity ranged from 6 (Rumble and 186 187 Gange, 2013) to 44 species (Joimel et al., 2018) in all sampled roofs. Questions then arise: 188 how did they get there and is it a one-off or a regular process?

189 Schrader and Böning (2006) already suggested that dispersion onto green roofs was a190 dynamic process over time. They found significant differences in species composition

between young (<4 years) and old (>8 years) roofs, as the result of several communities 191 192 successions. This comforts the hypothesis that Collembola can reach green roofs even after 193 their construction. Rumble et al. (2018) observed a Collembola species (Sminthurinus 194 *trinotatus*) that was neither present in the original substrate or in *Sedum* plugs before planting. 195 The recent use of functional traits in urban Collembola assemblages studies brought some 196 insights to understand their dispersion in cities (Joimel et al., 2018). The study of dominant 197 traits revealed that Collembola of productive and extensive green roofs were characterized by 198 a small size, a parthenogenetic reproduction and the presence of ocelli and furcula. These 199 observations concerning mainly surface Collembola suggest that wind is the principal medium 200 for dispersion onto green roofs. Productive roofs also housed Collembola without ocelli and 201 furcula, which is in agreement with dispersion through organic matter supply. The 202 configuration of green roofs-fitted buildings probably act as vertical barriers and makes active dispersion impossible. 203

204 Some results suggest that passive dispersion through human activities on a larger scale also 205 exists. An exotic species of Collembola, Coecobrya tenebricosa, has been collected in a 206 vivarium in Poland after being imported as food for frogs (Jaskuła et al., 2019). It is 207 reasonable to think that abiotic conditions would not enable this species to thrive and spread in a territory so different of its native environment. Nevertheless acclimation to particular 208 209 habitats within the area of introduction might be possible. *Paranurophorus simplex* is thought 210 to be an Asian species but was found in hothouses in Paris (Delamare Deboutteville, 1947) 211 and hothouses and flower pots in Poland (Pomorski and Skarzynski, 1996) and Ukraine 212 (Shrubovych, 2002). P. simplex could have been imported through exotic plants collections 213 and thrived in the particular environment of hothouses. This dispersion way has been 214 highlighted by Rumble et al. (2018) with Collembola species brought on a green roof 215 substrate via *Sedum spp* plugs. Despite a lack of knowledge in the original distribution range

of many Collembola species (Hopkin, 1997), those insights suggest that global human transferactivities may contribute to the large-scale dispersion of Collembola.

218

219 Sterzyńska et al. (2018) supported the idea that urban matrix can be an efficient barrier for dispersion of Collembola as they observed less species in urban than natural habitats. 220 221 However, some publications demonstrated the success of Collembola to disperse in cities as 222 they are able to colonise constructed soils (Burrow et al., 2018; Santorufo et al., 2021). If the 223 distance between the urban habitat and the nearest large habitat area is short, wind dispersion 224 might be a more likely and frequent event (Querner et al., 2018). Dispersion processes may 225 compensate the negative effects of intense anthropisation (Rzeszowski, 2017); the presence of 226 hedgerows enhances abundance and diversity during colonisation process (Burrow et al., 227 2018); close environment could play a role of source of Collembola for urban habitats (Braaker et al., 2014; Schindler et al., 2019). Yet road network seems to decrease Collembola 228 229 abundance (Eitminaviciute, 2006) and close building or canopy cover are thought to modify 230 species composition (Ge et al., 2019; Milano et al., 2017).

Investigation about other means of dispersion like phoresy (transport of one organism by another) or eggs transport need to be conduct and quantify the main source of collembolan in urban habitat. Most of dispersal processes are stochastic passive processes and will depend of urban habitat design and management (practices, high ground site, ...). In order to get to a discerning comprehension of colonisation in urban Collembola communities, much work has to be done.

A greater diversity of habitats such as parks, urban forests, domestic gardens, lawns and their spatial arrangement in cities need to be considered to understand their role in the connectivity and colonisation of soil organisms in the urban environment. Hence more works are necessary to reveal patterns of assemblages, notably in heterogeneous habitats such as cities where Collembola may display various responses to habitat fragmentation (Rantalainen et al., 2008; Sousa et al., 2004) and land use history (Ponge et al., 2006). Finally, genetic
tools, that have been shown efficient to demonstrate the effect of landscape on connectivity
between soil populations (Dupont et al., 2017), should be considered for urban Collembola
communities studies even if these methods have yet to be refined for Collembola group.

246 Concerning the colonisation, in the corpus papers, a serious lack of data dealing with the 247 temporal dynamics of Collembola colonisation in urban habitats was noticed. As studies were 248 not conducted over several years, whether colonisation onto green roofs is continuous over 249 time is not known yet. Communities are thought to change, specialists replacing ubiquists 250 (Schrader and Böning, 2006) but they also could gone extinct as reduced flow of immigrants 251 due to urban constraints and can lead to a decrease in abundance and diversity (Rumble et al., 252 2018). These evolutions are observed in the 4-years experiments of colonisation in 253 constructed Technosol (Santorufo et al., 2021).

- 254
- 255

256 4. Urban habitats and their characteristics

257

258 According to theoretical ecology, after a pool of species reaches a place through 259 dispersal, habitat is the second filter that explains communities assemblages (Drake, 1990; 260 Zobel, 1997). Several environmental factors can impact this filtering in soil communities: e.g. 261 habitat type, soil local properties(Giller, 1996; Nielsen et al., 2010). Environmental conditions 262 of each urban habitat could explain selection and survival of Collembola community. The effect of urban habitat on Collembola communities was by far the most studied factor among 263 264 the corpus since only 6 articles did not include it in their reflections. Investigated sites within 265 all studies (n=47) have been classified according to their function in the city (Tab.1). Although 12 on 15 different categories of habitats have been investigated among all papers, 266 267 only 10 have been so in more than or equal to 3 different articles. It shows how scarce knowledge about many urban habitats is, with a great disparity on the habitats studied. Indeed,
several "natural-like" habitats such as urban forests (52%) and parks (35%) are quite well
documented but with a wide diversity in each of the categories (surface, plant species...) and
few, even no study, focus on very anthropised sites like cemeteries or public (institutional,
educational) grounds.

Table 1: Distribution of habitats present in studies dealing with habitats as a filter of urban Collembola communities according to their function in cities

n°	urban land uses	%
1	Parks & gardens: country, urban and local parks, public gardens	35
2	street sides, rails, roads	23
3	Institutional grounds	0
4	Built-up areas residential land	23
5	Industrial and commercial site	8
6	educational establishments	0
7	Outdoor sports facilities : golfcourses, tennis, sport fields	2
8	Churchyards, cemeteries, burial grounds	0
9	Brownfield land: quarries, wastelands, landfills, vacant and derelict land	23
10	Community green spaces: allotments, community gardens, urban farms	15
11	Horticultural establishments	4
12	Natural & semi-natural green spaces: woodlands, forests, reserves, heathlands, grassland, meadow, conservation land	52
13	Avenue trees	4
14	Agricultural land, farms, ranches	6
15	green roofs and vertical forms	13

273

Moreover, it was very difficult to classify certain studies because the name mentioned in the publication could correspond to several categories and no reference on the vegetation or the use of the site for example was precised. Among the problematic terms, lawn (brownfield or park?), garden (allotment or domestic garden?) or park (square or a residual forest?), were especially identified.

The lack of consensus on habitat typology, even for the better documented habitats, 279 280 attempting to bring out general patterns of Collembola communities structural in response to 281 habitats turned out to be complicated. In addition, a variety of indices are used (e.g. species 282 diversity, families, QBS index, ratio) and not always comparable. Moreover, many publications do not give their raw values and rarely make taxonomic identifications on 283 springtails. Values of density and species richness were specified in 26 and 24 studies 284 respectively. Six different habitats, with their names provided by the authors, were 285 286 characterized by at least two values by index and are presented in Tab. 2 to highlight the variability of the indices. The lowest and highest densities of Collembola were both found in 287 lawns with 50 and over three millions ind/m^2 respectively (Eitminaviciute, 2006; Natuhara et 288 al., 1994), but all habitats display large differences in minimal and maximal densities. 289

Table 2: Range of values of density, mean richness (mean number of species among sites of the study) and total richness (total number of species among all sites of the study) recorded in literature. N is the number of studies providing indices values. a:Eitminaviciute 2006, b:Natuhara et al., 1994, c:Shrubovych 2002, d:Milano et al., 2017, e: Fountain & Hopkin 2004, f:Huang et al., 2013, h:Rumble et al., 2018,i:Schrader & Böning 2006, j:Santorufo et al., 2015, k:Vincent et al., 2018, l:Joimel et al., 2017, m:Joimel et al., 2019,n: Sterzynska 1990, o:Joimel 2018, p:Devigne et al., 2016, q: Rzeszowski et al., 2017a, r: Rzeszowski et al., 2017b, s:Santorufo et al., 2014, t:Sterzynska 1987

Habitat	Density $(.10^3)$			Mean richness			Total richness		
	Min	Max	N	Min	Max	Ν	Min	Max	Ν
Lawns	0,05 ^a	3 398 ^b	9	2 ^b	32 ⁿ	5	33 ^q	64 ^r	7
Parks	0,331 ^c	36,5 ^d	8	7^{d}	37 ⁿ	2	13 ^s	64 ^t	9
Urban forests	4,95 ^e	268 ^f	6	3 ^b	21 ^f	5	21 ^e	70 ^c	3
Green roofs	11 ^h	88 ⁱ	4	5 ^h	14 [°]	3	30 ⁱ	44°	2
Industrial	2,5 ^j	91 ^k	3	3 ^k	10 ^p	2	17 ^j	33 ^k	4
Gardens	7^1	73 ^m	3	6 ^m	33 ¹	3	28 ^m	41 ¹	4

290

A high variability was also observed between replicates within a same site: density can vary from none to hundred thousand individuals per square meter (Milano et al., 2017; Rumble and Gange, 2013). Specific richness is also highly variable between and within habitats, ranging from only two species in a lawn (Natuhara et al., 1994) to 37 in a park (Sterzynska, 1990). When total richness is considered, disparity again occurs. Less than 20 overall species were sampled in parks (Santorufo et al., 2014a) and in industrial sites

(Santorufo et al., 2015) and up to 70 overall species in urban forests (Shrubovych, 2002). 297 298 However it is obvious that urban sites can shelter enormous amounts of individuals (Huang et 299 al., 2013; Joimel et al., 2019; Natuhara et al., 1994). Besides species richness of urban lawns, 300 parks and forests can reach similar levels as non-anthropogenic sites (Joimel et al., 2017; 301 Ponge et al., 2003). This concurs with findings arguing that habitat type in urban area does not 302 always influence abundance or species richness, emphasising that biogeographic processes may be of greater significance (Joimel et al., 2018; Sterzyńska et al., 2018). But these 303 304 conclusions need to be nuanced because i) as the classification is coarse, each category may 305 group sites with different structural characteristics and ii) in addition sampling methods and 306 season or habitat size vary from one study to another one and can impact results and make 307 comparisons between studies difficult (Querner and Bruckner, 2010; Santorufo et al., 2014b; 308 Yi et al., 2012). Thus provided figures (Tab.2) only broadly inform on Collembola community 309 indices adopted in urban habitats.

310 So if disparities between habitats exist, no general pattern concerning abundance and 311 richness seems to emerge. And the local characteristics might be more relevant to explain 312 Collembola urban assemblages (Sterzynska, 1987). Habitat structure can be spatially and temporally defined (Byrne, 2007). Water content, pH, organic matter or nutrients content, 313 314 metal concentrations and presence or absence of a litter layer are the soil characteristics that 315 have been the most studied in relation to their effects on Collembola communities and they 316 are important drivers regarding their assemblages. Organic matter and water content - except 317 for anaerobic conditions - are almost always positively correlated with abundance, richness 318 and composition of Collembola (Ge et al., 2019; Islam et al., 2018; Milano et al., 2018, 2017; 319 Santorufo et al., 2015, 2014a). Soil pH and nutrients such as phosphorus and potassium seem rather influence communities composition (Milano et al., 2018, 2017; Rzeszowski et al., 320 321 2017). Soil concentrations of potassium and phosphorus can for instance be positively related

with the presence of *Pseudosinella alba* (Packard, 1873) and *Stenaphorura quadrispina* 322 (Börner, 1901) respectively and negatively with other species such as Entomobrya nicoleti 323 324 (Lubbock, 1868) or Isotoma anglicana (Lubbock, 1862; Rzeszowski et al., 2017). Metal concentrations can have different types of impact on Collembola (Fountain and Hopkin, 2004; 325 326 Joimel et al., 2019; Vincent et al., 2018). Some studies showed that the presence of metals 327 even at high concentrations could not affect widespread species such as *P. minuta* (Milano et al., 2017; Rota et al., 2015; Santorufo et al., 2015; Sterzyńska et al., 2018). However, high 328 329 metal concentrations could reduce Collembola abundance (Santorufo et al., 2014a, 2014b, 330 2012) or survival and reproduction (Fountain and Hopkin, 2004). Urban management practices could also play a role in structuring Collembola communities. The presence of a 331 litter layer increases Collembola indices such as abundances or number of species (Milano et 332 al., 2017; Natuhara et al., 1994; Santorufo et al., 2014a). Too little work addressed the effect 333 334 of other management practices such as pesticide application or mowing to draw conclusions, 335 but they could also have an impact (Byrne and Bruns, 2004; Jensen et al., 1973; Rochefort et 336 al., 2013). Furthermore, habitat characteristics may evolve with time and these effects on 337 Collembola communities remain unknown. Pioneer studies showed that site history (i.e age and previous land use) and long-term exposure to urbanisation may affect species community 338 and the number of Collembola species (Milano et al., 2017; Rzeszowski and Sterzyńska, 339 340 2016; Schrader and Böning, 2006).

341

342 5. Biotic interactions between Collembola and between Collembola and other soil
 343 organisms

344

Biotic interactions can be understood at different scales, between Collembola themselves and between Collembola and other soil organisms, but in all cases they have been little studied in urban areas.

348 Concerning Collembola interspecific relations, while the dominance of certain species 349 among communities has been evidenced (Shrubovych, 2002; Sterzynska, 1987, 1987), 350 whether the presence of species can exclude or favour other ones is still obscure. But Collembola are known to secrete pheromones involved in aggregation of individuals, either 351 352 inter or intra specific (Salmon et al., 2019), gathering up to millions of individuals. Their 353 interactions are proved to possibly be detrimental or beneficial depending on the species (Theenhaus et al., 1999). Interactions between Collembola are complex and can vary 354 355 according to habitat (Christiansen et al., 1992), which raises interrogation to what extent 356 urban habitat affects those interactions and whether it induces a shift in community 357 composition.

358 Although half of publications of the selected corpus used a multi-taxa approach, only a 359 few addressed questions of trophic relations, competition, facilitation, or co-occurrence. 360 When several taxa were studied simultaneously, it was generally to evaluate the effect of an 361 environmental parameter on the overall soil fauna community (McIntyre 2000, Maisto 2017, Mo 2018, Ge 2019). It is well established that Collembola interact with numerous other taxa 362 363 in different types of ways and habitats (Hopkins 1997). Many groups of the micro-flora and 364 fauna are part of their various diets, such as fungi, algae, bacteria and more rarely other 365 invertebrates (nematodes, rotifers and tardigrades). Collembola therefore play a crucial role in 366 soil communities structure, for instance by selectively eating certain fungi species or 367 regulating density of phytopathogens (Wardle 2006). Only two studies integrated both urban 368 Collembola and fungi in their investigations. Rumble and Gange (2013) found no correlation 369 between Collembola and arbuscular mycorrhizal fungi. It reinforces the hypothesis of a selective diet for non-arbuscular fungi (Gange 2000). Moreover, Joimel et al. (2018) showed
that lead contamination can modify fungal communities. It could then have an impact on
Collembola diet, especially in urban habitats where contaminations are widespread.
Collembola are also preyed on by soil predators including spiders, acarian, myriapodes or
even some vertebrates (Thibaud 2010). Predator-prey interactions in soil have been shown to
display functional responses influenced by habitat structure and prey aggregation (VucicPestic 2010). However no work has been conducted in urban habitats for now.

377 Interactions between earthworms and Collembola have been the object of more attention, but again very rarely in urban soils. Contrasted results were observed. Some 378 379 Collembola species seem to be preferentially present in soils richer in earthworms 380 (Wickenbrock 1997, Salmon 1999). An explanation could be that they feed on earthworm's excrements and use their burrows to move and hide (Salmon 2004). Earthworms can impact 381 382 density, richness and species succession of Collembola communities (Dunger 2004, Mudrak 383 2012). Interactions between these two groups of soil organisms appear to depend on 384 community composition and how it impacts ecosystem processes is still to be discovered 385 (Grubert 2016). But some trade-offs may also exist. Schrader and Boenig (2006) think that the absence of earthworms might favor the high densities of Collembola in green roofs substrates 386 387 where competition with earthworms does not exist. Joimel et al. (2018) also observed a higher 388 density of Collembola in Parisian urban farms soils where earthworms were absent.

Plants also appear to play a major role in the structure of Collembola communities. Generally, Collembola diversity is positively linked with above plants diversity in cities (Burrow 2017, Tresch 2019). As plants modify local soil properties (Qin 2019), it is possible that a diversity of plants may offer for Collembola a variety of microhabitats or food resources (e.g. by effect on microorganisms), explaining their higher abundance and richness in more complex environments. Moreover, as mentioned above, when looking at sprout plugs

meant to be planted, differences can be observed in the Collembola species present in the plug 395 396 in function of the plant species (Rumble 2018). In return Collembola may provide nutrients to 397 plants with their excrements, their role in organic matter decomposition and improve their 398 growth through mycorrhizal stimulation (Gange 2000). These insights support the idea that 399 multiple feedback interactions exist between Collembola and plants, stressing the necessity to 400 incorporate reflections about selected plants in urban landscapes in order to favour long-term 401 viable soil communities. It should even be considered with great care as cities are hotspots for 402 the introduction of alien species (Cadotte 2017, Jäkel 2005) that could threaten local 403 communities by changing soil characteristics (Qin 2019), or introducing predators and competitors (Jaskula 2019, Rowles 2019). Again, interactions between Collembola and 404 405 earthworms, plants and other soil dweller organisms deserve more interest, particularly in 406 constructed soils where their synergy could be very important in functional processes. These 407 biotic interactions can rapidly evolve and change biological interactions between Collembola 408 and the other organisms of macrofauna, especially in environments where dispersal can only 409 be achieved through voluntary inputs (e.g. compost for earthworms on roofs).

410

412

411 6. Synthesis and future research

This study, reviewing main data available about how urbanisation affects Collembola used as bioindicators, has provided an initial overview (descriptive study) but it shows that, although work has been developing increasingly in recent years, there is still a serious lack of data to understand, explain and prioritise (i) the processes controlling the dispersion of Collembola in urban soils and (ii) the relationships between Collembola and habitat characteristics on the one hand (impact of abiotic factors) and Collembola and other soil organisms on the other (impact of biotic factors). Thus, in order to reconcile human activities and ecosystem functionality and to contribute answering to the Sustainable Development Goal 11, "creating a more sustainable future for cities and other urban communities" (FAO, ITPS, GSBI, CBD and EC, 2020), we suggest some proposals for future research in urban soil biodiversity.

424 Concerning the methodology used, three points have been noticed: (i) the lack of standardisation for sampling, collecting and extraction of Collembola; there is no standardized 425 426 methods and strategy making it difficult inter- and intra-habitat comparisons and creating 427 references values. We encourage following international standard ISO 23611-2 on sampling 428 and extraction of microarthropods with soil cores; (ii) No clear classification of urban 429 habitats; indeed confusions always reside between their type (e.g. forest, lawn), function (e.g. 430 recreational, ornamental, productive), spatial arrangement (proximity of roads, building 431 density) and their management (mowing, litter removal, chemical inputs). Moreover, behind 432 the same land use can hide a variety of soil types. As there is no common classification, we 433 propose to always specify the reference terminology used to classify the habitat. More details 434 on the different cited parameters could be also useful for future meta-analyses. These 435 parameters needs to include the presence of irrigation, connection to other soils or not (i.e. height, above ground), soil depth and regular inputs of materials (e.g. compost) which directly 436 imply assumptions in main dispersion routes, selection and establishment processes of 437 438 Collembola community in urban habitat (iii) Density is not sufficient for highlighting 439 mechanisms underlying the assembly of communities. Species identification, which is only 440 present in less than half studies, is also necessary to better understand processes relatives to 441 homogeneisation and invasions by exotic species recorded in few studies. Moreover, functional traits seem to be more widely complementary to allow comparison between various 442 443 cities, obtain a better global comprehension of assemblage processes, and soil functioning. 444 These functional approaches can be facilitated by the existence of many open databases that

445 provide data not only on Collembola but also on other soil fauna organisms (e.g. BETSI, Pey 446 et al. 2014). Direct measurements of functional traits on organisms can also be useful in 447 regions of the world where the taxonomy is less well known by offering an approach that 448 allows a finer understanding without the need for specific identification.

A second point noted in this review is the limited worldwide recovery. Moreover, an international research in cities of varying sizes is necessary. In addition to new studies, the diffusion of data published in the authors' native language to a wider audience could lead to a rapid increase in global knowledge. Indeed, many data seems to be available in China and Russia but the majority of non-English publications do not allow drawing conclusions.

454

Finally, concerning knowledge acquired on urban Collembola, studies are recent in 455 456 lines with all studies on urban soils, and make it possible to identify many areas of research to 457 be carried out: (i) increase the number of studies between the different habitat categories and 458 within each category ; (ii) Take into account management practices;; iii) Take into account 459 local conditions such as environmental conditions and soil properties (e.g. moisture, organic 460 matter, nutrients and metal contents) that may be responsible for the wide variability in the 461 structural responses of Collembola communities to the effect of habitat; (iv) Integrating 462 temporal evolutions; to understand the role of colonisation of urban habitats. Nowadays many 463 hypotheses are proposed, but none was verified. Specific studies must therefore be built in order to understand precisely the mechanics of colonisation and dispersion of soil organisms 464 465 in cities; (v) Consider biotic interactions to better understand the functioning of urban soils 466 and be able to manage them in the long term; (vi) Take into account spatial organisation of 467 cities; Contemporary urbanization differs from former urbanization (Ramalho and Hobbs, 468 2012). Cities are larger, more rapidly urbanized, and more irregularly shaped (Batty, 2008; 469 Liu et al., 2014). Many questions therefore arise about the impact of the spatial organisation within cities on the dissemination of soils organisms in cities. In order to answer these 470

- 471 questions, it seems necessary to set up in situ studies with varying soil types, heights,
- 472 obstacles or connectivity in order to monitor taxonomic and functional biodiversity indicators
- 473 over time: from installation to the eventual stabilisation of the community.

474

475

482

476 **Declarations**

477 Funding

478 This work was supported by the lab recherche environnement VINCI ParisTech.

479 Authors' contributions

480 **SJ**: idea for the article. **AJ**, **SJ**: literature search and data analysis. **AJ**, **SJ**: drafted the 481 manuscript. **LVG** : critically revised the work.

483 References

- André, H.M., Noti, M.-I., Lebrun, P., 1994. The soil fauna: the other last biotic frontier.
 Biodivers Conserv 3, 45–56. https://doi.org/10.1007/BF00115332
- Auclerc, A., Ponge, J.F., Barot, S., Dubs, F., 2009. Experimental assessment of habitat
 preference and dispersal ability of soil springtails. Soil Biology and Biochemistry 41,
 1596–1604.
- Barot, S., Abbadie, L., Auclerc, A., Barthélémy, C., Bérille, E., Billet, P., Clergeau, P., Consales,
 J.-N., Deschamp-Cottin, M., David, A., Devigne, C., Dham, V., Dusza, Y., Gaillard, A.,
- 491 Gonzalez, E., Hédont, M., Labarraque, D., Le Bastard, A.-M., Morel, J.-L., Petit-
- Berghem, Y., Rémy, E., Rochelle-Newall, E., Veyrières, M., 2019. Urban ecology,
 stakeholders and the future of ecology. Science of The Total Environment 667, 475–
 484. https://doi.org/10.1016/j.scitotenv.2019.02.410
- Biffi, S., De Souza, C.M., Firbank, L.G., 2019. Epigeal fauna of urban food production sites
 show no obvious relationships with soil characteristics or site area. Agriculture
 Ecosystems & Environment 286, UNSP 106677.
- 498 https://doi.org/10.1016/j.agee.2019.106677
- Braaker, S., Ghazoul, J., Obrist, M.K., Moretti, M., 2014. Habitat connectivity shapes urban
 arthropod communities: the key role of green roofs. Ecology 95, 1010–1021.
 https://doi.org/10.1890/13-0705.1
- Burrow, C., Pernin, C., Lepretre, A., 2018. Influence of connectivity & topsoil management
 practices of a constructed technosol on pedofauna colonization: A field study.
 Applied Soil Ecology, HUMUSICA 3 Reviews, Applications, Tools 123, 416–419.
 https://doi.org/10.1016/j.apsoil.2017.12.001
- 506 Byrne, L.B., 2007. Habitat structure: A fundamental concept and framework for urban soil 507 ecology. Urban Ecosyst 10, 255–274. https://doi.org/10.1007/s11252-007-0027-6
- Byrne, L.B., Bruns, M.A., 2004. The Effects of Lawn Management on Soil Microarthropods.
 Journal of Agricultural and Urban Entomology 150–156.
- 510 Christiansen, K., Doyle, M., Kahlert, M., Gobaleza, D., 1992. Interspecific interactions
 511 between collembolan populations in culture. Pedobiologia.

512 Clobert, J., Baguette, M., Benton, T.G., Bullock, J.M. (Eds.), 2012. Dispersal Ecology and 513 Evolution. Oxford University Press, Oxford, New York. 514 Culik, M.P., Filho, D.Z., 2003. Diversity and distribution of Collembola (Arthropoda: 515 Hexapoda) of Brazil. Biodiversity and Conservation 12, 1119–1143. https://doi.org/10.1023/A:1023069912619 516 de Groot, R.S., Wilson, M.A., Boumans, R.M.J., 2002. A typology for the classification, 517 518 description and valuation of ecosystem functions, goods and services. Ecological 519 Economics 41, 393-408. https://doi.org/10.1016/S0921-8009(02)00089-7 520 Decaëns, T., Jiménez, J.J., Gioia, C., Measey, G.J., Lavelle, P., 2006. The values of soil animals 521 for conservation biology. European Journal of Soil Biology 42, Supplement 1, S23-522 S38. https://doi.org/10.1016/j.ejsobi.2006.07.001 523 Delamare Deboutteville, C., 1947. Note faunistique sur les Collemboles de France. Revue 524 française d'Entomologie 125–138. 525 Drake, J.A., 1990. Communities as assembled structures: Do rules govern pattern? Trends in 526 Ecology & Evolution 5, 159–164. https://doi.org/10.1016/0169-5347(90)90223-Z 527 Dunger, W., 1998. Immigration, Ansiedlung und Primärsukzession der Bodenfauna auf 528 jungen Kippböden, in: Pflug, W. (Ed.), Braunkohlentagebau und Rekultivierung: 529 Landschaftsökologie — Folgenutzung — Naturschutz. Springer, Berlin, Heidelberg, 530 pp. 635-644. https://doi.org/10.1007/978-3-642-58846-4_53 531 Dunger, W., Schulz, H.-J., Zimdars, B., 2002. Colonization behaviour of Collembola under 532 different conditions of dispersal: Proceedings of the Xth international Colloquium on 533 Apterygota, České Budějovice 2000: Apterygota at the Beginning of the Third 534 Millennium. Pedobiologia 46, 316–327. https://doi.org/10.1078/0031-4056-00139 535 Dupont, L., Torres-Leguizamon, M., René-Corail, P., Mathieu, J., 2017. Landscape features 536 impact connectivity between soil populations: a comparative study of gene flow in 537 earthworms. Molecular Ecology 26, 3128–3140. https://doi.org/10.1111/mec.14102 538 Eitminaviciute, I., 2006. Microarthropod communities in anthropogenic urban soils. 2. 539 Seasonal dynamics of microarthropod number in crossing curb soils. Zoologichesky 540 Zhurnal 1309–1320. 541 Fountain, M.T., Hopkin, S.P., 2004. A Comparative Study of the Effects of Metal 542 Contamination on Collembola in the Field and in the Laboratory. Ecotoxicology 13, 543 573-587. https://doi.org/10.1023/B:ECTX.0000037194.66321.2c 544 Ge, B., Mehring, A.S., Levin, L.A., 2019. Urbanization alters belowground invertebrate 545 community structure in semi-arid regions: A comparison of lawns, biofilters and sage 546 scrub. Landscape and Urban Planning 192, 103664. 547 https://doi.org/10.1016/j.landurbplan.2019.103664 548 Giller, P.S., 1996. The diversity of soil communities, the 'poor man's tropical rainforest.' 549 Biodivers Conserv 5, 135–168. https://doi.org/10.1007/BF00055827 550 Guilland, C., Maron, P.A., Damas, O., Ranjard, L., 2018. Biodiversity of urban soils for 551 sustainable cities. Environmental Chemistry Letters. https://doi.org/10.1007/s10311-552 018-0751-6 553 Hopkin, S.P., 1997. Biology of the Springtails : (Insecta: Collembola): (Insecta: Collembola). 554 Oxford University Press. 555 Huang, C., Wang, T., Luo, Y., Chen, S., Kim, K.C., 2013. Assessing Collembola Biodiversity 556 Under Human Influence at Three Gorges Area, China. Environ Entomol 42, 214–222. 557 https://doi.org/10.1603/EN12253

- 558 Islam, M.S., Hossain, M., Hossain, M., Yasmin, M., 2018. Impact of climatic factors and soil 559 quality on the abundance and population density of Collembola in the Rajshahi 560 University Campus, Bangladesh. Journal of Entomology and Zoology Studies 6, 1119-561 1125.
- 562 Janion-Scheepers, C., Deharveng, L., Bedos, A., Chown, S.L., 2015. Updated list of Collembola 563 species currently recorded from South Africa. ZooKeys 55. 564 https://doi.org/10.3897/zookeys.503.8966
- 565 Jaskuła, R., Sulikowska-Drozd, A., Jabłońska, A., Banaś, K., Rewicz, T., 2019. Undesirable 566 immigrants: hobbyist vivaria as a potential source of alien invertebrate species. PeerJ 567 7, e7617. https://doi.org/10.7717/peerj.7617
- Jensen, P., Jacobson, G.L., Willard, D.E., 1973. Effects of Mowing and Raking on Collembola. 568 Ecology 54, 564–572. https://doi.org/10.2307/1935341 569
- 570 Joimel, S., Baptiste Grard, Apolline Auclerc, Mickaël Hedde, Nolwenn Le Doaré, Sandrine 571 Salmon, Claire Chenu, 2018. Are Collembola "flying" onto green roofs? Ecological 572 Engineering 111, 117–124.
- 573 Joimel, S., Cortet, J., Jolivet, C.C., Saby, N.P.A., Chenot, E.D., Branchu, P., Consalès, J.N., 574 Lefort, C., Morel, J.L., Schwartz, C., 2016. Physico-chemical characteristics of topsoil 575 for contrasted forest, agricultural, urban and industrial land uses in France. Science of 576 The Total Environment 545–546, 40–47.
- 577 https://doi.org/10.1016/j.scitotenv.2015.12.035
- 578 Joimel, S., Schwartz, C., Hedde, M., Kiyota, S., Krogh, P.H., Nahmani, J., Pérès, G., Vergnes, A., 579 Cortet, J., 2017. Urban and industrial land uses have a higher soil biological quality 580 than expected from physicochemical quality. Science of The Total Environment 584-581 585, 614–621. https://doi.org/10.1016/j.scitotenv.2017.01.086
- 582 Joimel, S., Schwartz, C., Maurel, N., Magnus, B., Machon, N., Bel, J., Cortet, J., 2019. 583 Contrasting homogenization patterns of plant and collembolan communities in urban 584 vegetable gardens. Urban Ecosystems. https://doi.org/10.1007/s11252-019-00843-z
- 585 Lang, A., 2000. The pitfalls of pitfalls: a comparison of pitfall trap catches and absolute 586 density estimates of epigeal invertebrate predators in Arable Land. J. Pest Science 73, 587 99–106. https://doi.org/10.1007/BF02956438
- 588 Liu, Z., He, C., Zhou, Y., Wu, J., 2014. How much of the world's land has been urbanized, 589 really? A hierarchical framework for avoiding confusion. Landscape Ecology 29, 763-590 771. https://doi.org/10.1007/s10980-014-0034-y
- 591 Lortie, C.J., Brooker, R.W., Choler, P., Kikvidze, Z., Michalet, R., Pugnaire, F.I., Callaway, R.M., 592 2004. Rethinking plant community theory. Oikos 107, 433–438.
- 593 https://doi.org/10.1111/j.0030-1299.2004.13250.x
- Marzluff, J.M., 2019. A decadal review of urban ornithology and a prospectus for the future. 594 595 Ibis 1-13. https://doi.org/10.1111/ibi.12430@10.1111/(ISSN)1474-596

919X.highly cited us research

- 597 McKinney, M.L., 2008. Effects of urbanization on species richness: A review of plants and 598 animals. Urban Ecosystems 11, 161–176. https://doi.org/10.1007/s11252-007-0045-4
- 599 Milano, V., Cortet, J., Baldantoni, D., Bellino, A., Dubs, F., Nahmani, J., Strumia, S., Maisto, G., 600 2017. Collembolan biodiversity in Mediterranean urban parks: impact of history, 601 urbanization, management and soil characteristics. Applied Soil Ecology 119, 428-602 437. https://doi.org/10.1016/j.apsoil.2017.03.022
- 603 Milano, V., Maisto, G., Baldantoni, D., Bellino, A., Bernard, C., Croce, A., Dubs, F., Strumia, S., 604 Cortet, J., 2018. The effect of urban park landscapes on soil Collembola diversity: A

605 Mediterranean case study. Landscape and Urban Planning 180, 135–147. 606 https://doi.org/10.1016/j.landurbplan.2018.08.008 607 Morel, J.L., Chenu, C., Lorenz, K., 2014. Ecosystem services provided by soils of urban, 608 industrial, traffic, mining, and military areas (SUITMAs). J Soils Sediments 1–8. 609 https://doi.org/10.1007/s11368-014-0926-0 Natuhara, Y., Imai, C., Takeda, H., 1994. Classification and ordination of communities of soil 610 611 arthropods in an urban park of Osaka City. Ecological Research 9, 131–141. 612 https://doi.org/10.1007/BF02347488 613 Nielsen, U.N., Osler, G.H.R., Campbell, C.D., Burslem, D.F.R.P., Wal, R. van der, 2010. The 614 influence of vegetation type, soil properties and precipitation on the composition of 615 soil mite and microbial communities at the landscape scale. Journal of Biogeography 37, 1317–1328. https://doi.org/10.1111/j.1365-2699.2010.02281.x 616 617 Pomorski, R.J., Skarzynski, D.S., 1996. (Collembola) pól uprawnych, ziemi doniczrjwej, 618 szklarni i pieczarkarni – klucz do oznaczania // Diagnostyka czkodników roslin i ich 619 wrogów naturalnych. 620 Ponge, J.-F., 2013. Disturbances, organisms and ecosystems: a global change perspective. 621 Ecol. Evol. 3, 1113-1124. https://doi.org/10.1002/ece3.505 622 Ponge, J.-F., Dubs, F., Gillet, S., Sousa, J.P., Lavelle, P., 2006. Decreased biodiversity in soil 623 springtail communities: the importance of dispersal and landuse history in 624 heterogeneous landscapes. Soil Biology and Biochemistry 38, 1158–1161. 625 https://doi.org/10.1016/j.soilbio.2005.09.004 626 Ponge, J.-F., Gillet, S., Dubs, F., Fedoroff, E., Haese, L., Sousa, J.P., Lavelle, P., 2003. 627 Collembolan communities as bioindicators of land use intensification. Soil Biology 628 and Biochemistry 35, 813-826. https://doi.org/10.1016/S0038-0717(03)00108-1 629 Querner, P., Bruckner, A., 2010. Combining pitfall traps and soil samples to collect 630 Collembola for site scale biodiversity assessments. Applied Soil Ecology 45, 293–297. 631 https://doi.org/10.1016/j.apsoil.2010.05.005 632 Querner, P., Milasowszky, N., Zulka, K.P., Abensperg-Traun, M., Willner, W., Sauberer, N., 633 Jakomini, C., Wrbka, T., Schmitzberger, I., Zechmeister, H.G., 2018. Habitat Structure, 634 Quality and Landscape Predict Species Richness and Communities of Collembola in 635 Dry Grasslands in Austria. Insects 9, 81. https://doi.org/10.3390/insects9030081 636 Rantalainen, M.-L., Haimi, J., Fritze, H., Pennanen, T., Setälä, H., 2008. Soil decomposer 637 community as a model system in studying the effects of habitat fragmentation and 638 habitat corridors. Soil Biology and Biochemistry 40, 853–863. 639 https://doi.org/10.1016/j.soilbio.2007.11.008 640 Rochefort, S., Shetlar, D.J., Brodeur, J., 2013. Impact of four turf management regimes on 641 arthropod abundance in lawns. Pest Management Science 69, 54–65. 642 https://doi.org/10.1002/ps.3361 643 Rota, E., Caruso, T., Migliorini, M., Monaci, F., Agamennone, V., Biagini, G., Bargagli, R., 2015. 644 Diversity and abundance of soil arthropods in urban and suburban holm oak stands. 645 Urban Ecosyst 18, 715–728. https://doi.org/10.1007/s11252-014-0425-5 646 Rumble, H., Finch, P., Gange, A.C., 2018. Green roof soil organisms: Anthropogenic 647 assemblages or natural communities? Applied Soil Ecology 126, 11–20. 648 https://doi.org/10.1016/j.apsoil.2018.01.010 Rumble, H., Gange, A.C., 2013. Soil microarthropod community dynamics in extensive green 649 650 roofs. Ecological Engineering 57, 197-204. 651 https://doi.org/10.1016/j.ecoleng.2013.04.012

- Rusek, J., 1998. Biodiversity of Collembola and their functional role in the ecosystem.
 Biodiversity and Conservation 7, 1207–1219.
 https://doi.org/10.1023/A:1008887817883
- Rzeszowski, K., 2017. The effects of anthropogenic disturbances and hydrological activity of a
 river on soil Collembola communities in an urbanized zone. European Journal of Soil
 Biology 82, 116–120. https://doi.org/10.1016/j.ejsobi.2017.09.004
- Rzeszowski, K., Sterzyńska, M., 2016. Changes through time in soil Collembola communities
 exposed to urbanization. Urban Ecosyst 19, 143–158.
- 660 https://doi.org/10.1007/s11252-015-0478-0
- Rzeszowski, K., Zadrożny, P., Nicia, P., 2017. The effect of soil nutrient gradients on
 Collembola communities inhabiting typical urban green spaces. Pedobiologia 64, 15–
 24. https://doi.org/10.1016/j.pedobi.2017.06.003
- Salmon, S., Rebuffat, S., Prado, S., Sablier, M., D'Haese, C., Sun, J.-S., Ponge, J.-F., 2019.
 Chemical communication in springtails: a review of facts and perspectives. Biol Fertil
 Soils 55, 425–438. https://doi.org/10.1007/s00374-019-01365-8
- Santorufo, L., Cortet, J., Arena, C., Goudon, R., Rakoto, A., Morel, J.-L., Maisto, G., 2014a. An
 assessment of the influence of the urban environment on collembolan communities
 in soils using taxonomy- and trait-based approaches. Applied Soil Ecology 78, 48–56.
 https://doi.org/10.1016/j.apsoil.2014.02.008
- Santorufo, L., Cortet, J., Nahmani, J., Pernin, C., Salmon, S., Pernot, A., Morel, J.L., Maisto, G.,
 2015. Responses of functional and taxonomic collembolan community structure to
 site management in Mediterranean urban and surrounding areas. European Journal
 of Soil Biology 70, 46–57. https://doi.org/10.1016/j.ejsobi.2015.07.003
- Santorufo, L., Joimel, S., Auclerc, A., Deremiens, J., Grisard, G., Hedde, M., Nahmani, J.,
 Pernin, C., Cortet, J., 2021. Early colonization of constructed technosol by
 microarthropods. Ecological Engineering 162, 106174.
- 678 https://doi.org/10.1016/j.ecoleng.2021.106174
- Santorufo, L., Van Gestel, C.A.M., Maisto, G., 2014b. Sampling season affects conclusions on
 soil arthropod community structure responses to metal pollution in Mediterranean
 urban soils. Geoderma 226–227, 47–53.
- 682 https://doi.org/10.1016/j.geoderma.2014.02.001
- Santorufo, L., Van Gestel, C.A.M., Rocco, A., Maisto, G., 2012. Soil invertebrates as
 bioindicators of urban soil quality. Environmental Pollution 161, 57–63.
 https://doi.org/10.1016/j.envpol.2011.09.042
- Schindler, B.Y., Vasl, A., Blaustein, L., Gurevich, D., Kadas, G.J., Seifan, M., 2019. Fine-scale
 substrate heterogeneity does not affect arthropod communities on green roofs.
 PeerJ 7, e6445. https://doi.org/10.7717/peerj.6445
- Schrader, S., Böning, M., 2006. Soil formation on green roofs and its contribution to urban
 biodiversity with emphasis on Collembolans. Pedobiologia 50, 347–356.
 https://doi.org/10.1016/j.pedobi.2006.06.003
- Shochat, E., Warren, P.S., Faeth, S.H., Mcintyre, N.E., Hope, D., 2006. From patterns to
 emerging processes in mechanistic urban ecology. Trends in ecology & evolution 21,
 186–191.
- 695 Shrubovych, J., 2002. The fauna of springtails (Collembola) in Lviv. Vestnik Zoologii 36, 63–
 696 67.
- Sousa, J.P., da Gama, M.M., Pinto, C., Keating, A., Calhôa, F., Lemos, M., Castro, C., Luz, T.,
 Leitão, P., Dias, S., 2004. Effects of land-use on Collembola diversity patterns in a

699 Mediterranean landscape. Pedobiologia 48, 609–622. 700 https://doi.org/10.1016/j.pedobi.2004.06.004 Sterzynska, M., 1990. Communities of Collembola in Natural and Transformed Soils of the 701 702 Linden- Oak-Hornbeam Sites of the Mazovian Lowland Poland. Fragmenta Faunistica 703 (Warsaw) 34, 165-262. 704 Sterzynska, M., 1987. Structure of springtail (Collembola) communities in the urban green of 705 Warsaw. Memorabilia Zoologica 3–18. 706 Sterzyńska, M., Nicia, P., Zadrożny, P., Fiera, C., Shrubovych, J., Ulrich, W., 2018. Urban 707 springtail species richness decreases with increasing air pollution. Ecological 708 Indicators 94, 328–335. https://doi.org/10.1016/j.ecolind.2018.06.063 709 Sukopp, H., 2008. On the Early History of Urban Ecology in Europe, in: Marzluff, J.M., 710 Shulenberger, E., Endlicher, W., Alberti, M., Bradley, G., Ryan, C., Simon, U., 711 ZumBrunnen, C. (Eds.), Urban Ecology: An International Perspective on the 712 Interaction Between Humans and Nature. Springer US, Boston, MA, pp. 79–97. 713 https://doi.org/10.1007/978-0-387-73412-5 6 714 Theenhaus, A., Scheu, S., Schaefer, M., 1999. Contramensal interactions between two 715 collembolan species: effects on population development and on soil processes. 716 Functional Ecology 13, 238–246. https://doi.org/10.1046/j.1365-2435.1999.00303.x 717 Vincent, Q., Leyval, C., Beguiristain, T., Auclerc, A., 2018. Functional structure and 718 composition of Collembola and soil macrofauna communities depend on abiotic 719 parameters in derelict soils. Applied Soil Ecology 130, 259–270. 720 https://doi.org/10.1016/j.apsoil.2018.07.002 721 Yi, Z., Jinchao, F., Dayuan, X., Weiguo, S., Axmacher, J.C., 2012. A Comparison of Terrestrial 722 Arthropod Sampling Methods. jore 3, 174–182. https://doi.org/10.5814/j.issn.1674-723 764x.2012.02.010 724 Young, R.F., Wolf, S.A., 2006. Goal attainment in urban ecology research: A bibliometric 725 review 1975-2004. Urban Ecosyst 9, 179-193. https://doi.org/10.1007/s11252-006-726 8589-2 727 Zobel, M., 1997. The relative of species pools in determining plant species richness: an 728 alternative explanation of species coexistence? Trends in Ecology & Evolution 12, 729 266-269. https://doi.org/10.1016/S0169-5347(97)01096-3 730 731 Baguette, M., Van Dyck, H., 2007. Landscape connectivity and animal behavior: functional 732 grain as a key determinant for dispersal. Landscape Ecol 22, 1117–1129. 733 https://doi.org/10.1007/s10980-007-9108-4 734 Barot, S., Abbadie, L., Auclerc, A., Barthélémy, C., Bérille, E., Billet, P., Clergeau, P., Consales, 735 J.-N., Deschamp-Cottin, M., David, A., Devigne, C., Dham, V., Dusza, Y., Gaillard, A., 736 Gonzalez, E., Hédont, M., Labarraque, D., Le Bastard, A.-M., Morel, J.-L., Petit-737 Berghem, Y., Rémy, E., Rochelle-Newall, E., Veyrières, M., 2019. Urban ecology, 738 stakeholders and the future of ecology. Science of The Total Environment 667, 475-739 484. https://doi.org/10.1016/j.scitotenv.2019.02.410 740 Beninde, J., Veith, M., Hochkirch, A., 2015. Biodiversity in cities needs space: a meta-analysis 741 of factors determining intra-urban biodiversity variation. Ecology Letters 18, 581-742 592. https://doi.org/10.1111/ele.12427 743 Biffi, S., De Souza, C.M., Firbank, L.G., 2019. Epigeal fauna of urban food production sites 744 show no obvious relationships with soil characteristics or site area. Agriculture

745 Ecosystems & Environment 286, UNSP 106677. 746 https://doi.org/10.1016/j.agee.2019.106677 747 Braaker, S., Ghazoul, J., Obrist, M.K., Moretti, M., 2014. Habitat connectivity shapes urban 748 arthropod communities: the key role of green roofs. Ecology 95, 1010–1021. 749 https://doi.org/10.1890/13-0705.1 750 Burrow, C., Pernin, C., Lepretre, A., 2018. Influence of connectivity & topsoil management 751 practices of a constructed technosol on pedofauna colonization: A field study. 752 Applied Soil Ecology, HUMUSICA 3 - Reviews, Applications, Tools 123, 416–419. 753 https://doi.org/10.1016/j.apsoil.2017.12.001 754 Byrne, L.B., 2007. Habitat structure: A fundamental concept and framework for urban soil 755 ecology. Urban Ecosyst 10, 255–274. https://doi.org/10.1007/s11252-007-0027-6 756 Byrne, L.B., Bruns, M.A., 2004. The Effects of Lawn Management on Soil Microarthropods. 757 Journal of Agricultural and Urban Entomology 150–156. 758 Cadotte, M.W., 2006. Dispersal and Species Diversity: A Meta-Analysis. The American 759 Naturalist 167, 913–924. https://doi.org/10.1086/504850 760 Christiansen, K., Doyle, M., Kahlert, M., Gobaleza, D., 1992. Interspecific interactions 761 between collembolan populations in culture. Pedobiologia. 762 Clobert, J., Baguette, M., Benton, T.G., Bullock, J.M. (Eds.), 2012. Dispersal Ecology and 763 Evolution. Oxford University Press, Oxford, New York. 764 Culik, M.P., Filho, D.Z., 2003. Diversity and distribution of Collembola (Arthropoda: 765 Hexapoda) of Brazil. Biodiversity and Conservation 12, 1119–1143. 766 https://doi.org/10.1023/A:1023069912619 767 de Groot, R.S., Wilson, M.A., Boumans, R.M.J., 2002. A typology for the classification, 768 description and valuation of ecosystem functions, goods and services. Ecological 769 Economics 41, 393-408. https://doi.org/10.1016/S0921-8009(02)00089-7 770 Decaëns, T., Jiménez, J.J., Gioia, C., Measey, G.J., Lavelle, P., 2006. The values of soil animals 771 for conservation biology. European Journal of Soil Biology 42, Supplement 1, S23-772 S38. https://doi.org/10.1016/j.ejsobi.2006.07.001 773 Delamare Deboutteville, C., 1947. Note faunistique sur les Collemboles de France. Revue 774 française d'Entomologie 125–138. 775 Drake, J.A., 1990. Communities as assembled structures: Do rules govern pattern? Trends in 776 Ecology & Evolution 5, 159–164. https://doi.org/10.1016/0169-5347(90)90223-Z 777 Dunger, W., 1998. Immigration, Ansiedlung und Primärsukzession der Bodenfauna auf 778 jungen Kippböden, in: Pflug, W. (Ed.), Braunkohlentagebau und Rekultivierung: 779 Landschaftsökologie — Folgenutzung — Naturschutz. Springer, Berlin, Heidelberg, 780 pp. 635-644. https://doi.org/10.1007/978-3-642-58846-4 53 781 Dupont, L., Torres-Leguizamon, M., René-Corail, P., Mathieu, J., 2017. Landscape features 782 impact connectivity between soil populations: a comparative study of gene flow in 783 earthworms. Molecular Ecology 26, 3128–3140. https://doi.org/10.1111/mec.14102 784 Eitminaviciute, I., 2006. Microarthropod communities in anthropogenic urban soils. 2. 785 Seasonal dynamics of microarthropod number in crossing curb soils. Zoologichesky 786 Zhurnal 1309–1320. 787 Emilsson, T., 2008. Vegetation development on extensive vegetated green roofs: Influence of 788 substrate composition, establishment method and species mix. Ecological 789 Engineering 33, 265–277. https://doi.org/10.1016/j.ecoleng.2008.05.005

790 Fountain, M.T., Hopkin, S.P., 2004. A Comparative Study of the Effects of Metal 791 Contamination on Collembola in the Field and in the Laboratory. Ecotoxicology 13, 792 573-587. https://doi.org/10.1023/B:ECTX.0000037194.66321.2c 793 Ge, B., Mehring, A.S., Levin, L.A., 2019. Urbanization alters belowground invertebrate 794 community structure in semi-arid regions: A comparison of lawns, biofilters and sage 795 scrub. Landscape and Urban Planning 192, 103664. 796 https://doi.org/10.1016/j.landurbplan.2019.103664 797 Giller, P.S., 1996. The diversity of soil communities, the 'poor man's tropical rainforest.' 798 Biodivers Conserv 5, 135–168. https://doi.org/10.1007/BF00055827 799 Guilland, C., Maron, P.A., Damas, O., Ranjard, L., 2018. Biodiversity of urban soils for 800 sustainable cities. Environmental Chemistry Letters. https://doi.org/10.1007/s10311-801 018-0751-6 802 Hopkin, S.P., 1997. Biology of the Springtails : (Insecta: Collembola): (Insecta: Collembola). 803 Oxford University Press. 804 Huang, C., Wang, T., Luo, Y., Chen, S., Kim, K.C., 2013. Assessing Collembola Biodiversity 805 Under Human Influence at Three Gorges Area, China. Environ Entomol 42, 214–222. 806 https://doi.org/10.1603/EN12253 807 Islam, M.S., Hossain, M., Hossain, M., Yasmin, M., 2018. Impact of climatic factors and soil 808 quality on the abundance and population density of Collembola in the Rajshahi 809 University Campus, Bangladesh. Journal of Entomology and Zoology Studies 6, 1119-810 1125. 811 Janion-Scheepers, C., Deharveng, L., Bedos, A., Chown, S.L., 2015. Updated list of Collembola 812 species currently recorded from South Africa. ZooKeys 55. 813 https://doi.org/10.3897/zookeys.503.8966 814 Jaskuła, R., Sulikowska-Drozd, A., Jabłońska, A., Banaś, K., Rewicz, T., 2019. Undesirable 815 immigrants: hobbyist vivaria as a potential source of alien invertebrate species. PeerJ 816 7, e7617. https://doi.org/10.7717/peerj.7617 817 Jensen, P., Jacobson, G.L., Willard, D.E., 1973. Effects of Mowing and Raking on Collembola. 818 Ecology 54, 564-572. https://doi.org/10.2307/1935341 819 Joimel, S., Baptiste Grard, Apolline Auclerc, Mickaël Hedde, Nolwenn Le Doaré, Sandrine 820 Salmon, Claire Chenu, 2018. Are Collembola "flying" onto green roofs? Ecological 821 Engineering 111, 117–124. 822 Joimel, S., Cortet, J., Jolivet, C.C., Saby, N.P.A., Chenot, E.D., Branchu, P., Consalès, J.N., 823 Lefort, C., Morel, J.L., Schwartz, C., 2016. Physico-chemical characteristics of topsoil 824 for contrasted forest, agricultural, urban and industrial land uses in France. Science of 825 The Total Environment 545–546, 40–47. 826 https://doi.org/10.1016/j.scitotenv.2015.12.035 827 Joimel, S., Schwartz, C., Hedde, M., Kiyota, S., Krogh, P.H., Nahmani, J., Pérès, G., Vergnes, A., 828 Cortet, J., 2017. Urban and industrial land uses have a higher soil biological quality 829 than expected from physicochemical quality. Science of The Total Environment 584-830 585, 614–621. https://doi.org/10.1016/j.scitotenv.2017.01.086 831 Joimel, S., Schwartz, C., Maurel, N., Magnus, B., Machon, N., Bel, J., Cortet, J., 2019. 832 Contrasting homogenization patterns of plant and collembolan communities in urban 833 vegetable gardens. Urban Ecosystems. https://doi.org/10.1007/s11252-019-00843-z 834 Lang, A., 2000. The pitfalls of pitfalls: a comparison of pitfall trap catches and absolute 835 density estimates of epigeal invertebrate predators in Arable Land. J. Pest Science 73, 836 99–106. https://doi.org/10.1007/BF02956438

837 Lehmitz, R., Russell, D., Hohberg, K., Christian, A., Xylander, W.E.R., 2012. Active dispersal of 838 oribatid mites into young soils. Applied Soil Ecology 55, 10–19. 839 https://doi.org/10.1016/j.apsoil.2011.12.003 Liu, Z., He, C., Zhou, Y., Wu, J., 2014. How much of the world's land has been urbanized, 840 841 really? A hierarchical framework for avoiding confusion. Landscape Ecology 29, 763– 771. https://doi.org/10.1007/s10980-014-0034-y 842 843 Lortie, C.J., Brooker, R.W., Choler, P., Kikvidze, Z., Michalet, R., Pugnaire, F.I., Callaway, R.M., 844 2004. Rethinking plant community theory. Oikos 107, 433–438. 845 https://doi.org/10.1111/j.0030-1299.2004.13250.x 846 Marzluff, J.M., 2019. A decadal review of urban ornithology and a prospectus for the future. 847 Ibis 1-13. https://doi.org/10.1111/ibi.12430@10.1111/(ISSN)1474-848 919X.highly_cited_us_research 849 Mayrand, F., Clergeau, P., 2018. Green Roofs and Green Walls for Biodiversity Conservation: 850 A Contribution to Urban Connectivity? Sustainability 10, 985. 851 https://doi.org/10.3390/su10040985 852 McKinney, M.L., 2008. Effects of urbanization on species richness: A review of plants and 853 animals. Urban Ecosystems 11, 161–176. https://doi.org/10.1007/s11252-007-0045-4 854 Milano, V., Cortet, J., Baldantoni, D., Bellino, A., Dubs, F., Nahmani, J., Strumia, S., Maisto, G., 855 2017. Collembolan biodiversity in Mediterranean urban parks: impact of history, 856 urbanization, management and soil characteristics. Applied Soil Ecology 119, 428-857 437. https://doi.org/10.1016/j.apsoil.2017.03.022 858 Milano, V., Maisto, G., Baldantoni, D., Bellino, A., Bernard, C., Croce, A., Dubs, F., Strumia, S., 859 Cortet, J., 2018. The effect of urban park landscapes on soil Collembola diversity: A 860 Mediterranean case study. Landscape and Urban Planning 180, 135–147. 861 https://doi.org/10.1016/j.landurbplan.2018.08.008 862 Morel, J.L., Chenu, C., Lorenz, K., 2014. Ecosystem services provided by soils of urban, 863 industrial, traffic, mining, and military areas (SUITMAs). J Soils Sediments 1–8. 864 https://doi.org/10.1007/s11368-014-0926-0 865 Natuhara, Y., Imai, C., Takeda, H., 1994. Classification and ordination of communities of soil arthropods in an urban park of Osaka City. Ecological Research 9, 131–141. 866 867 https://doi.org/10.1007/BF02347488 868 Nielsen, U.N., Osler, G.H.R., Campbell, C.D., Burslem, D.F.R.P., Wal, R. van der, 2010. The 869 influence of vegetation type, soil properties and precipitation on the composition of 870 soil mite and microbial communities at the landscape scale. Journal of Biogeography 871 37, 1317–1328. https://doi.org/10.1111/j.1365-2699.2010.02281.x 872 Oberndorfer, E., Lundholm, J., Bass, B., Coffman, R.R., Doshi, H., Dunnett, N., Gaffin, S., 873 Köhler, M., Liu, K.K.Y., Rowe, B., 2007. Green Roofs as Urban Ecosystems: Ecological 874 Structures, Functions, and Services. BioScience 57, 823-833. 875 https://doi.org/10.1641/B571005 876 Pomorski, R.J., Skarzynski, D.S., 1996. (Collembola) pól uprawnych, ziemi doniczrjwej, 877 szklarni i pieczarkarni – klucz do oznaczania // Diagnostyka czkodników roslin i ich 878 wrogów naturalnych. 879 Ponge, J.-F., 2020. Move or change, an eco-evolutionary dilemma: The case of Collembola. 880 Pedobiologia 79, 150625. https://doi.org/10.1016/j.pedobi.2020.150625 881 Ponge, J.-F., 2013. Disturbances, organisms and ecosystems: a global change perspective. 882 Ecol. Evol. 3, 1113-1124. https://doi.org/10.1002/ece3.505

884 springtail communities: the importance of dispersal and landuse history in 885 heterogeneous landscapes. Soil Biology and Biochemistry 38, 1158–1161. 886 https://doi.org/10.1016/j.soilbio.2005.09.004 Ponge, J.-F., Gillet, S., Dubs, F., Fedoroff, E., Haese, L., Sousa, J.P., Lavelle, P., 2003. 887 Collembolan communities as bioindicators of land use intensification. Soil Biology 888 889 and Biochemistry 35, 813-826. https://doi.org/10.1016/S0038-0717(03)00108-1 890 Querner, P., Bruckner, A., 2010. Combining pitfall traps and soil samples to collect 891 Collembola for site scale biodiversity assessments. Applied Soil Ecology 45, 293–297. 892 https://doi.org/10.1016/j.apsoil.2010.05.005 893 Rantalainen, M.-L., Haimi, J., Fritze, H., Pennanen, T., Setälä, H., 2008. Soil decomposer 894 community as a model system in studying the effects of habitat fragmentation and 895 habitat corridors. Soil Biology and Biochemistry 40, 853-863. 896 https://doi.org/10.1016/j.soilbio.2007.11.008 897 Rochefort, S., Shetlar, D.J., Brodeur, J., 2013. Impact of four turf management regimes on 898 arthropod abundance in lawns. Pest Management Science 69, 54-65. 899 https://doi.org/10.1002/ps.3361 900 Rota, E., Caruso, T., Migliorini, M., Monaci, F., Agamennone, V., Biagini, G., Bargagli, R., 2015. 901 Diversity and abundance of soil arthropods in urban and suburban holm oak stands. 902 Urban Ecosyst 18, 715–728. https://doi.org/10.1007/s11252-014-0425-5 903 Rumble, H., Finch, P., Gange, A.C., 2018. Green roof soil organisms: Anthropogenic 904 assemblages or natural communities? Applied Soil Ecology 126, 11–20. 905 https://doi.org/10.1016/j.apsoil.2018.01.010 906 Rumble, H., Gange, A.C., 2013. Soil microarthropod community dynamics in extensive green 907 roofs. Ecological Engineering 57, 197-204. 908 https://doi.org/10.1016/j.ecoleng.2013.04.012 909 Rusek, J., 1998. Biodiversity of Collembola and their functional role in the ecosystem. 910 Biodiversity and Conservation 7, 1207–1219. 911 https://doi.org/10.1023/A:1008887817883 912 Rzeszowski, K., 2017. The effects of anthropogenic disturbances and hydrological activity of a 913 river on soil Collembola communities in an urbanized zone. European Journal of Soil 914 Biology 82, 116–120. https://doi.org/10.1016/j.ejsobi.2017.09.004 915 Rzeszowski, K., Sterzyńska, M., 2016. Changes through time in soil Collembola communities 916 exposed to urbanization. Urban Ecosyst 19, 143–158. 917 https://doi.org/10.1007/s11252-015-0478-0 918 Rzeszowski, K., Zadrożny, P., Nicia, P., 2017. The effect of soil nutrient gradients on 919 Collembola communities inhabiting typical urban green spaces. Pedobiologia 64, 15-920 24. https://doi.org/10.1016/j.pedobi.2017.06.003 Salmon, S., Rebuffat, S., Prado, S., Sablier, M., D'Haese, C., Sun, J.-S., Ponge, J.-F., 2019. 921 922 Chemical communication in springtails: a review of facts and perspectives. Biol Fertil 923 Soils 55, 425–438. https://doi.org/10.1007/s00374-019-01365-8 924 Santorufo, L., Cortet, J., Arena, C., Goudon, R., Rakoto, A., Morel, J.-L., Maisto, G., 2014a. An 925 assessment of the influence of the urban environment on collembolan communities 926 in soils using taxonomy- and trait-based approaches. Applied Soil Ecology 78, 48–56. 927 https://doi.org/10.1016/j.apsoil.2014.02.008 928 Santorufo, L., Cortet, J., Nahmani, J., Pernin, C., Salmon, S., Pernot, A., Morel, J.L., Maisto, G., 929 2015. Responses of functional and taxonomic collembolan community structure to

Ponge, J.-F., Dubs, F., Gillet, S., Sousa, J.P., Lavelle, P., 2006. Decreased biodiversity in soil

883

930 site management in Mediterranean urban and surrounding areas. European Journal 931 of Soil Biology 70, 46–57. https://doi.org/10.1016/j.ejsobi.2015.07.003 932 Santorufo, L., Joimel, S., Auclerc, A., Deremiens, J., Grisard, G., Hedde, M., Nahmani, J., 933 Pernin, C., Cortet, J., 2021. Early colonization of constructed technosol by 934 microarthropods. Ecological Engineering 162, 106174. 935 https://doi.org/10.1016/j.ecoleng.2021.106174 936 Santorufo, L., Van Gestel, C.A.M., Maisto, G., 2014b. Sampling season affects conclusions on 937 soil arthropod community structure responses to metal pollution in Mediterranean 938 urban soils. Geoderma 226–227, 47–53. 939 https://doi.org/10.1016/j.geoderma.2014.02.001 940 Santorufo, L., Van Gestel, C.A.M., Rocco, A., Maisto, G., 2012. Soil invertebrates as 941 bioindicators of urban soil quality. Environmental Pollution 161, 57-63. 942 https://doi.org/10.1016/j.envpol.2011.09.042 943 Schindler, B.Y., Vasl, A., Blaustein, L., Gurevich, D., Kadas, G.J., Seifan, M., 2019. Fine-scale 944 substrate heterogeneity does not affect arthropod communities on green roofs. 945 PeerJ 7, e6445. https://doi.org/10.7717/peerj.6445 946 Schrader, S., Böning, M., 2006. Soil formation on green roofs and its contribution to urban 947 biodiversity with emphasis on Collembolans. Pedobiologia 50, 347–356. 948 https://doi.org/10.1016/j.pedobi.2006.06.003 949 Shochat, E., Warren, P.S., Faeth, S.H., Mcintyre, N.E., Hope, D., 2006. From patterns to 950 emerging processes in mechanistic urban ecology. Trends in ecology & evolution 21, 951 186-191. 952 Shrubovych, J., 2002. The fauna of springtails (Collembola) in Lviv. Vestnik Zoologii 36, 63-953 67. 954 Sousa, J.P., da Gama, M.M., Pinto, C., Keating, A., Calhôa, F., Lemos, M., Castro, C., Luz, T., 955 Leitão, P., Dias, S., 2004. Effects of land-use on Collembola diversity patterns in a 956 Mediterranean landscape. Pedobiologia 48, 609-622. 957 https://doi.org/10.1016/j.pedobi.2004.06.004 958 Sterzynska, M., 1990. Communities of Collembola in Natural and Transformed Soils of the 959 Linden- Oak-Hornbeam Sites of the Mazovian Lowland Poland. Fragmenta Faunistica 960 (Warsaw) 34, 165–262. 961 Sterzynska, M., 1987. Structure of springtail (Collembola) communities in the urban green of 962 Warsaw. Memorabilia Zoologica 3–18. 963 Sterzyńska, M., Nicia, P., Zadrożny, P., Fiera, C., Shrubovych, J., Ulrich, W., 2018. Urban 964 springtail species richness decreases with increasing air pollution. Ecological 965 Indicators 94, 328–335. https://doi.org/10.1016/j.ecolind.2018.06.063 966 Sukopp, H., 2008. On the Early History of Urban Ecology in Europe, in: Marzluff, J.M., 967 Shulenberger, E., Endlicher, W., Alberti, M., Bradley, G., Ryan, C., Simon, U., 968 ZumBrunnen, C. (Eds.), Urban Ecology: An International Perspective on the 969 Interaction Between Humans and Nature. Springer US, Boston, MA, pp. 79–97. 970 https://doi.org/10.1007/978-0-387-73412-5 6 971 Theenhaus, A., Scheu, S., Schaefer, M., 1999. Contramensal interactions between two 972 collembolan species: effects on population development and on soil processes. 973 Functional Ecology 13, 238–246. https://doi.org/10.1046/j.1365-2435.1999.00303.x 974 Vincent, Q., Leyval, C., Beguiristain, T., Auclerc, A., 2018. Functional structure and 975 composition of Collembola and soil macrofauna communities depend on abiotic

- 976 parameters in derelict soils. Applied Soil Ecology 130, 259–270.
- 977 https://doi.org/10.1016/j.apsoil.2018.07.002
- Yi, Z., Jinchao, F., Dayuan, X., Weiguo, S., Axmacher, J.C., 2012. A Comparison of Terrestrial
 Arthropod Sampling Methods. jore 3, 174–182. https://doi.org/10.5814/j.issn.1674764x.2012.02.010
- Young, R.F., Wolf, S.A., 2006. Goal attainment in urban ecology research: A bibliometric
 review 1975–2004. Urban Ecosyst 9, 179–193. https://doi.org/10.1007/s11252-0068589-2
- Zobel, M., 1997. The relative of species pools in determining plant species richness: an
 alternative explanation of species coexistence? Trends in Ecology & Evolution 12,
 266–269. https://doi.org/10.1016/S0169-5347(97)01096-3
- 987