



**HAL**  
open science

# Single Cell Analysis Reveals a New Microsporidia-Host Association in a Freshwater Lake

Marina Chauvet, Arthur Monjot, Anne Mone, Cécile Lepère

► **To cite this version:**

Marina Chauvet, Arthur Monjot, Anne Mone, Cécile Lepère. Single Cell Analysis Reveals a New Microsporidia-Host Association in a Freshwater Lake. *Microbial ecology*, In press, 10.1007/s00248-022-02023-7. hal-03666982

**HAL Id: hal-03666982**

**<https://hal.science/hal-03666982v1>**

Submitted on 13 May 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



23 **Abstract**

24

25 Microsporidia are a large group of obligate intracellular eukaryotic parasites. Recent studies  
26 suggest that their diversity can be huge in freshwater lake ecosystems especially in the <150 µm  
27 size fraction. However, little is known about their hosts and therefore their impact on the trophic  
28 food web functioning. In this study, single cell analysis and fluorescence microscopy were used  
29 to detect new host-parasite association within rotifer communities in lake Aydat (France). Our  
30 analysis showed the existence of a potential new species belonging to the *Crispospora* genus  
31 able of infecting the rotifer *Kellicottia* with a high prevalence (42.5%) suggesting that  
32 Microsporidia could have a great impact on the rotifer populations regulation in lakes.

33

34 **Keywords**

35 Microsporidia, host-parasite association, single cell, TSA-FISH, lake, microbial ecology

36

37 Microsporidia are obligate intracellular eukaryotic parasites related to Fungi. They are mostly  
38 known to parasitize a broad range of hosts among the animal kingdom in different environments  
39 [1]. Recent metabarcoding studies revealed an unexpected diversity in lake ecosystems  
40 especially in the <150 µm size fraction suggesting that Microsporidia could have an impact on  
41 the planktonic communities at the base of the food web [2, 3]. However, they have rarely been  
42 described as parasites of protists and microzooplankton. In marine environments, they are  
43 mainly known to parasite gregarines (e.g. [4]), but they were also shown in association with  
44 Paramyxea [5]. In freshwaters, only a few observations showed Microsporidia parasiting  
45 ciliates with only two microsporidian species characterized by morphological and molecular  
46 approaches (*Euplotespora binucleata* and *Globosporidium paramecii* parasitizing the ciliates  
47 *Euplotes woodruffi* and *Paramecium aurelia*, respectively [6, 7]). Among microzooplankton,  
48 Microsporidia have been observed in the rotifers *Brachionus calyciflorus* as well as in  
49 *Asplanchna priodonta* and *A. brightwellii* in both river and freshwater lakes [8, 9] however  
50 these observations (no molecular data) remain sporadic and for most at low prevalence.  
51 Moreover, with co-occurrence network and microscopic observations, a more recent study  
52 suggested that Microsporidia could infect a broader range of rotifers [3]. We therefore used, in  
53 this study, single cell analysis as well as Tyramide Signal Amplification – Fluorescent *In Situ*  
54 Hybridization (TSA-FISH) to detect new host-parasite association within microzooplankton  
55 communities in a lake ecosystem (lake Aydat, Massif Central, France) and especially (i)  
56 confirm the hypothesis raised about the broader range of rotifers parasitized by Microsporidia  
57 and (ii) characterize by molecular information and microscopic observation both partners.

58 Zooplankton was sampled with vertical tows of 20 µm-mesh plankton net at the central point  
59 of the lake (depth = 14 m) on the 18/05/2021 (see Supplementary Material). 17 rotifers,  
60 identified morphologically as probably belonging to the genera *Kellicottia* (n = 13) and  
61 *Keratella* (n = 4), were isolated by micromanipulation and individually placed in a 96 wells

62 plate before Whole Genome Amplification (WGA) (see Supplementary Material). The Single  
63 Amplified Genomes (SAGs) were then screened for the presence of Microsporidia by the  
64 amplification of the V1-V3 region of the microsporidian gene coding for the SSU rRNA  
65 according to Chauvet *et al.* [3]. The sequences obtained after cloning and sequencing were  
66 compared to those available in public databases and included in a phylogenetic tree (see  
67 Supplementary Material). This analysis showed that on the 17 isolated rotifers, 6 *Kellicottia*  
68 had a positive PCR amplification signal with Microsporidia primers however no Microsporidia  
69 amplification was recorded in *Keratella*. All Microsporidia sequences were affiliated to the  
70 genus *Crispospora* (Fig. 1) which is affiliated to the clade IV2; Arlequin group (defined in  
71 Dubuffet *et al.* [2]), and well represented (~ 50% of the total OTUs) in metabarcoding studies  
72 conducted in lake Aydat [2, 3]. With 97% identity with *Crispospora chironomi* (the only known  
73 species of the genus so far) these sequences could represent a new species of the *Crispospora*  
74 genus (Fig. S2a). Indeed, although thresholds for species delineation may vary, an identity close  
75 to 100% is suggested for taxa separation at the species level for Microsporidia [2, 11].  
76 *Crispospora chironomi* is known to parasite chironomids eggs [12] that can be found in aquatic  
77 environments especially when they are eutrophic and anthropized [13].  
78 On the Microsporidia positive SAGs, rotifers were characterized by amplification of the COI  
79 using the 30F and 885R primers set according to Zhang *et al.* [14] with some modifications (see  
80 Supplementary Material). The rotifer affiliation analysis revealed that this potential new species  
81 of *Crispospora* is able to parasite the genus *Kellicottia* (Fig. S1). Even though *Kellicottia*  
82 sequences grouped into two clades in the phylogenetic analysis, species determination is  
83 challenging due to high intraspecific variation of identity percent which can be higher than 20%  
84 for rotifers (Fig. S2b) [15].  
85 To visualize and quantify Microsporidia-rotifer associations, TSA-FISH was performed on the  
86 rest of the plankton net sample, using a Microsporidia specific probe (USP01) according to

87 Chauvet *et al.* [3] (see Supplementary Material). Positive signals were obtained in *Kellicottia*  
88 rotifers with a surprisingly high prevalence (42.5%, total rotifers observed = 73) (Fig. 2).  
89 Microsporidia were so far recorded in a small number of rotifers [8, 9]. *Microsporidium* sp. and  
90 *Microsporidium asperspora* have been observed in *Brachionus calyciflorus*, and  
91 *Microsporidium polygona* in *Asplanchna priodonta* and *A. brightwellii* in freshwater  
92 ecosystems most of the time at low prevalence (*e.g.* *Brachionus calyciflorus* by *Microsporidium*  
93 sp. on 1.2% of the host population, [9]). However, these descriptions were only morphological  
94 and neither the site of infection nor the microsporidian spore or intracellular form were  
95 described. Here, we identified an infection which could be in the rotifer's stomach or  
96 pseudocoelom. The intracellular forms of the Microsporidia were on average 1.53  $\mu\text{m}$  long,  
97 most being oval, some spherical. These findings were consistent with the described shape of  
98 the only known *Crispospora* species (*i.e.* *Crispospora chironomi*), with two shapes identified:  
99 short oval and spherical, with a size of 1.5 x 2.5  $\mu\text{m}$  and 1.5 x 2  $\mu\text{m}$ , respectively [12].  
100 This finding is highly relevant since Microsporidia, along with hyphomycetes and oomycetes  
101 [17, 18] could be an important regulator of these populations in lakes. Rotifers are indeed an  
102 important component of plankton in aquatic ecosystems playing fundamental role in both  
103 microbial loop and classical food web and as a link in energy flow between primary producers  
104 (algae, cyanobacteria) and higher trophic levels (zooplankton, shellfish, fish) [16]. *Kellicottia*  
105 can reach 600 ind./l in lake Aydat and is the most abundant rotifers along with *Keratella* [10].  
106 Microsporidia sequences retrieved during this study had an identity >99% with an highly  
107 abundant OTU identified in the same lake in a previous study where the temporal dynamic of  
108 the Microsporidia diversity was analyzed over one year [3] (Fig. S2a). This *Crispospora* OTU  
109 and the only OTU affiliated to *Kellicottia* presented the same pattern of temporal variations  
110 (data not shown). This co-variation and the high prevalence of this association suggest an  
111 impact on the dynamic of *Kellicottia* in the lake however this needs to be confirmed for example

112 by the culture of both partners which will also allow the characterization of the developmental  
113 cycle of this potential new species of Microsporidia.

114

115 **References**

116

- 117 1. Murareanu BM, Sukhdeo R, Qu R, Jiang J, Reinke AW (2021) Generation of a  
118 Microsporidia Species Attribute Database and Analysis of the Extensive Ecological and  
119 Phenotypic Diversity of Microsporidia. *mBio* 12:e01490-21.  
120 <https://doi.org/10.1128/mBio.01490-21>
- 121 2. Dubuffet A, Chauvet M, Moné A, Debroas D, Lepère C (2021) A phylogenetic  
122 framework to investigate the microsporidian communities through metabarcoding and its  
123 application to lake ecosystems. *Environ Microbiol* 23:4344–4359.  
124 <https://doi.org/10.1111/1462-2920.15618>
- 125 3. Chauvet M, Debroas D, Moné A, Dubuffet A and Lepère C (2022) Temporal variations  
126 of Microsporidia diversity and discovery of new host-parasite interactions in a lake  
127 ecosystem. *Environ Microbiol*. <https://doi.org/10.1111/1462-2920.15950>
- 128 4. Mikhailov KV, Simdyanov TG, Aleoshin VV (2016) Genomic Survey of a  
129 Hyperparasitic Microsporidian *Amphiamblys* sp. (Metchnikovellidae). *Genome Biol*  
130 *Evol* 9:454–467. <https://doi.org/10.1093/gbe/evw235>
- 131 5. Stentiford GD, Ramilo A, Abollo E, Kerr R, Bateman KS, Feist SW, et al (2017)  
132 *Hyperspora aquatica* n.g.n., n.sp. (Microsporidia), hyperparasitic in *Marteilia cochillia*  
133 (Paramyxida), is closely related to crustacean-infecting microsporidian taxa. *Parasitology*  
134 144:186–199. <https://doi.org/10.1017/S0031182016001633>
- 135 6. Fokin SI, Di Giuseppe G, Erra F, Dini F (2008) *Euplotespora binucleata* n. gen., n. sp.  
136 (Protozoa: Microsporidia), a parasite infecting the hypotrichous ciliate *Euplotes*



- 137 *woodruffi*, with observations on microsporidian infections in ciliophora. J Eukaryot  
138 Microbiol 55:214–228. <https://doi.org/10.1111/j.1550-7408.2008.00322.x>
- 139 7. Yakovleva Y, Nasonova E, Lebedeva N, Lanzoni O, Petroni G, Potekhin A, et al (2020)  
140 The first case of microsporidiosis in *Paramecium*. Parasitology 147:957–971.  
141 <https://doi.org/10.1017/S0031182020000633>
- 142 8. Gorbunov AK, Kosova AA (2001) Parasites in rotifers from the Volga delta.  
143 Hydrobiologia 446:51–55. <https://doi.org/10.1023/A:1017569004998>
- 144 9. Wolska M, Mazurkiewicz-Zapalowicz K (2013) Parasites of zooplankton and periphyton  
145 assemblages in the littoral zone of lakes in Drawa National Park, Poland. Acta Mycol  
146 48:51–59. <https://doi.org/10.5586/am.2013.007>
- 147 10. Lair N, Oulad Ali H (1990) Grazing and assimilation rates of natural populations of  
148 planktonic rotifers *Keratella cochlearis*, *Keratella quadrata* and *Kellicottia longispina* in  
149 a eutrophic lake (Aydat, France). Hydrobiologia 194:119–131.  
150 <https://doi.org/10.1007/BF00028413>
- 151 11. Tokarev YS, Huang W-F, Solter LF, Malysh JM, Becnel JJ, Vossbrinck CR (2020) A  
152 formal redefinition of the genera *Nosema* and *Vairimorpha* (Microsporidia:  
153 Nosematidae) and reassignment of species based on molecular phylogenetics. J Invertebr  
154 Pathol 169:107279. <https://doi.org/10.1016/j.jip.2019.107279>
- 155 12. Tokarev YS, Voronin VN, Seliverstova EV, Pavlova OA, Issi IV (2010) Life cycle,  
156 ultrastructure, and molecular phylogeny of *Crispospora chironomi* g.n. sp.n.  
157 (Microsporidia: Terresporidia), a parasite of *Chironomus plumosus* L. (Diptera:

- 158 Chironomidae). *Parasitol Res* 107:1381–1389. <https://doi.org/10.1007/s00436-010-2012->  
159 x
- 160 13. Langdon PG, Ruiz Z, Brodersen KP, Foster IDL (2006) Assessing lake eutrophication  
161 using chironomids: understanding the nature of community response in different lake  
162 types. *Freshw Biol* 51:562–577. <https://doi.org/10.1111/j.1365-2427.2005.01500.x>
- 163 14. Zhang Y, Xu S, Sun C, Dumont H, Han B-P (2021) A new set of highly efficient primers  
164 for COI amplification in rotifers. *Mitochondrial DNA B* 6:636–640.  
165 <https://doi.org/10.1080/23802359.2021.1878951>
- 166 15. Zhang Y-N, Xu S-L, Huang Q, Liu P, Han B-P (2021) Application of COI Primers  
167 30F/885R in Rotifers to Regional Species Diversity in (Sub)Tropical China. *Diversity*  
168 13:390. <https://doi.org/10.3390/d13080390>
- 169 16. Rico-Martínez R, Arzate-Cárdenas MA, Robles D, Pérez-Legaspi I, Alvaro-Flores J,  
170 Santos Medrano GE (2016) Rotifers as Models in Toxicity Screening of Chemicals and  
171 Environmental Samples. In *Invertebrates - Experimental Models in Toxicity Screening*.  
172 InTech pp 57–99. <https://doi.org/10.5772/61771>
- 173 17. Barron GL, Szijarto E (1990) A New Genus of the Hyphomycetes Endoparasitic in  
174 Rotifers. *Mycologia* 82:134–137. <https://doi.org/10.2307/3759974>
- 175 18. Thomas SH, Housley JM, Reynolds AN, Penczykowski RM, Kenline KH, Hardegee N,  
176 et al (2011) The ecology and phylogeny of oomycete infections in *Asplanchna* rotifers:  
177 Oomycete infections in *Asplanchna*. *Freshw Biol* 56:384–394.  
178 <https://doi.org/10.1111/j.1365-2427.2010.02505.x>

179

180 **Statements and Declarations**

181

182 **Funding**

183 The authors declare that no funds, grants, or other support were received during the  
184 preparation of this manuscript.

185

186 **Competing Interests**

187 The authors declare they have no competing interests.

188

189 **Availability of data and material**

190 All data and material are available in this publication and additional information online  
191 (supplementary material and supplementary figures). Sequences are available on GenBank,  
192 accession numbers: from ON026091 to ON026096 for the microsporidian sequences and from  
193 ON033621 to ON033624 for the rotifers.

194 **Figure Legends**

195

196 **Fig. 1** Maximum Likelihood phylogeny of microsporidian sequences.

197 We included in this phylogeny several sequences from undescribed microsporidian species  
198 (found through BLAST search) and a sequence found extensively in a previous study (in italic,  
199 [3]), in addition to described species, all belonging to the Arlequin group (clade IV2).  
200 Sequences from this study are given in bold. Outgroup species are representative species of  
201 other groups belonging to clade IV2. Numbers next to branches indicate the ML bootstrap  
202 support (100 replicates).


203

204 **Fig. 2** Microsporidia (targeted by TSA-FISH with the USP01 probe, [3]) infecting a rotifer  
205 (genus *Kellicottia*).

206 Under UV showing nucleus after DAPI staining (a), under blue light showing Microsporidia in  
207 green fluorescence (b), overlay (c). Magnification on microsporidian cells inside a rotifer with  
208 a confocal microscope (UV (d), blue light (e), overlay (f)).

209 Scale bars, 20  $\mu\text{m}$

210



0.10

