



HAL
open science

Impact of rainfall spatial variability on Flash Flood Forecasting

Audrey Douinot, H el ene Roux, Pierre-Andr e Garambois, K evin Larnier

► **To cite this version:**

Audrey Douinot, H el ene Roux, Pierre-Andr e Garambois, K evin Larnier. Impact of rainfall spatial variability on Flash Flood Forecasting. European Geosciences Union General Assembly 2014, Apr 2014, Vienna, Austria. pp.0. hal-03667262

HAL Id: hal-03667262

<https://hal.science/hal-03667262v1>

Submitted on 13 May 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin ee au d ep ot et  a la diffusion de documents scientifiques de niveau recherche, publi es ou non,  emanant des  tablissements d'enseignement et de recherche fran ais ou  trangers, des laboratoires publics ou priv es.



Open Archive Toulouse Archive Ouverte (OATAO)

OATAO is an open access repository that collects the work of Toulouse researchers and makes it freely available over the web where possible.

This is an author-deposited version published in: <http://oatao.univ-toulouse.fr/>
Eprints ID: 11370

To cite this version:

Douinot, Audrey and Roux, H el ene and Garambois, Pierre-Andr e and Larnier, K evin *Impact of rainfall spatial variability on Flash Flood Forecasting*. (2014)
In: European Geosciences Union General Assembly 2014, 27 April 2014 - 02 May 2014 (Vienna, Austria).

Any correspondence concerning this service should be sent to the repository administrator:
staff-oatao@inp-toulouse.fr

Impact of rainfall spatial variability on Flash Flood Forecasting

Audrey Douinot, H el ene Roux, Pierre-Andr e Garambois, and Kevin Larnier

IMFT (Institut de M ecanique des Fluides de Toulouse), Universit e de Toulouse, Toulouse, France (audrey.douinot@imft.fr)

According to the United States National Hazard Statistics database, flooding and flash flooding have caused the largest number of deaths of any weather-related phenomenon over the last 30 years (Flash Flood Guidance Improvement Team, 2003). Like the storms that cause them, flash floods are very variable and non-linear phenomena in time and space, with the result that understanding and anticipating flash flood genesis is far from straightforward. In the U.S., the Flash Flood Guidance (FFG) estimates the average number of inches of rainfall for given durations required to produce flash flooding in the indicated county.

In Europe, flash flood often occurred on small catchments (approximately 100 km²) and it has been shown that the spatial variability of rainfall has a great impact on the catchment response (Le Lay and Saulnier, 2007). Therefore, in this study, based on the Flash flood Guidance method, rainfall spatial variability information is introduced in the threshold estimation. As for FFG, the threshold is the number of millimeters of rainfall required to produce a discharge higher than the discharge corresponding to the first level (yellow) warning of the French flood warning service (SCHAPI: Service Central d'Hydrom et eorologie et d'Appui   la Pr evision des Inondations). The indexes δ_1 and δ_2 of Zocatelli et al. (2010), based on the spatial moments of catchment rainfall, are used to characterize the rainfall spatial distribution. Rainfall spatial variability impacts on warning threshold and on hydrological processes are then studied.

The spatially distributed hydrological model MARINE (Roux et al., 2011), dedicated to flash flood prediction is forced with synthetic rainfall patterns of different spatial distributions. This allows the determination of a warning threshold diagram: knowing the spatial distribution of the rainfall forecast and therefore the 2 indexes δ_1 and δ_2 , the threshold value is read on the diagram. A warning threshold diagram is built for each studied catchment.

The proposed methodology is applied on three Mediterranean catchments often submitted to flash floods. The new forecasting method as well as the Flash Flood Guidance method (uniform rainfall threshold) are tested on 25 flash floods events that had occurred on those catchments.

Results show a significant impact of rainfall spatial variability. Indeed, it appears that the uniform rainfall threshold (FFG threshold) always overestimates the observed rainfall threshold. The difference between the FFG threshold and the proposed threshold ranges from 8% to 30%. The proposed methodology allows the calculation of a threshold more representative of the observed one. However, results strongly depend on the related event duration and on the catchment properties. For instance, the impact of the rainfall spatial variability seems to be correlated with the catchment size. According to these results, it seems to be interesting to introduce information on the catchment properties in the threshold calculation.

Flash Flood Guidance Improvement Team, 2003. River Forecast Center (RFC) Development Management Team. Final Report. Office of Hydrologic Development (OHD), Silver Spring, Maryland.

Le Lay, M. and Saulnier, G.-M., 2007. Exploring the signature of climate and landscape spatial variabilities in flash flood events: Case of the 8–9 September 2002 C evennes-Vivarais catastrophic event. *Geophysical Research Letters*, 34(L13401), doi:10.1029/2007GL029746.

Roux, H., Labat, D., Garambois, P.-A., Maubourguet, M.-M., Chorda, J. and Dartus, D., 2011. A physically-based parsimonious hydrological model for flash floods in Mediterranean catchments. *Nat. Hazards Earth Syst. Sci.* J1 - NHESS, 11(9), 2567-2582.

Zocatelli, D., Borga, M., Zanon, F., Antonescu, B. and Stancalie, G., 2010. Which rainfall spatial information for flash flood response modelling? A numerical investigation based on data from the Carpathian range, Romania. *Journal of Hydrology*, 394(1-2), 148-161.