



HAL
open science

A Consecutive Lehmer Code for Parabolic Quotients of the Symmetric Group

Wenjie Fang, Henri Mühle, Jean-Christophe Novelli

► **To cite this version:**

Wenjie Fang, Henri Mühle, Jean-Christophe Novelli. A Consecutive Lehmer Code for Parabolic Quotients of the Symmetric Group. *The Electronic Journal of Combinatorics*, 2021, 28 (3), 10.37236/10578 . hal-03667557

HAL Id: hal-03667557

<https://hal.science/hal-03667557v1>

Submitted on 21 Dec 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A Consecutive Lehmer Code for Parabolic Quotients of the Symmetric Group

Wenjie Fang

Laboratoire Informatique Gaspard Monge
Université Gustave Eiffel
CNRS, ESIEE Paris
F-77454 Marne-la-Vallée, France
wenjie.fang@u-pem.fr

Henri Mühle*

Institut für Algebra
Technische Universität Dresden
D-01062 Dresden, Germany
henri.muehle@tu-dresden.de

Jean-Christophe Novelli[†]

Laboratoire Informatique Gaspard Monge
Université Gustave Eiffel
CNRS, ESIEE Paris
F-77454 Marne-la-Vallée, France
novelli@univ-mlv.fr

Submitted: Jul 16, 2021; Accepted: Aug 19, 2021; Published: Sep 24, 2021

© The authors. Released under the CC BY-ND license (International 4.0).

Abstract

In this article we define an encoding for parabolic permutations that distinguishes between parabolic 231-avoiding permutations. We prove that the componentwise order on these codes realizes the parabolic Tamari lattice, and conclude a direct and simple proof that the parabolic Tamari lattice is isomorphic to a certain ν -Tamari lattice, with an explicit bijection. Furthermore, we prove that this bijection is closely related to the map Θ used when the lattice isomorphism was first proved in (Ceballos, Fang and Mühle, 2020), settling an open problem therein.

Keywords: parabolic quotient, symmetric group, Lehmer code, parabolic Tamari lattice, ν -Tamari lattice, bracket vector

Mathematics Subject Classifications: 05A19, 06B99

*HM has received funding from the European Research Council (Grant Agreement no. 681988, CSP-Infinity).

[†]JCN was partially supported by the CARPLO project of the Agence Nationale de la recherche (ANR-20-CE40-0007).

1 Introduction

A (right) *inversion* of a permutation w is a pair of indices (i, j) with $i < j$ such that $w(i) > w(j)$. The number of inversions of w can therefore be regarded as a degree of disorder of w . The *Lehmer code* associated with w , attributed to Lehmer [9] but can be traced back to Laisant [8], is the integer tuple whose i^{th} entry counts the number of inversions of w of the form (i, \cdot) .

Björner and Wachs defined a “consecutive” version of the Lehmer code in [2, Section 9], which we shall call the *BW-code* of w . This encoding associates an integer tuple with a permutation w whose i^{th} entry counts the length k of the longest sequence such that (i, j) is an inversion for all $j \in \{i+1, i+2, \dots, i+k\}$.

In contrast to the original Lehmer code, the BW-code no longer uniquely determines a permutation. However, the permutations with the same BW-code form an interval in the (left) weak order on the group of all permutations, the *symmetric group* [2, Proposition 9.10]. This (left) *weak order* is defined by containment of (right) inversion sets.

Another consequence of [2, Proposition 9.10] is that among all permutations with the same BW-code, there is a unique permutation w which *avoids* the pattern 231, *i.e.*, in which no three indices $i < j < k$ exist such that $w(k) < w(i) < w(j)$, and this permutation minimizes the number of inversions among all permutations with the same BW-code as w .

Let us denote the symmetric group of degree n by \mathfrak{S}_n , and its subset of all 231-avoiding permutations by $\mathfrak{S}_n(231)$. The (left) weak order on \mathfrak{S}_n is a lattice, *i.e.*, every two elements have a unique lower bound and a unique upper bound. This property was established independently by Guilbaud and Rosenstiehl [7] and Yanagimoto and Okamoto [15]. The restriction of this lattice to $\mathfrak{S}_n(231)$ was proved to be a sublattice by Björner and Wachs [2, Theorem 9.6(i)] and a quotient lattice by Reading [13, Theorem 5.1]. In fact, the resulting lattice incarnates the famous *Tamari lattice* denoted by \mathcal{T}_n , first defined by Tamari in [14]. We can thus see the BW-code as a concrete and simple way to quotient the weak order on \mathfrak{S}_n into \mathcal{T}_n .

An analogue of 231-avoiding permutations for parabolic quotients of \mathfrak{S}_n was introduced by Mühle and Williams in [10], and it was shown that these permutations constitute a quotient lattice (but no longer a sublattice) of the corresponding (left) weak order, the *parabolic Tamari lattice* [10, Theorem 1]. Since any parabolic quotient of \mathfrak{S}_n is naturally indexed by a composition α of n , we call the resulting lattice the *α -Tamari lattice* \mathcal{T}_α . The main purpose of this article is to define a parabolic analogue of the BW-code; see Definition 6. We prove that the componentwise order on these parabolic BW-codes is isomorphic to \mathcal{T}_α .

It was shown in [10, Theorem 1] that \mathcal{T}_α is in fact a quotient lattice of the weak order on \mathfrak{S}_α with respect to a certain equivalence relation R_α on \mathfrak{S}_α . The corresponding equivalence classes form intervals in the weak order and if two α -permutations are equivalent, then one can be obtained from the other by a sequence of certain local moves. Moreover, each equivalence class contains a unique $(\alpha, 231)$ -avoiding permutation which is also the smallest element in the corresponding weak order interval. Globally, determining whether two α -permutations are equivalent is quite tedious, because one essentially has to compare

the smallest elements in the corresponding equivalence classes. Our parabolic BW-code provides a simple criterion for equivalence, because we prove that two α -permutations are equivalent with respect to R_α if and only if they have the same code; see Lemma 17.

If we denote the set of parabolic BW-codes by \mathcal{C}_α and the componentwise order on integer tuples (of the same length) by \leq_{comp} , then our first main result follows immediately.

Theorem 1. *For every $n > 0$ and every integer composition α of n it holds that $\mathcal{T}_\alpha \cong (\mathcal{C}_\alpha, \leq_{\text{comp}})$.*

Originally, the Tamari lattice was defined in terms of a “rotation” operation on parenthesizations, binary trees or equivalently Dyck paths. A *northeast path* is a lattice path in \mathbb{N}^2 comprised of north steps (marked by N) and east steps (marked by E) of unit length. A *Dyck path* of semilength n is equivalent to a northeast path that stays weakly above the staircase path $(NE)^n$ and uses n north and n east steps.

A *rotation* of a northeast path exchanges two portions of the path under certain conditions, and \mathcal{T}_n arises as the rotation order on the set of Dyck paths of semilength n . An extension of this construction was introduced by Préville-Ratelle and Viennot in [12]. In that paper, the set of all northeast paths weakly above a fixed northeast path ν , which start and end at the same coordinates as ν , was considered. Ordering this set by rotation produces another lattice, the *ν -Tamari lattice* [12, Theorem 1.1].

For any composition $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_r)$ of n , we can define the α -bounce path $\nu_\alpha = N^{\alpha_1} E^{\alpha_1} N^{\alpha_2} E^{\alpha_2} \dots N^{\alpha_r} E^{\alpha_r}$. It was established by Ceballos, Fang and Mühle in [3, Theorem II] that \mathcal{T}_α is isomorphic to the ν_α -Tamari lattice. The proof of this result is rather technical, using some deep lattice-theoretic properties of \mathcal{T}_α , namely the fact that \mathcal{T}_α is extremal and is thus uniquely determined by its Galois graph. The second main contribution of this article is a much simpler and direct proof of this result.

In general, the ν -Tamari lattice admits a simple encoding as the componentwise order on so-called ν -bracket vectors, given by Ceballos, Padrol and Sarmiento in [4, Theorem 4.2]. If $\nu = \nu_\alpha$, then the corresponding bracket vectors can be converted in a simple way into parabolic BW-codes. Since both parabolic BW-codes and bracket vectors are ordered componentwise, the proof of the next result follows readily.

Theorem 2 ([3, Theorem II]). *For every $n > 0$ and every integer composition α of n , the ν_α -Tamari lattice is isomorphic to \mathcal{T}_α .*

The original proof of Theorem 2 in [3] did not provide an explicit map between the two lattices, but rather passed through their Galois graphs, whose elements are related by a map Θ between the two lattices. In [3], it was postulated as Open Problem 2.23 to prove that Θ extends to a full lattice isomorphism, not limited to elements of the Galois graphs. Using parabolic BW-codes, by introducing a stack processing procedure on $(\alpha, 231)$ -avoiding permutations, we settle this open problem affirmatively, while giving another interpretation of parabolic BW-codes; see Corollary 40.

In Section 2, we recall the basic definitions regarding parabolic quotients of the symmetric group, parabolic pattern avoidance and the weak order. In Section 3, we define the parabolic BW-codes and prove Theorem 1.

In Section 4, we first recall the definitions of Dyck paths and northeast paths, as well as ordinary Tamari lattices and ν -Tamari lattices. We then prove Theorem 2 in Section 4.3 by describing an explicit conversion from parabolic BW-codes to ν_α -bracket vectors. Finally, in Section 5, after introducing some classical constructions that we need, we give a combinatorial interpretation of the map Θ mentioned after Theorem 2 in terms of a certain stack-processing procedure, and relate the bijection between parabolic BW-codes and ν_α -bracket vectors to Θ , thus solving [3, Open Problem 2.23].

2 α -permutations and the α -Tamari lattice

Throughout this article, we fix an integer $n > 0$ and define $[n] \stackrel{\text{def}}{=} \{1, 2, \dots, n\}$.

2.1 α -permutations

Let $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_r)$ be a composition of n . For $a \in [r]$, we define

$$s_a \stackrel{\text{def}}{=} \alpha_1 + \alpha_2 + \dots + \alpha_a,$$

and we set $s_0 \stackrel{\text{def}}{=} 0$. The set $\{s_{a-1}+1, s_{a-1}+2, \dots, s_a\}$ is the a^{th} α -region.

The *indicator map* $\varrho_\alpha: [n] \rightarrow [r]$ is defined by $\varrho_\alpha(i) = a$ if and only if $s_{a-1} < i \leq s_a$. In other words, $\varrho_\alpha(i)$ is the index of the α -region containing i . When no confusion will arise, we will drop the subscript α . For three indices $i < j < k$ with $\varrho(i) < \varrho(k)$, we say that j is in an α -region *strictly between* i and k if $\varrho(i) < \varrho(j) < \varrho(k)$.

Let \mathfrak{S}_n denote the *symmetric group* of degree n . We are interested in the subset of *α -permutations*, defined by

$$\mathfrak{S}_\alpha \stackrel{\text{def}}{=} \{w \in \mathfrak{S}_n \mid \text{if } \varrho(i) = \varrho(i+1), \text{ then } w(i) < w(i+1)\}.$$

Clearly, if $\alpha = (1, 1, \dots, 1)$, then $\mathfrak{S}_\alpha = \mathfrak{S}_n$.

Remark 3. If we consider the subgroup $G \stackrel{\text{def}}{=} \mathfrak{S}_{|\alpha_1|} \times \mathfrak{S}_{|\alpha_2|} \times \dots \times \mathfrak{S}_{|\alpha_r|}$ of \mathfrak{S}_n , then we may identify \mathfrak{S}_α with the set of minimal-length representatives of the left cosets in \mathfrak{S}_n/G .

An α -permutation $w \in \mathfrak{S}_\alpha$ has an *$(\alpha, 231)$ -pattern* if there are three indices $i < j < k$ —each in different α -regions—such that $w_i < w_j$ and $w_i = w_k + 1$. If w does not have an $(\alpha, 231)$ -pattern, then w is *$(\alpha, 231)$ -avoiding*. Let $\mathfrak{S}_\alpha(231)$ denote the set of *$(\alpha, 231)$ -avoiding permutations*.

Remark 4. In the case $\alpha = (1, 1, \dots, 1)$, the $(\alpha, 231)$ -avoiding permutations are exactly the *classical 231-avoiding permutations*: one can either require $w_i = w_k + 1$ or not, since if w has any 231-pattern, then one can find one with the extra condition $w_i = w_k + 1$. In the general case, these notions differ since i could belong to the same α -region as j . For example, 3 24 1 belongs to $\mathfrak{S}_{(1,2,1)}(231)$ whereas it has a classical 231-pattern with elements in different α -regions.

2.2 The weak order

For $w \in \mathfrak{S}_\alpha$, we define its *(right) inversion set* by

$$\text{Inv}(w) \stackrel{\text{def}}{=} \{(i, j) \mid i < j \text{ and } w_i > w_j\}.$$

This enables us to define a partial order—the *(left) weak order*—on \mathfrak{S}_α by setting

$$u \leq_L v \quad \text{if and only if} \quad \text{Inv}(u) \subseteq \text{Inv}(v).$$

Two permutations $u, v \in \mathfrak{S}_\alpha$ form a *cover relation*—denoted by $u \triangleleft_L v$ —if $u <_L v$ and there is no $w \in \mathfrak{S}_\alpha$ with $u <_L w <_L v$. It follows from the analogous property of the weak order on the full group \mathfrak{S}_n that in \mathfrak{S}_α we have $u \triangleleft_L v$ if and only if there are two indices $i < j$ in different α -regions, such that $u_i = u_j - 1$, and

$$v_k = \begin{cases} u_j, & \text{if } k = i, \\ u_i, & \text{if } k = j, \\ u_k, & \text{otherwise.} \end{cases}$$

The partially ordered set $(\mathfrak{S}_\alpha, \leq_L)$ is a lattice by [15, Theorem 2.1]; see also [7]. For an arbitrary composition α of n , it follows from [1, Theorem 4.1] that $(\mathfrak{S}_\alpha, \leq_L)$ is an interval of (\mathfrak{S}_n, \leq_L) , and thus also a lattice.

The partially ordered set $\mathcal{T}_\alpha \stackrel{\text{def}}{=} (\mathfrak{S}_\alpha(231), \leq_L)$ is the *α -Tamari lattice*. This name is justified by the following result.

Theorem 5 ([10, Theorem 1]). *\mathcal{T}_α is a lattice for every $n > 0$ and every integer composition α of n .*

3 A generalized Lehmer code for \mathfrak{S}_α

3.1 Encoding α -permutations

We consider the following set of integer tuples.

Definition 6. Let \mathcal{C}_α denote the set of all integer tuples (c_1, c_2, \dots, c_n) with the following properties:

- (C1) $0 \leq c_i \leq n - s_{\varrho(i)}$ for all $i \in [n]$;
- (C2) $c_i \leq c_{i+1}$ for all $i \in [n - 1]$ such that $\varrho(i) = \varrho(i + 1)$;
- (C3) $c_{s_a} \leq c_i - s_a + s_{\varrho(i)}$ for all $i \in [s_{r-2}]$ and all $a \in \{\varrho(i)+1, \varrho(i)+2, \dots, r-1\}$ such that $c_i \geq s_a - s_{\varrho(i)}$.

The set $\mathcal{C}_{(1,1,\dots,1)}$ is precisely the set of integer tuples defined in [2, Definition 9.1].

Remark 7. The statement of (C3) is directly true if $a = r$ and trivial if $i > s_{r-2}$, hence the restriction to $i \in [s_{r-2}]$ and $a < r$.

Indeed, by (C1), $c_n = 0$ so that the implication required by (C3) is trivially satisfied when $a = r$. If $i > s_{r-2}$, then $\varrho(i) \geq r - 1$, so that the only case one could consider is again $a = r$.

For example, with $n = 3$ and $\alpha = (2, 1)$, all conditions boil down to $0 \leq c_1 \leq c_2 \leq 1$ and $c_3 = 0$, hence three solutions. With $n = 3$ and $\alpha = (1, 2)$, one gets $0 \leq c_1 \leq 2$ and $0 \leq c_2 \leq c_3 \leq 0$, again providing three solutions. One can check that they are indeed the codes obtained in Table 1 (right column).

To see all conditions of the definition play a role, one has to consider compositions of at least three parts and at least one greater than one. For example, if $\alpha = (1, 2, 1)$, one gets the following set of relations: $0 \leq c_1 \leq 3$, $0 \leq c_2 \leq c_3 \leq 1$, $c_4 = 0$, and the extra condition coming from (C3): $c_1 \geq 2 \Rightarrow c_3 \leq c_1 - 2$. In practice, we have twelve tuples satisfying all conditions except the last one and this last condition gets rid of $(2, 0, 1, 0)$ and $(2, 1, 1, 0)$, hence providing a total of ten solutions. One can check that these solutions are exactly the codes obtained in Figure 2 (bottom elements in each cell of the drawing).

Given two tuples $\mathbf{a} = (a_1, a_2, \dots, a_n)$ and $\mathbf{b} = (b_1, b_2, \dots, b_n)$ we write $\mathbf{a} \leq_{\text{comp}} \mathbf{b}$ if $a_i \leq b_i$ for all $i \in [n]$. We claim in Theorem 1 that the poset $(\mathcal{C}_\alpha, \leq_{\text{comp}})$ is isomorphic to \mathcal{T}_α . For example, one can check, again on Figure 2 that the bottom elements are indeed (partially) ordered by the componentwise order on their tuples.

As a first step towards proving Theorem 1, we associate an integer tuple with each $w \in \mathfrak{S}_\alpha$.

Definition 8. For $w \in \mathfrak{S}_\alpha$, we define its α -code by

$$\text{code}_\alpha(w) \stackrel{\text{def}}{=} (c_1, c_2, \dots, c_n),$$

where

$$c_i \stackrel{\text{def}}{=} \max\{k \mid w_i > w_{s_{\varrho(i)}+1}, w_i > w_{s_{\varrho(i)}+2}, \dots, w_i > w_{s_{\varrho(i)}+k}\}.$$

In other words, c_i counts the number of consecutive entries in the one-line notation of w that are smaller than w_i , starting from the first entry in the α -region immediately after that of i . For $\alpha = (1, 1, \dots, 1)$, Definition 8 agrees with [2, Definition 9.9].

If $\text{code}_\alpha(w) = (c_1, c_2, \dots, c_n)$, then we say that w_i *sees* w_k if $0 < k - s_{\varrho(i)} \leq c_i$. Clearly, if w_i sees w_k , then $(i, k) \in \text{Inv}(w)$, and w_i sees exactly c_i elements for each index i .

In terms of patterns, c_i is the number of 21-patterns where the 2 is at position i that are not 231-patterns. Examples of codes of α -permutations are shown in Table 1 and Figure 2.

3.2 Properties of the encoding

Lemma 9. For $w \in \mathfrak{S}_\alpha$ it holds that $\text{code}_\alpha(w) \in \mathcal{C}_\alpha$.

Proof. Let $w \in \mathfrak{S}_\alpha$ and $\text{code}_\alpha(w) = (c_1, c_2, \dots, c_n)$. Let $i \in [n]$. The maximal number of inversions of the form (i, k) is $n - s_{\varrho(i)}$, because $w_i < w_k$ for all $k \in \{i+1, i+2, \dots, s_{\varrho(i)}\}$.

Hence, $c_i \leq n - s_{\varrho(i)}$, which establishes (C1). If $\varrho(i) = \varrho(i + 1)$, then $w_i < w_{i+1}$ by construction, and thus $c_i \leq c_{i+1}$. This establishes (C2).

Now let $a \in \{\varrho(i)+1, \varrho(i)+2, \dots, r\}$ be such that $c_i \geq s_a - s_{\varrho(i)}$. In particular, w_i sees w_{s_a} , meaning that $w_i > w_{s_a}$. Since w_{s_a} is the rightmost, hence largest, element of its region, w_i also sees any w_j which is seen by w_{s_a} . This implies that $c_i \geq c_{s_a} + s_a - s_{\varrho(i)}$, which is (C3). \square

Theorem 1.1 in [10] establishes that the α -Tamari lattice arises as a quotient lattice of the weak order on \mathfrak{S}_α . This is established by proving that for every $w \in \mathfrak{S}_\alpha$ there exists a unique maximal $(\alpha, 231)$ -avoiding permutation below w in the weak order. The next lemma records this fact.

Lemma 10 ([10, Lemma 3.8]). *For $w \in \mathfrak{S}_\alpha$, the set $\{w' \in \mathfrak{S}_\alpha(231) \mid w' \leq_L w\}$ has a greatest element denoted by $\pi_\alpha^\downarrow(w)$.*

We may thus regard π_α^\downarrow as a map from \mathfrak{S}_α to $\mathfrak{S}_\alpha(231)$. The next lemma characterizes the preimages of this map.

Lemma 11 ([10, Lemma 3.16]). *Let $u, v \in \mathfrak{S}_\alpha$ with $u \leq_L v$. The following are equivalent.*

- (i) *There are indices $i < j < k$, each in different α -regions, such that $v_k < v_i < v_j$, $v_i = v_k + 1$ and $\text{Inv}(v) \setminus \text{Inv}(u) = \{(i, k)\}$,*
- (ii) $\pi_\alpha^\downarrow(u) = \pi_\alpha^\downarrow(v)$.

We now prove that code_α is an order-preserving map from $(\mathfrak{S}_\alpha, \leq_L)$ to $(\mathcal{C}_\alpha, \leq_{\text{comp}})$.

Lemma 12. *Let $u, v \in \mathfrak{S}_\alpha$ with $u \leq_L v$. Then $\text{code}_\alpha(u) \leq_{\text{comp}} \text{code}_\alpha(v)$, and these tuples differ by at most one element. Moreover, $\text{code}_\alpha(u) = \text{code}_\alpha(v)$ if and only if $\pi_\alpha^\downarrow(u) = \pi_\alpha^\downarrow(v)$.*

Proof. Let $u \leq_L v$ and $\text{code}_\alpha(u) = (a_1, a_2, \dots, a_n)$ and $\text{code}_\alpha(v) = (b_1, b_2, \dots, b_n)$.

By assumption, $\text{Inv}(v) \setminus \text{Inv}(u) = \{(i, k)\}$ for some indices $i < k$ in different α -regions such that $v_i = v_k + 1$. It follows that any entry which sees v_k must be greater than v_i , and any entry which does not see v_i must be smaller than v_k . Thus, $a_j = b_j$ for all $j \neq i$. By construction, $u_i = v_k$ and $u_k = v_i$. Since $u_i < u_k$, we conclude that u_i does not see u_k .

If v_i sees v_k , then $a_i < b_i$. This is the case precisely when every j in α -regions strictly between i and k satisfies $v_j < v_i$, which by Lemma 11 means that $\pi_\alpha^\downarrow(u) \neq \pi_\alpha^\downarrow(v)$.

If v_i does not see v_k , then there exists an index j in an α -region strictly between i and k such that $v_i < v_j$, which by Lemma 11 is equivalent to $\pi_\alpha^\downarrow(u) = \pi_\alpha^\downarrow(v)$. If we choose j as small as possible with this property, then any $j' < j$ in an α -region strictly between i and k satisfies $v_i > v_{j'}$, and thus $u_i = v_k > v_{j'} = u_{j'}$, which entails $a_i = b_i$. \square

Corollary 13. *If $u \leq_L v$, then $\text{code}_\alpha(u) \leq_{\text{comp}} \text{code}_\alpha(v)$.*

Proof. This follows from repeated application of Lemma 12. \square

Lemma 14. *Let $u, v \in \mathfrak{S}_\alpha(231)$. If $\text{code}_\alpha(u) \leq_{\text{comp}} \text{code}_\alpha(v)$, then $u \leq_L v$.*

Proof. Let $\text{code}_\alpha(u) = (a_1, a_2, \dots, a_n)$ and $\text{code}_\alpha(v) = (b_1, b_2, \dots, b_n)$ be chosen in such a way that $\text{code}_\alpha(u) \leq_{\text{comp}} \text{code}_\alpha(v)$.

Assume that there exists $(i, k) \in \text{Inv}(u) \setminus \text{Inv}(v)$, and among all these inversions choose (i, k) such that $u_i - u_k$ is minimal. Since (i, k) is not an inversion of v , we have $v_i < v_k$, so that v_i does not see v_k . Since $a_i \leq b_i$ it follows that u_i does not see u_k either. Since $u_i > u_k$, $\varrho(i) < \varrho(k)$ and there exists a smallest index j with $\varrho(i) < \varrho(j) < \varrho(k)$ and $u_i < u_j$. Since $u \in \mathfrak{S}_\alpha(231)$, we have that $u_i > u_k + 1$.

Now, there cannot be any element between $u_k + 1$ and $u_i - 1$ in the same α -region as u_j . Indeed, if this was the case, since $u_j > u_i$, u_{j-1} would be such an element. But, since it is seen by u_i by minimality of j , and since $b_i \geq a_i$, the value v_{j-1} would be seen by v_i , so that $v_k > v_i > v_{j-1}$. In that case, $(j-1, k)$ would be an inversion of u , not an inversion of v and would violate the minimality of (i, k) among such elements as defined earlier. So all elements between u_k and u_i belong to α -regions different from the α -region containing u_j . Thus, among those, there is a smallest one u_ℓ (which is not u_k but can be u_i) that is on the left of u_j . This element belongs to an $(\alpha, 231)$ -pattern in u : (ℓ, j, ℓ') , where ℓ' is the position of $u_\ell - 1$ in u , which is a contradiction.

Therefore, our assumption must have been wrong, and it follows $\text{Inv}(u) \subseteq \text{Inv}(v)$, thus $u \leq_L v$ by definition. \square

Note that we never used in the previous proof that v is $(\alpha, 231)$ -avoiding. This makes sense thanks to Lemma 10: its property has no equivalent going upwards so u and v do not play symmetrical roles.

3.3 Decoding α -codes

We proceed to prove that code_α is a bijection from $\mathfrak{S}_\alpha(231)$ to \mathcal{C}_α .

Lemma 15. *If $w \in \mathfrak{S}_\alpha(231)$, then the leftmost 0 in $\text{code}_\alpha(w)$ corresponds to the position of the 1 in the one-line notation of w .*

Proof. Let $\text{code}_\alpha(w) = (c_1, c_2, \dots, c_n)$, and let $j_o \in [n]$ be such that $w_{j_o} = 1$. Moreover, if $j = \min\{i \mid c_i = 0\}$, then $j \leq j_o$, since $c_{j_o} = 0$. Let $w_j = a$. Since the entries in an α -region are ordered increasingly, a is the smallest element in its α -region.

Now, define $m = \min(w_1, w_2, \dots, w_j)$. Then, if $m \neq a$, since m is strictly to the left of a in w , it cannot be 1 either, so that we have a 231 pattern with the values m , a , and $m - 1$ necessarily in that order in w and in different regions. Otherwise $m = a$. If $m \neq 1$, the region of a cannot be the rightmost region of w since $a - 1$ did not appear in this prefix of w . Let b be the smallest element in the $(\varrho(j) + 1)$ -st α -region, and let $k = s_{\varrho(j)} + 1$, i.e., $w_k = b$. Since $c_j = 0$, we have $a < b$. We then have a 231 pattern with the values a , b , and $a - 1$. So $a = 1$ and $j = j_o$. \square

Proposition 16. *For $\mathbf{c} \in \mathcal{C}_\alpha$ there exists a unique $w \in \mathfrak{S}_\alpha(231)$ such that $\text{code}_\alpha(w) = \mathbf{c}$.*

α	w	$\text{code}_\alpha(w)$
(1)	1	(0)
(2)	12	(0, 0)
(1, 1)	12	(0, 0)
	21	(1, 0)
(3)	123	(0, 0, 0)
(2, 1)	123	(0, 0, 0)
	132	(0, 1, 0)
	231	(1, 1, 0)
(1, 2)	123	(0, 0, 0)
	213	(1, 0, 0)
	312	(2, 0, 0)
(1, 1, 1)	123	(0, 0, 0)
	132	(0, 1, 0)
	213	(1, 0, 0)
	231	(0, 1, 0)
	312	(2, 0, 0)
	321	(2, 1, 0)

Table 1: The α -permutations for any composition α of $n \leq 3$ together with their corresponding α -codes.

Proof. We proceed by induction on n . For $n \leq 3$, the claim can be checked directly (see Table 1), which establishes the induction base. Assume that the claim holds for all compositions of $n' < n$.

Let $\mathbf{c} = (c_1, c_2, \dots, c_n) \in \mathcal{C}_\alpha$. By definition, $c_n = 0$, which enables us to define $j_o = \min\{j \in [n] \mid c_j = 0\}$. By (C2), $j_o = s_{a-1} + 1$ for some $a \in [r]$, meaning that j_o is the first element in the a^{th} α -region.

Let $\alpha' = (\alpha'_1, \alpha'_2, \dots, \alpha'_{r'})$ be the unique composition of $n - 1$ which is obtained by subtracting 1 from α_a . (If $\alpha_a = 1$, then we simply remove this part.) We define $s'_{a'} = \alpha'_1 + \alpha'_2 + \dots + \alpha'_{a'}$, and we obtain

$$s'_b = \begin{cases} s_b, & \text{if } b < a, \\ s_b - 1, & \text{if } b \geq a. \end{cases}$$

We define $\mathbf{c}' = (c'_1, c'_2, \dots, c'_{n-1})$ by setting

$$c'_i = \begin{cases} c_i, & i < j_o \text{ and } c_i < s_{a-1} - s_{\varrho_\alpha(i)}, \\ c_i - 1, & i < j_o \text{ and } c_i \geq s_{a-1} - s_{\varrho_\alpha(i)}, \\ c_{i+1}, & i \geq j_o. \end{cases}$$

It is straightforward to check that $\mathbf{c}' \in \mathcal{C}_{\alpha'}$. By induction hypothesis, there exists a unique $w' \in \mathfrak{S}_{\alpha'}(231)$ with $\text{code}_{\alpha'}(w') = \mathbf{c}'$.

We now “inject” 1 into w' to construct a permutation $w \in \mathfrak{S}_n$ via

$$w_i = \begin{cases} w'_i + 1, & \text{if } i < j_o, \\ 1, & \text{if } i = j_o, \\ w'_{i-1} + 1, & \text{if } i > j_o. \end{cases}$$

Since j_o is the first element in the a^{th} α -region, it follows that $w \in \mathfrak{S}_{\alpha}$. Assume that w has an $(\alpha, 231)$ -pattern (i, j, k) . Since $w' \in \mathfrak{S}_{\alpha'}(231)$, it must be that $k = j_o$, and $w_i = 2$. By construction, $w'_i = 1$, implying that $c'_i = 0$. Since $i < j_o$, it follows that $c_i = 0$, contradicting the choice of j_o . Thus, $w \in \mathfrak{S}_{\alpha}(231)$. By construction, it follows that w is the only $(\alpha, 231)$ -avoiding permutation with $\text{code}_{\alpha}(w) = \mathbf{c}$. \square




Figure 1: Decoding the $(2, 3, 2, 1)$ -code $(2, 6, 0, 1, 3, 1, 1, 0)$. The arrows indicate the leftmost zero in each step; the red digits indicate the positions that see the leftmost zero.

Figure 1 illustrates the procedure described in the proof of Proposition 16. We may now conclude the proof of our first main theorem.

Proof of Theorem 1. Proposition 16 establishes that $\mathfrak{S}_{\alpha}(231)$ and \mathcal{C}_{α} are in bijection, and Corollary 13 and Lemma 14 establish that for $u, v \in \mathfrak{S}_{\alpha}(231)$ we have $u \leq_L v$ if and only if $\text{code}_{\alpha}(u) \leq_{\text{comp}} \text{code}_{\alpha}(v)$. This finishes the proof. \square

In fact, the preimages of the maps $\text{code}_{\alpha}: \mathfrak{S}_{\alpha} \rightarrow \mathcal{C}_{\alpha}$ and $\pi_{\alpha}^{\downarrow}: \mathfrak{S}_{\alpha} \rightarrow \mathfrak{S}_{\alpha}(231)$ coincide.

Lemma 17. *For $u, v \in \mathfrak{S}_{\alpha}$ we have $\text{code}_{\alpha}(u) = \text{code}_{\alpha}(v)$ if and only if $\pi_{\alpha}^{\downarrow}(u) = \pi_{\alpha}^{\downarrow}(v)$.*

Proof. Let $u, v \in \mathfrak{S}_{\alpha}$. Let $\text{code}_{\alpha}(u) = (a_1, a_2, \dots, a_n)$ and $\text{code}_{\alpha}(v) = (b_1, b_2, \dots, b_n)$.

If $u \leq_L v$, then the desired equivalence follows by repeatedly applying Lemma 12.

Otherwise, u and v are incomparable. By [1, Theorem 4.1], $(\mathfrak{S}_\alpha, \leq_L)$ is a lattice and thus the meet $w = u \wedge_L v$ exists and satisfies $w \leq_L u$ and $w \leq_L v$. If $\pi_\alpha^\downarrow(u) = \pi_\alpha^\downarrow(v)$, then $\pi_\alpha^\downarrow(u) = \pi_\alpha^\downarrow(w)$, by Lemma 10. It follows that $\text{code}_\alpha(u) = \text{code}_\alpha(w) = \text{code}_\alpha(v)$ by Lemma 12.

Conversely, let $\text{code}_\alpha(u) = \text{code}_\alpha(v)$. Lemma 10 implies $\pi_\alpha^\downarrow(u) \leq_L u$ and $\pi_\alpha^\downarrow(v) \leq_L v$. In view of the previous reasoning we find $\text{code}_\alpha(\pi_\alpha^\downarrow(u)) = \text{code}_\alpha(u) = \text{code}_\alpha(v) = \text{code}_\alpha(\pi_\alpha^\downarrow(v))$. Proposition 16 thus implies $\pi_\alpha^\downarrow(u) = \pi_\alpha^\downarrow(v)$. \square

Figure 2 shows the weak order on $\mathfrak{S}_{(1,2,1)}$ with the preimages of the map $\pi_{(1,2,1)}^\downarrow$ indicated. The bottom elements in each highlighted region correspond exactly to the $((1, 2, 1), 231)$ -avoiding permutations. The elements are also labeled by their associated $(1, 2, 1)$ -codes.




Figure 2: The weak order on $\mathfrak{S}_{(1,2,1)}$, where the permutations are labeled by their $(1, 2, 1)$ -codes.

4 The α -Tamari lattices and the ν -Tamari lattices

Among other things, [10] introduces a bijection Θ from $\mathfrak{S}_\alpha(231)$ to a certain family of northeast paths, denoted by \mathcal{L}_{ν_α} . This map was simplified in [3] and used to show that the lattice \mathcal{T}_α is isomorphic to a certain lattice on \mathcal{L}_{ν_α} , denoted by \mathcal{T}_{ν_α} [3, Theorem II]. The proof of this result is only partially bijective, and relies on structural properties of both lattices, namely that they are both extremal and have isomorphic Galois graphs.

In the following, we show that our parabolic BW-codes induce a more direct and combinatorial proof of the isomorphism between \mathcal{T}_α and \mathcal{T}_{ν_α} (Theorem 2). We start with

the definitions of related objects, then we will show that our parabolic BW-codes for \mathcal{T}_α are in bijection with ν_α -bracket vectors in \mathcal{T}_{ν_α} , leading to the wanted isomorphism.

4.1 Dyck paths and the ordinary Tamari lattice

We first consider *up steps* (of the form $U \stackrel{\text{def}}{=} (1, 1)$) and *down steps* (of the form $D \stackrel{\text{def}}{=} (1, -1)$). A *Dyck path* of semilength n is a lattice path using only up and down steps, starting and ending on the x -axis and never going below it. Consequently, any Dyck path uses as many up steps as it uses down steps. Let \mathcal{D}_n denote the set of Dyck paths of semilength n . The *ordinate* of a lattice point on a Dyck path is simply the value of its y -coordinate.

If $P \in \mathcal{D}_n$, then any up step U on P has a *matching* down step: this is the first down step D on P whose starting point has the same ordinate as the ending point of U . In particular, the portion of P strictly between U and D is a Dyck path (of strictly smaller semilength) in its own right.

A *valley* of P is a lattice point V on P preceded by a down step and followed by an up step. The *rotation* of P by a valley V is the Dyck path $P' \in \mathcal{D}_n$ obtained by swapping the down step before V and the portion of P (weakly) between the up step after V and its matching down step. The reflexive and transitive closure of this operation yields a partial order on \mathcal{D}_n .

The set of Dyck paths ordered by this *rotation order* forms the (ordinary) *Tamari lattice* \mathcal{T}_n , first described in [14].

Remark 18. It may not be immediately clear from this definition, but the ordinary Tamari lattice is a particular instance of an α -Tamari lattice (see Section 2.2), namely when $\alpha = (1, 1, \dots, 1)$; see [2, Section 9].

4.2 Northeast paths and the ν -Tamari lattice

Now, we consider *north steps* (of the form $N \stackrel{\text{def}}{=} (0, 1)$) and *east steps* (of the form $E \stackrel{\text{def}}{=} (1, 0)$). A *northeast path* of length n is a lattice path using k north and $n - k$ east steps which starts on the x -axis. The *height* of a lattice point on a northeast path is the value of its y -coordinate. A *valley* of a northeast path μ is a lattice point V on μ preceded by an east step and followed by a north step.

Given a northeast path ν , a *ν -path* is a northeast path which shares the starting and ending points with ν and never goes below ν . Let \mathcal{L}_ν denote the set of ν -paths. The *horizontal distance* of a lattice point Q on $\mu \in \mathcal{L}_\nu$, denoted by $\text{horiz}_\nu(Q)$, is the largest number of east steps that can be added to Q without crossing to the other side of ν .

The *rotation* of μ by a valley V is the path $\mu' \in \mathcal{L}_\nu$ obtained by exchanging the east step before V with the portion of μ between V and the next lattice point W on μ satisfying $\text{horiz}_\nu(V) = \text{horiz}_\nu(W)$. In this situation, we write $\mu <_\nu \mu'$. See Figure 3 for an illustration. The reflexive and transitive closure of this operation yields a partial order on \mathcal{L}_ν .




Figure 3: Example of horizontal distance and cover relation in \mathcal{T}_ν for northeast paths.

For any northeast path ν , the set of ν -paths ordered by this *rotation order* forms the *ν -Tamari lattice* \mathcal{T}_ν introduced in [12].

Remark 19. Note that the definition implies that $\mathcal{T}_\nu \cong \mathcal{T}_n$ when $\nu = (NE)^n$. The isomorphism is given by substituting north steps by up steps and east steps by down steps.

4.3 The α -Tamari lattices are certain ν -Tamari lattices

In [4], it was shown that the ν -Tamari lattice can be realized using the componentwise order on so-called ν -bracket vectors. If ν is a northeast path of length n , then the *minimal ν -bracket vector* is the vector \mathbf{b}^{\min} consisting of $n+1$ entries, whose i -th entry is the height of the i -th lattice point on ν . If ν has k north steps, then the *fixed positions* are the entries f_0, f_1, \dots, f_k , where f_i is the position of the last appearance of i in \mathbf{b}^{\min} . An integer vector \mathbf{b} with $n+1$ entries is a *ν -bracket vector*, if it has the following three properties:

- (B1) for $0 \leq s \leq k$, we have $\mathbf{b}(f_s) = s$;
- (B2) for $1 \leq i \leq n+1$, we have $\mathbf{b}^{\min}(i) \leq \mathbf{b}(i) \leq k$;
- (B3) if $\mathbf{b}(i) = s$, then for all j with $i < j < f_s$, we have $\mathbf{b}(j) \leq s$.

The set of ν -bracket vectors is denoted by \mathcal{B}_ν . The following theorem was proven by means of an explicit bijection in [4, Section 4].

Theorem 20 ([4, Theorem 21]). *For any northeast path ν , the ν -Tamari lattice \mathcal{T}_ν is isomorphic to $(\mathcal{B}_\nu, \leq_{\text{comp}})$.*

It was shown in [3, Theorem II], that the α -Tamari lattice is isomorphic to a certain ν -Tamari lattice, namely, when ν is the *α -bounce path*, defined by

$$\nu_\alpha \stackrel{\text{def}}{=} N^{\alpha_1} E^{\alpha_1} N^{\alpha_2} E^{\alpha_2} \dots N^{\alpha_r} E^{\alpha_r}, \quad (1)$$

where $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_r)$. The proof given in [3] is rather indirect and exploits certain lattice-theoretic properties of the α - and the ν -Tamari lattices. In this section, we give a direct proof using the realization of \mathcal{T}_α as the componentwise order on α -codes established in Theorem 1.

In particular, we construct a bijection from the α -codes to the ν_α -bracket vectors. Let us adapt the definitions to the particular case of ν_α . The *minimal ν_α -bracket vector*, denoted by \mathbf{b}_α^{\min} , is defined by

$$\mathbf{b}_\alpha^{\min}(k) \stackrel{\text{def}}{=} \begin{cases} i + s_{a-1} - 1, & \text{if } k = 2s_{a-1} + i \text{ for } 0 < i \leq \alpha_a, \\ s_a, & \text{if } k = 2s_{a-1} + \alpha_a + i \text{ for } 0 < i \leq \alpha_a, \\ n, & \text{if } k = 2n + 1. \end{cases}$$

We write \mathcal{B}_α instead of \mathcal{B}_{ν_α} for the set of ν_α -bracket vectors.

For $\mathbf{b} \in \mathcal{B}_\alpha$, by (B1), there are $n + 1$ positions with fixed value in a vector of length $2n + 1$. For simplification, we define a reduced version of ν_α -bracket vectors. For $\mathbf{b} \in \mathcal{B}_\alpha$, we define its *ν_α -reduced vector* \mathbf{r} by

$$\mathbf{r}(s_{a-1} + i) \stackrel{\text{def}}{=} \mathbf{b}(2s_{a-1} + \alpha_a + i)$$

for $1 \leq i \leq \alpha_a$. It is clear that \mathbf{r} is obtained from \mathbf{b} by removing components whose indices are fixed positions. To recover \mathbf{b} from \mathbf{r} , we only need to fill in the positions of the fixed positions according to (B1). Let Λ_{red} denote the “reduction” map from \mathbf{b} to \mathbf{r} , and let Λ_{ext} be its inverse.

Such ν_α -reduced vectors thus inherit the following properties from ν_α -bracket vectors.

Proposition 21. *A vector $\mathbf{r} \in \mathbb{N}^n$ is a ν_α -reduced vector if, and only if:*

(R1) *for $1 \leq i \leq n$, we have $s_{\varrho(i)} \leq \mathbf{r}(i) \leq n$;*

(R2) *for all i, j with $i < j \leq s_{\varrho(\mathbf{r}(i)+1)-1}$, we have $\mathbf{r}(j) \leq \mathbf{r}(i)$.*

Proof. Let \mathbf{b} be the ν_α -bracket vector corresponding to \mathbf{r} . We only need to show that the conditions for \mathbf{r} are equivalent to those for \mathbf{b} .

Condition (B1) for \mathbf{b} is satisfied by construction. The equivalence between (B2) for \mathbf{b} and (R1) for \mathbf{r} is trivial given the definition of \mathbf{b}_α^{\min} .

Now for the equivalence between (B3) for \mathbf{b} and (R2) for \mathbf{r} , we observe that for (B3) to hold for \mathbf{b} , for each i with $\mathbf{b}(i) = k$, we only need to check for all j with $i < j \leq 2s_{\varrho(k+1)-1}$, since all indices from $2s_{\varrho(k+1)-1} + 1$ to f_k are fixed positions. \square

We can thus take Proposition 21 as the definition of ν_α -reduced vectors without passing through ν_α -bracket vectors, and we denote by \mathcal{R}_α the set of all ν_α -reduced vectors. By Proposition 21, $(\mathcal{R}_\alpha, \leq_{\text{comp}})$ is isomorphic to the ν_α -Tamari lattice. We also have the following property.

Proposition 22. *Given a ν_α -reduced vector \mathbf{r} , for any indices $i < j$ with $\varrho(i) = \varrho(j)$, we have $\mathbf{r}(i) \geq \mathbf{r}(j)$.*

Proof. Let $k = \mathbf{r}(i)$. By (R1), we have $k \geq s_{\varrho(i)}$, and thus $s_{\varrho(k+1)} - 1 \geq s_{\varrho(i)+1} - 1 \geq s_{\varrho(i)}$. Since $\varrho(i) = \varrho(j)$, we have $i < j \leq s_{\varrho(i)} \leq s_{\varrho(k+1)} - 1$. Then (R2) in Proposition 21 states that $\mathbf{r}(j) \leq k = \mathbf{r}(i)$. \square

$$\begin{array}{ccc}
 (2, 6, 0, 1, 3, 1, 1, 0) & \xrightarrow{\Gamma_{\mathcal{R}}} & (8, 4, 8, 6, 5, 8, 8, 8) & \xrightarrow{\Lambda_{\text{ext}}} & (0, 1, 8, 4, 2, 3, 4, 8, 6, 5, 5, 6, 8, 8, 7, 8, 8) \\
 \begin{array}{ccccccc}
 \color{red}{\rule{0.5em}{0.4pt}} & \color{orange}{\rule{1.5em}{0.4pt}} & \color{green}{\rule{0.5em}{0.4pt}} & \color{blue}{\rule{0.2em}{0.4pt}} & & & \\
 \text{m} & & & & & & \\
 \mathcal{C}_{(2,3,2,1)} & & & & & &
 \end{array} & & \begin{array}{ccccccc}
 \color{red}{\rule{0.5em}{0.4pt}} & \color{orange}{\rule{1.5em}{0.4pt}} & \color{green}{\rule{0.5em}{0.4pt}} & \color{blue}{\rule{0.2em}{0.4pt}} & & & \\
 \text{m} & & & & & & \\
 \mathcal{R}_{(2,3,2,1)} & & & & & &
 \end{array} & & \begin{array}{ccccccc}
 \color{red}{\rule{0.5em}{0.4pt}} & \color{orange}{\rule{1.5em}{0.4pt}} & \color{green}{\rule{0.5em}{0.4pt}} & \color{blue}{\rule{0.2em}{0.4pt}} & & & \\
 \text{m} & & & & & & \\
 \mathcal{B}_{(2,3,2,1)} & & & & & &
 \end{array}
 \end{array}$$

Figure 4: Illustration of the map $\Gamma_{\mathcal{R}}$.

$\mathbf{c} \in \mathcal{C}_{(1,2,1)}$	$\Gamma_{\mathcal{R}}(\mathbf{c}) \in \mathcal{R}_{(1,2,1)}$	$\Lambda_{\text{ext}} \circ \Gamma_{\mathcal{R}}(\mathbf{c}) \in \mathcal{B}_{(1,2,1)}$
(0, 0, 0, 0)	(1, 3, 3, 4)	(0, 1, 1, 2, 3, 3, 3, 4, 4)
(1, 0, 0, 0)	(2, 3, 3, 4)	(0, 2, 1, 2, 3, 3, 3, 4, 4)
(0, 0, 1, 0)	(1, 4, 3, 4)	(0, 1, 1, 2, 4, 3, 3, 4, 4)
(2, 0, 0, 0)	(3, 3, 3, 4)	(0, 3, 1, 2, 3, 3, 3, 4, 4)
(1, 0, 1, 0)	(2, 4, 3, 4)	(0, 2, 1, 2, 4, 3, 3, 4, 4)
(0, 1, 1, 0)	(1, 4, 4, 4)	(0, 1, 1, 2, 4, 4, 3, 4, 4)
(3, 0, 0, 0)	(4, 3, 3, 4)	(0, 4, 1, 2, 3, 3, 3, 4, 4)
(3, 0, 1, 0)	(4, 4, 3, 4)	(0, 4, 1, 2, 4, 3, 3, 4, 4)
(1, 1, 1, 0)	(2, 4, 4, 4)	(0, 2, 1, 2, 4, 4, 3, 4, 4)
(3, 1, 1, 0)	(4, 4, 4, 4)	(0, 4, 1, 2, 4, 4, 3, 4, 4)

Table 2: Illustration of the map $\Gamma_{\mathcal{R}}$ for $\alpha = (1, 2, 1)$.

For any composition α , we define a transform $\Delta_{\mathcal{R}}$ on \mathcal{R}_{α} such that

$$(\Delta_{\mathcal{R}}(\mathbf{r}))_i \stackrel{\text{def}}{=} \mathbf{r}(2s_{\varrho(i)} - \alpha_{\varrho(i)} - i + 1) - s_{\varrho(i)}.$$

More intuitively, to obtain $\Delta_{\mathcal{R}}(\mathbf{r})$, we first split \mathbf{r} into regions according to α , then reverse each region while subtracting s_k on the k^{th} region. We denote by $\Gamma_{\mathcal{R}}$ its inverse. Although both $\Delta_{\mathcal{R}}$ and $\Gamma_{\mathcal{R}}$ depend on α , the composition α should always be clear from the context. The transformation $\Delta_{\mathcal{R}}$ relates α -codes with \mathcal{R}_{α} .

Figure 4 illustrates the map $\Gamma_{\mathcal{R}}$ on the $(2, 3, 2, 1)$ -code $(2, 6, 0, 1, 3, 1, 1, 0)$, and Table 2 illustrates this bijection for $\alpha = (1, 2, 1)$.

Proposition 23. *Given a composition α of n , the transformation $\Delta_{\mathcal{R}}$ is a bijection from \mathcal{R}_{α} to \mathcal{C}_{α} .*

Proof. First, for $\mathbf{r} \in \mathcal{R}_{\alpha}$, let $\mathbf{c} = \Delta_{\mathcal{R}}(\mathbf{r})$ and let us check that \mathbf{c} satisfies the conditions in Definition 6 using those in Proposition 21 for \mathbf{r} . By (R1) for \mathbf{r} and the definition of $\Delta_{\mathcal{R}}$, clearly \mathbf{c} satisfies (C1). Proposition 22 and the definition of $\Delta_{\mathcal{R}}$ imply that \mathbf{c} satisfies (C2). To check (C3) for \mathbf{c} given that it satisfies (C2), we only need to show that, for any i and j such that $\varrho(i) < \varrho(j)$, if $c_i \geq s_{\varrho(j)} - s_{\varrho(i)}$, then we have $c_j + s_{\varrho(j)} \leq c_i + s_{\varrho(i)}$. Translating to \mathbf{r} , we need to check that, for any i' and j' with $\varrho(i') < \varrho(j')$, if $\mathbf{r}(i') \geq s_{\varrho(j')}$, then we have $\mathbf{r}(j') \leq \mathbf{r}(i')$. Now, suppose that $\mathbf{r}(i') \geq s_{\varrho(j')}$. We have $\varrho(\mathbf{r}(i') + 1) > \varrho(j')$ by the definition of ϱ . As the values are integers, we have $\varrho(j') \leq \varrho(\mathbf{r}(i') + 1) - 1$, which means $j' \leq s_{\varrho(j')} \leq s_{\varrho(\mathbf{r}(i') + 1) - 1}$, and by (R2), we have $\mathbf{r}(j') \leq \mathbf{r}(i')$. Therefore, \mathbf{c} also satisfies (C3).

Now for the reverse direction, given $\mathbf{c} \in \mathcal{C}_\alpha$, let $\mathbf{r} = \Gamma_{\mathcal{R}}(\mathbf{c})$. It is clear that (C1) translates directly to (R1). We only need to show that (R2) holds for \mathbf{r} . Suppose that $1 \leq i < j \leq s_{\varrho(\mathbf{r}(i)+1)-1}$. If $\varrho(i) = \varrho(j)$, by the definition of $\Gamma_{\mathcal{R}}$ and (C2) on \mathbf{c} , we have $\mathbf{r}(j) \leq \mathbf{r}(i)$. Now we check the case $\varrho(i) < \varrho(j)$. When translated to \mathbf{c} , (R2) in this case means that we need to check for any $i' < j'$ such that $\varrho(i') < \varrho(j') \leq \varrho(c_{i'} + s_{\varrho(i')} + 1) - 1$, we have $c_{j'} + s_{\varrho(j')} \leq c_{i'} + s_{\varrho(i')}$. By (C2), we may assume that $j' = s_a$ for some a . By the definition of ϱ , we see that $\varrho(s_a) \leq \varrho(c_{i'} + s_{\varrho(i')} + 1) - 1$ implies $s_a < c_{i'} + s_{\varrho(i')} + 1$, thus $s_a \leq c_{i'} + s_{\varrho(i')}$ since they are integers. By (C3), we have $c_{s_a} \leq c_{i'} - s_a + s_{\varrho(i')}$. Therefore, (R2) holds for \mathbf{r} , meaning that $\mathbf{r} \in \mathcal{R}_\alpha$. \square

This allows us to conclude to the announced simple proof of Theorem 2.

Proof of Theorem 2. Let \mathcal{T}_{ν_α} denote the ν_α -Tamari lattice. We have the following isomorphisms of lattices:

$$\mathcal{T}_\alpha \stackrel{\text{Thm. 1}}{\cong} (\mathcal{C}_\alpha, \leq_{\text{comp}}) \stackrel{\text{Prop. 23}}{\cong} (\mathcal{R}_\alpha, \leq_{\text{comp}}) \stackrel{\text{trivial}}{\cong} (\mathcal{B}_\alpha, \leq_{\text{comp}}) \stackrel{\text{Thm. 20}}{\cong} \mathcal{T}_{\nu_\alpha}. \quad \square$$

Note that the proof of Theorem 2 in [3] relies on lattice-theoretic properties of \mathcal{T}_α and \mathcal{T}_{ν_α} , and is only partially bijective. Our proof here is fully bijective, which gives a clearer vision of the isomorphism.

5 A combinatorial anti-isomorphism on the ν_α -Tamari lattice

As mentioned at the beginning of Section 4, the original proof of Theorem 2 used the bijection Θ first defined in [10], but only to show that it extends to an isomorphism between the Galois graphs of \mathcal{T}_α and \mathcal{T}_{ν_α} . It is then natural to ask whether Θ extends to all elements of \mathcal{T}_α and \mathcal{T}_{ν_α} , and this question was stated in [3] as Open Problem 2.23. Comparing with the bijection in our proof of Theorem 2 from \mathcal{T}_α to \mathcal{T}_{ν_α} , we may wonder if it is equivalent in some sense to Θ .

We now provide a direct combinatorial interpretation of this equivalence, using our parabolic BW-codes and constructions from [3]. This settles [3, Open Problem 2.23]. More precisely, using a stack-processing procedure and the α -code for \mathcal{T}_α , we will show that Θ is in fact an anti-isomorphism from \mathcal{T}_α to $\mathcal{T}_{\text{Flip}(\nu_\alpha)}$, where $\text{Flip}(\nu_\alpha)$ is essentially the northeast path ν_α read backwards. To this end, we will need to consider Dyck paths with certain sequence statistics, which are then used to prove that our α -codes are closely related to the bijection Θ .

5.1 ν -Tamari lattices are intervals of ordinary Tamari lattices

It was shown in [12, Theorem 3] in terms of binary trees that every ν -Tamari lattice is isomorphic to an interval in some ordinary Tamari lattice. This result was obtained through a bijection in [12, Section 2 and 3] between binary trees and pairs of non-crossing lattice paths, which is equivalent to the bijection β in [5] between Dyck paths and parallelogram




Figure 5: An example of the bijection **Dyck**.

polyominoes (see [5, Section 4]). We now restate the bijection β for our need, based on the definition of its inverse defined in [5, Section 4].

A northeast path is uniquely determined by the lengths of the *runs* of east steps at each height. In other words, if ν is a northeast path, then we can write it uniquely as

$$\nu = E^{a_0} N E^{a_1} N \dots E^{a_{k-1}} N E^{a_k}.$$

By abuse of notation we will also write $\nu = [a_0, a_1, \dots, a_k]$. Now, if $\mu \in \mathcal{L}_\nu$ with $\mu = [b_0, b_1, \dots, b_k]$, then we have $\sum_{i=0}^k a_i = \sum_{i=0}^k b_i = m$ and $\sum_{i=0}^j a_i \geq \sum_{i=0}^j b_i$ for all $0 \leq j \leq k$. We may have $a_i = 0$ or $b_i = 0$ for some indices i .

Construction 24. Given a northeast path ν composed of k north steps and $n - k$ east steps, and for any $\mu \in \mathcal{L}_\nu$, we write ν and μ under the form $\nu = [a_0, a_1, \dots, a_k]$ and $\mu = [b_0, b_1, \dots, b_k]$. We define the Dyck path $\mathbf{Dyck}(\nu, \mu)$ of semilength $n + 1$ by

$$\mathbf{Dyck}(\nu, \mu) \stackrel{\text{def}}{=} U^{a_0+1} D^{b_0+1} \dots U^{a_k+1} D^{b_k+1}.$$

Conversely, if $P \in \mathcal{D}_{n+1}$ we can recover the pair (ν, μ) satisfying $P = \mathbf{Dyck}(\nu, \mu)$ by looking at the lengths of the runs of up and down steps in P , which determine the a_i 's and b_i 's, respectively.

The map **Dyck** is a bijection from the set

$$\{(\nu, \mu) \mid \nu \text{ has length } n \text{ and } \mu \in \mathcal{L}_\nu\}$$

to \mathcal{D}_{n+1} , and is illustrated in Figure 5. It was proved in [12], in terms of binary trees, that **Dyck** is an order isomorphism.

Proposition 25 ([12, Theorem 3]). *Given a northeast path ν of length n , the map **Dyck** is an isomorphism from \mathcal{T}_ν to an interval I_ν of \mathcal{T}_{n+1} .*

5.2 Two sequence statistics of Dyck paths

In preparation of things to come, we now consider an anti-isomorphism **Conj** of \mathcal{T}_n in terms of Dyck paths, first defined by Deutsch in [6], that exchanges two particular sequence statistics on Dyck paths. The exchange property can be deduced from the recent work

of Pons on Tamari interval posets [11]; see the paragraph after Theorem 23 therein. We translate the appropriate specialization of her result to the framework of Dyck paths. We then relate **Conj** and these statistics to ν -Tamari lattices via the map **Dyck**.

A *rising contact* of a Dyck path P is an up step in P that starts on the x -axis. Every Dyck path P of length $n > 0$ can be uniquely decomposed into $P = P_\ell UP_r D$ with P_ℓ, P_r both Dyck paths, by taking P_ℓ to be the sub-path before the last rising contact of P . We denote by ϵ the empty Dyck path, and we define an involution **Conj** recursively by

$$\mathbf{Conj}(\epsilon) \stackrel{\text{def}}{=} \epsilon, \quad \mathbf{Conj}(P_\ell UP_r D) \stackrel{\text{def}}{=} \mathbf{Conj}(P_r) U \mathbf{Conj}(P_\ell) D. \quad (2)$$

The following is well-known and can be proven by induction using the decomposition $P = P_\ell UP_r D$.

Proposition 26. *The involution **Conj** is an anti-isomorphism of \mathcal{T}_n .*

We define $\mathbf{Conj}' \stackrel{\text{def}}{=} \mathbf{Dyck}^{-1} \circ \mathbf{Conj} \circ \mathbf{Dyck}$, which is simply **Conj** conjugated to the domain of ν -paths using **Dyck**. See Figure 6 for an example of **Conj** and **Conj'**.

Given a northeast path ν , we denote by **Flip**(ν) the northeast path obtained by reversing ν and exchanging north and east steps. Geometrically, **Flip**(ν) is ν reflected across a diagonal of slope -1 . It is known that, for $(\nu', \mu') = \mathbf{Conj}'(\nu, \mu)$, we have $\nu = \mathbf{Flip}(\nu')$ (see [12, Theorem 2 and 3]). We have the following corollary.

Corollary 27. *The bijection \mathbf{Conj}' is an anti-isomorphism between \mathcal{T}_ν and $\mathcal{T}_{\mathbf{Flip}(\nu)}$.*

We now consider two statistics on Dyck paths that are interchanged by **Conj**. Given a Dyck path P , we define its *descent run sequence*, denoted by **Drun**(P) and *indexed from 0 to n* , as follows. We write $P = D^{c_0} U D^{c_1} U \dots U D^{c_n}$ with some $c_i \geq 0$ (again, noting $c_0 = 0$), then we take $\mathbf{Drun}(P)_i = c_{n-i}$ for all $0 \leq i \leq n$. The map **Drun** is injective because P can be reconstructed from the c_i 's.

We now define another sequence statistic. The *contact sequence* of P , denoted by **Cont**(P) and *indexed from 0 to n* , is obtained by taking **Cont**(P)₀ the number of rising contacts of P , and **Cont**(P) _{i} the number of rising contacts of the sub-Dyck path strictly between the i -th up step and its matching down step.

Examples for **Cont** and **Drun** are given in Figure 6. The following result, also illustrated in Figure 6, is well-known and can be proven inductively. In terms of binary trees, **Cont** (resp. **Drun**) describes maximal left (resp. right) descending paths, and the counterpart of **Conj** on binary trees is taking the vertical mirror image.

Proposition 28 (See [11]). *For any Dyck path P , we have $\mathbf{Cont}(P) = \mathbf{Drun}(\mathbf{Conj}(P))$.*

As **Dyck** is bijective, let $(\nu, \mu) = \mathbf{Dyck}^{-1}(P)$, and define $\mathbf{Drun}(\nu, \mu) = \mathbf{Drun}(P)$ and $\mathbf{Cont}(\nu, \mu) = \mathbf{Cont}(P)$. Suppose that $\nu = [a_0, a_1, \dots, a_k]$ and $\mu = [b_0, b_1, \dots, b_k]$. Then we have

$$\mathbf{Drun}(\nu, \mu) = \mathbf{Drun}(P) = (b_k + 1, 0^{a_k}, b_{k-1} + 1, 0^{a_{k-1}}, \dots, b_0 + 1, 0^{a_0}, 0). \quad (3)$$




Figure 6: Example of the bijections \mathbf{Conj} and \mathbf{Conj}' and how they transform the statistics \mathbf{Cont} and \mathbf{Drun} . The reverse horizontal distances of points on the lower paths are also given.

Here “ 0^a ” means the entry 0 repeated a times. For an expression of \mathbf{Cont} , we need some more definitions. We define the *reverse horizontal distance* to μ of a lattice point Q on ν , denoted by $\text{horiz}'_{\mu}(Q)$, to be the number of west steps $(-1, 0)$ we can take from Q before crossing to the other side of μ . See Figure 6 for an example. This figure also illustrates the following result.

Proposition 29. *Let $P = \mathbf{Dyck}(\nu, \mu)$ with ν, μ northeast paths of length n . Take the sequence $(d_i)_{0 \leq i \leq n}$ with $d_i = \text{horiz}'_{\mu}(Q_i)$, where Q_i is the $(i + 1)$ -st lattice point of μ . Then we have*

$$\begin{aligned} \mathbf{Cont}(\nu, \mu)_0 &= |\{\ell \mid 0 \leq \ell \leq n, d_{\ell} = 0\}|; \\ \mathbf{Cont}(\nu, \mu)_i &= |\{\ell \mid i < \ell \leq n, d_{\ell} = d_i + 1, \forall i < m \leq \ell, d_m > d_i\}|. \end{aligned} \tag{4}$$

Proof. Assume that $\nu = [a_0, a_1, \dots, a_k]$ and $\mu = [b_0, b_1, \dots, b_k]$. For each $0 \leq i \leq k$, there are $a_i + 1$ lattice points of height i on ν , and the $(i + 1)$ -st consecutive run of up steps of P consists of $a_i + 1$ steps. According to the construction of \mathbf{Dyck} , the $(i + 1)$ -st up step in P corresponds to a lattice point Q_i on ν . Moreover, Q_i is the leftmost point with height j on ν if and only if the corresponding up step is the first in the j -th run of up steps in P .

We now prove that d_i is the ordinate of the starting point of the $(i + 1)$ -st up step in P . We proceed by induction. For $i = 0$, we have $d_0 = 0 = \text{horiz}'_{\mu}(Q_0)$, as ν and μ start at the origin. Now suppose that d_{i-1} is the ordinate of the starting point of the i -th up step in P . We have two cases.

- (i) If Q_i is not the leftmost lattice point with its height, then its corresponding up step in P directly comes after the one associated with Q_{i-1} . Thus, the ordinate of the starting point of this up step is $d_{i-1} + 1 = d_i$.
- (ii) If Q_i is the leftmost lattice point with height ℓ , then its corresponding up step is the first up step of the next run after the up step associated with Q_{i-1} . The ordinate of the up step associated with Q_i is thus $d_{i-1} + 1 - (b_{\ell} + 1)$, taking into account the length of the ℓ -th run of down steps in P . This is also equal to d_i , as we reach Q_i from Q_{i-1} by a north step, and it takes b_{ℓ} less west steps to cross μ from Q_i than from Q_{i-1} .

We thus conclude the induction. Now (4) is a translation of the definition of **Cont** in terms of the d_i 's. □

5.3 Two ways from $(\alpha, 231)$ -avoiding permutations to α -paths

Recall that we have fixed a composition $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_r)$ of n , and that $s_a = \alpha_1 + \alpha_2 + \dots + \alpha_a$ for $a \in [r]$. Moreover, recall the definition of the α -bounce path from (1). We usually say α -path rather than ν_{α} -path.

We now define two bijections from $(\alpha, 231)$ -avoiding permutations to α -paths. The first one uses the α -code from Section 3.1, and sends $w \in \mathfrak{S}_{\alpha}(231)$ to $\varphi(w) \in \mathcal{L}_{\nu_{\alpha}}$ satisfying $\varphi(w) = [f_0, f_1, \dots, f_n]$, where

$$f_i \stackrel{\text{def}}{=} \left| \{j \mid 1 \leq j \leq n, \text{code}_{\alpha}(w)_j + s_{\ell(j)} = i\} \right|. \tag{5}$$

For example, for $\alpha = (1, 3, 1, 2)$ and $w = \mathbf{5347126} \in \mathfrak{S}_{\alpha}(231)$, we have $\text{code}_{\alpha}(w) = (2, 2, 2, 3, 0, 0, 0)$ and $\varphi(w) = [0, 0, 0, 1, 0, 1, 2, 3]$. Note that the first entry of $\varphi(w)$ is always 0.

Proposition 30. *The map φ is an isomorphism from \mathcal{T}_{α} to $\mathcal{T}_{\nu_{\alpha}}$.*

Proof. We write $\varphi(w) = [f_0, f_1, \dots, f_n]$. By the definition of $\Delta_{\mathcal{R}}$, we have

$$f_i = \left| \{j \mid 1 \leq j \leq n, \Delta_{\mathcal{R}}^{-1}(\text{code}_{\alpha}(w))_j = i\} \right|.$$

We may rephrase this using ν_{α} -reduced vectors. Let $\mathbf{r} = \Gamma_{\mathcal{R}}(\text{code}_{\alpha}(w))$ denote the ν_{α} -reduced bracket vector associated with w . Then,

$$f_i = \left| \{j \mid 1 \leq j \leq n, \mathbf{r}_j = i\} \right| - 1.$$

Now, if $\mathbf{b} = \Lambda_{\text{ext}}(\mathbf{r})$ is the associated ν_α -bracket vector, then the number of entries equal to i in \mathbf{b} is $f_i + 1$. According to [4, Definition 26], there exists a unique ν_α -path with as many lattice points of height i as there are entries equal to i in \mathbf{b} . (See also item (ii) in the proof of [4, Proposition 27].) We conclude that the ν_α -path associated with \mathbf{b} is precisely $\varphi(w)$.

Therefore, φ is precisely the isomorphism used in the proof of Theorem 2. \square

The second bijection, denoted by Θ , from $(\alpha, 231)$ -avoiding permutations to α -paths was first defined in [10]. We will use an equivalent definition derived from [3], using a family of trees called α -trees, which we will not explicitly define.

Construction 31. Given $w \in \mathfrak{S}_\alpha(231)$, we construct a labeled plane tree $T(w)$ by an insertion procedure. We start with a node labeled $n + 1$ as the root, and we read the elements of w from left to right. Upon reading of an element $w(i)$, we start a walk from the root. When we reach a node v with label ℓ , if $w(i) < \ell$, then we move to the left-most child of v ; otherwise, we move to the first sibling of v on its right. When the destination node does not exist, we add it with label $w(i)$ and terminate the walk. The labeled plane tree thus obtained is denoted by $T(w)$.

Now we construct a northeast path P from $T(w)$. If the root of $T(w)$ has k children, then we start P with k north steps. Then, for each $a \in [r]$, we inspect the elements $w(i)$ in the a -th α -region from right to left, i.e., i runs from s_a down to $s_{a-1} + 1$. For each such $w(i)$, let v_i be the node with label $w(i)$ in $T(w)$, and we append EN^{k_i} to P , where k_i is the number of children of v_i . We define $\Theta(w)$ to be the path P thus obtained. See Figure 7 for an example.




Figure 7: Example of $\Theta(w)$ for $w \in \mathfrak{S}_\alpha(231)$ with $\alpha = (1, 3, 1, 2, 4, 3)$.

For $w \in \mathfrak{S}_\alpha(231)$, the tree $T(w)$ from Construction 31 has the following immediate property.

Proposition 32 ([3]). For $w \in \mathfrak{S}_\alpha(231)$, let $T(w)$ be the labeled plane tree constructed in Construction 31. Reading the labels of $T(w)$ in postorder (i.e., for each node u , the children of u are increasing in order from left to right, all greater than u) gives $1, 2, \dots, n$.

Remark 33. The map Θ in Construction 31 is in fact $\Theta^{-1} \circ \mathbf{Flip}$ in [3]. We have altered the definition here for simplicity. In Construction 31, the tree $T(w)$ is a labeled version of an α -tree, and the map from w to $T(w)$ is the map Λ_{perm} in [3]. Moreover, the map from $T(w)$ to $\Theta(w)$ is Ξ_{nn} in that article, but our definition here is adapted from Lemma 1.31 of the same article. The validity of our definition of Θ is ensured by [3, Propositions 1.33 and 1.34]. The property in Proposition 32 follows from [3, Construction 1.14].

5.4 A stack-processing procedure

We now give another combinatorial definition of φ . For $w \in \mathfrak{S}_\alpha(231)$, we define the *companion* of an element $w(i)$ in w to be the last element it sees, or $w(s_{\varrho(i)})$ when $w(i)$ sees no element. Then we can define $\varphi(w) = [f_0, f_1, \dots, f_n]$, where f_i is the number of elements in w with $w(i)$ as its companion. We check that this definition of f_i is the same as (5). We define the following *stack processing* that can be used to compute the companions of elements of w .

Construction 34. Given $w \in \mathfrak{S}_\alpha(231)$, we start with an empty stack S and then perform the following steps on the α -regions *in reverse order*, i.e., k runs from r down to 1.

- **(Popping)** For i from 1 to α_k , consider the i -th element $w(s_{k-1} + i)$ in region k . Pop elements from the stack until the top one is larger than $w(s_{k-1} + i)$.
- **(Pushing)** For i from 1 to α_k , push the element $w(s_k - i + 1)$ into the stack.

There are n elements in w , and each element passes through two steps, totaling to $2n$ steps. See Figure 8 for an example.

Remark 35. Note that in terms of popping elements we only need the popping step for the last element in each region, as it is also the largest. However, taking the popping step for each element into account is important to understand the link between φ and Θ . Namely, given $w \in \mathfrak{S}_\alpha(231)$, the number of elements popped out in the popping step of $w(i)$ is the number of children of the node with label $w(i)$ in $T(w)$ (Proposition 38), which is in turn the length of the corresponding run of north steps in $\Theta(w)$.

Lemma 36. For $w \in \mathfrak{S}_\alpha(231)$, at each step of the stack processing of w with the stack S , we have:

- (i) the elements of S are increasing from top to bottom;
- (ii) for elements in S , their indices in w are increasing from top to bottom.

Proof. For the first point, we proceed by induction on the number of steps. The claim is clearly satisfied at the beginning, when S is empty. The popping step maintains the claim.




Figure 8: An example of stack processing for $w = 5347126 \in \mathfrak{S}_\alpha(231)$ with $\alpha = (1, 3, 1, 2)$.

Suppose that we are now pushing an element $w(i)$ into S . If $w(i)$ is the last element in its region, then by induction hypothesis, all elements smaller than $w(i)$ should have been popped out; otherwise, the top of the stack is $w(i+1) > w(i)$. In both cases, pushing $w(i)$ maintains our claim. We thus conclude the induction.

For the second point, we observe that the claim is valid for the empty stack, and the pushing steps maintain the claim, since all the elements in the same region are pushed consecutively starting with the last element in the region. The popping steps clearly also maintains the claim. \square

Proposition 37. *For $w \in \mathfrak{S}_\alpha(231)$, we consider the popping step of an element $w(i)$ in the stack processing of w with stack S . If after that popping step S is not empty with top element $w(j)$, then the companion of $w(i)$ is $w(j-1)$; if S is empty, then the companion of $w(i)$ is $w(n)$.*

Proof. Assume that there is an index ℓ such that $s_{\varrho(i)} < \ell < j$ and $w(\ell) > w(i)$. We take the smallest such index ℓ . Then $w(\ell)$ cannot be popped out of S before the treatment of the region $\varrho(i)$, since an element $w(\ell')$ that pops $w(\ell)$ out must have $s_{\varrho(i)} < \ell' < \ell$ and $w(\ell') > w(\ell) > w(i)$, violating the minimality of ℓ . At this moment, $w(j)$ is also in S . By Lemma 36(ii), $w(j)$ is below $w(\ell)$. Since $w(j)$ is on top of S after the popping step of $w(i)$, there is some $w(i')$ in the same region of i with $i' < i$ that popped $w(\ell)$ out, meaning that $w(\ell) < w(i') < w(i)$, contradicting our hypothesis. Therefore, such $w(\ell)$ does not exist. From the definition of the popping step, we have $w(j) > w(i)$. Thus, the companion of $w(i)$ is $w(j-1)$. In the case of empty stack, it means that no element in previously inserted regions is larger than $w(i)$, thus the companion of $w(i)$ is $w(n)$. \square

5.5 Stack processing and the bijection Θ

We now describe a link between stack processing of $w \in \mathfrak{S}_\alpha(231)$ and the tree $T(w)$ in Construction 31. Given $1 \leq k \leq r$, the *nodes of region k* of $T(w)$ are those with labels corresponding to the values of w in the k -th α -region. We say that the root is of region 0, and that the region $r + 1$ is empty. From the insertion procedure, for a node of region k with its parent of region k' , we have $k > k'$. The *active nodes* for region k are nodes of region $k' \geq k$ whose parent is of region $k'' < k$.

Proposition 38. *For $w \in \mathfrak{S}_\alpha(231)$, consider the stack processing of w with stack S . For $1 \leq k \leq r + 1$, the elements in S after processing the k -th α -region are exactly the labels of the active nodes for region k in $T(w)$.*

Furthermore, for a node u in $T(w)$, the labels of its children are exactly the elements popped out by the label of u in the stack processing of w .

Proof. Let E_k be the set of elements in S after processing the k -th α -region, and let L_k be the set of labels of the active nodes of region k in $T(w)$. We show that $E_k = L_k$. We proceed by induction on k from $r + 1$ to 1. For $k = r + 1$, the set E_{r+1} is empty, and there are no active nodes for region $r + 1$, so we have $E_{r+1} = L_{r+1}$. Suppose that $E_{k+1} = L_{k+1}$. We observe that, by Construction 34 and Lemma 36(i), we have

$$E_k = (E_{k+1} \setminus R_k) \cup \{w(i) \mid \varrho(i) = k\}, \quad (6)$$

where $R_k = \{w(i) \mid \varrho(i) > k, w(i) < w(s_k)\}$.

Now, by the definition of active nodes, we split L_k into two parts, $L_k^{(1)}$ for nodes of region k , and $L_k^{(2)}$ for other nodes. It is clear that $L_k^{(1)} = \{w(i) \mid \varrho(i) = k\}$. A node with label in $L_k^{(2)}$ must be in some region $k' > k$, and its parent in some other region $k'' < k < k + 1$. Therefore, $L_k^{(2)} \subseteq L_{k+1}$. Conversely, let u be a node with labels in L_{k+1} , and let v be the parent of u in region k_v . If $k_v < k$, then u is also in $L_k^{(2)}$; otherwise, if $k_v = k$, then u is not in $L_k^{(2)}$. We thus have

$$L_k = (L_{k+1} \setminus R'_k) \cup \{w(i) \mid \varrho(i) = k\}. \quad (7)$$

Here, R'_k is the set of labels of nodes whose parents are of region k .

Now, by the construction of $T(w)$ in Construction 31, labels in R'_k must be in some region $k' > k$, and they are smaller than some element in region k of w , therefore smaller than $w(s_k)$. We thus have $R'_k \subseteq R_k$. Conversely, suppose that $R_k \setminus R'_k$ is not empty, and let $w(i) \in R_k \setminus R'_k$ and u the node in $T(w)$ with $w(i)$ as label. We have $\varrho(i) > k$ and $w(i) < w(s_k)$. Let v be the parent of u in $T(w)$, and $w(j)$ the label of v . As $w(i) \in R'_k$, we know that v is in some region $k_v < k$. Suppose that v' is the node with label $w(s_k)$. As $w(i) < w(s_k)$, by Proposition 32, u precedes v' in postorder. By the construction of $T(w)$, we know that the region of a node is always strictly smaller than that of its children, and weakly smaller than that of its siblings to the right. Therefore, the node v of region $k_v < k$ must not be a descendant of v' of region k . If v' is a descendant of v , as u precedes v' in postorder, meaning that v' must be a sibling of u to the right, or a descendant of

such a sibling, which is impossible because u is in region $\varrho(i) > k$. Therefore, v' and v are not comparable in $T(w)$, and along with the fact that u precedes v' in postorder, v also precedes v' in postorder. Now take the rightmost child of v , say u' with label $w(i')$, which also precedes v' in postorder. We have $w(j) = w(i') + 1$ and $w(s_k) > w(j)$ by Proposition 32. Furthermore, v is in region $k_v < k$, while v' is in region k and u' is in a region $k_{u'} \geq \varrho(i) > k$. We thus have an $(\alpha, 231)$ -pattern $w(j), w(s_k), w(i')$ in w , which is not possible. Therefore, $w(i)$ cannot exist, and we have $R'_k = R_k$.

Comparing (6) and (7), along with $R'_k = R_k$ and the induction hypothesis $E_{k+1} = L_{k+1}$, we have $E_k = L_k$, concluding the induction. Therefore, the first part of our claim holds for all $1 \leq k \leq r + 1$.

For the second part, let $w(i)$ be the label of u and $w(j)$ an element popped out by $w(i)$ in the stack processing, and v the node with $w(j)$ as label. Suppose that u is of region k_u . By the first part of our claim, v is an active node for region $k_u - 1$ but not for region k_u . Therefore, the parent of v is of region k_u . By Proposition 32, the label of the parent of v must be the first element in region k_u larger than $w(j)$, which is $w(i)$ according to the popping step of region k_u . Thus, u is the parent of v , and we have the second part of our claim. \square

We now prove that the isomorphism φ from \mathcal{T}_α to \mathcal{T}_{ν_α} is closely related to Θ defined in [3]; see Section 5.3

Theorem 39. *Let $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_r)$. For $w \in \mathfrak{S}_\alpha(231)$, we have*

$$(\nu_\alpha, \varphi(w)) = \mathbf{Conj}'(\mathbf{Flip}(\nu_\alpha), \mathbf{Flip}(\Theta(w))).$$

Proof. By definition of \mathbf{Conj}' and Proposition 28, we only need to show that

$$\mathbf{Drun}(\nu_\alpha, \varphi(w)) = \mathbf{Cont}(\mathbf{Flip}(\nu_\alpha), \mathbf{Flip}(\Theta(w))),$$

as \mathbf{Drun} is injective. For the pair $(\nu_\alpha, \varphi(w))$, we observe that

$$\nu_\alpha = [0, 0^{\alpha_1-1}, \alpha_1, 0^{\alpha_2-1}, \alpha_2, \dots, 0^{\alpha_r-1}, \alpha_r],$$

with 0^k standing for k entries of 0. Suppose that $\varphi(w) = [f_0, f_1, \dots, f_n]$. We have

$$\mathbf{Drun}(\nu_\alpha, \varphi(w)) = (f_{s_r} + 1, 0^{\alpha_r}, f_{s_{r-1}} + 1, f_{s_{r-2}} + 1, \dots, f_{s_{r-1}+1} + 1, 0^{\alpha_{r-1}}, \dots, f_{s_1-1} + 1, \dots, f_0 + 1, 0^{\alpha_1}, 0). \quad (8)$$

Now for the pair $(\nu_\alpha, \Theta(w))$, for $0 \leq i \leq 2n$, let Q'_i the $(i + 1)$ -st lattice point Q'_i on ν_α in reverse order. We define d'_i to be the number of north steps $(0, 1)$ we can take from Q'_i without crossing to the other side of $\Theta(w)$. It is clear that d'_i is also the reverse horizontal distance of the $(i + 1)$ -st lattice point of $\mathbf{Flip}(\nu_\alpha)$ with respect to $\mathbf{Flip}(\Theta(w))$. By Proposition 29,

$$\mathbf{Cont}(\mathbf{Flip}(\nu_\alpha), \mathbf{Flip}(\Theta(w)))_0 = |\{\ell \mid 1 \leq \ell \leq 2n, d'_\ell = 0\}|;$$

$$\mathbf{Cont}\left(\mathbf{Flip}(\nu_\alpha), \mathbf{Flip}(\Theta(w))\right)_i = |\{\ell \mid i < \ell \leq 2n, d'_\ell = d'_i + 1, \forall i < m \leq \ell, d'_m > d'_i\}|.$$

Consider the stack processing of w with stack S . We now show that the number of elements in the stack after i steps of stack processing is d'_i . We proceed by induction on the number of steps we have taken in the stack processing. In the initial stage, $d'_0 = 0$ agrees with the empty stack. When dealing with region k , we first perform the popping step. By the construction of $\Theta(w)$, for the i -th element in region k , the number of children of its correspondent node, which is also the number elements popped out by $w(s_{k-1} + i)$ by Proposition 38, is the number of north steps of $\Theta(w)$ on abscissa $s_k - i + 1$, which is exactly $d'_{2n-2s_k+i-1} - d'_{2n-2s_k+i}$. Then for the pushing step, the stack size increases by 1 at each step, just as when we pass from d'_{2n-s_k+i-1} to d'_{2n-s_k+i} . We thus conclude the induction.

We now show that $\mathbf{Drun}(\nu_\alpha, \varphi(w)) = \mathbf{Cont}\left(\mathbf{Flip}(\nu_\alpha), \mathbf{Flip}(\Theta(w))\right)$. First, we know that d'_i is weakly decreasing for i from $2(n - s_k) + 1$ to $2(n - s_k) + \alpha_k$ for all $1 \leq k \leq r$, and by definition, $\mathbf{Cont}\left(\mathbf{Flip}(\nu_\alpha), \mathbf{Flip}(\Theta(w))\right)$ takes the form

$$(g_{s_r}, 0^{\alpha_r}, g_{s_r-1}, g_{s_r-2}, \dots, g_{s_{r-1}}, 0^{\alpha_{r-1}}, \dots, g_{s_1-1}, \dots, g_0, 0^{\alpha_1}, 0).$$

Here, $(g_i)_{0 \leq i \leq n}$ is a sequence of positive integers. The last 0 is from the last point, because it does not have any lattice point after it. In comparison to $\mathbf{Drun}(\nu_\alpha, \varphi(w))$, it is clear that we only need to prove $g_\ell = f_\ell + 1$ for all ℓ .

For $\ell = n$, according to Proposition 37, an element $w(j)$ has $w(n)$ as its companion if and only if the stack is empty after its popping step, which is equivalent to $d'_j = 0$. Therefore, $g_n = f_n + 1$. For $\ell = 0$, it is clear that $g_0 = 1 = f_0 + 1$, since ν_α starts with a north step.

For $0 < \ell < n$, we know that f_ℓ is the number of nodes with $w(\ell)$ as companion, which is also the number of times we see $w(\ell + 1)$ at the top of the stack during a popping step according to Proposition 37. Suppose that $w(\ell + 1)$ is the i -th element in region k , thus $\ell + 1 = s_{k-1} + i$. We know that $w(\ell + 1)$ is pushed into the stack at step $2(n - s_{k-1}) - i + 1$. Suppose that there are $p = d'_{2(n-s_{k-1})-i}$ elements before $w(\ell + 1)$ is pushed down, we have $d'_{2(n-s_{k-1})-i+1} = p + 1$. Then $w(\ell + 1)$ is popped out once $d'_j \leq p$. When we see $w(\ell + 1)$ on top of the stack, we must have $d'_j = p + 1$ before it is popped. This is exactly the definition of $\mathbf{Cont}(\mathbf{Flip}(\nu_\alpha), \mathbf{Flip}(\Theta_1(w)))_{2(n-s_{k-1})-i}$, which is also $g_{s_{k-1}+i-1}$. We thus know that $g_{s_{k-1}+i-1}$ is the number of times we see $w(\ell + 1)$ on the top of the stack, the first time it is pushed, the other times we have an element whose companion is $w(\ell)$. We thus have $g_\ell = g_{s_{k-1}+i-1} = f_\ell + 1$. It follows that $\mathbf{Drun}(\nu_\alpha, \varphi(w)) = \mathbf{Cont}(\mathbf{Flip}(\nu_\alpha), \mathbf{Flip}(\Theta(w)))$, which concludes the proof. \square

An example of the proof of Theorem 39 can be seen in Figure 9. We thus solve [3, Open Problem 2.23].

Corollary 40. *The map $\mathbf{Flip} \circ \Theta$ is an anti-isomorphism between \mathcal{T}_α and $\mathcal{T}_{\mathbf{Flip}(\nu_\alpha)}$.*

Proof. This is a consequence of Theorem 39, and Propositions 27 and 30. \square

5 3 4 10 1 2 7 6 9 13 14 8 11 12 $\in \mathfrak{S}_\alpha(231)$

Companion index: 3 | 6, 6, 9 | 5 | 7, 8 | 11, 12, 14, 14 | 14, 14, 14




Figure 9: An example of the transfer of statistics in the two ways of looking at parabolic objects.

Remark 41. There is a typo in [3, Open Problem 2.23]. The statement should include “lattice anti-isomorphism” instead of “lattice isomorphism”, as we can also see in Figure 11 therein.

References

- [1] Anders Björner and Michelle L. Wachs, *Generalized quotients in Coxeter groups*, Transactions of the American Mathematical Society **308** (1988), 1–37.
- [2] Anders Björner and Michelle L. Wachs, *Shellable nonpure complexes and posets II*, Transactions of the American Mathematical Society **349** (1997), 3945–3975.
- [3] Cesar Ceballos, Wenjie Fang, and Henri Mühle, *The Steep-Bounce zeta map in Parabolic Cataland*, Journal of Combinatorial Theory, Series A **172** (2020), Research paper 105210, 59 pages.

- [4] Cesar Ceballos, Arnau Padrol, and Camilo Sarmiento, *The ν -Tamari lattice via ν -trees, ν -bracket vectors, and subword complexes*, The Electronic Journal of Combinatorics **27** (2020), #P1.14.
- [5] Marie-Pierre Delest and Xavier Gérard Viennot, *Algebraic languages and polyominoes enumeration*, Theoretical Computer Science **34** (1984), 169–206.
- [6] Emeric Deutsch, *An involution on Dyck paths and its consequences*, Discrete Mathematics **204** (1999), 163–166.
- [7] Georges-Théodule Guilbaud and Pierre Rosenstiehl, *Analyse algébrique d'un scrutin* (Marc Barbut, ed.), Gauthier-Villars, Paris, 1971.
- [8] Charles-Ange Laisant, *Sur la numération factorielle, application aux permutations*, Bulletin de la Société Mathématique de France **16** (1888), 176–183.
- [9] Derrick H. Lehmer, *Teaching combinatorial tricks to a computer* (Richard Bellman and Marshall Hall Jr, eds.), Vol. X, Proceedings of Symposia in Applied Mathematics, Providence, RI, 1960.
- [10] Henri Mühle and Nathan Williams, *Tamari lattices for parabolic quotients of the symmetric group*, The Electronic Journal of Combinatorics **26** (2019), #P4.34.
- [11] Viviane Pons, *The Rise-Contact Involution on Tamari intervals*, The Electronic Journal of Combinatorics **26** (2019), #P2.32.
- [12] Louis-François Prévaille-Ratelle and Xavier Viennot, *The enumeration of generalized Tamari intervals*, Transactions of the American Mathematical Society **369** (2017), 5219–5239.
- [13] Nathan Reading, *Cambrian lattices*, Advances in Mathematics **205** (2006), 313–353.
- [14] Dov Tamari, *The algebra of bracketings and their enumeration*, Nieuw Archief voor Wiskunde **10** (1962), 131–146.
- [15] Takemi Yanagimoto and Masashi Okamoto, *Partial orderings of permutations and monotonicity of a rank correlation statistic*, Annals of the Institute of Statistical Mathematics **21** (1969), 489–506.