

Interaction of two oscillating bubbles rising in a thin gap Audrey Filella, Véronique Roig, Patricia Ern

▶ To cite this version:

Audrey Filella, Véronique Roig, Patricia Ern. Interaction of two oscillating bubbles rising in a thin gap. American Physical Society DFD 2013, Nov 2013, Pittsburgh, United States. pp.0. hal-03668786

HAL Id: hal-03668786 https://hal.science/hal-03668786v1

Submitted on 16 May 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Open Archive TOULOUSE Archive Ouverte (OATAO)

OATAO is an open access repository that collects the work of Toulouse researchers and makes it freely available over the web where possible.

This is an author-deposited version published in : <u>http://oatao.univ-toulouse.fr/</u> <u>Eprints ID</u> : 9842

To cite this version : Filella, Audrey and Roig, Véronique and Ern, Patricia *Interaction of two oscillating bubbles rising in a thin gap.* In: American Physical Society DFD 2013, 24 November 2013 - 26 November 2013 (Pittsburgh, United States). (Unpublished)

Any correspondance concerning this service should be sent to the repository administrator: staff-oatao@listes-diff.inp-toulouse.fr

Interaction of two oscillating bubbles rising in a thin gap A. Filella, V. Roig, P. Ern

We investigate experimentally different mechanisms of hydrodynamic interaction between two oscillating bubbles rising in a liquid at rest confined in a thin gap cell. In order to understand the relation between the motion of a single bubble and its wake as well as the hydrodynamic interaction between two oscillating bubbles, we characterize the kinematics of the bubbles interaction using high speed cameras and we measure the associated liquid velocity by High-Frequency PIV. The motion of the trailing bubble is modified while passing through the unstable wake of the leading bubble: depending on the relative size of the bubbles and their relative position, we can observe horizontal attraction to the center of the leading bubble's wake, vertical entrainment, ejection of the second bubble by a vortex of the wake or a sequential combination of the three mechanisms.