



RECURRENCE PROPERTIES OF SEQUENCES OF INTEGERS

Fan Ai-Hua, Dominique Schneider

► To cite this version:

Fan Ai-Hua, Dominique Schneider. RECURRENCE PROPERTIES OF SEQUENCES OF INTEGERS. 2022. hal-03672933

HAL Id: hal-03672933

<https://hal.science/hal-03672933>

Preprint submitted on 19 May 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

RECURRENCE PROPERTIES OF SEQUENCES OF INTEGERS

AI-HUA FAN AND DOMINIQUE SCHNEIDER

ABSTRACT. In order to study the recurrence of sequences of integers, we investigate their L^2 -exactness and Θ -Hartman property (Θ being a set of rational numbers). Two classes of sequences of integers are well studied, which are return times relative to a weakly mixing system and Bernoulli random sequences.

1. INTRODUCTION

Let $\Lambda = \{u_n\}_{n \geq 1} \subset \mathbb{N}^* = \{1, 2, \dots\}$ be a subset of integers, or a strictly increasing sequence of integers. Λ is called a *Poincaré set* or *1-recurrent set* or simply *recurrent set* if for any measure preserving system (Y, \mathcal{B}, ν, S) and any $A \in \mathcal{B}$ with $\nu(A) > 0$ there exists $n \in \Lambda$ such that $\nu(A \cap S^{-n}A) > 0$. Poincaré recurrence theorem says that \mathbb{N}^* is a recurrent set. Λ is said to be L^2 -good if for any measure preserving system (Y, \mathcal{B}, ν, S) and any $g \in L^2(\nu)$, the follow average

$$(1.1) \quad A_N^\Lambda g(x) := \frac{1}{N} \sum_{n=1}^N g(S^{u_n}x)$$

L^2 -converges as $N \rightarrow \infty$. If furthermore,

$$L^2\text{-}\lim_{N \rightarrow \infty} A_N^\Lambda g := \mathbb{E}^{\mathcal{J}_S}(g),$$

we say that Λ is L^2 -exact, where $\mathbb{E}^{\mathcal{J}_S}$ denotes the conditional expectation with respect to the invariant σ -field \mathcal{J}_S of S . These notions are different. It is well known that $2\mathbb{N}^*$ is a recurrent set, but $2\mathbb{N}^* + 1$ is not. However both are L^2 -good. On the other hand, $\mathbb{N}^{*2} := \{1, 2^2, 3^2, \dots\}$ is L^2 -good, but it is not L^2 -exact.

The L^2 -good and L^2 -exact sequences can be characterized as follows. Let

$$(1.2) \quad \hat{A}_N^\Lambda(t) = \frac{1}{N} \sum_{n=1}^N e^{2\pi i u_n t}.$$

Theorem A. *A sequence $\Lambda = \{u_n\}_{n \geq 1} \subset \mathbb{N}^*$ is L^2 -good iff the limit $\lim_{N \rightarrow \infty} \hat{A}_N^\Lambda(t)$ exists for every $0 \leq t < 1$. It is L^2 -exact iff the limit exists and is equal to zero for every $0 < t < 1$.*

So the study of L^2 -goodness and the L^2 -exactness is a part of Fourier analysis. The L^2 -goodness doesn't imply the recurrence, but it will be proved that the L^2 -exactness does.

Theorem B. *An L^2 -exact sequence is recurrent.*

Let us present a class of L^2 -exact sequences. Let (X, \mathcal{A}, μ, T) be a measure preserving system. Let $A \in \mathcal{A}$. For $x \in X$, we define the *set of return times* of x into A by

$$R_A(x) = \{n \geq 1 : T^n x \in A\}.$$

If the system is weakly mixing, we have the following result.

Theorem C. *Suppose (X, \mathcal{A}, μ, T) is a weakly mixing measure preserving system and $A \in \mathcal{A}$ with $\mu(A) > 0$. Then for μ -a.e. $x \in X$, $R_A(x)$ is L^2 -exact. If T is weakly mixing and uniquely ergodic, then for every $x \in X$, $R_A(x)$ is L^2 -exact.*

Without the weakly mixing condition in the theorem, the conclusion doesn't hold in general. In fact, let $0 < \alpha < 1$ be an irrational number and $I = (-\delta, \delta)$ be an interval in the circle $\mathbb{T} = \mathbb{R}/\mathbb{Z}$ ($0 < \delta < 1/4$). The set $R_I(0)$ of n such that $n\alpha \in I$ is a set of return times (called a Sturmian sequence), which is not L^2 -exact because

$$\liminf \operatorname{Re} \frac{1}{N} \sum_{n=1}^N 1_I(n\alpha) e^{2\pi i n \alpha} \geq 2\delta \cos(2\pi\delta) > 0.$$

So, the condition in the definition of L^2 -exactness is not satisfied with S to be the α -rotation and $g(x) = e^{2\pi i x}$.

Let us present another class of recurrent sets. Let $\Theta \subset \mathbb{Q}/\mathbb{Z} \subset [0, 1)$ be a set of rational numbers containing 0. Define

$$D_\Theta = \left\{ m \geq 1 : m = [q_1, \dots, q_r] \text{ for some } \frac{p_1}{q_1}, \dots, \frac{p_r}{q_r} \in \Theta \right\}$$

where $[q_1, \dots, q_r]$ denotes the least common multiple of q_1, \dots, q_r . A set $\Lambda \subset \mathbb{N}^*$ is called a Θ -Hartman sequence if for each $m \in D_\Theta$ there is a subsequence $\{u_{n_k}\} \subset \Lambda$ such that (i) $\{u_{n_k}\} \subset m\mathbb{Z} \cap \Lambda$ and (ii)

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{k=1}^N e^{2\pi i \alpha u_{n_k}} = 0 \quad \forall \alpha \in (0, 1) \setminus \Theta.$$

If $\Theta = \{0\}$, the sequence $\{u_{n_k}\}$ satisfying (ii) is said to be Hartman uniformly distributed, i.e. uniformly distributed in the Bohr compactification of \mathbb{Z} . Therefore a $\{0\}$ -Hartman sequence is nothing but a sequence containing a Hartman uniformly distributed subsequence. Remark that the limit in (ii) is ensured if $\{\alpha u_{n_k}\}$ is uniformly distributed modulo 1. Also remark that $D_\Theta = \{1\}$ for $\Theta = \{0\}$, $D_\Theta = \{2^n : n \geq 0\}$ for $\Theta = \{\frac{k}{2^n} : k, n \in \mathbb{N}\}$, $D_\Theta = \mathbb{N}$ for $\Theta = \mathbb{Q}/\mathbb{Z}$. The second assertion in Theorem A means that Hartman uniformly distributed sequences are just L^2 -exact sequences. The following result shows that the Θ -Hartman property implies the recurrence.

Theorem D. *Let $\Theta \subset \mathbb{Q}/\mathbb{Z}$ contain 0. Every Θ -Hartman sequence is recurrent.*

Now let us describe the class of Bernoulli random sequences of integers. Let $\{p_n\}_{n \geq 1}$ be a sequence of real numbers such that

$$0 \leq p_n \leq 1, \quad \sum_{n=1}^{\infty} p_n = \infty.$$

Let $\{\xi_n\}_{n \geq 1}$ be a sequence of independent Bernoulli random variables such that

$$P(\xi_n = 1) = p_n = 1 - P(\xi_n = 0).$$

We are interested in the random set

$$W(\omega) = \{n \geq 1 : \xi_n(\omega) = 1\}.$$

This is almost surely (a.s. for short) an infinite subset of integers.

Theorem E. *The sequence $W(\omega)$ is a.s. L^2 -exact (equivalently Hartman uniformly distributed) if the following conditions are satisfied:*

$$(1) \quad \log N = o\left(\sum_{n=1}^N p_n\right), \quad (2) \quad \sum_{n=1}^N |p_n - p_{n+1}| = o\left(\sum_{n=1}^N p_n\right).$$

Without the variation condition (2), the conclusion of the theorem is not true. For example, if $p_{2n} = 0$ and $p_{2n+1} = 1$, we have the deterministic set $W(\omega) = 2\mathbb{N}^* + 1$ (the set of odd integers) which is even not recurrent. If p_n is decreasing, the condition (2) becomes redundant.

Kahane and Katznelson [9] obtained the following conditions for $W(\omega)$ to be a.s. Hartman uniformly distributed (i.e. L^2 -exact)

$$\lim np_n = \infty; \quad n^{1+\delta}p_n \uparrow, \quad n^{1-\delta}p_n \downarrow \quad \forall 0 < \delta < 1.$$

The condition stated in the above theorem seems a little bit more flexible.

Theorem E implies that if $p_n = n^{-\tau}$ with $0 < \tau < 1$, then $W(\omega)$ is a.s. Hartman uniformly distributed. This is to be compared with the following Kahane and Katznelson's result: if $p_n = n^{-1}$, then $W(\omega)$ is a.s. a Sidon set; if $p_n = n^{-\tau}$ with $\tau < 1$, then $W(\omega)$ is a.s. not a Sidon set [8]. We will discuss the L^2 -universality of $W(\omega)$.

2. PRELIMINARIES

We present in this section some preliminaries including van der Corput inequality, spectral lemma and Kronecker factor.

2.1. van der Corput inequality. We will use a simple form the van der Corput inequality (see [12]):

Lemma 2.1. *For $N \geq 1$ complex numbers a_0, a_1, \dots, a_{N-1} and any integer H with $1 \leq H \leq N$, we have*

$$\left| \frac{1}{N} \sum_{n=0}^{N-1} a_n \right|^2 \leq \frac{C}{H} \left(\frac{1}{N} \sum_{n=0}^{N-1} |a_n|^2 + \left| \sum_{h=1}^H \frac{1}{N} \sum_{n=0}^{N-h} a_n \overline{a_{n+h}} \right| \right)$$

where C is a absolute constant.

2.2. Spectral lemma. The spectral lemma holds for contracting operators on Hilbert space (see [11]). Let (X, \mathcal{A}, μ, T) is a measure preserving system. The Koopman map $f \mapsto f \circ T$, also denoted by T , is an isometry on $L^2(\mu)$. The *spectral measure* μ_f of $f \in L^2(\mu)$ is characterized by its Fourier coefficients as follows

$$\widehat{\mu}_f(n) = \int f \circ T^n \cdot f d\mu \text{ for } n \geq 0; \quad \widehat{\mu}_f(n) = \overline{\widehat{\mu}_f(-n)} \text{ for } n < 0.$$

Lemma 2.2. Let $p(z) = \sum a_n z^n$ be a polynomial. We have

$$\|p(T)f\|_{L^2(T)}^2 = \int_{\mathbb{R}/\mathbb{Z}} |p(e^{2\pi i t})|^2 d\mu_f(t)$$

2.3. Kronecker factor. Let (X, \mathcal{A}, μ, T) is a measure preserving system. The *Kronecker factor* \mathcal{K} of the system is by definition the closed linear span in $L^2(\mu)$ of the eigenfunctions for T . There is a sub σ -algebra \mathcal{A}' of \mathcal{A} such that $\mathcal{K} = L^2(\mathcal{A}')$. The orthogonal complementary space \mathcal{K}^\perp consists of all functions $f \in L^2(\mu)$ whose spectral measures μ_f are continuous.

If $f = f_1 + f_2$ with $f_1 \in \mathcal{K}$ and $f_2 \in \mathcal{K}^\perp$, then $\mu_f = \mu_{f_1} + \mu_{f_2}$. Furthermore, μ_{f_2} is continuous, μ_{f_1} is discrete and $\mu_{f_1}(\{\theta\}) \neq 0$ if and only if $e^{2\pi i \theta}$ is an eigenvalue and f is not orthogonal to the corresponding eigenfunction.

For any $\theta \in \mathbb{R}/\mathbb{Z}$, denote

$$\mathcal{K}_\theta = \{f \in H : f(Tx) = e^{2\pi i \theta} f(x)\}.$$

That \mathcal{K}_θ is not trivial means that $e^{2\pi i \theta}$ is an eigenvalue for T . Thus \mathcal{K} is the direct sum of all these non-trivial \mathcal{K}_θ with $\theta \in \mathbb{R}/\mathbb{Z}$. If the system is ergodic, all eigenfunctions are in $L^\infty(\mu)$ and each eigenspace \mathcal{K}_θ is one-dimensional.

The subspace $L_0^2(\mu)$ of $L^2(\mu)$ consists of those f such that $\int f d\mu = 0$. Let $\mathcal{K}^* = \mathcal{K} \cap L_0^2(\mu)$. We have the orthogonal decomposition

$$L^2(\mu) = \mathbb{C} \oplus L_0^2(\mu) = \mathbb{C} \oplus \mathcal{K}^* \oplus \mathcal{K}^\perp.$$

For any $\Theta \subset \mathbb{R}/\mathbb{Z}$, we denote

$$\mathcal{K}_\Theta = \bigoplus_{\theta \in \Theta} \mathcal{K}_\theta.$$

In the following, we only consider the case where Θ is a set of rational numbers.

3. RECURRENCE, L^2 -GOODNESS AND L^2 -EXACTNESS

3.1. Recurrent sequences. Bertrand-Mathis [1] proved that Λ is a recurrent set if and only if it is a *set of intersectivity*, that is to say, $\Lambda \cap (S - S) \neq \emptyset$ for any set $S \subset \mathbb{N}^*$ with positive Banach density, i.e.

$$B(S) = \sup \lim_{n \rightarrow \infty} \frac{|S \cap I_n|}{|I_n|}$$

where the supremum is take over all sequence of intervals $\{I_n\}$ with length $|I_n|$ tending to infinity as n does. As was pointed out in [1], this result was also proved independently by P. Liardet, J. F. M  la and Y. Katznelson.

A necessary condition for Λ to be recurrent is that $m\mathbb{Z} \cap \Lambda \neq \emptyset$ for any $m \in \mathbb{N}^*$, which is easy to see from the definition. It follows that the set $2\mathbb{N}^* + 1$ of odd integers is not recurrent.

Kamae and Mendes France [10] proved that every van der Corput sequence is intersective and then a recurrent set. Recall the definition of a van der Corput sequence Λ : for any sequence of real numbers $\{x_n\}_{n \geq 1}$ such that $\{x_{n+h} - x_n\}_{n \geq 1}$ is uniformly distributed (mod 1) for each $h \in \Lambda$, all sequences $\{x_{an+b}\}_{n \geq 1}$ are uniformly distributed (mod 1) for all integers $a \geq 1, b > 0$.

Furstenberg proved that if $p \in \mathbb{Q}[x]$ is a polynomial taking values in \mathbb{N} such that $p(0) = 0$, then the image $p(\mathbb{N}^*)$ is a recurrent sequence. In particular, so are $2\mathbb{N}^*$ and $\mathbb{N}^{*2} = \{1, 2^2, 3^2, \dots\}$.

Bourgain [2] proved the pointwise convergence of $A_N^{p(\mathbb{N})}g$. Bourgain, Furstenberg, Katznelson and Ornstein proved the pointwise converges of $A_N^{R_A(x)}g$ for the return times $R_A(x)$ in an ergodic system (see [2]).

The L^2 -goodness doesn't imply the recurrence. But the L^2 -exactness does.

Theorem 3.1. *Suppose that $\Lambda \subset \mathbb{N}^*$ is an L^2 -exact sequence. Let (Y, \mathcal{B}, ν, S) be a measure preserving system and let $g \in L^2(\nu)$.*

- (1) *If $g \perp g \circ S^n$ for all $n \in \Lambda$, then $g \perp L^2(\mathcal{I}_S)$.*
- (2) *Λ is a recurrent sequence.*

Proof. (1) It suffices to prove $g \perp \mathbb{E}^{\mathcal{I}_S}g$. Let $\Lambda = \{u_n\}$. Recall that

$$A_N^\Lambda g(y) := \frac{1}{N} \sum_{n=1}^N g(S^{u_n}y) \rightarrow \mathbb{E}^{\mathcal{I}_S}g.$$

Write

$$(3.1) \quad \left| \int g \cdot \mathbb{E}_\nu^{\mathcal{I}_S}g d\nu \right| \leq \left| \int g[\mathbb{E}_\nu^{\mathcal{I}_S}g - A_N^\Lambda g] d\nu \right| + \left| \int g \cdot A_N^\Lambda g d\nu \right|$$

By the hypothesis, g is orthogonal to $A_N^\Lambda g$. So, it follows from (3.1) that

$$\begin{aligned} \left| \int g \cdot \mathbb{E}_\nu^{\mathcal{I}_S}g d\nu \right| &\leq \|g[\mathbb{E}_\nu^{\mathcal{I}_S}g - A_N^\Lambda g]\|_{L^1(\nu)} \\ &\leq \|g\|_{L^2(\nu)} \|\mathbb{E}_\nu^{\mathcal{I}_S}g - A_N^\Lambda g\|_{L^2(\nu)}. \end{aligned}$$

However the last term tends to zero when $n \rightarrow \infty$, because $A_N^\Lambda g \rightarrow \mathbb{E}^{\mathcal{I}_S}g$.

(2) Otherwise, there exist a measure preserving system (Y, \mathcal{B}, ν, S) and a $B \in \mathcal{B}$ with $\nu(B) > 0$ such that

$$\int 1_B \cdot 1_B \circ S^n d\nu = \nu(B \cap S^{-n}B) = 0, \quad \forall n \in \Lambda.$$

Applying (1) to 1_B , we get $1_B \perp L^2(\mathcal{I}_S)$. In particular, $1_B \perp 1$ which means $\nu(B) = 0$. \square

3.2. Characterizations of L^2 -goodness and L^2 -exactness.

Theorem 3.2. *A sequence $\Lambda = \{u_n\} \subset \mathbb{N}$ is L^2 -good iff the following limit exists*

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N e^{2\pi i u_n t}$$

for all $t \in (0, 1)$. The sequence is L^2 -exact iff not only the above limit exists but the limit is zero.

Proof. We use the notation defined by (1.1) and (1.2).

L^2 -goodness. By considering the rotation $x \mapsto x + t$ on \mathbb{R}/\mathbb{Z} , we see that the condition is necessary. Let (Y, \mathcal{B}, S, ν) be an arbitrary measure preserving system and $g \in L^2(\nu)$. We prove the sufficiency by showing that $A_N^\Lambda g$ is a Cauchy sequence in $L^2(\nu)$. In fact, this is a consequence of the spectral lemma which implies

$$\|A_p^\Lambda g - A_q^\Lambda g\|_{L^2(\nu)}^2 \leq \int |\widehat{A}_p^\Lambda(t) - \widehat{A}_q^\Lambda(t)|^2 d\mu_g(t) \quad (\forall 1 \leq p < q),$$

and the Lebesgue dominated convergence theorem.

L^2 -exactness. To get the necessity, we consider the irrational rotation $x \mapsto x + t$ on \mathbb{R}/\mathbb{Z} if t is irrational, or the ergodic rotation $x \mapsto x + 1$ on $\mathbb{Z}/r\mathbb{Z}$ when $t = \frac{s}{r}$ is rational with $(s, r) = 1$. To prove the sufficiency, we have only to show that $A_N^\Lambda g$ tends to zero when $g \in \mathcal{K}_0^\perp$ (\mathcal{K}_0 being the space of invariant functions). It is true because

$$\|A_p^\Lambda g\|_{L^2(\nu)}^2 \leq \int |\widehat{A}_p^\Lambda(t)|^2 d\mu_g(t)$$

and the spectral measure μ_g doesn't charge the point 0. \square

The theorem shows that Λ is L^2 -exact sequences means the sequence $n \mapsto 1$ or 0 according to $n \in \Lambda$ or not is almost periodic sequences in the sense of Bohr and that L^2 -good sequences correspond to generalized almost periodic sequences in the sense of Hartman (see [6], p.72).

3.3. Some remarks. Suppose that $\Lambda = \{u_n\}_{n \geq 1}$ is L^2 -good. Let

$$c_\Lambda(t) = \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N e^{2\pi i u_n t}.$$

Suppose that $c_\Lambda(t) = 0$ for all but at most countably many points t . Let (Y, \mathcal{B}, ν, S) be a measure preserving system. Any $g \in L^2(\nu)$ can be decomposed into

$$g = \sum a_\theta g_\theta + g_{K^\perp} \quad (a_\theta \in \mathbb{C})$$

where the sum is taken over all θ 's such that $e^{2\pi i \theta}$ is an eigenvalue of S , f_θ is the orthogonal projection of g onto the eigenspace corresponding to the eigenvalue $e^{2\pi i \theta}$ and g_{K^\perp} is the projection on the orthogonal of the

Kronecker factor \mathcal{K} . In this case, we can identify the L^2 -limit of $A_N^\Lambda g$ as follows

$$\lim_{N \rightarrow \infty} A_N^\Lambda g = \sum a_\theta c_\Lambda(\theta) g_\theta.$$

We have only to show this for each component of g in the above decomposition. In fact, if $g \in K^\perp$, then the spectral measure of g is continuous, which implies $\lim_{N \rightarrow \infty} A_N^\Lambda g = 0$ through the spectral lemma and the fact that $c_\Lambda(t) \neq 0$ for at most countably many t 's. If $g = g_\theta$, it suffices to notice that

$$A_N^\Lambda g_\theta = \hat{A}_N^\Lambda(\theta) g_\theta.$$

A set $\Lambda = \{u_n\}_{n \geq 1}$ of integers is said to be r -recurrent ($r \geq 1$) if for any measure-preserving system (Y, \mathcal{B}, ν, S) and any $A \in \mathcal{B}$ with $\nu(A) > 0$ there exists $n \in \Lambda$ such that

$$\nu(A \cap S^{-n}A \cap \cdots \cap S^{-rn}A) > 0.$$

Furstenberg proved that \mathbb{N}^* is r -recurrent for every $r \geq 1$ by showing

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N \nu(A \cap S^{-n}A \cap \cdots \cap S^{-rn}A) > 0.$$

We remark that any set $\Lambda = \{u_n\}_{n \geq 1}$ such that $\lim_{n \rightarrow \infty} \frac{u_n}{n} = 1$ is also r -recurrent for every $r \geq 1$. This is a consequence of the positivity of the above liminf and the following fact: Assume $\lim_{n \rightarrow \infty} \frac{u_n}{n} = 1$. For any bounded sequence of numbers $\{a_n\}$ we have

$$(3.2) \quad \lim_{N \rightarrow +\infty} \left| \frac{1}{N} \sum_{n=1}^N a_{u_n} - \frac{1}{N} \sum_{n=1}^N a_n \right| = 0.$$

As a consequence, Furstenberg's argument proves an improvement of Szemerédi theorem: Let $\Lambda = \{u_k\} \subset \mathbb{N}^*$ be a strictly increasing sequence such that $\frac{u_k}{k} \rightarrow 1$ as $k \rightarrow \infty$. Let $S \subset \mathbb{N}^*$ be a set with strictly positive upper Banach density. For any integer $\ell \geq 2$, there exist $h \geq 1$ and $u \in \Lambda$ such that

$$h \in S, \quad h + u \in S, \quad h + 2u \in S, \quad \dots, \quad h + \ell u \in S.$$

Let $\Lambda = \{u_k\}$ be a given sequence with some property like L^2 -goodness or L^2 -exactness (but not the recurrence). By (3.2), a sequence $\{u_k + \delta_k\}$ which is a perturbation of $\{u_k\}$ by $\{\delta_k\}$ such that $\delta_k = o(k)$ inherits the same property from Λ . For example, a sequence $\{u_k\}$ such that $u_k = \zeta k + o(k)$ with $\zeta > 0$ is L^2 -exact; a sequence $\{u_k\}$ such that $u_k = k^2 + o(k)$ is L^2 -good and has the same c_Λ function as $\{k^2\}$.

4. WIENER-WINTNER THEOREM AND RETURN TIMES

Let $\Lambda \subset \mathbb{N}^*$. One way to study the recurrence of Λ is to investigate the limit of the average $A_N^\Lambda g$. This section is devoted to the L^2 -exactness of the set of return times in a weakly mixing system. We are actually led to recognize the limit of the Birkhoff average of a product system.

Given two measure preserving systems (X, \mathcal{A}, μ, T) and (Y, \mathcal{B}, ν, S) . Their product is the measure preserving system $(X \times Y, \mathcal{A} \otimes \mathcal{B}, \mu \otimes \nu, T \times S)$. In general, the ergodicities of both systems don't imply the ergodicity of their product. It is easy to see that

$$\mathcal{J}_T \otimes \mathcal{J}_S \subset \mathcal{J}_{T \times S}.$$

The above inclusion maybe strict. We will prove that we do get an equality when T is weakly mixing and that in this case, for any $F \in L^2(\mu \times \nu)$ we have

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=0}^{N-1} F(T^n x, S^n y) = \mathbb{E}_\nu^{\mathcal{J}_S} \int F(x, y) d\mu(x).$$

Moreover, for a fixed $f \in L^2(\mu)$, we can find a measurable set X_f with full μ -measure such that for any $x \in X_f$, for any measure preserving system (Y, \mathcal{B}, ν, S) and for any $g \in L^2(\nu)$ we have

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=0}^{N-1} f(T^n x) g(S^n y) = \int f d\mu \cdot \mathbb{E}_\nu(g | \mathcal{J}_S)$$

in $L^2(\nu)$. Bourgain-Furstenberg-Katznelson-Ornstein (see [2]) proved the pointwise convergence without identifying the limit but under the condition that T is only ergodic. With the limit identified in the last equality, we can prove that if T is weakly mixing, for any set A with $\mu(A) > 0$, the set of return times $R_A(x)$ is L^2 -exact for μ -a.e. x .

4.1. Some lemmas. Let (X, \mathcal{A}, μ, T) is a measure preserving system. If $f \in L^2(\mu)$ and $1 \leq h \leq N$, write

$$J_{N,h}(x) = \frac{1}{N} \sum_{n=0}^{N-h} f(T^n x) \overline{f(T^{n+h} x)}.$$

Lemma 4.1. *Let (X, \mathcal{A}, μ, T) is a measure preserving system. If $f \in L^2(\mu)$ and $1 \leq H \leq N$, for any Borel measure ν on $[0, 1)$ we have*

$$\left\| \frac{1}{N} \sum_{n=0}^{N-1} f(T^n x) e^{2\pi i n t} \right\|_{L^2(\nu)}^2 \leq \frac{C}{H} \left(J_{N,0}(x) + \int \left| \sum_{h=1}^H e^{2\pi i h t} J_{N,h}(x) \right| d\nu(t) \right)$$

where C is the constant in the van der Corput inequality.

Proof. Apply the van der Corput inequality to $a_n = f(T^n x) e^{2\pi i n t}$, we get

$$(4.1) \quad \left| \frac{1}{N} \sum_{n=0}^{N-1} f(T^n x) e^{2\pi i n t} \right|^2 \leq \frac{C}{H} \left(J_{N,0}(x) + \left| \sum_{h=1}^H e^{2\pi i h t} J_{N,h}(x) \right| \right).$$

Integrate (4.1) with respect to ν to obtain the desired result. \square

Remark that the functions $J_{N,h}$ are independent of t . If the system is ergodic, by the Birkhoff ergodic theorem, for μ -a.e. x and for all $h \geq 0$ we have

$$(4.2) \quad \lim_{N \rightarrow \infty} J_{N,h}(x) = \int f \cdot \overline{f \circ T^h} d\mu.$$

The following lemma will allow us to apply the spectral lemma. The point (2) is due to Bourgain.

Lemma 4.2. *Let (X, \mathcal{A}, μ, T) is an ergodic measure preserving system.*

(1) *If $f \in \mathcal{K}^*$, for μ -a.e. x and for any Borel probability measure ν on $[0, 1)$ we have*

$$\overline{\lim}_{N \rightarrow \infty} \left\| \frac{1}{N} \sum_{n=0}^{N-1} f(T^n x) e^{2\pi i n t} \right\|_{L^2(\nu)}^2 \leq C \|f\|_{L^2(\mu)} \overline{\lim}_{N \rightarrow \infty} \left\| \frac{1}{N} \sum_{n=0}^{N-1} f(T^n x) e^{2\pi i n t} \right\|_{L^2(\mu \times \nu)}$$

where C is a constant.

(2) *If $f \in \mathcal{K}^\perp$, for μ -a.e. x we have*

$$\lim_{N \rightarrow \infty} \sup_{t \in [0,1)} \left| \frac{1}{N} \sum_{n=0}^{N-1} f(T^n x) e^{2\pi i n t} \right| = 0.$$

Proof. Since the system is ergodic, for μ -a.e. x and for all $h \geq 0$ we have

$$\lim_{N \rightarrow \infty} J_{N,h}(x) = \int f \cdot \overline{f \circ T^h} d\mu.$$

In (4.1), let $N \rightarrow \infty$ then let $H \rightarrow \infty$. We obtain that for μ -a.e. x the limit

$$\overline{\lim}_{N \rightarrow \infty} \left\| \frac{1}{N} \sum_{n=0}^{N-1} f(T^n x) e^{2\pi i n t} \right\|_{L^2(\nu)}^2$$

is bounded up to the multiplicative constant C by

$$\begin{aligned} & \overline{\lim}_{H \rightarrow \infty} \int d\nu(t) \left| \int f(x) \left(\frac{1}{H} \sum_{h=1}^H e^{2\pi i h t} \overline{f \circ T^h(x)} \right) d\mu(x) \right| \\ & \leq \|f\|_{L^2(\mu)} \overline{\lim}_{H \rightarrow \infty} \left\| \frac{1}{H} \sum_{h=1}^H e^{2\pi i h t} \overline{f \circ T^h(x)} \right\|_{L^2(\mu \times \nu)}. \end{aligned}$$

For the last inequality, we have used the Cauchy-Schwarz inequality twice (first with respect to μ , then with respect to ν). Thus we have proved (1)

To prove (2), first we deduce from (4.1) the uniform estimate

$$\sup_{t \in [0,1)} \left| \frac{1}{N} \sum_{n=0}^{N-1} f(T^n x) e^{2\pi i n t} \right| \leq \frac{C}{H} \left(J_{N,0}(x) + \sum_{h=1}^H |J_{N,h}(x)| \right).$$

Then using (4.2) and the fact $\int f \cdot \overline{f \circ T^h} d\mu = \widehat{\mu}_f(h)$, we get that μ -a.e.

$$\overline{\lim}_{N \rightarrow \infty} \sup_{t \in [0,1)} \left| \frac{1}{N} \sum_{n=0}^{N-1} f(T^n x) e^{2\pi i n t} \right|^2 \leq C \lim_{H \rightarrow \infty} \sqrt{\frac{1}{H} \sum_{h=1}^H |\widehat{\mu}_f(h)|^2}.$$

The last limit is zero according to Wiener theorem because the spectral measure μ_f is continuous for $f \in \mathcal{K}^\perp$. \square

4.2. Wiener-Wintner theorem. We are now ready to prove the following theorem, a kind of Wiener-Wintner theorem which provides a well defined expression for the ergodic limit.

Recall that \mathcal{K} denotes the Kronecker factor of the system (X, \mathcal{A}, μ, T) . If the system is ergodic, the space \mathcal{K}_0 of invariant functions consists of constant functions. Also recall that $L_0^2(\mu)$ is the subspace of $L^2(\mu)$ consisting of functions f such that $\int f d\mu = 0$. We have the decomposition

$$L_0^2(\mu) = \mathcal{K}^* \oplus \mathcal{K}^\perp \quad \text{with} \quad \mathcal{K}^* = \mathcal{K} \ominus \mathcal{K}_0.$$

When T is weakly mixing, $\mathcal{K} = \mathcal{K}_0$, i.e. $L_0^2(\mu) = \mathcal{K}^\perp$.

Let (Y, \mathcal{B}, ν, S) be another measure preserving system. The theorem below provides us a kind of approximation of the projection $\mathbb{E}_\nu^{\mathcal{J}_S} g$.

Theorem 4.3. *Suppose that (X, \mathcal{A}, μ, T) is a weakly mixing measure preserving system. Let $f \in L^2(\mu)$. There exists a measure set X_f with full μ -measure such that for any $x \in X_f$, for any measure preserving system (Y, \mathcal{B}, ν, S) and for any $g \in L^2(\nu)$ we have*

$$(4.3) \quad \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=0}^{N-1} f(T^n x) g(S^n y) = \int f d\mu \cdot \mathbb{E}_\nu(g | \mathcal{J}_S)$$

in $L^2(\nu)$. For any $F \in L^2(\mu \times \nu)$, we have

$$(4.4) \quad \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=0}^{N-1} F(T^n x, S^n y) = \mathbb{E}_\nu^{\mathcal{J}_S} \int F(x, y) d\mu(x)$$

in $L^2(\mu \times \nu)$. The results remain true if we assume that T is ergodic and $\mathcal{J}_{T \times S} = \mathcal{J}_T \otimes \mathcal{J}_S$ instead of assuming T is weakly mixing.

Proof. Let us first prove (4.3). By the linearity on f of both sides of the equality (4.3) and the von Neumann theorem applied to (Y, \mathcal{B}, ν, S) , in order to prove (4.3), we can assume that $f \in L_0^2(\mu)$. Since $L_0^2(\mu) = \mathcal{K}^* + \mathcal{K}^\perp$, it suffice to prove it for f in \mathcal{K}^* or in \mathcal{K}^\perp .

The spectral lemma gives

$$\left\| \frac{1}{N} \sum_{n=0}^{N-1} f(T^n x) g(S^n y) \right\|_{L^2(\nu)}^2 \leq \int \left| \frac{1}{N} \sum_{n=0}^{N-1} f(T^n x) e^{2\pi i n t} \right|^2 d\nu_g(t).$$

Then for $f \in \mathcal{K}^\perp$, we get the desired result immediately from Lemma 4.2 (2). For $f \in \mathcal{K}^*$, there is nothing to prove when T is weakly mixing which

implies $\mathcal{K}^* = \{0\}$. Assume that T is ergodic and $\mathcal{J}_{T \times S} = \mathcal{J}_T \otimes \mathcal{J}_S$. By Lemma 4.2 (1), we get

$$\overline{\lim}_{N \rightarrow \infty} \left\| \frac{1}{N} \sum_{n=0}^{N-1} f(T^n x) g(S^n y) \right\|_{L^2(\nu)}^2 \leq C \|f\|_{L^2} \overline{\lim}_{N \rightarrow \infty} \left\| \frac{1}{N} \sum_{n=0}^{N-1} f(T^n x) e^{2\pi i n t} \right\|_{L^2(\mu \times \nu_g)}.$$

Notice that, again by the spectral lemma,

$$\left\| \frac{1}{N} \sum_{n=0}^{N-1} f(T^n x) e^{2\pi i n t} \right\|_{L^2(\mu \times \nu_g)} = \left\| \frac{1}{N} \sum_{n=0}^{N-1} f(T^n x) g(S^n y) \right\|_{L^2(\mu \times \nu)}.$$

Now we can conclude with von Neumann theorem applied to the product system $T \times S$ to get that

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=0}^{N-1} f(T^n x) g(S^n y) = \mathbb{E}^{\mathcal{J}_T} f \mathbb{E}^{\mathcal{J}_S} g = 0.$$

Now prove (4.4). Both sides of (4.4) define bounded operators on $L^2(\mu \times \nu)$. So, we have to check (4.4) for F of the form of finite sum $\sum c_i 1_{A_i}(x) 1_{B_i}(y)$ with $A_i \in \mathcal{A}$ and $B_i \in \mathcal{B}$. For such simple function, the already proved (4.3) implies that the limit on the left hand side of (4.4) is equal to

$$\sum c_i \mu(A_i) \mathbb{E}^{\mathcal{J}_S} 1_{B_i} = \mathbb{E}^{\mathcal{J}_S} \sum c_i \mu(A_i) 1_{B_i} = \mathbb{E}^{\mathcal{J}_S} \int F d\mu.$$

□

4.3. Finite systems. By studying finite systems, we will show that Theorem 4.3 doesn't hold without the assumption that T is weakly mixing and that the L^2 -good sequence $\Lambda = \{n^2\}$ is not L^2 -exact (this can also be shown by computing $c_\Lambda(t)$).

For any integer $m \geq 2$, we denote by $T_m : \mathbb{Z}/m\mathbb{Z} \rightarrow \mathbb{Z}/m\mathbb{Z}$ the translation on the group $\mathbb{Z}/m\mathbb{Z}$ defined by

$$T_m x = x + 1 \pmod{m}.$$

T_m is a uniquely ergodic transformation with the unique invariant probability measure $\frac{1}{m} \sum_{x \in \mathbb{Z}/m\mathbb{Z}} \delta_x$. Let m_1, m_2 be two integers. Their least common multiple is denoted $[m_1, m_2]$.

Proposition 4.4. *Let $m_1, m_2 \geq 2$ be two integers. Consider the two translations T_{m_1}, T_{m_2} and their product $T_{m_1} \times T_{m_2}$.*

(1) *The orbit $O(x, y)$ of any point $(x, y) \in \mathbb{Z}/m_1\mathbb{Z} \times \mathbb{Z}/m_2\mathbb{Z}$ is a cycle of length $[m_1, m_2]$ consisting of $(x, y) + (k, k)$ with $0 \leq k < [m_1, m_2]$.*

(2) *Two orbits $O(x, y)$ and $O(x', y')$ are equal iff*

$$x' - x = y' - y \pmod{[m_1, m_2]}.$$

Therefore there are $\frac{m_1 m_2}{[m_1, m_2]}$ different cycles.

(3) *A set in $\mathbb{Z}/m_1\mathbb{Z} \times \mathbb{Z}/m_2\mathbb{Z}$ is $T_{m_1} \times T_{m_2}$ -invariant iff it is a finite union of cycles.*

Proof. (1) Since the finite set $O(x, y)$ verifies $T_{m_1} \times T_{m_2} O(x, y) \subset O(x, y)$ and $T_{m_1} \times T_{m_2}$ is bijective, we have $T_{m_1} \times T_{m_2} O(x, y) = O(x, y)$. This implies that $O(x, y)$ is a cycle. Let ℓ be the length of the cycle. Then

$$(T_{m_1} \times T_{m_2})^\ell(x, y) = (x, y),$$

i.e. $x + \ell = x \pmod{m_1}$ and $y + \ell = y \pmod{m_2}$. It follows that $\ell = [m_1, m_2]$.

(2) According to (1), $O(x, y) = O(x', y')$ means $(x', y') = (x, y) + (k, k)$ for some $0 \leq k < [m_1, m_2]$. In other words, we have $x' - x = k \pmod{m_1}$ and $y' - y = k \pmod{m_2}$ for some $0 \leq k < [m_1, m_2]$. Equivalently $x' - x = y' - y \pmod{[m_1, m_2]}$.

(3) It is a consequence of (1). □

It follows immediately from the proposition that for any function f defined on $\mathbb{Z}/m_1\mathbb{Z} \times \mathbb{Z}/m_2\mathbb{Z}$ we have

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} f(T_{m_1}^k x, T_{m_2}^k y) = \frac{1}{[m_1, m_2]} \sum_{(a,b) \in O(x,y)} f(a, b).$$

Remark that the limit only depend on the cycle.

For $f(x, y) = 1_{\{(i,j)\}}(x, y) = 1_{\{i\}}(x)1_{\{j\}}(y)$, the characteristic function of the point $(i, j) \in \mathbb{Z}/m_1\mathbb{Z} \times \mathbb{Z}/m_2\mathbb{Z}$, we get

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{k=0}^{n-1} 1_{\{i\}}(T_{m_1}^k x) 1_{\{j\}}(T_{m_2}^k y) = \begin{cases} \frac{1}{[m_1, m_2]} & \text{if } (x, y) \in O(i, j) \\ 0 & \text{if } (x, y) \notin O(i, j). \end{cases}$$

This shows that Theorem 4.3 doesn't hold without the assumption that T is weakly mixing.

Let $m \geq 2$ be an integer. For $0 \leq i < m$, write

$$R_i = \{n \geq 1 : n^2 = i \pmod{m}\}.$$

Notice that R_i is a disjoint union of arithmetic sequences:

$$R_i = \bigsqcup_{0 \leq k < m, k^2 \equiv i \pmod{m}} m\mathbb{Z} + k.$$

So the density of R_i exists:

$$d_i = \lim_{N \rightarrow \infty} \frac{\#\{1 \leq n \leq N : n \in R_i\}}{N} = \frac{\#\{0 \leq k < m : k^2 \equiv i \pmod{m}\}}{m}.$$

It follows the following proposition.

Proposition 4.5. *Consider the unique ergodic transformation $T_m : \mathbb{Z}/m\mathbb{Z} \rightarrow \mathbb{Z}/m\mathbb{Z}$ defined by $T_m x = x + 1 \pmod{m}$, where $m \geq 2$ is an integer. For any $j \in \mathbb{Z}/m\mathbb{Z}$ and any $x \in \mathbb{Z}/m\mathbb{Z}$, the following limit exists*

$$L_j(x) := \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N 1_{\{j\}}(T_m^{n^2} x) = d_{j-x}.$$

For example, when $m = 3$, we have $0^2 \equiv 0, 1^2 \equiv 1$ and $2^2 \equiv 1$, so that $d_0 = 1/3, d_1 = 2/3, d_2 = 0$. Then

$$L_0(0) = \frac{1}{3}, \quad L_0(1) = 0, \quad L_0(2) = \frac{2}{3}.$$

The limit function L_3 is not constant, however $\mathbb{E}(1_{\{0\}}|\mathcal{J}) = 1/3$ is constant because T_3 is uniquely ergodic. This shows that $\{n^2\}$ is not L^2 -exact.

The Proposition 4.5 still holds if we replace n^2 by a polynomial $P(n)$ such that $P(\mathbb{N}) \subset \mathbb{N}$. But d_i must be replaced by

$$d_i = \frac{\#\{0 \leq k < m : P(k) \equiv i \pmod{m}\}}{m}.$$

5. Θ -HARTMAN SEQUENCES ARE RECURRENT

Let $p \in \mathbb{Q}[x]$ be a polynomial such that $p(\mathbb{N}^*) \subset \mathbb{N}^*$ and $p(0) = 0$. For any irrational number t , $\{p(n)t\}$ is uniformly distributed mod 1 (Weyl theorem). It follows that $p(\mathbb{N}^*)$ is \mathbb{Q} -Hartman (see [4]).

Theorem 5.1. *Let $\Theta \subset \mathbb{Q}/\mathbb{Z}$ contain 0. Every Θ -Hartman sequence is recurrent.*

Since $D_{\mathbb{Q}/\mathbb{Z}} = \mathbb{N}$ and $D_{\{0\}} = \{1\}$, we have the following corollaries.

Corollary 5.2. *A subset $\Lambda = \{u_k\}$ of integers recurrent if the following condition is satisfied: for any integer m there is a subsequence $\{u_{n_k}\} \subset m\mathbb{Z} \cap \Lambda$ such that*

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{k=1}^N e^{2\pi i \alpha u_{n_k}} = 0, \quad \forall \alpha \in (0, 1) \setminus \mathbb{Q}.$$

Corollary 5.3. *A subset $\Lambda = \{u_k\}$ of integers recurrent if Λ contains a subsequence which is Hartman uniformly distributed. That is to say, there is a subsequence $\{u_{n_k}\} \subset \Lambda$ such that*

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{k=1}^N e^{2\pi i \alpha u_{n_k}} = 0, \quad \forall \alpha \in (0, 1).$$

5.1. Proof of Theorem 5.1. The proof of Theorem 5.1, inspired by [4], is divided into several lemmas, in the statements of which we will not repeat the conditions stated in Theorem 5.1.

The first lemma says that every function f in \mathcal{K}_Θ can be approximated by periodic elements under T with periods in D_Θ .

Lemma 5.4. *For any $f \in \mathcal{K}_\Theta$ and any $\epsilon > 0$, there exist an integer $m \in D_\Theta$ and a function $f' \in \mathcal{K}_\Theta$ with $T^m f' = f'$ such that $\|f - f'\| < \epsilon$.*

Proof. By the definition of \mathcal{K}_Θ , $\|f - f'\| < \epsilon$ holds for some finite linear combinations of eigenfunctions:

$$f' = \sum \alpha_j f_j, \quad \text{with } f_j \in \mathcal{K}_{\theta_j}, \theta_j \in \Theta$$

(α_j being a finite sequence of scalars). These numbers θ_j are rational. If we take m to be the least common multiple of the denominators of θ_j 's, we will have

$$T^m f' = \sum \alpha_j e^{2\pi i m \theta_j} f_j = \sum \alpha_j f_j = f'.$$

□

We denote by P_Θ the orthogonal projection onto \mathcal{K}_Θ . The next lemma show that the projection $P_\Theta f$ can be arbitrarily approximated by ergodic averages along with subsequences of Λ .

Lemma 5.5. *For any $f \in L^2(\mu)$ and any $\epsilon > 0$, there exists a subsequence $\{u_{n_k}\} \subset \Lambda$ such that*

$$\left\| \frac{1}{N} \sum_{k=1}^N T^{u_{n_k}} f - P_\Theta f \right\| < \epsilon$$

for sufficiently large N .

Proof. Write $f = f_\Theta + f_\Theta^\perp$ with $f_\Theta = P_\Theta f \in \mathcal{K}_\Theta$ and $f_\Theta^\perp \in \mathcal{K}_\Theta^\perp$. Apply Lemma 5.4 to f_Θ and $\epsilon/4$. Then there exist $m \in D_\Theta$ and $f'_\Theta \in \Theta$ such that

$$T^m f'_\Theta = f'_\Theta, \quad \|f_\Theta - f'_\Theta\| < \frac{\epsilon}{4}.$$

By the hypothesis on Θ and Lemma 5.4, we can find a subsequence $\{u_{n_k}\} \subset \Lambda$ which has the properties

- (i) $T^{u_{n_k}} f'_\Theta = f'_\Theta$;
- (ii) for all $\alpha \in \Theta^c$,

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{k=1}^N e^{2\pi i \alpha u_{n_k}} = 0.$$

By (i) and the contractivity of T , we have

$$\|T^{u_{n_k}} f_\Theta - f_\Theta\| \leq \|T^{u_{n_k}} (f_\Theta - f'_\Theta) - (f_\Theta - f'_\Theta)\| \leq \frac{\epsilon}{2}.$$

It follows immediately

$$\left\| \frac{1}{N} \sum_{k=1}^N T^{u_{n_k}} f_\Theta - f_\Theta \right\| \leq \frac{\epsilon}{2}.$$

On the other hand, by (ii) and the spectral lemma, we have

$$\left\| \frac{1}{N} \sum_{k=1}^N T^{u_{n_k}} f_\Theta^\perp \right\|^2 \leq \int_{\Theta^c} \left| \frac{1}{N} \sum_{k=1}^N e^{2\pi i \alpha u_{n_k}} \right|^2 d\mu_{f_\Theta^\perp} < \frac{\epsilon^2}{4}$$

if N is sufficiently large. Thus we conclude by combining the last two estimations. □

For $f \in L^2(\mu)$, let H_f^Λ be the subspace spanned by $T^u f$ with $u \in \Lambda$.

Lemma 5.6. *If $f \perp H_f^\Lambda$, then $f \perp \mathcal{K}_\Theta$.*

Proof. What we have to prove is that $\langle f, f_\Theta \rangle = 0$ where $f_\Theta = P_\Theta f$. For any $\epsilon > 0$, take the sequence $\{u_{n_k}\}$ in Lemma 5.5 such that $\|A_N f - f_\Theta\| < \epsilon$ for large N where $A_N f = N^{-1} \sum_{k=1}^N T^{u_{n_k}} f$. Write

$$\langle f, f_\Theta \rangle = \langle f, A_N f \rangle + \langle f, f_\Theta - A_N f \rangle.$$

Since the hypothesis of the lemma implies $\langle f, A_N f \rangle = 0$, we have

$$|\langle f, f_\Theta \rangle| \leq \|f\| \cdot \|f_\Theta - A_N f\| < \|f\| \epsilon.$$

□

Proof of Theorem 5.1. We prove the theorem by contradiction. Suppose that Λ is not a Poincaré set. Then there exists a measure preserving system (X, \mathcal{A}, T, μ) and a measurable set $A \in \mathcal{A}$ such that

$$\mu(A) > 0, \quad \mu(A \cap T^{-u} A) = 0 \quad (\forall u \in \Lambda).$$

Thus we have $1_A \perp H_f^\Lambda$ because

$$\mu(A \cap T^{-u} A) = \int 1_A \cdot 1_A \circ T^u d\mu.$$

By Lemma 5.6, we have $1_A \perp \mathcal{K}_\Theta$. Since $1 \in \mathcal{K}_\Theta$ (Θ contains 0), we have $\mu(A) = \langle 1_A, 1 \rangle = 0$, which is a contradiction. □

6. RANDOM SEQUENCES

Let $\{p_n\}_{n \geq 1}$ be a sequence of real numbers such that

$$0 \leq p_n \leq 1, \quad \sum_{n=1}^{\infty} p_n = \infty.$$

Let $\{\xi_n\}_{n \geq 1}$ be a sequence of independent Bernoulli random variables such that

$$P(\xi_n = 1) = p_n = 1 - P(\xi_n = 0).$$

The study object in this section is the random set of integers

$$W(\omega) = \{n \geq 1 : \xi_n(\omega) = 1\}.$$

This is almost surely (a.s. for short) an infinite subset of integers. The following is the main theorem in this section.

Theorem 6.1. *The sequence $W(\omega)$ is a.s. Hartman uniformly distributed if the following conditions are satisfied:*

$$(1) \quad \log N = o\left(\sum_{n=1}^N p_n\right), \quad (2) \quad \sum_{n=1}^N |p_n - p_{n+1}| = o\left(\sum_{n=1}^N p_n\right).$$

Let us first present some elementary properties of $W(\omega)$.

6.1. Basic properties of $W(\omega)$. The classical law of large numbers of Kolmogorov concerns independent sequences of identically distributed variables. The following proposition is a kind of law of large numbers. It can be proved by Doob's martingale convergence theorem.

Proposition 6.2. *Almost surely*

$$\lim_{N \rightarrow \infty} \frac{\sum_{n=1}^N \xi_n}{\sum_{n=1}^N p_n} = 1.$$

Proof. Let $s_n = \sum_{k=1}^n p_k$ and $s_0 = p_1$. Consider the martingale

$$M_n = \sum_{k=1}^n \frac{\xi_k - p_k}{s_k}.$$

It is easy to compute that

$$\mathbb{E}M_n^2 = \sum_{k=1}^n \frac{p_k(1-p_k)}{s_k^2} \leq \sum_{k=1}^n \frac{p_k}{s_k s_{k-1}} = \frac{1}{p_1} + \sum_{k=2}^n \left(\frac{1}{s_{k-1}} - \frac{1}{s_k} \right) \leq \frac{2}{p_1}.$$

By the martingale convergence theorem of Doob, the L^2 -bounded martingale M_n converges almost surely. In other words, the series $\sum_{k=1}^{\infty} \frac{\xi_k - p_k}{s_k}$ converges. We conclude by the Kronecker lemma. \square

Consequently, $W(\omega)$ admits a.s. its upper density

$$\overline{D}(W(\omega)) = \overline{\lim}_{N \rightarrow \infty} \frac{\#W(\omega) \cap [1, N]}{N} = \overline{\lim}_{N \rightarrow \infty} \frac{\sum_{n=1}^N p_n}{N}.$$

The density of $W(\omega)$ exists iff $\lim_{N \rightarrow \infty} N^{-1} \sum_{n=1}^N p_n$ exists.

An increasing sequence $\{u_n\} \subset \mathbb{N}^*$ is said to be *syndetic* if

$$\sup_n (u_{n+1} - u_n) < \infty.$$

Proposition 6.3. *Almost surely $W(\omega)$ is syndetic iff there exists an integer $\ell \geq 1$ such that*

$$\sum_{n=1}^{\infty} \prod_{j=0}^{\ell-1} (1 - p_{n+j}) < \infty.$$

Proof. Notice that $P(n \notin W(\omega)) = P(\xi_n = 0) = 1 - p_n$. Let

$$A_{n,\ell} = \{\omega : n \notin W(\omega), n+1 \notin W(\omega), \dots, n+\ell-1 \notin W(\omega)\}.$$

Then $P(A_{n,\ell}) = \prod_{j=0}^{\ell-1} (1 - p_{n+j})$. So, the convergence of the series, together with the Borel-Cantelli lemma, shows that $P(\overline{\lim}_n A_{n,\ell}) = 0$. That means, almost sure when n is sufficiently large, each interval $[n, n+\ell-1]$ contains a term of $W(\omega)$. So, $W(\omega)$ is syndetic.

Now suppose que the series diverges for each $\ell \geq 1$. Then, for each fixed ℓ , there exists $0 \leq r < \ell$ such that

$$\sum_{k=1}^{\infty} \prod_{j=0}^{\ell-1} (1 - p_{r+k\ell+j}) = \infty.$$

Observe that the events $A_{r+k\ell}$ ($k = 1, 2, \dots$) are independent and $P(A_{r+k\ell}) = \prod_{j=0}^{\ell-1} (1 - p_{r+k\ell+j})$. Then the divergence of the last series, together with the Borel-Cantelli lemma, implies $P(\overline{\lim}_k A_{r+k\ell}) = 1$. That means, almost sure there exists k (actually infinitely many), such that the interval $[r + k\ell, r + k\ell\ell - 1]$ contains no term of $W(\omega)$. Since ℓ is arbitrary, $W(\omega)$ is not syndetic. \square

6.2. Uniform norm of some random trigonometric polynomials.

The following theorem on the uniform norm of a random trigonometric polynomials will be useful in the proof of Theorem 6.1.

Theorem 6.4. *Let $\{\xi_n\}$ be a sequence of independent Bernoulli variables such that $P(\xi_n = 1) = p_n = 1 - P(\xi_n = 0)$. Suppose that*

$$\log N = o\left(\sum_{n=1}^N p_n\right).$$

There exists a constant $C > 0$ such that

$$P\left(\max_{t \in [0,1)} \left| \sum_{n=1}^N (\xi_n - p_n) e^{2\pi i n t} \right| \geq C \sqrt{\sum_{n=1}^N p_n \log N}\right) \leq \frac{1}{N^2}.$$

Proof. Similar estimates were first obtained by Salem and Zygmund when Rademacher sequence or Steinhaus sequence take the place of $\{\xi_n - p_n\}$. Kahane considered subnormal or subgaussian sequences. Fan and Schneider have replaced the subgaussian condition by another integrability condition. But for the centralized but biased Bernoulli sequence $\{\xi_n - p_n\}$, we can not directly apply these known results in [6, 3]. However, we can imitate [6].

First estimate the Laplace transform of $\xi_n - p_n$:

$$\mathbb{E} e^{\lambda(\xi_n - p_n)} = e^{-\lambda p_n} (1 + p_n(e^{-\lambda} - 1)) \leq e^{p_n(e^{-\lambda} - 1 - \lambda)}.$$

Using the fact that $e^{-\lambda} - 1 - \lambda \leq A\lambda^2/2$ for $|\lambda| \leq 1$ (A being a constant). Thus we have

$$\mathbb{E} e^{\lambda(\xi_n - p_n)} \leq e^{A p_n \lambda^2 / 2} \quad \text{if } |\lambda| \leq 1,$$

which could be called a local subnormal property. As in [6], we will take (with the notation in [6])

$$\lambda^2 = \frac{\log(2\rho\kappa)}{Ar} = \frac{\log(4\pi N^4)}{A \sum_{n=1}^N p_n},$$

We should notice that $\lambda \rightarrow 0$ when $N \rightarrow \infty$. \square

6.3. Proof of Theorem 6.1.

What we have to prove is

$$a.s. \quad \lim_{N \rightarrow \infty} \frac{\sum_{n=1}^N \xi_n e^{2\pi i n t}}{\sum_{n=1}^N \xi_n} = 0 \quad \forall t \in (0, 1).$$

By Lemma 6.3, we have only to show

$$(6.1) \quad a.s. \quad \lim_{N \rightarrow \infty} \frac{\sum_{n=1}^N (\xi_n - p_n) e^{2\pi i n t}}{\sum_{n=1}^N p_n} = 0 \quad \forall t \in (0, 1);$$

$$(6.2) \quad \lim_{N \rightarrow \infty} \frac{\sum_{n=1}^N p_n e^{2\pi i n t}}{\sum_{n=1}^N p_n} = 0 \quad \forall t \in (0, 1).$$

By Theorem 6.4, we have

$$(6.3) \quad a.s. \quad \max_{t \in [0, 1)} \left| \sum_{n=1}^N (\xi_n - p_n) e^{2\pi i n t} \right| = O \left(\sqrt{\sum_{n=1}^N p_n \log N} \right).$$

So, (6.1) follows immediately from (6.3) and the hypothesis $\log N = o(\sum_{n=1}^N p_n)$.

An Abel summation leads to

$$\left| \sum_{n=1}^N p_n e^{2\pi i n t} \right| \leq \frac{1}{|\sin \pi t|} \left(\sum_{n=1}^{N-1} |p_n - p_{n+1}| + p_N \right).$$

Then (6.2) follows from this estimate and the hypothesis $\sum_{n=1}^N |p_n - p_{n+1}| = o(\sum_{n=1}^N p_n)$. \square

Assume that $\sum_{n=1}^N |p_n - p_{n+1}| = o(\sum_{n=1}^N p_n)$. It is interesting to find a condition for $W(\omega)$ to be a.s. non L^2 -exact. In other words, we would like to find conditions such that a.s. there exists t such that

$$(6.4) \quad \lim_{N \rightarrow \infty} \frac{\sum_{n=1}^N (\xi_n - p_n) e^{2\pi i u_n t}}{\sum_{n=1}^N p_n} \neq 0.$$

Notice that for t fixed, the above limit is a.s. equal to zero (the same proof for Proposition 6.3 works). So, the point t in (6.4) must be a random point depending on ω . Such point does exist when $p_n = 1/n$ [8]. See [13, 14] for the study of $W(\omega)$ as thin sets in harmonic analysis. Another interesting problem is to study the multiple recurrence of $W(\omega)$.

6.4. L^2 -universality of $W(\omega)$. A sequence $\Lambda = \{u_k\} \in \mathbb{N}^*$ is said to be L^2 -universal if for any measure-preserving system (X, \mathcal{A}, μ, T) and any $f \in L^2(\mu)$,

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{k=1}^N f \circ T^{u_k} \quad \text{exists } \mu\text{-a.e.}$$

Bourgain and Boshernitzan proved the L^2 -universality of $W(\omega)$ under the conditions that the sequence $\{p_k\}$ decreasing and for any geometric sequence $\mathcal{N}_\rho = \{[\rho^k], k \geq 1\}$ ($\rho > 1$), we have

$$(6.5) \quad \sum_{N \in \mathcal{N}_\rho} \frac{\log N}{\sum_{k=1}^N p_k} < +\infty.$$

We remark that the monotonicity of $\{p_k\}$ can be replaced by the weaker condition

$$(6.6) \quad \overline{\lim}_{N \rightarrow +\infty} \frac{1}{\sum_{k=1}^N p_k} \left(N p_N + \sum_{k=1}^{N-1} k |p_k - p_{k+1}| \right) < \infty.$$

Recall that $\sum_1^n \xi_k \sim \sum_1^n p_k$ (Lemma 6.3). Write

$$\frac{1}{\sum_{k=1}^N p_k} \sum_{k=1}^N \xi_k f \circ T^k = \frac{1}{\sum_{k=1}^N p_k} \sum_{k=1}^N (\xi_k - p_k) f \circ T^k + \frac{1}{\sum_{k=1}^N p_k} \sum_{k=1}^N p_k f \circ T^k.$$

The condition (6.5) ensures the convergence of the first term on the right hand side. The monotonicity was used to prove the convergence of the second term as a consequence of the Birkhoff theorem. The monotonicity can be replaced by (6.6) because under the condition (6.6) we have

$$\lim_{n \rightarrow \infty} \frac{1}{N} \sum_{n=1}^N a_n = \ell \Rightarrow \lim_{n \rightarrow \infty} \frac{\sum_{n=1}^N p_n a_n}{\sum_{n=1}^N p_n} = \ell$$

for any sequence (a_n) .

REFERENCES

- [1] A. Bertrand-Mathis. Ensembles intersectifs et récurrence de Poincaré. *Israel J. Math.* **55** (1986), 184–198.
- [2] J. Bourgain. Pointwise ergodic theorems for arithmetic sets. With an appendix by the author, Harry Furstenberg, Yitzhak Katznelson and Donald S. Ornstein. *Inst. Hautes Etudes Sci. Publ. Math.* No. 69 (1989), 5–45.
- [3] A. H. Fan and D. Schneider. Une inégalité de Littlewood-Salem. *Ann. Inst. H. Poincaré Probab. Statist.* **39** (2003), no. 2, 193–216.
- [4] H. Furstenberg. Recurrence in Ergodic Theory and Combinatorial Number Theory. *Princeton University Press*, 1981.
- [5] S. Hartman. Remarks on equidistribution on non-compact groups. *Compositio* **16** (1964), 66–71.
- [6] J. P. Kahane. Some Series of Random Functions. *Cambridge University Press*, 1985.
- [7] J.P. Kahane and Y. Katznelson. Entiers aléatoires, ensembles de Sidon, densité dans le groupe de Bohr et ensembles d'analyticité. *C. R. Math. Acad. Sci. Paris* **345** (2007), no. 1, 21–24.
- [8] J. P. Kahane and Y. Katznelson. Entiers aléatoires et analyse harmonique. *J. Anal. Math.* 105 (2008), 363–378.
- [9] J. P. Kahane and Y. Katznelson. Distribution uniforme de certaines suites d'entiers aléatoires dans le groupe de Bohr. *J. Anal. Math.* 105 (2008), 379–382.
- [10] K. Kamae and M. Mendès France. van der Corput's difference theorem. *Israel J. Math.* **31** (1978), no. 3-4, 335–342.
- [11] U. Krengel. Ergodic Theorems. *W. de Gruyter*, 1989.
- [12] L. Kuipers and H. Niederreiter. Uniform distribution of sequences. Pure and Applied Mathematics. *Wiley-Interscience*, New York-London-Sydney, 1974.
- [13] D. Li, H. Queffélec and L. Rodríguez-Piazza. Some new thin sets of integers in harmonic analysis. *J. Anal. Math.* 86 (2002), 105–138.
- [14] D. Li, H. Queffélec and L. Rodríguez-Piazza. On some random thin sets of integers. *Proc. Amer. Math. Soc.* 136 (2008), no. 1, 141–150.

A. H. FAN: LAMFA, UMR 4160, CNRS, UNIVERSITY OF PICARDIE, 33 RUE SAINT LEU, 80039 AMIENS, FRANCE

E-mail address: `ai-hua.fan@u-picardie.fr`

URL: `http://www.mathinfo.u-picardie.fr/fan/`

D. SCHNEIDER: EA 2597, LABORATOIRE DE MATHÉMATIQUES PURES ET APPLIQUÉES, UNIVERSITE DE LITTORAL, CALAIS, FRANCE

E-mail address: `dominique.schneider@lmpa.univ-littoral.fr`