



HAL
open science

Hydrogels Based on Polysaccharides Grafted Ferulic Acid: A Biomimetic Approach

Ahdi Hadrich, Lecerf Didier, Virginie Dulong, Luc Picton

► **To cite this version:**

Ahdi Hadrich, Lecerf Didier, Virginie Dulong, Luc Picton. Hydrogels Based on Polysaccharides Grafted Ferulic Acid: A Biomimetic Approach. Priochem 2021, 2021, Bucarest, Romania. pp.69, <10.3390/chemproc2022007069>. <hal-03674089>

HAL Id: hal-03674089

<https://hal.science/hal-03674089v1>

Submitted on 12 May 2025

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire HAL, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons CC BY 4.0 - Attribution - International License

Abstract

Hydrogels Based on Polysaccharides Grafted Ferulic Acid: A Biomimetic Approach †

Ahdi Hadrich, Didier Le Cerf , Virginie Dulong  and Luc Picton * 

Polymères, Biopolymères, Surfaces (PBS) UMR6270 CNRS, Université de Rouen Normandie, 76821 Mont Saint Aignan, France; hadrich.ahdi@hotmail.fr (A.H.); didier.lecerf@univ-rouen.fr (D.L.C.); virginie.dulong@univ-rouen.fr (V.D.)

* Correspondence: luc.picton@univ-rouen.fr

† Presented at the 17th International Symposium “Priorities of Chemistry for a Sustainable Development” PRIOCHEM, Bucharest, Romania, 27–29 October 2021.

Keywords: hydrogels; biomimetic approach; artificial polysaccharides

Some cereal seeds present mucilage composed of polysaccharides bearing ferulic acid (FA) groups, which are capable of reacting in the presence of an enzyme such as laccase to generate chemical hydrogels [1]. The main idea of this work, following a biomimetic approach, is to elaborate artificial polysaccharides grafted with ferulic acid groups [2]. Various polysaccharides have been studied: anionic ones (carboxymethylpullulane (CMP) as a model, hyaluronan (HA) for application) and neutral ones (pullulan (P)). We report here the availability of the grafting, the evidence of crosslinking (Figure 1) leading to hydrogels in the presence of laccase (as a function of FA, polymer and laccase amount) and evidence of antioxidant properties of such derivatives [3].



Citation: Hadrich, A.; Le Cerf, D.; Dulong, V.; Picton, L. Hydrogels Based on Polysaccharides Grafted Ferulic Acid: A Biomimetic Approach. *Chem. Proc.* **2022**, *7*, 69. <https://doi.org/10.3390/chemproc2022007069>

Academic Editors: Mihaela Doni, Florin Oancea, Zina Vuluga and Radu Claudiu Fierăscu

Published: 28 April 2022

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).




Figure 1. G' vs. time for CMP-FA for various DSFA. Laccase at 2 nkat, 20 g L⁻¹ in polymer, 25 °C in citrate/phosphate buffer (0.1 mol L⁻¹ pH 5.5).

Author Contributions: Conceptualization, all; methodology, all; software, L.P.; validation, all; formal analysis, all; investigation, all; resources, all; data curation, all; writing—original draft preparation, all; writing—review and editing, L.P.; visualization, L.P.; supervision, L.P.; project administration, V.D. and L.P.; funding acquisition, V.D. and L.P. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Carvajal-Millan, E.; Landillon, V.; Morel, M.H.; Rouau, X.; Doublier, J.L.; Micard, V. Arabinoxylan gels: Impact of the feruloylation degree on their structure and properties. *Biomacromolecules* **2005**, *6*, 309–317. [[CrossRef](#)] [[PubMed](#)]
2. Dulong, V.; Hadrich, A.; Picton, L.; Le Cerf, D. Enzymatic cross-linking of carboxymethylpullulan grafted with ferulic acid. *Carbohydr. Polym.* **2016**, *151*, 78–87. [[CrossRef](#)] [[PubMed](#)]
3. Mathew, S.; Abraham, T.E. Ferulic acid: An antioxidant found naturally in plant cell walls and feruloyl esterases involved in its release and their applications. *Crit. Rev. Biotechnol.* **2004**, *24*, 59–83. [[CrossRef](#)]