

Brain mechanisms for foveating a visual target here-and-now (i.e., where it is and when it is there)

Laurent Goffart


CNRS

Laurent.Goffart@univ-amu.fr

Two types of goal-directed foveating movements

rapid : saccade

slow : smooth pursuit




Two types of goal-directed foveating movements

rapid : saccade

slow : smooth pursuit


Ultimately, they involve separate populations of neurons



The saccade :


a rapid and accurate transition between *dynamic equilibrium states*

Spatial trajectory




The saccade

a reaction shared by most animals



The saccade :


A transition which is *irreversible* and *goal-directed*




Sparks & Mays J Neurophysiol 1983

**observation made
in a primate (monkey)**


Extra-ocular muscles




Horizontal rotation




Upward rotation




Downward rotation




Vector decomposition and synergy




Synergy between H and V components



The horizontal saccade generator




Relaxation of antagonist muscles



Contraction of agonist muscles

LR : Lateral Rectus
MR : Medial Rectus


Relaxation of antagonist muscles



Contraction of agonist muscles


nVI = Abducens Nucleus
nIII = Oculomotor Nucleus

Phasic command




EBN = Excitatory
Burst Neurons
(Paramedian Pontine RF)


Phasic command




Phasic command




Phasic command




Phasic command




Reciprocal innervation (push-pull)




Reciprocal innervation (push-pull)




Reciprocal innervation (push-pull)




Bilateral phasic command




Bilateral phasic command




Bilateral phasic command







Tonic command (gaze holding)




Tonic command (gaze holding)






Fixation instability : saccade intrusions (unstable equilibrium)




L.Averbuch-Heller
A.A.Konnerot
K.G.Rothschild
L.E.Dell'Osso
B.F.Remmer
R.J.Leigh

Problem : no fixation instability in the monkey after RIP lesion/inactivation

OPN : a solution for synchronizing H and V components ?



Reduction of inhibition during pursuit




Missal & Keller JNP 2004


Generating a new saccade

Transmit a goal to EBN


Remove OPN inhibition




How to generate different amplitudes ?



Burst neuron (temporal code)




How to generate different amplitudes ?




How is the burst duration specified ?


The deep Superior Colliculus




For a given saccade, a population of active neurons




For a given saccade, a population of active neurons




For a given saccade, a population of active neurons




For a given saccade, a population of active neurons



For a given saccade, a population of active neurons




Topography of saccades evoked by electrical microstimulation




Gaze-related command (eye + head)

Microstimulation in the head-unrestrained monkey



Topographical encoding of goal for gaze

Muscimol injection (Gaba-A agonist)



No saccade with amplitude and direction corresponding to vector represented at inactivated site

Increase of latency for saccades represented at adjacent sites


The spatio-temporal transformation

HOW is
the position of active cells
transformed into a
a duration of burst ?

The dichotomist hypothesis


FIXATION / SACCADE ZONE

(MUNOZ and colleagues 1989;1993)



Mutual inhibition


The dichotomist hypothesis



The dichotomist hypothesis

FIXATION / SACCADE ZONE


(MUNOZ and colleagues 1989;1993)




Mutual inhibition

Dichotomy : MOVE versus DO NOT MOVE


The "moving hill" hypothesis (Munoz, Pélisson & Guitton *Science* 1991)




The "moving hill" hypothesis (Munoz, Pélisson & Guitton *Science* 1991)




The "moving hill" hypothesis (Munoz, Pélisson & Guitton *Science* 1991)




The "moving hill" hypothesis (Munoz, Pélisson & Guitton *Science* 1991)




The "moving hill" hypothesis (Munoz, Pélisson & Guitton *Science* 1991)




The "moving hill" hypothesis (Munoz, Pélisson & Guitton *Science* 1991)




The "spreading wave" hypothesis (Munoz & Wurtz *J. Neurophysiol.* 1993)




The "spreading wave" hypothesis (Munoz & Wurtz *J. Neurophysiol.* 1993)




The "spreading wave" hypothesis (Munoz & Wurtz *J. Neurophysiol.* 1993)



The "spreading wave" hypothesis (Munoz & Wurtz *J. Neurophysiol.* 1993)




“moving hill” or “spreading wave” hypotheses
=> assume a dichotomy FIX/SAC



1. Recording of FIX
→ sustained activity during fixation
2. Microstimulation of FIX
→ delays the onset of saccades (+ hypometria)
3. Inactivation of FIX
→ irrepressible saccades


dichotomy

IF assumption (dichotomy) is wrong
THEN WHAT ?




dichotomy


IF assumption (dichotomy) is wrong
THEN the SC is a homogeneous map




IF assumption (dichotomy) is wrong
THEN the SC is a homogeneous map











Fixation is not a "rigid" / "frozen" state







Fixation is the maintenance of gaze within a spatio-temporal window












Fastigio-reticular projections and the cerebellar control of saccade amplitude




Bilateral cerebellar control of the dynamic balance between EBN and IBN during the saccadic transition






Does the sustained activity in cFN influence anything ?



Deficit in fixation after cFN inactivation



Ipsilesional fixation offset



J Neurophysiol 103: 1988–2001, 2010.
First published February 3, 2010; doi:10.1152/jn.00771.2009.

Fastigial Oculomotor Region and the Control of Foveation During Fixation


Lorenzo Guerrasio,¹ Julie Quinet,² Ulrich Büttner,¹ and Laurent Goffart³

¹Department of Neurology, Klinikum Großhadern, LMU, München, Germany; ²Unité 534, Institut National de la Santé et de la Recherche Médicale/Université Claude Bernard-Lyon 1, IFR 19 Institut Fédératif des Neurosciences de Lyon, Bron, France; and ³Institut de Neurosciences Cognitives de la Méditerranée, Unité Mixte Recherche 6193, Centre National de la Recherche Scientifique, Aix-Marseille Universités, Marseille, France


Journal of Neurophysiology 2010

"Given the general role of the cerebellum in motor learning (Ito 1984), the FORs could play a specific role in the learning phase of the foveating behavior, compensating for idiosyncratic differences in neuromuscular morphology and anisotropies in the mechanisms triggering and executing fixational saccades."

Subcortical control of foveation



IF fixation is the maintenance of gaze within a spatio-temporal window
THEN how do we foveate a moving target ?




The *hic-et-nunc* (here-and-now) of saccade endpoints

Animals can make saccadic eye movements to visually intercept the trajectory of a moving object and foveate it *at the right place and time*, and initiate its pursuit.


Such interceptive saccades indicate that, in spite of variable sensorimotor delays, the brain is able to estimate the ***hic-et-nunc* (here-and-now)** coordinates of a target at least at the time of saccade landing.

How is this possible ?


From the behavior to its analysis




From the behavior to its analysis



From the behavior to its analysis




The dual-drive hypothesis



ASSUMPTIONS :

1. Snapshot of Eye Position Error (EPE)
→ discrete sampling


The dual-drive hypothesis




ASSUMPTIONS :

1. Snapshot of Eye Position Error (EPE)
→ discrete sampling
2. Estimate of Target Displacement (TD)
→ time window delimited by discrete events

The dual-drive hypothesis




The dual-drive hypothesis




The dual-drive hypothesis

Keller et al. JNP 1996
Optican & Quaiia ANYAS 2002
Optican Encyclopedia of Neurosc. 2010



An alternative hypothesis is possible

'Dual drive' hypothesis



The "dynamic remapping" hypothesis

Saccadic Foveation of a Moving Visual Target in the Rhesus Monkey

Jérôme Fleuriet, Sandrine Hugues, Laurent Perrinet, and Laurent Goffart
Institut de Neurosciences Cognitives de la Méditerranée, Unité Mixte de Recherche 6193, Centre National de la Recherche Scientifique, Aix-Marseille Universités, Marseille, France

Journal of Neurophysiology 2011

The "*hic-et-nunc*" hypothesis


Saccadic Interception of a Moving Visual Target after a Spatiotemporal Perturbation

Jérôme Fleuriet and Laurent Goffart
Institut de Neurosciences Cognitives de la Méditerranée, UMR 6193, Centre National de la Recherche Scientifique, Aix-Marseille Université, 13393 Marseille Cedex 03, France


Journal of Neuroscience 2012

An alternative hypothesis

'Dual drive' hypothesis




'Hic-et-nunc' hypothesis




Experimental test in the monkey

Perturb the interceptive saccade and test the ability to keep track of the moving target and intercept it




Predictions


IF the saccade goal = *hic-et-nunc* target position
THEN change in eye position and added delay should be taken into account (accurate saccades)




Visual feedback is removed



Saccadic interception of an occluded target




Perturbation of interceptive saccades



Fleuriet & Goffart JN 2012

Perturbation of interceptive saccades


Microstimulation in the right deep Superior Colliculus (30ms 400Hz 12µA)



Fleuriet & Goffart JN 2012

Perturbation of interceptive saccades

Microstimulation in the right deep Superior Colliculus (30ms 400Hz 12 μ A)



Conclusions

In spite of the "spatio-temporal blurring" (Pellioniz & Llinas 1982) that a visual event undergoes when the retinal activity propagates toward the oculomotor system, *in fine*, the brain is able to estimate the here-and-now (*hic-et-nunc*) coordinates of the target.

Fleuriet & Goffart (2012) showed that this estimate is *robust* enough for animals to capture the target even after an unexpected spatiotemporal perturbation (i.e., a change in eye position and associated delay).

Working hypothesis

Gaze direction is driven by a *flow of expected target positions*

This flow is an *internal model* that has been *learned* and *embodied*.

- There is no need for encoding the location of the target in the future.
- The brain needs to build up an estimate of *expected here-and-now* target position.

Obviously, this estimate is not stationary, the oculomotor system reads out a signal out of a memory (register) that *temporally unfolds*.

This dynamic signal feeds the same local feedback loop involved in the generation of saccades toward a stationary target.

On-going work

Investigations are being performed

with **Julie Quinet** and **Clara Bourrelly** to study :

- the build-up of this "dynamic attractor" and its long-term storage
- its spatiotemporal conformation with the on-going event (*dynamic morphing*)




with **Neeraj Gandhi**, University of Pittsburgh to test :

- whether an encoding of the "here-and-now" coordinates can be found in the deep superior colliculus

with **Yoshifumi Yamawaki**, Kyushu University to understand :


- the neurobiological foundations of *what is going on here-and-now* in a much simpler organism (praying mantis)

Building-up an internal model of expected target positions




Blocked trials

Building-up an internal model of expected target positions




Blocked trials

Building-up an internal model of expected target positions




Building-up an internal model of expected target positions



The putative "internal model" can not be reduced to a velocity / trajectory storage

Do we really need an internal model to explain this?



What internal model are we looking for ?


Our working hypothesis : Gaze direction is driven by a **flow of expected target positions here-and-now**

This flow is an internal model that has been *learned* (requires training) and *embodied* (some long-term maintenance)


- an estimate of *expected here-and-now* target position,
- a non stationary estimate that is read out by the oculomotor system and that *temporally unfolds* (observable).

Does such an internal model exist ?


Evidence for an internal model of here-and-now




Evidence for an internal model of here-and-now



Evidence for an internal model of here-and-now



Evidence for an internal model of here-and-now



*"L'expérience ne nous donne qu'un certain nombre de points isolés ;
il faut les réunir par un trait continu"*

Henri Poincaré



*Il faut les réunir en se gardant de ne pas faire trop d'extrapolations
lorsqu'il s'agit d'envisager l'avenir*

.....
hic-et-nunc