



HAL
open science

Report SEME for SCALIAN

Grégoire Barrué, Riccardo Di Dio, Zakarya El Khiyati, Yann Guguen, Roméo Tayewo

► **To cite this version:**

Grégoire Barrué, Riccardo Di Dio, Zakarya El Khiyati, Yann Guguen, Roméo Tayewo. Report SEME for SCALIAN. [Research Report] IRMAR - Université Rennes 1; Université Bretagne Sud; Université Cote d'Azur. 2022, pp.1-4. hal-03674181

HAL Id: hal-03674181

<https://hal.science/hal-03674181v1>

Submitted on 20 May 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Report SEME for SCALIAN

Grégoire Barrué¹, Riccardo Di Dio^{2, 3}, Zakarya El Khiyati³, Yann Guguen¹, and Roméo Tayewo⁴

¹*ENS Rennes*

²*Université Côte d'Azur, CNRS, LJAD, VADER Center, France*

³*INRIA, Sophia Antipolis, France*

⁴*Université de Bretagne Sud*

May 20, 2022

In this report we present the procedure that we set up to try to solve a mathematical problem proposed by Scalian. We work in the context of reduced order models for wind turbine applications. The objective is to improve a part of the algorithm developed by the company, in order to make the wind turbine blades more reactive.

The equation and its discretization

The reduced order model under location uncertainty used for this problem gives us stochastic differential equations for resolved modes denoted by $b = (b_i)_{i \in [1, n]}$:

$$\begin{aligned} db_i(t) &= F_i(b(t)) + (\alpha_{\cdot i} \cdot dB_t)^T b(t) + (\theta_{\cdot i} \cdot dB_t) \\ &= F_i(b(t)) + \sum_{j=1}^n \left(\sum_{k=1}^n \alpha_{kij} b_k(t) + \theta_{ij} \right) dB_t^j. \end{aligned} \quad (1)$$

In this equation F_i is a polynomial function of order two, and $B_t = (B_t^j)_{j \in [1, n]}$ is a vector composed of Brownian motions. In the algorithm of the company, these stochastic differential equations are solved using a Euler-Maruyama numerical scheme. In order to improve the algorithm, we chose to change the numerical scheme implemented for these equations, and to implement a Milstein scheme. We rewrite the (1) under a more general form:

$$db(t) = F(b(t)) + \sigma(b(t))dB_t, \quad (2)$$

where $F(b) = (F_i(b))_{i \in [1, n]}$ and

$$\sigma(b) = \left(\sum_{k=1}^n \alpha_{kij} b_k(t) + \theta_{ij} \right)_{i, j \in [1, n]} = (\sigma_{ij})_{i, j \in [1, n]}.$$

We discretize t in the interval $[0, T]$ and consider $N \in \mathbb{N}$, and $\Delta t = \frac{T}{N}$. Let $(\gamma_k)_{k \in [0, N]}$ be a sequence of vector $(\gamma_k^j)_{j \in [1, n]}$ such that all the γ_k^j are independent identically distributed standard Gaussian random variables. The Milstein numerical scheme is the following:

$$\begin{aligned} b(t_{k+1}) = & b(t_k) + F(b(t_k))\Delta t + \sigma(b(t_k))\gamma_k\sqrt{\Delta t} \\ & + \frac{1}{2}(\sigma, \partial\sigma)(b(t_k))\gamma_k^2\Delta t - \frac{1}{2}(\sigma, \partial\sigma)(b(t_k))\mathbb{1}\Delta t. \end{aligned} \quad (3)$$

In this scheme, $\mathbb{1}$ denotes a vector with 1 as coefficients, and $(\sigma, \partial\sigma)(b(t_k))$ denotes a matrix such that

$$(\sigma, \partial\sigma)(b(t_k))_{ij} = \sum_{k=1}^n \sigma_{ik} \frac{\partial\sigma_{ij}}{\partial\sigma_{ik}}.$$

Procedure and results

The aim is to compare the efficiency of the scheme (3) with the classical Euler-Maruyama scheme. First, we need to be sure that the Milstein scheme approximates the solution of (2), so we compare it with a reference solution, obtained using the Euler-Maruyama scheme with Δt very small and a very large number of simulations. We denote this reference solution b^{ref} . The idea is to compare the strong error between b^{ref} and the Euler-Maruyama scheme for larger values of Δt

$$\mathbb{E} \left[\sup_{t_k} \|b^{ref}(t_k) - b^{EM}(t_k)\|_2 \right]$$

to the strong error between b^{ref} and the Milstein scheme for the same values of Δt

$$\mathbb{E} \left[\sup_{t_k} \|b^{ref}(t_k) - b^{Mil}(t_k)\|_2 \right].$$

This comparison allows us to find the time-step size where the Milstein scheme becomes closer to the reference solution than the Euler-Maruyama scheme. Then we compare the execution time for the two schemes, for several values of Δt . We compute the strong errors with 100 simulations for each scheme, and Figure 1 shows that the Milstein scheme is slower than the Euler-Maruyama scheme. It makes sense since this scheme is more complex than the intuitive Euler-Maruyama scheme. The strategy is to think that we will need a larger Δt for the Milstein scheme than for the Euler-Maruyama scheme, and that the execution time of the Milstein scheme for this larger Δt will be smaller than the execution time of the Euler-Maruyama scheme with the smaller Δt .

The strong error is represented by Figure 2. On this figure we can not observe any visible difference between the two schemes. Hence it does not prove our theory, but it can be due to the number of simulations which is only 100, and it could be good to increase it or to run other tests on these schemes. It could also be interesting to try other schemes like the Runge-Kutta or the multi-step

scheme. The other possibility is that the regime that we studied in Figure 2 is already a regime where Euler-Maruyama and Milstein are close, so it could be interesting to compute the strong error with $N \in \llbracket 0, 50 \rrbracket$. Finally, we do not control the influence of the variance, which creates a Monte-Carlo error in addition to the discretization error. This could be a problem and one can try to use variance reduction methods to improve the algorithm.




Figure 1: Execution time for the Milstein and Euler-Maruyama schemes, for 100 simulations and several values of Δt .




Figure 2: Strong error for the Milstein and Euler-Maruyama schemes, for 100 simulations and several values of Δt .

For variance reduction techniques one can refer to [3] [2] [7]

References

- [1] Sigal Gottlieb. On high order strong stability preserving Runge-Kutta and multi step time discretizations. *J. Sci. Comput.*, 25(1-2):105–128, 2005.
- [2] Oindrila Kanjilal and CS Manohar. Girsanov’s transformation based variance reduced monte carlo simulation schemes for reliability estimation in nonlinear stochastic dynamics. *Journal of Computational Physics*, 341:278–294, 2017.
- [3] Ahmed Kebaier. Statistical romberg extrapolation: a new variance reduction method and applications to option pricing. *The Annals of Applied Probability*, 15(4):2681–2705, 2005.
- [4] G. N. Mil štein. Approximate integration of stochastic differential equations. *Teor. Verojatnost. i Primenen.*, 19:583–588, 1974.
- [5] Valentin Resseguier, Matheus Ladvig, and Dominique HEITZ. Real-time estimation and prediction of unsteady flows using reduced-order models coupled with few measurements. working paper or preprint, March 2022.
- [6] Valentin Resseguier, Agustin M. Picard, Etienne Memin, and Bertrand Chapron. Quantifying truncation-related uncertainties in unsteady fluid dynamics reduced order models. *SIAM/ASA J. Uncertain. Quantif.*, 9(3):1152–1183, 2021.
- [7] JGM Schoenmakers, AW Heemink, K Ponnambalam, and PE Kloeden. Variance reduction for monte carlo simulation of stochastic environmental models. *Applied Mathematical Modelling*, 26(8):785–795, 2002.