



**HAL**  
open science

# Occurrence, origin and potential ecological risk of dissolved polycyclic aromatic hydrocarbons and organochlorines in surface waters of the Gulf of Gabès (Tunisia, Southern Mediterranean Sea)

Hatem Zaghden, Badreddine Barhoumi, Lobna Jlaiel, Catherine Guigue, Lassaad Chouba, Soufiane Touil, Sami Sayadi, Marc Tedetti

## ► To cite this version:

Hatem Zaghden, Badreddine Barhoumi, Lobna Jlaiel, Catherine Guigue, Lassaad Chouba, et al.. Occurrence, origin and potential ecological risk of dissolved polycyclic aromatic hydrocarbons and organochlorines in surface waters of the Gulf of Gabès (Tunisia, Southern Mediterranean Sea). *Marine Pollution Bulletin*, 2022, 180, pp.113737. 10.1016/j.marpolbul.2022.113737 . hal-03675064

**HAL Id: hal-03675064**

**<https://hal.science/hal-03675064v1>**

Submitted on 15 Feb 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Occurrence, origin and potential ecological risk of dissolved polycyclic aromatic**  
2 **hydrocarbons and organochlorines in surface waters of the Gulf of Gabès (Tunisia,**  
3 **Southern Mediterranean Sea)**

4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

5 Hatem Zaghden<sup>a,b\*</sup>, Badreddine Barhoumi<sup>a,c\*</sup>, Lobna Jlaiel<sup>a</sup>, Catherine Guigue<sup>d</sup>, Lassaad  
6 Chouba<sup>e</sup>, Soufiane Touil<sup>c</sup>, Sami Sayadi<sup>f\*</sup>, Marc Tedetti<sup>a,d\*</sup>

8 <sup>a</sup> Laboratory of Environmental Bioprocesses, Centre of Biotechnology of Sfax, BP 1177,  
9 3018 Sfax, Tunisia

10 <sup>b</sup> Laboratory of Plant Molecular Physiology, Centre of Biotechnology of Borj-Cédria, 2050  
11 Hammam-Lif, Tunisia

12 <sup>c</sup> Laboratory of Hetero-Organic Compounds and Nanostructured Materials (LR18ES11),  
13 Department of Chemistry, Faculty of Sciences of Bizerte, University of Carthage, 7021  
14 Zarzouna, Tunisia

15 <sup>d</sup> Aix Marseille Univ., Université de Toulon, CNRS, IRD, MIO UM 110, 13288, Marseille,  
16 France

17 <sup>e</sup> Laboratory of Marine Environment, National Institute of Marine Science and Technology  
18 (INSTM), Goulette, Tunisia

19 <sup>f</sup> Biotechnology Program, Center for Sustainable Development, College of Arts and Sciences,  
20 Qatar University, Doha 2713, Qatar

21  
22 \*Corresponding authors:

23 E-mail: [hatem.zaghden@gmail.com](mailto:hatem.zaghden@gmail.com) (H. Zaghden)

24 E-mail: [barhoumibadredine@yahoo.fr](mailto:barhoumibadredine@yahoo.fr) (B. Barhoumi)

25 E-mail: [sami.sayadi@gmail.com](mailto:sami.sayadi@gmail.com) (S. Sayadi)

26 E-mail: [marc.tedetti@mio.osupytheas.fr](mailto:marc.tedetti@mio.osupytheas.fr) (M. Tedetti)

27 **Abstract**

28 We investigated the occurrence, origin, and potential ecological risk of dissolved polycyclic  
29 aromatic hydrocarbons (PAHs), polychlorinated biphenyl (PCBs) and organochlorine  
30 pesticides (OCPs) in 27 surface water samples collected from a highly anthropized and  
31 industrialized area in the Gulf of Gabès (Tunisia, Southern Mediterranean Sea) in October-  
32 November 2017. The results demonstrated a wide range of concentrations (ng L<sup>-1</sup>) with the  
33 following decreasing order:  $\Sigma 16$  PAHs (17.6-71.2) >  $\Sigma 20$  PCBs (2.9-33.7) >  $\Sigma 6$  DDTs (1.1-  
34 12.1) >  $\Sigma 4$  HCHs (1.1-14.8). Selected diagnostic ratios indicated a mixture of both pyrolytic  
35 and petrogenic sources of PAHs, with a predominance of petrogenic sources. PCB  
36 compositions showed distinct contamination signatures for tetra- to hepta-chlorinated PCBs,  
37 characteristic of contamination by commercial (Aroclor) PCB mixtures. The dominant OCP  
38 congeners were  $\gamma$ -HCH, 2,4'-DDD and 2,4'-DDE, reflecting past use of Lindane and DDTs in  
39 the study area. Agricultural, industrial and domestic activities, as well as atmospheric  
40 transport are identified as potential sources of PAHs, PCBs and OCPs in surface waters of the  
41 Gulf of Gabès. Toxic equivalents (TEQs) suggested a low carcinogenic potential for PAHs in  
42 seawater samples (mean of 0.14 ng TEQ L<sup>-1</sup>). Evaluation of risk coefficients revealed low risk  
43 for PAHs and PCBs, and moderate to severe risk for OCPs.

44 **Keywords:** PAHs, PCBs, OCPs, ecological risk, Gulf of Gabès, Mediterranean Sea

## 52 1. Introduction

53 Polycyclic aromatic hydrocarbons (PAHs), polychlorinated biphenyls (PCBs) and  
54 organochlorine pesticides (OCPs) are widespread in the environment (e.g. water, particulate  
55 matter, sediment, soil and organisms) and have been a focus of attention due to their  
56 persistence nature, bioaccumulation, toxicity, carcinogenicity and long range transport (Lee et  
57 al., 2016; Miraji et al., 2021; Nguyen et al., 2014; Nyarko et al., 2011). They are a serious  
58 threat to both ecosystems and human health. As a consequence, many of these compounds are  
59 listed as priority pollutants by international organizations (Ashraf, 2017; Guo et al., 2019;  
60 Jones and de Voogt, 1999; Muir and Howard, 2006). PCBs and OCPs have been widely used  
61 for industrial and agricultural purposes, while PAHs are accidental byproducts of combustion.  
62 Therefore, they can be used as chemical tracers for their possible sources (Lin et al., 2013).  
63 These pollutants can enter the ocean by various vectors including rivers, effluents, runoff and  
64 the atmosphere (Barhoumi et al., 2018; Moreno-González et al., 2013; Tolosa et al., 1997). In  
65 marine waters, they can undergo diverse processes, such as adsorption onto particles and  
66 subsequent sinking to sediments, adsorption on/absorption in biota, and accumulation into the  
67 food chains (Jamieson et al., 2017; Luo et al., 2014; Ma et al., 2015). Generally, their fate in  
68 the environment is strongly influenced by their physical–chemical properties (Mackay et al.,  
69 1992). Due to their hydrophobicity, PAHs, PCBs and OCPs have a tendency to partitioning  
70 between the dissolved phase of seawater/coastal waters and particles/sediment (Mhadhbi et  
71 al., 2019; Zhang et al., 2013; Rabodonirina et al., 2015). When sediment resuspension occurs  
72 due to changes in hydrodynamic processes such as waves, tides and currents, trawling and  
73 ship traffic, deposited contaminants in sediments may be again reintroduced to the overlying  
74 water, thus causing a secondary contamination (Zhao et al., 2021). In the dissolved state, these  
75 compounds may be subjected to quite strong dynamics, i.e., exchange with the atmosphere  
76 (deposition, volatilization), photodegradation, interactions with dissolved and particulate

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

77 organic matter, or transport with water masses. In addition, from their dissolved form, they  
78 may bioconcentrate in a variety of micro- and macro-organisms *via* respiration or passive  
79 diffusion (Dachs et al., 2002; Dachs and Méjanelle, 2010; Daewel et al., 2020; Duran and  
80 Cravo-Laureau, 2016; Montuori et al., 2014; Salvadó et al., 2019). Therefore, the knowledge  
81 of the distribution of dissolved PAHs, PCBs and OCPs is an essential step to better  
82 understand and quantify their global budget and their potential impact on a marine  
83 environment.

84 The Mediterranean Sea, a semi-closed aquatic ecosystem, is particularly vulnerable to  
85 chemical pollution due mainly to the long history of industrialization of its basin, its high  
86 population density, and its natural characteristics (such as large watersheds, high continental  
87 inputs, intense and extreme meteorological events, thermohaline circulation) (The Mermex  
88 group, 2011). As consequence, a multitude of chemical pollutants derived from urban,  
89 agricultural and industrial activities find their way into the Mediterranean waters (Campillo et  
90 al., 2019; Gómez-Gutiérrez et al., 2007; Köck-Schulmeyer et al., 2021; Sánchez-Avila et al.,  
91 2012). The Gulf of Gabès (Southeastern Tunisia) is one particular ecoregion of interest within  
92 the Mediterranean Sea basin due to its hydrodynamics, biogeochemical and biological  
93 features (Ayata et al., 2018; D’Ortenzio and Ribera d’Alcalà, 2009). It is one of the most  
94 productive areas of the Mediterranean, characterized by hydrodynamics driven by tides and  
95 low bathymetry (Béjaoui et al., 2019; Hattab et al., 2013). While the Gulf of Gabès  
96 contributes approximately 40% of the national fish production in Tunisia (DGPA, 2015), the  
97 coastal zone of the Gulf of Gabès is subjected to high anthropogenic pressures due to  
98 numerous industrial and agricultural activities within the zone. The latter are mainly  
99 concentrated in great coastal urban centers (Gabès and Sfax cities), which host fishing and  
100 commercial harbours, phosphoric acid production plants, municipal landfills, and various  
101 industries (Fourati et al., 2018; Nedja et al., 2011; Zaghden et al., 2007). Despite its

102 ecological and biogeochemical importance within the Mediterranean Sea, and at the same  
103 time, its high level of anthropogenic pressure, the Gulf of Gabès remains very poorly  
104 documented in terms of concentrations of organic pollutants in the dissolved phase of  
105 seawater, in particular organochlorine (OCs) compounds such as PCBs and OCPs. In this  
106 context, the objectives of this work were i) to describe the spatial distribution of dissolved  
107 PAHs, PCBs and OCPs (concentrations, molecular patterns) in surface coastal waters of the  
108 Ghannouch-Gabès area (Gulf of Gabès, Tunisia) and compare this distribution with that of  
109 core physico-chemical and biogeochemical parameters, ii) to assess the main origins/sources  
110 of these organic contaminants, and iii) to determine their potential ecological risk. To our  
111 knowledge, this work is the first to provide together levels of dissolved PAHs, PCBs and  
112 OCPs in coastal marine waters of Tunisia.

113

## 2. Material and Methods

### 2.1. Study area

114 The Gulf of Gabès is located in the Southeastern part of Tunisia (Southern Mediterranean  
115 Sea), and spreads over 400 km of coastline (more than 50% of the Tunisian coastline)  
116 between the city of Chebba in the North and the Tunisian-Libyan border in the South (Fig. 1).  
117 The Gulf of Gabès, which accounts for ~ 30% of the Tunisian coastal waters, is characterized  
118 by a wide continental shelf, shallow waters (isobaths 50-m deep are reached 110-km  
119 offshore), high amplitude (up to 2-m) tides, and hydrodynamics driven by tides and  
120 anticyclonic winds (Béjaoui et al., 2019; Zayen et al., 2020). The Gulf of Gabès hosts coastal  
121 cities, such as Sfax and Gabès, with concentrated industrial, harbour and agriculture activities.  
122 The Ghannouch-Gabès area in the central part of the Gulf, holds one of the biggest industrial  
123 zones in Tunisia. This industrial complex includes the Ghannouch commercial harbour, the  
124 phosphoric acid industry of the Tunisian Chemical Group, the Gabès Fluorine Chemical

127 Industry, a thermal power plant, petrochemical infrastructures, as well as industries related to  
128 agribusiness, building materials, energy production and textiles (El Zrelli et al., 2015, 2018;  
129 Fourati et al., 2018). The phosphoric acid industry of the Ghannouch industrial complex has  
130 been discharging the phosphogypsum, an acidic by-product of phomanufacturing, into the  
131 Gabès coastal waters over the past several decades. The solid phosphogypsum issued from the  
132 production of phosphoric acid, diluted with seawater to form gypsum water, is conveyed by  
133 several pipes to the South of the Ghannouch commercial harbour where the gypsum water  
134 mixes with the cooling water discharged by all the chemical industry units, and finally ends  
135 up at sea through a unique canal (Darmoul et al., 1980). Between 1000 and 13,000 t of  
136 phosphogypsum is released daily this way into the sea (Béjaoui et al., 2004; El Zrelli et al.,  
137 2015). In addition to this phosphogypsum discharges, several wadis carry a variety of urban,

699

## 2.2. Sampling and in situ measurements

701 Samples were collected at 27 stations in the Ghannouch-Gabès coastal area during the  
702 MERITE cruise from 30 October to 3 November 2017. Stations were distributed along  
703 coastal-offshore transects (also corresponding to North-South transects), i.e., S1-S3, S20 in  
704 front of El Melah wadi (Northern Ghannouch), S4-S6, S21 in front of El Demna wadi  
705 (Northern Ghannouch), S7-S12, S23 in front of Ghannouch industrial complex/commercial  
706 harbour and phosphogypsum effluent outlet, S13-S15 in front of fishing harbour and Gabès  
707 wadi, and S16-S18, S27 in front of Mersit wadi (Southern Gabès). Also, S19 was positioned  
708 as the north most station, S22 close to the Ghannouch industrial complex, in front of Ettine  
709 wadi/fluorine discharges, S24 and S25 between the Ghannouch industrial complex and the  
710 fishing harbour, and S26 in front of Griaa wadi (Fig. 1, Table S1). Stations S1-S18 (> 5-m  
711 depth) were sampled on board the vessel “Rahma” on 30 October (S1-S6), 1 November (S7-  
712 S12) and 3 November (S13-S18). The most coastal stations S19-S27 (< 5-m depth) were

1  
2 714 sampled on board a zodiac-type vessel on 31 October (S23-S27) and 2 November (S19-S22).  
3  
4 715 Stations S1-S11 and S16 were sampled at high tide, while stations S12-S15 and S17-S27 were  
5  
6 716 sampled at low tide. Sunny weather and very calm sea conditions prevailed during the cruise,  
7  
8 717 except during the sampling of stations S13-S18 (3 November) for which windy weather and  
9  
10 rough sea conditions were encountered (Fig. 1, Table S1).

11  
12 718 At each station, seawater samples for the analyses of dissolved organic pollutants (PAHs,  
13  
14 719 PCBs and OCPs) and other biogeochemical parameters were collected at ~ 0.5-m depth using  
15  
16 720 a LIMNOS water sampler (Hydro-Bios) equipped with two Schott® glass bottles of 1-L  
17  
18 721 volume each. This sampler allowed the bottles to go down in water closed and thus avoid the  
19  
20 722 sampling of the surface microlayer. The glass bottles were rinsed several times with sea water  
21  
22 723 before filling. Once filled, they were unscrewed from the sampler, closed with their Teflon  
23  
24 724 lined caps, and stored in the dark at 4 °C. Before sampling, the glass bottles were extensively  
25  
26 725 washed with 1 M hydrochloric acid (HCl) and ultrapure (resistivity 18.2 MΩ cm) water, and  
27  
28 726 baked at 450 °C for 6 h.

29  
30  
31  
32 727 Besides sampling, *in situ* profiles of temperature, salinity, concentration of total  
33  
34 728 chlorophyll *a* (TChla) and turbidity were conducted in the water column (from 0.5-m depth to  
35  
36 729 the bottom) using a 19 *plus* SBE conductivity-temperature-depth (CTD) profiler (SeaBird  
37  
38 730 Electronics, Inc.) equipped with a WETStar Chla fluorometer (WETLabs, Inc.) and a STM  
39  
40 731 turbidity meter (Seapoint, Inc.). Moreover, a Tripod multiparameter probe (Aqualabo) was  
41  
42 732 deployed in surface waters (at ~ 0.5-m depth) for the measurement of temperature, salinity,  
43  
44 733 pH, dissolved oxygen (O<sub>2</sub>) concentration and redox potential (redox). Sensors were calibrated  
45  
46 734 just before the cruise according to manufacturer's procedures.

### 53 54 735 **2.3. Filtration of samples**

55  
56  
57 736 Once in the laboratory, seawater samples were immediately filtered under a low vacuum (<  
58  
59 737 50 mm Hg) through pre-baked (450 °C, 6 h) GF/F (~ 0.7 μm) glass fiber filters (25- or 47-mm



738 diameter, Whatman) using glassware filtration systems. For dissolved organic pollutants  
739 (PAHs, PCBs, OCPs), 1 L of sample was filtered through 47-mm diameter GF/F filters.  
740 Filtered samples were stored in 2.5 L glass bottles with 50 mL of dichloromethane (CH<sub>2</sub>Cl<sub>2</sub>)  
741 at 4 °C in the dark before solvent extraction.

742 For the determination of the concentrations of suspended particulate matter (SPM),  
743 particulate organic carbon and nitrogen (POC and PON), as well as δ<sup>13</sup>C and δ<sup>15</sup>N isotopic  
744 ratios, between 250 and 1100 mL of sample were filtered with pre-weighted 25-mm diameter  
745 GF/F filters (the same filter was used for SPM, POC, PON, δ<sup>13</sup>C and δ<sup>15</sup>N analyses). After  
746 seawater filtration, filters were rinsed with ultrapure water to remove salts, and then were  
747 dried at 60 °C for 24 h and stored in aluminum foil until analysis.

#### 748 ***2.4. Analytical procedures for PAHs, PCBs and OCPs***

749 The analyzed compounds included 16 PAHs, 20 PCB congeners and 10 OCPs (Text S1,  
750 Table S2). The targeted PAHs were naphthalene (Nap), acenaphthylene (Acy), acenaphthene  
751 (Ace), fluorene (Flu), phenanthrene (Phe), anthracene (Ant), fluoranthene (Flt), pyrene (Pyr),  
752 benz[a]anthracene (BaA), chrysene (Chr), benzo[b]fluoranthene (BbF), benzo[k]fluoranthene  
753 (BkF), benzo[a]pyrene (BaP), dibenz[a,h]anthracene (DahA), benzo[g,h,i]perylene (BghiP),  
754 and indeno[1,2,3-cd]pyrene (IcdP). The 20 PCBs congeners were PCB 8, 18, 28, 52, 44, 66,  
755 77, 101, 105, 118, 126, 128, 138, 153, 170, 180, 195, 206, 187 and 209. The 10 OCPs were  
756 four hexachlorocyclohexanes (α-HCH, β-HCH, γ-HCH, δ-HCH), as well as two  
757 dichlorodiphenyltrichloroethanes (2,4'-DDT, 4,4'-DDT) and their metabolites,  
758 dichlorodiphenyldichloroethylenes (2,4'-DDE and 4,4'-DDE) and  
759 dichlorodiphenyldichloroethanes (2,4'-DDD and 4,4'-DDD). Details on analytical procedures  
760 for the determination of dissolved PAHs, PCBs and OCPs including chemicals,  
761 extraction/cleanup and instrumental analysis are presented in Text S1. Briefly, water samples  
762 (dissolved phase) were spiked with a multi standard mixture containing surrogate standards

763 for PAHs, PCBs and OCPs, and extracted by Liquid–Liquid extraction (LLE). The extracts  
764 were then purified using silica-alumina columns, internal standards were added before the  
765 analysis which was subsequently performed using gas chromatography coupled with a mass  
766 spectrometer (GC–MS) and an electron capture detector (GC-ECD) (Text S1, Table S2).

## 767 **2.5. Quality assurance/Quality control**

768 All data were guaranteed through the implementation of strict QA/QC procedures. Before  
769 use, all glassware was intensively cleaned and baked at 450 °C for 6 h. All the material that  
770 could not be baked was washed with 1 M HCl, ultrapure water, *n*-hexane:acetone mixture 1:1  
771 (v/v) and then dried at room temperature. Procedural blanks (to determine the contamination  
772 introduced during extraction and cleanup) were processed in the same way as the real samples  
773 and were included with each batch of 5 samples. Most of the analytes were not detected in  
774 blanks, except Nap, Acy, Ace, Flu, Phe, Ant, Flt, Pyr, BaA and Chr. Their amounts were not  
775 significant relative to that of the water samples (< 7%). The recoveries of target compounds  
776 were checked using standard mixtures spiked in ultrapure water (50 ng for PAHs, 20 ng for  
777 PCBs and 30 ng for OCPs). They ranged from 63 to 114% (mean of 87%), 7 to 117% (mean  
778 of 85%) and 20 to 115% (mean of 79%) for PAHs, PCBs and OCPs, respectively (Table S2).  
779 The surrogate standard (added to perform the quantification) recoveries ranged from 69 to  
780 92% and from 66 to 84% for PAHs and OCs, respectively (Table S3). Method detection limits  
781 (MDLs) defined as mean blank value + 3 × standard deviation (SD) ranged from 0.04 to 2.66  
782 ng L<sup>-1</sup>, 0.03 to 0.42 ng L<sup>-1</sup> and 0.10 to 0.40 ng L<sup>-1</sup> for PAHs, PCBs and OCPs, respectively  
783 (Table S2). For compounds that were not detected in blanks, instrument detection limits  
784 (IDLs), determined as a signal-to-noise ratio (S/N) of 3, were substituted. If the concentration  
785 of a given compound in a sample was below its MDL/IDL, thus this compound was  
786 considered as not detected in the sample, and was assumed zero. All data reported here were  
787 blank corrected.

## 788 2.6. Data handling and risk assessment

1  
2 789 The low molecular weight (LMW) PAHs (Nap, Acy, Ace, Flu, Phe, Ant) are petrogenic  
3  
4 790 (non-combustion processes) origin while the high molecular weight (HMW) PAHs (Flt, Pyr,  
5  
6  
7 791 BaA, Chr, BbF, BkF, BaP, DahA, BghiP, IcdP) are rather associated with incomplete  
8  
9 792 pyrogenic processes origin. In addition, two molecular PAH ratios, i.e., Flt/(Flt+Pyr) and  
10  
11 793 Ant/(Ant+Phe) were used to identify the possible PAH sources (petrogenic *versus* pyrogenic)  
12  
13 794 (Table S4). According to their degree of chlorination, the 20 PCBs can be classified into 9  
14  
15 795 groups: di-(PCB-8), tri- (PCB-18 and -28), tetra- (PCB-52, -44, -66 and -77), penta- (PCB-  
16  
17 796 101, -105, -118 and -126), hexa- (PCB-128, -138 and -153), hepta- (PCB-170 and -180), octa-  
18  
19 797 (PCB-195), nona- (PCB-206) and deca-PCBs (PCB-187 and -209). Among the 20 PCB  
20  
21 798 congeners targeted, seven PCBs (PCB-28, -52, -101, -118, -138, -153, and -180), are named  
22  
23 799 indicators PCBs ( $\Sigma 7$  IN-PCBs). Among the 10 targeted OCPs,  $\Sigma 4$  HCHs represent the sum of  
24  
25 800  $\alpha$ -,  $\beta$ -,  $\gamma$ - and  $\delta$ -HCH and  $\Sigma 6$  DDTs represent the sum of 2,4'-DDT, 4,4'-DDT, 2,4'-DDE, 4,4'  
26  
27 801 '-DDE, 2,4'-DDD and 4,4'-DDD. Two ratios, (DDE+DDD)/ $\Sigma$ DDTs and DDD/DDE, were  
28  
29 802 used to determine the recent/historical character of DDT contamination and to determine the  
30  
31 803 type of biodegradation mechanism (aerobic or anaerobic) to which DDT is subjected in the  
32  
33 804 study area (Table S4), respectively.

34  
35 805 To estimate the environmental risk posed by PAHs in the study aquatic system, toxicity  
36  
37 806 equivalents (TEQs) was calculated using Toxicity Equivalent Factors (TEFs) (Nisbet and  
38  
39 807 LaGoy, 1992), as follows:

$$40 \quad \text{TEQs} = \text{TEF} \times \text{PAH concentration}$$

41  
42 808 The risk quotients of negligible concentrations  $\text{RQ}_{(\text{NCs})}$  and maximum permissible  
43  
44 809 concentrations  $\text{RQ}_{(\text{MPCs})}$  (Cao et al., 2010) were calculated as follows:

$$45 \quad \text{RQ}_{(\text{NCs})} = C_{\text{PAHs}}/C_{(\text{NCs})}$$

$$46 \quad \text{RQ}_{(\text{MPCs})} = C_{\text{PAHs}}/C_{(\text{MPCs})}$$

813 Finally, the risk of PAHs on benthic species endemic in the water column ( $\sum RQs$ ) was  
 814 calculated by the sum of RQs for individual PAHs, as follows:

$$RQ_{\Sigma PAHs(NCs)} = \sum_{i=1}^{16} RQ_i \left( RQ_{(NCs)} \geq 1 \right)$$

$$RQ_{\Sigma PAHs(MPCs)} = \sum_{i=1}^{16} RQ_i \left( RQ_{(MPCs)} \geq 1 \right)$$

817 The interpretation of the risk classification of individual PAHs and  $\Sigma PAHs$  are as follows  
 818 (Cao et al., 2010). In principle,  $RQ_{(MPCs)} > 1$  indicates that contamination by the single PAH is  
 819 sever, whereas an  $RQ_{(NCs)} < 1$  indicates that a single PAHs is probably of negligible concern.  
 820 In the case of  $RQ_{(NCs)} > 1$  and  $RQ_{(MPCs)} < 1$ , contamination of the single PAH might be  
 821 considered moderate. For  $RQ_{\Sigma PAHs}$ ,  $RQ_{\Sigma PAHs(MPCs)} < 1$  indicates PAH contamination is low-  
 822 level, while  $RQ_{\Sigma PAHs(NCs)} > 800$  would indicate severe PAH contamination. If  $RQ_{\Sigma PAHs(NCs)} \geq 1$   
 823 or  $< 800$  and  $\sum RQ_{(MPCs)} > 1$ , it likely indicates moderate PAH contamination.

824 Regarding OCP and PCBs, the risk assessment was estimated similarly using the following  
 825 equations:

$$RQ = C_{TCS} / C_{QV}$$

$$RQ_{CCCs} = C_{TCS} / C_{QV (CCCs)}$$

$$RQ_{CMCs} = C_{TCS} / C_{QV (CMCs)}$$

829 where,  $C_{TCS}$  is the concentration of targeted contaminants,  $C_{QV}$ , is the corresponding quality  
 830 values of certain OCPs and PCBs,  $RQ_{CCCs}$  is the risk quotient for continuous criteria  
 831 concentrations (CCCs), whereas  $RQ_{CMCs}$  is the risk quotient for maximum criteria  
 832 concentrations (CMCs) (USEPA, 2006).

## 833 2.7. Other analyses

834 After 24 h in the desiccator, 25-mm GF/F filters were reweighed by the same balance that  
 835 was used for pre-weighting. The SPM concentration was then calculated as the difference

836 between filter weight before and after sample filtration, normalized to the filtration volume  
1  
2 837 (Neukermans et al., 2012). POC and PON concentrations, and  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  stable isotopic  
3  
4 838 ratios were determined by high combustion method with mass spectrometer (Raimbault et al.,  
5  
6  
7 839 2008). Measurements of fluorescent dissolved organic matter (FDOM) at 254-nm excitation  
8  
9 840 wavelength and 360-nm emission wavelength were performed on unfiltered (raw) seawater  
10  
11 841 samples (~ 500 mL) with a UV EnviroFlu-HC fluorometer (TriOS Optical sensors) (Tedetti et  
12  
13 842 al., 2010). Here, fluorescence data are provided in relative unit (RU).

## 17 843 **2.8. Statistics**

20 844 Statistical analyses, including Pearson correlations and a principal component analysis  
21  
22 845 (PCA) based on Spearman's rank order correlation matrix, were performed using XLSTAT  
23  
24 846 2013.5.01. Colour maps of physico-chemical parameters and pollutant concentrations were  
25  
26 847 carried out using Ocean Data View (ODV) software version 4.6.5 (<http://odv.awi.de>). The  
27  
28 848 spatial interpolation/gridding of data was conducted using Data-Interpolating Variational  
29  
30 849 Analysis (DIVA) (Barth et al., 2010; Troupin et al., 2012).

35 850

## 37 851 **3. Results**

### 41 852 **3.1. Spatial distribution of physico-chemical and biogeochemical parameters**

43 853 The spatial distribution of temperature, salinity, pH, FDOM,  $\text{O}_2$  concentration, and redox  
44  
45 854 potential in surface waters in the study area (at ~ 0.5-m depth) is shown Fig. 2. Temperature  
46  
47 855 did not display any strong variability (from 21.48 °C at S4 to 23.43 °C at S22), nor any clear  
48  
49 856 trend relative to the phosphogypsum effluent outlet (S23, Fig. 2a). However, salinity, pH,  
50  
51 857 FDOM and  $\text{O}_2$  concentration exhibited a distribution pattern related to the effluent. Indeed,  
52  
53 858 the most coastal stations, particularly stations S23, S24 and S25 (the three stations located  
54  
55 859 between the two harbours and directly under the influence of the effluent) presented the

1 860 highest salinity values (40.11-40.95), the lowest pH values (7.49-7.82), the highest FDOM  
2 861 values (1089-1410 RU), and the highest O<sub>2</sub> concentrations (9.6-10.6 mg L<sup>-1</sup>) (Figs. 2b-e).  
3  
4 862 Stations S22 and S26, situated North and South of these 3 stations, respectively, also showed  
5  
6  
7 863 higher values of FDOM (634-824 RU), O<sub>2</sub> concentration (8.6-9.8 mg L<sup>-1</sup>), and lower values of  
8  
9 864 pH (8.17-8.29) relative to the other stations. In contrast, the lowest values of salinity (<  
10  
11 865 39.40), FDOM (< 600 RU), O<sub>2</sub> concentration (< 8.0 mg L<sup>-1</sup>), and the highest values of pH (>  
12  
13 866 8.30) were encountered mostly in the Northern and offshore sites (S1-S11) (Figs. 2b-e).  
14  
15  
16 867 Redox potential displayed the lowest values (< 200 mV) along the transect in front of  
17  
18 868 Ghannouch commercial harbour/phosphogypsum discharges (S7, S8, S10, S11), whereas the  
19  
20  
21 869 highest values (> 260 mV) were found in the Southern part (S13-S18) (Fig. 2f).  
22  
23

24 870 The coastal stations S23-S25 were also characterized by the highest POC and PON  
25  
26 871 concentrations (73.2-89.6 μM and 8.9-10.9 μM, respectively), the highest SPM values (15.0-  
27  
28 872 16.8 mg L<sup>-1</sup>), and the lowest δ<sup>15</sup>N values (-1.2-1.9‰) (Table S5). On the other side, the  
29  
30  
31 873 lowest concentrations of POC, PON and SPM were recorded in the Northern and offshore  
32  
33 874 sites (S3, S4, S8, S11) (Table S5). Also, salinity, FDOM, O<sub>2</sub> concentration, POC, PON and  
34  
35 875 SPM concentrations were positively (linearly) correlated to each other (r = 0.65-0.97, n = 15,  
36  
37 876 p < 0.0001-0.009), while these latter were all negatively correlated to pH (r = -0.70-0.93, n =  
38  
39 877 15, p < 0.0001-0.004). Turbidity and TChl<sub>a</sub> concentration, albeit not measured in stations  
40  
41 878 S19-S27, showed higher values in the Southern stations (S13-S18) than in the Northern ones  
42  
43  
44 879 (Table S5). Overall, these results highlight coast-offshore and South-North gradients of the  
45  
46  
47 880 different physico-chemical and biogeochemical parameters, with a substantial influence of the  
48  
49  
50 881 phosphogypsum effluent outlet on the local biogeochemistry.  
51  
52

### 53 882 ***3.2. Spatial distribution of total dissolved PAHs, PCBs and OCPs***

54 883 The spatial distribution of concentrations of Σ16 PAHs (sum of 16 compounds), Σ20 PCBs  
55  
56 884 (sum of 20 congeners) and Σ10 OCPs (sum of 10 compounds) in surface waters (at ~ 0.5-m  
57  
58  
59  
60  
61  
62  
63  
64  
65

885 depth) is provided [Fig. 3](#). Detailed information about individual compounds is given in [Table](#)  
1  
2 886 [S6](#). Concentration of  $\Sigma 16$  PAHs was the highest in the vicinity of Ghannouch industrial  
3  
4 887 complex: at stations S8 and S9 (62.7-71.2 ng L<sup>-1</sup>), in front of Ghannouch  
5  
6 888 harbour/phosphogypsum discharges, at station S22 (64.9 ng L<sup>-1</sup>), in front of Ettine  
7  
8 889 wadi/fluorine discharges, at stations S4, S19, S20 (40.5-59.8 ng L<sup>-1</sup>), Northern Ghannouch,  
9  
10 890 and at station S24 (46.1 ng L<sup>-1</sup>), between Ghannouch and Gabès harbours ([Fig. 3a](#), [Table 1](#)).  
11  
12 891 Concentration of  $\Sigma 20$  PCBs showed a quite different distribution with the highest values  
13  
14 892 found off Ghannouch-Gabès area (at S9: 33.7 ng L<sup>-1</sup> and S11: 31.4 ng L<sup>-1</sup>), Northern  
15  
16 893 Ghannouch (at S1: 22.1 ng L<sup>-1</sup>), and in front of fishing harbour and Gabès wadi (at S13: 28.3  
17  
18 894 ng L<sup>-1</sup>) ([Fig. 3b](#), [Table 1](#)).

24 895 For OCPs, concentration of  $\Sigma 4$  HCHs displayed a distribution pattern relatively similar to  
25  
26 896 that of PAHs, with the highest values observed around the Ghannouch industrial complex and  
27  
28 897 Northern Ghannouch: at stations S8 (14.8 ng L<sup>-1</sup>), S22 (9.4 ng L<sup>-1</sup>), S19 and S20 (6.4-7.0 ng  
29  
30 898 L<sup>-1</sup>) ([Fig. 3c](#), [Table 1](#)). Interestingly,  $\Sigma 16$  PAH and  $\Sigma 4$  HCH concentrations were positively  
31  
32 899 correlated ( $r = 0.72$ ,  $n = 27$ ,  $p < 0.001$ ). The highest concentration of  $\Sigma 6$  DDTs observed was  
33  
34 900 in Northern Ghannouch (at S4-S6 and S19: 11.3-13.2 ng L<sup>-1</sup>), and in front of phosphogypsum  
35  
36 901 discharges (at S8: 12.0 ng L<sup>-1</sup>), as well as Southern Gabès, in front of Griaa wadi (at S26: 10.9  
37  
38 902 ng L<sup>-1</sup>) and in front of Mersit wadi (at S16: 11.7 ng L<sup>-1</sup>) ([Fig. 3d](#), [Table 1](#)). It appeared that the  
39  
40 903 spatial distributions of PAHs, PCBs, HCHs and DDTs were relatively different from those of  
41  
42 904 the physical and physico-chemical parameters described above (i.e., salinity, pH, FDOM, O<sub>2</sub>  
43  
44 905 concentration and redox potential) because the contaminant concentrations exhibited less  
45  
46 906 pronounced coast-offshore gradients, and were less driven by the phosphogypsum effluent  
47  
48 907 outlet. PAHs and OCPs (particularly HCHs) nonetheless seemed to show more similarities  
49  
50 908 with physico-chemical parameters than PCBs, especially regarding their higher concentrations  
51  
52 909 in the Ghannouch/effluent zone.

### 3.3. Molecular distribution of dissolved PAHs, PCBs and OCPs

Among the 16 PAHs targeted in this study, four were not detected in the samples: BaP, DahA, IcdP, and BghiP (Fig. 4a). The most abundant PAHs were Nap (on average 63% of total PAHs, ranging from 27% at S7 to 82% at S19) followed by Phe (on average 12%, ranging from 7% at S8, S19, S22 to 25% at S4), and Flt (on average 7%, ranging from 1% at S8, S22 to 36% at S7), while other compounds only accounted for 0-5% on average of total PAHs (Fig. 4a). With respect to size, the concentrations of LMW-PAHs in surface waters of the Gulf of Gabès ranged from 15.8 to 62.4 ng L<sup>-1</sup> and for HMW-PAHs, the range was 1.4 to 23.3 ng L<sup>-1</sup>. The Flt/(Flt+Pyr) ratio was 0.56 on average and ranging in values from 0.40 (S10) to 0.90 (S7), while the Ant/(Ant+Phe) ratio was 0.07 on average with values ranging from 0.01 (S1) to 0.21 (S8) (Table 1).

Of the 20 targeted PCB congeners, 11 were not detected in the samples. These were PCB-8, -44, -66, -77, -105, -126, -170, -187, -195, -206, -209 (Fig. 4b). The relative abundance of the 9 congeners detected was highly variable from station to station. PCB-180 alone accounted for 27% of the PCBs on average, with minimal contribution of 1% (at S10) and maximal contribution of 71% (at S26). PCB-18 and -52 were also detected in large amounts. The two congeners represented 18 and 19% of PCBs on average, respectively, with minimal values of 0% (at S3 and S26 for PCB-18, and at S12, S14 and S15 for PCB-52) and maximal values of 41% (at S20 for PCB-52) or 59% (at S14 for PCB-18). PCB-101, -28 and -153 displayed lower mean relative abundances representing 14, 10 and 6% (respectively) while the sum of the remaining congeners (PCB-118, -128, -138) only depicted 6% of total PCBs (Fig. 4b).

Among OCPs of interest, 2,4'-DDT, 4,4'-DDT and 4,4'-DDD were not at all detected. 2,4'-DDD and 2,4'-DDE compounds were major species, with average contributions to total OCPs of 31 and 28%, respectively, minimal contributions of 0% (at S12, S15) and 4% (at S7, S18),



1  
2 936 respectively, and maximal contributions of 64% (at S7) and 51% (at S11), respectively (Fig.  
3  
4  
5 937 4c). They were followed by  $\gamma$ -HCH, which represented on average 19% with minimal and  
6  
7 938 maximal values of 4% (at S11, S14) and 51% (at S12), respectively. The sum of the  
8  
9 939 remaining compounds ( $\alpha$ -HCH,  $\beta$ -HCH,  $\delta$ -HCH, 4,4'-DDE) accounted on average for 22% of  
10  
11 940 total OCPs (Fig. 4c). The DDD/DDE ratio was quite variable, ranging from 0.21 (S11) to 3.74  
12  
13 941 (S7), with mean and median values of 1.30 and 0.80, respectively (Table 1). Since 2,4'-DDT  
14  
15 942 and 4,4'-DDT were not detected, the (DDE+DDD)/ $\Sigma$ DDTs ratios were all equal to 1 (data not  
16  
17 943 reported in Table 1).

### 18 19 20 943 **3.4. PCA classification of stations**

21  
22  
23 944 Spearman PCA was applied on various parameters including salinity, pH, O<sub>2</sub>  
24  
25 945 concentration, FDOM, as well as concentrations of  $\Sigma$ 16 PAHs,  $\Sigma$ 20 PCBs and  $\Sigma$ 10 OCPs (Fig.  
26  
27 946 5). The first principal component (PC1), which explained 39% of total variability within  
28  
29 947 samples, mainly received the contribution of variables salinity, pH, FDOM and in a lesser  
30  
31 948 extent O<sub>2</sub> concentration and  $\Sigma$ 10 OCPs. The second principal component (PC2) accounted for  
32  
33 949 21% of total variability within samples and was mostly related to variables  $\Sigma$ 16 PAHs,  $\Sigma$ 20  
34  
35 950 PCBs, followed by O<sub>2</sub> concentration and  $\Sigma$ 10 OCPs. PCA allowed the discrimination of  
36  
37 951 different groups of stations. Group n°1 (S2, S3, S5, S7, S10, S12, S14, S17, S18, S21 and  
38  
39 952 S27) is rather characterized by high values of pH, low values of salinity, O<sub>2</sub> concentration and  
40  
41 953 FDOM, and low concentrations of total dissolved PAHs, PCBs and OCPs. Group n°2 (S1, S6  
42  
43 954 and S11) is depicted by quite high values of pH, low values of salinity, O<sub>2</sub> concentration and  
44  
45 955 FDOM, and high concentrations of total dissolved PAHs, PCBs and OCPs. Group n°3 (S4,  
46  
47 956 S8, S9, S13, S16, S19, S20, S22) is defined by lower values of pH, higher values of salinity,  
48  
49 957 O<sub>2</sub> concentration and FDOM, and high concentrations of total dissolved PAHs, PCBs and  
50  
51 958 OCPs. Finally, group n°4 is characterized by roughly low values of pH, high values of  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2 960 salinity, O<sub>2</sub> concentration and FDOM, and low concentrations of total dissolved PAHs, PCBs  
3 and OCPs (Fig. 5).

4  
5 961

## 6 7 962 **4. Discussion**

### 8 9 10 963 **4.1. Concentration levels**

11  
12  
13 964 The concentrations of total dissolved ( $\Sigma 16$ ) PAHs (17.6-71.2 ng L<sup>-1</sup>) recorded here in the  
14  
15 965 surface waters of the Ghannouch-Gabès area (Fig. 3a, Table 1) were of the same order of  
16  
17  
18 966 magnitude but were lower than those recently measured in the neighboring area by Fourati et  
19  
20 967 al. (2018) (i.e., 9-198 ng L<sup>-1</sup>; Table S7). This difference may be due to the fact that this latter  
21  
22 968 study targeted more compounds ( $\Sigma 29$  PAHs) including alkylated compounds, which are  
23  
24  
25 969 recognized to account for a large part of the pool of dissolved PAHs in natural waters (Guigue  
26  
27 970 et al., 2014; Wang et al., 1999). Also, dissolved PAH concentrations were in the lower range  
28  
29  
30 971 of those reported in coastal Bohai Bay, China (48-607 ng L<sup>-1</sup> for  $\Sigma 16$  PAHs) (Tong et al.,  
31  
32 972 2019), various coastal areas of the Southern, Northern and Northwestern Mediterranean Sea,  
33  
34  
35 973 such as the Gulf of Tunis, Tunisia (139-1008 ng L<sup>-1</sup> for  $\Sigma 24$  PAHs) (Mzoughi and Chouba,  
36  
37 974 2011), the Marseilles coastal area, France (1.9-151 ng L<sup>-1</sup> for  $\Sigma 17$  PAHs) (Guigue et al., 2011,  
38  
39 975 2014) and the Alexandria coast, Egypt (13-120 ng L<sup>-1</sup> for  $\Sigma 7$  PAHs) (El Nemr and Abd-Allah,  
40  
41 976 2003), but were higher than those observed in the Open Western and Eastern Mediterranean  
42  
43  
44 977 Sea (0.16-9 ng L<sup>-1</sup> for  $\Sigma 19$  PAHs) (Berrojalbiz et al., 2011), the Northwest Atlantic coast of  
45  
46  
47 978 the Iberian Peninsula (9.3 ± 0.6 ng L<sup>-1</sup> for  $\Sigma 16$  PAHs) (Rocha and Rocha, 2021), the European  
48  
49 979 Atlantic coastline, Portugal (8.3-9.5 ng L<sup>-1</sup> for  $\Sigma 16$  PAHs) (Rocha et al., 2021) and Barcelona,  
50  
51  
52 980 Spain and Banyuls-sur-Mer, France (0.4-8.6 ng L<sup>-1</sup> for  $\Sigma 15$  PAHs) (Garcia-Flor et al., 2005)  
53  
54 981 (Table S7).

55  
56  
57 982 The concentrations of total dissolved ( $\Sigma 20$ ) PCBs (2.9-33.6 ng L<sup>-1</sup>) we determined here in  
58  
59 983 the Gulf of Gabès (Fig. 3b, Table 1) were in most cases higher than those recorded in other  
60  
61

1 984 environments, such as the Open Baltic Sea ( $0.02 \pm 0.01 \text{ ng L}^{-1}$  for  $\Sigma 7$  PCBs) (Cornelissen et  
2 985 al., 2008) and the Tokyo Bay, Japan ( $0.04\text{-}0.6 \text{ ng L}^{-1}$  for  $\Sigma 209$  PCBs) (Kobayashi et al., 2010).  
3  
4 986 However, they were of the same order of magnitude than those recorded in the Singapore's  
5  
6  
7 987 northeastern and southwestern coastlines ( $0.22\text{-}20.41 \text{ ng L}^{-1}$  for  $\Sigma 8$  PCBs) (Basheer et al.,  
8  
9 988 2003), and they were lower than those found in the Daya Bay, China ( $91.1\text{-}1355.3 \text{ ng L}^{-1}$  for  
10  
11  
12 989  $\Sigma 12$  PCBs) (Zhou et al., 2001), and the Panama Canal and California Coast, USA (*bdl*- $18,450$   
13  
14 990  $\text{ ng L}^{-1}$  for  $\Sigma 19$  PCBs) (Menzies et al., 2013) (Table S7). Urbanization, human activities and  
15  
16 991 particularly, intense industrialization in this area may explain the higher PCB concentrations  
17  
18  
19 992 compared to the other environments.

20  
21  
22 993 The concentrations of total dissolved ( $\Sigma 10$ ) OCPs ( $5.3\text{-}26.7 \text{ ng L}^{-1}$ ) found here (Fig. 3c and  
23  
24 994 d, Table 1) were in the same range than those measured in Hangzhou Bay, China ( $1.3\text{-}26 \text{ ng}$   
25  
26 995  $\text{ L}^{-1}$  for  $\Sigma 10$  OCPs) (Li et al., 2016) and Singapore's northeastern and southwestern coastlines  
27  
28 996 ( $4.9\text{-}22 \text{ ng L}^{-1}$  for  $\Sigma 12$  OCPs) (Basheer et al., 2003), but lower than that recorded in Jinzhou  
29  
30  
31 997 Bay, China ( $4.2\text{-}136.8 \text{ ng L}^{-1}$  for  $\Sigma 19$  OCPs) (Yao et al., 2013) (Table S7). Large differences  
32  
33  
34 998 in pesticide concentrations in global surface waters may be explained mainly by differences in  
35  
36 999 agricultural activities, given the strong connection between OCPs and agriculture (Montory et  
37  
38  
39 1000 al., 2017). Although all these comparisons should be taken with caution due to the different  
40  
41 1001 number of compounds analyzed, the tendency that emerges is that the concentrations we  
42  
43  
44 1002 measured in this work in the Ghannouch-Gabès area (Southern Gulf of Gabès) were in the  
45  
46 1003 mid-range of those reported in other coastal environments (Table S7).

#### 49 1004 **4.2. Sources/origins**

50  
51  
52 1005 Compositional profiles for PAHs in water samples are illustrated in Fig. 4a. LMW-PAHs  
53  
54 1006 were abundant at all sampling sites, representing on average  $> 80\%$  of  $\Sigma 16$  PAHs; HMW-  
55  
56  
57 1007 PAHs, on the other hand, were present in low concentrations and contributions to  $\Sigma 16$  PAHs.  
58  
59  
60  
61  
62  
63  
64  
65

1008 This is typical for surface waters because of the relatively high solubility of LMW-PAHs (log  
1  
21009  $K_{ow}$  is 3-5 for LMW and 5-7 for HMW) (Moeckel et al., 2014; Shen et al., 2021).  
3  
4  
51010 The evaluation of PAH sources by sampling site was achieved by considering published  
6  
71011 indicators (Table 1) and the analysis of bivariate cross plots, which reflect characteristic  
8  
9  
101012 PAHs ratios (Fig. S1). Because PAHs naturally partitioned between dissolved and particulate  
11  
121013 water phases in line with their molecular weight/hydrophobicity, the dominance of the LMW  
13  
141014 compounds in the present samples (especially Nap and Phe), which are much more soluble  
15  
16  
171015 than the HMW ones, cannot be used as an origin/source criterion. For that, isomeric  
18  
191016 compound ratios, Ant/(Ant+Phe) and Flt/(Flt+Pyr) are more reliable, although they might be  
20  
21  
221017 affected by weathering processes between the emission points and the sampling area  
23  
241018 (Katsoyiannis and Breivik, 2014). According to Table 1 and Table S4, PAHs were found to be  
25  
26  
271019 both pyrolytic and petrogenic origin, with a predominance of petrogenic sources (63% of the  
28  
291020 stations had a ratio of Flt/(Flt+Pyr) between 0.4 and 0.5, and 89% of the stations had a ratio of  
30  
31  
321021 Ant/(Ant+Phe) < 0.1) (Ravindra et al., 2006; Soclo et al., 2000; Yunker et al., 2002). The plot  
33  
341022 of isomeric ratios, Ant/(Ant+Phe) versus Flt/(Flt+Pyr) (Fig. S1), confirm the presence of the  
35  
361023 last referred sources in surface waters. This reflects the multitude of sources of PAHs in this  
37  
38  
391024 anthropized coastal area of the Gulf of Gabès through the utilization of petroleum products  
40  
411025 and incomplete combustion of fossil fuels and biomass (Stogiannidis and Laane, 2015).  
42  
43  
441026 Petrogenic PAHs could be introduced *via* accidental oil spills, discharges from routine ship  
45  
461027 and harbour activities/operations, and urban runoff, while pyrogenic PAHs may enter *via*  
47  
48  
491028 exhausts and residues issued from the combustion related to the various urban, industrial and  
50  
511029 agriculture activities occurring in this region. Moreover, PAHs, particularly LMW-PAHs  
52  
53  
541030 prevailing in this study, can reach surface waters of the Gulf of Gabès *via* atmospheric  
55  
561031 transport over long distances from the Gabès city and other areas. Interestingly, the stations  
57  
581032 located either in front of (S7-S9, S23) or close to (S22, S24) Ghannouch industrial  
59  
60  
61  
62  
63  
64  
65

1033 complex/commercial harbour and phosphogypsum effluent outlet (Fig. 1, Table S1) presented  
1  
2 1034 dissolved PAHs mainly of pyrogenic (combustion) origin, as seen from their Ant/(Ant+Phe)  
3  
4  
5 1035 (0.12-0.21) and Flu/(Flu+Pyr) ratios (0.43 and 0.90) (Table 1, Table S4). Moreover, these  
6  
7 1036 stations (S7-S9, S22-S24) were marked by higher concentrations of total dissolved PAHs  
8  
9  
10 1037 (19.5-71.2 ng L<sup>-1</sup>; mean and median: 49 and 56 ng L<sup>-1</sup>) compared to the other stations (17.6-  
11  
12 1038 59.8 ng L<sup>-1</sup>; mean and median: 31 and 28 ng L<sup>-1</sup>). This could be the sign of a specific PAH  
13  
14 1039 fingerprint due to the whole Ghannouch industrial complex activity.

16  
17 1040 The dissolved phase was mostly enriched by PCBs displaying a low degree of chlorination,  
18  
19 1041 i.e., tri- (CB-18, CB-28; 28% of total PCBs on average), tetra- (CB-52; 19% of total PCBs on  
20  
21  
22 1042 average), penta- (CB-101; 19% of total PCBs on average), and hexa-CBs (CB-153; 19% of  
23  
24 1043 total PCBs on average) (Table 1, Fig. 4b), which is in agreement with previous observations  
25  
26  
27 1044 and may be explained by their higher solubility in water compared to PCBs of higher degree  
28  
29 1045 of chlorination (Li et al., 2016; Montuori et al., 2020; Moret et al., 2005). Less chlorinated  
30  
31  
32 1046 PCBs are more amenable to long range atmospheric transport than heavier ones (Wania and  
33  
34 1047 MacKay, 1995). This suggests that atmospheric transport could be the most important PCB  
35  
36 1048 source in the studied region. Furthermore, Tri-CBs, which are mainly used as an insulating  
37  
38  
39 1049 liquid for transformer oils and power capacitors (Tang et al., 2020), can leaked into the soil  
40  
41 1050 and end up in surface waters *via* surface runoff and land use (Duan et al., 2013). Surprisingly,  
42  
43  
44 1051 CB-180 which is a heptachloro-congener, characterized by a higher number of chlorine atoms  
45  
46 1052 (and thus a lower solubility in water), was one of the most abundant PCBs in our samples  
47  
48  
49 1053 (27% of total PCBs on average). Generally, CB-180 is not detected or detected at very low  
50  
51 1054 levels in the dissolved phase of natural waters (Montuori et al., 2020; Moret et al., 2005;  
52  
53 1055 Rabodonirina et al., 2015). Indicator PCBs were detected in the surface waters of the Gulf of  
54  
55  
56 1056 Gabès with detection frequency varied from 19 to 100%, and concentrations ranged from 1.8  
57  
58 1057 to 28.4 ng L<sup>-1</sup>. The levels of Σ7 IN-PCBs were strongly correlated with total PCBs (Σ20  
59  
60  
61  
62  
63  
64  
65

1058 PCBs) (Pearson  $r = 0.97$ ,  $p < 0.0001$ ) and a conversion factor of 4 can be used to estimate  
1  
21059 total PCB concentrations ( $\Sigma 209$  PCBs) from 7 IN-PCBs (i.e.,  $\Sigma 209$  PCBs =  $4 \times 7$  IN-PCBs).  
3  
4  
51060 Similar conversion factor was applied by previous studies for sediment and road dust (Anh et  
6  
71061 al., 2019; Hoai et al., 2010). PCBs are used in industry as a heat exchanger liquid in  
8  
9  
101062 transformers and capacitors, and as additives in paints, carbonless copy papers and plastics.  
11  
121063 As in numerous other countries, equipment containing PCBs have been widely used in  
13  
141064 Tunisia from the 1970s to late 1980s (an import ban of those equipment being occurred in  
15  
16  
171065 1986). Generally, sources of PCBs in marine environment are wastes of electrical  
18  
191066 transformers, oil spillage, and any historical use of PCB (Mitra et al., 2019). Recently, PCBs  
20  
21  
221067 have been recorded in Tunisian (Bizerte) coastal areas: in surface sediments, mussels, fishes  
23  
241068 and atmospheric particles (Barhoumi et al., 2014a, b, c; Barhoumi et al., 2018; Ben Ameur et  
25  
26  
271069 al., 2013). In these compartments (sediments, fishes, atmosphere), the presence of tetrachloro-  
28  
291070 , pentachloro-, hexachloro- and even heptachloro-congeners was consistent with a  
30  
31  
321071 contribution of commercial (Aroclor) PCB mixtures (Takasuga et al., 2006). In the same way,  
33  
341072 our PCB molecular distribution may reflect the influence of commercial PCB mixtures.

361073 Various approaches based on the ratio of parent DDTs and their metabolites have been  
37  
38  
391074 used to assess whether the DDT contamination in a given area can be considered recent, and  
40  
411075 whether the degradation of DDTs occurred rather under aerobic or anaerobic condition (Hitch  
42  
43  
441076 and Day, 1992; Hong et al., 2003; Lee et al., 2001; Liu et al., 2008) (Table S4). Here, since  
45  
461077 parent DDTs were not detected at all in our samples, the ratios (DDE+DDD)/ $\Sigma$ DDTs were  
47  
48  
491078 always equal to 1. This point outs the historical/ancient character of the DDT contamination  
50  
511079 in the Gulf of Gabès. Moreover, more than half of stations displayed DDD/DDE ratio  $< 1$   
52  
53  
541080 (Table 1), which suggests that DDT was rather biotransformed into DDE under aerobic  
55  
561081 conditions (Galanopoulou et al., 2005; Mostafa et al., 2007). This is in accordance with the  
57  
581082 oxidizing conditions prevailing in surface waters of the Gulf of Gabès, as indicated by  $O_2$

1083 concentrations and redox values (Table S5). Therefore, our results indicate that DDTs origin  
1  
21084 from historical load since only their metabolites have been detected. Prior to their ban in  
3  
41085 Tunisia early 1984s (APEK, 2005), DDTs were largely used as pesticides in agriculture in the  
5  
6  
71086 environment of the Gulf of Gabès, thus the current flux may come from places where they  
8  
9  
101087 have remained *via* runoff and wadis or the sediment resuspension. The highest DDT  
11  
121088 concentrations measured at stations S4 (12.1 ng L<sup>-1</sup>), S5 (11.35 ng L<sup>-1</sup>), S6 (13.23 ng L<sup>-1</sup>) and  
13  
141089 S19 (11.5 ng L<sup>-1</sup>) located in front of El Demna and El Melah wadis (Northern Ghannouch),  
15  
16  
171090 and station S16 (11.66 ng L<sup>-1</sup>) located in front of Mersit wadi (Southern Gabès), may confirms  
18  
191091 the last statement. In addition, antifouling paints used in boat maintenance could be another  
20  
21  
221092 source of DDTs in surface water of the Gulf of Gabès, evidenced by the high DDT  
23  
241093 concentration measured at station S8 (11.97 ng L<sup>-1</sup>) located in front of Ghannouch  
25  
26  
271094 harbour/phosphogypsum discharges.

291095 HCHs usually have two formulations: technical-grade (predominantly constituted by  $\alpha$ -  
30  
311096 HCH) and lindane (> 99%  $\gamma$ -HCH) (Iwata et al., 1993). The isomer ratio  $\alpha$ -/ $\gamma$ -HCH can be  
32  
33  
341097 used as an indicator of the HCH sources, i.e., technical HCH has  $\alpha$ -HCH/ $\gamma$ -HCH ratios  
35  
361098 ranging between 3 and 7, whereas those of lindane are < 1.0 or close to 1.0 (Willett et al.,  
37  
38  
391099 1998).  $\alpha$ -HCH was found in only three sites, in the Gulf of Gabès, whereas  $\gamma$ -HCH was  
40  
411100 detected at all sites (Table S6), thus yielding ratios lower than technical HCHs. This reflects  
42  
43  
441101 the fact that HCHs were predominantly of lindane origin. Lindane was usually used in  
45  
461102 agriculture for seed treatment for beans, barley and wheat, as well as a lesser degree as a  
47  
48  
491103 pharmaceutical to treat lice and scabies (Barhoumi et al., 2016). Given their high vapour  
50  
511104 pressure (Shunthirasingham et al., 2010),  $\gamma$ -HCH was probably entered the Gulf *via*  
52  
53  
541105 atmospheric transport. Additionally, high detection frequency (85%) of  $\beta$ -HCH was observed,  
55  
561106 suggested that HCHs were applied in these areas in the past and had degraded, because  $\beta$ -  
57  
581107 HCH is more stable than other HCH isomers and accumulates in the environment over time  
59  
60  
61  
62  
63  
64  
65

1108 (Malik et al., 2009). Direct discharges, surface runoff from surrounding agricultural soils,  
1  
2 1109 along with long-range transport may account for the HCHs pollution in this area.  
3

4  
5 1110 As shown in Fig. 4c, DDTs and HCHs displayed different patterns. This could indicate  
6  
7 1111 different types of human activities (agriculture) prevailed in the area, as well as different input  
8  
9  
10 1112 sources, e.g., wet and dry deposition sources, surface runoff, and atmospheric transport. DDT  
11  
12 1113 and HCH patterns could be also influenced by the physico-chemical properties of isomer  
13  
14 1114 compounds along with environmental conditions. In addition, redox potential, topography,  
15  
16  
17 1115 and hydrodynamic condition together with other factors play important roles in controlling the  
18  
19 1116 distribution and fate of DDTs and HCHs (Su et al., 2006).  
20  
21

#### 22 1117 ***4.3. Potential influence of the industrial effluent and ecological risk assessment*** 23

24  
25 1118 Our results show a significant influence of the phosphogypsum effluent on the local  
26  
27  
28 1119 physico-chemistry and biogeochemistry, with the highest values of salinity, FDOM, O<sub>2</sub>, SPM,  
29  
30 1120 POC, PON, and the lowest values of pH observed in the most coastal stations located between  
31  
32 1121 the Ghannouch and Gabès harbours (S23-S25; Fig. 1), and the opposite pattern (i.e., the  
33  
34  
35 1122 lowest values of salinity, FDOM, O<sub>2</sub>, SPM, POC, PON, and the highest values of pH) found  
36  
37 1123 in the Northern and offshore sites, leading to marked coast-offshore and South-North  
38  
39  
40 1124 gradients (Fig. 2, Table S5). Hence, the influence of the phosphogypsum effluent mostly  
41  
42 1125 appeared South of the Ghannouch industrial complex, in the inter-harbour (Ghannouch-  
43  
44  
45 1126 Gabès) zone. These observations are in accordance with recent works, which pointed out a  
46  
47 1127 substantial effect of the Ghannouch phosphogypsum effluent on seawater or surface sediment  
48  
49  
50 1128 chemistry, with lower pH values and higher concentrations in various major, trace and metal  
51  
52 1129 elements in stations situated close to/South of the discharge outlet, and in the inter-harbour  
53  
54  
55 1130 area (El Kateb et al., 2018; El Zrelli et al., 2015, 2018). The fact that the phosphogypsum  
56  
57 1131 followed a North to South direction, and accumulate in the area between Ghannouch and  
58  
59 1132 Gabès harbours is consistent with results from Zayen et al. (2020), who reported, during our  
60  
61  
62  
63  
64  
65



1133 sampling period and in our sampling area, the occurrence of a tidally-averaged coastal current  
1  
21134 flowing from the North to the South, very likely in relation with wind and the regional  
3  
4  
51135 current. Dissolved organic contaminants (PAHs, PCBs and OCPs) did not closely follow the  
6  
71136 spatial distribution of salinity, FDOM, O<sub>2</sub>, SPM, POC and PON (Fig. 3), neither were  
8  
9  
101137 correlated to these parameters as seen from the PCA (Fig. 5). They displayed less marked  
11  
121138 coast-offshore and North-South gradients (particularly for PCBs and DDTs), although PAHs  
13  
141139 and HCHs presented higher concentrations in the Ghannouch/effluent zone (Fig. 3). This was  
15  
16  
171140 due to the fact that dissolved PAHs, PCBs and OCPs in coastal surface waters of the Gulf of  
18  
191141 Gabès very likely originated from various sources including probably the phosphogypsum  
20  
21  
221142 effluent but also many others as industrial effluents, wadis, runoffs, sediment resuspension,  
23  
241143 and the atmosphere with aerosol deposition and gas exchanges (Barhoumi et al., 2018; Fourati  
25  
26  
271144 et al., 2018; Zouch et al., 2018).

291145 Concentrations of the targeted PAH, PCB and OCP molecules in surface waters of the Gulf  
30  
311146 of Gabès were generally low. As a consequence of their environmental persistence,  
32  
33  
341147 bioaccumulation and toxicity, the continued exposure to these POPs can pose adverse effects  
35  
361148 to animals and human health, even at very low levels (Kim et al., 2018). Therefore, the  
37  
38  
391149 potential risk of those chemicals on aquatic organisms was evaluated. TEFs were applied here  
40  
411150 to quantify the current mixture of PAHs carcinogenicity in seawater samples (Table 2). The  
42  
43  
441151 results show that in terms of total TEQ<sub>ΣPAHs</sub>, the measured values were found in the range  
45  
461152 between 0.04 and 0.51 ng TEQ L<sup>-1</sup> (mean of 0.14 ng TEQ L<sup>-1</sup>) which were lower than the  
47  
48  
491153 threshold values established for BaP in water, suggesting that at the ongoing concentrations  
50  
511154 the evaluated mixture does not seem able to start carcinogenic responses in aquatic organisms  
52  
531155 (Abdel-Shafy and Mansour, 2016). TEQ<sub>ΣPAHs</sub> observed in surface waters of the Gulf of  
54  
55  
561156 Gabès is comparable to that found in the Atlantic Iberian northwest coastline (Rocha and  
57  
581157 Rocha, 2021).

1158 It is also important to evaluate if the measured concentrations of PAHs pose ecological  
1  
21159 risks. As shown in Table 2,  $RQ_{(MPCs)}$  of individual PAHs in waters from the study area were  
3  
4  
51160 all  $< 1$ , indicating relatively low ecosystem risk (Cao et al., 2010).  $RQ_{(NCs)}$  of Chr was  $< 1$   
6  
71161 indicating almost no risk to the ecosystem, whereas  $RQ_{(NCs)}$  of the other PAHs was  $> 1$  in at  
8  
9  
101162 least 7% of samples indicating moderate risks. Concerning the mix of the twelve PAHs  
11  
121163 ( $RQ_{\Sigma PAHs}$ ), as the value of  $RQ_{\Sigma PAHs(NCs)}$  are  $\geq 1$  or  $< 800$  and those of  $RQ_{\Sigma PAHs(MPCs)}$  are  $\approx 0$ , it  
13  
141164 is concluded that, overall, the analyzed PAHs have a low risk to aquatic organisms.

16  
171165 Regarding organochlorines, it can be seen that all  $RQ_{CMCs}$  values for individual OCPs were  
18  
191166  $< 1$  and  $RQ_{CCCs} > 1$  suggesting that risk from measured OCPs is moderate to severe, however,  
20  
21  
221167  $RQ_{CCCs}$  for PCBs are  $< 1$  indicating a low risk (Table 3).

## 241168 25 26 271169 **5. Conclusion**

291170 The results obtained in this study provided the first detailed analysis of dissolved PAH and  
30  
31  
321171 OCs (including PCBs and OCPs) distributions in surface water collected from 27 stations in  
33  
341172 the Gulf of Gabès (Tunisia). PAH levels were lower than those found in other coastal waters  
35  
361173 around the world, while OC levels were moderated. The tendency that emerges is that the  
37  
38  
391174 concentrations we measured in this work in the Ghannouch-Gabès area (Southern Gulf of  
40  
411175 Gabès) were in the mid-range of those reported in other coastal environments. They displayed  
42  
43  
441176 less marked coast-offshore and North-South gradients (particularly for PCBs and DDTs),  
45  
461177 although PAHs and HCHs presented higher concentrations in the Ghannouch/effluent zone.  
47  
48  
491178 Risk assessment based on toxic equivalents (TEQs) indicated that DDTs could induce the  
50  
511179 highest risks for benthic organisms, whereas ecological risks from PAHs and PCBs were  
52  
53  
541180 potentially lower. This work provides baseline information for further studies about this Gulf.  
55  
561181 Since PAHs and OCs levels were only studied in autumn, more studies throughout the whole

1182 year with increasing samples sizes are needed to fully understand the levels of PAHs and OCs  
1  
21183 and the ecological risk assessment of this area.  
3  
4  
5

## 61184 **Funding**

7

8  
91185 This work was financially supported by the CNRS-INSU MISTRALS MERMEX-  
10  
111186 MERITE project and the IRD French-Tunisian International Joint Laboratory (LMI)  
12  
13  
141187 “COSYS-Med”.

## 171188 **CRedit authorship contribution statement**

18

19  
201189 **Hatem Zaghden, Badreddine Barhoumi and Marc Tedetti:** Conceptualization,  
21  
221190 Investigation, Methodology, Data curation, Writing - original draft, and Writing - review &  
23  
24  
251191 editing. **Lobna Jlaiel:** Methodology and Data curation. **Catherine Guigue:** Writing – review  
26  
271192 & editing. **Lassaad Chouba:** Conceptualization and Data curation. **Soufiane Touil and Sami**  
28  
29  
301193 **Sayadi:** Supervision

## 331194 **Declaration of competing interest**

34

351195 Publishing of this paper will cause no conflict of interest with any people, institutions or  
36  
37  
381196 legislation.

## 411197 **Acknowledgements**

42

431198 We are grateful to the National Institute of Marine Science and Technology (INSTM) for  
44  
45  
461199 their logistic support, as well as the captain and the crew of the vessel “*Rhama*” for their  
47  
481200 technical help during the cruise. We acknowledge P. Raimbault, N. Garcia, and the  
49  
50  
511201 Plateforme Analytique de Chimie des Environnements Marins (PACEM platform) of the  
52  
531202 Mediterranean Institute of Oceanography (MIO) for POC, PON and isotopic analyses, as well  
54  
551203 as C. Sammari (INSTM) for the cruise organization and management. We warmly thank  
56  
57  
581204 M.M. Serbaji from the National Engineering School of Sfax (ENIS) for the production of the  
59  
601205 map (Figure 1).  
61

1206 **References**

- 1  
2 1207 Abdel-Shafy, H.I., Mansour, M.S.M., 2016. A review on polycyclic aromatic hydrocarbons:  
3  
4  
5 1208 source, environmental impact, effect on human health and remediation. Egypt. J. of Pet.  
6  
7 1209 25, 107–123. <https://doi.org/10.1016/j.ejpe.2015.03.011>.  
8  
9  
10 1210 Anh, H.Q., Watanabe, I., Tomioka, K., Minh, T.B., Takahashi, S., 2019. Characterization of  
11  
12 1211 209 polychlorinated biphenyls in street dust from northern Vietnam: contamination  
13  
14 1212 status, potential sources, and risk assessment. Sci. of the Tot. Environ. 652, 345–355.  
15  
16  
17 1213 <https://doi.org/10.1016/j.scitotenv.2018.10.240>.  
18  
19 1214 APEK, 2005. Tunisia country situation report. Association for environmental protection in  
20  
21  
22 1215 Kerkennah, Tunisia. <http://www.ipen.org>.  
23  
24 1216 Ashraf, M.A., 2017. Persistent organic pollutants (POPs): A global issue, a global challenge.  
25  
26  
27 1217 Environ. Sci. and Pollut. Res. 24, 4223–4227. [https://doi.org/ 10.1007/s11356-015-](https://doi.org/10.1007/s11356-015-5225-9)  
28  
29 1218 [5225-9](https://doi.org/10.1007/s11356-015-5225-9).  
30  
31  
32 1219 Ayata, S.D., Irisson, J.O., Aubert, A., Berline, L., Dutay, J.C., Mayot, N., Nieblas, A.E.,  
33  
34 1220 D’Ortenzio, F., Palmiéri, J., Reygondeau, G., Rossi, V., Guieu, C., 2018.  
35  
36 1221 Regionalisation of the Mediterranean basin, a MERMEX synthesis. Prog. in Oceanog.  
37  
38  
39 1222 163, 7–20. [https://doi.org/ 10.1016/j.pocean.2017.09.016](https://doi.org/10.1016/j.pocean.2017.09.016).  
40  
41 1223 Barhoumi, B., Castro-Jiménez, J., Guigue, C., Goutx, M., Sempéré, R., Derouiche, A.,  
42  
43  
44 1224 Achour, A., Touil, S., Driss, M.R., Tedetti, M., 2018. Levels and risk assessment of  
45  
46 1225 hydrocarbons and organochlorines in aerosols from a North African coastal city  
47  
48  
49 1226 (Bizerte, Tunisia). Environ. Pollut. 240, 422–431. [https://doi.org/](https://doi.org/10.1016/j.envpol.2018.04.109)  
50  
51 1227 [10.1016/j.envpol.2018.04.109](https://doi.org/10.1016/j.envpol.2018.04.109).  
52  
53 1228 Barhoumi, B., Clerandau, C., Gourves, P.Y., Le Menach, K., El Megdiche, Y., Peluhet, L.,  
54  
55  
56 1229 Budzinski, H., Baudrimont, M., Driss, M.R., Cachot, J., 2014c. Pollution bio-  
57  
58 1230 monitoring in the Bizerte lagoon (Tunisia), using combined chemical and biomarker  
59  
60  
61  
62  
63  
64  
65

- 1231 analyses in grass goby, *Zosterisessor ophiocephalus* (Teleostei, Gobiidae). Mar.  
1  
21232 Environ. Res. 101, 184–195. <https://doi.org/10.1016/j.marenvres.2014.07.002>.  
3  
4  
51233 Barhoumi, B., Jouili, S., Elbarhoumi, A., Derouiche, A., El Megdiche, Y., Bouabdallah, S.,  
6  
71234 Touil, S., Driss, M.R., 2016. Organochlorines in the surface sediments of the Bahiret el  
8  
91235 Bibane lagoon (Tunisia, Southwestern Mediterranean Sea). Chem. Ecol. 9, 884–897.  
10  
11  
121236 <https://doi.org/10.1080/02757540.2016.1195817>.  
13  
14  
151237 Barhoumi, B., Le Menach, K., Clerandau, C., Ben Ameer, W., Budzinski, H., Driss, M.R.,  
16  
171238 Cachot, J., 2014b. Assessment of pollution in the Bizerte lagoon (Tunisia) by the  
18  
191239 combined use of chemical and biochemical markers in mussels, *Mytilus*  
20  
21  
221240 galloprovincialis. Mar. Pollut. Bull. 84, 379–390.  
23  
241241 <https://doi.org/10.1016/j.marpolbul.2014.05.002>.  
25  
26  
271242 Barhoumi, B., Le Menach, K., Devier, M.H., El Megdiche, Y., Hammami, B., Ben Ameer,  
28  
291243 W., Ben Hassine, S., Cachot, J., Budzinski, H., Driss, M.R., 2014a. Distribution and  
30  
311244 ecological risk of polychlorinated biphenyls (PCBs) and organochlorine pesticides  
32  
33  
341245 (OCPs) in surface sediments from the Bizerte lagoon, Tunisia. Environ. Sci. Pollut. Res.  
35  
361246 21, 6290–6302. <https://doi.org/10.1007/s11356-013-1709-7>.  
37  
38  
391247 Barth, A., Alvera-Azcárate, A., Troupin, C., Ouberdous, M., Beckers, J.-M., 2010. A web  
40  
411248 interface for gridding arbitrarily distributed in situ data based on Data-Interpolating  
42  
43  
441249 Variational Analysis (DIVA). Adv. Geosci. 28, 29–37. [https://doi.org/10.5194/adgeo-](https://doi.org/10.5194/adgeo-28-29-2010)  
45  
461250 [28-29-2010](https://doi.org/10.5194/adgeo-28-29-2010).  
47  
48  
491251 Basheer, C., Obbard, J.P., Lee, H.K., 2003. Persistent Organic Pollutants in Singapore's  
50  
511252 Coastal Marine Environment: Part I, Seawater. Water Air Soil Pollut. 149, 295–313.  
52  
53  
541253 <https://doi.org/10.1023/A:1025689600993>.  
55  
561254 Béjaoui, B., Ben Ismail, S., Othmani, A., Ben Abdallah-Ben Hadj Hamida, O., Chevalier, C.,  
57  
581255 Feki-Sahnoun, W., Harzallah, A., Ben Hadj Hamida, N., Bouaziz, R., Dahech, S., Diaz,  
59  
60  
61  
62  
63  
64  
65

- 1256 F., Tounsi, K., Sammari, C., Pagano, M., Bel Hassen, M., 2019. Synthesis review of the  
1  
2 1257 Gulf of Gabes (eastern Mediterranean Sea, Tunisia): morphological, climatic, physical  
3  
4  
5 1258 oceanographic, biogeochemical and fisheries features. *Est. Coast. Shelf Sci.* 219, 395–  
6  
7 1259 408. [https://doi.org/ 10.1016/j.ecss.2019.01.006](https://doi.org/10.1016/j.ecss.2019.01.006).  
8
- 9  
10 1260 Béjaoui, B., Rais, S., Koutitonsky, V., 2004. Modélisation de la dispersion du phosphogypse  
11  
12 1261 dans le golfe de gabès. *Bulletin de l’Institut National des Sciences et Technologie de la*  
13  
14 1262 *Mer de Salammbô*, 31, 103–109.  
15
- 16  
17 1263 Ben Ameer, W., Trabelsi, S., El Megdiche, Y., Ben Hassine, S., Barhoumi, B., Hammami, B.,  
18  
19 1264 Eljarrat, E., Barceló, D., Driss, M.R., 2013. Concentration of polychlorinated biphenyls  
20  
21 1265 and organochlorine pesticides in mullet (*Mugil cephalus*) and sea bass (*Dicentrarchus*  
22  
23 1266 *labrax*) from Bizerte Lagoon (Northern Tunisia). *Chemosphere* 90, 2372–2380.  
24  
25  
26 1267 <https://doi.org/10.1016/j.chemosphere.2012.10.028>.  
27
- 28  
29 1268 Berrojalbiz, N., Dachs, J., Ojeda, M.J., Valle, M.C., Castro-Jiménez, J., Wollgast, J., Ghiani,  
30  
31 1269 M., Hanke, G., Zaldivar, J.M., 2011. Biogeochemical and physical controls on  
32  
33 1270 concentrations of polycyclic aromatic hydrocarbons in water and plankton of the  
34  
35 1271 Mediterranean and Black Seas. *Glo. Biogeo. Cycl.* 25, GB4003. [https:// doi.org/  
36  
37 10.1029/2010GB003775](https://doi.org/10.1029/2010GB003775).  
38  
39 1272
- 40  
41 1273 Campillo, J.A., Santos-Echeandía, J., Fernández, B., 2019. The hydrological regime of a large  
42  
43 1274 Mediterranean river influences the availability of pollutants to mussels at the adjacent  
44  
45 1275 marine coastal area: Implications for temporal and spatial trends. *Chemosphere* 237,  
46  
47 1276 124492. [https://doi.org/ 10.1016/j.chemosphere.2019.124492](https://doi.org/10.1016/j.chemosphere.2019.124492).  
48  
49
- 50  
51 1277 Cao, Z., Liu, J., Luan, Y., Li, Y., Ma, M., Xu, J., Han, S., 2010. Distribution and ecosystem  
52  
53 1278 risk assessment of polycyclic aromatic hydrocarbons in the Luan River, China.  
54  
55 1279 *Ecotoxicology* 19, 827–837. <https://doi.org/10.1007/s10646-010-0464-5>.  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1280 Cornelissen, G., Wiberg, K., Broman, D., Arp, H.P.H., Persson, Y., Sundqvist, K., Jonsson,  
1  
21281 P., 2008. Freely Dissolved Concentrations and Sediment-Water Activity Ratios of  
3  
41282 PCDD/Fs and PCBs in the Open Baltic Sea. *Environ. Sci. Technol.* 42, 23, 8733–8739.  
5  
6  
71283 [https:// doi.org/10.1021/es8018379](https://doi.org/10.1021/es8018379).  
8  
9  
101284 Dachs, J., Lohmann, R., Ockenden, W.A., Méjanelle, L., Eisenreich, S.J., Jones, K.C., 2002.  
11  
121285 Oceanic biogeochemical controls on global dynamics of persistent organic pollutants.  
13  
141286 *Environ. Sci. Technol.* 36, 4229–4237. [https://doi.org/ 10.1021/es025724k](https://doi.org/10.1021/es025724k).  
15  
16  
171287 Dachs, J., Méjanelle, L., 2010. Organic pollutants in coastal waters, sediments, and biota: A  
18  
191288 relevant driver for ecosystems during the anthropocene. *Estuar. Coast.* 33, 1–14.  
20  
21  
221289 <https://doi.org/10.1007/s12237-009-9255-8>.  
23  
241290 Daewel, U., Yakushev, E.V., Schrum, C., Nizzetto, L., Mikheeva, E., 2020. Understanding  
25  
261291 the Role of Organic Matter Cycling for the Spatio- Temporal Structure of PCBs in the  
27  
28  
291292 North Sea. *Water* 12, 817. <https://doi.org/10.3390/w12030817>.  
30  
31  
321293 Darmoul, B., Hadj Ali Salem, M., Vitiello, P., 1980. Effets des rejets industriels de la région  
33  
341294 de Gabès (Tunisie) sur le milieu marin récepteur. *Bulletin de l’Institut National*  
35  
361295 *Scientifique et Technique d’Océanographie et de Pêche de Salammbô*, 7, 5–61.  
37  
38  
391296 DGPA, 2015. *Annuaire des Statistiques des Pêches en Tunisie (Année 2015)*. Direction  
40  
411297 Générale de la Pêche et de l’Aquaculture, Tunis.  
42  
43  
441298 D’Ortenzio, F., d’Alcalà, M.R., 2009. On the trophic regimes of the Mediterranean Sea: A  
45  
461299 satellite analysis. *Biogeosciences* 6, 139–148. <https://doi.org/10.5194/bg-6-139-2009>.  
47  
48  
491300 Duan, X., Li, Y., Li, X., Li, M., Zhang, D., 2013. Distributions and sources of polychlorinated  
50  
511301 biphenyls in the coastal East China Sea sediments. *Sci. of the Tot. Environ.* 463-464C,  
52  
531302 894–903. <https://doi.org/10.1016/j.scitotenv.2013.06.102>.  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1303 Duran, R., Cravo-Laureau, C., 2016. Role of environmental factors and microorganisms in  
1  
21304 determining the fate of polycyclic aromatic hydrocarbons in the marine environment.  
3  
4  
51305 FEMS Microb. Lett. 40, 814–830. <https://doi.org/10.1093/femsre/fuw031>.  
6
- 71306 El Kateb, A., Stalder, C., Rüggeberg, A., Neururer, C., Spangenberg, J.E., Spezzaferri, S.,  
8  
9  
101307 2018. Impact of industrial phosphate waste discharge on the marine environment in the  
11  
121308 Gulf of Gabes (Tunisia). PLoS ONE 13, e0197731. [https://doi.org/](https://doi.org/10.1371/journal.pone.0197731)  
13  
141309 [10.1371/journal.pone.0197731](https://doi.org/10.1371/journal.pone.0197731).  
15
- 16  
171310 El Nemr, A., Abd-Allah, A.M.A., 2003. Contamination of polycyclic aromatic hydrocarbons  
18  
191311 (PAHs) in microlayer and subsurface waters along Alexandria coast, Egypt. Chemosphere  
20  
21  
221312 52, 1711–1716. [https://doi.org/10.1016/S0045-6535\(03\)00300-X](https://doi.org/10.1016/S0045-6535(03)00300-X).  
23
- 241313 El Zrelli, R., Courjault-Radé, P., Rabaoui, L., Castet, S., Michel, S., Bejaoui, N., 2015. Heavy  
25  
26  
271314 metal contamination and ecological risk assessment in the surface sediments of the  
28  
291315 coastal area surrounding the industrial complex of Gabes city, Gulf of Gabes, SE  
30  
311316 Tunisia. Mar. Pollut. Bull. 101, 922–929. [https://doi.org/](https://doi.org/10.1016/j.marpolbul.2015.10.047)  
32  
331317 [10.1016/j.marpolbul.2015.10.047](https://doi.org/10.1016/j.marpolbul.2015.10.047).  
34  
35
- 361318 El Zrelli, R., Rabaoui, L., Ben Alaya, M., Daghbouj, N., Castet, S., Besson, P., Michel, S.,  
37  
38  
391319 Bejaoui, N., Courjault-Radé, P., 2018. Seawater quality assessment and identification of  
40  
411320 pollution sources along the central coastal area of Gabes Gulf (SE Tunisia): Evidence of  
42  
43  
441321 industrial impact and implications for marine environment protection. Mar. Pollut. Bull.  
45  
461322 127, 445–452. [https://doi.org/ 10.1016/j.marpolbul.2017.12.012](https://doi.org/10.1016/j.marpolbul.2017.12.012).  
47
- 48  
491323 Fourati, R., Tedetti, M., Guigue, C., Goutx, M., Garcia, N., Zaghden, H., Sayadi, S., Elleuch,  
50  
511324 B., 2018. Sources and spatial distribution of dissolved aliphatic and polycyclic aromatic  
52  
531325 hydrocarbons in surface coastal waters from the Gulf of Gabès (Tunisia, Southern  
54  
55  
561326 Mediterranean Sea). Progr. Oceanogr. 163, 232–247. [https://doi.org/](https://doi.org/10.1016/j.pocean.2017.02.001)  
57  
581327 [10.1016/j.pocean.2017.02.001](https://doi.org/10.1016/j.pocean.2017.02.001).  
59  
60  
61  
62  
63  
64  
65



- 1328 Galanopoulou, S., Vgenopoulos, A., Conispoliatis, N., 2005. DDTs and other chlorinated  
1 pesticides and polychlorinated biphenyls pollution in the surface sediments of Kerstini  
2 1329 Harbor, Saronikos gulf, Greece. *Mar. Pollut. Bull.* 50, 520–525.  
3  
4 1330 <https://doi.org/10.1016/j.marpolbul.2004.11.043>.  
5  
6  
7 1331  
8  
9 1332 Garcia-Flor, N., Guitart, C., Abalos, M., Dachs, J., Bayona, J.M., Albaiges, J., 2005.  
10  
11 Enrichment of organochlorine contaminants in the sea surface microlayer: An organic  
12 1333 carbon-driven process. *Mar. Chem.* 96, 331–345.  
13  
14 1334 <https://doi.org/10.1016/j.marchem.2005.01.005>.  
15  
16  
17 1335  
18  
19 1336 Gómez-Gutiérrez, A., Garnacho, E., Bayona, J.M., Albaigés, J., 2007. Screening ecological  
20  
21 risk assessment of persistent organic pollutants in Mediterranean Sea sediments.  
22 1337 *Environ. Int.* 33, 867–876. <https://doi.org/10.1016/j.envint.2007.04.002>.  
23  
24 1338  
25  
26 1339 Guigue, C., Tedetti, M., Ferretto, N., Garcia, N., Méjanelle, L., Goutx, M., 2014. Spatial and  
27  
28 seasonal variabilities of dissolved hydrocarbons in surface waters from the  
29 1340 Northwestern Mediterranean Sea: results from one year intensive sampling. *Sci. Total*  
30  
31 1341 *Environ.* 466–467, 650–662. <https://doi.org/10.1016/j.scitotenv.2013.07.082>.  
32  
33  
34 1342  
35  
36 1343 Guigue, C., Tedetti, M., Giorgi, S., Goutx, M., 2011. Occurrence and distribution of  
37  
38 hydrocarbons in the surface microlayer and subsurface water from the urban coastal  
39 1344 marine area off Marseilles, Northwestern Mediterranean Sea. *Mar. Pollut. Bull.* 62,  
40  
41 1345 2741–2752. <https://doi.org/10.1016/j.marpolbul.2011.09.013>.  
42  
43  
44 1346  
45  
46 1347 Guo, W., Pan, B., Sakkiah, S., Yavas, G., Ge, W., Zou, W., Tong, W., Hong, H., 2019.  
47  
48 Persistent Organic Pollutants in Food: Contamination Sources, Health Effects and  
49 1348 Detection Methods. *Int. J. Environ. Res. Public Health* 16, 4361.  
50  
51 1349 <https://doi.org/10.3390/ijerph16224361>.  
52  
53 1350  
54  
55 1351 Hattab, T., Ben Rais Lasram, F., Albouy, C., Romdhane, M.S., Jarboui, O., Halouani, G.,  
56  
57 Cury, P., Le Loc'h, F., 2013. An ecosystem model of an exploited southern  
58 1352  
59  
60  
61  
62  
63  
64  
65

- 1353 Mediterranean shelf region (Gulf of Gabes, Tunisia) and a comparison with other  
1  
21354 Mediterranean ecosystem model properties. *J. Mar. Syst.* 128, 159–174.  
3  
4  
51355 <https://doi.org/10.1016/j.jmarsys.2013.04.017>.  
6  
71356 Hitch, R.K., Day, H.R., 1992. Unusual persistence of DDT in some Western USA soils. *Bull.*  
8  
9  
101357 *Environ. Contam. Toxicol.* 48, 259–264. <https://doi.org/10.1007/BF00194381>.  
11  
121358 Hoai, P.M., Ngoc, N.T., Minh, N.H., Viet, P.H., Berg, M., Alder, A.C., Giger, W., 2010.  
13  
141359 Recent levels of organochlorine pesticides and polychlorinated biphenyls in sediments  
15  
16  
171360 of the sewer system in Hanoi, Vietnam. *Environ. Pollut.* 158, 913–920. [https://doi.org/](https://doi.org/10.1016/j.envpol.2009.09.018)  
18  
191361 [10.1016/j.envpol.2009.09.018](https://doi.org/10.1016/j.envpol.2009.09.018).  
20  
21  
221362 Hong, S.H., Yim, U.H., Shim, W.J., Oh, J.R., Lee, I.S., 2003. Horizontal and vertical  
23  
241363 distribution of PCBs and chlorinated pesticides in sediments from Masan Bay, Korea.  
25  
26  
271364 *Mar. Pollut. Bul.* 46, 244–253. [https://doi.org/10.1016/s0025-326x\(02\)00399-5](https://doi.org/10.1016/s0025-326x(02)00399-5).  
28  
291365 Iwata, H., Tanabe, S., Tatsukawa, R., 1993. A new view on the divergence of HCH isomer  
30  
31  
321366 compositions in oceanic air. *Mar. Pollut. Bull.* 26, 302–305. [https://doi.org/](https://doi.org/10.1016/0025-326X(93)90571-Z)  
33  
341367 [10.1016/0025-326X\(93\)90571-Z](https://doi.org/10.1016/0025-326X(93)90571-Z).  
35  
361368 Jamieson, A., Malkocs, T., Piertney, S., Fujii, T., Zhang, Z., 2017. Bioaccumulation of  
37  
38  
391369 persistent organic pollutants in the deepest ocean fauna. *Nat. Ecol. Evol.* 1, 0051.  
40  
411370 <https://doi.org/10.1038/s41559-016-0051>.  
42  
43  
441371 Jones, K.C., de Voogt, P., 1999. Persistent organic pollutants (POPs): State of the science.  
45  
461372 *Environ. Pollut.* 100, 209–221. [https://doi.org/10.1016/s0269-7491\(99\)00098-6](https://doi.org/10.1016/s0269-7491(99)00098-6).  
47  
48  
491373 Katsoyiannis, A., Breivik, K., 2014. Model-based evaluation of the use of polycyclic aromatic  
50  
511374 hydrocarbons molecular diagnostic ratios as a source identification tool. *Environ. Pollut.*  
52  
531375 184, 488–494. <https://doi.org/10.1016/j.envpol.2013.09.028>.  
54  
55  
561376 Kim, S.A., Lee, Y.M., Choi, J.Y., Jacobs, D., Lee, D., 2018. Evolutionarily adapted hormesis-  
57  
581377 inducing stressors can be a practical solution to mitigate harmful effects of chronic  
59  
60  
61  
62  
63  
64  
65

- 1378 exposure to low dose chemical mixtures. *Environ. Pollut.* 233, 725–734. [https://](https://doi.org/10.1016/j.envpol.2017.10.124)  
1  
21379 [doi.org/10.1016/j.envpol.2017.10.124](https://doi.org/10.1016/j.envpol.2017.10.124).  
3  
4  
51380 Kobayashi, J., Serizawa, S., Sakurai, T., Imaizumi, Y., Suzuki, N., Horiguchi, T., 2010.  
6  
71381 Spatial distribution and partitioning of polychlorinated biphenyls in Tokyo Bay, Japan.  
8  
9  
101382 *J. Environ. Monit.* 12, 838–845. <https://doi.org/10.1039/B925541A>.  
11  
121383 Köck-Schulmeyer, M., Ginebreda, A., Petrovic, M., Giulivo, M., Aznar-Aleman, Ò.,  
13  
141384 Eljarrat, E., Valle-Sistac, J., Molins-Delgado, D., Diaz-Cruz, M.S., Monllor-Alcaraz,  
15  
16 L.S., Guillem-Argiles, N., Martínez, E., de Alda Miren, L., Llorca, M., Farré, M., Peña,  
171385 J.M., Mandaric, L., Pérez, S., Majone, B., Bellin, A., Kalogianni, E., Skoulikidis, N.  
18  
191386 Th., Milačič, R., Barceló, D., 2021. Priority and emerging organic microcontaminants in  
20  
21  
221387 three Mediterranean river basins: Occurrence, spatial distribution, and identification of  
23  
241388 river basin specific pollutants. *Sci. Total Environ.* 754, 142344. [https://doi.org/](https://doi.org/10.1016/j.scitotenv.2020.142344)  
25  
261389 [10.1016/j.scitotenv.2020.142344](https://doi.org/10.1016/j.scitotenv.2020.142344).  
27  
28  
291390  
30  
311391 Lee, C.L., Huang, H.C., Wang, C.C., Sheu, C.C., Wu, C.C., Leung, S.Y., Lai, R.S., Lin, C.C.,  
32  
33  
341392 Wei, Y.F., Lai, I.C., Jiang, H., Choug, W.L., Chung, W.Y., Huang, M.S., Huang, S.K.,  
35  
361393 2016. A new grid-scale model simulating the spatiotemporal distribution of PM<sub>2.5</sub>-PAHs  
37  
38 for exposure assessment. *J. Hazard Mater.* 314, 286–294.  
391394 <https://doi.org/10.1016/j.jhazmat.2016.04.047>.  
40  
411395  
42  
431396 Lee, K.T., Tanabe, S., Koh, C.H., 2001. Distribution of organochlorine pesticides in  
44  
45 sediments from Kyeonggi Bay and nearby areas, Korea. *Environ. Pollut.* 114, 207–213.  
461397 [https://doi.org/10.1016/S0269-7491\(00\)00217-7](https://doi.org/10.1016/S0269-7491(00)00217-7).  
47  
48  
491398  
50  
511399 Li, W., Yang, H., Jiang, X., Liu, Q., Sun, Y., Zhou, J., 2016. Residues and distribution of  
52  
53 organochlorine pesticides in water and suspended particulate matter from Hangzhou  
541400 Bay, East China Sea. *Bull. Environ. Contam. Toxicol.* 96, 295–302.  
55  
561401 <https://doi.org/10.1007/s00128-016-1739-1>.  
57  
581402  
59  
60  
61  
62  
63  
64  
65

- 1403 Lin, B.S., Brimblecombe, P., Lee, C.L., Liu, J.T., 2013. Tracing typhoon effects on particulate  
1  
21404 transport in a submarine canyon using polycyclic aromatic hydrocarbons. *Mar. Chem.*  
3  
4  
51405 157, 1–11. <https://doi.org/10.1016/j.marchem.2013.07.004>.  
6
- 71406 Liu, M., Cheng, S., Ou, D., Yang, Y., Liu, H., Hou, L., Gao, L., Xu, S., 2008. Organochlorine  
8  
9  
101407 pesticides in surface sediments and suspended particulate matters from the Yangtze  
11  
121408 estuary, China. *Environ. Pollut.* 156, 168–173. [https://doi.org/](https://doi.org/10.1016/j.envpol.2007.12.015)  
13  
14  
151409 [10.1016/j.envpol.2007.12.015](https://doi.org/10.1016/j.envpol.2007.12.015).  
16
- 171410 Luo, Y., Guo, W., Ngo, H.H., Nghiem, L.D., Hai, F.I., Zhang, J., Liang, S., Wang, X.C.,  
18  
191411 2014. A review on the occurrence of micropollutants in the aquatic environment and  
20  
21  
221412 their fate and removal during wastewater treatment. *Sci. Total Environ.* 473, 619–641.  
23  
241413 [https:// doi.org/10.1016/j.scitotenv.2013.12.065](https://doi.org/10.1016/j.scitotenv.2013.12.065).  
25
- 26  
271414 Ma, Y., Halsall, C.J., Crosse, J.D., Graf, C., Cai, M., He, J., Gao, G., Jones, K., 2015.  
28  
291415 Persistent organic pollutants in ocean sediments from the North Pacific to the Arctic  
30  
31  
321416 *Ocean. J. Geophys. Res. Oceans* 120, 2723–2735.  
33  
341417 <https://doi.org/10.1002/2014JC010651>.  
35
- 361418 Mackay, D., Shiu, W.Y., Ma, K.C., 1992. *Illustrated handbook of physical–chemical.*  
37  
38  
391419 *Monoaromatic Hydrocarbons, Chlorobenzenes and PCBs*, vol. I. Lewis Publishers,  
40  
411420 Chelsea, MI, pp. 697.  
42
- 43  
441421 Malik, A., Ojha, P., Singh, K.P., 2009. Levels and distribution of persistent organochlorine  
45  
461422 pesticide residues in water and sediments of Gomti River (India)-a tributary of the  
47  
48  
491423 Ganges River. *Environ. Monit. Assess.* 148, 421–435. [https://doi.org/ 10.1007/s10661-](https://doi.org/10.1007/s10661-008-0172-2)  
50  
511424 [008-0172-2](https://doi.org/10.1007/s10661-008-0172-2).  
52
- 53  
541425 Menzies, R., Quinete, N.S., Gardinali, P., Seba, D., 2013. Baseline occurrence of  
55  
561426 organochlorine pesticides and other xenobiotics in the marine environment: Caribbean  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1427 and Pacific collections. *Mar. Pollut. Bull.* 70, 289–295. [https://](https://doi.org/10.1016/j.marpolbul.2013.03.003)  
1  
21428 [doi.org/10.1016/j.marpolbul.2013.03.003](https://doi.org/10.1016/j.marpolbul.2013.03.003).  
3  
4  
51429 Mhadhbi, T., Pringault, O., Nouri, H., Spinelli, S., Beyrem, H., Gonzalez, C., 2019.  
6  
71430 Evaluating polar pesticide pollution with a combined approach: a survey of agricultural  
8  
9  
101431 practices and POCIS passive samplers in a Tunisian lagoon watershed. *Environ. Sci.*  
11  
121432 *Pollut. Res.* 26, 342–361. [https:// doi.org/10.1007/s11356-018-3552-3](https://doi.org/10.1007/s11356-018-3552-3).  
13  
14  
151433 Miraji, H., Ripanda, A., Moto, E., 2021. A review on the occurrences of persistent organic  
16  
171434 pollutants in corals, sediments, fish and waters of the Western Indian Ocean. *Egypt. J.*  
18  
191435 *Aquat. Res.* 47, 373–379. <https://doi.org/10.1016/j.ejar.2021.08.003>.  
20  
21  
221436 Mitra, S., Corsolini, S., Pozo, K., Audy, O., Sarkar, S.K., Biswas, J.K., 2019.  
23  
241437 Characterization, source identification and risk associated with polyaromatic and  
25  
26  
271438 chlorinated organic contaminants (PAHs, PCBs, PCBzs and OCPs) in the surface  
28  
291439 sediments of Hooghly estuary, India. *Chemosphere* 221, 154–165. [https://doi.org/](https://doi.org/10.1016/j.chemosphere.2018.12.173)  
30  
311440 [10.1016/j.chemosphere.2018.12.173](https://doi.org/10.1016/j.chemosphere.2018.12.173).  
32  
33  
341441 Moeckel, C., Monteith, D.T., Llewellyn, N.R., Henrys, P.A., Pereira, M.G., 2014.  
35  
361442 Relationship between the concentrations of dissolved organic matter and polycyclic  
37  
38  
391443 aromatic hydrocarbons in a typical UK upland stream. *Environ. Sci. Technol.* 48, 130–  
40  
411444 138. [https://doi.org/ 10.1021/es403707q](https://doi.org/10.1021/es403707q).  
42  
43  
441445 Montory, M., Ferrer, J., Rivera, D., Villouta, M.V., Grimalt, J.O., 2017. First report on  
45  
461446 organochlorine pesticides in water in a highly productive agro-industrial basin of the  
47  
48  
491447 Central Valley, Chile. *Chemosphere* 174, 148–156. [https://doi.org/](https://doi.org/10.1016/j.chemosphere.2016.12.125)  
50  
511448 [10.1016/j.chemosphere.2016.12.125](https://doi.org/10.1016/j.chemosphere.2016.12.125).  
52  
53  
541449 Montuori, P., Cirillo, T., Fasano, E., Nardone, A., Esposito, F., Triassi, M., 2014. Spatial  
55  
561450 distribution and partitioning of polychlorinated biphenyl and organochlorine pesticide in  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1451 water and sediment from Sarno River and Estuary, Southern Italy. *Environ. Sci. Pollut.*  
1  
21452 Res. 21, 5023–5035. <https://doi.org/10.1007/s11356-013-2419-x>.  
3  
4  
51453 Montuori, P., De Rosa, E., Sarnacchiaro, P., Di Duca, F., Provisiero, D.P., Nardone, A.,  
6  
71454 Triassi, M., 2020. Polychlorinated biphenyls and organochlorine pesticides in water and  
8  
91455 sediment from Volturno River, Southern Italy: occurrence, distribution and risk  
10  
11  
121456 assessment. *Environ. Sci. Euro.* 32, 123. <https://doi.org/10.1186/s12302-020-00408-4>.  
13  
14  
151457 Moreno-González, R., Campillo, J.A., García, V., León, V.M., 2013. Seasonal input of  
16  
171458 regulated and emerging organic pollutants through surface watercourses to a  
18  
191459 Mediterranean coastal lagoon. *Chemosphere* 92, 247–257. [https://doi.org/](https://doi.org/10.1016/j.chemosphere.2012.12.022)  
20  
21  
221460 [10.1016/j.chemosphere.2012.12.022](https://doi.org/10.1016/j.chemosphere.2012.12.022).  
23  
241461 Moret, I., Gambaro, A., Piazza, R., Ferrari, S., Manodori, L., 2005. Determination of  
25  
261462 polychlorobiphenyl congeners (PCBs) in the surface water of the Venice lagoon. *Mar.*  
27  
28  
291463 *Pollut. Bull.* 50, 167–174. <https://doi.org/10.1016/j.marpolbul.2004.10.005>.  
30  
31  
321464 Mostafa, A.R., Wade, T.L., Sweet, S.T., Al-Alimi, A.K.A., Barakat, A.O., 2007. Assessment  
33  
341465 of persistent organochlorine residues in sediments of Hadramout coastal area, Gulf of  
35  
361466 Aden, Yemen. *Mar. Pollut. Bull.* 54, 1053–1058. [https://doi.org/](https://doi.org/10.1016/j.marpolbul.2007.03.018)  
37  
38  
391467 [10.1016/j.marpolbul.2007.03.018](https://doi.org/10.1016/j.marpolbul.2007.03.018).  
40  
411468 Muir, D.C.G., Howard, P.H., 2006. Are there other persistent organic pollutants? A challenge  
42  
43  
441469 for environmental chemists. *Environ. Sci. Technol.* 40, 7157–7166. [https://](https://doi.org/10.1021/es061677a)  
45  
461470 [doi.org/10.1021/es061677a](https://doi.org/10.1021/es061677a).  
47  
48  
491471 Mzoughi, N., Chouba, L., 2011. Distribution and partitioning of aliphatic hydrocarbons and  
50  
511472 polycyclic aromatic hydrocarbons between water, suspended particulate matter, and  
52  
531473 sediment in harbours of the West coastal of the Gulf of Tunis (Tunisia). *J. Environ.*  
54  
55  
561474 *Monit.* 13, 689–698. <https://doi.org/10.1039/c0em00616e>.  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1475 Nedia, G., Chafai, A., Moncef, S.M., Chokri, Y., 2011. Spatial distribution of heavy metals in  
1  
21476 the coastal zone of “sfax-Kerkennah” plateau, Tunisia. *Environ Prog. Susta. Ener.* 30,  
3  
4  
51477 221–233. <https://doi.org/10.1002/ep.10462>.  
6
- 71478 Neukermans, G., Ruddick, K., Loisel, H., Roose, P., 2012. Optimization and quality control  
8  
9  
101479 of suspended particulate matter concentration measurement using turbidity  
11  
121480 measurements. *Limno. Oceanogr. Meth.* 10, 1011–1023. [https://doi.org/](https://doi.org/10.4319/lom.2012.10.1011)  
13  
141481 [10.4319/lom.2012.10.1011](https://doi.org/10.4319/lom.2012.10.1011).  
15
- 16  
171482 Nguyen, T.C., Loganathan, P., Nguyen, T.V., Vigneswaran, S., Kandasamy, J., Stevenson,  
18  
191483 D.S.G., Naidu, R., 2014. Polycyclic aromatic hydrocarbons in road-deposited  
20  
21  
221484 sediments, water sediments, and soils in Sydney, Australia: comparisons of  
23  
241485 concentration distribution, sources and potential toxicity. *Ecotoxicol. Environ. Saf.* 104,  
25  
26  
271486 339–348. <https://doi.org/10.1016/j.ecoenv.2014.03.010>.  
28
- 291487 Nisbet, I.C.T., LaGoy, P.K., 1992. Toxic equivalency factors (TEFs) for polycyclic aromatic  
30  
31  
321488 hydrocarbons (PAHs). *Regul. Toxicol. Pharmac.* 16, 290–300.  
33  
341489 [https://doi.org/10.1016/0273-2300\(92\)90009-X](https://doi.org/10.1016/0273-2300(92)90009-X).  
35
- 361490 Nyarko, E., Botwe, B.O., Klubi, E., 2011. Polycyclic aromatic hydrocarbons (PAHs) levels in  
37  
38  
391491 two commercially important fish species from the coastal waters of Ghana and their  
40  
411492 carcinogenic health risks. *West African J. App. Ecol.* 19, 53–66.  
42
- 43  
441493 Rabodonirina, S., Net, S., Ouddane, B., Merhaby, D., Dumoulin, D., 2105. Distribution of  
45  
461494 persistent organic pollutants (PAHs, Me-PAHs, PCBs) in dissolved, particulate and  
47  
48  
491495 sedimentary phases in freshwater systems. *Environ. Pollut.* 206, 38–48. [https://doi.org/](https://doi.org/10.1016/j.envpol.2015.06.023)  
50  
511496 [10.1016/j.envpol.2015.06.023](https://doi.org/10.1016/j.envpol.2015.06.023).  
52
- 531497 Raimbault, P., Garcia, N., Cerrutti, F., 2008. Distribution of inorganic and organic nutrients in  
54  
55  
561498 the South Pacific Ocean. Evidence for long-term accumulation of organic matter in  
57  
58  
59  
60  
61  
62  
63  
64  
65

1499 nitrogen-depleted waters. *Biogeosciences* 5, 281–298. [https://doi.org/ 10.5194/bg-5-](https://doi.org/10.5194/bg-5-)  
1  
21500 [281-2008](https://doi.org/10.5194/bg-5-281-2008).  
3  
4  
51501 Ravindra, K., Bencs, L., Wauters, E., Hoog, J.D., Deutsch, F., Roekens, E., Bleux, N.,  
6  
71502 Berghmans, P., Grieken, R.V., 2006. Seasonal and site specific variation in vapour and  
8  
9  
101503 aerosol phase PAHs over Flanders (Belgium) and their relation with anthropogenic  
11  
121504 activities. *Atmos. Environ.* 40, 771–785. [https://doi.org/](https://doi.org/10.1016/j.atmosenv.2005.10.011)  
13  
141505 [10.1016/j.atmosenv.2005.10.011](https://doi.org/10.1016/j.atmosenv.2005.10.011).  
15  
16  
171506 Rocha, M.J., Ribeiro, A.B., Campos, D., Rocha, E., 2021. Temporal-spatial survey of PAHs  
18  
191507 and PCBs in the Atlantic Iberian northwest coastline, and evaluation of their sources and  
20  
21  
221508 risks for both humans and aquatic organisms. *Chemosphere* 279, 130506.  
23  
241509 <https://doi.org/10.1016/j.chemosphere.2021.130506>.  
25  
26  
271510 Rocha, M.J., Rocha, E., 2021. Concentrations, sources and risks of PAHs in dissolved and  
28  
291511 suspended material particulate fractions from the Northwest Atlantic Coast of the  
30  
31  
321512 Iberian Peninsula. *Mar. Pollut. Bull.* 165, 112143. [https://doi.org/](https://doi.org/10.1016/j.marpolbul.2021.112143)  
33  
341513 [10.1016/j.marpolbul.2021.112143](https://doi.org/10.1016/j.marpolbul.2021.112143).  
35  
361514 Salvadó, J.A., Grimalt, J.O., López, J.F., Palanques, A., Canals, M., 2019. Influence of deep  
37  
38  
391515 water formation by open-sea convection on the transport of low hydrophobicity organic  
40  
411516 pollutants in the NW Mediterranean Sea. *Sci. Total Environ.* 647, 597–605.  
42  
43  
441517 <https://doi.org/10.1016/j.scitotenv.2018.07.458>.  
45  
461518 Sánchez-Avila, J., Tauler, R., Lacorte, S., 2012. Organic micropollutants in coastal waters  
47  
48  
491519 from NW Mediterranean Sea: sources distribution and potential risk. *Environ. Intern.*  
50  
511520 46, 50–62. <https://doi.org/10.1016/j.envint.2012.04.013>.  
52  
53  
541521 Shen, B., Wu, J., Zhan, S., Jin, M., 2021. Residues of organochlorine pesticides (OCPs) and  
55  
561522 polycyclic aromatic hydrocarbons (PAHs) in waters of the Ili-Balkhash Basin, arid  
57  
58  
59  
60  
61  
62  
63  
64  
65



- 1523 Central Asia: Concentrations and risk assessment. *Chemosphere* 273, 129705.  
1  
2 1524 <https://doi.org/10.1016/j.chemosphere.2021.129705>.  
3  
4  
5 1525 Shunthirasingham, C., Oyiliagu, C.E., Cao, X., Gouin, T., Wania, F., Lee, S.-C., Pozo, K.,  
6  
7 1526 Harner, T., Muir, D.C.G., 2010. Spatial and temporal pattern of pesticides in the global  
8  
9  
10 1527 atmosphere. *J. Environ. Monit.* 12, 1650–1657. <https://doi.org/10.1039/c0em00134a>.  
11  
12 1528 Soclo, H.H., Garrigues, P., Ewald, M., 2000. Origin of polycyclic aromatic hydrocarbons  
13  
14 1529 (PAHs) in coastal marine sediments: case studies in Cotonou (Benin) and Aquitaine  
15  
16  
17 1530 (France) areas. *Mar. Pollut. Bull.* 40, 387–396. [https://doi.org/10.1016/S0025-](https://doi.org/10.1016/S0025-326X(99)00200-3)  
18  
19 1531 [326X\(99\)00200-3](https://doi.org/10.1016/S0025-326X(99)00200-3).  
20  
21  
22 1532 Stogiannidis E., Laane, R., 2015. Source Characterization of Polycyclic Aromatic  
23  
24 1533 Hydrocarbons by Using Their Molecular Indices: An Overview of Possibilities. In:  
25  
26  
27 1534 Whitacre D. (eds) *Reviews of Environ. Cont. Toxicol. Continuation of Residue*  
28  
29 1535 *Reviews*. Vol 234. Springer, Cham. [https://doi.org/10.1007/978-3-319-10638-0\\_2](https://doi.org/10.1007/978-3-319-10638-0_2).  
30  
31  
32 1536 Su, Q.K., Qi, S.H., Wu, C.X., Julia, E.B., Liu, H.F., Fang, M., Li, J., Zhang, G., 2006.  
33  
34 1537 Organochlorine pesticides in marine environment of quanzhou bay, southeast China.  
35  
36 1538 *Chin. J. Geochem.* 25 (B08), 190. <https://doi.org/10.1007/BF02840111>.  
37  
38  
39 1539 Takasuga, T., Senthilkumar, K., Matsumura, T., Shiozaki, K., Sakai, S.I., 2006. Isotope  
40  
41 1540 dilution analysis of polychlorinated biphenyls (PCBs) in transformer oil and global  
42  
43  
44 1541 commercial PCB formulations by high resolution gas chromatography–high resolution  
45  
46 1542 mass spectrometry. *Chemosphere* 62, 469–484. [https://doi.org/](https://doi.org/10.1016/j.chemosphere.2005.04.034)  
47  
48 1543 [10.1016/j.chemosphere.2005.04.034](https://doi.org/10.1016/j.chemosphere.2005.04.034).  
49  
50  
51 1544 Tang, D., Liu, X., He, H., Cui, Z., Xia, Z., 2020. Distribution, sources and ecological risks of  
52  
53 1545 organochlorine compounds (DDTs, HCHs and PCBs) in surface sediments from the  
54  
55  
56 1546 Pearl River Estuary, China. *Mar. Pollut. Bull.* 152, 110942. [https://doi.org/](https://doi.org/10.1016/j.marpolbul.2020.110942)  
57  
58 1547 [10.1016/j.marpolbul.2020.110942](https://doi.org/10.1016/j.marpolbul.2020.110942).  
59  
60  
61  
62  
63  
64  
65

- 1548 Tedetti, M., Guigue, C., Goutx, M., 2010. Utilization of a submersible UV fluorometer for  
1  
2 1549 monitoring anthropogenic inputs in the Mediterranean coastal waters. *Mar. Pollut. Bull.*  
3  
4  
5 1550 60, 350–362. [https://doi.org/ 10.1016/j.marpolbul.2009.10.018](https://doi.org/10.1016/j.marpolbul.2009.10.018).  
6
- 7 1551 The MerMex Group, 2011. Marine ecosystems responses to climatic and anthropogenic  
8  
9  
10 1552 forcings in the Mediterranean. *Progr. Oceanogr.* 91, 97–166. [https://doi.org/](https://doi.org/10.1016/j.pocean.2011.02.003)  
11  
12 1553 [10.1016/j.pocean.2011.02.003](https://doi.org/10.1016/j.pocean.2011.02.003).  
13
- 14 1554 Tolosa, I., Readman, J.W., Fowler, S.W., Villeneuve, J.P., Dachs, J., Bayona, J.M., Albaigés,  
15  
16  
17 1555 J., 1997. PCBs in the western Mediterranean. Temporal trends and mass balance  
18  
19 1556 assessment. *Deep Sea Res. Part II Top. Stud. Oceanogr.* 44, 907–928.  
20  
21  
22 1557 [https://doi.org/10.1016/S0967-0645\(96\)00104-X](https://doi.org/10.1016/S0967-0645(96)00104-X).  
23
- 24 1558 Tong, Y., Chen, L., Liu, Y., Wang, Y., Tian, S., 2019. Distribution, sources and ecological  
25  
26  
27 1559 risk assessment of PAHs in surface seawater from coastal Bohai Bay, China. *Mar. Pollut.*  
28  
29 1560 *Bull.* 142, 520–524. <https://doi.org/10.1016/j.marpolbul.2019.04.004>.  
30
- 31 1561 Troupin, C., Barth, A., Sirjacobs, D., Ouberdous, M., Brankart, J.-M., Brasseur, P., Rixen, M.,  
32  
33  
34 1562 Alvera Azcarate, A., Belounis, M., Capet, A., Lenartz, F., Toussaint, M.-E., Beckers, J.-  
35  
36 1563 M., 2012. Generation of analysis and consistent error fields using the Data Interpolating  
37  
38  
39 1564 Variational Analysis (Diva). *Ocean Model.* 52–53, 90–101.  
40  
41 1565 <https://doi.org/10.1016/j.ocemod.2012.05.002>.  
42
- 43 1566 United States Environmental Protection Agency (USEPA), 2006. National Recommended  
44  
45  
46 1567 Water Quality Criteria (NRWQC) (4304T).  
47
- 48 1568 Wang, Z., Fingas, M., Page, D.S., 1999. Oil spill identification. *J. Chromato. A* 843, 369–411.  
49  
50  
51 1569 [https://doi.org/10.1016/S0021-9673\(99\)00120-X](https://doi.org/10.1016/S0021-9673(99)00120-X).  
52
- 53 1570 Wania, F., Mackay, D., 1995. A global distribution model for persistent organic chemicals.  
54  
55  
56 1571 *Sci. Total Environ.* 160, 211–232. [https://doi.org/10.1016/0048-9697\(95\)04358-8](https://doi.org/10.1016/0048-9697(95)04358-8).  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1572 Willett, K.L., Ulrich, E.M., Hites, R.A., 1998. Differential toxicity and environmental fates of  
1  
21573 hexachlorocyclohexane isomers. *Environ. Sci. Technol.* 32, 2197–2207. [https://](https://doi.org/10.1021/es9708530)  
3  
41574 [doi.org/10.1021/es9708530](https://doi.org/10.1021/es9708530).  
5  
6  
71575 Yao, T., He, C., Zhang, P., Gao, H., Zhou, C., 2013. Distribution and sources of  
8  
91576 polychlorinated biphenyls (PCBs) and organochlorine pesticides (OCPs) in surface waters  
10  
11 of Jinzhou Bay in China. *Procedia Environ. Sci.* 18, 317–322.  
121577  
13  
141578 <https://doi.org/10.1016/j.proenv.2013.04.041>.  
15  
16  
171579 Yunker, M.B., Macdonald, R.W., Vingarzan, R., Mitchell, R.H., Goyette, D., Sylvestre, S.,  
18  
191580 2002. PAHs in the Fraser River basin: a critical appraisal of PAH ratios as indicators of  
20  
21 PAH source and composition. *Org. Geochem.* 33, 489–515.  
221581  
23  
241582 [https://doi.org/10.1016/S0146-6380\(02\)00002-5](https://doi.org/10.1016/S0146-6380(02)00002-5).  
25  
26  
271583 Zaghden, H., Kallel, M., Elleuch, B., Oudot, J., Saliot, A., 2007. Sources and distribution of  
28  
291584 aliphatic and polyaromatic hydrocarbons in sediments of Sfax, Tunisia, Mediterranean  
30  
311585 Sea. *Mar. Chem.* 105, 70–89. <https://doi.org/10.1016/j.marchem.2006.12.016>.  
32  
33  
341586 Zayen, A., Sayadi, S., Chevalier, C., Boukthir, M., Ben Ismail, S., Tedetti, M., 2020.  
35  
361587 Microplastics in surface waters of the Gulf of Gabes, southern Mediterranean Sea:  
37  
38 Distribution, composition and influence of hydrodynamics. *Estuar. Coast. Shelf Sci.*  
391588  
40  
411589 242, 106832. <https://doi.org/10.1016/j.ecss.2020.106832>.  
42  
43  
441590 Zhang, K., Wei, Y.L., Zeng, E.Y., 2013. A review of environmental and human exposure to  
45  
461591 persistent organic pollutants in the Pearl River Delta, South China. *Sci. Total Environ.*  
47  
48 463–464, 1093–1110. <https://doi.org/10.1016/j.scitotenv.2012.10.104>.  
491592  
50  
511593 Zhao, Z., Gong, X., Zhang, L., Jin, M., Cai, Y., Wang, X., 2021. Riverine transport and  
52  
531594 water-sediment exchange of polycyclic aromatic hydrocarbons (PAHs) along the middle-  
54  
55 lower Yangtze River, China. *J. Hazard. Mater.* 403, 123973.  
561595  
57  
581596 <https://doi.org/10.1016/j.jhazmat.2020.123973>.  
59  
60  
61  
62  
63  
64  
65

1597 Zhou, J.L., Maskaoui, K., Qiu, Y.W., Hong, H.S., Wang, Z.D., 2001. Polychlorinated  
1 biphenyl congeners and organochlorine insecticides in the water column and sediments of  
2 1598 Daya Bay, China. Environ. Pollut. 113, 373–384. [https://doi.org/10.1016/S0269-](https://doi.org/10.1016/S0269-7491(00)00180-9)  
3  
4 1599  
5  
6  
7 1600  
8  
9  
10 1601 Zouch, H., Cabrol, L., Chifflet, S., Tedetti, M., Karray, F., Zaghden, H., Sayadi, S.,  
11  
12 1602 Quéméneur, M., 2018. Effect of acidic industrial effluent release on microbial diversity  
13  
14 1603 and trace metal dynamics during resuspension of coastal sediment. Front. Microbiol. 9,  
15  
16 1604 3103. <https://doi.org/10.3389/fmicb.2018.03103>.  
17  
18  
19 1605  
20  
21 1606  
22  
23 1607  
24  
25 1608  
26  
27 1609  
28  
29 1610  
30  
31 1611  
32  
33 1612  
34  
35 1613  
36  
37 1614  
38  
39 1615  
40  
41 1616  
42  
43 1617  
44  
45 1618  
46  
47 1619  
48  
49 1620  
50  
51 1621  
52  
53 1622  
54  
55 1623  
56  
57 1624  
58  
59 1625  
60  
61  
62  
63  
64  
65

1626 **Figure captions**

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Figure 1.** Location of the study stations in the Ghannouch-Gabès coastal area in the Gulf of Gabès (Tunisia, Southern Mediterranean Sea). The detailed characteristics of the stations are given in [Table 1](#).

**Figure 2.** Spatial distribution of a) temperature, b) salinity, c) pH, d) fluorescent dissolved organic matter (FDOM) at 254-nm excitation wavelength and 360-nm emission wavelength (in relative unit, RU), e) dissolved oxygen (O<sub>2</sub>) concentration, and f) redox potential in coastal surface waters (at ~ 0.5-m depth) of the Gulf of Gabès. Spatial interpolation was made using Data-Interpolating Variational Analysis (DIVA) method from Ocean Data View (ODV) software version 4.6.5. <http://odv.awi.de>.

**Figure 3.** Spatial distribution of concentrations of total dissolved a) PAHs (sum of 16 compounds), b) PCBs (sum of 20 congeners), c) HCHs (sum of 4 compounds), and d) DDTs (sum of 6 compounds) (in ng L<sup>-1</sup>) in coastal surface waters (at ~ 0.5-m depth) of the Gulf of Gabès. Spatial interpolation was made using Data-Interpolating Variational Analysis (DIVA) method from Ocean Data View (ODV) software version 4.6.5. <http://odv.awi.de>.

**Figure 4.** Relative abundance (in %) of a) 12 PAHs, b) 9 PCBs congeners, and c) 7 OCPs in each station.

**Figure 5.** Principal component analysis (PCA), based on the Spearman 's correlation matrix, applied on various parameters measured in coastal surface waters (at ~ 0.5-m depth) of the Gulf of Gabès, i.e., salinity (Sal), pH, dissolved oxygen concentration (O<sub>2</sub>), fluorescent dissolved organic matter (FDOM), and concentrations of total dissolved PAHs ( $\Sigma$  PAHs), PCBs ( $\Sigma$  PCBs) and OCPs ( $\Sigma$  OCPs). Projection of variables (green lines) and samples (blue circles) on the first factorial plane (PC1 *versus* PC2). Four groups of stations have been highlighted (G1-G4).

1651 **Table captions**

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65


**Table 1.** Total concentrations of dissolved PAHs, PCBs, OCPs, HCHs, DDTs and associated ratios from seawater samples collected at ~ 0.5-m depth in coastal area of the Gulf of Gabès (Tunisia, Southern Mediterranean Sea).

**Table 2.** Toxic equivalence quantities (TEQs) and mean values of  $RQ_{(NCs)}$  and  $RQ_{(MPCs)}$  of PAHs from seawater samples collected at ~ 0.5-m depth in coastal area of the Gulf of Gabès (Tunisia, Southern Mediterranean Sea).


**Table 3.** Mean values of RQ (CCCs) and RQ (CMCs) for OCPs and PCBs from seawater samples collected at ~ 0.5-m depth in coastal area of the Gulf of Gabès (Tunisia, Southern Mediterranean Sea).

16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1677 **Figure 1**



1695 **Figure 2**




1  
21696  
3  
41697


5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
521698  
53  
541699  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65




1700 **Figure 3**



1707 **Figure 4**



1736 **Figure 5**



1744 **Table 1.** Total concentrations of dissolved PAHs, PCBs, OCPs, HCHs, DDTs and associated  
 1745 ratios from seawater samples collected at ~ 0.5-m depth in coastal area of the Gulf of Gabès  
 1746 (Tunisia, Southern Mediterranean Sea).

| Stations      | PAHs                              |               |               | OCs                               |                                   |                                  |                                  |             |
|---------------|-----------------------------------|---------------|---------------|-----------------------------------|-----------------------------------|----------------------------------|----------------------------------|-------------|
|               | Σ16 PAHs<br>(ng L <sup>-1</sup> ) | Flt/(Flt+Pyr) | Ant/(Ant+Phe) | Σ20 PCBs<br>(ng L <sup>-1</sup> ) | Σ10 OCPs<br>(ng L <sup>-1</sup> ) | Σ4 HCHs<br>(ng L <sup>-1</sup> ) | Σ6 DDTs<br>(ng L <sup>-1</sup> ) | DDD/ DDE    |
| S1            | 32.7                              | 0.71          | 0.01          | 22.0                              | 11.0                              | 4.3                              | 6.6                              | 0.77        |
| S2            | 27.7                              | 0.43          | nd            | 2.9                               | 6.8                               | 1.3                              | 5.5                              | 1.21        |
| S3            | 25.0                              | 0.43          | nd            | 4.3                               | 12.7                              | 4.1                              | 8.6                              | 0.78        |
| S4            | 59.8                              | 0.79          | 0.02          | 11.6                              | 16.6                              | 4.5                              | 12.1                             | 1.48        |
| S5            | 27.1                              | 0.50          | nd            | 4.6                               | 13.9                              | 2.6                              | 11.3                             | 0.80        |
| S6            | 32.8                              | 0.73          | 0.04          | 6.9                               | 16.1                              | 2.9                              | 13.2                             | 0.69        |
| S7            | 19.5                              | 0.90          | 0.16          | 8.1                               | 6.0                               | 1.1                              | 4.9                              | 3.74        |
| S8            | 71.2                              | 0.49          | 0.21          | 11.2                              | 26.7                              | 14.8                             | 12.0                             | 0.71        |
| S9            | 62.7                              | 0.47          | 0.15          | 33.6                              | 13.5                              | 3.0                              | 10.5                             | 0.64        |
| S10           | 17.6                              | 0.40          | nd            | 7.3                               | 6.7                               | 2.2                              | 4.6                              | 0.63        |
| S11           | 35.3                              | 0.54          | 0.05          | 31.4                              | 12.3                              | 4.0                              | 8.2                              | 0.21        |
| S12           | 28.3                              | 0.52          | 0.04          | 9.6                               | 5.6                               | 4.5                              | 1.1                              | nd          |
| S13           | 33.5                              | 0.55          | 0.03          | 28.3                              | 6.1                               | 2.3                              | 3.8                              | 2.24        |
| S14           | 26.8                              | 0.44          | 0.06          | 4.3                               | 5.3                               | 2.3                              | 2.9                              | 2.38        |
| S15           | 27.5                              | 0.51          | 0.03          | 4.7                               | 11.8                              | 5.6                              | 6.2                              | nd          |
| S16           | 27.4                              | 0.49          | 0.03          | 13.3                              | 16.7                              | 5.1                              | 11.7                             | 1.13        |
| S17           | 29.0                              | 0.55          | 0.03          | 6.7                               | 8.9                               | 3.8                              | 5.0                              | 3.41        |
| S18           | 20.6                              | 0.48          | nd            | 4.6                               | 5.9                               | 1.6                              | 4.3                              | 3.25        |
| S19           | 45.0                              | 0.53          | nd            | 12.7                              | 18.5                              | 7.0                              | 11.5                             | 1.13        |
| S20           | 40.5                              | nd            | 0.02          | 8.5                               | 13.7                              | 6.4                              | 7.3                              | 0.85        |
| S21           | 27.7                              | 0.81          | 0.08          | 10.1                              | 8.2                               | 3.9                              | 4.3                              | 0.57        |
| S22           | 64.9                              | 0.48          | 0.18          | 4.6                               | 19.0                              | 9.4                              | 9.7                              | 0.31        |
| S23           | 30.9                              | 0.61          | 0.15          | 9.4                               | 10.4                              | 1.5                              | 8.9                              | 0.87        |
| S24           | 46.1                              | 0.43          | 0.12          | 6.9                               | 13.7                              | 4.6                              | 9.1                              | 2.93        |
| S25           | 28.1                              | 0.73          | 0.04          | 10.7                              | 10.7                              | 1.5                              | 9.2                              | 0.57        |
| S26           | 22.9                              | 0.72          | 0.04          | 7.6                               | 13.3                              | 2.3                              | 10.9                             | 0.77        |
| S27           | 25.3                              | 0.47          | 0.02          | 4.8                               | 12.3                              | 3.6                              | 8.7                              | 0.42        |
| <b>Min</b>    | <b>17.6</b>                       | <b>0.40</b>   | <b>0.01</b>   | <b>2.9</b>                        | <b>5.3</b>                        | <b>1.1</b>                       | <b>1.1</b>                       | <b>0.21</b> |
| <b>Max</b>    | <b>71.2</b>                       | <b>0.90</b>   | <b>0.21</b>   | <b>33.6</b>                       | <b>26.7</b>                       | <b>14.8</b>                      | <b>13.2</b>                      | <b>3.74</b> |
| <b>Mean</b>   | <b>34.7</b>                       | <b>0.56</b>   | <b>0.07</b>   | <b>10.8</b>                       | <b>11.9</b>                       | <b>4.1</b>                       | <b>7.9</b>                       | <b>1.30</b> |
| <b>Median</b> | <b>28.3</b>                       | <b>0.51</b>   | <b>0.04</b>   | <b>8.1</b>                        | <b>12.3</b>                       | <b>3.8</b>                       | <b>8.6</b>                       | <b>0.80</b> |

nd: not determined.

16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Table 2.** Toxic equivalence quantities (TEQs) and mean values of  $RQ_{(NCs)}$  and  $RQ_{(MPCs)}$  of PAHs from seawater samples collected at ~ 0.5-m depth in coastal area of the Gulf of Gabès (Tunisia, Southern Mediterranean Sea).

| PAHs                  | TEQs (ng L <sup>-1</sup> ) |       |       |       | RQs for PAHs          |       |        |        |                    |                        |       |       |       |
|-----------------------|----------------------------|-------|-------|-------|-----------------------|-------|--------|--------|--------------------|------------------------|-------|-------|-------|
|                       | TEFs                       | Min   | Max   | Mean  | NECs                  | Min   | Max    | Mean   | $RQ_{(NCs)} > 1^*$ | MPCs                   | Min   | Max   | Mean  |
| Nap                   | 0.001                      | 0.005 | 0.039 | 0.021 | 12                    | 0.436 | 3.236  | 1.776  | 93                 | 1200                   | 0.004 | 0.032 | 0.018 |
| Acy                   | 0.001                      | 0.000 | 0.002 | 0.001 | 0.7                   | 0.000 | 3.469  | 0.741  | 30                 | 70                     | 0.000 | 0.035 | 0.007 |
| Ace                   | 0.001                      | 0.000 | 0.010 | 0.002 | 0.7                   | 0.000 | 13.770 | 2.830  | 81                 | 70                     | 0.000 | 0.138 | 0.028 |
| Flu                   | 0.001                      | 0.000 | 0.009 | 0.002 | 0.7                   | 0.332 | 12.960 | 2.708  | 56                 | 70                     | 0.003 | 0.130 | 0.027 |
| Phe                   | 0.001                      | 0.002 | 0.015 | 0.004 | 3                     | 0.592 | 4.949  | 1.389  | 59                 | 300                    | 0.006 | 0.050 | 0.014 |
| Ant                   | 0.001                      | 0.000 | 0.002 | 0.001 | 0.7                   | 0.000 | 2.304  | 0.409  | 15                 | 70                     | 0.000 | 0.023 | 0.004 |
| Flt                   | 0.01                       | 0.005 | 0.185 | 0.028 | 3                     | 0.154 | 6.159  | 0.766  | 19                 | 300                    | 0.002 | 0.062 | 0.008 |
| Pyr                   | 0.001                      | 0.000 | 0.005 | 0.001 | 0.7                   | 0.000 | 6.849  | 1.654  | 81                 | 70                     | 0.000 | 0.069 | 0.017 |
| BaA                   | 0.1                        | 0.000 | 0.078 | 0.027 | 0.1                   | 0.000 | 7.771  | 2.655  | 89                 | 10                     | 0.000 | 0.078 | 0.027 |
| Chr                   | 0.01                       | 0.000 | 0.011 | 0.004 | 3.4                   | 0.000 | 0.311  | 0.104  | 0                  | 340                    | 0.000 | 0.003 | 0.001 |
| BbF                   | 0.1                        | 0.000 | 0.310 | 0.030 | 0.1                   | 0.000 | 31.000 | 2.082  | 7                  | 10                     | 0.000 | 0.310 | 0.021 |
| BkF                   | 0.1                        | 0.000 | 0.277 | 0.029 | 0.4                   | 0.000 | 6.926  | 0.531  | 11                 | 40                     | 0.000 | 0.069 | 0.005 |
| TEQs <sub>∑PAHs</sub> |                            | 0.040 | 0.510 | 0.137 | $RQ_{\sum PAHs(NCs)}$ | 6.550 | 73.170 | 17.646 |                    | $RQ_{\sum PAHs(MPCs)}$ | 0.070 | 0.730 | 0.176 |

Percent of stations with  $RQ_{(NCs)}$  of PAHs >1.

1753 **Table 3.** Mean values of RQ (CCCs) and RQ (CMCs) for OCPs and PCBs from seawater  
 1754 samples collected at ~ 0.5-m depth in coastal area of the Gulf of Gabès (Tunisia, Southern  
 1755 Mediterranean Sea).

|                            | $\gamma$ -HCH | DDEs  | DDDs  | PCBs   |
|----------------------------|---------------|-------|-------|--------|
| CMCs (ng L <sup>-1</sup> ) | 950           | 1100  | 1100  | -      |
| CCCs (ng L <sup>-1</sup> ) | -             | 1     | 1     | 4      |
| MEC*                       | 2.286         | 4.295 | 3.568 | 10.771 |
| RQ <sub>CMCs</sub>         | 0.002         | 0.004 | 0.003 | -      |
| RQ <sub>CCCs</sub>         | -             | 4.295 | 3.568 | 0.769  |

\*Average Measured Concentration for OCPs and PCBs in this study (ng L<sup>-1</sup>).