



HAL
open science

Variability of bone preservation in a confined environment: the case of the catacomb of Sts Peter and Marcellinus (Rome, Italy)

Salesse Kevin, Élise Dufour, Matthieu Lebon, Chris M Wurster, Dominique Castex, Jaroslav Bruzek, Antoine Zazzo

► To cite this version:

Salesse Kevin, Élise Dufour, Matthieu Lebon, Chris M Wurster, Dominique Castex, et al.. Variability of bone preservation in a confined environment: the case of the catacomb of Sts Peter and Marcellinus (Rome, Italy). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 2014. hal-03677853

HAL Id: hal-03677853

<https://hal.science/hal-03677853v1>

Submitted on 25 May 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Variability of bone preservation in a confined environment: the case of the**
2 **catacomb of Sts Peter and Marcellinus (Rome, Italy)**

3
4 Salesse K.^{1*}, Dufour E.², Lebon M.³, Wurster C.⁴, Castex D.¹, Bruzek J.^{1,5}, Zazzo A.²

5
6 ¹Univ. Bordeaux, CNRS, UMR 5199: “PACEA – De la Préhistoire à l’Actuel: Culture,
7 Environnement et Anthropologie”, Bâtiment B8, Allée Geoffroy S^t Hilaire, CS 50023, F-
8 33615 Pessac cedex, France.

9 ²Muséum national d'Histoire naturelle, CNRS, Département Ecologie et Gestion de la
10 Biodiversité, USM 303/UMR 7209: “Archéozoologie, Archéobotanique : Sociétés, Pratiques
11 et Environnements”, CP 56, 55 rue Buffon, F-75231 Paris cedex 05, France.

12 ³Muséum national d'Histoire naturelle, CNRS, Département de Préhistoire, UMR 7194:
13 “Histoire naturelle de l’Homme préhistorique”, CP 57, 57 rue Cuvier, F-75231 Paris cedex
14 05, France.

15 ⁴Centre for Tropical Environmental and Sustainability Science, School of Earth and
16 Environmental Sciences, James Cook University, Cairns QLD 4870, Australia.

17 ⁵Department of Anthropology and Human Genetics, Faculty of Science, Charles University,
18 Viničná 7, 128 43 Praha 2, Czech Republic.

19
20 *corresponding author: K. Salesse, Tel: +33 (0)5-40-00-37-39, Email: kevin.salesse@u-bordeaux.fr

21
22 **Keywords:** diagenesis; stable isotopes, collagen; apatite; radiocarbon; Fourier-transform
23 infrared spectroscopy

24
25 **Abstract**

26 Most of the studies investigating the diagenetic trajectory of fossil bones focus on
27 open-air sites and very little work have been published in confined environments such as
28 catacombs. While the stable thermal history of catacombs should favor bone preservation, the
29 accumulation of corpses over a short period of time could favor their destruction. The aim of
30 this study is to describe the diagenetic trajectory of 128 human bone samples coming from six
31 different burial chambers of the catacomb of Sts Peter and Marcellinus (SSPM, Rome, Italy).
32 A multi-proxy approach was undertaken to provide an assessment of the molecular
33 preservation as well as a direct record of the isotopic composition itself. Collagen yield,
34 carbon and nitrogen abundances, C:N ratio, FT-IR based collagen and carbonate contents and
35 crystallinity index, radiocarbon dating and stable isotope analysis of bone collagen and
36 carbonate indicate that both the mineral and the organic fractions are impacted by diagenesis
37 to various degrees, and that bones originating from the small burial chambers are more
38 affected than those coming from the large ones. While some of the bones were strongly
39 recrystallized, the impact of bone diagenesis on the stable isotope values of bone carbonate
40 was limited. Comparison with contemporary sites from the Latium showed that conditions
41 prevailing in catacombs seem overall to favor, rather than disadvantage bone preservation.

42

43

44 **1. Introduction**

45 Excavations conducted from 2004 to 2010 in the central area of the catacomb of Sts
46 Peter and Marcellinus (hereafter the SSPM catacomb) located in the south-east of Rome
47 revealed several mass graves dated to between the first and the third century AD based on
48 radiocarbon and artefact (coins) dating (Castex et al., 2007; Castex et al., 2011). They
49 contained tens to hundreds of articulated human skeletons laid together according to well-
50 reasoned management (Blanchard et al., 2007; Castex et al., 2007; Castex et al., 2009; Castex

51 et al., 2011). A significant number of these individuals received a specific funerary treatment
52 characterized, in particular, by the use of gypsum to partially or entirely cover the corpses
53 (Castex and Blanchard, 2011; Castex et al., 2009; Devière et al., 2010; Vanhove, 2006). A
54 mortality crisis following an epidemic episode is thought to be at the origin of these mass
55 graves suggesting that they have been used for a relatively limited period of time (Blanchard
56 et al., 2007; Castex and Blanchard, 2011; Castex et al., 2009; Castex et al., 2011; Kacki et al.,
57 2013). Unfortunately, the study of the biological identity (sex, age at death, stature, non-
58 metric traits, pathological conditions, *etc.*) of the individuals was restricted due to the state of
59 skeletal preservation (Blanchard et al., 2007; Castex and Blanchard, 2011; Castex et al., 2007;
60 Castex et al., 2009). Many bones and teeth have lost their anatomical integrity even though
61 the preservation quality differs strongly among burial chambers, different deposit levels
62 within a chamber, and different skeletons within the same level (Blanchard et al., 2007;
63 Castex and Blanchard, 2011; Castex et al., 2007; Castex et al., 2009). Stable isotope
64 approaches offer invaluable assistance when the macroscopic anthropological observations
65 are limited by the poor quality of skeletal preservation (*e.g.* Katzenberg et al., 2009; Oelze et
66 al., 2012; Wright et al., 2010). Measurements of the stable isotope composition of the SSPM
67 population would allow us to better understand their life history, provided that *in vivo* isotopic
68 compositions are preserved.

69 Checking whether the original isotope compositions of biomineral fractions are not
70 altered by diagenesis is an essential prerequisite to interpret isotope analyses (Bocherens et
71 al., 2005; Bocherens et al., 2008; Brady et al., 2008; Shin and Hedges, 2012). Most of the
72 studies investigating the diagenetic trajectory of fossil bones focus on open-air sites. To our
73 knowledge, very little work has been published in confined environments. The advantage of
74 such an approach was demonstrated by Bocherens et al. (2008) in Chauvet cave. In this
75 Palaeolithic site, the authors showed that the preservation of bear bones exposed on the floor

76 of the cave varied widely probably due to spatial variation in local conditions. The Chauvet
77 cave results cannot be directly translated to man-made cavities such as catacombs because the
78 age and taphonomic contexts are quite different. Like in caves, the thermal history of
79 catacombs is more stable than in open-air sites which should *a priori* favor their preservation.
80 On the other hand, the accumulation of corpses over a short period of time in catacombs could
81 have an effect on the soil pH and therefore favor the destruction of human bones. The aim of
82 this study is to describe the diagenetic trajectory of the human bone samples from the SSPM
83 catacomb in order to assess their preservation and compare it with the preservation observed
84 in contemporary open-air sites from the region of Rome. A multi-proxy approach
85 (measurement of classical collagen preservation indicators; Fourier-transform infrared
86 spectroscopy (FT-IR) based collagen preservation indicators, carbonate content and
87 crystallinity, ^{14}C measurements, and stable isotope analysis of the collagen and carbonate
88 fractions) was undertaken to provide a record of the preservation of the molecules. Our results
89 show that both the mineral and the organic fractions are affected by diagenesis to various
90 degrees, and that bones originating from the small burial chambers are the most diagenetically
91 altered. While some of the bones were strongly recrystallized, the stable isotope values of
92 bone carbonate remained reliable. Comparison with contemporary sites from the Latium
93 showed that conditions prevailing in catacombs seem overall to favor, rather than
94 disadvantage bone preservation

95

96 **2. Study area and material**

97 A total of 128 individuals were selected for this study. The material comes from six of
98 the seven burial chambers discovered in 2003 in a previously unexplored area (region X) of
99 the SSPM catacomb (Blanchard et al., 2007; Castex et al., 2007) (Figure 1). These burial
100 chambers are part of a vast funerary complex that covers approximately 3 ha with 4.5 km of

101 underground galleries (Guyon, 1987). Four of them (X78/T15, X80/T16, X81 and X82/T18)
102 are small in size ($< 7\text{m}^2$), while the remaining two (X83 and X84) are larger in size ($> 9\text{m}^2$)
103 (Castex et al., 2007; Castex et al., 2009). Different phases of multiple deposits separated by a
104 layer of sediment were identified in the small chambers (Blanchard et al., 2007; Castex et al.,
105 2011; Kacki et al., 2013; Sachau-Carcel et al., 2013). At least three phases were recognized in
106 X78/T15 (Pagni and Burdassi, 2004) whereas 9 and 12 phases were distinguished in X80/T16
107 and X82/T18, respectively (Blanchard et al., 2007; Castex et al., 2007). In contrast, no
108 stratum of soil was discovered between the skeletons in X81, X83 and X84 (Castex and
109 Blanchard, 2011; Castex et al., 2009; Castex et al., 2011).

110 For each individual, a fragment of long (*i.e.* humerus, radius, ulna, metacarpus, femur
111 and tibia), flat (*i.e.* cranial vault, rib and hip bone), irregular (*i.e.* mandible and maxilla) or
112 short (calcaneus) bone was sampled. Due to the poor skeletal representation of bone remains
113 of five individuals from the burial chamber X82/T18, several fragments of different skeletal
114 parts were selected in order to obtain enough material to achieve the required analyses
115 (hereafter named miscellaneous samples). Modern reference material required for FT-IR
116 analysis was composed of bones of modern ox, calf, horse and sheep in order to take into
117 account variation due to species and age at death.

118

119 **3. Methods**

120 3.1. Bone carbonate preparation and analysis

121 Between 30 and 36 mg of bone powder (grain size $< 0.3\text{ mm}$) was first soaked in 2 to
122 3% sodium hypochlorite at room temperature for 48 h to remove organic matter then rinsed
123 five times with distilled water. The remaining fraction was treated with 1 M acetic acid for 1 h
124 to remove exogenous carbonate then rinsed five times with distilled water. Sodium
125 hypochlorite and acetic acid solutions were renewed half-way through each process. Samples

126 were oven-dried for 18 h at 65°C following pretreatment. The purification process resulted in
127 a weight loss comprising between 10 and 80%. Bone carbonate $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ values were
128 measured at the Service de Spectrométrie de Masse Isotopique of the MNHN (SSMIM).
129 Samples weighing from 580 to 630 mg were reacted for 360 s with 100% phosphoric acid at
130 70°C in individual vessels in an automated cryogenic distillation system (Kiel IV device),
131 interfaced with a ThermoScientific Delta V Advantage isotope ratio mass spectrometer. The
132 stable isotope results are reported in per mil (‰) relative to VPDB. Over the period of
133 analysis, the analytical precision calculated based on repeated measurements of our internal
134 carbonate standard (Marble LM) was 0.02‰ for $\delta^{13}\text{C}_{\text{carb}}$ and 0.05‰ for $\delta^{18}\text{O}_{\text{carb}}$ values.

135

136 3.2. Bone collagen extraction and analysis

137 Bone collagen extraction followed Longin's (1971) protocol modified by DeNiro and
138 Epstein (1981) and Bocherens et al. (1988; 1991). Between 0.1 and 1.1 g of bone powder
139 (grain size comprised between 0.3 and 0.7 mm) was soaked in 40 ml of 1 M hydrochloric acid
140 for 20 min at room temperature then filtered for decalcification using 5 μm cellulose filters.
141 The residue was soaked into 0.125 M sodium hydroxide for 20 h at room temperature to
142 oxidize fulvic and humic acids then filtered. The gelatin was then solubilized in 0.01 M
143 hydrochloric acid at 100°C for 17 h, and filtered again to remove impurities. The solubilized
144 collagen was then freeze-dried for 48 to 72 h and extraction yields calculated. Stable isotope
145 composition was measured at the environmental isotope laboratory at James Cook
146 University's Advanced Analytical Centre (Cairns, Australia) using samples weighing between
147 200 and 400 μg . Carbon and nitrogen abundance and isotope composition ($\delta^{13}\text{C}_{\text{col}}$ and $\delta^{15}\text{N}_{\text{col}}$
148 values) were determined using a Costech Elemental Analyzer 4010 fitted with a zero-blank
149 auto-sampler coupled via a ConFlo IV to a Thermo Scientific Delta V^{PLUS} using Continuous-
150 Flow Isotope Ratio Mass Spectrometry. The $\delta^{13}\text{C}_{\text{col}}$ and $\delta^{15}\text{N}_{\text{col}}$ values are reported in per mil

151 (‰) deviation relative to VPDB and AIR, respectively. Over the period of analysis, the
152 carbon and nitrogen abundances were $\pm 5\%$ of the value for laboratory internal standards and
153 the analytical precision for $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ was better than $\pm 0.1\%$ and $\pm 0.2\%$, respectively.
154 Weight percent carbon and nitrogen were used to calculate atomic C:N ratios.

155

156 3.3. FT-IR analysis

157 Attenuated total reflexion mode (ATR) FT-IR analysis was performed on bone powder
158 (grain size < 0.3 mm) before and after pretreatment. Infrared spectra were collected by
159 accumulation of 64 scans with a spectral resolution of 2 cm^{-1} on a Bruker Vector 22 FT-IR
160 spectrometer equipped with a diamond-ATR accessory (Specac Golden Gate). Spectral data
161 analyses were performed using OPUS software (Bruker). Crystallinity was evaluated using
162 the Infrared Splitting Factor (IRSF) (Figure 2). The sum of the two $\nu_4\text{PO}_4$ bands at 565 and
163 600 cm^{-1} were divided by the valley at 590 cm^{-1} following the protocol described by Weiner
164 (1990). Relative carbonate content was measured by the intensity ratio of the $\nu_3\text{CO}_3$ band at
165 1415 cm^{-1} over the $\nu_4\text{PO}_4$ band at 600 cm^{-1} (adapted from Wright and Schwarcz 1996).
166 Relative collagen content was measured by the intensity ratio of the Amide I band at 1660 cm^{-1}
167 over the $\nu_4\text{PO}_4$ band at 600 cm^{-1} (adapted from Trueman et al., 2004). The precipitation of
168 secondary calcite was monitored by the presence of its specific band at 712 cm^{-1} .

169

170 3.4. Radiocarbon dating

171 The collagen and apatite fraction of five individuals from burial chambers X80/T16,
172 X82/T18 and X83 were radiocarbon dated. An aliquot of collagen (between 8 and 13 mg) was
173 combusted for 15 to 60 min at 800°C . Carbonate was purified following a modified version of
174 Krueger et al. (1991). About 2 g of bone powder (grain size < 0.1 mm) was first soaked in 2

175 to 3% sodium hypochlorite at room temperature for six days to oxidize the organic matter.
176 Bone powder was then rinsed several times with distilled water and dried for 3 h at 100°C. It
177 was then placed in 100 ml of 2 M acetic acid for 24 h at room temperature under weak
178 vacuum. The evolution of fine gas bubbles usually slows down drastically after 1 h, and stops
179 after 10+ h, suggesting that all secondary calcite as well as the smallest apatite crystals had
180 been dissolved. The acetic acid treatment resulted in weight losses between 40 and 80%. An
181 aliquot of purified bioapatite weighing between 220 and 280 mg was then reacted in
182 orthophosphoric acid for 12 min at 70°C under vacuum. CO₂ derived from the collagen and
183 apatite samples was cryogenically trapped and about 1 mg C was sealed in a glass tube for
184 radiocarbon dating. Graphitization and AMS measurement were conducted at Queen's
185 University's ¹⁴Chrono Centre (Belfast, Northern Ireland).

186

187 **4. Bone diagenesis evaluation**

188 4.1. Macroscopic preservation

189 Bone samples were classified into three categories according to their macroscopic
190 characteristics as well as their strength and resistance when cutting: (1) “good preservation”
191 characterizes a solid and heavy bone difficult to cut and crush; (2) “poor preservation”
192 corresponds to a bone which is fragile, soft and easy to cut and crush; and finally (3) “medium
193 preservation” refers to a bone with intermediate characteristics (Figure 3).

194

195 4.2. Collagen quality

196 Bone collagen preservation was evaluated based on three criteria: bone collagen
197 extraction yield (wt%), carbon and nitrogen abundances in collagen (%), and atomic C:N ratio
198 (Table 1). Collagen concentration in modern bone is around 20.4 wt% (SD ±3.9 wt%)
199 (Bocherens et al., 1991), and values <1 wt% are usually considered as being unreliable

200 (Dobberstein et al., 2009; van Klinken, 1999). In modern bones, carbon and nitrogen
201 abundances range from 13 to 47% and from 4.8 to 17.3% respectively (Ambrose, 1990), and
202 values below or above these thresholds indicate alteration or contamination (Garvie-Lok,
203 2001; Iacumin et al., 1998; Reitsema, 2012, van Klinken, 1999). Average atomic C:N ratio in
204 modern bones is 3.2 to 3.3 (Ambrose, 1990; van Klinken, 1999), and vary between 2.9 and
205 3.6 (DeNiro, 1985). Atomic ratios below or above these thresholds indicate alteration or
206 contamination (Ambrose, 1990; DeNiro and Weiner, 1988; Grupe et al., 2001). Samples that
207 fail to pass any of these three criteria were considered to be poorly-preserved.

208

209 4.3. Infrared spectroscopy

210 Bone recrystallization promotes the formation of a more stable mineral phase
211 characterized by a higher crystallinity (*i.e.* crystal size and lattice perfection) and lower
212 carbonate content. These changes are related to both microbial activities (bioerosion) and
213 interaction of bone mineral with the burial environment (*i.e.* pH) that control dissolution-
214 recrystallization processes (Berna et al., 2004; Hedges and Millard, 1995; Smith et al., 2007).
215 Crystallinity and carbonate content can be evaluated by FT-IR using the IRSF and CO₃/PO₄
216 ratio, respectively. IRSF values vary between 2.5 and 3.3 for modern bones (Stiner et al.,
217 1995; Wright and Schwarcz, 1996; Nielsen-Marsh and Hedges, 2000; Berna et al., 2004;
218 Munro et al., 2007; Asscher et al., 2011) and usually reaches values between 3.4 and 4.5 for
219 archaeological samples (Weiner and Bar-Yosef, 1990; Stiner et al., 1995; Wright and
220 Schwarcz, 1996; Munro et al., 2007). Bones are usually considered altered for values above
221 3.8 to 4.0 (Shemesh, 1990; Stiner et al., 1995; Berna et al., 2004). In modern animal bones,
222 CO₃/PO₄ ratio is comprised between 0.7 and 1.0. This ratios decreases to less than 0.2 for
223 altered samples (Lebon, 2008). The fate of the mineral phase is mainly dependent on the
224 degradation of the organic phase (Collins et al., 2002; Lebon et al., 2011; Trueman et al.,

225 2008). FT-IR analyses offer also the ability to measure the relative collagen content of a bone
226 by measuring the AmideI/PO₄ ratio. This ratio is comprised between 0.9 and 1.2 for modern
227 bones and decreases as collagen decays. The AmideI/PO₄ and CO₃/PO₄ ratios are usually
228 correlated (Lebon, 2008; Lebon et al., 2011).

229

230 4.4. Radiocarbon dating of bone collagen-carbonate pairs

231 The ¹⁴C/¹²C and ¹³C/¹²C ratios differ by several orders of magnitude. This results in a
232 higher sensitivity of the ¹⁴C/¹²C ratio over the ¹³C/¹²C ratio to post-mortem isotope exchange.
233 Therefore, ¹⁴C is a useful proxy to assess the influence of diagenesis on the carbon stable
234 isotope composition of carbonate in apatite (Shin and Hedges, 2012; Zazzo, in press). To
235 quantify carbon isotope exchange between bone carbonate and dissolved inorganic carbon
236 during diagenesis, we measured the radiocarbon age difference between the collagen and
237 carbonate fraction (¹⁴C_{col-carb}) from the same bone. If bone collagen passes the three tests
238 outlined above, any deviation should be the result of carbon isotope exchange in the carbonate
239 fraction. This difference can then be transformed in percent of carbon isotope exchange
240 assuming that exchange takes place with modern carbon (Hedges et al., 1995a). This is
241 probably a reasonable approximation as an extensive review of the literature and a recent ¹⁴C
242 dating program of several archaeological sites worldwide demonstrated that diagenetically
243 altered apatites are always younger than the reference age for the site (Zazzo, in press; Zazzo
244 and Saliège, 2011). Radiocarbon ages are presented as BP and fraction modern (F_m).

245

246 **5. Results**

247 5.1. Collagen preservation

248 Collagen preservation varies greatly between different samples (Table 2). Collagen
249 yields differ according to bone location within the catacomb, the anatomical part category and

250 the macroscopic aspect (Figure 4). In total, 29 (23%) bone collagen samples do not pass at
251 least one of the three tests defined (collagen yield, carbon and nitrogen abundances, and C:N
252 ratio (Table 3). Within this group, all but two were classified as macroscopically poorly-
253 preserved (Table 3). These samples all originate from three of the the four lowest levels of the
254 chamber X80/T18 and from all the levels of the chamber X82/T18 (Table 3). They consist
255 largely of long and flat bones (31 and 38%, respectively). Additionally, 46% of the flat bones
256 and 100% of the miscellaneous samples have poorly-preserved collagen (Table 3).
257 Macroscopic preservation was variable among the 99 (77%) bone collagen samples with
258 reliable collagen (*e.g.* see Figure 5 for the variations of atomic carbon and nitrogen
259 abundances). Poor scores were obtained for macroscopically poorly-preserved specimens
260 coming from two small chambers (X80/T16 and X82/T18).

261

262 5.2. FT-IR

263 The modern animal reference samples have relatively low IRSF values (Mean = 3.4;
264 SD \pm 0.1) and high CO₃/PO₄ ratios (Mean = 0.85; SD \pm 0.10), consistent with values
265 previously published (Lebon et al., 2010). With the exception of three samples, all the
266 archaeological human bone samples present higher IRSF values and lower CO₃/PO₄ ratios
267 than modern animal reference samples (Figure 6). IRSF values measured on the
268 archaeological samples before pretreatment range from 3.6 to 5.1 (Mean = 4.3; SD \pm 0.4),
269 while CO₃/PO₄ ratios vary between 0.15 and 1.01 (Mean = 0.49; SD \pm 0.20). IRSF values are
270 negatively correlated with bone CO₃/PO₄ ratios ($R^2 = 0.75$) (Figure 6). No trace of calcite was
271 detected (detection limit 1.5 %) in bone samples before pretreatment. Amide I/PO₄ ratios vary
272 from 0.07 to 0.86 (Mean = 0.35; SD \pm 0.25) and are lower for the archaeological bones than
273 for the modern animal reference samples (Figure 7). Carbonate and collagen relative contents
274 are positively correlated ($R^2 = 0.76$) (Figure 7). Small chamber samples are significantly

275 different from large chambers samples with respect to IRSF values, CO₃/PO₄ and Amide
276 I/PO₄ ratios (Mann-Whitney U tests; p-values < 0.03). Samples from small chambers have
277 lower relative carbonate content, lower relative collagen content, and higher IRSF values
278 compared to samples from larger chambers. Following chemical pretreatment of bone
279 carbonate, IRSF values increase (Mean = 4.6; SD ±0.3) and relative carbonate content
280 decrease (Mean = 0.28; SD ±0.10). CO₃/PO₄ ratios remain negatively correlated to IRSF
281 values (R² = 0.59) (Figure 8).

282

283 5.3. Stable isotope analysis

284 Bone collagen carbon and nitrogen isotope values were measured for all specimens
285 that yielded a collagen extract (Table 2). There was no significant difference in δ¹³C_{col}
286 between the well-preserved and the poorly-preserved collagen samples (Mann-Whitney U
287 test; p-value = 0.52). In contrast, the well- and poorly-preserved collagen samples have
288 significantly different δ¹⁵N_{col} values (Mann-Whitney U test; p-value = 0.00). There was no
289 relationship between the δ¹³C_{col} or δ¹⁵N_{col} values and the collagen yield (R² < 0.05), the
290 atomic carbon and nitrogen abundances (R² < 0.2) or the atomic C:N ratios (R² < 0.07). There
291 was no significant difference between the isotope values and the burial chambers, the skeletal
292 part categories, or the macroscopic appearances when either one of the two groups or the
293 totality of the corpus is considered (Mann-Whitney U tests; p-values >> 0.05). We found one
294 exception, however. When samples with well-preserved collagen were considered (n = 99),
295 δ¹⁵N_{col} values measured in macroscopically poorly- and well-preserved samples differed
296 significantly (Mann-Whitney U test; p-value = 0.04).

297 Bone carbonate carbon and oxygen stable isotope values were measured for the 128
298 individuals (Table 2). Bone δ¹⁸O_{carb} values in X82/T18 differed significantly from bone
299 δ¹⁸O_{carb} values in X80/T16 and X84 (Mann-Whitney U test; p-value = 0.01). When the totality

300 of the corpus was considered, $\delta^{13}\text{C}_{\text{carb}}$ values of irregular and flat bones, irregular and long
301 bone, and long and miscellaneous bones differed significantly (Mann-Whitney U tests; p-
302 values < 0.04). The same conclusion was reached (excepted for long vs. miscellaneous bones)
303 when only samples with well-preserved collagen were considered (Mann-Whitney U tests; p-
304 values < 0.04). Moreover, when the totality of the corpus was considered bone
305 macroscopically identified as well-preserved had $\delta^{13}\text{C}_{\text{carb}}$ and $\delta^{18}\text{O}_{\text{carb}}$ values that differed
306 significantly from those measured in medium and poorly-preserved bones, respectively
307 (Mann-Whitney U tests; p-values = < 0.015). The samples which yielded well- (n = 99) and
308 poorly-preserved (n = 29) collagen samples also had significantly different $\delta^{13}\text{C}_{\text{carb}}$ and
309 $\delta^{18}\text{O}_{\text{carb}}$ values (Mann-Whitney U test; p-value = 0.01 and 0.02, respectively). There was no
310 relationship between the $\delta^{13}\text{C}_{\text{carb}}$ or $\delta^{18}\text{O}_{\text{carb}}$ values and the IRSF values ($R^2 < 0.01$), the
311 CO_3/PO_4 ratio ($R^2 < 0.01$) or the AmideI/ PO_4 ratio ($R^2 < 0.01$) before or after bone carbonate
312 pretreatment.

313

314 5.4. Radiocarbon dating

315 The $^{14}\text{C}_{\text{col}}$ ages range from 1779 ± 27 BP to 1965 ± 31 BP whereas the $^{14}\text{C}_{\text{carb}}$ ages
316 range from 1690 ± 27 BP to 1848 ± 22 BP for the five individuals (Table 4). The $^{14}\text{C}_{\text{col-carb}}$ age
317 differences vary between -1 ± 40 and 153 ± 46 ^{14}C yr (Table 4). Except for one individual in
318 chamber X83 where the age measured in the two fractions are identical, the $^{14}\text{C}_{\text{carb}}$ age is
319 always younger than the $^{14}\text{C}_{\text{col}}$ age. The $^{14}\text{C}_{\text{col}}$ ages are positively correlated with the $\delta^{13}\text{C}_{\text{col}}$
320 values ($R^2 = 0.95$), negatively correlated with the $\delta^{13}\text{C}_{\text{carb}}$ values ($R^2 = 0.68$) but are not
321 correlated with the $\delta^{15}\text{N}_{\text{col}}$ values ($R^2 = 0.24$). The $^{14}\text{C}_{\text{carb}}$ ages are negatively correlated with
322 the $\delta^{13}\text{C}_{\text{carb}}$ values ($R^2 = 0.86$) but are not correlated with the $\delta^{18}\text{O}_{\text{carb}}$ values ($R^2 = 0.01$). The
323 $^{14}\text{C}_{\text{col-carb}}$ age difference is not correlated with the $\delta^{13}\text{C}_{\text{col}}$, $\delta^{13}\text{C}_{\text{carb}}$, or $\delta^{18}\text{O}_{\text{carb}}$ values ($R^2 <$
324 0.42), but is negatively correlated with bone $\delta^{15}\text{N}_{\text{col}}$ values ($R^2 = 0.88$). In terms of ^{14}C

325 activity, Fm values of bone collagen varies between 78.30 ± 0.31 and 80.13 ± 0.26 pMC
326 while bone apatite is about 1% more active on average and varies between 79.45 ± 0.22 and
327 81.02 ± 0.27 pMC (Table 4).

328

329 **6. Discussion**

330 6.1. Diagenetic trajectory

331 The high IRSF values displayed by bone apatite from the SSPM catacomb, ranging
332 between 3.6 and 5.1, are indicative of a moderate to high degree of recrystallization for these
333 samples (Table 2, Figure 6). The presence of a PO_4 peak at 1093 cm^{-1} , and of shoulders at
334 3570 cm^{-1} and 630 cm^{-1} in the most severely altered samples argue for a partial
335 recrystallization in hydroxylapatite. The inverse relationship between mineral crystallinity and
336 carbonate content is typical of dissolution/recrystallization processes that take place following
337 burial (Nielsen-Marsh and Hedges, 2000). Such an alteration of the mineral phase is
338 frequently linked to the loss of the organic matter. This is confirmed by the correlation
339 observed between collagen and carbonate relative contents (Figure 7). This relationship can
340 be explained by the intimate association of the apatite crystals and collagen fibrils at the
341 nanoscale in the bone structure. Mineral dissolution exposes the collagen to microbial or
342 physico-chemical alterations while, in turn, the loss of the organic matter increases micro-
343 porosities and interactions of bone minerals with surrounding water (Collins et al., 2002). The
344 combination of collagen loss, decrease in carbonate content and increase in crystallinity could
345 have been driven by collagen hydrolysis and/or microbial activity (Smith et al., 2007).
346 Bacterial attacks generally result in a loss of the chemical integrity of collagen and a loss of
347 collagen in bone (Balzer et al., 1997; Child, 1995; Grupe, 2001; Harbeck and Grupe, 2009)
348 and begin quickly following burial (Hedges, 2002). Chemical hydrolysis of collagen is
349 another, slower mechanism (Child, 1995; Smith et al., 2002). The lack of data regarding the

350 histological preservation and the porosity of samples does not allow us to distinguish between
351 the two processes. However, archaeological evidence suggests that both could have occurred.
352 The simultaneous or quasi-simultaneous decomposition of the high number of bodies in the
353 mass graves from the SSPM catacomb could have been conducive to microorganism
354 development. Moreover, the burial chambers from the SSPM catacomb were carved in a
355 pozzolan (Funicicillo et al., 2008), which presents an alkaline pH ranging from 7.5 to 7.9
356 (measurements carried out by Zitelli (2013) on samples from ceiling of X82/T18 and X83).
357 These results are consistent with the pH of environmental water recorded in other catacombs
358 in Rome varying between 7.5 and 7.8 (Sánchez-Moral et al., 2005a). However, a much lower
359 pH ranging from 5.9 to 6.4 has been found at the floor level of the X82/T18 and X83
360 chambers (Zitelli 2013). This decrease of the rock pH at the bottom of the stratigraphic
361 sequences could be caused by the decomposition fluids (principally organic acids; Vass et al.
362 (2002)) released from the corpses in putrefaction (Carter and Tibbett, 2008; Gill-King, 1997;
363 Schotsmans et al., 2014). In addition, the gypsum used as part of the funerary practices inside
364 these graves could have also contributed to the acidification of the burial environment. In
365 environmental management, gypsum is frequently incorporated alone or with other
366 amendments in alkaline soils in order to reduce soil pH (Carter, 1986; Courtney and Kirwan,
367 2012; Kordlaghari and Rowell, 2006; Wong et al., 2009). Bone apatite is known to have a low
368 solubility in alkaline environments ($\text{pH} > 7.5$) while its solubility becomes relatively high in
369 acidic environments ($\text{pH} < 6.5$) (Berna et al., 2004; Nielsen-Marsh and Hedges, 2000; Pate et
370 al., 1989). The drop in pH of the burial environment in the chambers could have favored
371 dissolution/recrystallization processes leading to the high IRSF values that we have measured.
372 Finally, the high porosity of the pozzolan (Sánchez-Moral et al., 2005a; Sánchez-Moral et al.,
373 2005b) could have facilitated water drainage. These particular conditions, associated with the
374 absence of sediments between the different strata of corpses in certain burial chambers may

375 have limited the capacity of the burial environment to buffer water and increase interaction of
376 bone with groundwater (Nielsen-Marsh et al., 2007; Smith et al., 2002). These conditions
377 could then have accelerated the recrystallization of bone apatite and the subsequent alteration
378 of the collagen matrix.

379

380 6.2. Spatial heterogeneity of bone preservation

381 Bone preservation in the SSPM catacomb is highly variable. While some of the
382 archaeological samples present IRSF values and CO_3/PO_4 ratios very close to those observed
383 for the modern samples, most of them display higher IRSF values and lower CO_3/PO_4 ratios.
384 Collagen extraction yields are also variable (from 0 to 100%). Time, burial environment
385 (temperature, hydrogeology, soil pH, etc.) and taphonomic processes (inhumation type, filling
386 of the grave, anthropogenic or animal disturbance, etc.) are the three main parameters
387 controlling the state of preservation of bones in archaeological contexts (Collins et al., 2002;
388 Hedges, 2002). These different parameters interplay in a complex manner between and within
389 archaeological sites (Hedges et al., 1995b; Nielsen-Marsh and Hedges, 2000; Pestle and
390 Colvard, 2012). In the SSPM catacomb the organization and taphonomic evolution of the
391 deposition levels suggests massive accumulation of corpses (Castex et al., 2007; Castex et al.,
392 2009; Castex et al., 2011). Despite the fact that contemporary inhumation of 2000-3000
393 corpses *stricto sensu* is unlikely given the dimensions of the burial chambers (Sachau-Carcel,
394 2012; Sachau-Carcel et al., 2013), sepulchral areas appear to have been used during a
395 relatively short period of time (Castex and Blanchard, 2011; Kacki et al., 2013). Therefore,
396 time cannot be invoked to account for the strong heterogeneity of the conservation of the
397 remains and other parameters need to be discussed.

398 The heterogeneity of bone preservation is not randomly distributed. With the
399 exception of the small chamber X81 in which bone samples were well-preserved, most of the

400 well-preserved samples come from the large chambers (X83 and X84) while the poorly-
401 preserved samples are found mostly in the small chambers (X80/T16 and X82/T18). In
402 chamber X78/T15, the limited sampling (n= 2) did not allow us to conclude on the quality of
403 collagen preservation in this room. This result is in keeping with anthropological observations
404 performed during the excavation although local variations can occur depending on the
405 disposition of the bones within a chamber (see below). Like in other great Roman catacombs,
406 the SSPM catacomb has a relatively stable climate throughout the year. Sánchez-Moral et al.
407 (2005a) reported a number of micro-environmental parameters in the catacombs of Domitilla
408 and St. Callistus that can easily be extrapolated to the SSPM catacomb given the similar
409 characteristics and geographical proximity (a few kilometers away) between these sepulchral
410 complexes. A high humidity close to saturation ($\geq 97\%$) and a constant temperature of 15-
411 17°C were recorded in the two catacombs. Given the small size of the central sector of the
412 SSPM catacomb and the existence of large openings between chambers, differences in
413 microclimate between rooms are highly unlikely. During the excavation, visual inspection of
414 bones in the large chambers indicated that bones close to the chamber walls were more altered
415 than in the center (Castex et al., 2009). We propose that runoff against the walls and higher
416 humidity locally could explain this difference in preservation between the center and the
417 periphery of the large chambers. In the smaller chambers, all the bones remain relatively close
418 to the walls and would be equally affected by the water runoff. In the large chambers,
419 buffering conditions (constant humidity and temperature) would have favored exceptional
420 bone collagen preservation for bones situated further away from the walls. Due to its central
421 position, the small chamber X81 has several openings (Figure 1). This has probably restricted
422 runoff against the walls, which could explain the better preservation of the bones in this
423 chamber in general. Finally, trampling of the human remains by the grave diggers, families of
424 the deceased and/or pilgrims along the walls of the burial chambers, especially in the large

425 ones, could be another explanation. A circulation area along the west wall of the chamber
426 X83 has been identified causing the loss of the anatomical integrity of the bone in this sector
427 of the chamber (Castex et al., 2009).

428

429 6.3. Testing for the preservation of the isotopic composition of bone apatite using radiocarbon 430 dating

431 A crucial point for the use of bone apatite isotope values to infer diet and mobility
432 patterns of past populations is to establish whether bone recrystallization was associated with
433 carbon and oxygen isotope exchange. We found apatite to be slightly more ^{14}C active
434 (between 0 and 1.5 pMC) than collagen from the same individual. This is in keeping with
435 differences calculated for bones in archaeological sites ranging in age between 0 and 9.5 kyr
436 BP (Zazzo, in press). The positive $^{14}\text{C}_{\text{col-carb}}$ age difference together with the correlation
437 between $^{14}\text{C}_{\text{carb}}$ and $\delta^{13}\text{C}_{\text{carb}}$ values, indicate that carbon isotope exchange between bone
438 apatite and dissolved inorganic carbon took place with a younger source of carbon with higher
439 $\delta^{13}\text{C}$ values. The $^{14}\text{C}_{\text{col}}$ age offset can be used to calculate a percent contamination of the
440 carbonate fraction with modern carbon if this value represents the true age of an individual.
441 However, the positive correlation found between $\delta^{13}\text{C}_{\text{col}}$ and $^{14}\text{C}_{\text{col}}$ values suggests that the
442 individuals with the highest $\delta^{13}\text{C}$ values could have consumed marine foods as these usually
443 have more positive $\delta^{13}\text{C}$ values than terrestrial foods (Craig et al., 2013; Richards et al., 2005;
444 Schoeninger et al., 1983). The lack of correlation between the $\delta^{15}\text{N}_{\text{col}}$ values and the $^{14}\text{C}_{\text{col}}$
445 ages could be explained by the consumption of low trophic level fish (e.g., anchovies,
446 sardines, mackerel, *etc.*) or bivalves (e.g., oysters, cockles, clams, *etc.*) (Carlier et al., 2007;
447 Garvie-Lok, 2001). This result is coherent with the significant (ca. 15-20%) proportion of
448 marine food found in the diet of the victims of the 79 AD Vesuvius eruption at Herculaneum
449 (Craig et al., 2013). According to the equation in Figure 3A from Craig et al. (2013), $\delta^{13}\text{C}_{\text{col}}$

450 values measured for the five individuals could correspond to ^{14}C age excesses between 60 (for
451 Individual US216/Mand1) and 133 (for Individual 64) ^{14}C yr. This would affect the collagen
452 age more than the apatite age because collagen records the protein fraction of the diet, while
453 apatite carbon records the whole diet, leading to an overestimation of the age differences
454 between the two fractions. A large dependence on marine food in individual 64 could explain
455 the large age difference (180 ± 40 ^{14}C yr) measured between the two individuals dated in the
456 burial chamber X82/T18 (Table 4). This hypothesis is also supported by the fact that in
457 chamber X80/T16, the lack of $^{14}\text{C}_{\text{col}}$ age difference between individuals 24 and 79
458 corresponds to identical $\delta^{13}\text{C}_{\text{col}}$ values for these specimens, suggesting a similar amount of
459 marine food in their diet. For these reasons, the value of 1.5 pMC contamination must be
460 considered as a maximum estimate.

461 The central area of the entire SSPM catacomb and numerous others roman catacombs
462 (e.g. the Domitilla and St Callixtus catacombs) is carved through pozzolan, a tuff layer of the
463 Villa Senni formation resulting from an eruptive event of the Colli Albani volcano
464 (Funicicillo et al., 2008; Giordano, 2010). Bulk measurements on samples coming from the
465 Domitilla and St Callixtus catacombs indicate that pozzolan can contain 1-5% calcite
466 (Sánchez-Moral et al., 2005a; Sánchez-Moral et al., 2005b). To our knowledge, there is no
467 published stable isotope data for this ignimbrite. However, $\delta^{13}\text{C}$ of carbonate in lavas and
468 ejecta from the same volcanic complex have been measured and range from -19.8 to 3.5‰
469 (Fornaseri and Turi, 1969). If we use the highest $\delta^{13}\text{C}$ value as an estimate for the diagenetic
470 endmember, a simple mass balance calculation taking into account the intra-individual
471 differences in ^{14}C activity allows us to estimate a $\delta^{13}\text{C}$ values shift between 0.00‰ (SD
472 ± 0.07 ‰) and 0.27‰ (SD ± 0.08 ‰) as a result of diagenesis. Therefore, the intake of
473 exogenous carbon in bone carbonate is limited and does not modify its carbon isotope
474 composition to any great extent. This is in keeping with the slightly acidic conditions in the

475 SSPM catacomb. As previously mentioned, the correction corresponds to a maximum
476 estimate. Thus, the isotopic composition of the carbonate fraction of bone can be considered
477 as reliable as that of the collagen fraction and can be used to reconstruct the life histories of
478 the individuals buried in the central area of the catacomb. This is beyond the scope of this
479 study and will be the focus of another article.

480 Finally, the significant carbon and oxygen isotope difference of apatite found between
481 samples whose collagen was well- and poorly-preserved is intriguing and deserves attention.
482 Samples which did not pass the collagen preservation tests had $\delta^{13}\text{C}_{\text{carb}}$ values 0.3‰ lower
483 and $\delta^{18}\text{O}_{\text{carb}}$ 0.3‰ higher on average than those with well-preserved collagen (Table 2). The
484 difference between the two groups, although small, was significant for $\delta^{13}\text{C}_{\text{carb}}$ and $\delta^{18}\text{O}_{\text{carb}}$
485 (Mann-Whitney U tests; p-values $\ll 0.05$). Because of the negative correlation found
486 between $^{14}\text{C}_{\text{carb}}$ ages and $\delta^{13}\text{C}_{\text{carb}}$ values, lower $\delta^{13}\text{C}_{\text{carb}}$ are an unlikely result of a contaminant
487 with a more negative $\delta^{13}\text{C}$ value. We propose a different interpretation for the difference
488 found between the two groups. It is striking that the direction and magnitude of the isotopic
489 shifts that we observed are similar to those measured by Koch et al. (1997) between untreated
490 and acetic-acid treated modern enamel and bone samples. The two sub-groups of bones from
491 the SSPM mimic the untreated and treated samples from the Koch et al. (1997) experiment.
492 Moreover, bones with low collagen yields are also highly recrystallized and have lost part of
493 their structural carbonate. Carbonates in apatite are present in two different positions, A and
494 B, A-bonds being the weakest. We found OH⁻ ions in some of the spectra, possibly
495 substituting for A-carbonates. As postulated by Koch et al. (1997) it is plausible to assume
496 that these two sites with different chemical bonds discriminate stable isotopes differently.
497 Since A/B type carbonate ratios are likely different in recrystallized and preserved apatites,
498 the slight isotopic difference measured at the population level could well reflect this

499 difference in isotope fractionation. This difference would then result from recrystallization,
500 not isotope exchange.

501

502 6.4. Regional variability in bone preservation: catacombs vs. open-air sites

503 Rutgers et al. (2009) hypothesized that the microclimate of catacombs affects
504 negatively bone collagen preservation. To further investigate this issue, we compared bone
505 collagen preservation in the catacombs and in contemporary open air sites from Italy (Figure
506 9). The choice of sites included in this comparison is dependent on the collagen extraction
507 protocol used on each site. Some methods, such as ultra-filtration, are known to drastically
508 reduce collagen extraction yields. Because of this possible methodological bias, our
509 comparison only includes sites from central Italy – all located in Rome’s region, also known
510 as Latium - (Killgrove and Tykot, 2013; Prowse et al., 2004; Prowse, 2001; Prowse et al.,
511 2007; Prowse et al., 2005; Rutgers et al., 2009), but excludes sites from southern Italy (Craig
512 et al. 2009; Craig et al. 2013). With an average collagen extraction yield close to 50%, bones
513 from the SSPM appear to be better preserved than most of the contemporary sites of the
514 Latium. Comparison indicates that the quantity of extracted collagen is similar (50% or
515 below) in the small chambers (with the exception of X81) but much higher (between 60-90%
516 on average) in the large chambers. It is also interesting to note that the two sites presenting the
517 highest (although highly variable) average extraction yields in the Latium are catacombs.
518 Despite the limitations described above, the relatively stable climatic conditions prevailing in
519 catacombs seem overall to have favored, rather than disadvantaged bone collagen
520 preservation and therefore contradicts Rutgers et al. (2009)’s assumption. However, an
521 opposite pattern seems to emerge if one looks at the mineral fraction of bone. The comparison
522 of the crystallinity values calculated for the bone samples from the SSPM catacomb with
523 those published for open air sites from different Roman Empire provinces clearly shows that

524 the mineral phase of bones is much more altered in the SSPM catacomb than in the other sites
525 (Figure 10). Although we are aware that the comparison of IRSF values reported in different
526 studies can be problematic due to variances in sample preparation protocols and measurement
527 methods (KBr vs. ATR mode), the IRSF values calculated for the SSPM catacomb are similar
528 to those observed for highly degraded bones (Berna et al., 2004; Lebon et al., 2010). Overall,
529 bones found in catacomb contexts seem to present *a Janus face*¹, showing both a well-
530 preserved organic and a poorly-preserved mineral phase. But as previously shown, this high
531 degree of recrystallization is not correlated with isotope exchange of the mineral phase with
532 the surrounding environment, possibly due to the relatively low pH reigning in the catacomb.
533 Based on this extensive study, we can conclude that it will be possible to examine traits of the
534 life history of the individuals buried in the catacomb based on the combined analysis of bone
535 collagen (when preserved) and apatite isotope values.

536

537 **7. Conclusion**

538 This study - the first focusing on diagenesis in a catacomb context – illustrates the
539 effect of post-depositional processes on bone preservation. Our multi-proxy approach showed
540 that extreme differences in preservation can be found between contemporary individuals
541 within a small area, highlighting the influence of localized variations on taphonomy. We
542 found that the heterogeneity of bone preservation was not randomly distributed in the SSPM
543 catacomb and that bone was better preserved in the large chambers than in the small ones.
544 This is in keeping with anthropological observations performed during the excavation and we
545 propose that the combination of acidic conditions, together with condensation of air humidity
546 and runoff against the walls could explain this differential preservation. Comparison with
547 contemporary sites from the Latium indicates the two sites presenting the highest (although

¹ named after the Roman deity Janus who had two faces, one looking forward, the other looking backward. By extension, this term refers to a deceitful person or an object having or containing contrasting characteristics.

548 highly variable) average extraction yields are catacombs, and that bone collagen in large
549 chambers is much better preserved than in other open air contemporary sites. Therefore, the
550 relatively stable climatic conditions prevailing in catacombs seem overall to have favored,
551 rather than disadvantaged bone preservation.

552 Despite high rates of recrystallization observed for some of the bone samples, coherent
553 radiocarbon ages measured in the carbonate and collagen fractions of several individuals
554 demonstrates a lack of significant (>1%) isotopic exchange. We noticed a slight (0.3‰)
555 difference in both $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ values between the well-preserved and the most severely
556 recrystallized samples. This difference is similar in magnitude and direction with differences
557 observed on modern samples pretreated with acetic acid and could be due to selective
558 leaching of A vs. B-type carbonate in apatite during recrystallization. This shift is close to
559 analytical precision and does not preclude reliable reconstructions of the individual history of
560 these specimens based on stable isotope analysis of the carbonate fraction. Therefore, further
561 dietary reconstructions can include all the individuals of the population, even those for which
562 bone collagen is no longer preserved.

563

564 **8. Acknowledgements**

565 The authors wish to express their gratitude to R. Giuliani, chief inspector of the
566 Vatican's Pontifical Commission for Sacred Archaeology. The study has profited largely from
567 extensive discussions with the members of the scientific team of the SSPM catacomb project,
568 P. Blanchard, G. Sachau-Carcel, S. Kacki and E. Schotsmans. We thank O. Tombret (Labex
569 BCDiv) and J. Ughetto-Monfrin (SSMIM) for their help with the FT-IR and stable isotope
570 analysis. Analyses benefited from the financial help of the Action Thématique du Muséum
571 (ATM) "Biomineralization" and from Archeosphère Inc. (Bordeaux, France) thanks to F.
572 Lacrampe-Cuyaubère and C. Beauval. We are also grateful to the Maison des Sciences de

573 l'Homme d'Aquitaine for their support. K. Salesse benefitted from a scholarship by the
574 Ministère de l'Enseignement Supérieur et de la Recherche allocated to Bordeaux University.
575 Finally, we thank the Guest Editor V. Balter and the two anonymous reviewers for their
576 constructive criticism and their help to clarify several points of this manuscript.

577
578

579 **9. References**

580 Alfano, D., Alburnia, A.R., Motta, O., Proto, A., 2009. Detection of diagenetic alterations by
581 spectroscopic analysis on archaeological bones from the necropolis of Poseidonia
582 (Paestum): a case study. *Journal of Cultural Heritage* 10, 509-513.

583 Ambrose, S.H., 1990. Preparation and characterization of bone and tooth collagen for isotopic
584 analysis. *Journal of Archaeological Science* 17, 431-451.

585 Asscher, Y., Weiner, S., Boaretto, E., 2011. Variations in atomic disorder in biogenic
586 carbonate hydroxyapatite using the infrared spectrum grinding curve method. *Advanced*
587 *Functional Materials* 21, 3308-3313.

588 Balzer, A., Gleixner, G., Grupe, G., Schmidt, H.L., Schramm, S., Turban-Just, S., 1997. In
589 vitro decomposition of bone collagen by soil bacteria: the implications for stable isotope
590 analysis in archaeometry. *Archaeometry* 39, 415-429.

591 Berna, F., Matthews, A., Weiner, S., 2004. Solubilities of bone mineral from archaeological
592 sites: the recrystallization window. *Journal of Archaeological Science* 31, 867-882.

593 Blanchard, P., Castex, D., Coquerelle, M., Giuliani, R., Ricciardi, M., 2007. A mass grave
594 from the catacomb of Saints Peter and Marcellinus in Rome, second-third century AD.
595 *Antiquity* 81, 989-998.

596 Bocherens, H., Drucker, D., Billiou, D., Moussa, I., 2005. Une nouvelle approche pour
597 évaluer l'état de conservation de l'os et du collagène pour les mesures isotopiques

598 (datation au radiocarbone, isotopes stables du carbone et de l'azote). *L'Anthropologie*
599 109, 557-567.

600 Bocherens, H., Drucker, D.G., Billiou, D., Geneste, J.M., Kervazo, B., 2008. Grotte Chauvet
601 (Ardèche, France): a "natural experiment" for bone diagenesis in karstic context.
602 *Palaeogeography, Palaeoclimatology, Palaeoecology* 266, 220-226.

603 Bocherens, H., Fizet, M., Cuif, J.P., Jaeger, J.-J., Michard, J.-G., Mariotti, A., 1988.
604 Premières mesures d'abondances isotopiques naturelles en ^{13}C et en ^{15}N de la matière
605 organique fossile de dinosaure. Application à l'étude du régime alimentaire du genre
606 *Anatosaurus* (*Ornithischia*, *Hadrosauridae*). *Comptes Rendus de l'Académie des*
607 *Sciences* 306, 1521-1526.

608 Bocherens, H., Fizet, M., Mariotti, A., Lange-Badre, B., Vandermeersch, B., Borel, J.P.,
609 Bellon, G., 1991. Isotopic biogeochemistry (^{13}C , ^{15}N) of fossil vertebrate collagen:
610 application to the study of a past food web including Neandertal man. *Journal of Human*
611 *Evolution* 20, 481-492.

612 Brady, A.L., White, C.D., Longstaffe, F.J., Southam, G., 2008. Investigating intra-bone
613 isotopic variations in bioapatite using IR-laser ablation and micromilling: implications
614 for identifying diagenesis? *Palaeogeography, Palaeoclimatology, Palaeoecology* 266,
615 190-199.

616 Carlier, A., Riera, P., Amouroux, J.-M., Bodiou, J.-Y., Grémare, A., 2007. Benthic trophic
617 network in the Bay of Banyuls-sur-Mer (northwest Mediterranean, France): an
618 assessment based on stable carbon and nitrogen isotopes analysis. *Estuarine, Coastal*
619 *and Shelf Science* 72, 1-15.

620 Carter, D.O., Tibbett, M., 2008. Cadaver decomposition and soil: processes, in: Tibbett, M.,
621 Carter, D.O. (Eds.), *Soil analysis in forensic taphonomy*. CRC Press, Boca Raton, pp.
622 29-52.

623 Carter, M.R., 1986. Microbial biomass and mineralizable nitrogen in solonchic soils:
624 influence of gypsum and lime amendments. *Soil Biology and Biochemistry* 18, 531-
625 537.

626 Castex, D., Blanchard, A., 2011. Témoignages archéologiques de crise(s) épidémique(s) : la
627 catacombe des Saints Marcellin et Pierre (Rome, fin I^{er}-III^e s.), in: Castex, D., Courtaud,
628 P., Duday, H., Le Mort, F., Tillier, A.-M. (Eds.), *Le regroupement des morts. Genèse et*
629 *diversité archéologique*. Maison des Sciences de l'Homme d'Aquitaine, Bordeaux, pp.
630 281-293.

631 Castex, D., Blanchard, A., Giuliani, R., Ricciardi, M., 2007. Les ensembles funéraires du
632 secteur central de la catacombe des Saints Pierre et Marcellin (Rome, I^{er}-III^e siècle) :
633 caractérisation, hypothèses d'interprétations et perspectives de recherches. *Mélanges de*
634 *l'École française de Rome - Antiquité* 119, 274-282.

635 Castex, D., Blanchard, A., Réveillias, H., Kacki, S., Giuliani, R., 2009. Les sépultures du
636 secteur central de la catacombe des Saints Pierre-et-Marcellin (Rome). État des analyses
637 bio-archéologiques et perspectives *Mélanges de l'École française de Rome - Antiquité*
638 121, 287-297.

639 Castex, D., Blanchard, P., Kacki, S., Réveillias, H., Giuliani, R., 2011. Le secteur central de la
640 catacombe des Sts Pierre-et-Marcellin (Rome, I^{er}-III^e siècle). *Indices archéologiques*
641 *d'une crise brutale de mortalité*. *Mélanges de l'École Française de Rome – Chronique*
642 123, 274-280.

643 Child, A.M., 1995. Towards an understanding of the microbial decomposition of
644 archaeological bone in the burial environment. *Journal of Archaeological Science* 22,
645 165-174.

646 Collins, M.J., Nielsen-Marsh, C.M., Hiller, J., Smith, C.I., Roberts, J.P., Prigodich, R.V.,
647 Weiss, T.J., Csapo, J., Millard, A.R., Turner-Walker, G., 2002. The survival of organic
648 matter in bone: a review. *Archaeometry* 44, 383-394.

649 Courtney, R., Kirwan, L., 2012. Gypsum amendment of alkaline bauxite residue – Plant
650 available aluminium and implications for grassland restoration. *Ecological Engineering*
651 42, 279-282.

652 Craig, O.E., Biazzo, M., O'Connell, T.C., Garnsey, P., Martinez-Labarga, C., Lelli, R.,
653 Salvadei, L., Tartaglia, G., Nava, A., Renò, L., Fiammenghi, A., Rickards, O., Bondioli,
654 L., 2009. Stable isotopic evidence for diet at the imperial Roman coastal site of Velia
655 (1st and 2nd Centuries AD) in southern Italy. *American Journal of Physical*
656 *Anthropology* 139, 572-583.

657 Craig, O.E., Bondioli, L., Fattore, L., Higham, T., Hedges, R., 2013. Evaluating marine diets
658 through radiocarbon dating and stable isotope analysis of victims of the AD79 eruption
659 of Vesuvius. *American Journal of Physical Anthropology* 152, 345-352.

660 DeNiro, M.J., 1985. Postmortem preservation and alteration of in vivo bone collagen ratios in
661 relation to palaeodietary reconstruction. *Nature* 317, 806-809.

662 DeNiro, M.J., Epstein, S., 1981. Influence of diet on the distribution of nitrogen isotopes in
663 animals. *Geochimica et Cosmochimica Acta* 45, 341-351.

664 DeNiro, M.J., Weiner, S., 1988. Chemical, enzymatic and spectroscopic characterization of
665 “collagen” and other organic fractions from prehistoric bones. *Geochimica et*
666 *Cosmochimica Acta* 52, 2197-2206.

667 Devières, T., Vanhove, C., Chapoulie, R., Blanchard, P., Colombini, M.P., Regert, M., Castex,
668 D., 2010. Détermination et fonction des substances organiques et des matières minérales
669 exploitées dans les rites funéraires de la catacombe des Saints Pierre-et-Marcellin à
670 Rome (Ier-IIIe siècle), in: Cartron, I., Castex, D., Georges, P., Vivas, M., Charageat, M.

671 (Eds.), *De corps en corps. Traitement et devenir du cadavre*. Maison des Sciences de
672 l'Homme d'Aquitaine, Bordeaux, pp. 115-139.

673 Dobberstein, R.C., Collins, M.J., Craig, O.E., Taylor, G., Penkman, K.E.H., Ritz-Timme, S.,
674 2009. Archaeological collagen: why worry about collagen diagenesis? *Archaeological*
675 *and Anthropological Sciences* 1, 31-42.

676 Dupras, T.L., 1999. *Dining in the Dakhleh Oasis, Egypt: determination of diet using*
677 *documents and stable isotope analysis*, Department of Anthropology. McMaster
678 University, Hamilton, p. 345.

679 Dupras, T.L., Schwarcz, H.P., 2001. Strangers in a strange land: stable isotope evidence for
680 human migration in the Dakhleh Oasis, Egypt. *Journal of Anthropological Archaeology*
681 28, 1199-1208.

682 Fornaseri, M., Turi, B., 1969. Carbon and oxygen isotopic composition of carbonates in lavas
683 and ejectites from the Alban Hills, Italy. *Contributions to Mineralogy and Petrology* 23,
684 244-256.

685 Funicicillo, R., Praturlon, A., Giordano, G., 2008. *La geologia di Roma dal centro storico alla*
686 *periferia*. S.E.L.C.A., Firenze, p. 660.

687 Garvie-Lok, S.J., 2001. *Loaves and fishes: a stable isotope reconstruction of diet in medieval*
688 *Greece*, Department of Archaeology. University of Calgary, Calgary, p. 525.

689 Gill-King, H., 1997. Chemical, ultrastructural aspects of decomposition, in: Haglund, W.D.,
690 Sorg, M.H. (Eds.), *Forensic taphonomy: the post-mortem fate of human remains*. CRC
691 Press, Boca Raton, pp. 93-108.

692 Giordano, A., 2010. Stratigraphy, volcano tectonics and evolution of the Colli Albani
693 volcanic field, in: Funicicillo, R., Giordani, A. (Eds.), *The Colli Albani volcano*.
694 Geological Society, London, pp. 43-98.

695 Grupe, G., 2001. Archaeological microbiology, in: Brothwell, D.R., Pollard, A.M. (Eds.),
696 Handbook of archaeological sciences. Wiley, New York, pp. 351-363.

697 Grupe, G., Balzer, A., Turban-Just, S., 2001. Modeling protein diagenesis in ancient bone:
698 towards a validation of stable isotope data, in: Ambrose, S.H., Katzenberg, M.A. (Eds.),
699 Biogeochemical approaches to paleodietary analysis. Kluwer Academic, Plenum
700 Publishers, New York, pp. 173-188.

701 Guyon, J., 1987. Le cimetière aux deux lauriers. Recherches sur les catacombes romaines.
702 École française de Rome, Rome, p. 556.

703 Harbeck, M., Grupe, G., 2009. Experimental chemical degradation compared to natural
704 diagenetic alteration of collagen: implications for collagen quality indicators for stable
705 isotope analysis. *Archaeological and Anthropological Sciences* 1, 43-57.

706 Hare, P.E., Estep, M.L.F., 1982. Carbon and nitrogen isotopic composition of amino acids in
707 modern and fossil collagens. *Carnegie Institute of Washington Yearbook* 82, 410-414.

708 Hare, P.E., Fogel, M.L., Stafford Jr, T.W., Mitchell, A.D., Hoering, T.C., 1991. The isotopic
709 composition of carbon and nitrogen in individual amino acids isolated from modern and
710 fossil proteins. *Journal of Archaeological Science* 18, 277-292.

711 Hedges, R.E.M., 2002. Bone diagenesis: an overview of processes. *Archaeometry* 44, 319-
712 328.

713 Hedges, R.E.M., Lee-Thorp, J.A., Tuross, N.C., 1995a. Is tooth enamel carbonate a suitable
714 material for radiocarbon dating? *Radiocarbon* 37, 285-290.

715 Hedges, R.E.M., Millard, A.R., 1995. Bones and groundwater: towards the modelling of
716 diagenetic processes. *Journal of Archaeological Science* 22, 155-164.

717 Hedges, R.E.M., Millard, A.R., Pike, A.W.G., 1995b. Measurements and relationships of
718 diagenetic alteration of bone from three archaeological sites. *Journal of Archaeological*
719 *Science* 22, 201-209.

720 Iacumin, P., Bocherens, H., Chaix, L., Mariotti, A., 1998. Stable carbon and nitrogen isotopes
721 as dietary indicators of ancient Nubian populations (northern Sudan). *Journal of*
722 *Anthropological Archaeology* 25, 293-301.

723 Kacki, S., Réveillas, H., Sachau-Carcel, G., Giuliani, R., Blanchard, P., Castex, D., 2013.
724 Réévaluation des arguments de simultanéité des dépôts de cadavres: l'exemple des
725 sépultures plurielles de la catacombe des Saints Pierre-et-Marcellin (Rome). *Bulletins et*
726 *Mémoires de la Société d'Anthropologie de Paris*, 1-10.

727 Katzenberg, M.A., Goriunova, O., Weber, A., 2009. Paleodiet reconstruction of Bronze age
728 Siberians from the mortuary site of Khuzhir-Nuge XIV, Lake Baikal. *Journal of*
729 *Archaeological Science* 36, 663–674.

730 Keenleyside, A., Schwarcz, H.P., Stirling, L., Ben Lazreg, N., 2009. Stable isotopic evidence
731 for diet in a Roman and late Roman population from Leptiminus, Tunisia. *Journal of*
732 *Archaeological Science* 36, 51-63.

733 Killgrove, K., 2010. Migration and mobility in imperial Rome, Department of Anthropology.
734 University of North Carolina, Chapel Hill, p. 390.

735 Killgrove, K., Tykot, R.H., 2013. Food for Rome: a stable isotope investigation of diet in the
736 Imperial period (1st–3rd centuries AD). *Journal of Anthropological Archaeology* 32, 28-
737 38.

738 Koch, P.L., Tuross, N., Fogel, M.L., 1997. The effects of sample treatment and diagenesis on
739 the isotopic integrity of carbonate in biogenic hydroxylapatite. *Journal of*
740 *Archaeological Science* 24, 417-429.

741 Kordlaghari, M.P., Rowell, D.L., 2006. The role of gypsum in the reactions of phosphate with
742 soils. *Geoderma* 132, 105-115.

743 Krueger, H.W., 1991. Exchange of carbon with biological apatite. *Journal of Archaeological*
744 *Science* 18, 355-361.

745 Lebon, M., Muller, K., Bahain, J.-J., Frohlich, F., Falgueres, C., Bertrand, L., Sandt, C.,
746 Reiche, I., 2011. Imaging fossil bone alterations at the microscale by SR-FTIR
747 microspectroscopy. *Journal of Analytical Atomic Spectrometry* 26, 922-929.

748 Lebon, M., 2008. Caractérisation par spectroscopie infrarouge à transformée de Fourier des
749 ossements chauffés en contexte archéologique - Comparaison entre référentiel moderne
750 et matériel fossile. Implications diagénétiques, Spécialité : Préhistoire. Muséum
751 National d'Histoire Naturelle, Paris, p. 334.

752 Lebon, M., Reiche, I., Bahain, J.J., Chadeaux, C., Moigne, A.M., Fröhlich, F., Sémah, F.,
753 Schwarcz, H.P., Falguères, C., 2010. New parameters for the characterization of
754 diagenetic alterations and heat-induced changes of fossil bone mineral using Fourier
755 transform infrared spectrometry. *Journal of Archaeological Science* 37, 2265-2276.

756 Longin, R., 1971. New method of collagen extraction for radiocarbon dating. *Nature* 230,
757 241-242.

758 Munro, L.E., Longstaffe, F.J., White, C.D., 2007. Burning and boiling of modern deer bone:
759 effects on crystallinity and oxygen isotope composition of bioapatite phosphate.
760 *Palaeogeography, Palaeoclimatology, Palaeoecology* 249, 90-102.

761 Nielsen-Marsh, C.M., Hedges, R.E.M., 2000. Patterns of Diagenesis in Bone I: the Effects of
762 Site Environments. *Journal of Archaeological Science* 27, 1139-1150.

763 Nielsen-Marsh, C.M., Smith, C.I., Jans, M.M.E., Nord, A., Kars, H., Collins, M.J., 2007.
764 Bone diagenesis in the European Holocene II: taphonomic and environmental
765 considerations. *Journal of Archaeological Science* 34, 1523-1531.

766 Oelze, V.M., Koch, J.K., Kupke, K., Nehlich, O., Zauner, S., Wahl, J., Weise, S.M.,
767 Rieckhoff, S., Richards, M.P., 2012. Multi-isotopic analysis reveals individual mobility
768 and diet at the early Iron age monumental tumulus of Magdalenenberg, Germany.
769 *American Journal of Physical Anthropology* 148, 406-421.

770 Pagni, G., Burdassi, L., 2004. Gli inumati nella catacomba dei SS Pietro e Marcellino.
771 Intervento di scavo antropologico. Excavation report, p. 10.

772 Pate, F.D., Hutton, J.T., Norrish, K., 1989. Ionic exchange between soil solution and bone:
773 toward a predictive model. *Applied Geochemistry* 4, 303-316.

774 Pestle, W.J., Colvard, M., 2012. Bone collagen preservation in the tropics: a case study from
775 ancient Puerto Rico. *Journal of Archaeological Science* 39, 2079-2090.

776 Prowse, T., Schwarcz, H.P., Saunders, S., Macchiarelli, R., Bondioli, L., 2004. Isotopic
777 paleodiet studies of skeletons from the imperial Roman-age cemetery of Isola Sacra,
778 Rome, Italy. *Journal of Archaeological Science* 31, 259-272.

779 Prowse, T.L., 2001. Isotopic and dental evidence for diet from the necropolis of Isola Sacra
780 (1st-3rd centuries AD), Italy, Department of Anthropology. McMaster University,
781 Hamilton, p. 356.

782 Prowse, T.L., Schwarcz, H.P., Garnsey, P., Knyf, M., Macchiarelli, R., Bondioli, L., 2007.
783 Isotopic evidence for age-related immigration to imperial Rome. *American Journal of*
784 *Physical Anthropology* 132, 510-519.

785 Prowse, T.L., Schwarcz, H.P., Saunders, S., Macchiarelli, R., Bondioli, L., 2005. Isotopic
786 evidence for age-related variation in diet from Isola Sacra, Italy. *American Journal of*
787 *Physical Anthropology* 128, 2-13.

788 Reitsema, L.J., 2012. Stable carbon and nitrogen isotope analysis of human diet change in
789 prehistoric and historic Poland, Department of Anthropology. The Ohio State
790 University, Columbus, p. 377.

791 Richards, M.P., Jacobi, R., Cook, J., Pettitt, P.B., Stringer, C.B., 2005. Isotope evidence for
792 the intensive use of marine foods by late Upper Palaeolithic humans. *Journal of Human*
793 *Evolution* 49, 390-394.

794 Rutgers, L.V., van Strydonck, M., Boudin, M., van der Linde, C., 2009. Stable isotope data
795 from the early Christian catacombs of ancient Rome: new insights into the dietary habits
796 of Rome's early Christians. *Journal of Archaeological Science* 36, 1127–1134.

797 Sachau-Carcel, G., 2012. Apport de la modélisation tridimensionnelle à la compréhension du
798 fonctionnement des sépultures multiples. L'exemple du secteur central de la catacombe
799 des Saints Pierre-et-Marcellin (Rome, Italie). (Ier-milieu IIIe s. ap. J.-C.), Spécialité :
800 Histoire ancienne. Université Bordeaux 3, Bordeaux, p. 505.

801 Sachau-Carcel, G., Castex, D., Vergnieux, R., 2013. Sites à stratification complexe et
802 modélisation tridimensionnelle : vers une nouvelle approche des sépultures multiples.
803 *ArchéoSciences* 37, 89-104.

804 Sánchez-Moral, S., Luque, L., Cañaveras, J.-C., Soler, V., Garcia-Guinea, J., Aparicio, A.,
805 2005a. Lime–pozzolana mortars in Roman catacombs: composition, structures and
806 restoration. *Cement and Concrete Research* 35, 1555-1565.

807 Sánchez-Moral, S., Luque, L., Cuezva, S., Soler, V., Benavente, D., Laiz, L., Gonzalez, J.M.,
808 Saiz-Jimenez, C., 2005b. Deterioration of building materials in Roman catacombs: the
809 influence of visitors. *Science of the Total Environment* 349, 260-276.

810 Schoeninger, M.J., DeNiro, M.J., Tauber, H., 1983. Stable nitrogen isotope ratios of bone
811 collagen reflect marine and terrestrial components of Prehistoric human diet. *Science*
812 220, 1381-1383.

813 Schotsmans, E.M.J., Fletcher, J.N., Denton, J., Janaway, R.C., Wilson, A.S., 2014. Long-term
814 effects of hydrated lime and quicklime on the decay of human remains using pig
815 cadavers as human body analogues: field experiments. *Forensic Science International*
816 238, 141.e1–141.e13.

817 Shemesh, A., 1990. Crystallinity and diagenesis of sedimentary apatites. *Geochimica et*
818 *Cosmochimica Acta* 54, 2433-2438.

819 Shin, J.Y., Hedges, R.E.M., 2012. Diagenesis in bone and enamel apatite carbonate; the
820 potential of density separation to assess the original composition. *Journal of*
821 *Archaeological Science* 39, 1123-1130.

822 Smith, C.I., Nielsen-Marsh, C.M., Jans, M.M.E., Collins, M.J., 2007. Bone diagenesis in the
823 European Holocene : patterns and mechanisms. *Journal of Archaeological Science* 34,
824 1485-1493.

825 Smith, C.I., Nielsen-Marsh, C.M., Jans, M.M.E., Arthur, P., Nord, A.G., Collins, M.J., 2002.
826 The strange case of Apigliano: early ‘fossilization’ of medieval bone in southern Italy.
827 *Archaeometry* 44, 405-415.

828 Stafford, J.T.W., Hare, P.E., Currie, L., Jull, A.J.T., Donahue, D.J., 1991. Accelerator
829 radiocarbon dating at the molecular level. *Journal of Archaeological Science* 18, 35-72.

830 Stiner, M.C., Kuhn, S.L., Weiner, S., Bar-Yosef, O., 1995. Differential burning,
831 recrystallization, and fragmentation of archaeological bone. *Journal of Archaeological*
832 *Science* 22, 223-237.

833 Trueman, C.N.G., Behrensmeyer A.K., Tuross, N., Weiner, S., 2004. Mineralogical and
834 compositional changes in bones exposed on soil surfaces in Amboseli National Park,
835 Kenya: diagenetic mechanisms and the role of sediment pore fluids. *Journal of*
836 *Archaeological Science* 31, 721-739.

837 Trueman, C.N.G., Privat, K., Field, J., 2008. Why do crystallinity values fail to predict the
838 extent of diagenetic alteration of bone mineral? *Palaeogeography, Palaeoclimatology,*
839 *Palaeoecology* 266, 160-167.

840 van Klinken, G.J., 1999. Bone collagen quality indicators for palaeodietary and radiocarbon
841 measurements. *Journal of Archaeological Science* 26, 687-695.

842 Vanhove, C., 2006. Pratiques et rites funéraires du monde romain méditerranéen aux II^{ème} et
843 III^{ème} siècles de notre ère. Caractérisation du matériel de la catacombe des Saints Pierre

844 et Marcellin à Rome. Master's thesis directed by R. Chapoulie (IRAMAT-CRP2A).
845 Université Bordeaux Montaigne, Bordeaux, p. 36.

846 Vass, A.A., Barshick, S.-A., Segal, G., Caton, J., Skeen, J.T., Love, J.C., Synstelien, J.A.,
847 2002. Decomposition chemistry of human remains: a new methodology for determining
848 the postmortem interval. *Journal of Forensic Sciences* 47, 542-553.

849 Weiner, S., Bar-Yosef, O., 1990. States of preservation of bones from prehistoric sites in the
850 Near East: a survey. *Journal of Archaeological Science* 17, 187-196.

851 Wong, V.N.L., Dalal, R.C., Greene, R.S.B., 2009. Carbon dynamics of sodic and saline soils
852 following gypsum and organic material additions: a laboratory incubation. *Applied Soil
853 Ecology* 41, 29-40.

854 Wright, L.E., Schwarcz, H.P., 1996. Infrared and isotopic evidence for diagenesis of bone
855 apatite at Dos Pilas, Guatemala: palaeodietary implications. *Journal of Archaeological
856 Science* 23, 933-944.

857 Wright, L.E., Valdés, J.A., Burton, J.H., Douglas Price, T., Schwarcz, H.P., 2010. The
858 children of Kaminaljuyu: isotopic insight into diet and long distance interaction in
859 Mesoamerica. *Journal of Anthropological Archaeology* 29, 155-178.

860 Zazzo, A., in press. Bone and enamel carbonate diagenesis: a radiocarbon prospective.
861 *Palaeogeography, Palaeoclimatology, Palaeoecology*, DOI:
862 10.1016/j.palaeo.2014.05.006

863 Zazzo, A., Saliège, J.F., 2011. Radiocarbon dating of biological apatites: a review.
864 *Palaeogeography, Palaeoclimatology, Palaeoecology* 310, 52-61.

865 Zitelli, F., 2013. Étude de la cinétique de dépôt de corps par analyses physico-chimiques des
866 parois de tombes. Cas de la catacombe des Saints Pierre et Marcellin (Rome Ier-IIIème
867 s. ap. J.-C.). ENSCBP internship report directed by R. Chapoulie (IRAMAT-CRP2A).
868 Université Bordeaux Montaigne, Bordeaux, p. 30.

869

870 **Figure Captions**

871

872 **Figure 1:** Plan of the burial chambers in the region X from the SSPM catacomb. After M.

873 Ricciardi, modified.

874

875 **Figure 2:** FT-IR spectrum of a fresh bone showing the regions of interest as well as the IRSF

876 calculation procedure.

877

878 **Figure 3:** Photographs showing examples of good (A), medium (B) and poor (C) preservation

879 of the SSPM bones, based on their macroscopic characteristics

880

881 **Figure 4:** Collagen yield of the SSPM bone samples grouped by the burial chamber, skeletal

882 part category and macroscopic appearance. Boxplots represent the 25th, 50th (median) and 75th

883 percentiles in the distribution of scores. Whiskers indicate the variability outside the upper

884 and lower quartiles. Outliers are identified by individual points. The shaded area corresponds

885 to the collagen yield measured in modern bones (after Bocherens et al. 1991).

886

887 **Figure 5:** Carbon and nitrogen abundances of the SSPM bone samples. Samples which fail

888 one of the collagen preservation criteria defined in Table 1 are considered poorly-preserved

889 (closed symbols) while those passing all the preservation criteria are considered well-

890 preserved (open symbols).

891

892 **Figure 6:** IRSF plotted against CO₃/PO₄ ratio of the SSPM bone samples. Modern bone

893 samples are shown for reference.

894

895 **Figure 7:** CO₃/PO₄ ratio plotted against Amide1/ PO₄ ratio of the SSPM bone samples.

896 Modern bone samples are shown for reference.

897

898 **Figure 8:** IRSF plotted against Amide1/ PO₄ ratio of the SSPM bone samples. Modern bone
899 samples are shown for reference.

900

901 **Figure 9:** Inter-site comparison of the collagen extraction yields measured on contemporary
902 human bone samples from the Latium. The Isola Sacra necropolis is located near Ostia and
903 Portus Romae and dated from the 1st to 3rd c. AD (Prowse et al., 2004; Prowse, 2001; Prowse
904 et al., 2007; Prowse et al., 2005). The ANAS cemetery is located between Ostia and Rome
905 and dated from the 1st to 3rd c. AD (Prowse et al., 2004; Prowse, 2001). The necropolises of
906 Casal Bertone and Castellaccio Europarco are located in Rome and dated from the 2nd to 3rd
907 and from the 1st to 2nd c. AD, respectively (Killgrove, 2010; Killgrove and Tykot, 2013). The
908 St Callixtus catacomb is located in Rome and dated from the 3rd to 5th c. AD (Rutgers et al.,
909 2009). Boxplots represent the 25th, 50th (median) and 75th percentiles in the distribution of
910 scores. Whiskers indicate the variability outside the upper and lower quartiles. Outliers are
911 identified by individual points. The shaded area corresponds to the collagen yield measured in
912 modern bones (after Bocherens et al. 1991). OAS stands for open air site and SS stands for
913 subterranean site.

914

915 **Figure 10:** Inter-site comparison of the IRSF values measured on human bone apatite
916 samples from the Roman Empire. The Kellis 2 cemetery is located near Mut in the Dakhleh
917 Oasis in Egypt and dated from the 1st to 5th c. AD (Dupras, 1999; Dupras and Schwarcz,
918 2001). The Leptiminus cemeteries are located in the modern day town of Lamta in Tunisia
919 and dated from the 2nd to 6th century AD (Keenleyside et al. 2009). The Poseidonia

920 necropolises are located near Capaccio-Paestum in Italy and dated from the 7th c. BC to the
921 2nd c. AD (Alfano et al. 2009). The Isola Sacra necropolis is located near Ostia and Portus
922 Romae and dated from the 1st to 3rd c. AD (Prowse et al., 2004; Prowse, 2001; Prowse et al.,
923 2007; Prowse et al., 2005). Boxplots represent the 25th, 50th (median) and 75th percentiles in
924 the distribution of scores. Whiskers indicate the variability outside the upper and lower
925 quartiles. Outliers are identified by individual points. The hatched area corresponds to the
926 IRSF values found for modern bones. The shaded area corresponds to the samples mildly and
927 highly recrystallized. OAS stands for open air site and SS for subterranean site.
928