
Local Identifiability of Deep ReLU Neural Networks: the Theory

Joachim Bona-Pellissier^{ab*}, François Malgouyres^b, François Bachoc^b

Institut de Mathématiques de Toulouse ; UMR 5219

Université de Toulouse ; CNRS

^a UT1, F-31042 Toulouse, France

^b UPS, F-31062 Toulouse Cedex 9, France

*Corresponding author: Joachim Bona-Pellissier; joachim.bona-pellissier@univ-toulouse.fr

Abstract

Is a sample rich enough to determine, at least locally, the parameters of a neural network? To answer this question, we introduce a new local parameterization of a given deep ReLU neural network by fixing the values of some of its weights. This allows us to define local lifting operators whose inverses are charts of a smooth manifold of a high dimensional space. The function implemented by the deep ReLU neural network composes the local lifting with a linear operator which depends on the sample. We derive from this convenient representation a geometric necessary and sufficient condition of local identifiability. Looking at tangent spaces, the geometric condition provides: 1/ a sharp and testable necessary condition of identifiability and 2/ a sharp and testable sufficient condition of local identifiability. The validity of the conditions can be tested numerically using backpropagation and matrix rank computations.

1 Introduction

1.1 Context and motivations

Neural networks are famous for their capacity to perform complex tasks in a wide variety of domains such as image classification [20], object recognition [33, 34], speech recognition [17, 36, 16], natural language processing [27, 26, 19], anomaly detection [32] or climate sciences [1].

The following properties of the parameters of neural networks have recently drawn attention: identifiability, inverse stability and stable recovery; from weaker to stronger. Let $f_\theta(X)$ be the outputs of a network parameterized by the parameters θ , for given inputs X . Global identifiability means that if $f_\theta(X) = f_{\tilde{\theta}}(X)$ then $\theta = \tilde{\theta}$, up to identified invariances, for instance neuron permutation and rescaling for ReLU networks. Local identifiability restricts this analysis for θ and $\tilde{\theta}$ sufficiently close. Then, inverse stability means that the distance between θ and $\tilde{\theta}$ (up to invariances) is bounded by a function of the distance between $f_\theta(X)$ and $f_{\tilde{\theta}}(X)$. Finally, stable recovery consists in obtaining an algorithm to approximately recover θ from a noisy version of $f_\theta(X)$, with quantitative guarantees. In all cases, we must distinguish between statements for X being a finite list of inputs, in which case we would like X to be small, and for infinite X (for instance determining θ from the entire function f_θ).

Identifiability from finite X , which is the focus of this paper, is important for different reasons. In the first place, model extraction attacks for neural networks have been a growing topic over the last years. Indeed, some algorithms are able to recover in practice the parameters of a neural network

from queries [9, 35]. This can be a concern since neural network providers may wish to keep these parameters secret, for security [21], for privacy [13, 8], or for intellectual property [42].

A way of preventing such a recovery can be by guaranteeing that identifiability does not hold, that is to check that a necessary condition of identifiability is not met. On the opposite side, guaranteeing that identifiability holds is interesting in the position of an attacker. If the attacker has access to X , to $f_\theta(X)$, and is able to compute a $\tilde{\theta}$ such that $f_{\tilde{\theta}}(X) = f_\theta(X)$, the question then becomes: does this guarantee that $\tilde{\theta} = \theta$ or shall the attacker expand X with new queries? The attacker needs a sufficient condition of identifiability.

Another important motivation for identifiability is having a better understanding and control of neural networks. Indeed, if the learning sample has the form $(X, f_\theta(X))$, with θ the parameters of a teaching network, global identifiability from X means that the global minimizer of the empirical risk is unique. In this case, if the global minimizer is reached, there will typically be no variability due to the optimization parameters (choice of the algorithm, number of epochs,...) and to stochasticity (for stochastic optimizers). Even if very recent works on double descent phenomena, e.g. [4], highlight a benefit of overparameterization (thus absence of identifiability) for increasing prediction performances, a user may be interested in a small enough number of parameters to retain identifiability, if the loss of performance is mild compared to overparameterization.

Note that, of course, global identifiability is more relevant than local identifiability to the above motivations. This work nevertheless focuses on local identifiability, which is a necessary condition for global identifiability, and which analysis can be a first step to analyzing global identifiability. Local identifiability is also arguably insightful on the geometry of the relationship between the parameter space of θ and its image $\{f_\theta(X), \theta \text{ varies}\}$. Note that most existing identifiability, inverse stability and stable recovery results (see the next section) are also local.

1.2 Existing work on identifiability, inverse stability, stable recovery and attacks

Identifiability: Even though it has regained interest recently, the question of identifiability for neural networks is not new. Indeed, in the 1990s, some positive results of identifiability for networks with smooth activation functions (tanh, logistic sigmoid or Gaussian for instance) have been established [40, 2, 22, 18, 12]. These results are mainly theoretical, they concern activation functions which are not the most used nowadays (in particular, they do not apply to ReLU networks), and assume full knowledge of the function f_θ implemented by the network, which is impossible in practice.

When it comes to ReLU, for shallow [30, 38] as well as deep [31, 5] neural networks, some positive results of identifiability have been recently established. They show that under some conditions on the architecture and parameters of the network, the function implemented by the network uniquely characterizes its parameters, up to neuron permutation and rescaling operations. Although they apply to ReLU networks, these results share a limitation with those of previous paragraph: they assume the function implemented by the network to be known on the whole input space, or at least on an open subset of it.

As far as we know, there exists only one identifiability result for deep ReLU networks assuming the knowledge of f_θ on a *finite* sample only. Stock and Gribonval [39] give a theoretical condition for the existence of a finite set which locally identifies the parameters of a deep neural network. It is an *existence* result: it does not concretely provide such a finite set, nor does it allow to test local identifiability for any finite sample, as we propose in this work. The construction in [39] shares similarities with previous works on deep structured matrix factorization [23, 24, 25]. The present article also lies in this line of research.

Inverse stability and stable recovery: Closely related to identifiability are the topics of inverse stability and stable recovery of the parameters of a network. Some negative [29] as well as positive [11, 23, 24, 25] results of inverse stability exist. The articles [23, 24, 25] examine the case of structured networks with the identity as activation function. Only [25] considers a finite X . The authors of [11] consider a general class of networks amongst which ReLU networks, but the result only holds for one-hidden-layer neural networks. Furthermore this result also requires the knowledge of f_θ on a whole domain.

Several stable recovery algorithms have also been proposed, for one-hidden-layer neural networks in a first place, for smooth activation function [14], as well as ReLU in the fully-connected case




Figure 1: The local intersection between the affine space $N(X, \theta)$ (in green) and the smooth manifold Σ_1^* (color gradient). We also represent in red the tangent space to Σ_1^* at $\phi(\theta)$. Left: The identifiable case. The intersection is reduced to $\{\phi(\theta)\}$. Right: The non identifiable case. The intersection, represented with a dashed white line, is not reduced to $\{\phi(\theta)\}$.

[15, 44, 45, 46] or in the convolutional case [6, 43]. These references consider a finite X but provide a large sample complexity under which a smartly constructed initialization followed by a first order algorithm allows to stably recover the parameters of the network.

For deep networks, some stable recovery algorithms also exist, for instance for Heavyside activation function [3], or for only recovering the first layer with sparsity assumptions [37] in the ReLU case, but to the best of our knowledge there does not exist any algorithm recovering fully a deep ReLU network from a finite sample.

Model inversion attacks: For deep ReLU networks, when one has full access to the function implemented by the network, a practical algorithm [35] sequentially constructs a sample X and approximately recovers the architecture and the parameters modulo permutation and rescaling. Similarly, formulating the problem as a cryptanalytic problem, [9] reconstructs a functionally equivalent network with fewer requests. As mentioned in Section 1.1, these two references are related to identifiability, but consider a different setting. In this article we consider an arbitrary given X , while they work mostly on its construction.

1.3 Contributions

1/ We establish a necessary and sufficient geometric condition of local identifiability from a finite sample X for deep fully-connected ReLU networks. The condition is that the intersection between an affine space and a smooth manifold is reduced to a single point. See Figure 1 for an illustration.

2/ Considering tangent spaces, we then provide a computable necessary condition of local identifiability from a finite sample X . Since global identifiability implies local identifiability, it is also a computable necessary condition of global identifiability.

3/ We also establish a computable sufficient condition of local identifiability, which is close to the necessary condition. To the best of our knowledge, these are the first testable conditions of local identifiability for any finite input sample. In particular, [39] provides a theoretical condition equivalent to the existence of a finite sample for which local identifiability holds, but does not provide the sample explicitly, nor does it characterize local identifiability for any arbitrary sample.

4/ To prove these results, we develop geometric tools which can be of independent interest for theoretically understanding deep ReLU networks as well as for possible applications. Namely, we introduce local reparameterizations ρ_θ of the network by fixing some weight values as constants. Building on these local parameterizations, we introduce local lifting operators ψ^θ and we decompose the function implemented by the network $f_\theta(x)$ as a composition of ψ^θ , which only depends on the parameters, and a piecewise constant operator α which depends on θ and the inputs x^i . For almost any parameterization θ , the operator α is constant in a neighborhood of θ and consists in applying a linear function to ψ^θ . We show that in fact, the operators ψ^θ are the inverses of coordinate charts of a smooth manifold Σ_1^* , contained in a high dimensional space. We find Σ_1^* to be of particular interest

in representing geometrically some properties of the network parameters (in particular to establish 1/, 2/ and 3/ above).

1.4 Overview of the article

This work is structured as follows. We start by introducing basic tools and already known results, and we state the definition of local identifiability in Section 2. We then introduce the local parameterizations ρ_θ and the set Σ_1^* , and we show that the latter is a smooth manifold in Section 3. This allows us to state our main results in Section 4, that is the geometric and the numerically testable conditions of local identifiability. Finally we discuss in Section 5 the numerical computations needed to test the latter conditions. All the proofs are provided in the appendices.

2 ReLU networks, lifting operator and rescaling of the parameters

2.1 ReLU networks

Let us introduce our notations for deep fully-connected ReLU networks. In this paper, a network is a graph (E, V) of the following form.

- V is a set of neurons, which is divided in $L + 1$ layers, with $L \geq 2$: $V = (V_l)_{l \in \llbracket 0, L \rrbracket}$. V_0 is the input layer, V_L the output layer and the layers V_l with $1 \leq l \leq L - 1$ are the hidden layers. Using the notation $|C|$ for the cardinal of a finite set C , we denote, for all $l \in \llbracket 0, L \rrbracket$, $N_l = |V_l|$ the size of the layer V_l .
- E is the set of all oriented edges $v \rightarrow v'$ between neurons in consecutive layers, that is

$$E = \{v \rightarrow v', v \in V_l, v' \in V_{l+1}, \text{ for } l \in \llbracket 0, L - 1 \rrbracket\}.$$

A network is parameterized by weights and biases, gathered in its parameterization θ , with

$$\theta = ((w_{v \rightarrow v'})_{v \rightarrow v' \in E}, (b_v)_{v \in B}) \in \mathbb{R}^E \times \mathbb{R}^B,$$

where $B = \bigcup_{l=1}^L V_l$. It is also convenient to consider the weights and biases in matrix/vector form: for a given θ , we denote, for $l \in \llbracket 1, L \rrbracket$,

$$W_l = (w_{v \rightarrow v'})_{v' \in V_l, v \in V_{l-1}} \in \mathbb{R}^{N_l \times N_{l-1}} \quad \text{and} \quad b_l = (b_v)_{v \in V_l} \in \mathbb{R}^{N_l}.$$

When dealing with two parameterizations θ and $\tilde{\theta} \in \mathbb{R}^E \times \mathbb{R}^B$, we take as a convention that $w_{v \rightarrow v'}$ and b_v as well as W_l and b_l denote the weights and biases associated to θ , and $\tilde{w}_{v \rightarrow v'}$ and \tilde{b}_v as well as \tilde{W}_l and \tilde{b}_l denote those associated to $\tilde{\theta}$.

The activation function, denoted σ , is always ReLU: for any $p \in \mathbb{N}^*$ and any vector $x = (x_1, \dots, x_p)^T \in \mathbb{R}^p$, it is defined as $\sigma(x) = (\max(x_1, 0), \dots, \max(x_p, 0))^T$.

For a given θ , we define recursively $f_l : \mathbb{R}^{V_0} \rightarrow \mathbb{R}^{V_l}$ (we omit the dependency in θ in the notation for simplicity), for $l \in \llbracket 0, L \rrbracket$, by

- $\forall x \in \mathbb{R}^{V_0}, \quad f_0(x) = x$;
- $\forall l \in \llbracket 1, L - 1 \rrbracket, \forall x \in \mathbb{R}^{V_0}, \quad f_l(x) = \sigma(W_l f_{l-1}(x) + b_l)$;
- $\forall x \in \mathbb{R}^{V_0}, \quad f_L(x) = W_L f_{L-1}(x) + b_L$.

We define the function $f_\theta : \mathbb{R}^{V_0} \rightarrow \mathbb{R}^{V_L}$ implemented by the network of parameter θ as $f_\theta = f_L$.

2.2 The lifting operator ϕ and the activation operator α

For a fixed $x \in \mathbb{R}^{V_0}$, the value of $f_\theta(x)$ is a non-linear function of θ . The goal of this section is to obtain a higher-dimensional representation of θ , that will be written $\phi(\theta)$, and such that $f_\theta(x)$ is locally a linear function of $\phi(\theta)$. This will be achieved with Proposition 1. The function ϕ is called a lifting operator, a wording borrowed from category theory and commonly used in compressed sensing and dictionary learning, for instance in [7]. The components of $\phi(\theta)$ will be associated to paths in the neural network. Linearity in Proposition 1 will correspond to summing over these paths.

We now introduce the paths notations. For all $l \in \llbracket 0, L-1 \rrbracket$, we define

$$\mathcal{P}_l = V_l \times \cdots \times V_{L-1},$$

which is the set of all paths in the network starting from layer l and ending in layer $L-1$. We consider an additional element β which can be interpreted as an empty path and whose role will be clear once ϕ has been defined and Proposition 1 stated. We define

$$\mathcal{P} = \left(\bigcup_{l=0}^{L-1} \mathcal{P}_l \right) \cup \{\beta\}.$$

In a similar way to [39], we can now define the above-mentioned ‘lifting operator’

$$\begin{aligned} \phi : \mathbb{R}^E \times \mathbb{R}^B &\longrightarrow \mathbb{R}^{\mathcal{P} \times V_L} \\ \theta &\longmapsto (\phi_{p,v}(\theta))_{p \in \mathcal{P}, v \in V_L} \end{aligned} \quad (1)$$

by:

- for all $l \in \llbracket 0, L-1 \rrbracket$ and all $p = (v_l, \dots, v_{L-1}) \in \mathcal{P}_l$, and for all $v_L \in V_L$,

$$\phi_{p,v_L}(\theta) = \begin{cases} \prod_{l'=0}^{L-1} w_{v_{l'} \rightarrow v_{l'+1}} & \text{if } l = 0 \\ b_{v_l} \prod_{l'=l}^{L-1} w_{v_{l'} \rightarrow v_{l'+1}} & \text{if } l \geq 1; \end{cases}$$

- for $p = \beta$ and $v_L \in V_L$, $\phi_{\beta,v_L}(\theta) = b_{v_L}$.

To define the activation operator, we first define, for all $l \in \llbracket 1, L-1 \rrbracket$, all $v \in V_l$, all $\theta \in \mathbb{R}^E \times \mathbb{R}^B$ and $x \in \mathbb{R}^{V_0}$,

$$a_v(x, \theta) = \begin{cases} 1 & \text{if } (W_l f_{l-1}(x) + b_l)_v \geq 0 \\ 0 & \text{otherwise,} \end{cases}$$

which is the activation indicator of neuron v . We then define the ‘activation operator’

$$\begin{aligned} \alpha : \mathbb{R}^{V_0} \times (\mathbb{R}^E \times \mathbb{R}^B) &\longrightarrow \mathbb{R}^{1 \times \mathcal{P}} \\ (x, \theta) &\longmapsto (\alpha_p(x, \theta))_{p \in \mathcal{P}} \end{aligned} \quad (2)$$

by:

- for all $l \in \llbracket 0, L-1 \rrbracket$ and all $p = (v_l, \dots, v_{L-1}) \in \mathcal{P}_l$:

$$\alpha_p(x, \theta) = \begin{cases} x_{v_0} \prod_{l'=1}^{L-1} a_{v_{l'}}(x, \theta) & \text{if } l = 0 \\ \prod_{l'=l}^{L-1} a_{v_{l'}}(x, \theta) & \text{if } l \geq 1; \end{cases}$$

- for $p = \beta$, $\alpha_\beta(x, \theta) = 1$.

We then have the announced linear representation of the function f_θ implemented by the network.

Proposition 1. For all $\theta \in \mathbb{R}^E \times \mathbb{R}^B$ and all $x \in \mathbb{R}^{V_0}$, $f_\theta(x)^T = \alpha(x, \theta)\phi(\theta)$.

This result, which is proven in Appendix B, is for instance also stated in [39, Sec. 4] with slightly different notations. Note that each component of the vector $f_\theta(x)$ above is written as a sum over a (very large) number of paths.

Let us reformulate Proposition 1 with several inputs. We consider, for some $n \in \mathbb{N}^*$, some given inputs $x^i \in \mathbb{R}^{V_0}$, with $i \in \llbracket 1, n \rrbracket$. We denote by $X \in \mathbb{R}^{n \times V_0}$ the matrix whose lines are the transpose $(x^i)^T$ of the inputs. For all $\theta \in \mathbb{R}^E \times \mathbb{R}^B$, we denote by $f_\theta(X) \in \mathbb{R}^{n \times V_L}$ the matrix whose lines are the transpose $f_\theta(x^i)^T$ of the corresponding outputs. We also denote by $\alpha(X, \theta) \in \mathbb{R}^{n \times \mathcal{P}}$ the matrix whose lines are the line vectors $\alpha(x^i, \theta)$. Using Proposition 1 for all the x^i , we have the relation

$$f_\theta(X) = \alpha(X, \theta)\phi(\theta). \quad (3)$$

We prove in Appendix B the next proposition, which states that $\theta \mapsto \alpha(X, \theta)$ is piecewise constant.

Proposition 2. For all $n \in \mathbb{N}^*$, for all $X \in \mathbb{R}^{n \times V_0}$, the mapping

$$\begin{aligned} \alpha_X : \mathbb{R}^E \times \mathbb{R}^B &\longrightarrow \mathbb{R}^{n \times \mathcal{P}} \\ \theta &\longmapsto \alpha(X, \theta) \end{aligned}$$

is piecewise-constant, with a finite number of pieces. Furthermore, the boundary of each piece has Lebesgue measure zero. We call Δ_X the union of all these boundaries. The set $\Delta_X \subset \mathbb{R}^E \times \mathbb{R}^B$ is closed and has Lebesgue measure zero.

As discussed before, for a given $X \in \mathbb{R}^{n \times V_0}$, when studying the function $\theta \mapsto f_\theta(X)$, Proposition 2 alongside (3) shows that on a piece over which α_X is constant, $f_\theta(X)$ depends linearly on $\phi(\theta)$. Since Δ_X is closed with measure zero, for almost all $\tilde{\theta} \in \mathbb{R}^E \times \mathbb{R}^B$, there exists a neighborhood of $\tilde{\theta}$ over which α_X is constant. As noted for instance by Stock and Gribonval [39, Sec. 2], for any θ in such a neighborhood, we thus have

$$f_\theta(X) - f_{\tilde{\theta}}(X) = \alpha(X, \tilde{\theta}) (\phi(\theta) - \phi(\tilde{\theta})). \quad (4)$$

Hence, studying ϕ will allow us to understand better how $f_\theta(X)$ locally depends on θ .

2.3 Invariant rescaling operations on θ

Some well-known rescaling operations on the parameters θ do not affect the value of $\phi(\theta)$. Before detailing them, let us define, for all $t \in \mathbb{R}$, the sign indicator $\text{sign}(t)$ as 1, 0 or -1 depending on whether $t > 0$, $t = 0$ or $t < 0$ respectively. For any $\theta \in \mathbb{R}^E \times \mathbb{R}^B$, we then define

$$\text{sign}(\theta) = \left((\text{sign}(w_{v \rightarrow v'}))_{v \rightarrow v' \in E}, (\text{sign}(b_v))_{v \in B} \right) \in \{-1, 0, 1\}^E \times \{-1, 0, 1\}^B.$$

We can now describe the rescaling operations.

Definition 3. Let $\theta \in \mathbb{R}^E \times \mathbb{R}^B$ and $\tilde{\theta} \in \mathbb{R}^E \times \mathbb{R}^B$.

- We say that θ is equivalent to $\tilde{\theta}$ modulo rescaling, and we write $\theta \stackrel{R}{\sim} \tilde{\theta}$ iff there exists a family of vectors $(\lambda^0, \dots, \lambda^L) \in (\mathbb{R}^*)^{V_0} \times \dots \times (\mathbb{R}^*)^{V_L}$, with $\lambda^0 = \mathbf{1}_{V_0}$ and $\lambda^L = \mathbf{1}_{V_L}$, such that, for all $l \in \llbracket 1, L \rrbracket$,

$$\begin{cases} W_l = \text{Diag}(\lambda^l) \tilde{W}_l \text{Diag}(\lambda^{l-1})^{-1} \\ b_l = \text{Diag}(\lambda^l) \tilde{b}_l. \end{cases} \quad (5)$$

- We say that θ is equivalent to $\tilde{\theta}$ modulo positive rescaling, and we write $\theta \sim \tilde{\theta}$ iff

$$\theta \stackrel{R}{\sim} \tilde{\theta} \quad \text{and} \quad \text{sign}(\theta) = \text{sign}(\tilde{\theta}).$$

For all $l \in \llbracket 1, L \rrbracket$, to satisfy (5) is equivalent to satisfy, for all $(v_{l-1}, v_l) \in V_{l-1} \times V_l$,

$$\begin{cases} w_{v_{l-1} \rightarrow v_l} = \frac{\lambda_{v_l}^l}{\lambda_{v_{l-1}}^{l-1}} \tilde{w}_{v_{l-1} \rightarrow v_l} \\ b_{v_l} = \lambda_{v_l}^l \tilde{b}_{v_l}. \end{cases} \quad (6)$$

The relations $\stackrel{R}{\sim}$ and \sim are equivalence relations on the set of parameters $\mathbb{R}^E \times \mathbb{R}^B$. The equivalence modulo positive rescaling \sim is a well-known invariant for ReLU networks [38, 39, 5, 28, 41]. We have indeed the following property: if $\theta \sim \tilde{\theta}$, for all $x \in \mathbb{R}^{V_0}$,

$$f_\theta(x) = f_{\tilde{\theta}}(x). \quad (7)$$

One of the interests of the operator ϕ is that it captures this invariant, as described by Stock and Gribonval [39, Sec. 2.4]. Propositions 4 and 5 are similar to their results and are restated here and proven in Appendix B for completeness. Indeed, combining the definition of ϕ with (6), we have the following property.

Proposition 4. For all $\theta, \tilde{\theta} \in \mathbb{R}^E \times \mathbb{R}^B$, we have

$$\theta \stackrel{R}{\sim} \tilde{\theta} \implies \phi(\theta) = \phi(\tilde{\theta}),$$

and thus in particular

$$\theta \sim \tilde{\theta} \implies \phi(\theta) = \phi(\tilde{\theta}).$$

The reciprocal of Proposition 4 holds provided we exclude some degenerate cases. Let us denote, for any $l \in \llbracket 1, L-1 \rrbracket$ and any $v \in V_l$, by $w_{\bullet \rightarrow v}$ the vector $(w_{v' \rightarrow v})_{v' \in V_{l-1}} \in \mathbb{R}^{V_{l-1}}$ and by $w_{v \rightarrow \bullet}$ the vector $(w_{v \rightarrow v'})_{v' \in V_{l+1}} \in \mathbb{R}^{V_{l+1}}$. We define the following set, which is close to the notion of ‘non admissible parameter’ in [39]:

$$S = \{\theta \in \mathbb{R}^E \times \mathbb{R}^B, \exists v \in V_1 \cup \dots \cup V_{L-1}, w_{v \rightarrow \bullet} = 0 \text{ or } (w_{\bullet \rightarrow v}, b_v) = (0, 0)\}.$$

When $w_{v \rightarrow \bullet} = 0$, all the outward weights of v are zero. When $(w_{\bullet \rightarrow v}, b_v) = (0, 0)$, all the inward weights as well as the bias of v are zero, so for any input the information flowing through neuron v is always zero. In both cases, the neuron v does not contribute to the output and could be removed from the network without changing the function f_θ . Since the set S is a finite union of linear subspaces of codimension larger than 1, it is closed and has Lebesgue measure zero. We can thus exclude the degenerate cases in S without loss of generality. Proposition 5 states that the reciprocal of Proposition 4 holds over $(\mathbb{R}^E \times \mathbb{R}^B) \setminus S$.

Proposition 5. For all $\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S$, for all $\tilde{\theta} \in \mathbb{R}^E \times \mathbb{R}^B$,

$$\phi(\theta) = \phi(\tilde{\theta}) \implies \theta \stackrel{R}{\sim} \tilde{\theta}.$$

2.4 Local identifiability

We have now introduced all the concepts used in the formal definition of ‘local identifiability’.

Definition 6. Let $X \in \mathbb{R}^{n \times V_0}$ and $\theta \in \mathbb{R}^E \times \mathbb{R}^B$. We say that θ is *locally identifiable from X* if there exists $\epsilon > 0$ such that for all $\tilde{\theta} \in \mathbb{R}^E \times \mathbb{R}^B$, if $\|\theta - \tilde{\theta}\|_\infty < \epsilon$,

$$f_\theta(X) = f_{\tilde{\theta}}(X) \implies \theta \sim \tilde{\theta}.$$

3 The smooth manifold Σ_1^*

We explained in the previous section that studying ϕ allows to better understand how the output $f_\theta(X)$ locally depends on θ . The image of ϕ is of particular interest in this study and is the subject of this section. We define

$$\Sigma_1^* = \{\phi(\theta), \theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S\}.$$

The main result of this section, Theorem 7, states that Σ_1^* is a smooth manifold. This result is a key element of the article. Indeed, it allows to consider tangent spaces to Σ_1^* , and by doing so, to linearize the geometric characterization of Theorem 8 illustrated in Figure 1. Instead of considering the intersection between a smooth manifold and an affine space as in Theorem 8, this instead allows to consider the intersection between two affine spaces, which can be characterized with rank computations as in Theorems 9 and 10.

To show this result, we need local injectivity. In this aim, let us consider a fixed θ and analyze the functions $u \mapsto f_{\theta+u}(X)$ and $u \mapsto \phi(\theta+u)$ for u around 0. We can select $N_1 + \dots + N_{L-1}$ scalar scaling parameters (each in a neighborhood of 1), and use them to ‘rescale’ $\theta+u$ as in Definition 3, leaving $f_{\theta+u}(X)$ and $\phi(\theta+u)$ unchanged ((7) and Proposition 4). Locally, at first order, this means that there are $N_1 + \dots + N_{L-1}$ linear combinations of u which leave $f_{\theta+u}(X)$ and $\phi(\theta+u)$ invariant. In order to obtain injectivity with respect to u , locally around 0, we will fix $N_1 + \dots + N_{L-1}$ components of u as follows.

For each neuron v in a hidden layer, we choose the outward edge $v \rightarrow v'$ whose weight $w_{v \rightarrow v'}$ has largest (absolute) value (if there are several such edges, we choose one arbitrarily). We denote by $s_{\max}^\theta(v)$ such a neuron v' . For each neuron v in a hidden layer V_l , there is exactly one neuron $s_{\max}^\theta(v)$ in the layer V_{l+1} , and one corresponding edge $v \rightarrow s_{\max}^\theta(v)$. See Figure 2 for an illustration. We will set to 0 the components of u corresponding to all the edges of the form $v \rightarrow s_{\max}^\theta(v)$. Intuitively,




Figure 2: Left: The outward edges of a hidden neuron v and their weights. In this example, $v_1 = s_{\max}^\theta(v)$, so the weight of the edge in red, $v \rightarrow v_1$, has its value fixed as $w_{v \rightarrow v_1}$. The weights of the remaining edges, $\tau_{v \rightarrow v_2}$ and $\tau_{v \rightarrow v_3}$, are free to vary. Right: In red, all the edges whose weights are fixed. The remaining edges, in black, constitute the set F_θ .

it will not limit the set of functions $f_{\tilde{\theta}}$, in the vicinity of f_θ ; but will permit to obtain a one-to-one correspondence between u and $f_{\theta+u}$.

More precisely, let us denote by $F_\theta \subset E$ the set of remaining edges, which is formally defined as¹

$$F_\theta = E \setminus \left(\bigcup_{l=1}^{L-1} \{(v, s_{\max}^\theta(v)), v \in V_l\} \right). \quad (8)$$

The mapping from the space of restricted parameters $\mathbb{R}^{F_\theta} \times \mathbb{R}^B$ to the parameter space $\mathbb{R}^E \times \mathbb{R}^B$ locally around θ is simply given by the following application

$$\rho_\theta : \mathbb{R}^{F_\theta} \times \mathbb{R}^B \longrightarrow \mathbb{R}^E \times \mathbb{R}^B$$

$$\tau \longmapsto \tilde{\theta} \quad \text{such that} \quad \begin{cases} \forall (v, v') \in F_\theta, & \tilde{w}_{v \rightarrow v'} = \tau_{v \rightarrow v'} \\ \forall (v, v') \in E \setminus F_\theta, & \tilde{w}_{v \rightarrow v'} = w_{v \rightarrow v'} \\ \forall v \in B, & \tilde{b}_v = \tau_v. \end{cases} \quad (9)$$

In particular, if we define $\tau_\theta \in \mathbb{R}^{F_\theta} \times \mathbb{R}^B$ by $(\tau_\theta)_{v \rightarrow v'} = w_{v \rightarrow v'}$ and $(\tau_\theta)_v = b_v$, we have $\rho_\theta(\tau_\theta) = \theta$. The function ρ_θ is affine and injective. We define

$$U_\theta = \rho_\theta^{-1} \left((\mathbb{R}^E \times \mathbb{R}^B) \setminus S \right), \quad (10)$$

which is an open set of $\mathbb{R}^{F_\theta} \times \mathbb{R}^B$. We define, for all $\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S$, the local lifting operator

$$\psi^\theta : \begin{array}{l} U_\theta \longrightarrow \mathbb{R}^{\mathcal{P} \times V_L} \\ \tau \longmapsto \phi \circ \rho_\theta(\tau). \end{array} \quad (11)$$

One can show that ψ^θ is C^∞ and that it is a homeomorphism from U_θ onto its image (see the proofs in Appendix C), which we denote V_θ and is thus an open subset of Σ_1^* (with the topology induced on Σ_1^* by the standard topology on $\mathbb{R}^{\mathcal{P} \times V_L}$). In particular, since $\rho_\theta(\tau_\theta) = \theta$, we have $\phi(\theta) = \psi^\theta(\tau_\theta) \in V_\theta$. We have the following fundamental result that will allow us to consider and make use the tangent spaces of Σ_1^* .

Theorem 7. Σ_1^* is a smooth manifold of $\mathbb{R}^{\mathcal{P} \times V_L}$ of dimension

$$|F_\theta| + |B| = N_0 N_1 + N_1 N_2 + \cdots + N_{L-1} N_L + N_L,$$

and the family $(V_\theta, (\psi^\theta)^{-1})_{\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S}$ is an atlas.

Theorem 7 is proven in Appendix C. Besides being key in Section 4, Theorem 7 (both the smooth manifold nature of Σ_1^* and the explicit atlas $(V_\theta, (\psi^\theta)^{-1})_{\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S}$) may also be considered of more general independent interest. To our knowledge, such a result has not been established elsewhere in the literature. Notice that, as announced, despite the use of restricted parameters in $\mathbb{R}^{F_\theta} \times \mathbb{R}^B$, we can represent the *whole* tangent space at any point of Σ_1^* . The only consequence of the restriction is the uniqueness of the representation of the elements of tangent spaces.

¹Note, in the definition of F_θ , the index l starting at $l = 1$ and not $l = 0$.

4 Main results: necessary and sufficient conditions for local identifiability

The main results of this paper rely on the decomposition (4) introduced in Section 2. To reformulate (4), let us introduce the linear operator $A(X, \theta)$, which simply corresponds to the matrix product with $\alpha(X, \theta)$:

$$\begin{aligned} A(X, \theta) : \mathbb{R}^{\mathcal{P} \times V_L} &\longrightarrow \mathbb{R}^{n \times V_L} \\ \eta &\longmapsto \alpha(X, \theta)\eta, \end{aligned}$$

where $\alpha(X, \theta)\eta$ is the matrix product between $\alpha(X, \theta) \in \mathbb{R}^{n \times \mathcal{P}}$ and $\eta \in \mathbb{R}^{\mathcal{P} \times V_L}$. The operator $A(X, \theta)$ inherits the properties of $\alpha(X, \theta)$, in particular those stated in Proposition 2. Using $A(X, \theta)$, the relation (4) satisfied by $\tilde{\theta}$ in the neighborhood of θ becomes

$$f_\theta(X) - f_{\tilde{\theta}}(X) = A(X, \theta) \cdot (\phi(\theta) - \phi(\tilde{\theta})). \quad (12)$$

Let us also define the affine space (set-sum of a fixed point and a vector space)

$$N(X, \theta) = \phi(\theta) + \text{Ker } A(X, \theta).$$

If a parameterization $\tilde{\theta} \in \mathbb{R}^E \times \mathbb{R}^B$ is such that $f_{\tilde{\theta}}(X) = f_\theta(X)$ and (12) holds, then $\phi(\theta) - \phi(\tilde{\theta}) \in \text{Ker } A(X, \theta)$, so by definition $\phi(\tilde{\theta}) \in N(X, \theta)$. Since for $\tilde{\theta}$ in the neighborhood of θ , we also have $\phi(\tilde{\theta}) \in \Sigma_1^*$, we see that local identifiability is closely related to the nature of the intersection between the smooth manifold Σ_1^* and the affine subspace $N(X, \theta)$.

Indeed, let us denote by $B_\infty(\phi(\theta), \epsilon) = \{\eta \in \mathbb{R}^{\mathcal{P} \times V_L}, \|\phi(\theta) - \eta\|_\infty < \epsilon\}$ the ball of center $\phi(\theta)$ and of radius $\epsilon > 0$. We have the following geometric necessary and sufficient condition of local identifiability, which states that local identifiability of θ holds if and only if the intersection between Σ_1^* and $N(X, \theta)$ is locally reduced to the single point $\{\phi(\theta)\}$.

Theorem 8. *For any $X \in \mathbb{R}^{n \times V_0}$ and $\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus (S \cup \Delta_X)$, the two following statements are equivalent.*

- i) θ is locally identifiable from X .
- ii) There exists $\epsilon > 0$ such that $B_\infty(\phi(\theta), \epsilon) \cap \Sigma_1^* \cap N(X, \theta) = \{\phi(\theta)\}$.

Theorem 8 is proven in Appendix D, and is illustrated in Figure 1. This geometric condition is crucial for showing the next two results which give testable conditions of identifiability. Theorems 9 and 10 rely on the rank of $A(X, \theta)$ and of another linear operator $\Gamma(X, \theta)$, which we now define. Since, as we said, the function ψ^θ is C^∞ , let us denote by $D\psi^\theta(\tau) : \mathbb{R}^{F_\theta} \times \mathbb{R}^B \rightarrow \mathbb{R}^{\mathcal{P} \times V_L}$ its differential at the point τ , for any $\tau \in U_\theta$. We define the linear operator $\Gamma(X, \theta) : \mathbb{R}^{F_\theta} \times \mathbb{R}^B \rightarrow \mathbb{R}^{n \times V_L}$ by

$$\Gamma(X, \theta) = A(X, \theta) \circ D\psi^\theta(\tau_\theta). \quad (13)$$

We denote $R_A = \text{rank}(A(X, \theta))$ and $R_\Gamma = \text{rank}(\Gamma(X, \theta))$. Since $\Gamma(X, \theta)$ is defined on $\mathbb{R}^{F_\theta} \times \mathbb{R}^B$, we have $0 \leq R_\Gamma \leq |F_\theta| + |B|$, and the expression (13) shows that we also have $0 \leq R_\Gamma \leq R_A$. We can now define the two following conditions.

Condition C_N . Condition C_N is satisfied by (θ, X) iff $R_\Gamma < R_A$ or $R_\Gamma = |F_\theta| + |B|$.

Condition C_S . Condition C_S is satisfied by (θ, X) iff $R_\Gamma = |F_\theta| + |B|$.

The following result states that C_N is necessary for local and therefore global identifiability.

Theorem 9 (Necessary condition of identifiability). *Let $X \in \mathbb{R}^{n \times V_0}$ and $\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus (S \cup \Delta_X)$. If C_N is not satisfied, then θ is not locally identifiable from X (thus not globally identifiable).*

The following result states that C_S is a sufficient condition of local identifiability.

Theorem 10 (Sufficient condition of local identifiability). *Let $X \in \mathbb{R}^{n \times V_0}$ and $\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus (S \cup \Delta_X)$. If C_S is satisfied, then θ is locally identifiable from X .*

Both theorems are proven in Appendix D. To discuss these two results, let us point out that the output spaces of $\Gamma(X, \theta)$ and $A(X, \theta)$ have the same dimension, equal to nN_L . Each new input adds N_L to this dimension. One can verify that $R_A - R_\Gamma$ is initially 0 and cannot decrease when new inputs are

added. If a new input leads to $R_A > R_\Gamma$, it can be discarded to preserve $R_A = R_\Gamma$. Moreover, such an input seems unlikely when $R_A < |F_\theta| + |B|$. If the equality $R_\Gamma = R_A$ is enforced, the condition $R_\Gamma = |F_\theta| + |B|$ is both necessary and sufficient. Finally, to satisfy $R_\Gamma = |F_\theta| + |B|$, the dimensions must satisfy $nN_L \geq |F_\theta| + |B|$. The general belief is that the latter is the condition of identifiability since nN_L is the number of scalar measurements and $|F_\theta| + |B|$ is the number of independent free parameters, see Theorem 7.

5 Checking the conditions numerically

The key benefit of the conditions C_N and C_S , compared to the existing literature, is that they can be numerically tested for any fixed finite sample. They need the computation of the rank of two linear operators, namely $\Gamma(X, \theta)$ and $A(X, \theta)$. The operator $\Gamma(X, \theta)$ satisfies the following:

Proposition 11. *Let $X \in \mathbb{R}^{n \times V_0}$ and $\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus (S \cup \Delta_X)$. The function $\tau \mapsto f_{\rho_\theta(\tau)}(X)$, for $\tau \in U_\theta$ is differentiable in a neighborhood of τ_θ , and we denote by $D_\tau f_{\rho_\theta(\tau)}(X)$ its differential at τ_θ . We have*

$$D_\tau f_{\rho_\theta(\tau)}(X) = \Gamma(X, \theta). \quad (14)$$

The proof of Proposition 11 is in Appendix E. Since the reparameterization with ρ_θ simply consists in fixing the weights of the edges $v \rightarrow s_{\max}^\theta(v)$ to the value $w_{v \rightarrow s_{\max}^\theta(v)}$, (59) shows that the coefficients of $\Gamma(X, \theta)$ can be computed by a classic backpropagation algorithm N_L times for each input x^i , simply omitting the derivatives with respect to the edges of the form $v \rightarrow s_{\max}^\theta(v)$. An explicit expression of the coefficients of $\Gamma(X, \theta)$ is given in the Appendix E.

To be satisfied, C_S needs the dimensions of $\Gamma(X, \theta)$ to satisfy $nN_L \geq |F_\theta| + |B|$. One then needs to compute the rank R_Γ of $\Gamma(X, \theta)$, which means computing the rank of a $nN_L \times (|F_\theta| + |B|)$ matrix. Existing algorithms allow to do this with a complexity $O(nN_L(|F_\theta| + |B|)^\omega)$ (up to polylog terms), where ω is the matrix multiplication exponent and satisfies $\omega < 2.38$ [10].

When it comes to C_N , one needs in addition to know the rank R_A of $A(X, \theta)$, which, as Proposition 12 states, requires to compute the rank of $\alpha(X, \theta)$.

Proposition 12. *Let $X \in \mathbb{R}^{n \times V_0}$ and $\theta \in \mathbb{R}^E \times \mathbb{R}^B$. We have $R_A = N_L \text{rank}(\alpha(X, \theta))$.*

The dimensions of $\alpha(X, \theta)$ are sensibly larger, with $|\mathcal{P}|$ columns and n lines, and typically $|\mathcal{P}| \gg n$. However it may have some sparsity properties, as its entries consist in products of activation indicators (with possibly one input $x_{v_0}^i$), any one of them being zero causing many entries to vanish. The question of the efficient computation of R_A still needs to be explored and is left as open for future work.

6 Conclusion

This paper is the first to characterize local identifiability for deep ReLU networks for any given finite sample, with testable conditions. The practical use of these conditions deserves follow-up research, and so does an extension of our approach to inverse stability. The role of ReLU is crucial in our approach, especially for the necessary condition of local identifiability and with the linear representation (Proposition 1). In the end, from Theorem 10 and Proposition 11, the sufficient condition for local indentifiability is expressed from the Jacobian matrix of the neural network function with respect to its parameters. Extending this to other activation functions than ReLU is an interesting perspective.

Acknowledgments and Disclosure of Funding

The authors would like to thank Pierre Stock and Rémi Gribonval for the fruitful discussions around this work, notably regarding the construction of ϕ and its link to the question of local identifiability.

This work has benefited from the AI Interdisciplinary Institute ANITI. ANITI is funded by the French ‘‘Investing for the Future – PIA3’’ program under the Grant agreement n°ANR-19-PI3A-0004.

The authors gratefully acknowledge the support of the DEEL project.²

²<https://www.deel.ai/>

References

- [1] Rilwan A Adewoyin, Peter Dueben, Peter Watson, Yulan He, and Ritabrata Dutta. Tru-net: a deep learning approach to high resolution prediction of rainfall. *Machine Learning*, 110(8): 2035–2062, 2021.
- [2] Francesca Albertini, Eduardo D Sontag, and Vincent Maillot. Uniqueness of weights for neural networks. *Artificial Neural Networks for Speech and Vision*, pages 115–125, 1993.
- [3] Sanjeev Arora, Aditya Bhaskara, Rong Ge, and Tengyu Ma. Provable bounds for learning some deep representations. In Eric P. Xing and Tony Jebara, editors, *Proceedings of the 31st International Conference on Machine Learning*, volume 32 of *Proceedings of Machine Learning Research*, pages 584–592, Beijing, China, 22–24 Jun 2014. PMLR.
- [4] Mikhail Belkin, Daniel Hsu, and Ji Xu. Two models of double descent for weak features. *SIAM Journal on Mathematics of Data Science*, 2(4):1167–1180, 2020.
- [5] Joachim Bona-Pellissier, François Bachoc, and François Malgouyres. Parameter identifiability of a deep feedforward ReLU neural network. *arXiv preprint arXiv:2112.12982*, 2021.
- [6] Alon Brutzkus and Amir Globerson. Globally optimal gradient descent for a ConvNet with Gaussian inputs. In *Proceedings of the 34th International Conference on Machine Learning—Volume 70*, pages 605–614, 2017.
- [7] Emmanuel J Candes, Yonina C Eldar, Thomas Strohmer, and Vladislav Voroninski. Phase retrieval via matrix completion. *SIAM review*, 57(2):225–251, 2015.
- [8] Nicholas Carlini, Chang Liu, Úlfar Erlingsson, Jernej Kos, and Dawn Song. The secret sharer: Evaluating and testing unintended memorization in neural networks. In *28th {USENIX} Security Symposium ({USENIX} Security 19)*, pages 267–284, 2019.
- [9] Nicholas Carlini, Matthew Jagielski, and Ilya Mironov. Cryptanalytic extraction of neural network models. In *Annual International Cryptology Conference*, pages 189–218. Springer, 2020.
- [10] Ho Yee Cheung, Tsz Chiu Kwok, and Lap Chi Lau. Fast matrix rank algorithms and applications. *Journal of the ACM (JACM)*, 60(5):1–25, 2013.
- [11] Dennis Maximilian Elbrächter, Julius Berner, and Philipp Grohs. How degenerate is the parametrization of neural networks with the ReLU activation function? In *Advances in Neural Information Processing Systems*, volume 32, 2019.
- [12] Charles Fefferman. Reconstructing a neural net from its output. *Revista Matemática Iberoamericana*, 10(3):507–555, 1994.
- [13] Matt Fredrikson, Somesh Jha, and Thomas Ristenpart. Model inversion attacks that exploit confidence information and basic countermeasures. In *Proceedings of the 22nd ACM SIGSAC Conference on Computer and Communications Security*, pages 1322–1333, 2015.
- [14] Haoyu Fu, Yuejie Chi, and Yingbin Liang. Guaranteed recovery of one-hidden-layer neural networks via cross entropy. *IEEE Transactions on Signal Processing*, 68:3225–3235, 2020.
- [15] Rong Ge, Jason D Lee, and Tengyu Ma. Learning one-hidden-layer neural networks with landscape design. In *6th International Conference on Learning Representations, ICLR 2018*, 2018.
- [16] Awni Hannun, Carl Case, Jared Casper, Bryan Catanzaro, Greg Diamos, Erich Elsen, Ryan Prenger, Sanjeev Satheesh, Shubho Sengupta, Adam Coates, et al. Deep speech: Scaling up end-to-end speech recognition. *arXiv preprint arXiv:1412.5567*, 2014.
- [17] Geoffrey Hinton, Li Deng, Dong Yu, George E. Dahl, Abdel-rahman Mohamed, Navdeep Jaitly, Andrew Senior, Vincent Vanhoucke, Patrick Nguyen, Tara N. Sainath, and Brian Kingsbury. Deep neural networks for acoustic modeling in speech recognition: The shared views of four research groups. *IEEE Signal Processing Magazine*, 29(6):82–97, 2012.

- [18] Paul C Kainen, Věra Kůrková, Vladik Kreinovich, and Ongard Sirisaengtaksin. Uniqueness of network parametrization and faster learning. *Neural, Parallel & Scientific Computations*, 2(4): 459–466, 1994.
- [19] Nal Kalchbrenner and Phil Blunsom. Recurrent continuous translation models. In *Proceedings of the 2013 conference on empirical methods in natural language processing*, pages 1700–1709, 2013.
- [20] Alex Krizhevsky, Ilya Sutskever, and Geoffrey E Hinton. Imagenet classification with deep convolutional neural networks. *Advances in neural information processing systems*, 25:1097–1105, 2012.
- [21] Alexey Kurakin, Ian Goodfellow, and Samy Bengio. Adversarial examples in the physical world. *International Conference on Learning Representations*, 2017.
- [22] Věra Kůrková and Paul C Kainen. Functionally equivalent feedforward neural networks. *Neural Computation*, 6(3):543–558, 1994.
- [23] François Malgouyres and Joseph Landsberg. On the identifiability and stable recovery of deep/multi-layer structured matrix factorization. In *IEEE, Info. Theory Workshop*, Sept. 2016.
- [24] François Malgouyres and Joseph Landsberg. Multilinear compressive sensing and an application to convolutional linear networks. *SIAM Journal on Mathematics of Data Science*, 1(3):446–475, 2019.
- [25] Francois Malgouyres. On the stable recovery of deep structured linear networks under sparsity constraints. In *Mathematical and Scientific Machine Learning*, pages 107–127. PMLR, 2020.
- [26] Tomas Mikolov, Martin Karafiát, Lukas Burget, Jan Cernocký, and Sanjeev Khudanpur. Recurrent neural network based language model. In *Interspeech*, volume 2, pages 1045–1048, 2010.
- [27] Tomáš Mikolov, Kai Chen, Greg Corrado, and Jeffrey Dean. Efficient estimation of word representations in vector space. In Yoshua Bengio and Yann LeCun, editors, *1st International Conference on Learning Representations, ICLR 2013, Scottsdale, Arizona, USA, May 2-4, 2013, Workshop Track Proceedings*, 2013.
- [28] Behnam Neyshabur, Russ R Salakhutdinov, and Nati Srebro. Path-SGD: Path-normalized optimization in deep neural networks. *Advances in neural information processing systems*, 28, 2015.
- [29] Philipp Petersen, Mones Raslan, and Felix Voigtlaender. Topological properties of the set of functions generated by neural networks of fixed size. *Foundations of Computational Mathematics*, 21:375–444, 2021.
- [30] Henning Petzka, Martin Trimmel, and Cristian Sminchisescu. Notes on the symmetries of 2-layer ReLU-networks. In *Proceedings of the Northern Lights Deep Learning Workshop*, volume 1, pages 6–6, 2020.
- [31] Mary Phuong and Christoph H. Lampert. Functional vs. parametric equivalence of ReLU networks. In *International Conference on Learning Representations*, 2020.
- [32] José Pedro Pinto, André Pimenta, and Paulo Novais. Deep learning and multivariate time series for cheat detection in video games. *Machine Learning*, 110(11):3037–3057, 2021.
- [33] Joseph Redmon, Santosh Divvala, Ross Girshick, and Ali Farhadi. You only look once: Unified, real-time object detection. In *Proceedings of the IEEE conference on computer vision and pattern recognition*, pages 779–788, 2016.
- [34] Shaoqing Ren, Kaiming He, Ross Girshick, and Jian Sun. Faster R-CNN: Towards real-time object detection with region proposal networks. *Advances in neural information processing systems*, 28:91–99, 2015.

- [35] David Rolnick and Konrad Kording. Reverse-engineering deep ReLU networks. In Hal Daumé III and Aarti Singh, editors, *Proceedings of the 37th International Conference on Machine Learning*, volume 119 of *Proceedings of Machine Learning Research*, pages 8178–8187, 13–18 Jul 2020.
- [36] Haşim Sak, Andrew Senior, and Françoise Beaufays. Long short-term memory recurrent neural network architectures for large scale acoustic modeling. In *Fifteenth Annual Conference of the International Speech Communication Association*, 2014.
- [37] Hanie Sedghi and Anima Anandkumar. Provable methods for training neural networks with sparse connectivity. In *Deep Learning and representation learning workshop: NIPS*, 2014.
- [38] Pierre Stock. *Efficiency and Redundancy in Deep Learning Models : Theoretical Considerations and Practical Applications*. PhD thesis, Université de Lyon, April 2021. URL <https://tel.archives-ouvertes.fr/tel-03208517>.
- [39] Pierre Stock and Rémi Gribonval. An Embedding of ReLU Networks and an Analysis of their Identifiability. *Constructive Approximation*, 2022. URL <https://hal.archives-ouvertes.fr/hal-03292203>.
- [40] Héctor J Sussmann. Uniqueness of the weights for minimal feedforward nets with a given input-output map. *Neural networks*, 5(4):589–593, 1992.
- [41] Mingyang Yi, Qi Meng, Wei Chen, Zhi-ming Ma, and Tie-Yan Liu. Positively scale-invariant flatness of ReLU neural networks. *arXiv preprint arXiv:1903.02237*, 2019.
- [42] Jialong Zhang, Zhongshu Gu, Jiyong Jang, Hui Wu, Marc Ph Stoecklin, Heqing Huang, and Ian Molloy. Protecting intellectual property of deep neural networks with watermarking. In *Proceedings of the 2018 on Asia Conference on Computer and Communications Security*, pages 159–172, 2018.
- [43] Shuai Zhang, Meng Wang, Jinjun Xiong, Sijia Liu, and Pin-Yu Chen. Improved linear convergence of training CNNs with generalizability guarantees: A one-hidden-layer case. *IEEE Transactions on Neural Networks and Learning Systems*, 32(6):2622–2635, 2020.
- [44] Xiao Zhang, Yaodong Yu, Lingxiao Wang, and Quanquan Gu. Learning one-hidden-layer ReLU networks via gradient descent. In *The 22nd International Conference on Artificial Intelligence and Statistics*, pages 1524–1534. PMLR, 2019.
- [45] Kai Zhong, Zhao Song, Prateek Jain, Peter L Bartlett, and Inderjit S Dhillon. Recovery guarantees for one-hidden-layer neural networks. In *Proceedings of the 34th International Conference on Machine Learning-Volume 70*, pages 4140–4149, 2017.
- [46] Mo Zhou, Rong Ge, and Chi Jin. A local convergence theory for mildly over-parameterized two-layer neural network. *arXiv preprint arXiv:2102.02410*, 2021.

Checklist

1. For all authors...
 - (a) Do the main claims made in the abstract and introduction accurately reflect the paper’s contributions and scope? [Yes] The outline in Section 1.4 (“Overview of the article”) provides pointers to where the claimed contributions of the paper are provided.
 - (b) Did you describe the limitations of your work? [Yes] Section 5 acknowledges the open problem of an efficient computation of the rank of $\alpha(X, \theta)$ and Section 6 describes other remaining open questions.
 - (c) Did you discuss any potential negative societal impacts of your work? [N/A] This is a theoretical/foundation work that adds to the theory and methodology of deep learning. As for any such contributions, the positive or negative societal impact will depend on the application case. We do not promote any harmful use of this theory, but we expand on the existing knowledge.

- (d) Have you read the ethics review guidelines and ensured that your paper conforms to them? [Yes] See previous question.
- 2. If you are including theoretical results...
 - (a) Did you state the full set of assumptions of all theoretical results? [Yes] All our results explicitly refer to their required assumptions. Some general assumptions that hold throughout the paper are also stated at the beginning.
 - (b) Did you include complete proofs of all theoretical results? [Yes] All the proofs are provided in the supplement.
- 3. If you ran experiments...
 - (a) Did you include the code, data, and instructions needed to reproduce the main experimental results (either in the supplemental material or as a URL)? [N/A] We did not run experiments.
 - (b) Did you specify all the training details (e.g., data splits, hyperparameters, how they were chosen)? [N/A]
 - (c) Did you report error bars (e.g., with respect to the random seed after running experiments multiple times)? [N/A]
 - (d) Did you include the total amount of compute and the type of resources used (e.g., type of GPUs, internal cluster, or cloud provider)? [N/A]
- 4. If you are using existing assets (e.g., code, data, models) or curating/releasing new assets...
 - (a) If your work uses existing assets, did you cite the creators? [N/A] We did not use existing assets (code, data or models) nor cure/release new assets (code, data or models).
 - (b) Did you mention the license of the assets? [N/A]
 - (c) Did you include any new assets either in the supplemental material or as a URL? [N/A]
 - (d) Did you discuss whether and how consent was obtained from people whose data you're using/curating? [N/A]
 - (e) Did you discuss whether the data you are using/curating contains personally identifiable information or offensive content? [N/A]
- 5. If you used crowdsourcing or conducted research with human subjects...
 - (a) Did you include the full text of instructions given to participants and screenshots, if applicable? [N/A] We did not use crowdsourcing nor conducted research with human subjects.
 - (b) Did you describe any potential participant risks, with links to Institutional Review Board (IRB) approvals, if applicable? [N/A]
 - (c) Did you include the estimated hourly wage paid to participants and the total amount spent on participant compensation? [N/A]

A Notations

In this section, we define notations, many of which are standard, that are useful in the proofs.

We denote by \mathbb{N} the set of all natural numbers, including 0, and by \mathbb{N}^* the set \mathbb{N} without 0. We denote by \mathbb{Z} the set of all integers. For any $a, b \in \mathbb{Z}$, we denote by $\llbracket a, b \rrbracket$ the set of all integers $k \in \mathbb{Z}$ satisfying $a \leq k \leq b$. For any finite set A , we denote by $|A|$ the cardinal of A .

For $n, N \in \mathbb{N}^*$, we denote by \mathbb{R}^N the N -dimensional real vector space and by $\mathbb{R}^{n \times N}$ the vector space of real matrices with n lines and N columns. For a vector $x = (x_1, \dots, x_N)^T \in \mathbb{R}^N$, we use the norm $\|x\|_\infty = \max_{i \in \llbracket 1, N \rrbracket} |x_i|$. For $x \in \mathbb{R}^N$ and $r > 0$, we denote $B_\infty(x, r) = \{y \in \mathbb{R}^N, \|y - x\|_\infty < r\}$.

For any vector $x = (x_1, \dots, x_N)^T \in \mathbb{R}^N$, we define $\text{sign}(x) = (\text{sign}(x_1), \dots, \text{sign}(x_N))^T \in \{-1, 0, 1\}^N$ as the vector whose i^{th} component is equal to

$$\text{sign}(x_i) = \begin{cases} 1 & \text{if } x_i > 0 \\ 0 & \text{if } x_i = 0 \\ -1 & \text{if } x_i < 0. \end{cases}$$

For any matrix $M \in \mathbb{R}^{n \times N}$, for all $i \in \llbracket 1, n \rrbracket$, we denote by $M_{i,\cdot}$ the i^{th} line of M . The vector $M_{i,\cdot}$ is a line vector whose j^{th} component is $M_{i,j}$. Similarly, for $j \in \llbracket 1, N \rrbracket$, we denote by $M_{\cdot,j}$ the j^{th} column of M , which is the column vector whose i^{th} component is $M_{i,j}$. For any matrix $M \in \mathbb{R}^{n \times N}$, we denote by $M^T \in \mathbb{R}^{N \times n}$ the transpose matrix of M .

We denote by Id_N the $N \times N$ identity matrix and by $\mathbf{1}_N$ the vector $(1, 1, \dots, 1)^T \in \mathbb{R}^N$. If $\lambda \in \mathbb{R}^N$ is a vector of size N , for some $N \in \mathbb{N}^*$, we denote by $\text{Diag}(\lambda)$ the $N \times N$ matrix defined by:

$$\text{Diag}(\lambda)_{i,j} = \begin{cases} \lambda_i & \text{if } i = j \\ 0 & \text{otherwise.} \end{cases}$$

If X and Y are two sets and $h : X \rightarrow Y$ is a function, for a subset $A \subset Y$, we denote by $h^{-1}(A)$ the preimage of A under f , that is

$$h^{-1}(A) = \{x \in X, h(x) \in A\}.$$

Note that this does not require the function h to be injective.

For any $n, N \in \mathbb{N}^*$ and any differentiable function $f : \mathbb{R}^n \rightarrow \mathbb{R}^N$, for all $x \in \mathbb{R}^n$, we denote by $Df(x)$ its differential at the point x , i.e. the linear application $Df(x) : \mathbb{R}^n \rightarrow \mathbb{R}^N$ satisfying, for all $h \in \mathbb{R}^n$,

$$f(x + h) = f(x) + Df(x) \cdot h + o(h).$$

If we denote by x_j and h_j the components of x and h , for $j \in \llbracket 1, n \rrbracket$, we have

$$Df(x) \cdot h = \sum_{j=1}^n \frac{\partial f}{\partial x_j}(x) h_j,$$

where for all j , $\frac{\partial f}{\partial x_j}(x) \in \mathbb{R}^N$. If $f : \mathbb{R}^n \rightarrow \mathbb{R}^N$ is a linear application, we denote by $\text{Ker } f$ the set $\{x \in \mathbb{R}^n, f(x) = 0\}$, which is a linear subset of \mathbb{R}^n .

B The lifting operator ϕ

Let us introduce the notion of ‘path’, extending the definition in Section 2.2. A path is a sequence of neurons $(v_k, v_{k+1}, \dots, v_l) \in V_k \times V_{k+1} \times \dots \times V_l$, for integers k, l satisfying $0 \leq k \leq l \leq L$. In particular, for all $l \in \llbracket 0, L - 1 \rrbracket$, the set \mathcal{P}_l defined in Section 2.2 contains all the paths starting from layer l and ending in layer $L - 1$. We recall

$$\mathcal{P} = \left(\bigcup_{l=0}^{L-1} \mathcal{P}_l \right) \cup \{\beta\}.$$

If $k, l, m \in \mathbb{N}$ are three integers satisfying $0 \leq k < l \leq m \leq L$, and $p = (v_k, \dots, v_{l-1}) \in V_k \times \dots \times V_{l-1}$ and $p' = (v_l, \dots, v_m) \in V_l \times \dots \times V_m$ are two paths such that p ends in the layer preceding the starting layer of p' , we define the union of the paths by

$$p \cup p' = (v_k, \dots, v_{l-1}, v_l, \dots, v_m) \in V_k \times \dots \times V_m.$$

Before proving Proposition 1, let us compare briefly our construction to [39]. The lifting operator ϕ introduced in Section 2.2 is similar to the operator Φ in [39], except that Φ does not take a matrix form. The operator $\alpha(x, \theta)$ introduced in Section 2.2 corresponds partly to the object $\bar{\alpha}(\theta, x)$ in [39]. One of the differences is that $\bar{\alpha}(\theta, x)$ does not include any product with x_{v_0} in its entries, as does $\alpha(x, \theta)$. Finally, a similar statement to Proposition 1 and a similar proof can be found in [39]. However, one of the present contributions is to simplify the construction.

Let us now prove Proposition 1, which we restate here.

Proposition 13. For all $\theta \in \mathbb{R}^E \times \mathbb{R}^B$ and all $x \in \mathbb{R}^{V_0}$,

$$f_\theta(x)^T = \alpha(x, \theta)\phi(\theta).$$

Proof. Let us prove first the following expression, for all $v_L \in V_L$:

$$\begin{aligned} f_\theta(x)_{v_L} = & \left(\sum_{v_0 \in V_0} x_{v_0} w_{v_0 \rightarrow v_1} \prod_{l=1}^{L-1} a_{v_l}(x, \theta) w_{v_l \rightarrow v_{l+1}} \right) \\ & \vdots \\ & v_{L-1} \in V_{L-1} \\ & + \left(\sum_{l=1}^{L-1} \sum_{v_l \in V_l} b_{v_l} \prod_{l'=l}^{L-1} a_{v_{l'}}(x, \theta) w_{v_{l'} \rightarrow v_{l'+1}} \right) + b_{v_L}. \end{aligned} \quad (15)$$

We prove this by induction on the number L of layers of the network.

Initialization ($L = 2$). Let $v_2 \in V_2$.

$$\begin{aligned} f_\theta(x)_{v_2} &= (W_2)_{v_2, :} \sigma(W_1 x + b_1) + b_{v_2} \\ &= \left(\sum_{v_1 \in V_1} w_{v_1 \rightarrow v_2} [\sigma(W_1 x + b_1)]_{v_1} \right) + b_{v_2} \\ &= \left(\sum_{v_1 \in V_1} w_{v_1 \rightarrow v_2} \sigma((W_1)_{v_1, :} x + b_{v_1}) \right) + b_{v_2} \\ &= \left(\sum_{v_1 \in V_1} w_{v_1 \rightarrow v_2} a_{v_1}(x, \theta) \left(\sum_{v_0 \in V_0} w_{v_0 \rightarrow v_1} x_{v_0} + b_{v_1} \right) \right) + b_{v_2} \\ &= \left(\sum_{\substack{v_0 \in V_0 \\ v_1 \in V_1}} w_{v_1 \rightarrow v_2} a_{v_1}(x, \theta) w_{v_0 \rightarrow v_1} x_{v_0} \right) + \left(\sum_{v_1 \in V_1} w_{v_1 \rightarrow v_2} a_{v_1}(x, \theta) b_{v_1} \right) + b_{v_2} \\ &= \left(\sum_{\substack{v_0 \in V_0 \\ v_1 \in V_1}} x_{v_0} w_{v_0 \rightarrow v_1} a_{v_1}(x, \theta) w_{v_1 \rightarrow v_2} \right) + \left(\sum_{v_1 \in V_1} b_{v_1} a_{v_1}(x, \theta) w_{v_1 \rightarrow v_2} \right) + b_{v_2} \end{aligned}$$

which proves (15), when $L = 2$.

Now let $L \geq 3$ and suppose (15) holds for all ReLU networks with $L - 1$ layers. Let us consider a network with L layers.

Let us denote by $g_\theta(x)$ the output of the $L - 1$ first layers of the network pre-activation (before applying the ReLUs of the layer $L - 1$). The function g_θ is that of a ReLU network with $L - 1$ layers, and we have

$$f_\theta(x) = W_L \sigma(g_\theta(x)) + b_L.$$

Let $v_L \in V_L$. We thus have

$$f_\theta(x)_{v_L} = \sum_{v_{L-1} \in V_{L-1}} w_{v_{L-1} \rightarrow v_L} \sigma(g_\theta(x)_{v_{L-1}}) + b_{v_L}. \quad (16)$$

By the induction hypothesis, for all $v_{L-1} \in V_{L-1}$, $g_\theta(x)_{v_{L-1}}$ can be expressed with (15). Considering that $\sigma(g_\theta(x)_{v_{L-1}}) = a_{v_{L-1}}(x, \theta) g_\theta(x)_{v_{L-1}}$ and replacing $g_\theta(x)_{v_{L-1}}$ by its expression using (15), (16) becomes

$$\begin{aligned} f_\theta(x)_{v_L} &= \sum_{v_{L-1} \in V_{L-1}} w_{v_{L-1} \rightarrow v_L} a_{v_{L-1}}(x, \theta) \left[\left(\sum_{v_0 \in V_0} x_{v_0} w_{v_0 \rightarrow v_1} \prod_{l=1}^{L-2} a_{v_l}(x, \theta) w_{v_l \rightarrow v_{l+1}} \right) \right. \\ &\quad \vdots \\ &\quad \left. + \left(\sum_{l=1}^{L-2} \sum_{v_l \in V_l} b_{v_l} \prod_{l'=l}^{L-2} a_{v_{l'}}(x, \theta) w_{v_{l'} \rightarrow v_{l'+1}} \right) + b_{v_{L-1}} \right] + b_{v_L} \\ &\quad \vdots \\ &\quad \left. + \left(\sum_{v_{L-2} \in V_{L-2}} w_{v_{L-2} \rightarrow v_L} a_{v_{L-1}}(x, \theta) x_{v_0} w_{v_0 \rightarrow v_1} \prod_{l=1}^{L-2} a_{v_l}(x, \theta) w_{v_l \rightarrow v_{l+1}} \right) \right. \\ &\quad \vdots \\ &\quad \left. + \left(\sum_{l=1}^{L-2} \sum_{v_l \in V_l} w_{v_{L-1} \rightarrow v_L} a_{v_{L-1}}(x, \theta) b_{v_l} \prod_{l'=l}^{L-2} a_{v_{l'}}(x, \theta) w_{v_{l'} \rightarrow v_{l'+1}} \right) \right. \\ &\quad \vdots \\ &\quad \left. + \left(\sum_{v_{L-1} \in V_{L-1}} w_{v_{L-1} \rightarrow v_L} a_{v_{L-1}}(x, \theta) b_{v_{L-1}} \right) + b_{v_L} \right. \\ &= \left(\sum_{v_0 \in V_0} x_{v_0} w_{v_0 \rightarrow v_1} \prod_{l=1}^{L-1} a_{v_l}(x, \theta) w_{v_l \rightarrow v_{l+1}} \right) \\ &\quad \vdots \\ &\quad \left. + \left(\sum_{l=1}^{L-1} \sum_{v_l \in V_l} b_{v_l} \prod_{l'=l}^{L-1} a_{v_{l'}}(x, \theta) w_{v_{l'} \rightarrow v_{l'+1}} \right) + b_{v_L}, \right. \\ &\quad \vdots \\ &\quad \left. + \left(\sum_{v_{L-1} \in V_{L-1}} w_{v_{L-1} \rightarrow v_L} a_{v_{L-1}}(x, \theta) b_{v_{L-1}} \right) + b_{v_L}, \right. \\ &= \left(\sum_{v_0 \in V_0} x_{v_0} w_{v_0 \rightarrow v_1} \prod_{l=1}^{L-1} a_{v_l}(x, \theta) w_{v_l \rightarrow v_{l+1}} \right) \\ &\quad \vdots \\ &\quad \left. + \left(\sum_{l=1}^{L-1} \sum_{v_l \in V_l} b_{v_l} \prod_{l'=l}^{L-1} a_{v_{l'}}(x, \theta) w_{v_{l'} \rightarrow v_{l'+1}} \right) + b_{v_L}, \right. \\ &\quad \vdots \\ &\quad \left. + \left(\sum_{v_{L-1} \in V_{L-1}} w_{v_{L-1} \rightarrow v_L} a_{v_{L-1}}(x, \theta) b_{v_{L-1}} \right) + b_{v_L}, \right. \end{aligned}$$

which proves (15) holds for ReLU networks with L layers. This ends the induction, and we conclude that (15) holds for all ReLU networks.

We can now use this expression to prove Proposition 13. The first sum in (15) is taken over all the paths $p = (v_0, \dots, v_{L-1}) \in \mathcal{P}_0$, and each summand can be written as

$$x_{v_0} w_{v_0 \rightarrow v_1} \prod_{l=1}^{L-1} a_{v_l}(x, \theta) w_{v_l \rightarrow v_{l+1}} = \left(x_{v_0} \prod_{l=1}^{L-1} a_{v_l}(x, \theta) \right) \left(\prod_{l=0}^{L-1} w_{v_l \rightarrow v_{l+1}} \right) = \alpha_p(x, \theta) \phi_{p, v_L}(\theta).$$

For all $l \in \llbracket 1, L-1 \rrbracket$, the inner sum of the double sum in (15) is taken over all the paths $p = (v_l, \dots, v_{L-1}) \in \mathcal{P}_l$, and each summand can be written as

$$b_{v_l} \prod_{l'=l}^{L-1} a_{v_{l'}}(x, \theta) w_{v_{l'} \rightarrow v_{l'+1}} = \left(\prod_{l'=l}^{L-1} a_{v_{l'}}(x, \theta) \right) \left(b_{v_l} \prod_{l'=l}^{L-1} w_{v_{l'} \rightarrow v_{l'+1}} \right) = \alpha_p(x, \theta) \phi_{p, v_L}(\theta).$$

And finally, we can also write

$$b_{v_L} = \alpha_\beta(x, \theta) \phi_{\beta, v_L}(\theta).$$

Joining all these sums and denoting $\phi_{\cdot, v_L}(\theta) = (\phi_{p, v_L}(\theta))_{p \in \mathcal{P}} \in \mathbb{R}^{\mathcal{P}}$, we have

$$f_\theta(x)_{v_L} = \sum_{p \in \mathcal{P}} \alpha_p(x, \theta) \phi_{p, v_L}(\theta) = \alpha(x, \theta) \phi_{\cdot, v_L}(\theta),$$

so in other words,

$$f_\theta(x)^T = \alpha(x, \theta) \phi(\theta).$$

□

We restate here and prove Proposition 2.

Proposition 14. *For all $n \in \mathbb{N}^*$, for all $X \in \mathbb{R}^{n \times V_0}$, the mapping*

$$\begin{aligned} \alpha_X : \mathbb{R}^E \times \mathbb{R}^B &\longrightarrow \mathbb{R}^{n \times \mathcal{P}} \\ \theta &\longmapsto \alpha(X, \theta) \end{aligned}$$

appearing in (3) is piecewise-constant, with a finite number of pieces. Furthermore, the boundary of each piece has Lebesgue measure zero. We call Δ_X the union of all the boundaries. The set Δ_X is closed and has Lebesgue measure zero.

Proof. Let us first notice that for any $i \in \llbracket 1, n \rrbracket$, for any $l \in \llbracket 1, L-1 \rrbracket$,

$$(a_v(x^i, \theta))_{v \in V_1 \cup \dots \cup V_{l-1}} \in \{0, 1\}^{V_1 \cup \dots \cup V_{l-1}}$$

takes at most $2^{N_1 + \dots + N_{l-1}}$ distinct values, so the mapping $\theta \mapsto (a_v(x^i, \theta))_{v \in V_1 \cup \dots \cup V_{l-1}}$ is piecewise constant, with a finite number of pieces.

Let $i \in \llbracket 1, n \rrbracket$. Let $l \in \llbracket 1, L-1 \rrbracket$ and $v \in V_l$. Recall the definition of f_{l-1} , as given in Section 2.1. The function $\theta \rightarrow a_v(x^i, \theta)$ takes only two values, 1 or 0, and its values are determined by the sign of

$$\sum_{v' \in V_{l-1}} w_{v' \rightarrow v} f_{l-1}(x^i)_{v'} + b_v. \quad (17)$$

For all $v' \in V_{l-1}$, the value of $f_{l-1}(x^i)_{v'}$ depends on θ . On a piece $P \subset \mathbb{R}^E \times \mathbb{R}^B$ such that $(a_{v''}(x^i, \theta))_{v'' \in V_1 \cup \dots \cup V_{l-1}}$ is constant, this dependence is polynomial. Thus, on P , the value of (17) is a polynomial function of θ , and since the coefficient applied to b_v is equal to 1, the corresponding polynomial is non constant. Since the values of $a_v(x^i, \theta)$ are determined by the sign of (17), inside P , the boundary between $\{\theta \in \mathbb{R}^E \times \mathbb{R}^B, a_v(x^i, \theta) = 0\}$ and $\{\theta \in \mathbb{R}^E \times \mathbb{R}^B, a_v(x^i, \theta) = 1\}$ is included in the set of θ for which (17) equals 0. This piece of boundary is thus contained in a level set of a non constant polynomial, whose Lebesgue measure is zero.

Since there is a finite number of pieces P , the Lebesgue measure of the boundary between $\{\theta \in \mathbb{R}^E \times \mathbb{R}^B, a_v(x^i, \theta) = 0\}$ and $\{\theta \in \mathbb{R}^E \times \mathbb{R}^B, a_v(x^i, \theta) = 1\}$, which is contained in the union of the boundaries on all the pieces P , is thus equal to 0.

Since this is true for all $l \in \llbracket 1, L-1 \rrbracket$ and all $v \in V_l$, the boundary of a piece over which $(a_v(x^i, \theta))_{v \in V_1 \cup \dots \cup V_{L-1}}$ is constant also has Lebesgue measure zero.

Now since, for all x^i , the value of $\alpha(x^i, \theta)$ only depends on $(a_v(x^i, \theta))_{v \in V_1 \cup \dots \cup V_{L-1}}$ and since $\alpha_X(\theta)$ is a matrix whose lines are the vectors $\alpha(x^i, \theta)$, we can conclude that

$\alpha_X : \mathbb{R}^E \times \mathbb{R}^B \longrightarrow \mathbb{R}^{n \times \mathcal{P}}$
 $\theta \longmapsto \alpha(X, \theta)$ is piecewise-constant, with a finite number of pieces, and that the boundary of each piece has Lebesgue measure zero.

A boundary is, by definition, closed. Finally, a finite union of closed sets with Lebesgue measure 0, as Δ_X is, is closed and has Lebesgue measure 0. □

For convenience, we introduce the two following notations. Let $l \in \llbracket 0, L \rrbracket$. For any $l' \in \llbracket 0, l \rrbracket$ and any path $p_i = (v_{l'}, \dots, v_l) \in V_{l'} \times \dots \times V_l$, we denote

$$\theta_{p_i} = \begin{cases} \prod_{k=0}^{l'-1} w_{v_k \rightarrow v_{k+1}} & \text{if } l' = 0 \\ b_{l'} \prod_{k=l'}^{l-1} w_{v_k \rightarrow v_{k+1}} & \text{if } l' \geq 1, \end{cases} \quad (18)$$

where as a classic convention, an empty product is equal to 1. In particular, if $l = 0$, for any $p_i = (v_0) \in V_0$, we have $\theta_{p_i} = 1$. For any path $p_o = (v_l, \dots, v_L) \in V_l \times \dots \times V_L$, we denote

$$\theta_{p_o} = \prod_{k=l}^{L-1} w_{v_k \rightarrow v_{k+1}}, \quad (19)$$

with again the convention that an empty product is equal to 1, so if $l = L$, $\theta_{p_o} = 1$.

Some attention must be paid to the fact that for any $l' \in \llbracket 1, L \rrbracket$, if we take p_i in the case $l = L$ and p_o in the case $l = l'$, it is possible to have

$$p_i = (v_{l'}, \dots, v_L) = p_o,$$

but in that case we DO NOT have $\theta_{p_i} = \theta_{p_o}$, since $\theta_{p_i} = b_{l'} \prod_{k=l'}^{L-1} w_{v_k \rightarrow v_{k+1}}$ and $\theta_{p_o} = \prod_{k=l'}^{L-1} w_{v_k \rightarrow v_{k+1}}$. We will always denote the paths p_i and p_o with an i (as in ‘input’) or an o (as in ‘output’) to clarify which definition is used.

When considering another parameterization $\tilde{\theta} \in \mathbb{R}^E \times \mathbb{R}^B$, we denote by $\tilde{\theta}_{p_i}$ and $\tilde{\theta}_{p_o}$ the corresponding objects.

We establish different characterizations of the set S defined in Section 2.3 that will be useful in the proofs. As mentioned in Section 2.3, the subset of parameters $(\mathbb{R}^E \times \mathbb{R}^B) \setminus S$ is close to the notion of ‘admissible’ parameter in [39], but is slightly larger since the condition $w_{\bullet \rightarrow v} \neq 0$ is replaced by $(w_{\bullet \rightarrow v}, b_v) \neq (0, 0)$, for each hidden neuron v .

Proposition 15. *Let $\theta \in \mathbb{R}^E \times \mathbb{R}^B$. The following statements are equivalent.*

- i) $\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S$.
- ii) For all $l \in \llbracket 1, L-1 \rrbracket$ and all $v_l \in V_l$, there exist $l' \in \llbracket 0, l \rrbracket$, a path $p_i = (v_{l'}, \dots, v_l) \in V_{l'} \times \dots \times V_l$ and a path $p_o = (v_l, \dots, v_L) \in V_l \times \dots \times V_L$ such that

$$\theta_{p_i} \neq 0 \quad \text{and} \quad \theta_{p_o} \neq 0.$$
- iii) For all $l \in \llbracket 1, L-1 \rrbracket$ and all $v_l \in V_l$, there exist $l' \in \llbracket 0, l \rrbracket$, a path $p = (v_{l'}, \dots, v_l, \dots, v_{L-1}) \in \mathcal{P}_{l'}$ and $v_L \in V_L$ such that

$$\phi_{p, v_L}(\theta) \neq 0.$$

Proof. Let us show successively that i) \Rightarrow ii), ii) \Rightarrow iii) and iii) \Rightarrow i).

i) \rightarrow ii) Let $\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S$. Let us show ii) holds.

Let $l \in \llbracket 1, L \rrbracket$ and $v_l \in V_l$. To form a path p_i satisfying the condition, we follow the procedure:

```

 $p_i \leftarrow (v_l)$ 
 $k \leftarrow l$ 
while  $k \geq 1$  and  $b_k = 0$  do
   $\exists v_{k-1} \in V_{k-1}, w_{v_{k-1} \rightarrow v_k} \neq 0$ 
   $p_i \leftarrow (v_{k-1}, p_i)$ 
   $k \leftarrow k - 1$ 
end while
 $l' \leftarrow k$ 

```

The existence of v_{k-1} in the loop is guaranteed by the fact that $\theta \notin S$ and $b_k = 0$ in the condition of the while loop. In the end, we obtain a path $p_i = (v_{l'}, \dots, v_l)$ with either $l' > 0$ and $b_{l'} \neq 0$, or $l' = 0$. In both cases, we have by construction

$$\theta_{p_i} \neq 0.$$

We do similarly the other way to form a path $p_o = (v_l, \dots, v_L)$. We follow the procedure:

$p_o \leftarrow (v_l)$
 $k \leftarrow l$
while $k \leq L - 1$ **do**
 $\exists v_{k+1} \in V_{k+1}, w_{v_k \rightarrow v_{k+1}} \neq 0$
 $p_o \leftarrow (p_o, v_{k+1})$
 $k \leftarrow k + 1$
end while

The existence of v_{k+1} in the loop is guaranteed by the fact that $\theta \notin S$. In the end, we obtain a path $p_o = (v_l, \dots, v_L)$ satisfying by construction

$$\theta_{p_o} \neq 0.$$

ii) \rightarrow iii) Let $l \in \llbracket 1, L - 1 \rrbracket$ and $v_l \in V_l$. There exist $l' \in \llbracket 0, l \rrbracket$, a path $p_i = (v_{l'}, \dots, v_l) \in V_{l'} \times \dots \times V_l$ and a path $p_o = (v_l, \dots, v_L) \in V_l \times \dots \times V_L$ such that

$$\theta_{p_i} \neq 0 \quad \text{and} \quad \theta_{p_o} \neq 0.$$

Denoting $p = (v_{l'}, \dots, v_l, \dots, v_{L-1})$, we have

$$\phi_{p, v_L}(\theta) = \theta_{p_i} \theta_{p_o} \neq 0.$$

iii) \rightarrow i) Let us show the contrapositive: let $\theta \in S$, and let us show the statement iii) is not true. Indeed, if $\theta \in S$, there exist $l \in \llbracket 1, L - 1 \rrbracket$ and $v_l \in V_l$ such that $(w_{\bullet \rightarrow v_l}, b_{v_l}) = (0, 0)$ or $w_{v_l \rightarrow \bullet} = 0$. Consider a path $p = (v_{l'}, \dots, v_l, \dots, v_{L-1})$ and $v_L \in V_L$. We have

$$\phi_{p, v_L}(\theta) = \begin{cases} b_{v_{l'}} w_{v_{l'} \rightarrow v_{l'+1}} \dots w_{v_{l-1} \rightarrow v_l} w_{v_l \rightarrow v_{l+1}} \dots w_{v_{L-1} \rightarrow v_L} & \text{if } l' \geq 1 \\ w_{v_0 \rightarrow v_1} \dots w_{v_{l-1} \rightarrow v_l} w_{v_l \rightarrow v_{l+1}} \dots w_{v_{L-1} \rightarrow v_L} & \text{if } l' = 0. \end{cases}$$

If $(w_{\bullet \rightarrow v_l}, b_{v_l}) = (0, 0)$, either $l' = l$ and $b_{v_{l'}} = 0$ so $\phi_{p, v_L}(\theta) = 0$, or $l' < l$ and since $w_{v_{l-1} \rightarrow v_l} = 0$, we have $\phi_{p, v_L}(\theta) = 0$.

If $w_{v_l \rightarrow \bullet} = 0$, $w_{v_l \rightarrow v_{l+1}} = 0$ so $\phi_{p, v_L}(\theta) = 0$. Thus iii) is not satisfied. \square

We restate and prove Proposition 4.

Proposition 16. For all $\theta, \tilde{\theta} \in \mathbb{R}^E \times \mathbb{R}^B$, we have

$$\theta \stackrel{R}{\sim} \tilde{\theta} \implies \phi(\theta) = \phi(\tilde{\theta}),$$

and thus in particular

$$\theta \sim \tilde{\theta} \implies \phi(\theta) = \phi(\tilde{\theta}).$$

Proof. Let $\theta, \tilde{\theta} \in \mathbb{R}^E \times \mathbb{R}^B$ such that $\theta \stackrel{R}{\sim} \tilde{\theta}$. There exists a family $(\lambda^0, \dots, \lambda^L) \in (\mathbb{R}^*)^{V_0} \times \dots \times (\mathbb{R}^*)^{V_L}$, with $\lambda^0 = \mathbf{1}_{V_0}$ and $\lambda^L = \mathbf{1}_{V_L}$, such that for all $l \in \llbracket 1, L \rrbracket$, for all $(v_{l-1}, v_l) \in V_{l-1} \times V_l$, (6) holds. We consider first a path $p = (v_0, \dots, v_{L-1}) \in \mathcal{P}_0$ and $v_L \in V_L$. Using (6) and the fact that $\lambda_{v_0}^0 = \lambda_{v_L}^L = 1$, we have

$$\phi_{p, v_L}(\theta) = \prod_{l=1}^L w_{v_{l-1} \rightarrow v_l} = \prod_{l=1}^L \frac{\lambda_{v_l}^l}{\lambda_{v_{l-1}}^{l-1}} \tilde{w}_{v_{l-1} \rightarrow v_l} = \frac{\lambda_{v_L}^L}{\lambda_{v_0}^0} \prod_{l=1}^L \tilde{w}_{v_{l-1} \rightarrow v_l} = \phi_{p, v_L}(\tilde{\theta}).$$

Similarly, for $l \in \llbracket 1, L - 1 \rrbracket$ and a path $p = (v_l, \dots, v_{L-1}) \in \mathcal{P}_l$, and for all $v_L \in V_L$, we have, using (6) and the fact that $\lambda_{v_L}^L = 1$,

$$\begin{aligned} \phi_{p, v_L}(\theta) &= b_{v_l} \prod_{l'=l+1}^L w_{v_{l'-1} \rightarrow v_{l'}} = \lambda_{v_l}^l \tilde{b}_{v_l} \prod_{l'=l+1}^L \frac{\lambda_{v_{l'}}^{l'}}{\lambda_{v_{l'-1}}^{l'-1}} \tilde{w}_{v_{l'-1} \rightarrow v_{l'}} = \lambda_{v_L}^L \tilde{b}_{v_l} \prod_{l'=l+1}^L \tilde{w}_{v_{l'-1} \rightarrow v_{l'}} \\ &= \phi_{p, v_L}(\tilde{\theta}). \end{aligned}$$

Finally, for $p = \beta$ and $v_L \in V_L$, we have

$$\phi_{p, v_L}(\theta) = b_{v_L} = \lambda_{v_L}^L \tilde{b}_{v_L} = \tilde{b}_{v_L} = \phi_{p, v_L}(\tilde{\theta}).$$

This shows $\phi(\theta) = \phi(\tilde{\theta})$.

For the second implication, we simply use the fact that if $\theta \sim \tilde{\theta}$, in particular, $\theta \stackrel{R}{\sim} \tilde{\theta}$. \square

Corollary 17. The set $(\mathbb{R}^E \times \mathbb{R}^B) \setminus S$ is stable by rescaling equivalence: if $\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S$, and $\tilde{\theta} \in \mathbb{R}^E \times \mathbb{R}^B$ satisfies $\theta \stackrel{R}{\sim} \tilde{\theta}$, then $\tilde{\theta} \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S$.

Proof. Let $\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S$ and $\tilde{\theta} \in \mathbb{R}^E \times \mathbb{R}^B$ such that $\theta \stackrel{R}{\sim} \tilde{\theta}$. Proposition 16 shows that $\phi(\tilde{\theta}) = \phi(\theta)$.

Let $l \in \llbracket 1, L \rrbracket$ and $v \in V_l$. Since $\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S$, according to Proposition 15 there exists $l' \in \llbracket 0, l \rrbracket$, a path $p = (v_{l'}, \dots, v_l, \dots, v_{L-1})$ and $v_L \in V_L$ such that $\phi_{p, v_L}(\theta) \neq 0$. We have

$$\phi_{p, v_L}(\tilde{\theta}) = \phi_{p, v_L}(\theta) \neq 0,$$

and since this is true for any $l \in \llbracket 1, L \rrbracket$ and $v \in V_l$, Proposition 15 shows that $\tilde{\theta} \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S$. \square

We restate and prove Proposition 5.

Proposition 18. For all $\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S$, for all $\tilde{\theta} \in \mathbb{R}^E \times \mathbb{R}^B$,

$$\phi(\theta) = \phi(\tilde{\theta}) \implies \theta \stackrel{R}{\sim} \tilde{\theta}.$$

Proof. Let us choose $(\lambda^0, \dots, \lambda^L) \in (\mathbb{R}^*)^{V_0} \times \dots \times (\mathbb{R}^*)^{V_L}$ as follows. For all $l \in \llbracket 1, L-1 \rrbracket$ and all $v_l \in V_l$, since $\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S$, Proposition 15 shows that there exists a path $p_o(v_l) = (v_l, \dots, v_L) \in V_l \times \dots \times V_L$ such that $\theta_{p_o(v_l)} \neq 0$. Let us define $\lambda^0 = \mathbf{1}_{V_0}$, $\lambda^L = \mathbf{1}_{V_L}$ and for all $l \in \llbracket 1, L-1 \rrbracket$,

$$\lambda_{v_l}^l = \frac{\tilde{\theta}_{p_o(v_l)}}{\theta_{p_o(v_l)}}.$$

The value of $\lambda_{v_l}^l$ a priori depends on the choice of the path $p_o(v_l)$, but the first of the two following facts, that we are going to prove, shows it only depends on v_l , since in (20), p_i does not depend on $p_o(v_l)$.

- For all $l \in \llbracket 0, L \rrbracket$, for all $v_l \in V_l$, for any $l' \in \llbracket 0, l \rrbracket$ and any $p_i = (v_{l'}, \dots, v_l) \in V_{l'} \times \dots \times V_l$,

$$\theta_{p_i} = \lambda_{v_l}^l \tilde{\theta}_{p_i}. \quad (20)$$

- For all $l \in \llbracket 0, L \rrbracket$, for all $v_l \in V_l$, $\lambda_{v_l}^l \neq 0$.

Indeed, let $l \in \llbracket 0, L \rrbracket$ and let us consider $l' \in \llbracket 0, l \rrbracket$ and a path $p_i = (v_{l'}, \dots, v_l) \in V_{l'} \times \dots \times V_l$. Let $v_{l+1}, \dots, v_L \in V_{l+1} \times \dots \times V_L$ such that $p_o(v_l) = (v_l, v_{l+1}, \dots, v_L)$. Let $p = (v_{l'}, \dots, v_l, \dots, v_{L-1}) \in \mathcal{P}_{l'}$ so that $p_i \cup p_o(v_l) = p \cup (v_L)$. We have by hypothesis

$$\theta_{p_i} \theta_{p_o(v_l)} = \phi_{p, v_L}(\theta) = \phi_{p, v_L}(\tilde{\theta}) = \tilde{\theta}_{p_i} \tilde{\theta}_{p_o(v_l)},$$

thus

$$\theta_{p_i} = \frac{\tilde{\theta}_{p_o(v_l)}}{\theta_{p_o(v_l)}} \tilde{\theta}_{p_i} = \lambda_{v_l}^l \tilde{\theta}_{p_i},$$

which proves the first point. To prove the second point, we simply use Proposition 15 to consider a path p_i such that $\theta_{p_i} \neq 0$, and (20) shows that $\lambda_{v_l}^l \neq 0$.

Let us now prove the rescaling equivalence. Let $l \in \llbracket 1, L \rrbracket$, and let $(v_{l-1}, v_l) \in V_{l-1} \times V_l$. Let us consider, thanks to Proposition 15, $l' \in \llbracket 0, l-1 \rrbracket$ and a path $p_i = (v_{l'}, \dots, v_{l-1}) \in V_{l'} \times \dots \times V_{l-1}$ such that $\theta_{p_i} \neq 0$. The relation (20) shows we also have $\tilde{\theta}_{p_i} \neq 0$. Let $p'_i = p_i \cup (v_l)$. Using (20) with $\theta_{p'_i}$ we have

$$\theta_{p_i} w_{v_{l-1} \rightarrow v_l} = \theta_{p'_i} = \lambda_{v_l}^l \tilde{\theta}_{p'_i} = \lambda_{v_l}^l \tilde{\theta}_{p_i} \tilde{w}_{v_{l-1} \rightarrow v_l}.$$

At the same time, using (20) with θ_{p_i} we have,

$$\theta_{p_i} w_{v_{l-1} \rightarrow v_l} = \lambda_{v_{l-1}}^{l-1} \tilde{\theta}_{p_i} w_{v_{l-1} \rightarrow v_l},$$

so combining both equalities, we have

$$\lambda_{v_l}^l \tilde{\theta}_{p_i} \tilde{w}_{v_{l-1} \rightarrow v_l} = \lambda_{v_{l-1}}^{l-1} \tilde{\theta}_{p_i} w_{v_{l-1} \rightarrow v_l}.$$

Using the fact that $\tilde{\theta}_{p_i} \neq 0$ and $\lambda_{v_{l-1}}^{l-1} \neq 0$, we finally obtain, for all $l \in \llbracket 1, L \rrbracket$ and all $(v_{l-1}, v_l) \in V_{l-1} \times V_l$:

$$w_{v_{l-1} \rightarrow v_l} = \frac{\lambda_{v_l}^l}{\lambda_{v_{l-1}}^{l-1}} \tilde{w}_{v_{l-1} \rightarrow v_l}.$$

For all $l \in \llbracket 1, L \rrbracket$ and all $v_l \in V_l$, using (20) with $p_i = (v_l)$, we obtain

$$b_{v_l} = \lambda_{v_l}^l \tilde{b}_{v_l}.$$

This shows that (6) is satisfied for all $(v_{l-1}, v_l) \in V_{l-1} \times V_l$, and thus $\theta \stackrel{R}{\sim} \tilde{\theta}$. \square

The following proposition is useful in the proof of Theorem 26 and allows to improve identifiability modulo rescaling into identifiability modulo positive rescaling.

Proposition 19. *For all $\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S$, there exists $\epsilon > 0$ such that for all $\tilde{\theta} \in \mathbb{R}^E \times \mathbb{R}^B$,*

$$\|\theta - \tilde{\theta}\|_\infty < \epsilon \text{ and } \theta \stackrel{R}{\sim} \tilde{\theta} \implies \theta \sim \tilde{\theta}.$$

Proof. Let $\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S$. We define

$$\epsilon = \min \left(\{|w_{v \rightarrow v'}|, v \rightarrow v' \in E \text{ and } w_{v \rightarrow v'} \neq 0\} \cup \{|b_v|, v \in B \text{ and } b_v \neq 0\} \right).$$

Let $\tilde{\theta} \in \mathbb{R}^E \times \mathbb{R}^B$ such that $\|\theta - \tilde{\theta}\|_\infty < \epsilon$ and $\theta \stackrel{R}{\sim} \tilde{\theta}$. To prove $\theta \sim \tilde{\theta}$, we simply have to prove $\text{sign}(\theta) = \text{sign}(\tilde{\theta})$. There exists $(\lambda^0, \dots, \lambda^L) \in (\mathbb{R}^*)^{V_0} \times \dots \times (\mathbb{R}^*)^{V_L}$, with $\lambda^0 = \mathbf{1}_{V_0}$ and $\lambda^L = \mathbf{1}_{V_L}$, such that, for all $l \in \llbracket 1, L \rrbracket$, for all $(v_{l-1}, v_l) \in V_{l-1} \times V_l$, (6) holds. Let us show that $\text{sign}(\theta) = \text{sign}(\tilde{\theta})$.

Indeed, let $l \in \llbracket 1, L \rrbracket$, and let $(v, v') \in V_{l-1} \times V_l$. If $w_{v \rightarrow v'} \neq 0$, then since $|w_{v \rightarrow v'} - \tilde{w}_{v \rightarrow v'}| < \epsilon$ and by definition $\epsilon \leq |w_{v \rightarrow v'}|$, we have $\text{sign}(w_{v \rightarrow v'}) = \text{sign}(\tilde{w}_{v \rightarrow v'})$. Otherwise, if $w_{v \rightarrow v'} = 0$, (6) shows that we have

$$\tilde{w}_{v \rightarrow v'} = \frac{\lambda_v^{l-1}}{\lambda_{v'}^l} w_{v \rightarrow v'} = 0,$$

so we still have $\text{sign}(w_{v \rightarrow v'}) = \text{sign}(\tilde{w}_{v \rightarrow v'})$.

Now let $l \in \llbracket 1, L \rrbracket$ and let $v \in V_l$. Similarly, if $b_v \neq 0$, we have $|b_v - \tilde{b}_v| < \epsilon \leq |b_v|$, so $\text{sign}(b_v) = \text{sign}(\tilde{b}_v)$, and if $b_v = 0$, we have

$$\tilde{b}_v = \frac{b_v}{\lambda_v^l} = 0,$$

so again $\text{sign}(b_v) = \text{sign}(\tilde{b}_v)$.

This shows $\text{sign}(\theta) = \text{sign}(\tilde{\theta})$, so $\theta \sim \tilde{\theta}$. \square

C The smooth manifold structure of Σ_1^*

In this section, we prove Theorem 7, which is restated as Theorem 25. Before doing so, we establish intermediary results, some of which are evoked in Section 3.

Let us discuss the cardinal of F_θ defined in Section 3. The set F_θ is obtained by removing the edges of the form $v \rightarrow s_{\max}^\theta(v)$ for $v \in V_1 \cup \dots \cup V_{L-1}$. Note that we do not remove the edges of the form $v \rightarrow s_{\max}^\theta(v)$ for $v \in V_0$. For all $l \in \llbracket 1, L-1 \rrbracket$, there are precisely N_l edges of the form $(v, s_{\max}^\theta(v))$ with $v \in V_l$, so

$$\begin{aligned} |F_\theta| &= |E| - (N_1 + \dots + N_{L-1}) \\ &= N_0 N_1 + \dots + N_{L-1} N_L - N_1 - \dots - N_{L-1}. \end{aligned}$$

As a consequence, since $|B| = N_1 + \dots + N_L$, we have in particular

$$\begin{aligned} |F_\theta| + |B| &= N_0 N_1 + \dots + N_{L-1} N_L - N_1 - \dots - N_{L-1} + N_1 + \dots + N_L \\ &= N_0 N_1 + \dots + N_{L-1} N_L + N_L. \end{aligned} \quad (21)$$

The following proposition is a first step towards Proposition 21, which states that ψ^θ is a homeomorphism.

Proposition 20. *For all $\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S$, the function $\psi^\theta : U_\theta \rightarrow \mathbb{R}^{\mathcal{P} \times V_L}$ is injective.*

Proof. Let $\tau, \tilde{\tau} \in U_\theta$ such that $\psi^\theta(\tau) = \psi^\theta(\tilde{\tau})$. Let us show $\tau = \tilde{\tau}$. We have $\phi(\rho_\theta(\tau)) = \phi(\rho_\theta(\tilde{\tau}))$ and by definition of U_θ , $\rho_\theta(\tau) \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S$, so by Proposition 18 we have the rescaling equivalence

$$\rho_\theta(\tau) \stackrel{R}{\sim} \rho_\theta(\tilde{\tau}).$$

By definition of the rescaling equivalence, in its formulation (6), there exists $(\lambda^0, \dots, \lambda^L) \in (\mathbb{R}^*)^{V_0} \times \dots \times (\mathbb{R}^*)^{V_L}$, with $\lambda^0 = \mathbf{1}_{V_0}$ and $\lambda^L = \mathbf{1}_{V_L}$, such that, for all $l \in \llbracket 1, L \rrbracket$, for all $(v_{l-1}, v_l) \in V_{l-1} \times V_l$,

$$\begin{cases} \rho_\theta(\tau)_{v_{l-1} \rightarrow v_l} = \frac{(\lambda^l)_{v_l}}{(\lambda^{l-1})_{v_{l-1}}} \rho_\theta(\tilde{\tau})_{v_{l-1} \rightarrow v_l} \\ b_{v_l} = \lambda_{v_l}^l \tilde{b}_{v_l}. \end{cases} \quad (22)$$

Let $l \in \llbracket 2, L \rrbracket$ and let $v_{l-1} \in V_{l-1}$. Let $v_l = s_{\max}^\theta(v_{l-1})$. According to (22) we have

$$\rho_\theta(\tau)_{v_{l-1} \rightarrow v_l} = \frac{(\lambda^l)_{v_l}}{(\lambda^{l-1})_{v_{l-1}}} \rho_\theta(\tilde{\tau})_{v_{l-1} \rightarrow v_l}.$$

But since $v_l = s_{\max}^\theta(v_{l-1})$ and $v_{l-1} \in V_{l-1}$ with $l-1 \in \llbracket 1, L-1 \rrbracket$, we have $v_{l-1} \rightarrow v_l \in E \setminus F_\theta$, so by definition of ρ_θ in (9),

$$\rho_\theta(\tau)_{v_{l-1} \rightarrow v_l} = w_{v_{l-1} \rightarrow v_l} = \rho_\theta(\tilde{\tau})_{v_{l-1} \rightarrow v_l} \neq 0,$$

so $\frac{(\lambda^l)_{v_l}}{(\lambda^{l-1})_{v_{l-1}}} = 1$.

We have shown that for all $l \in \llbracket 2, L \rrbracket$, for all $v_{l-1} \in V_{l-1}$, there exists $v_l \in V_l$ such that

$$(\lambda^{l-1})_{v_{l-1}} = (\lambda^l)_{v_l}.$$

As a consequence, if l is such that $\lambda^l = \mathbf{1}_{V_l}$, then $\lambda^{l-1} = \mathbf{1}_{V_{l-1}}$.

Starting from $\lambda^L = \mathbf{1}_{V_L}$, this shows by induction that for all $l \in \llbracket 1, L \rrbracket$,

$$\lambda^l = \mathbf{1}_{V_l}.$$

By hypothesis we also have $\lambda^0 = \mathbf{1}_{V_0}$. Using (22), this shows that

$$\rho_\theta(\tau) = \rho_\theta(\tilde{\tau}).$$

The injectivity of ρ_θ allows us to conclude that

$$\tau = \tilde{\tau}.$$

□

The following proposition shows, as mentioned in Section 3, that ψ^θ is a homeomorphism. This is a necessary step to prove that $(V_\theta, (\psi^\theta)^{-1})_{\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S}$ is an atlas of Σ_1^* .

Proposition 21. *For all $\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S$, ψ^θ is a homeomorphism from U_θ onto its image V_θ .*

Proof. We already know from Proposition 20 that ψ^θ is injective, so we need to prove that ψ^θ is continuous and its inverse is continuous. The function ρ_θ is affine and ϕ is a polynomial function, so the function $\psi^\theta = \phi \circ \rho_\theta$ is a polynomial function, and in particular it is continuous.

To prove that $(\psi^\theta)^{-1}$ is continuous, we consider a sequence (τ_n) taking values in U_θ and $\tau \in U_\theta$ such that $\psi^\theta(\tau_n) \rightarrow \psi^\theta(\tau)$, and we want to show that $\tau_n \rightarrow \tau$.

Let us first show that for all $v \in B$, $(\tau_n)_v \rightarrow \tau_v$. Indeed, let $l \in \llbracket 1, L \rrbracket$ and let $v_l \in V_l$, so that v_l is an arbitrary element of B . Let us define $v_{l+1} = s_{\max}^\theta(v_l)$, then $v_{l+2} = s_{\max}^\theta(v_{l+1})$ and so on up to $v_L = s_{\max}^\theta(v_{L-1})$. Since for all $l' \in \llbracket l, L-1 \rrbracket$, $v_{l'+1} = s_{\max}^\theta(v_{l'})$, by definition of F_θ and ρ_θ (see (8) and (9)), we have

$$\rho_\theta(\tau_n)_{v_{l'} \rightarrow v_{l'+1}} = w_{v_{l'} \rightarrow v_{l'+1}}, \quad (23)$$

and

$$\rho_\theta(\tau)_{v_{l'} \rightarrow v_{l'+1}} = w_{v_{l'} \rightarrow v_{l'+1}}. \quad (24)$$

In particular, since $\theta \notin S$, for all $l' \in \llbracket l, L-1 \rrbracket$ we have $w_{v_{l'} \rightarrow \bullet} \neq 0$, so by definition of s_{\max}^θ , $w_{v_{l'} \rightarrow v_{l'+1}} \neq 0$. We thus have

$$w_{v_l \rightarrow v_{l+1}} \cdots w_{v_{L-1} \rightarrow v_L} \neq 0. \quad (25)$$

If we denote $p = (v_l, \dots, v_{L-1})$, we have, using the definition of ϕ and (23),

$$\psi_{p, v_L}^\theta(\tau_n) = (\tau_n)_{v_l} w_{v_l \rightarrow v_{l+1}} \cdots w_{v_{L-1} \rightarrow v_L}$$

and using (24),

$$\psi_{p, v_L}^\theta(\tau) = (\tau)_{v_l} w_{v_l \rightarrow v_{l+1}} \cdots w_{v_{L-1} \rightarrow v_L}.$$

Using (25) and the fact that

$$\psi^\theta(\tau_n) \rightarrow \psi^\theta(\tau),$$

we conclude that

$$(\tau_n)_{v_l} \rightarrow \tau_{v_l}.$$

Let us now prove that for all $(v, v') \in E$, $(\tau_n)_{v \rightarrow v'} \rightarrow \tau_{v \rightarrow v'}$. Let us show by induction on $l \in \llbracket 1, L \rrbracket$ the following hypothesis

$$\forall l' \in \llbracket 1, l \rrbracket, \quad \forall (v, v') \in (V_{l'-1} \times V_{l'}) \cap F_\theta, \quad (\tau_n)_{v \rightarrow v'} \rightarrow \tau_{v \rightarrow v'}. \quad (H_l)$$

Initialization. Let $(v_0, v_1) \in (V_0 \times V_1) \cap F_\theta$. We define $v_2 = s_{\max}^\theta(v_1)$, then we define $v_3 = s_{\max}^\theta(v_2)$, and so on up to $v_L = s_{\max}^\theta(v_{L-1})$. Let $p = (v_0, \dots, v_{L-1}) \in \mathcal{P}$.

As above, using the definition of ρ_θ , F_θ and ϕ , we have

$$\psi_{p, v_L}^\theta(\tau_n) = (\tau_n)_{v_0 \rightarrow v_1} w_{v_1 \rightarrow v_2} \cdots w_{v_{L-1} \rightarrow v_L}$$

and

$$\psi_{p, v_L}^\theta(\tau) = (\tau)_{v_0 \rightarrow v_1} w_{v_1 \rightarrow v_2} \cdots w_{v_{L-1} \rightarrow v_L},$$

and since $\theta \notin S$, we also have, as above,

$$w_{v_1 \rightarrow v_2} \cdots w_{v_{L-1} \rightarrow v_L} \neq 0. \quad (26)$$

Since

$$\psi^\theta(\tau_n) \rightarrow \psi^\theta(\tau)$$

we conclude using (26) that

$$(\tau_n)_{v_0 \rightarrow v_1} \rightarrow \tau_{v_0 \rightarrow v_1}.$$

We have shown H_1 .

Induction step. Let $l \in \llbracket 2, L \rrbracket$ and let us assume that H_{l-1} holds.

Let $(v_{l-1}, v_l) \in (V_{l-1} \times V_l) \cap F_\theta$. We define $v_{l+1} = s_{\max}^\theta(v_l)$, $v_{l+2} = s_{\max}^\theta(v_{l+1})$, and so on up to $v_L = s_{\max}^\theta(v_{L-1})$. Let us denote $p_o = (v_l, \dots, v_L)$. Recalling the notation defined in (19), we have

$$\rho_\theta(\tau_n)_{p_o} = w_{v_l \rightarrow v_{l+1}} \cdots w_{v_{L-1} \rightarrow v_L} = \rho_\theta(\tau)_{p_o} \neq 0. \quad (27)$$

At the same time, since $\tau \in U_\theta$, Proposition 15 shows there exist $l' \in \llbracket 0, l-1 \rrbracket$ and a path $p_i = (v_{l'}, \dots, v_{l-2}, v_{l-1})$ such that

$$\rho_\theta(\tau)_{p_i} \neq 0. \quad (28)$$

If $l' \geq 1$, we have shown in the first part of the proof that $(\tau_n)_{v_{l'}} \rightarrow \tau_{v_{l'}}$. Moreover, whatever the value of l' is, for $k \in \llbracket l', l-2 \rrbracket$, if $(v_k, v_{k+1}) \in E \setminus F_\theta$,

$$\rho_\theta(\tau_n)_{v_k \rightarrow v_{k+1}} = w_{v_k \rightarrow v_{k+1}} = \rho_\theta(\tau)_{v_k \rightarrow v_{k+1}},$$

and if $(v_k, v_{k+1}) \in F_\theta$, according to H_{l-1} ,

$$\rho_\theta(\tau_n)_{v_k \rightarrow v_{k+1}} = (\tau_n)_{v_k \rightarrow v_{k+1}} \longrightarrow \tau_{v_k \rightarrow v_{k+1}} = \rho_\theta(\tau)_{v_k \rightarrow v_{k+1}}.$$

We therefore have

$$\rho_\theta(\tau_n)_{p_i} \longrightarrow \rho_\theta(\tau)_{p_i}, \quad (29)$$

and in particular, since $\rho_\theta(\tau)_{p_i} \neq 0$, there exists $n_0 \in \mathbb{N}$ such that for all $n \geq n_0$,

$$\rho_\theta(\tau_n)_{p_i} \neq 0. \quad (30)$$

We can write

$$\psi_{p, v_L}^\theta(\tau_n) = \rho_\theta(\tau_n)_{p_i} (\tau_n)_{v_{l-1} \rightarrow v_l} \rho_\theta(\tau_n)_{p_o}$$

and

$$\psi_{p, v_L}^\theta(\tau) = \rho_\theta(\tau)_{p_i} (\tau)_{v_{l-1} \rightarrow v_l} \rho_\theta(\tau)_{p_o},$$

so using (27), (30) and (29), we have

$$(\tau_n)_{v_{l-1} \rightarrow v_l} = \frac{\psi_{p, v_L}^\theta(\tau_n)}{\rho_\theta(\tau_n)_{p_i} \rho_\theta(\tau_n)_{p_o}} \longrightarrow \frac{\psi_{p, v_L}^\theta(\tau)}{\rho_\theta(\tau)_{p_i} \rho_\theta(\tau)_{p_o}} = \tau_{v_{l-1} \rightarrow v_l}.$$

We have shown H_l , which concludes the induction step.

In particular, H_L is satisfied, and finally $\tau_n \rightarrow \tau$.

This shows that ψ^θ is a homeomorphism. \square

The following lemma is necessary for the proof of Proposition 23.

Lemma 22. Let $\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S$. Let $(v, v') \in E$ (resp. $v \in B$). If $w_{v \rightarrow v'} \neq 0$ (resp. $b_v \neq 0$), then there exists $\epsilon > 0$ such that for all $\tilde{\theta} \in \mathbb{R}^E \times \mathbb{R}^B$, if $\|\phi(\theta) - \phi(\tilde{\theta})\|_\infty < \epsilon$, then $\tilde{w}_{v \rightarrow v'} \neq 0$ (resp. $\tilde{b}_v \neq 0$).

Proof. Let $\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S$ and $(v, v') \in E$ such that $w_{v \rightarrow v'} \neq 0$. Denote $l \in \llbracket 0, L-1 \rrbracket$ such that $v \in V_l$. If $l = 0$, we take $p_i = (v)$ so that by convention $\theta_{p_i} = 1 \neq 0$, and if $l \geq 1$, we use Proposition 15 which states that there exists $l' \in \llbracket 0, l-1 \rrbracket$ and a path $p_i = (v_{l'}, \dots, v_{l-2}, v)$ such that $\theta_{p_i} \neq 0$. Similarly, if $l = L-1$, we take $p_o = (v')$ so that by convention $\theta_{p_o} = 1 \neq 0$ and if $l < L-1$, we use Proposition 15 which states that there exists a path $p_o = (v', v_{l+1}, \dots, v_L)$ such that $\theta_{p_o} \neq 0$. If we denote

$$p = \begin{cases} (v, v', v_{l+2}, \dots, v_{L-1}) & \text{if } l = 0 \\ (v_{l'}, \dots, v_{l-1}, v, v') & \text{if } l = L-1 \\ (v_{l'}, \dots, v_{l-1}, v, v', v_{l+2}, \dots, v_{L-1}) & \text{otherwise,} \end{cases}$$

we have

$$\phi_{p, v_L}(\theta) = \theta_{p_i} w_{v \rightarrow v'} \theta_{p_o} \neq 0.$$

We define $\epsilon = |\phi_{p, v_L}(\theta)| > 0$. For all $\tilde{\theta} \in \mathbb{R}^E \times \mathbb{R}^B$ such that $\|\phi(\tilde{\theta}) - \phi(\theta)\|_\infty < \epsilon$ we have

$$\phi_{p, v_L}(\tilde{\theta}) \neq 0.$$

Since $\phi_{p, v_L}(\tilde{\theta}) = \tilde{\theta}_{p_i} \tilde{w}_{v \rightarrow v'} \tilde{\theta}_{p_o}$, this implies in particular that

$$\tilde{w}_{v \rightarrow v'} \neq 0.$$

The proof is similar in the case $v \in B$ and $b_v \neq 0$. \square

The following proposition, which states that for any $\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S$, $V_\theta = \psi^\theta(U_\theta)$ is open with respect to the topology induced on Σ_1^* by the standard topology of $\mathbb{R}^{\mathcal{P} \times V_L}$, is necessary to show that $(V_\theta, (\psi^\theta)^{-1})_{\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S}$ is an atlas of Σ_1^* .

Proposition 23. For any $\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S$, for any $\tau \in U_\theta$, there exists $\epsilon > 0$ such that

$$\Sigma_1^* \cap B_\infty(\psi^\theta(\tau), \epsilon) \subset V_\theta.$$

Proof. Let us first construct ϵ and then consider an element of the set on the left of the inclusion and prove it belongs to V_θ . Let $\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S$ and $\tau \in U_\theta$. For all $l \in \llbracket 1, L-1 \rrbracket$, for all $v \in V_l$, by definition of F_θ and ρ_θ , we have $\rho_\theta(\tau)_{v \rightarrow s_{\max}^\theta(v)} = w_{v \rightarrow s_{\max}^\theta(v)}$, and since $\theta \notin S$, by definition of s_{\max}^θ , $w_{v \rightarrow s_{\max}^\theta(v)} \neq 0$, so according to Lemma 22 there exists $\epsilon_v > 0$ such that for all $\tilde{\theta} \in \mathbb{R}^E \times \mathbb{R}^B$,

$$\|\phi(\rho_\theta(\tau)) - \phi(\tilde{\theta})\|_\infty < \epsilon_v \implies \tilde{w}_{v \rightarrow s_{\max}^\theta(v)} \neq 0.$$

Let $\epsilon = \min_{v \in V_1 \cup \dots \cup V_{L-1}} \epsilon_v$.

Let us now show the inclusion: let $\tilde{\theta} \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S$ such that $\|\phi(\rho_\theta(\tau)) - \phi(\tilde{\theta})\|_\infty < \epsilon$, and let us show that $\phi(\tilde{\theta}) \in V_\theta$. Notice first that for all $l \in \llbracket 1, L-1 \rrbracket$ and $v \in V_l$, by definition of ϵ , $w_{v \rightarrow s_{\max}^\theta(v)} \neq 0$ and $\tilde{w}_{v \rightarrow s_{\max}^\theta(v)} \neq 0$. We are going to define $\tilde{\tau} \in U_\theta$ such that $\rho_\theta(\tilde{\tau}) \stackrel{R}{\sim} \tilde{\theta}$, so that using Proposition 16, $\psi^\theta(\tilde{\tau}) = \phi(\tilde{\theta})$.

Let us define recursively a family $(\lambda^0, \dots, \lambda^L) \in (\mathbb{R}^*)^{V_0} \times \dots \times (\mathbb{R}^*)^{V_L}$ as follows:

- we define $\lambda^L = \mathbf{1}_{V_L}$;
- for all $l \in \llbracket 1, L-1 \rrbracket$, for all $v \in V_l$, we define

$$\lambda_v^l = \frac{\tilde{w}_{v \rightarrow s_{\max}^\theta(v)}}{w_{v \rightarrow s_{\max}^\theta(v)}} \lambda_{s_{\max}^\theta(v)}^{l+1}. \quad (31)$$

- we define finally $\lambda^0 = \mathbf{1}_{V_0}$.

Note that for all $l \in \llbracket 0, L \rrbracket$ and for all $v \in V_l$, $\lambda_v^l \neq 0$. Also note that for all $l \in \llbracket 2, L \rrbracket$, for all $v \in V_{l-1}$, reformulating (31) in a way that will be useful later, we have

$$\frac{\lambda_{s_{\max}^\theta(v)}^l}{\lambda_v^{l-1}} = \frac{w_{v \rightarrow s_{\max}^\theta(v)}}{\tilde{w}_{v \rightarrow s_{\max}^\theta(v)}}. \quad (32)$$

We then define $\tilde{\tau} \in \mathbb{R}^{F_\theta} \times \mathbb{R}^B$ by:

- for all $l \in \llbracket 1, L \rrbracket$, for all $(v, v') \in (V_{l-1} \times V_l) \cap F_\theta$,

$$\tilde{\tau}_{v \rightarrow v'} = \frac{\lambda_{v'}^l}{\lambda_v^{l-1}} \tilde{w}_{v \rightarrow v'}; \quad (33)$$

- for all $l \in \llbracket 1, L \rrbracket$, for all $v \in V_l$,

$$\tilde{\tau}_v = \lambda_v^l \tilde{b}_v. \quad (34)$$

Let us show $\rho_\theta(\tilde{\tau}) \stackrel{R}{\sim} \tilde{\theta}$. Indeed, let $l \in \llbracket 1, L \rrbracket$ and let $(v, v') \in V_{l-1} \times V_l$. If $v \in V_0$ or $v \in V_1 \cup \dots \cup V_{L-1}$ and $v' \neq s_{\max}^\theta(v)$, then by definition (8) of F_θ , we have $v \rightarrow v' \in F_\theta$, so using (9) and (33) we have

$$\rho_\theta(\tilde{\tau})_{v \rightarrow v'} = \tilde{\tau}_{v \rightarrow v'} = \frac{\lambda_{v'}^l}{\lambda_v^{l-1}} \tilde{w}_{v \rightarrow v'}. \quad (35)$$

If $v \in V_1 \cup \dots \cup V_{L-1}$ and $v' = s_{\max}^\theta(v)$, then by definition (8) of F_θ , we have $v \rightarrow v' \in E \setminus F_\theta$, and since in that case, $l \geq 2$, using (9) and (32), we see that

$$\rho_\theta(\tilde{\tau})_{v \rightarrow v'} = w_{v \rightarrow v'} = \frac{\lambda_{v'}^l}{\lambda_v^{l-1}} \tilde{w}_{v \rightarrow v'}. \quad (36)$$

If $v \in B$, using (9) and (34), we have

$$\rho_\theta(\tilde{\tau})_v = \tilde{\tau}_v = \lambda_v^l \tilde{b}_v. \quad (37)$$

Equations (35), (36) and (37) prove that

$$\rho_\theta(\tilde{\tau}) \stackrel{R}{\sim} \tilde{\theta}.$$

Using Corollary 17, since $\tilde{\theta} \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S$ and $\rho_\theta(\tilde{\tau}) \stackrel{R}{\sim} \tilde{\theta}$, we also have $\rho_\theta(\tilde{\tau}) \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S$. Since, by definition, $U_\theta = \rho_\theta^{-1}((\mathbb{R}^E \times \mathbb{R}^B) \setminus S)$, we have $\tilde{\tau} \in U_\theta$. We have shown

$$\Sigma_1^* \cap B_\infty(\psi^\theta(\tau), \epsilon) \subset V_\theta.$$

□

The following proposition is necessary in order to show that $(V_\theta, (\psi^\theta)^{-1})_{\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S}$ is an atlas of Σ_1^* .

Proposition 24. *For all $\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S$, the function ψ^θ is C^∞ and its differential $D\psi^\theta(\tau)$ is injective for all $\tau \in U_\theta$.*

Proof. Let $\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S$. First of all, ψ^θ is a polynomial function as a composition of ϕ and ρ_θ which are both polynomial functions. So, ψ^θ is C^∞ .

In order to show the injectivity of the differential $D\psi^\theta(\tau)$ for all $\tau \in U_\theta$, let us compute the partial derivatives of $\psi_{p, v_L}^\theta(\tau)$. Let $p \in \mathcal{P}$ and $v_L \in V_L$. Using the definition of ψ^θ and ϕ , three cases are possible.

Case 1. The path p is of the form $(v_0, v_1, \dots, v_{L-1})$. We have

$$\psi_{p, v_L}^\theta(\tau) = \rho_\theta(\tau)_{v_0 \rightarrow v_1} \cdots \rho_\theta(\tau)_{v_{L-1} \rightarrow v_L}.$$

Case 2. The path p is of the form (v_l, \dots, v_{L-1}) with $l \in \llbracket 1, L-1 \rrbracket$. We have, for all $\tau \in U_\theta$,

$$\psi_{p, v_L}^\theta(\tau) = \tau_{v_l} \rho_\theta(\tau)_{v_l \rightarrow v_{l+1}} \cdots \rho_\theta(\tau)_{v_{L-1} \rightarrow v_L}.$$

Case 3. For $p = \beta$, we have, for all $\tau \in U_\theta$,

$$\psi_{p, v_L}^\theta(\tau) = \tau_{v_L}.$$

Let $(v, v') \in F_\theta$, and let us compute $\frac{\partial \psi_{p, v_L}^\theta}{\partial \tau_{v \rightarrow v'}}(\tau)$.

Case 1. We have $p = (v_0, \dots, v_{L-1}) \in \mathcal{P}_0$. If $\{v, v'\} \subset \{v_0, \dots, v_L\}$, there exists $l \in \llbracket 0, L-1 \rrbracket$ such that $(v, v') = (v_l, v_{l+1})$, in which case, since $(v, v') \in F_\theta$, $\rho_\theta(\tau)_{v_l \rightarrow v_{l+1}} = \tau_{v_l \rightarrow v_{l+1}}$ and

$$\frac{\partial \psi_{p, v_L}^\theta}{\partial \tau_{v \rightarrow v'}}(\tau) = \prod_{\substack{k \in \llbracket 0, L-1 \rrbracket \\ k \neq l}} \rho_\theta(\tau)_{v_k \rightarrow v_{k+1}}. \quad (38)$$

Otherwise if $\{v, v'\} \not\subset \{v_0, \dots, v_L\}$,

$$\frac{\partial \psi_{p, v_L}^\theta}{\partial \tau_{v \rightarrow v'}}(\tau) = 0.$$

Case 2. We have $p = (v_l, \dots, v_{L-1}) \in \mathcal{P}_l$, for $l \in \llbracket 1, L-1 \rrbracket$. If $\{v, v'\} \subset \{v_l, \dots, v_L\}$, there exists $l' \in \llbracket l, L-1 \rrbracket$ such that $(v, v') = (v_{l'}, v_{l'+1})$, in which case, since $(v, v') \in F_\theta$, $\rho_\theta(\tau)_{v_{l'} \rightarrow v_{l'+1}} = \tau_{v_{l'} \rightarrow v_{l'+1}}$ and

$$\frac{\partial \psi_{p, v_L}^\theta}{\partial \tau_{v \rightarrow v'}}(\tau) = \tau_{v_l} \prod_{\substack{k \in \llbracket l, L-1 \rrbracket \\ k \neq l'}} \rho_\theta(\tau)_{v_k \rightarrow v_{k+1}}. \quad (39)$$

Otherwise if $\{v, v'\} \not\subset \{v_l, \dots, v_L\}$,

$$\frac{\partial \psi_{p, v_L}^\theta}{\partial \tau_{v \rightarrow v'}}(\tau) = 0.$$

Case 3. We have $p = \beta$. In that case, we have

$$\frac{\partial \psi_{p,v_L}^\theta}{\partial \tau_{v \rightarrow v'}}(\tau) = 0.$$

Now let $v \in B$, and let us compute $\frac{\partial \psi_{p,v_L}^\theta}{\partial \tau_v}(\tau)$.

Case 1. We have $p = (v_0, \dots, v_{L-1}) \in \mathcal{P}_0$ and

$$\frac{\partial \psi_{p,v_L}^\theta}{\partial \tau_v}(\tau) = 0.$$

Case 2. We have $p = (v_l, \dots, v_{L-1}) \in \mathcal{P}_l$ for $l \in \llbracket 1, L-1 \rrbracket$. If $v = v_l$, then

$$\frac{\partial \psi_{p,v_L}^\theta}{\partial \tau_v}(\tau) = \prod_{k \in \llbracket l, L-1 \rrbracket} \rho_\theta(\tau)_{v_k \rightarrow v_{k+1}}.$$

If $v \neq v_l$,

$$\frac{\partial \psi_{p,v_L}^\theta}{\partial \tau_v}(\tau) = 0.$$

Case 3. We have $p = \beta$ and

$$\frac{\partial \psi_{p,v_L}^\theta}{\partial \tau_v}(\tau) = \begin{cases} 1 & \text{if } v = v_L \\ 0 & \text{if } v \neq v_L. \end{cases}$$

Now that we know the partial derivatives, let us show $D\psi^\theta(\tau)$ is injective for all $\tau \in U_\theta$. Let $\tau \in U_\theta$ and let $h \in \mathbb{R}^{F_\theta} \times \mathbb{R}^B$ such that

$$D\psi^\theta(\tau) \cdot h = 0.$$

We need to prove that $h = 0$.

Let us show first that for all $v \in B$, $h_v = 0$. Let $l \in \llbracket 1, L-1 \rrbracket$, and let $v_l \in V_l$ so that v_l is arbitrary in $B \setminus V_L$. Let us define $v_{l+1} = s_{\max}^\theta(v_l)$, then $v_{l+2} = s_{\max}^\theta(v_{l+1})$, and so on up to $v_L = s_{\max}^\theta(v_{L-1})$. Let us denote $p = (v_l, \dots, v_{L-1})$. We have

$$\psi_{p,v_L}^\theta(\tau) = \tau_{v_l} w_{v_l \rightarrow v_{l+1}} \cdots w_{v_{L-1} \rightarrow v_L},$$

so

$$[D\psi^\theta(\tau) \cdot h]_{p,v_L} = \frac{\partial \psi_{p,v_L}^\theta}{\partial \tau_{v_l}}(\tau) h_{v_l} = w_{v_l \rightarrow v_{l+1}} \cdots w_{v_{L-1} \rightarrow v_L} h_{v_l}.$$

Since $[D\psi^\theta(\tau) \cdot h]_{p,v_L} = 0$ and $w_{v_l \rightarrow v_{l+1}} \cdots w_{v_{L-1} \rightarrow v_L} \neq 0$, we conclude that $h_{v_l} = 0$. Now let $v_L \in V_L$. We consider $p = \beta$ and we have

$$[D\psi^\theta(\tau) \cdot h]_{p,v_L} = h_{v_L}.$$

Since $[D\psi^\theta(\tau) \cdot h]_{p,v_L} = 0$, we also conclude in that case that $h_{v_L} = 0$.

Let us now show that for all $(v, v') \in F_\theta$, $h_{v \rightarrow v'} = 0$. Let $l \in \llbracket 1, L \rrbracket$ and let $(v_{l-1}, v_l) \in (V_{l-1} \times V_l) \cap F_\theta$ so that (v_{l-1}, v_l) is arbitrary in F_θ . If $l = 1$, we define $p_i = (v_{l-1})$ and we have by convention $\theta_{p_i} = 1 \neq 0$. If $l > 1$, using Proposition 15 there exist $l' \in \llbracket 0, l-1 \rrbracket$ and a path $p_i = (v_{l'}, \dots, v_{l-1})$ such that $\rho_\theta(\tau)_{p_i} \neq 0$. If $l < L$, we define $v_{l+1} = s_{\max}^\theta(v_l)$, then $v_{l+2} = s_{\max}^\theta(v_{l+1})$, and so on up to $v_L = s_{\max}^\theta(v_{L-1})$, and we denote $p = p_i \cup (v_{l-1}, v_l, \dots, v_{L-1})$. If $l = L$, we denote $p = p_i$. Let us show the following expression.

$$[D\psi^\theta(\tau) \cdot h]_{p,v_L} = \sum_{\substack{k \in \llbracket l', l-1 \rrbracket \\ (v_k, v_{k+1}) \in F_\theta}} \frac{\partial \psi_{p,v_L}^\theta}{\partial \tau_{v_k \rightarrow v_{k+1}}}(\tau) h_{v_k \rightarrow v_{k+1}} \quad (40)$$

Indeed, if $l' \geq 1$, we have

$$\psi_{p,v_L}^\theta(\tau) = \tau_{v_{l'}} \prod_{k=l'}^{l-1} \rho_\theta(\tau)_{v_k \rightarrow v_{k+1}} \prod_{k=l}^{L-1} w_{v_k \rightarrow v_{k+1}},$$

with the classical convention that if $l = L$, the product on the right is empty thus equal to 1. We thus have

$$\begin{aligned} [D\psi^\theta(\tau) \cdot h]_{p,v_L} &= \frac{\partial \psi_{p,v_L}^\theta}{\partial \tau_{v_{l'}}}(\tau) h_{v_{l'}} + \sum_{\substack{k \in \llbracket l', l-1 \rrbracket \\ (v_k, v_{k+1}) \in F_\theta}} \frac{\partial \psi_{p,v_L}^\theta}{\partial \tau_{v_k \rightarrow v_{k+1}}}(\tau) h_{v_k \rightarrow v_{k+1}} \\ &= \sum_{\substack{k \in \llbracket l', l-1 \rrbracket \\ (v_k, v_{k+1}) \in F_\theta}} \frac{\partial \psi_{p,v_L}^\theta}{\partial \tau_{v_k \rightarrow v_{k+1}}}(\tau) h_{v_k \rightarrow v_{k+1}}, \end{aligned}$$

since we have already shown that $h_{v_{l'}} = 0$.

If $l' = 0$, we have

$$\psi_{p,v_L}^\theta(\tau) = \prod_{k=0}^{l-1} \rho_\theta(\tau)_{v_k \rightarrow v_{k+1}} \prod_{k=l}^{L-1} w_{v_k \rightarrow v_{k+1}},$$

with the same convention that when $l = L$ the product on the right is equal to 1, so again

$$[D\psi^\theta(\tau) \cdot h]_{p,v_L} = \sum_{\substack{k \in \llbracket 0, l-1 \rrbracket \\ (v_k, v_{k+1}) \in F_\theta}} \frac{\partial \psi_{p,v_L}^\theta}{\partial \tau_{v_k \rightarrow v_{k+1}}}(\tau) h_{v_k \rightarrow v_{k+1}}.$$

This concludes the proof of (40).

We can now show by induction the following statement, for $l \in \llbracket 0, L \rrbracket$.

$$\forall l' \in \llbracket 1, l \rrbracket, \forall (v, v') \in (V_{l'-1} \times V_{l'}) \cap F_\theta, h_{v \rightarrow v'} = 0. \quad (H_l)$$

Since $\llbracket 1, 0 \rrbracket = \emptyset$, H_0 is trivially true. Now let $l \in \llbracket 1, L \rrbracket$ and suppose H_{l-1} is true. We consider $(v_{l-1}, v_l) \in (V_{l-1} \times V_l) \cap F_\theta$, and $l' \in \llbracket 0, l \rrbracket$, p_i and p just as before. Since for all $k \in \llbracket 0, l-2 \rrbracket$, the induction hypothesis guarantees that $h_{v_k \rightarrow v_{k+1}} = 0$, (40) becomes

$$[D\psi^\theta(\tau) \cdot h]_{p,v_L} = \frac{\partial \psi_{p,v_L}^\theta}{\partial \tau_{v_{l-1} \rightarrow v_l}}(\tau) h_{v_{l-1} \rightarrow v_l}.$$

Using (38) and (39), we obtain

$$[D\psi^\theta(\tau) \cdot h]_{p,v_L} = \begin{cases} \rho_\theta(\tau)_{p_i} w_{v_l \rightarrow v_{l+1}} \cdots w_{v_{L-1} \rightarrow v_L} h_{v_{l-1} \rightarrow v_l} & \text{if } l < L \\ \rho_\theta(\tau)_{p_i} h_{v_{l-1} \rightarrow v_l} & \text{if } l = L. \end{cases}$$

Since $\rho_\theta(\tau)_{p_i} \neq 0$, and for $l < L$, $w_{v_l \rightarrow v_{l+1}} \cdots w_{v_{L-1} \rightarrow v_L} \neq 0$, we conclude that $h_{v_{l-1} \rightarrow v_l} = 0$ and that H_l holds.

This induction leads to the conclusion that $h = 0$ and $D\psi^\theta(\tau)$ is injective. \square

We are now equipped to prove Theorem 7, which we restate here.

Theorem 25. Σ_*^1 is a smooth manifold of $\mathbb{R}^{\mathcal{P} \times V_L}$ of dimension

$$|F_\theta| + |B| = N_0 N_1 + N_1 N_2 + \cdots + N_{L-1} N_L + N_L,$$

and the family $(V_\theta, (\psi^\theta)^{-1})_{\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S}$ is an atlas.

Proof. Our goal is to show that the family $(V_\theta, (\psi^\theta)^{-1})_{\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S}$ is a smooth atlas, which will show that Σ_*^1 is a smooth manifold.

We already know from Proposition 23 that for any $\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S$, V_θ is an open subset of Σ_*^1 and from Proposition 21 that $(\psi^\theta)^{-1}$ is a homeomorphism from V_θ onto U_θ . Since for any




Figure 3: The points η_0, τ_0, τ_1 and the inverse charts ψ^θ and $\psi^{\tilde{\theta}}$.

$\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S$, $\tau_\theta \in U_\theta$, we have $\phi(\theta) = \psi^\theta(\tau_\theta) \in V_\theta$ which shows that $(V_\theta)_{\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S}$ covers Σ_1^* .

Let $\theta, \tilde{\theta} \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S$, let us show that the transition map

$$(\psi^\theta)^{-1} \circ \psi^{\tilde{\theta}} : (\psi^{\tilde{\theta}})^{-1}(V_\theta \cap V_{\tilde{\theta}}) \rightarrow (\psi^\theta)^{-1}(V_\theta \cap V_{\tilde{\theta}})$$

is smooth.

Let $\tau_0 \in U_{\tilde{\theta}}$ such that $\tau_0 \in (\psi^{\tilde{\theta}})^{-1}(V_\theta \cap V_{\tilde{\theta}})$. We are going to show that the function $(\psi^\theta)^{-1} \circ \psi^{\tilde{\theta}}$ is C^∞ in a neighborhood of τ_0 .

For ease of reading, let us denote $\psi^{\tilde{\theta}}(\tau_0)$ by η_0 . By definition, $\eta_0 \in V_\theta \cap V_{\tilde{\theta}}$. In particular, since $\eta_0 \in V_\theta$, we can define $\tau_1 = (\psi^\theta)^{-1}(\eta_0)$. See Figure 3 for a representation.

Let $T = \text{Im } D\psi^\theta(\tau_1)$, and let us consider a linear subspace G such that $T \oplus G = \mathbb{R}^{\mathcal{P} \times V_L}$. Let $N_C = |\mathcal{P}|N_L - |F_\theta| - |B| = \dim(G)$. Let $i : \mathbb{R}^{N_C} \rightarrow G$ be linear and invertible. Let us consider the function

$$\begin{aligned} \varphi_\theta : U_\theta \times \mathbb{R}^{N_C} &\longrightarrow \mathbb{R}^{\mathcal{P} \times V_L} \\ (\tau, x) &\longmapsto \psi^\theta(\tau) + i(x). \end{aligned}$$

We are going to show that there exist an open neighborhood \tilde{U} of $(\tau_1, 0)$ in $(\mathbb{R}^{F_\theta} \times \mathbb{R}^B) \times \mathbb{R}^{N_C}$ and an open neighborhood \tilde{V} of η_0 in $\mathbb{R}^{\mathcal{P} \times V_L}$ such that φ_θ is a C^∞ diffeomorphism from \tilde{U} onto \tilde{V} satisfying

$$\varphi_\theta \left([(\mathbb{R}^{F_\theta} \times \mathbb{R}^B) \times \{0\}^{N_C}] \cap \tilde{U} \right) = \Sigma_1^* \cap \tilde{V}.$$

Let us first show that φ_θ is a C^∞ -diffeomorphism from a neighborhood of $(\tau_1, 0)$ in $(\mathbb{R}^{F_\theta} \times \mathbb{R}^B) \times \mathbb{R}^{N_C}$ onto a neighborhood of η_0 in $\mathbb{R}^{\mathcal{P} \times V_L}$. As shown in Proposition 24, ψ^θ is C^∞ and i is a linear function, so φ_θ is C^∞ . Let us prove that the differential $D\varphi_\theta(\tau_1, 0)$ is injective. For all $(\tau, x) \in (\mathbb{R}^{F_\theta} \times \mathbb{R}^B) \times \mathbb{R}^{N_C}$,

$$D\varphi_\theta(\tau_1, 0) \cdot (\tau, x) = D\psi^\theta(\tau_1) \cdot \tau + i(x).$$

Since $D\psi^\theta(\tau_1) \cdot \tau \in T$, $i(x) \in G$, and T and G are in direct sum, if $D\varphi_\theta(\tau_1, 0) \cdot (\tau, x) = 0$, then we have

$$\begin{cases} D\psi^\theta(\tau_1) \cdot \tau = 0 \\ i(x) = 0. \end{cases}$$

Since as shown in Proposition 24 $D\psi^\theta(\tau_1)$ is injective, and since i is invertible, we have

$$(\tau, x) = (0, 0).$$

Hence, $D\varphi_\theta(\tau_1, 0)$ is injective. Since $\dim((\mathbb{R}^{F_\theta} \times \mathbb{R}^B) \times \mathbb{R}^{N_C}) = |F_\theta| + |B| + N_C = |\mathcal{P}|N_L$, the differential $D\varphi_\theta(\tau_1, 0)$ is bijective. Using the inverse function theorem, there exists an open set $U \subset U_\theta \times \mathbb{R}^{N_C}$ containing $(\tau_1, 0)$, an open set $V \subset \mathbb{R}^{\mathcal{P} \times V_L}$ containing η_0 such that φ_θ is a C^∞ -diffeomorphism from U onto V .

We have

$$\varphi_\theta \left([(\mathbb{R}^{F_\theta} \times \mathbb{R}^B) \times \{0\}^{N_C}] \cap U \right) \subset V_\theta \cap V.$$

In fact, if V is small enough, this inclusion is an equality. We are going to construct open subsets $\tilde{U} \subset U$ and $\tilde{V} \subset V$ so that it is the case. Let us define

$$O = \{\tau \in U_\theta, (\tau, 0) \in U\}.$$

Since U is an open set containing $(\tau_1, 0)$, O is an open set containing $\tau_1 = (\psi^\theta)^{-1}(\eta_0)$. Since, according to Proposition 21, ψ^θ is a homeomorphism, $\psi^\theta(O)$ is an open subset of V_θ so there exists $\epsilon > 0$ such that

$$V_\theta \cap B_\infty(\eta_0, \epsilon) \subset \psi^\theta(O). \quad (41)$$

We can now define $\tilde{V} = V \cap B_\infty(\eta_0, \epsilon)$, and $\tilde{U} = \{(\tau, x) \in U, \varphi_\theta(\tau, x) \in \tilde{V}\}$, which are open sets such that $(\tau_1, 0) \in \tilde{U}$, $\eta_0 \in \tilde{V}$, and φ_θ is a C^∞ -diffeomorphism from \tilde{U} onto \tilde{V} . Let us show that

$$\varphi_\theta \left([(\mathbb{R}^{F_\theta} \times \mathbb{R}^B) \times \{0\}^{N_C}] \cap \tilde{U} \right) = V_\theta \cap \tilde{V}. \quad (42)$$

The direct inclusion is immediate: if $(\tau, 0) \in [(\mathbb{R}^{F_\theta} \times \mathbb{R}^B) \times \{0\}^{N_C}] \cap \tilde{U}$, then

$$\varphi_\theta(\tau, 0) = \psi^\theta(\tau) \in V_\theta \cap \tilde{V}.$$

For the reciprocal inclusion, if $\tau \in U_\theta$ is such that $\psi^\theta(\tau) \in V_\theta \cap \tilde{V}$, then by definition of ϵ and \tilde{V} , (41) guarantees, since ψ^θ is injective, that $\tau \in O$. By definition of O , we have $(\tau, 0) \in U$, and since

$$\varphi_\theta(\tau, 0) = \psi^\theta(\tau) \in \tilde{V},$$

this shows $(\tau, 0) \in \tilde{U}$. This shows the reciprocal inclusion, and thus (42) holds.

Let us now define

$$P_\theta : \mathbb{R}^{F_\theta} \times \mathbb{R}^B \times \mathbb{R}^{N_C} \longrightarrow \mathbb{R}^{F_\theta} \times \mathbb{R}^B \\ (\tau, x) \longmapsto \tau$$

the restriction to the first component, and let us observe that over $V_\theta \cap \tilde{V}$, we have

$$P_\theta \circ (\varphi_\theta)^{-1} = (\psi^\theta)^{-1}. \quad (43)$$

Indeed, if $\eta \in V_\theta \cap \tilde{V}$, then by (42) there exists $\tau \in U_\theta$ such that $(\tau, 0) \in \tilde{U}$ and $\varphi_\theta(\tau, 0) = \eta$. Since $\varphi_\theta(\tau, 0) = \psi^\theta(\tau)$, this shows that $\tau = (\psi^\theta)^{-1}(\eta)$ and thus

$$(\psi^\theta)^{-1}(\eta) = P_\theta(\tau, 0) = P_\theta \circ (\varphi_\theta)^{-1}(\eta).$$

Now recall that $\eta_0 = \psi^{\tilde{\theta}}(\tau_0)$. By continuity of $\psi^{\tilde{\theta}}$, there exists $\epsilon' > 0$ such that $B_\infty(\tau_0, \epsilon') \subset (\psi^{\tilde{\theta}})^{-1}(V_\theta \cap V_{\tilde{\theta}})$ and

$$\psi^{\tilde{\theta}}(B_\infty(\tau_0, \epsilon')) \subset \tilde{V}.$$

For any $\tau \in B_\infty(\tau_0, \epsilon')$, we have $\psi^{\tilde{\theta}}(\tau) \in V_\theta \cap \tilde{V}$ so, as we just proved with (43), $(\psi^\theta)^{-1} \circ \psi^{\tilde{\theta}}(\tau) = P_\theta \circ (\varphi_\theta)^{-1} \circ \psi^{\tilde{\theta}}(\tau)$. Since the functions $\psi^{\tilde{\theta}}$, $(\varphi_\theta)^{-1}$ and P_θ are all C^∞ , we conclude that the transition map $(\psi^\theta)^{-1} \circ \psi^{\tilde{\theta}}$ is C^∞ over $B_\infty(\tau_0, \epsilon')$, for all $\tau_0 \in (\psi^{\tilde{\theta}})^{-1}(V_\theta \cap V_{\tilde{\theta}})$. We conclude that $(\psi^\theta)^{-1} \circ \psi^{\tilde{\theta}}$ is C^∞ over $(\psi^{\tilde{\theta}})^{-1}(V_\theta \cap V_{\tilde{\theta}})$.

We have showed that $(V_\theta, (\psi^\theta)^{-1})_{\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S}$ is a smooth atlas, and thus that Σ_1^* is a smooth submanifold of $\mathbb{R}^{\mathcal{P} \times V_L}$. As computed in (21), its dimension is

$$|F_\theta| + |B| = N_0 N_1 + N_1 N_2 + \cdots + N_{L-1} N_L + N_L.$$

□

D Conditions of local identifiability

Let us restate (using Definition 6) and prove Theorem 8.

Theorem 26. *For any $X \in \mathbb{R}^{n \times V_0}$ and $\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus (S \cup \Delta_X)$, the two following statements are equivalent.*

- i) θ is locally identifiable from X .
- ii) There exists $\epsilon > 0$ such that $B_\infty(\phi(\theta), \epsilon) \cap \Sigma_1^* \cap N(X, \theta) = \{\phi(\theta)\}$.

Proof.

$i) \Rightarrow ii)$ Suppose $i)$ is satisfied for some $\epsilon_1 > 0$. We first construct $\epsilon' > 0$ and then consider $\eta \in B_\infty(\phi(\theta), \epsilon') \cap \Sigma_1^* \cap N(X, \theta)$, and we prove that $\eta = \phi(\theta)$. Since $\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus (S \cup \Delta_X)$ and since, according to Proposition 14, Δ_X is closed, there exists $\epsilon_2 > 0$ such that for any $\tilde{\theta} \in B_\infty(\theta, \epsilon_2)$,

$$\alpha(X, \theta) = \alpha(X, \tilde{\theta}),$$

i.e.

$$A(X, \theta) = A(X, \tilde{\theta}).$$

Consider $\epsilon = \min(\epsilon_1, \epsilon_2)$. Since, according to Proposition 21, $\rho_\theta \circ (\psi^\theta)^{-1}$ is continuous at $\phi(\theta) \in \psi^\theta(U_\theta)$, and since $\rho_\theta \circ (\psi^\theta)^{-1}(\phi(\theta)) = \rho_\theta(\tau_\theta) = \theta$, there exists $\epsilon' > 0$ such that for all $\tau \in U_\theta$,

$$\|\psi^\theta(\tau) - \phi(\theta)\|_\infty < \epsilon' \implies \|\rho_\theta(\tau) - \theta\|_\infty = \|\rho_\theta \circ (\psi^\theta)^{-1}(\psi^\theta(\tau)) - \rho_\theta \circ (\psi^\theta)^{-1}(\phi(\theta))\|_\infty < \epsilon. \quad (44)$$

Since $\phi(\theta) = \psi^\theta(\tau_\theta)$, Proposition 23 guarantees that, modulo a decrease of ϵ' , we can assume that

$$B_\infty(\phi(\theta), \epsilon') \cap \Sigma_1^* \subset \psi^\theta(U_\theta). \quad (45)$$

Now let $\eta \in B_\infty(\phi(\theta), \epsilon') \cap \Sigma_1^* \cap N(X, \theta)$. Let us prove that $\eta = \phi(\theta)$. Using (45), there exists $\tau \in U_\theta$ such that $\eta = \psi^\theta(\tau)$. Since $\|\phi(\theta) - \eta\|_\infty < \epsilon'$, we have using (44)

$$\|\rho_\theta(\tau) - \theta\|_\infty < \epsilon. \quad (46)$$

Since $\epsilon < \epsilon_2$, we have

$$A(X, \theta) = A(X, \rho_\theta(\tau)). \quad (47)$$

Since $\psi^\theta(\tau) = \eta \in N(X, \theta)$, we have by definition of $N(X, \theta)$ that $\psi^\theta(\tau) - \phi(\theta) \in \text{Ker } A(X, \theta)$, so

$$A(X, \theta) \cdot \psi^\theta(\tau) = A(X, \theta) \cdot \phi(\theta) \quad (48)$$

Using successively (3), (47), (48) and (3) again, we have

$$\begin{aligned} f_{\rho_\theta(\tau)}(X) &= A(X, \rho_\theta(\tau)) \cdot \phi(\rho_\theta(\tau)) \\ &= A(X, \theta) \cdot \phi(\rho_\theta(\tau)) \\ &= A(X, \theta) \cdot \phi(\theta) \\ &= f_\theta(X). \end{aligned}$$

Since the hypothesis $i)$ holds for ϵ_1 , using (46) and the fact that $\epsilon < \epsilon_1$, we have

$$\theta \sim \rho_\theta(\tau).$$

We conclude using Proposition 16 that

$$\eta = \phi(\rho_\theta(\tau)) = \phi(\theta),$$

which shows

$$B_\infty(\phi(\theta), \epsilon') \cap \Sigma_1^* \cap N(X, \theta) \subset \{\phi(\theta)\}.$$

The converse inclusion trivially holds and therefore $ii)$ holds.

$ii) \Rightarrow i)$ Suppose $ii)$ is satisfied for some $\epsilon' > 0$.

We first construct ϵ and prove $i)$ holds. Since $\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus (S \cup \Delta_X)$, using Proposition 14, there exists $\epsilon_1 > 0$ such that for all $\tilde{\theta} \in B_\infty(\theta, \epsilon_1)$,

$$\alpha(X, \theta) = \alpha(X, \tilde{\theta}),$$

i.e.

$$A(X, \theta) = A(X, \tilde{\theta}). \quad (49)$$

Since ϕ is continuous, there exists $\epsilon_2 > 0$ such that

$$\|\theta - \tilde{\theta}\|_\infty < \epsilon_2 \implies \|\phi(\theta) - \phi(\tilde{\theta})\|_\infty < \epsilon'.$$

Using Proposition 19, there exists $\epsilon_3 > 0$ such that

$$\theta \stackrel{R}{\sim} \tilde{\theta} \text{ and } \|\theta - \tilde{\theta}\|_\infty < \epsilon_3 \implies \theta \sim \tilde{\theta}.$$

Since $\theta \notin S$ and S is closed, there exists $\epsilon_4 > 0$ such that for all $\tilde{\theta} \in \mathbb{R}^E \times \mathbb{R}^B$, if $\|\theta - \tilde{\theta}\|_\infty < \epsilon_4$, then

$$\tilde{\theta} \notin S.$$

Let $\epsilon = \min(\epsilon_1, \epsilon_2, \epsilon_3, \epsilon_4)$. Let $\tilde{\theta} \in B_\infty(\theta, \epsilon)$, and suppose

$$f_\theta(X) = f_{\tilde{\theta}}(X).$$

Let us prove that $\theta \sim \tilde{\theta}$. Reformulating the above equality using (3) for both sides, and using the definition of A given in the beginning of Section 4, we have

$$A(X, \theta) \cdot \phi(\theta) = A(X, \tilde{\theta}) \cdot \phi(\tilde{\theta}).$$

Since $\|\theta - \tilde{\theta}\|_\infty < \epsilon \leq \epsilon_1$, we have the equality (49) and thus

$$A(X, \theta) \cdot \phi(\theta) = A(X, \theta) \cdot \phi(\tilde{\theta}).$$

In other words, $\phi(\tilde{\theta}) - \phi(\theta) \in \text{Ker } A(X, \theta)$. Since $\epsilon < \epsilon_4$, $\phi(\tilde{\theta}) \in \Sigma_1^*$. Since $\epsilon < \epsilon_2$, $\phi(\tilde{\theta}) \in B_\infty(\phi(\theta), \epsilon')$. Summarizing,

$$\phi(\tilde{\theta}) \in B_\infty(\phi(\theta), \epsilon') \cap \Sigma_1^* \cap N(X, \theta),$$

and using the hypothesis $ii)$, we conclude that

$$\phi(\tilde{\theta}) = \phi(\theta).$$

By Proposition 18, we have $\theta \stackrel{R}{\sim} \tilde{\theta}$, and since $\epsilon < \epsilon_3$, we conclude that

$$\theta \sim \tilde{\theta}.$$

□

We are now going to prove Theorems 9 and 10, which we restate as Theorems 27 and 28 respectively (using Definition 6).

Theorem 27 (Necessary condition). *Let $X \in \mathbb{R}^{n \times V_0}$ and $\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus (S \cup \Delta_X)$. If C_N is not satisfied, then θ is not locally identifiable from X (thus not globally identifiable).*

Theorem 28 (Sufficient condition). *Let $X \in \mathbb{R}^{n \times V_0}$ and $\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus (S \cup \Delta_X)$. If C_S is satisfied, then θ is locally identifiable from X .*

To prove the theorems, we need to prove first the following lemmas.

Lemma 29. Let us denote by $T = \text{Im } D\psi^\theta(\tau_\theta)$ the direction of the tangent plane to Σ_1^* at $\phi(\theta)$. Let us denote by H the intersection $\text{Ker } A(X, \theta) \cap T$. We have

$$\dim(H) = |F_\theta| + |B| - R_\Gamma. \quad (50)$$

Proof. Let $\eta \in T$. There exists $h \in \mathbb{R}^{F_\theta} \times \mathbb{R}^B$ such that $\eta = D\psi^\theta(\tau_\theta) \cdot h$. We have the following equivalence:

$$\begin{aligned} \eta \in \text{Ker } A(X, \theta) &\iff A(X, \theta) \cdot \eta = 0 \\ &\iff A(X, \theta) \circ D\psi^\theta(\tau_\theta) \cdot h = 0 \\ &\iff \Gamma(X, \theta) \cdot h = 0 \\ &\iff h \in \text{Ker } \Gamma(X, \theta). \end{aligned}$$

This shows that $D\psi^\theta(\tau_\theta)^{-1}(\text{Ker } A(X, \theta) \cap T) = \text{Ker } \Gamma(X, \theta) \subset \mathbb{R}^{F_\theta} \times \mathbb{R}^B$.

Since $D\psi^\theta(\tau_\theta)$ is injective, we thus have

$$\dim(H) = \dim(\text{Ker } \Gamma(X, \theta)) = |F_\theta| + |B| - R_\Gamma.$$

□

Lemma 30. Let G be a supplementary subspace of $\text{Ker } A(X, \theta)$ such that

$$H \oplus G = \text{Ker } A(X, \theta). \quad (51)$$

If $R_\Gamma = R_A$, there exist an open set $\mathcal{O} \subset U_\theta \times G$ containing $(\tau_\theta, 0)$ and an open set $\mathcal{V} \subset \mathbb{R}^{\mathcal{P} \times V_L}$ containing $\phi(\theta)$ such that

$$\xi : \begin{array}{ccc} \mathcal{O} & \longrightarrow & \mathcal{V} \\ (\tau, g) & \longmapsto & \psi^\theta(\tau) + g \end{array}$$

is a diffeomorphism from \mathcal{O} onto \mathcal{V} .

Proof. Let us first show that

$$T \oplus G = \mathbb{R}^{\mathcal{P} \times V_L}. \quad (52)$$

Indeed, since $\text{Ker } A(X, \theta) = H \oplus G$ and $T \cap \text{Ker } A(X, \theta) = H$, we have $T \cap G = \{0\}$. We of course have

$$T \oplus G \subset \mathbb{R}^{\mathcal{P} \times V_L}. \quad (53)$$

Let us show that $\dim(G) = \dim(\mathbb{R}^{\mathcal{P} \times V_L}) - \dim(T)$. First note that we have

$$\dim(\text{Ker } A(X, \theta)) = \dim(\mathbb{R}^{\mathcal{P} \times V_L}) - \text{rank}(A(X, \theta)) = |\mathcal{P}|N_L - R_A. \quad (54)$$

Using (51) and (54), we have

$$\begin{aligned} \dim(G) &= \dim(\text{Ker } A(X, \theta)) - \dim(H) \\ &= |\mathcal{P}|N_L - R_A - \dim(H). \end{aligned}$$

Using (50) and the hypothesis $R_\Gamma = R_A$ we thus have

$$\begin{aligned} \dim(G) &= |\mathcal{P}|N_L - R_A + R_\Gamma - |F_\theta| - |B| \\ &= |\mathcal{P}|N_L - |F_\theta| - |B| \\ &= |\mathcal{P}|N_L - \dim(T), \end{aligned}$$

where the last equality comes from the injectivity of $D\psi^\theta(\tau_\theta)$, shown in Proposition 24. Together with (53), this proves (52).

Let us now consider the function

$$\xi : \begin{array}{ccc} U_\theta \times G & \longrightarrow & \mathbb{R}^{\mathcal{P} \times V_L} \\ (\tau, g) & \longmapsto & \psi^\theta(\tau) + g. \end{array}$$

For all $(h, g) \in (\mathbb{R}^{F_\theta} \times \mathbb{R}^B) \times G$, we have

$$D\xi(\tau_\theta, 0) \cdot (h, g) = D\psi^\theta(\tau_\theta)h + g.$$

The differential $D\xi(\tau_\theta, 0)$ is injective. Indeed, if

$$D\xi(\tau_\theta, 0) \cdot (h, g) = 0,$$

then since $D\psi^\theta(\tau_\theta)h \in T$ and $g \in G$, we have

$$\begin{cases} D\psi^\theta(\tau_\theta)h = 0 \\ g = 0, \end{cases}$$

and since $D\psi^\theta(\tau_\theta)$ is injective, $h = 0$ and $D\xi(\tau_\theta, 0)$ is injective. Since, using (52),

$$\dim(\mathbb{R}^{F_\theta} \times \mathbb{R}^B) + \dim(G) = |\mathcal{P}|N_L,$$

$D\xi(\tau_\theta, 0)$ is bijective.

We can thus apply the inverse function theorem: there exists an open set $\mathcal{O} \subset U_\theta \times G$ containing $(\tau_\theta, 0)$, an open set $\mathcal{V} \subset \mathbb{R}^{\mathcal{P} \times V_L}$ containing $\phi(\theta)$ such that ξ is a diffeomorphism from \mathcal{O} into \mathcal{V} . \square

We can now prove the theorems.

Proof of Theorem 27. If C_N is not satisfied, then we have $R_\Gamma = R_A < |F_\theta| + |B|$. We can thus apply Lemma 30: there exist an open set $\mathcal{O} \subset U_\theta \times G$ containing $(\tau_\theta, 0)$ and an open set $\mathcal{V} \subset \mathbb{R}^{\mathcal{P} \times V_L}$ containing $\phi(\theta)$ such that

$$\xi : \begin{array}{ccc} \mathcal{O} & \longrightarrow & \mathcal{V} \\ (\tau, g) & \longmapsto & \psi^\theta(\tau) + g \end{array}$$

is a diffeomorphism from \mathcal{O} onto \mathcal{V} .

Consider $\epsilon > 0$. We define the open set $\tilde{\mathcal{O}} = \mathcal{O} \cap (\psi^\theta)^{-1}(B(\phi(\theta), \epsilon) \times G)$ and its image $\tilde{\mathcal{V}} = \xi(\tilde{\mathcal{O}})$.

Using the computation of $\dim(H)$ shown in Lemma 29, we have

$$\dim(H) = |F_\theta| + |B| - R_\Gamma > 0,$$

so there exists a nonzero $w \in H$ such that $\phi(\theta) + w \in \tilde{\mathcal{V}}$. Since ξ induces a diffeomorphism from $\tilde{\mathcal{O}}$ onto $\tilde{\mathcal{V}}$, there exists $(\tau, g) \in \tilde{\mathcal{O}}$ such that

$$\phi(\theta) + w = \psi^\theta(\tau) + g$$

i.e.

$$\psi^\theta(\tau) - \phi(\theta) = w - g. \quad (55)$$

Let us denote $\tilde{\theta} = \rho_\theta(\tau)$ and let us show that Theorem 26.ii) does not hold. By definition, $\phi(\tilde{\theta}) = \psi^\theta(\tau)$ and since $(\tau, g) \in \tilde{\mathcal{O}}$, $\|\phi(\theta) - \phi(\tilde{\theta})\|_\infty < \epsilon$. Since $H \cap G = \{0\}$, $w \in H$, $g \in G$ and $w \neq 0$, (55) shows that

$$\phi(\tilde{\theta}) - \phi(\theta) \neq 0.$$

Furthermore, since $w \in H \subset \text{Ker } A(X, \theta)$ and $g \in G \subset \text{Ker } A(X, \theta)$, (55) shows that

$$\phi(\tilde{\theta}) - \phi(\theta) \in \text{Ker } A(X, \theta),$$

so

$$\phi(\tilde{\theta}) \in N(X, \theta).$$

Summarizing, for any $\epsilon > 0$ there exists $\tilde{\theta} \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus S$ such that $\phi(\tilde{\theta}) \in B_\infty(\phi(\theta), \epsilon) \cap \Sigma_1^* \cap N(X, \theta) \setminus \{\phi(\theta)\}$. The second item of Theorem 26 does not hold. Since it is equivalent, the first item of Theorem 26 does not hold either. In other words, the conclusion of Theorem 27 holds. \square

Proof of Theorem 28. Suppose that C_S is satisfied. Using Lemma 29 and using C_S , we obtain

$$\dim(T \cap \text{Ker } A(X, \theta)) = |F_\theta| + |B| - R_\Gamma = 0.$$

We thus have

$$T \cap \text{Ker } A(X, \theta) = \{0\}. \quad (56)$$

In order to apply Theorem 26, let us show by contradiction that there exists $\epsilon > 0$ such that

$$B_\infty(\phi(\theta), \epsilon) \cap \Sigma_1^* \cap N(X, \theta) = \{\phi(\theta)\}. \quad (57)$$

More precisely, we suppose that for all $n \in \mathbb{N}^*$, there exists $\phi_n \in N(X, \theta) \cap \Sigma_1^*$ such that $\phi_n \neq \phi(\theta)$ and $\|\phi(\theta) - \phi_n\|_\infty < \frac{1}{n}$ and prove that it leads to $T \cap \text{Ker } A(X, \theta) \neq \{0\}$, which contradicts (56).

Using Proposition 23, there exists $n_0 \in \mathbb{N}^*$ such that for all $n \geq n_0$, there exists $\tau_n \in U_\theta$ such that $\phi_n = \psi^\theta(\tau_n)$. Since ψ^θ is a homeomorphism and $\psi^\theta(\tau_\theta) = \phi(\theta)$,

$$\phi_n \rightarrow \phi(\theta)$$

implies that

$$\tau_n \rightarrow \tau_\theta.$$

Moreover, for all $n \geq n_0$, $\tau_n \neq \tau_\theta$.

When n tends to infinity, we can thus write

$$\phi_n - \phi(\theta) = \psi^\theta(\tau_n) - \psi^\theta(\tau_\theta) = D\psi^\theta(\tau_\theta) \cdot (\tau_n - \tau_\theta) + o(\tau_n - \tau_\theta).$$

Let us apply $A(X, \theta)$ and divide by $\|\tau_n - \tau_\theta\|$.

$$\frac{1}{\|\tau_n - \tau_\theta\|} A(X, \theta) \cdot (\phi_n - \phi(\theta)) = A(X, \theta) \circ D\psi^\theta(\tau_\theta) \cdot \left(\frac{\tau_n - \tau_\theta}{\|\tau_n - \tau_\theta\|} \right) + \frac{1}{\|\tau_n - \tau_\theta\|} A(X, \theta) \circ o(\tau_n - \tau_\theta). \quad (58)$$

Since $\phi_n \in N(X, \theta)$ for all $n \in \mathbb{N}^*$,

$$\frac{1}{\|\tau_n - \tau_\theta\|} A(X, \theta) \cdot (\phi_n - \phi(\theta)) = 0.$$

Since $\frac{\tau_n - \tau_\theta}{\|\tau_n - \tau_\theta\|}$ belongs to the unit sphere, we can extract a subsequence that converges to a limit h with norm equal to 1. Taking the limit in (58) according to this subsequence, we obtain

$$0 = A(X, \theta) \circ D\psi^\theta(\tau_\theta) \cdot h,$$

which shows that $D\psi^\theta(\tau_\theta) \cdot h \in \text{Ker } A(X, \theta)$. Since $h \neq 0$ and $D\psi^\theta(\tau_\theta)$ is injective, $D\psi^\theta(\tau_\theta)h \neq 0$ and

$$T \cap \text{Ker } A(X, \theta) \neq \{0\}.$$

This is in contradiction with (56).

We have proven (57). We can now conclude thanks to Lemma 26: there exists $\epsilon' > 0$ such that for any $\tilde{\theta} \in \mathbb{R}^E \times \mathbb{R}^B$, if $\|\theta - \tilde{\theta}\| < \epsilon'$, then

$$f_\theta(X) = f_{\tilde{\theta}}(X) \implies \theta \sim \tilde{\theta}.$$

□

E Checking the conditions numerically

We restate and prove Proposition 12.

Proposition 31. *Let $X \in \mathbb{R}^{n \times V_0}$ and $\theta \in \mathbb{R}^E \times \mathbb{R}^B$. We have*

$$R_A = N_L \text{rank}(\alpha(X, \theta)).$$

Proof. Let $\eta \in \mathbb{R}^{\mathcal{P} \times V_L}$. We have

$$A(X, \theta) \cdot \eta = \alpha(X, \theta)\eta.$$

If we denote by $\eta^1, \dots, \eta^{N_L} \in \mathbb{R}^{\mathcal{P}}$ the N_L columns of η , the columns of $A(X, \theta) \cdot \eta$ are $\alpha(X, \theta)\eta^1, \dots, \alpha(X, \theta)\eta^{N_L}$. If we consider the matrix η as a family of N_L vectors of $\mathbb{R}^{\mathcal{P}}$ and the matrix $A(X, \theta) \cdot \eta$ as a family of N_L vectors of \mathbb{R}^n , the operator $A(X, \theta)$ can then be equivalently described as

$$A(X, \theta) : \begin{array}{ccc} (\mathbb{R}^{\mathcal{P}})^{N_L} & \longrightarrow & (\mathbb{R}^n)^{N_L} \\ (\eta^1, \dots, \eta^{N_L}) & \longmapsto & (\alpha(X, \theta)\eta^1, \dots, \alpha(X, \theta)\eta^{N_L}). \end{array}$$

The rank of such an operator is $N_L \text{rank}(\alpha(X, \theta))$.

□

We restate and prove Proposition 11.

Proposition 32. *Let $X \in \mathbb{R}^{n \times V_0}$ and $\theta \in (\mathbb{R}^E \times \mathbb{R}^B) \setminus (S \cup \Delta_X)$. The function*

$$\begin{aligned} U_\theta &\longrightarrow \mathbb{R}^{n \times V_L} \\ \tau &\longmapsto f_{\rho_\theta(\tau)}(X) \end{aligned}$$

is differentiable in a neighborhood of τ_θ , and we denote by $D_\tau f_{\rho_\theta(\tau_\theta)}(X)$ its differential at τ_θ . We have

$$D_\tau f_{\rho_\theta(\tau_\theta)}(X) = \Gamma(X, \theta). \quad (59)$$

Proof. Using (3) at $\rho_\theta(\tau)$ and the definition of ψ^θ in (11), we have

$$f_{\rho_\theta(\tau)}(X) = A(X, \theta) \cdot \psi^\theta(\tau).$$

Taking the differential of

$$\begin{aligned} U_\theta &\longrightarrow \mathbb{R}^{n \times V_L} \\ \tau &\longmapsto f_{\rho_\theta(\tau)}(X) \end{aligned}$$

at τ_θ , and using (13), we obtain

$$D_\tau f_{\rho_\theta(\tau_\theta)}(X) = A(X, \theta) \circ D\psi^\theta(\tau_\theta) = \Gamma(X, \theta). \quad \square$$

To finish with, the following proposition gives explicit expressions of the coefficients of $\Gamma(X, \theta)$. These expressions are given for the sake of theoretical completeness. Note that when it comes to computing $\Gamma(X, \theta)$ in practice (in order to compute R_Γ), the correct approach is using backpropagation as described in Section 5 rather than evaluating the expressions in Proposition 33 which involve sums with very large numbers of summands.

Proposition 33. *If we decompose it in the canonical bases of $\mathbb{R}^{F_\theta} \times \mathbb{R}^B$ and $\mathbb{R}^{\llbracket 1, n \rrbracket \times V_L}$, $\Gamma(X, \theta)$ is a $(nN_L) \times (|F_\theta| + |B|)$ matrix. For lighter notations, let us drop the dependency in (X, θ) and denote by γ^{i, v_L} the lines of $\Gamma(X, \theta)$, for $i \in \llbracket 1, n \rrbracket$ and $v_L \in V_L$, which satisfy $(\gamma^{i, v_L})^T \in \mathbb{R}^{F_\theta} \times \mathbb{R}^B$. For any $(i, v_L) \in \llbracket 1, n \rrbracket \times V_L$, let us express the coefficients of γ^{i, v_L} , i.e. express $\gamma_{v_l \rightarrow v_{l+1}}^{i, v_L}$ for any $v_l \rightarrow v_{l+1} \in F_\theta$ and express $\gamma_{v_l}^{i, v_L}$ for any $v_l \in B$.*

- For any $l \in \llbracket 0, L-1 \rrbracket$ and any $(v_l, v_{l+1}) \in V_l \times V_{l+1}$ such that $v_l \rightarrow v_{l+1} \in F_\theta$,

$$\begin{aligned} \gamma_{v_l \rightarrow v_{l+1}}^{i, v_L} &= \sum_{\substack{v_0 \in V_0 \\ \vdots \\ v_{l-1} \in V_{l-1} \\ v_{l+2} \in V_{l+2} \\ \vdots \\ v_{L-1} \in V_{L-1}}} x_{v_0}^i \bar{w}_{v_0 \rightarrow v_1} \bar{a}_{v_l}(x^i, \theta) \prod_{\substack{1 \leq k \leq L-1 \\ k \neq l}} a_{v_k}(x^i, \theta) w_{v_k \rightarrow v_{k+1}} \\ &\quad + \sum_{l'=1}^L \sum_{\substack{v_{l'} \in V_{l'} \\ \vdots \\ v_{l-1} \in V_{l-1} \\ v_{l+2} \in V_{l+2} \\ \vdots \\ v_{L-1} \in V_{L-1}}} b_{v_{l'}} \bar{a}_{v_l}(x^i, \theta) \prod_{\substack{l' \leq k \leq L-1 \\ k \neq l}} a_{v_k}(x^i, \theta) w_{v_k \rightarrow v_{k+1}}, \quad (60) \end{aligned}$$

where $\bar{w}_{v_0 \rightarrow v_1} = w_{v_0 \rightarrow v_1}$ and $\bar{a}_{v_l}(x^i, \theta) = a_{v_l}(x^i, \theta)$ except when $l = 0$ in which case $\bar{w}_{v_0 \rightarrow v_1} = 1$ and $\bar{a}_{v_l}(x^i, \theta) = 1$.

- For any $l \in \llbracket 1, L \rrbracket$ and any $v_l \in V_l$,

$$\gamma_{v_l}^{i, v_L} = \sum_{\substack{v_{l+1} \in V_{l+1} \\ \vdots \\ v_{L-1} \in V_{L-1}}} \prod_{l \leq k \leq L-1} a_{v_k}(x^i, \theta) w_{v_k \rightarrow v_{k+1}}. \quad (61)$$

Proof. Let $(i, v_L) \in \llbracket 1, n \rrbracket \times V_L$.

Let us compute $\gamma_{v_l \rightarrow v_{l+1}}^{i, v_L}$, for $l \in \llbracket 0, L-1 \rrbracket$ and $(v_l, v_{l+1}) \in V_l \times V_{l+1}$ such that $v_l \rightarrow v_{l+1} \in F_\theta$. $\gamma_{v_l \rightarrow v_{l+1}}^{i, v_L}$ is the coefficient corresponding to the line (i, v_L) and the column $(v_l \rightarrow v_{l+1})$ of $\Gamma(X, \theta)$. Let us denote by $h^{v_l \rightarrow v_{l+1}} \in \mathbb{R}^{F_\theta} \times \mathbb{R}^B$ the vector whose component indexed by $v_l \rightarrow v_{l+1}$ is equal to 1 and whose other components are zero. Let us denote by $d^{i, v_L} \in \mathbb{R}^{n \times V_L}$ the element whose component indexed by (i, v_L) is equal to 1 and whose other components are zero. Let us denote by $\langle \cdot, \cdot \rangle_{\mathbb{R}^{n \times V_L}}$ the scalar product of the euclidean space $\mathbb{R}^{n \times V_L}$. We have

$$\begin{aligned} \gamma_{v_l \rightarrow v_{l+1}}^{i, v_L} &= \left\langle d^{i, v_L}, \Gamma(X, \theta) \cdot h^{v_l \rightarrow v_{l+1}} \right\rangle_{\mathbb{R}^{n \times V_L}} \\ &= \left\langle d^{i, v_L}, A(X, \theta) \circ D\psi^\theta(\tau_\theta) \cdot h^{v_l \rightarrow v_{l+1}} \right\rangle_{\mathbb{R}^{n \times V_L}} \\ &= \left\langle d^{i, v_L}, A(X, \theta) \cdot \frac{\partial \psi^\theta}{\partial \tau_{v_l \rightarrow v_{l+1}}}(\tau_\theta) \right\rangle_{\mathbb{R}^{n \times V_L}} \\ &= \left\langle d^{i, v_L}, \alpha(X, \theta) \frac{\partial \psi^\theta}{\partial \tau_{v_l \rightarrow v_{l+1}}}(\tau_\theta) \right\rangle_{\mathbb{R}^{n \times V_L}} \\ &= \left[\alpha(X, \theta) \frac{\partial \psi^\theta}{\partial \tau_{v_l \rightarrow v_{l+1}}}(\tau_\theta) \right]_{i, v_L}, \end{aligned}$$

where $\left[\alpha(X, \theta) \frac{\partial \psi^\theta}{\partial \tau_{v_l \rightarrow v_{l+1}}}(\tau_\theta) \right]_{i, v_L}$ denotes the coefficient (i, v_L) of the product $\alpha(X, \theta) \frac{\partial \psi^\theta}{\partial \tau_{v_l \rightarrow v_{l+1}}}(\tau_\theta)$. Let us remind the dimensions in this product. For the left factor, recalling the definition given in the beginning of Section D, we have $\alpha(X, \theta) \in \mathbb{R}^{n \times \mathcal{P}}$. Concerning the right factor, since for any $\tau \in U_\theta$, we have $\psi^\theta(\tau) \in \mathbb{R}^{\mathcal{P} \times V_L}$, the partial derivative satisfies $\frac{\partial \psi^\theta}{\partial \tau_{v_l \rightarrow v_{l+1}}}(\tau_\theta) \in \mathbb{R}^{\mathcal{P} \times V_L}$. Hence, the dimension of the product is

$$\alpha(X, \theta) \frac{\partial \psi^\theta}{\partial \tau_{v_l \rightarrow v_{l+1}}}(\tau_\theta) \in \mathbb{R}^{n \times V_L}.$$

To obtain the coefficient (i, v_L) of this product, we keep the i^{th} line of the left factor, which is equal to $\alpha(x^i, \theta)$, and the column v_L of the right factor, which is equal to $\frac{\partial \psi_{v_L}^\theta}{\partial \tau_{v_l \rightarrow v_{l+1}}}(\tau_\theta)$. We thus have

$$\left[\alpha(X, \theta) \frac{\partial \psi^\theta}{\partial \tau_{v_l \rightarrow v_{l+1}}}(\tau_\theta) \right]_{i, v_L} = \alpha(x^i, \theta) \frac{\partial \psi_{v_L}^\theta}{\partial \tau_{v_l \rightarrow v_{l+1}}}(\tau_\theta) = \sum_{p \in \mathcal{P}} \alpha_p(x^i, \theta) \frac{\partial \psi_{p, v_L}^\theta}{\partial \tau_{v_l \rightarrow v_{l+1}}}(\tau_\theta).$$

Let $p \in \mathcal{P}$. If $p = (v_0, \dots, v_L) \in \mathcal{P}_0$, looking at the case 1 in the proof of Proposition 24, we have

$$\frac{\partial \psi_{p, v_L}^\theta}{\partial \tau_{v_l \rightarrow v_{l+1}}}(\tau_\theta) = \mathbf{1}_{\{v_l \rightarrow v_{l+1} \in p\}} \prod_{\substack{k \in \llbracket 0, L-1 \rrbracket \\ k \neq l}} w_{v_k \rightarrow v_{k+1}}.$$

Recalling the definition of $\alpha_p(x^i, \theta)$ in the case $p \in \mathcal{P}_0$, given in (2), we also have

$$\alpha_p(x^i, \theta) = x_{v_0}^i \prod_{k=1}^{L-1} a_{v_k}(x^i, \theta),$$

and thus

$$\alpha_p(x^i, \theta) \frac{\partial \psi_{p, v_L}^\theta}{\partial \tau_{v_l \rightarrow v_{l+1}}}(\tau_\theta) = \mathbf{1}_{\{v_l \rightarrow v_{l+1} \in p\}} x_{v_0}^i \prod_{k=1}^{L-1} a_{v_k}(x^i, \theta) \prod_{\substack{k \in \llbracket 0, L-1 \rrbracket \\ k \neq l}} w_{v_k \rightarrow v_{k+1}}. \quad (62)$$

Now if $p = (v_{l'}, \dots, v_L) \in \mathcal{P}_{l'}$, for $l' \in \llbracket 1, \dots, L-1 \rrbracket$, looking at the case 2 in the proof of Proposition 24, we have

$$\frac{\partial \psi_{p, v_L}^\theta}{\partial \tau_{v_l \rightarrow v_{l+1}}}(\tau_\theta) = \mathbf{1}_{\{v_l \rightarrow v_{l+1} \in p\}} b_{v_{l'}} \prod_{\substack{k \in \llbracket l', L-1 \rrbracket \\ k \neq l}} w_{v_k \rightarrow v_{k+1}}.$$

Recalling the definition of $\alpha_p(x^i, \theta)$ in the case $p \in \mathcal{P}_{l'}$, given in (2), we also have

$$\alpha_p(x^i, \theta) = \prod_{k=l'}^{L-1} a_{v_k}(x^i, \theta),$$

and thus

$$\alpha_p(x^i, \theta) \frac{\partial \psi_{p, v_L}^\theta}{\partial \tau_{v_l \rightarrow v_{l+1}}}(\tau_\theta) = \mathbf{1}_{\{v_l \rightarrow v_{l+1} \in p\}} b_{v_{l'}} \prod_{k=l'}^{L-1} a_{v_k}(x^i, \theta) \prod_{\substack{k \in [l', L-1] \\ k \neq l}} w_{v_k \rightarrow v_{k+1}}. \quad (63)$$

Finally, if $p = \beta$, looking at the case 3 in the proof of Proposition 24, we have

$$\frac{\partial \psi_{p, v_L}^\theta}{\partial \tau_{v_l \rightarrow v_{l+1}}}(\tau_\theta) = 0,$$

and thus

$$\alpha_p(x^i, \theta) \frac{\partial \psi_{p, v_L}^\theta}{\partial \tau_{v_l \rightarrow v_{l+1}}}(\tau_\theta) = 0. \quad (64)$$

Assembling (62), (63) and (64), we can sum over all $p \in \mathcal{P}$, and obtain

$$\begin{aligned} \gamma_{v_{l+1} \rightarrow v_l}^{i, v_L} &= \sum_{\substack{p \in \mathcal{P}_0 \\ p = (v_0, \dots, v_{L-1})}} \mathbf{1}_{\{v_l \rightarrow v_{l+1} \in p\}} x_{v_0}^i \prod_{k=1}^{L-1} a_{v_k}(x^i, \theta) \prod_{\substack{k \in [0, L-1] \\ k \neq l}} w_{v_k \rightarrow v_{k+1}} \\ &\quad + \sum_{l'=1}^L \sum_{\substack{p \in \mathcal{P}_{l'} \\ p = (v_{l'}, \dots, v_{L-1})}} \mathbf{1}_{\{v_l \rightarrow v_{l+1} \in p\}} b_{v_{l'}} \prod_{k=l'}^{L-1} a_{v_k}(x^i, \theta) \prod_{\substack{k \in [l', L-1] \\ k \neq l}} w_{v_k \rightarrow v_{k+1}} \end{aligned}$$

which can be reformulated, getting rid of the zero sums when $v_l \rightarrow v_{l+1} \notin p$, as

$$\begin{aligned} \gamma_{v_{l+1} \rightarrow v_l}^{i, v_L} &= \sum_{\substack{v_0 \in V_0 \\ \vdots \\ v_{l-1} \in V_{l-1} \\ v_{l+2} \in V_{l+2} \\ \vdots \\ v_{L-1} \in V_{L-1}}} x_{v_0}^i \bar{w}_{v_0 \rightarrow v_1} a_{v_l}(x^i, \theta) \prod_{\substack{k \in [1, L-1] \\ k \neq l}} a_{v_k}(x^i, \theta) w_{v_k \rightarrow v_{k+1}} \\ &\quad + \sum_{l'=1}^L \sum_{\substack{v_{l'} \in V_{l'} \\ \vdots \\ v_{l-1} \in V_{l-1} \\ v_{l+2} \in V_{l+2} \\ \vdots \\ v_{L-1} \in V_{L-1}}} a_{v_l}(x^i, \theta) b_{v_{l'}} \prod_{\substack{k \in [l', L-1] \\ k \neq l}} a_{v_k}(x^i, \theta) w_{v_k \rightarrow v_{k+1}}, \end{aligned}$$

which shows (60).

The proof of (61) is similar to the one of (60). \square