



HAL
open science

A bulky and stable copper(I)-phenanthroline complex: impact of steric strain and symmetry on the excited state properties

Lea Gimeno, Brian Phelan, Emily Sprague-Klein, Thierry Roisnel, Errol Blart, Christophe Gourlaouen, Lin Chen, Yann Pellegrin

► To cite this version:

Lea Gimeno, Brian Phelan, Emily Sprague-Klein, Thierry Roisnel, Errol Blart, et al.. A bulky and stable copper(I)-phenanthroline complex: impact of steric strain and symmetry on the excited state properties. *Inorganic Chemistry*, 2022, 61 (19), pp.7296-7307. 10.1021/acs.inorgchem.1c03901 . hal-03690046

HAL Id: hal-03690046

<https://hal.science/hal-03690046>

Submitted on 7 Jun 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A bulky and stable copper(I)-phenanthroline complex: impact of steric strain and symmetry on the excited state properties.

Lea Gimeno,¹ Brian T. Phelan,² Emily A. Sprague-Klein,² Thierry Roisnel,³ Errol Blart,¹ Christophe Gourlaouen,^{4*} Lin X. Chen,^{2,5*} Yann Pellegrin^{1*}

¹Université de Nantes, CNRS, CEISAM UMR6230, F-44000 Nantes, France

²Chemical Science and Engineering Division, Argonne National Laboratory, Lemont, Illinois 60439, United States

³Université de Rennes CNRS, Institut des Sciences Chimiques de Rennes, UMR6226, F-35000 Rennes, France

⁴Laboratoire de Chimie Quantique Institut de Chimie UMR 7177 CNRS-Université de Strasbourg, 4, Rue Blaise Pascal CS 90032, F-67081 Strasbourg Cedex, France

⁵Department of Chemistry, Northwestern University, Evanston Illinois 60208, United States

Corresponding authors Email addresses:

gourlaouen@unistra.fr

lchen@anl.gov

Yann.pellegrin@univ-nantes.fr

Abstract

The steric strain around copper(I) in typical $[\text{Cu}(\text{NN}_R)_2]^+$ complexes, where NN_R is a diimine ligand substituted in α of the nitrogen atoms by R, is known to strongly impact the excited-state properties. Generally speaking, the larger R, the longer the emission lifetime and higher quantum yield. However, the stability of the coordination scaffold can be at stake if the steric strain imposed by R is too large. In this work, we explore a way of fine-tuning the steric strain around Cu(I) to reach a balance between high emission quantum yield and stability in a highly bulky copper(I) complex. Taking stable $[\text{Cu}(\text{dipp})_2]^+$ and unstable $[\text{Cu}(\text{dtbp})_2]^+$ (where dipp and dtbp are respectively 2,9-diisopropyl-1,10-phenanthroline and 2,9-di-*tert*-butyl-1,10-phenanthroline) as boundary of two least and most sterically strained structures, we designed and characterized non-symmetrical ligand 2-isopropyl-9-*tert*-butyl-1,10-phenanthroline (**L1**) and corresponding complex $[\text{Cu}(\text{L1})_2]^+$ (**Cu1**).

The key experimental findings are that **Cu1** exhibits a rigid tetrahedral geometry in the ground state, close to that of $[\text{Cu}(\text{dtbp})_2]^+$ and with an intermediate stability between that of $[\text{Cu}(\text{dipp})_2]^+$

and $[\text{Cu}(\text{dtbp})_2]^+$. Conversely, the non-symmetrical nature of ligand **L1** leads to a shorter emission lifetime and smaller quantum yield than either $[\text{Cu}(\text{dipp})_2]^+$ or $[\text{Cu}(\text{dtbp})_2]^+$. This peculiar behavior is rationalized through the in depth analysis of the ultrafast dynamics of the excited state measured with optical transient absorption spectroscopy and theoretical calculations performed on the ground and excited state of **Cu1**. Our main findings are that the obtained complex is significantly more stable than $[\text{Cu}(\text{dtbp})_2]^+$ despite the sterically strained coordination sphere. The non-symmetrical nature of the ligand translates into a strongly distorted structure in the excited state. The distortion can be described as a rocking motion of one ligand, entailing the premature extinction of the excited state *via* several deactivation channels.

Introduction

Copper(I) complexes are drawing increased attention as photosensitizers, efficiently replacing heavy metal complexes in lighting devices,¹ dye sensitized solar cells,² solar fuel production³ and organic photochemistry.⁴ Homoleptic copper(I) complexes of the general formula $[\text{CuL}_2]^+$ where L is a diimine ligand are particularly interesting. They absorb over a large wavelength range (400-550 nm or farther in the red depending on L)^{5,6} through a metal to ligand charge transfer transition (MLCT) and emit in the red to near infrared, with respectable luminescence quantum yields Φ (up to 0.063)⁷ and lifetime τ up to a few microseconds.⁸ Importantly, it is necessary to tether sterically challenging groups in positions 2 and 9 of 1,10-phenanthroline (phen) to promote advantageous excited-state properties.^{9,10} The rationale is that photo-excitation of the MLCT transition entails electron transfer from copper-centered d orbitals towards π^* ligand-centered orbitals. Thus, $[\text{Cu}^{\text{I}}\text{L}_2]^+$ transiently turns into $[\text{Cu}^{\text{II}}(\text{L}^\bullet)\text{L}]^+$, leading to a drastic geometry change from pseudo-tetrahedral D_{2d} towards square planar D_2 . In other words, the coordination sphere flattens to make the transient Cu^{II} center in $[\text{Cu}^{\text{II}}(\text{L}^\bullet)\text{L}]^+$ more exposed to the solvent molecules, enabling fast quenching of the excited state assuming the solvent-solute interactions lower the energy gap between the MLCT triplet and the ground states.

Grafting bulky groups in α -positions of the nitrogen atoms on phen hinders the flattening of the copper(I) coordination sphere in the MLCT state and shields the Cu^{II} center from solvent quenching, thus enhancing the excited-state lifetime as well as other photochemical properties. In addition, increasing the steric bulkiness around copper(I) elevates the energy of the excited state,^{11,12} improving its reactivity from a thermodynamic point of view. Thus, larger steric strain around copper(I) generally leads to larger fluorescence yield Φ and longer excited-state lifetime τ .^{7,8,13,14} In dichloromethane (DCM), for example, complexes like $[\text{Cu}(\text{dipp})_2]^+$ ¹³ (where

dipp stands for 2,9-diisopropyl-1,10-phenanthroline) or $[\text{Cu}(\text{dsbtmp})_2]^+$ ⁷ (where dsbtmp stands for 2,9-disec-butyl-3,4,7,8-tetramethyl-1,10-phenanthroline) exhibit Φ of 0.4 and 6.3 %, and τ of 0.37 and 2.8 μs , respectively. $[\text{Cu}(\text{dmp})_2]^+$ (where dmp stands for 2,9-dimethyl-1,10-phenanthroline) features more modest Φ (0.02%) and τ (ca. 55 ns),¹⁰ while $[\text{Cu}(\text{phen})_2]^+$ is non luminescent at all.

On the other hand, increasing the steric bulk around copper(I) too much can make it difficult to prepare the homoleptic complex.^{15,16} For example, the synthesis of $[\text{Cu}(\text{dtbp})_2]^+$ (dtbp stands for 2,9-di-*tert*-butyl-1,10-phenanthroline) is very challenging,¹⁵ and the resulting complex is highly unstable.⁸ Nevertheless, even if $[\text{Cu}(\text{dtbp})_2]^+$ is an unstable complex, it exhibits fascinating properties, notably the longest luminescence lifetime recorded for a homoleptic copper(I) complex in solution (ca. 3.3 μs in DCM).⁸




Figure 1. Complex **Cu1** and models

Structurally, stable $[\text{Cu}(\text{dipp})_2]^+$ differs from unstable $[\text{Cu}(\text{dtbp})_2]^+$ by only two methyl groups on the substituents of each phenanthroline ligand. With this in mind, we were curious if removing only one methyl group from dtbp, yielding the non-symmetrical phenanthroline ligand 2-*tert*-butyl-9-isopropyl-1,10-phenanthroline (**L1**), would be sufficient to afford a stable copper(I) complex $[\text{Cu}(\text{L1})_2]^+$ in the usual mild conditions used for the synthesis of these coordination scaffolds (Figure 1). In addition, we were interested to study the optical and electrochemical properties of the corresponding complex in both the ground and excited states. In this contribution, we report the synthesis of ligand **L1** and associated complex $[\text{Cu}(\text{L1})_2]^+$ (named **Cu1**) as well as a detailed optical and electrochemical characterization. The stability and luminescence properties of **Cu1** demonstrate that steric strain is not the sole factor governing the excited-state properties of copper(I) complexes.^{12,17}

Results and discussion

Synthesis

The synthesis of ligand **L1** is performed in two steps (Figure S1). First, nucleophilic attack of anhydrous phen by *tert*-butyllithium followed by re-aromatization with MnO₂ afforded mono-substituted 2-*tert*-butyl-1,10-phenanthroline. The organolithium solution was carefully added (over 20 min) to limit the formation of the bis-substituted ligand. 2-*tert*-butyl-1,10-phenanthroline was then subjected to the same series of reactions with isopropyllithium as the organolithium reagent, affording **L1** in 77% yield over all steps. The homoleptic copper(I) complex [Cu(**L1**)₂]⁺ was finally assembled by mixing two equivalents of the ligand with one equivalent of copper(I) tetrakis acetonitrile hexafluorophosphate in degassed DCM at room temperature.

Structural properties

After solvent evaporation under reduced pressure, the obtained red powder was recrystallized from hot toluene. We obtained monocrystals which were suitable for a structural determination by X-ray scattering (Figure 2 and Figure S2). Relevant angles and distances are reported in Table 1 and nomenclature for the different atoms is given in figure S3. The structure reveals that the copper(I) ion is in a nearly tetrahedral environment. Cu-N distances are significantly influenced by the nature of the substituent in α of the corresponding nitrogen atom: *tert*-butyl groups impose a longer Cu-N distance (ca. 2.16 Å on average) whereas isopropyl allow for a shorter Cu-N bond length (2.04 Å on average) and likely a tighter coordination bond. Interestingly, the phenanthroline plane is slightly farther away from the bulkier *tert*-butyl group than from the less challenging isopropyl one (Figure S4), translating in a small wagging distortion (see figure S5 for the definition of all distortions in homoleptic copper(I) complexes). Nevertheless, the structures of **Cu1** and [Cu(dtbp)₂]⁺ are very close as evidenced by the calculation of the rocking, wagging and flattening angles (respectively θ_x , θ_y and θ_z) which are close to 90° in all cases (Figures S6 and S7). For example θ_x , θ_y and θ_z values are significantly different from 90° for [Cu(dipp)₂]⁺ (Figure S7) showing that the addition of one extra methyl on one isopropyl group of each phen ligand has substantially brought the structure closer to a tetrahedral geometry, as desired to promote the excited-state properties. This is confirmed by the τ_4 parameter ($\tau_4 = \frac{360^\circ - (\alpha + \beta)}{141^\circ}$, where α and β are the two largest N-Cu-N angles in the structure of **Cu1**, namely N1A-Cu-N2B and N2A-Cu-N2B, cf. Table 1) which is 0.66, 0.76 and 0.79 for [Cu(dipp)₂]⁺, **Cu1** and [Cu(dtbp)₂]⁺ respectively, the expected value for a perfect tetrahedron being 0.93.¹⁸

Table 1. Bond lengths and angles of interest for complex **Cu1** and models $[\text{Cu}(\text{dipp})_2]^+$ and $[\text{Cu}(\text{dtbp})_2]^+$ obtained by X-ray scattering.

Parameter	$[\text{Cu}(\text{dipp})_2]^+(\text{BF}_4)^{19}$	$[\text{Cu}(\text{dtbp})_2]^+(\text{BF}_4)^{15}$	Cu1 (PF ₆)
Distances (Angström)			
Cu-N1A	2.083	2.104(2)	2.057(4)
Cu-N2A	2.017	2.111(2)	2.122(4)
Cu-N1B	2.067	2.108(2)	2.031(4)
Cu-N2B	2.030	2.105(2)	2.207(4)
Angles (deg)			
N1A-Cu-N2A	82.25	84.53	82.09(14)
N1A-Cu-N1B	107.35	123.26	120.32(15)
N1A-Cu-N2B	118.71	122.92	126.36(14)
N2A-Cu-N1B	128.97	121.37	126.10(15)
N2A-Cu-N2B	138.38	124.93	126.94(13)
N1B-Cu-N2B	81.52	84.82	81.06(16)
θ_x	82.2	88.6	87.2
θ_y	72.5	90.0	87.4
θ_z	85.4	91.8	87.3
$\tau_4^{[18]}$	0.66	0.79	0.76




Figure 2. Structure of complex **Cu1**. Hydrogen atoms and counter-anion have been removed for the sake of clarity.

One last point concerns the orientation of the *tert*-Butyl groups of each ligand. As shown on Figure 2, *tert*-butyl groups can be in a staggered or eclipsed conformation (figure S8). The two

phen ligands are thus not equivalent in the crystal phase, possibly a result from crystal packing forces. However, the ^1H NMR spectrum of **Cu1** clearly shows that both coordinated ligands are equivalent in solution phase. DFT calculations are a powerful tool to assess the behavior of the complexes in solution, by recomposing the structures with computational chemistry means and defining DCM as the medium.

We computed the structure of the different possible conformers for **Cu1** (Figure 3). The first one (**Cu1_{sym1}**) is symmetric and corresponds to the case where both of the *tert*-Butyl groups on each ligand are in the staggered conformation. The second one (**Cu1_{nosym}**) is not symmetric, corresponding to the X-ray experimental structure displayed in Figure 2, namely one *tert*-Butyl is in a staggered conformation and the other one in an eclipsed conformation. Finally, the last calculated conformation (**Cu1_{sym2}**) corresponds to a symmetric structure where both *tert*-Butyl groups are in an eclipsed conformation. All relevant parameters (bond length, angles for each structure) are given in Table ST1. **Cu1_{sym1}** and **Cu1_{sym2}** are less stable than **Cu1_{nosym}** (Figure S9), though the structure of **Cu1_{sym2}** is very close in energy. The associated transition states are very low in energy. The barrier from **Cu1_{nosym}** to **Cu1_{sym1}** is 4.1 kcal mol⁻¹ and that from **Cu1_{nosym}** to **Cu1_{sym2}** is only 1.4 kcal mol⁻¹. These results show that there is free rotation of the *tert*-Butyl group at room temperature, as observed in the NMR spectra. The orientation of the *tert*-Butyl group impacts the structural parameters especially the Cu-N bond length. The difference between Cu-N2A or Cu-N2B (nitrogen atom close to *tert*-Butyl group) and Cu-N1A or Cu-N1B (nitrogen atom close to isopropyl groups) distances in the same ligand increases from **Cu1_{sym1}** (0.066 Å) to **Cu1_{sym2}** (0.175 Å), in accordance with the trend observed in the crystal structure.




Figure 3: structures of the three most stable rotamers of **Cu1**. Only the hydrogen of the tBu groups pointing toward the copper are shown. The relative Gibbs Free energies from GAUSSIAN calculations are in kcal/mol.

Absorption properties of complex **Cu1**

Cu1 was dissolved in DCM, yielding orange solutions which remained stable over several days when stored in the dark (Figure S10). The UV-Vis absorption spectrum of **Cu1** (Figure 4a) is dominated by a strong transition in the UV composed of multiple ligand-centered and high energy MLCT transitions while a broad lower energy MLCT transition spans between ca. 350 and 500 nm. The maximum absorption wavelength of the MLCT transition is 438 nm, between the values measured for $[\text{Cu}(\text{dipp})_2]^+$ (453 nm) and $[\text{Cu}(\text{dtbp})_2]^+$ (425 nm).⁸ This is in line with the steric strain of the coordination sphere in **Cu1** which is intermediate between $[\text{Cu}(\text{dipp})_2]^+$ and $[\text{Cu}(\text{dtbp})_2]^+$.




Figure 4. a) UV-Vis (black line) and b) emission (red line) spectra ($\lambda_{\text{exc}} = 438$ nm) of complex **Cu1** in DCM. Inset: zoom on the MLCT transition.




Figure 5. Theoretical (black line) and experimental (red line) absorption spectra of **Cu1_{nosym}** in DCM computed with ADF software. In inset a zoom on the 350-600 nm domain.

The theoretical spectra (Figure 5) of **Cu1_{nosym}** reproduces the features of the experimental one. The spectra of the two symmetric rotamers **Cu1_{sym1}** and **Cu1_{sym2}** are very similar (Figure S11). The nature of the electronic transitions was determined by mean of TheoDORE analysis (Figure 6). In this approach, the **Cu1_{nosym}** complex is divided into three fragments: the metallic cation and the two ligands. The nature of each transition is determined by the contribution of each fragment to either the emptied orbital or the filled one. In the computed spectra, several weak absorption bands are present between 350 and 550 nm corresponding to the broad and weak experimental bands observed in the same domain. The spectrum is dominated by an intense band at 270 nm (exp. 275 nm). The detailed nature of the transitions can be found in Table ST2. The low energy domain (316 to 526 nm) is composed only of MLCT states (S_1 to S_{23} , Figure 6). The lowest singlet-singlet transition $S_0 \rightarrow S_1$ is expected at 525 nm but has no oscillator strength, as is commonly observed for Cu(I) complexes with a nearly tetrahedral geometry in the ground state in agreement with the experimental results.

The first absorption band is due to $S_0 \rightarrow S_3$ transition (490 nm, MLCT transition). Several transitions ($S_0 \rightarrow S_6$ at 440 nm, $S_0 \rightarrow S_7$ at 433 nm and $S_0 \rightarrow S_8$ at 430 nm) generate the weaker band and explain the high-energy tail observed in the experimental spectra between 350 to 450 nm. The small experimental shoulder around 330 nm appearing on the computed spectra at 318 nm can be attributed to the MLCT transition $S_0 \rightarrow S_{23}$. The most intense band (245-300 nm, exp. 255 to 315 nm) is too often assigned to $\pi\text{-}\pi^*$ LC transitions, whereas it is the

combination of numerous electronic transitions (see comments below Table ST2). The nature of the lowest triplet states is identical to that of the corresponding singlet states.




Figure 6. Nature of the transitions generating the theoretical absorption spectra. Horizontal axis: numbering of the singlet states S_n for the corresponding $S_0 \rightarrow S_n$ transition. Vertical axis: proportions (between 0 and 1) of Metal Centered states (MC, black), ligand centered $\pi-\pi^*$ transitions (on the same ligand) (LC, red), Metal-to-Ligand Charge transfer (MLCT, blue), Ligand-to-Metal Charge transfer (LMCT, yellow) and Ligand-to-Ligand Charge transfer (LLCT, green) to the overall calculated $S_0 \rightarrow S_n$ transitions.

UV-Vis monitoring of the stability of Cu1 in presence of acetonitrile

As mentioned above, **Cu1** remained stable in aerated solution despite its crowded coordination sphere. In order to appreciate and compare the stabilities of **Cu1** and $[\text{Cu}(\text{dtbp})_2]^+$, we studied the evolution of the UV-Vis spectrum of **Cu1** in DCM when aliquots of acetonitrile were added in the medium (figure S12). Indeed, acetonitrile (ACN) is a weak ligand which nevertheless proved capable to destroy $[\text{Cu}(\text{dtbp})_2]^+$ efficiently following the equation:



The MLCT transition of **Cu1** collapses upon addition of ACN whereas model $[\text{Cu}(\text{dipp})_2]^+$ is unaffected (figure S12) showing that the coordination sphere of **Cu1** is bulky enough to favor ligand exchange with weakly coordinating acetonitrile for steric relief. On the other hand, **Cu1** is much more stable than $[\text{Cu}(\text{dtbp})_2]^+$ since ca. 5×10^4 equivalents of acetonitrile were necessary to destroy half the quantity of **Cu1** whereas only one equivalent was sufficient to quantitatively disrupt the coordination scaffold of $[\text{Cu}(\text{dtbp})_2]^+$.^{8,15} Conclusively, the removal of one mere methyl group *per* phenanthroline from the structure of $[\text{Cu}(\text{dtbp})_2]^+$ is enough to significantly improve the stability of the complex, although the coordination sphere is weakened because of the strong steric strain imposed by the remaining *tert*-butyl group, as previously observed.¹²

Study of the photoluminescence properties of complex Cu1

The strong steric bulk surrounding the copper(I) ion is expected to be a positive boon for the promotion of the photo-luminescence properties. Indeed, solutions of **Cu1** in DCM are brightly luminescent upon excitation of the MLCT transition. The usual broad emission band can be

monitored at 678 nm (Figure 4b, red line). The corresponding excitation spectrum (emission at 678 nm) is given in Figure S13 and reproduces the absorption features of the MLCT, confirming that the luminescence originates indeed from the light induced population of MLCT excited states. We retrieved $E^{00} = 2.14$ eV from the emission spectrum (using the tangent method, figure S14). This value is very close to $E^{00}([\text{Cu}(\text{dipp})_2]^+)$ despite **Cu1** be more sterically constrained than $[\text{Cu}(\text{dipp})_2]^+$. In addition, the emission quantum yield and lifetime are more than twice lower than those of $[\text{Cu}(\text{dipp})_2]^+$ (Table 2 and Figure S15): these results are in contradiction with the expected trend, and show that steric strain is not the sole parameter to influence the excited-state properties of copper(I) complexes. In previous works, it has been noticed that copper(I) complexes built with phenanthroline ligands which are substituted by two different bulky groups in positions 2 and 9 show decreased luminescence performances due to a high non-radiative rate constant k_{nr} .^{5,12,20} Likewise, a large k_{nr} of $7.86 \times 10^6 \text{ s}^{-1}$ was calculated in the case of **Cu1** (Table 2). Nevertheless, **Cu1** is the most luminescent complex reported in the series of homoleptic copper(I) complexes bearing non symmetrical phenanthroline ligands; it is as well the most sterically challenged, underlining that steric strain remains an important factor to take into consideration to increase Φ , τ and E^{00} , all things otherwise being equal.

Table 2. Ground state and excited state parameters for molecules **M** = **Cu1**, $[\text{Cu}(\text{dipp})_2]^+$ and $[\text{Cu}(\text{dtbp})_2]^+$. Measurements are performed in DCM unless otherwise mentioned.

M	$\lambda_{\text{abs}} / \text{nm}$ ($\epsilon / \text{M}^{-1} \cdot \text{cm}^{-1}$)	$E(\text{M}^*/\text{M})^{\text{a}}$ (V)	$E(\text{M}/\text{M}^{\cdot-})^{\text{b}}$ (V)	$\lambda_{\text{em}} / \text{nm}$ ($\Phi/\%$) ^a	$E(\text{M}^*/\text{M}^{\cdot-})^{\text{c}}$	$\tau / \text{ns}^{\text{a}}$	$k_{\text{r}}^{\text{d}} / \text{s}^{-1}$	$k_{\text{nr}}^{\text{d}} / \text{s}^{-1}$
Cu1	438 (4460)	0.60 (120)	-2.13 (100)	678 (0.17)	0.01	127	1.34×10^4	7.86×10^6
$[\text{Cu}(\text{dipp})_2]^+$	453 (7260)	0.65	-2.10	679 (0.4)	0.03	342	1.1×10^4	2.73×10^6
$[\text{Cu}(\text{dtbp})_2]^+ \cdot 8$	425 (3100)	-	-	599 (5.6)	-	3260	1.72×10^4	0.3×10^6

[a] degassed medium. [b] in degassed DMF. [c] Estimated using the Rehm and Weller equation. Potentials given vs. Fc^+/Fc couple. [d] $k_{\text{r}} = \Phi/\tau$ and $k_{\text{nr}} = 1/\tau - k_{\text{r}}$.

The emission properties were computed and thus the lowest excited singlet and triplet states were optimized for the three rotamers **Cu1**_{sym1}, **Cu1**_{sym2} and **Cu1**_{nosym} (Table 3). The excited states (S_1 or T_1) are MLCT states and therefore can be seen as intramolecular charge transfer states where a hole is mostly located on copper and one electron is localized either on phen_A or phen_B. This results in two degenerate electronic states with C_1 symmetry for **Cu1**_{sym1} and **Cu1**_{sym2} and nearly degenerate electronic states with C_1 symmetry for **Cu1**_{nosym}, corresponding crudely to $[\text{Cu}^{\text{II}}(\text{phen}_A^{\cdot-})(\text{phen}_B)]^+$ and $[\text{Cu}^{\text{II}}(\text{phen}_A)(\text{phen}_B^{\cdot-})]^+$. Shifting from one potential well to the other is possible through population of a transient state formally described as $[\text{Cu}^{\text{II}}(\text{phen}_A^{(0.5\cdot-)})(\text{phen}_B^{(0.5\cdot-)})]^+$ (Figure S16). The latter has a C_2 symmetry in the case of **Cu1**_{sym1} and **Cu1**_{sym2} (in the case of **Cu1**_{nosym}, the symmetry of the intermediate state remains in C_1 symmetry because of the different orientations of the *tert*-butyl groups).

In the C_2 point group the electron density of the excited state is delocalized on both ligands phen_A and phen_B. Breaking the symmetry leads to its localization on one of the phen, which is associated to a breathing of the Cu-N distance without rotation of the *tert*-Butyl fragment. In the ground state, **Cu1**_{nosym} was the lowest energy structure. In the excited states, the most stable geometries (in both S_1 and T_1) were generated by **Cu1**_{sym2}. All the computed emission wavelengths λ_{em} from the T_1 states (Table 3) are inconsistent with the experimental maxima at 678 nm. This shows that like other similar copper complexes, the emission likely stems from the singlet S_1 state, through thermally activated delayed fluorescence (TADF), where the luminescent S_1 state is populated from the weakly luminescent T_1 state by thermally activated reverse intersystem crossing. One should notice the strong influence of the *tert*-Butyl group orientations on the T_1 state properties (Table 3). This means that a simple rotation of the *tert*-butyl group (an easy motion at room temperature given the low energy barriers between rotamers) leads to the existence of many different minima on the lowest excited singlet or triplet potential energy surface (PES), each of them with its own set of properties. For example, the spin orbit coupling (SOC) varies impressively from one rotamer to the other. This value is known to strongly impact intersystem crossing and therefore the probability that T_1 either repopulates S_1 (leading to TADF) or decays to S_0 (either radiatively or non-radiatively).

We first computed the lowest excited S_1 singlet state retaining the C_2 symmetry for **Cu1**_{sym1} and **Cu1**_{sym2} and then we did the calculation without any symmetry (C_1 point group) for the three rotamers. The λ_{em} computed for the S_1 states in C_1 symmetry are too red-shifted (Table 4). However, the λ_{em} of the S_1 state of C_2 symmetry of rotamers **Cu1**_{sym1} (663 nm) or **Cu1**_{sym2} (706 nm) agree well with the experimental values. In what follows, we will not consider rotamer **Cu1**_{sym1} because its energy is significantly higher than **Cu1**_{sym2}, which means that this conformation will be minor in solution compared to the others. As mentioned above, the optimized S_1 state of C_1 symmetry of **Cu1**_{sym2} rotamer is the most stable structure, though that of **Cu1**_{nosym} is close in energy. Importantly, the **Cu1**_{sym2} rotamer S_1 state of C_2 symmetry (transition state with the electron delocalized over the two phenanthroline ligands, Figure S16) is also quite low in energy, being only 0.118 eV above the structure without symmetry **Cu1**_{nosym}.

From these data we can deduce the emission properties. After excitation, the complex relaxes in the lowest triplet PES where it hops between the different energetically accessible minima generated by two kinds of motion. The first one is due to the rotation of the *tert*-butyl groups and the second is the hopping of the exciton between the two ligands. The S_1 PES is thermally repopulated, being accessible due to the small S_1 - T_1 energy difference (Table 3) and significant SOC (as expected for copper(I) complexes with nearly tetrahedral geometries) whatever the geometry considered.²¹ From the S_1 PES, the complex radiatively relaxes to the ground state, generating this broad emission band due to contributions from the various rotamers (Table 4).

These multiple transition structures entail that the barriers between the PES minima are low and that the complex evolves freely between them, populating each of these structures with almost equal probability. This tends to “dilute” the excited state over several states, enabling accelerated deactivation channels, as was observed previously.¹²

In addition the SOC is significantly smaller for **Cu1** than $[\text{Cu}(\text{dipp})_2]^+$ (table ST3). This means that the thermally activated reverse intersystem crossing is less efficient for **Cu1** than $[\text{Cu}(\text{dipp})_2]^+$. The TADF process is therefore less likely for **Cu1**, likely participating to the lower luminescence quantum yield and lifetime of the latter compared to model $[\text{Cu}(\text{dipp})_2]^+$.

One last point concerns the global shape of the T_1 and S_1 structures. The tetrahedral arrangement around the copper atom is almost lost in the excited state; the coordination sphere approached a trigonal bipyramid with the copper at the center of the equatorial plane and one methyl group of a *tert*-Butyl group occupying an axial position (see file adf_struct.xyz in the SI). Such distortion is particularly pronounced for complexes bearing *tert*-Butyl substituents in α of the nitrogen atoms. It is well known that extended distortions of the copper(I) complexes coordination sphere lead to high non-radiative rates k_{nr} .

Table 3. Emission from the lowest T_1 state (E_{em}), distortion energy E_{dist} (defined as the difference between the energy of the ground state at its optimum and at the excited state geometry) and stabilization energy E_{stab} ($E_{stab} = E_{dist} + E_{em}$) energies in eV, emission wavelength in nm (λ_{em}), S_1 - T_1 energy splitting (ΔE_{ST}) in eV at T_1 geometry and S_1 - T_1 spin orbit coupling in cm^{-1} of the lowest triplet states T_1 at T_1 geometry for the three rotamers and group symmetry. a) The two phen ligand of **Cu1** being inequivalent, the emission properties from both have been computed.

Structure	Cu1_{sym1}		Cu1_{nosym}		Cu1_{sym2}	
	C_2	C_1	C_1	C_1^a	C_2	C_1
E_{em}	1.668	1.507	1.460	1.449	1.544	1.405
λ_{em}	743	823	849	855	803	882
E_{dist}	0.281	0.371	0.365	0.387	0.333	0.404
E_{stab}	1.949	1.878	1.824	1.837	1.877	1.810
ΔE_{ST}	0.222	0.197	0.211	0.184	0.266	0.232
SOC	0.1	19.0	33.6	15.0	0.0	25.8
	20.2	34.2	37.8	27.0	20.3	37.9
	20.2	34.2	37.8	27.0	20.3	37.9

Table 4. Emission (E_{em}), distortion (E_{dist}) and stabilization (E_{stab}) energies in eV, emission wavelength in nm (λ_{em}) and oscillator strength (f_{osc}) of the lowest S_1 states. a) The two phen ligands of **Cu1** being inequivalent, the emission properties from each of them have been computed.

Structure	Cu1_{sym1}		Cu1_{nosym}		Cu1_{sym2}	
	C_2	C_1	C_1	C_1^a	C_2	C_1
Symmetry						

E_{em}	1.870	1.650	1.589	1.597	1.756	1.530
E_{dist}	0.280	0.396	0.404	0.399	0.344	0.452
E_{stab}	2.149	2.046	1.992	1.996	2.100	1.982
λ_{em}	663	751	780	776	706	810
f_{osc}	$3.58 \cdot 10^{-2}$	$3.45 \cdot 10^{-3}$	$3.70 \cdot 10^{-3}$	$2.54 \cdot 10^{-3}$	$3.17 \cdot 10^{-2}$	$1.70 \cdot 10^{-3}$

Optical Transient Absorption Spectroscopy

Optical transient absorption (TA) spectroscopy experiments were performed on solutions of **Cu1** prepared in DCM at a concentration of about 0.5 mM (Figure 7A). Following excitation with 415 nm laser pulses (500 fs, 500 nJ), the difference spectra comprise negative features at wavelengths less than 480 nm and positive features at wavelengths greater than 480 nm, with excited-state absorption (ESA) maxima between 540-580 nm. At short time delays—0 to 2 ps—the ESA is broad and featureless, centered at 560 nm, and undergoes a subtle red-shift leading to a small decrease in the ESA intensity at 550 nm and a small increase in the ESA intensity at wavelengths greater than 560 nm. The broad ESA feature then becomes more structured over the time delay range of 2-50 ps, with strong maxima at 537 and 574 nm and weak maxima at 678 and 748 nm. The negative and positive features then decay concomitantly to zero over the time delay range of 50 ps to 1 μ s. Importantly, the absorbance of the samples did not change during photolysis, proving that **Cu1** is photo-stable during photolysis in degassed samples (Figure S17). The transient absorption data were analyzed using global-kinetic analysis to fit select kinetic traces with a first-order sequential kinetic model, $A \rightarrow B \rightarrow C \rightarrow GS$ (see Experimental Section and Figure S18 for details). The evolution-associated spectra (EAS) were extracted from the raw data using the resultant time-dependent model populations (Figure 7B and S18D).




Figure 7. (A) Optical transient absorption spectra at select time delays following excitation by 415 nm laser pulses for a solution of **Cu1** in DCM and (B) the evolution-associated spectra obtained via global-kinetic fitting of the raw data to a first-order sequential kinetic model, $A \rightarrow B \rightarrow C \rightarrow GS$.

According to the calculations, the 415 nm laser pulses were resonant with the $S_0 \rightarrow S_n$ ($n = 3, 6-8$, see calculations) MLCT transitions that are optically-allowed in the D_{2d} structure adopted by copper(I) diimine complexes in the ground state.²² Prior time-resolved spectroscopy experiments with shorter time resolution and theoretical studies demonstrated that the initial excited-state population rapidly relaxes from S_n to S_1 via internal conversion (IC), with a portion of the population undergoing intersystem crossing (ISC), on the 100 fs timescale.²³ The subtle red-shift of the broad ESA feature observed here occurs with $\tau = 1.2 \pm 0.6$ ps. This process is typically assigned to the pseudo Jahn-Teller flattening distortion commonly observed for photoexcited copper(I) diimine complexes bearing symmetrical phenanthroline ligands: as explained in the introduction, copper(I) is transiently oxidized into copper(II) upon excitation, thus moving from tetrahedral to square planar geometries, and leading to a flattening of the coordination scaffold. However, our calculations of the excited state of **Cu1** and of copper(I) and copper(II) variants of **Cu1** indicate that this excited-state structural distortion rather corresponds to the rocking motion of a phenanthroline ligand and leads to formation of the trigonal bipyramidal structure (calculated structures for the excited state of **Cu1** and the oxidized state of **Cu1** are available in the `adf_struc.xyz` and `gauss_struc.xyz` files, respectively, in SI. Also, see Table ST4 for relevant structural data on the excited states). This rocking motion is well illustrated in Figure 8, where a view of excited **Cu1** is given. The flattening distortion was observed on the 200-600 fs timescale in $[Cu(tbp)_2]^+$ (tbp stands for 2-*tert*-butyl-1,10-phenanthroline) and 700-800 fs timescale in $[Cu(dipp)_2]^+$, but was not observed in $[Cu(dtbp)_2]^+$ due to the increased steric bulk of the *tert*-butyl substituents in the 2 and 9 positions of the phenanthroline ligands.^{13,24} The $\tau \sim 1$ ps time constant for structural distortion in **Cu1** is consistent with the intermediate steric bulk of **L1** and our calculations, which predicted structural distortion upon oxidation of the copper.




Figure 8. Structure of **Cu1** in the excited state (example of **Cu1**_{nosym} in S₁ state)

Following structural distortion, the ESA blue-shifts and the various maxima become more pronounced with $\tau = 8.5 \pm 0.6$ ps, which is assigned to ISC from the ¹MLCT (S₁) to ³MLCT (T₁) state. ISC in **Cu1** occurs slightly faster than in [Cu(tbp)₂]⁺, [Cu(dipp)₂]⁺, and [Cu(dsbp)₂]⁺ (dsbp stands for 2,9-di-*sec*-butyl-1,10-phenanthroline), but slower than in the more sterically hindered [Cu(dtbp)₂]⁺, [Cu(diptmp)₂]⁺ (diptmp stands for 2,9-diisopropyl-3,4,7,8-tetramethyl-1,10-phenanthroline), and [Cu(dsbtmp)₂]⁺.^{13,24,25} Nozaki et al. demonstrated that SOC values for copper(I) bisimine complexes decrease significantly with the degree of the flattening distortion, which explains the trend among the copper(I) complexes with symmetrically substituted phenanthrolines.²¹ However, our calculations reveal that **Cu1** flattens only slightly (89.9° to 86.0°, see Table ST4), but distorts significantly via the rocking motion (160.4° to 148.6°, see Table ST4). Thus, distortions along nuclear coordinates other than flattening similarly decrease the rate of ISC and probably the magnitude of SOC in copper(I) bisphenanthroline complexes. This is corroborated by the lower calculated SOC value for **Cu1** compared to [Cu(dipp)₂]⁺ (Table ST3).

While the picosecond dynamics of **Cu1** suggest intermediate effects of the structural distortions compared to those of the less sterically hindered [Cu(dipp)₂]⁺ and more sterically hindered [Cu(dtbp)₂]⁺, the ³MLCT lifetime of **Cu1**, $\tau = 127$ ns, measured with TCSPC, is the shortest of this series of complexes (Table 2). As discussed above, the duration of the ³MLCT state typically decreases with increased flattening in the excited state for two reasons: the flattening distortion (1) enables quenching caused by exciplex-like solvent interactions with the transient copper(II) center and (2) decreases the energy gap between the T₁ and S₀ states. While **Cu1** does not flatten significantly, the rocking distortion has the same two effects on the excited-state lifetime. Given that the picosecond dynamics and absorbance and emission properties correlate with the steric hindrance of the phenanthroline ligands, an additional factor

beyond the steric hindrance and structural distortion likely contributes to the abbreviated $^3\text{MLCT}$ lifetime in copper(I) complexes bearing different substituents at the 2 and 9 positions. Figure S19 shows optical transient absorption spectra carried out on the nanosecond to microsecond timescales, with a measured $^3\text{MLCT}$ lifetime of 126 ns that is in agreement with the TCSPC results ($\tau = 127$ ns). The non-radiative decay rate constant of the $^3\text{MLCT}$ state—calculated from the emission decay rate and yield—is larger than that of $[\text{Cu}(\text{dipp})_2]^+$ and $[\text{Cu}(\text{dtbp})_2]^+$, which could result from the decreased symmetry of **Cu1** and may explain the relatively short excited-state lifetimes of the copper(I) complexes with non-symmetric substituents. This is in line with the distorted structures calculated for the S_1 and T_1 states. A secondary non-dominant decay component was also observed ($\tau = 5 \pm 1$ ns), which may correspond to solvent interactions following the excited-state rocking distortion that increases the exposure of the transient Cu(II) center to solvent molecules in the excited state. This 5-ns component is reproducible across two separate measurements of the **Cu1** compound in DCM (Figure S19c). Furthermore, global analysis of the ESA and ground-state bleach (GSB) showed that this secondary decay component ($\tau = 5$ ns) was present with similar relative amplitudes at both features. We assign this secondary decay component to the quenching of the $^3\text{MLCT}$ lifetime due to inner shell penetration of DCM and subsequent interaction with the Cu(II) center, drawing from our previous studies of the solute-solvent interactions of $[\text{Cu}^{\text{I}}(\text{phen})_2]^+$ in both acetonitrile and DCM.²⁶ In this earlier study, the MLCT triplet state lifetimes of the least sterically strained $[\text{Cu}^{\text{I}}(\text{phen})_2]^+$ were significantly shortened in DCM to ~ 140 ps and acetonitrile to 16 ps compared to 100 ns and ~ 1 ns for $[\text{Cu}^{\text{I}}(\text{dmp})_2]^+$, where methyl groups are attached to the 2,9 positions of the phen ligands. Therefore, when the transient Cu(II) of the MLCT state is exposed to DCM, the lifetime will also be shortened even though the solvent is considered non-coordinating. Thus, we hypothesize that the observation of biexponential $^3\text{MLCT}$ quenching results from distinct populations of **Cu1** with different orientations of the phenanthroline ligands/substituents that permit different degrees of solute-solvent (i.e. Cu(II)-DCM) interactions. The accelerated decay component ($\tau = 5$ ns) corresponds to the loss of a fraction of the $^3\text{MLCT}$ population through enhanced solute-solvent interactions. The relative amplitudes of the 5 ns and 126 ns decay components indicate about 1/5 to 1/4 of the **Cu1** $^3\text{MLCT}$ population adopted a structure that enabled closer approach of the DCM solvent molecules to the Cu(II) center. Because we observe the ~ 5 ns decay component in the GSB, we assume this process corresponds to **Cu1** returning back to the ground state without any intervening processes. Thus, we conclude that although excited-state **Cu1** was expected to be sterically hindered to solute-solvent interactions, it is still possible to observe large variations in excited-state quenching resulting from a distribution of coordination environments.

Electrochemistry

The electrochemical behavior of **Cu1** was investigated by cyclic voltammetry in DCM (anodic part, figure S20) and dimethylformamide (Figure S21). The voltammograms feature a reversible oxidation wave at 0.60 V and a reversible reduction wave at -2.10 V vs. Fc⁺/Fc, which is consistent with the values usually recorded for this kind of complex. E(**Cu1**⁺/**Cu1**) is slightly below the oxidation potential of [Cu(dipp)₂]⁺ despite the fact that **Cu1** is more sterically strained (which should imply the stabilization of the HOMO). Electronic factors could partly explain this result, since *tert*-butyl is more electron donating than isopropyl: **L1** is thus slightly electron richer than dipp and could better stabilize Cu(II). In addition, the coordination scaffold is sterically ill-balanced, and oxidized **Cu1** (i.e. [Cu^{II}(**L1**)₂]²⁺) is certainly more warped than [Cu^{II}(dipp)₂]²⁺ hence the observed trend of E(Cu^{II}/Cu^I). This is confirmed by DFT calculations (Table ST4 in supporting information): upon oxidation, the phen-Cu-phen angle in [Cu^{II}(dipp)₂]²⁺ shifts from 90.0° to 70.3° evidencing that this complex endures some flattening of about 20°. In addition, all four Cu-N distances shorten equivalently and the X-Cu-X angle stays at 180° (where X is the calculated centroid of the central C₆ ring of phenanthroline, Table ST3). On the contrary, the distortions are stronger in oxidized **Cu1**. Indeed, due to the imbalance, there is a rocking of one of the phen plane with an X-Cu-X angle reduced to 148.6°. The coordination tends towards a trigonal bipyramid (Figure 9) with the copper atom at the center of the structure and one methyl group of the *tert*-Butyl occupying an axial position.




Figure 9. Structure of [Cu^{II}(**L1**)₂]²⁺ (oxidized **Cu1**) (calculated)

The structure of oxidized **Cu1** is similar to the calculated structures of **Cu1** in the S₁ and T₁ states (see file adf_struct.xyz in SI), which is normal since the excited state of copper(I) complexes can be formally described by a copper(II) ion coordinated to a reduced ligand. Such distortions are limited by the four *tert*-butyl groups in [Cu(dtbp)₂]²⁺. The calculated structures

for $[\text{Cu}^{\text{I}}(\text{dipp})_2]^+$, $[\text{Cu}^{\text{II}}(\text{dipp})_2]^{2+}$, **Cu1**⁺, **Cu1**²⁺, $[\text{Cu}^{\text{I}}(\text{dtbp})_2]^+$ and $[\text{Cu}^{\text{II}}(\text{dtbp})_2]^{2+}$ are given in the SI (file gauss_struct.xyz) and relevant structural parameters are gathered in Table ST4.

Reductive quenching of the excited state

Knowing E^{00} and $E(\mathbf{Cu1}/\mathbf{Cu1}^{\cdot-})$, one can estimate the photo-oxidizing power $E(\mathbf{Cu1}^*/\mathbf{Cu1}^{\cdot-})$ at ca. 0.01 V vs. Fc^+/Fc (table 1). This value is modest and compares well with the usual values estimated for homoleptic Cu(I) complexes. Increasing the steric bulk around the Cu(I) ion did not entail the expected rise of E^{00} , and therefore $E(\mathbf{Cu1}^*/\mathbf{Cu1}^{\cdot-})$ and $E([\text{Cu}(\text{dipp})_2]^{*+}/[\text{Cu}(\text{dipp})_2]^+)$ are similar. **Cu1** was put to the test in a reductive quenching experiment in the presence of decamethylferrocene dmFc as the electron donor.²⁷ The quenching of the emission of **Cu1** was monitored for different concentrations of dmFc. Plotting τ_0/τ (where τ_0 and τ are the emission lifetimes in the absence and presence of dmFc, respectively) as a function of [dmFc] afforded a straight line (figure S22) whose slope gives the Stern-Volmer constant K_{SV} . The latter is equal to $k_{\text{Q}} \times \tau_0$ where k_{Q} is the bimolecular quenching rate. As stated in previous work,^{12,28} the quenching of the excited state of a copper complex by dmFc may occur via *two* mechanisms: energy transfer (“EnT” from the MLCT excited state to low lying iron(II) based (d,d) states with kinetic rate constant $k_{\text{EnT}} = 2.1 \times 10^8 \text{ s}^{-1}$)¹² and electron transfer (from dmFc to excited copper complexes with kinetic rate constant k_{RQ}). The quenching rate constants were anticipated to be rather large²⁷ and diffusion-limited processes (kinetic rate constant $k_{\text{d}} = 7 \times 10^9 \text{ M}^{-1} \cdot \text{s}^{-1}$ for ferrocene in DCM) cannot be ignored. Thus, the equation $\frac{1}{k_{\text{Q}}} = \frac{1}{k_{\text{d}}} + \frac{1}{k_{\text{RQ}} + k_{\text{EnT}}}$ applies.²⁷ From the Stern-Volmer plots, we retrieved $k_{\text{RQ}} = 3.9 \times 10^9 \text{ M}^{-1} \cdot \text{s}^{-1}$. Interestingly, $k_{\text{RQ}}(\mathbf{Cu1})$ is higher than $k_{\text{RQ}}([\text{Cu}(\text{dpp})_2]^+) = 8 \times 10^8 \text{ M}^{-1} \cdot \text{s}^{-1}$ (where dpp is 2,9-diphenyl-1,10-phenanthroline).²⁸ this is consistent with $E(\mathbf{Cu1}^*/\mathbf{Cu1}^{\cdot-})$ being significantly larger than $E([\text{Cu}(\text{dpp})_2]^{*+}/[\text{Cu}(\text{dpp})_2]^0)$ (0.01 V vs. -0.07 V vs. Fc^+/Fc respectively).^{12,28} However, $k_{\text{RQ}}(\mathbf{Cu1})$ is smaller than $k_{\text{RQ}}([\text{Cu}(\text{dipp})_2]^+)$ ($5.9 \times 10^9 \text{ M}^{-1} \cdot \text{s}^{-1}$) recorded in the same conditions, despite their similar photo-oxidative powers. This evidences that thermodynamic factors are not sufficient to rationalize the reactivity of **Cu1** and kinetic factors must be taken into account.²⁹ The non-symmetrical nature of ligand **L1** entails a large reorganization energy, as demonstrated by the distorted calculated structures of oxidized $[\text{Cu}^{\text{II}}(\mathbf{L1})_2]^{2+}$ (Figure 8) and S_1 or T_1 excited states (Tables 3 and 4). Interestingly, **Cu1**_{sym2} exhibits the highest E_{dist} in the series **Cu1**, $[\text{Cu}(\text{dipp})_2]^+$ and $[\text{Cu}(\text{dtbp})_2]^+$ (Tables 4 and ST3). The reorganization energy offsets the thermodynamic driving force and decreases the electron-transfer kinetics, following Marcus theory.

Conclusion

A sterically strained homoleptic copper(I) complex $[\text{Cu}(\mathbf{L1})_2]^+$ or **Cu1** has been designed, showing an improved stability compared to $[\text{Cu}(\text{dtbp})_2]^+$ in DCM solutions. The design is based on the synthesis of a non-symmetrical phenanthroline ligand where positions in α of the nitrogen atoms are occupied by two encumbering branched aliphatic chains, namely *tert*-butyl and isopropyl groups. The complex has been characterized in the ground and excited states. Early after excitation, structural distortions are observed by ultrafast transient absorption spectroscopy, in contrast to previously studied $[\text{Cu}(\text{dtbp})_2]^+$. Despite the increased steric strain imposed by the two substituents, the excited-state **Cu1** exhibit a shorter lifetime and a weaker emission quantum yield than those of $[\text{Cu}(\text{dipp})_2]^+$. The rationale behind this observation is two-fold. First, many energy minima exist on the potential energy surfaces (triplet and singlet) and the energy barriers between them is small. Statistically, several of these minima can be populated with almost equal probability and decay to the ground state, providing multiple deactivation channels. Second, because the vicinal isopropyl group is smaller than the *tert*-butyl group, the excited-state distortion corresponds to rocking motion of the ligand, which leads to a trigonal bipyramid structure of the excited state of **Cu1** (as verified by calculations of the structures of the excited state and of the oxidized state of **Cu1**) opening new deactivation channels.

Experimental section

General: chemicals were purchased from Sigma-Aldrich or Fisher Scientific and used as received. 2,9-bis-isopropyl-1,10-phenanthroline (dipp),³⁰ $[\text{Cu}(\text{dipp})_2]^{+13}$ and 2-*tert*-butyl-1,10-phenanthroline¹² were synthesized as previously published. Thin-layer chromatography (TLC) was performed on aluminium sheets precoated with Merck 5735 Kieselgel 60F254. Column chromatography was carried out with Merck 5735 Kieselgel 60F (0.040-0.063 mm mesh). ¹H NMR spectra were recorded on an AVANCE 300 UltraShield BRUKER. Chemical shifts for ¹H NMR spectra are referenced relative to residual protium in the deuterated solvent (CDCl_3 δ = 7.26 ppm). NMR spectra were recorded at room temperature, chemical shifts are written in ppm and coupling constants in Hz. Mass spectrometry was performed with a JEOL JMS-700 B/E spectrometer. Electrochemical measurements were made under an argon atmosphere in CH_2Cl_2 or in DMF with 0.1 M Bu_4NPF_6 . Cyclic voltammetry experiments were performed by using an Autolab PGSTAT 302N potentiostat/galvanostat. A standard three-electrode electrochemical cell was used. Potentials were referenced vs. reversible ferrocenium/ferrocene couple. The working electrode was a glassy carbon disk and the auxiliary electrode was a Pt wire. UV-visible absorption spectra were recorded on an Analytik

Jena spectrophotometer, using 1 cm path length cells. Emission spectra were recorded on a Fluoromax-3 Horiba spectrofluorimeter (1 cm quartz cells). E^{00} values were calculated by the tangent method: a tangent to the emission spectrum was drawn on the blue side of the spectrum. The intersection with the baseline gives the wavelength λ^{00} , used to calculate $E^{00} = 1240/\lambda^{00}$. Luminescence decays were recorded with a DELTAFLEX time correlated single photon counting system (HORIBA) on degassed DCM solutions.

A suitable crystal for X-ray diffraction single crystal experiment was selected and mounted with a cryoloop on the goniometer head of a D8 VENTURE Bruker AXS diffractometer equipped with a (CMOS) PHOTON 100 detector, using Mo-K radiation ($\lambda = 0.71073 \text{ \AA}$, multilayer monochromator) at $T = 150 \text{ K}$. Crystal structure has been described in monoclinic symmetry and P 21/c space group. Cell parameters were found as follows: $a = 16.0924(13)$, $b = 15.3035(11)$, $c = 15.6444(11) \text{ \AA}$, $\beta = 110.423(2)^\circ$, $V = 3610.6(5) \text{ \AA}^3$. Number of formula unit Z is equal to 4 and calculated density d and absorption coefficient μ values are 1.408 g.cm^{-3} and 0.714 mm^{-1} respectively. The structure was solved by dual-space algorithm using the SHELXT program,³¹ and then refined with full-matrix least-squares methods based on F^2 (SHELXL program).³² All non-hydrogen atoms were refined with anisotropic atomic displacement parameters. H atoms were finally included in their calculated positions and treated as riding on their parent atom with constrained thermal parameters. A final refinement on F^2 with 8267 unique intensities and 470 parameters converged at $\omega R(F^2) = 0.2218$ ($R_F = 0.0869$) for 7275 observed reflections with ($I > 2\sigma$).

Synthesis: **2-(tert-butyl)-9-isopropyl-1,10-phenanthroline :** 2-(tert-butyl)-1,10-phenanthroline (226 mg, 0.96 mmol) is dissolved in dry toluene (15 mL). The solution is cooled to 0°C before dropwise addition of isopropylolithium (0.7 M, 2.1 mL). The resulting solution is allowed to warm to room temperature and is stirred under argon for 18 h. Water (10 mL) is then added followed by a DCM extraction. Organic phases are stirred with activated manganese dioxide (2.15 g, 24 mmol) for 6h. The mixture is filtrated on celite, dried over magnesium sulfate and evaporated. The orange oil is purified by chromatography on silica (eluent: 0 to 5% of methanol in DCM). The title compound is obtained in quantitative yield (265 mg). ^1H NMR spectrum (300MHz, CDCl_3) $\delta = 8.14$ (m, 2H), 7.73 (m, 3H), 7.54 (d, $^3J_{H,H} = 8.4\text{Hz}$, 1H), 3.45 (sept, $^3J_{H,H} = 6.9\text{Hz}$, 1H), 1.60 (s, 9H), 1.55 (d, $^3J_{H,H} = 6.9\text{Hz}$, 6H). ^{13}C NMR spectrum (CDCl_3 , 75 MHz) $\delta = 169.29$ (s), 167.36 (s), 136.15 (s), 135.96 (s), 127.20 (s), 126.84 (s), 125.50 (s), 125.31 (s), 120.74 (s), 119.79 (s), 38.73 (s), 37.04 (s), 30.30 (s), 22.48 (s). HRMS (ESI⁺) for $\text{C}_{19}\text{H}_{22}\text{N}_2$ [$\text{M}+\text{H}$]⁺, m/z 279.1870 found, 279.1861 calc.

Complex **Cu1**: under argon atmosphere, $[\text{Cu}(\text{ACN})_4]\text{PF}_6$ (74 mg, 0.20 mmol) was dissolved in 5 mL of degassed DCM. This solution was transferred into a vial that contains **L1** (110 mg,

0.40 mmol). The red resulting solution was stirred overnight at room temperature. The mixture was precipitated in hexane to obtain a red powder. ^1H NMR spectrum (CDCl_3 , 300MHz) δ = 8.58 (m, 4H), 8.13 (d, $^3J_{\text{H,H}} = 8.7\text{Hz}$, 2H), 8.07 (s, 4H), 7.76 (d, $^3J_{\text{H,H}} = 8.7\text{Hz}$, 2H), 3.04 (sept, $^3J_{\text{H,H}} = 6.9\text{Hz}$, 2H), 1.38 (s, 18H), 0.93 (d, $^3J_{\text{H,H}} = 6.9\text{Hz}$, 4H), 0.66 (d, $^3J_{\text{H,H}} = 6.9\text{Hz}$, 4H). ^{13}C NMR spectrum (CDCl_3 , 300MHz) δ = 138.55 (s), 138.41 (s), 128.66 (s), 127.03 (s), 126.82 (s), 123.94 (s), 122.02 (s), 38.76 (s), 38.01 (s), 30.66 (s), 22.13 (s), 22.01 (s). HRMS (ESI⁺) for $\text{C}_{38}\text{H}_{44}\text{CuN}_4^+$ m/z 619.2861 found, 619.2862 calc.

Optical transient absorption spectroscopy: Femtosecond transient absorption (fsTA) spectroscopy experiments were performed on a previously described Ti:sapphire laser system (MaiTai oscillator / Spitfire Pro XP amplifier, Spectra-Physics) outputting a 1 kHz train of 830 nm, 100 fs, 2.6 mJ laser pulses.³³ 0.6 mJ of the fundamental were used to perform the fsTA experiments. A 10% reflective beamsplitter directed 60 μJ of the fundamental towards an optical delay line within the transient absorption spectrometer (Helios, Ultrafast Systems). The 0.54 mJ of transmitted fundamental were telescoped to about 2 mm diameter and frequency-doubled using a 2 mm thick type I lithium triborate crystal ($\theta = 90^\circ$, $\phi = 28^\circ$). The second harmonic was directed through a $\lambda/2$ waveplate and wiregrid polarizer to attenuate the pump pulse energy and set the relative polarization between the pump and probe to magic angle (54.7°). A mechanical chopper operating at 500 Hz was used to block every other pump pulse and the pump pulses were then focused to $\sim 200\text{-}300\ \mu\text{m}$ diameter at the sample with 500 nJ/pulse. The probe was generated by focusing about 3-4 μJ of the fundamental into a 3 mm sapphire crystal after the optical delay line. The probe pulses (450-800 nm) were set to horizontal polarization using a wiregrid polarizer and focused to about 200 μm at the sample. After interacting with the sample, the probe and signal were coupled into a fiber optic cable connected to a CCD detector. The probe spectra and difference signal were recorded and calculated using the Helios software (Ultrafast Systems). The fsTA spectra were measured with 3 s total averaging at each time delay.

Nanosecond transient absorption (nsTA) spectroscopy experiments were performed using a previously described Nd:YAG laser system (PL2210 oscillator and amplifier, Ekspla) coupled to a supercontinuum laser (STM-1-UV, Leukos).³³ The amplifier outputs a 0.5 kHz train of 355 nm, 25 ps, 0.3 mJ pulses which pumped an optical parametric generator (PG403, Ekspla) tuned to output 415 nm, 25 ps, 25 μJ pulses. The supercontinuum laser was triggered by the 1 kHz master trigger from the amplifier and output a 1 kHz train of broadband (400-1600 nm), 600 ps, 15 μJ pulses. The 415 nm pump was attenuated to 300 nJ/pulse and focused to about 200-300 μm at the sample. The broadband probe was focused to about 150 μm at the sample. After interacting with the sample, the probe and signal were detected using an SP-2150i Acton Series spectrograph (Princeton Instruments) coupled with a Spyder3 SG-14 (Teledyne

DALSA) CCD camera. A laboratory-written application (LabVIEW 2014, National Instruments) provided the time delays to the supercontinuum laser via DG535 delay generators (Stanford Research Systems) externally triggered by the 1 kHz master trigger, recorded the probe spectra, and calculated the difference signal.

The fsTA and nsTA data had the background signal (scattered pump photons and long-lived, >1 ms, transient signals) subtracted and were corrected for group delay dispersion in the probe pulses. The nsTA data were corrected for scan-to-scan drift in the pump-probe timing provided by the DG535 delay generators. Samples for the fsTA and nsTA spectroscopy experiments were prepared in DCM with an absorbance of about 0.5 (about 0.5 mM) at the peak of the MLCT transition. Three freeze-pump-thaw cycles were performed on the nsTA samples to degas the solutions and eliminate the effects of oxygen on the measured excited-state dynamics.

Global-kinetic analysis of the transient absorption data was performed using MATLAB programs written by Matthew D. Krzyaniak in Michael R. Wasielewski's group. Here, the program simultaneously fit raw kinetic traces at select probe wavelengths (450-700 nm at 10 nm increments) where the time constants ($\tau_i = 1/k_i$), time zero, and width of the instrument response function were shared and only the amplitudes varied between the kinetic traces. The program numerically solved the first-order kinetic equations for the corresponding rate matrix **K** (Figure S13) using matrix methods and convolved the solutions with a Gaussian instrument response function. The program employed least-squares fitting with the Levenberg-Marquardt method to find the parameters that best described the data, which were then fed back into the kinetic equations to solve for the time-dependent populations of each state in the kinetic model. The evolution-associated spectra were then extracted from the raw data using the time-dependent populations for each state in the proposed kinetic model.

Computational Details. Two sets of calculation were performed. A first set were performed with ADF 2019³⁴ at DFT level of theory (B3LYP functional). All atoms were described by TZP basis set. Scalar relativistic effects were introduced through ZORA Hamiltonian and solvent (DCM) taken into account through a PCM. The UV-Visible absorption spectrum was computed by TD-DFT on the basis of ADF optimized structures. The transitions were analyzed by mean of TheoDORE tool (TheoDORE: Theoretical Density, Orbital Relaxation and Exciton analysis). The emission wavelength of lowest excited singlet and triplet states was computed on the basis of their optimized structures.

A second set of calculations was performed with GAUSSIAN09 at DFT level of theory (B3LYP functional) including Grimme's dispersion corrections. All atoms were described by the 6-31+G** basis set and the solvent (DCM) introduced through a PCM. All structures were fully

optimized and the nature of the stationary point encountered determined by frequency calculations within the harmonic approximation. The minima were characterized by a full set of real frequencies.

Supporting information

ORTEP picture of the structure of **Cu1**, graphical illustrations, additional spectra and voltammograms, full data tables related to quantum calculations.

Acknowledgements

The optical transient spectroscopy part of the work is funded by Chemical, Biological and Geological Sciences Division, Basic Energy Science, Office of Science, the US Department of Energy under Contract No. DE-AC02-06CH11357.

The authors wish to thank the ANR (PERCO program n°ANR-16-CE07-0012-01) for financial support, the CEISAM (Nantes University) analysis platforms.

The calculations were carried out in part at the IDRIS computer centers through a grant of computer time from GENCI (A0050810629) and in part in the Strasbourg HPC. We would also like to thank Michael Eberhart for his assistance in the freeze-pump-thaw sample preparation procedure.

References

- (1) Roth, H. G.; Romero, N. A.; Nicewicz, D. A. *Synlett* **2016**, 27, 714, ; Keller, S.; Prescimone, A.; La Placa, M.-G.; Junquera-Hernández, J. M.; Bolink, H. J.; Constable, E. C.; Sessolo, M.; Ortí, E.; Housecroft, C. E. The shiny side of copper: bringing copper(i) light-emitting electrochemical cells closer to application. *Rsc Advances* **2020**, 10, 22631-22644, 10.1039/d0ra03824e; Fresta, E.; Weber, M. D.; Fernandez-Cestau, J.; Costa, R. D. White Light-Emitting Electrochemical Cells Based on Deep-Red Cu(I) Complexes. *Advanced Optical Materials* **2019**, 7, 1900830, <https://doi.org/10.1002/adom.201900830>; Marion, R.; Sguerra, F.; Di Meo, F.; Sauvageot, E.; Lohier, J.-F.; Daniellou, R.; Renaud, J.-L.; Linares, M.; Hamel, M.; Gaillard, S. NHC Copper(I) Complexes Bearing Dipyrindylamine Ligands: Synthesis, Structural, and Photoluminescent Studies. *Inorg. Chem.* **2014**, 53, 9181-9191, 10.1021/ic501230m; Zhang, Q.; Komino, T.; Huang, S.; Matsunami, S.; Goushi, K.; Adachi, C. Triplet Exciton Confinement in Green Organic Light-Emitting Diodes Containing Luminescent Charge-Transfer Cu(I) Complexes. *Adv. Funct. Mater.* **2012**, 22, 2327-2336, <https://doi.org/10.1002/adfm.201101907>.
- (2) Sandroni, M.; Favereau, L.; Planchat, A.; Akdas-Kilig, H.; Szuwarski, N.; Pellegrin, Y.; Blart, E.; Le Bozec, H.; Boujtita, M.; Odobel, F. Heteroleptic copper(i)-polypyridine complexes as efficient sensitizers for dye sensitized solar cells. *Journal of Materials Chemistry A* **2014**, 2, 9944-9947, 10.1039/c4ta01755b; Malzner, F. J.; Willgert, M.; Constable, E. C.; Housecroft, C. E. The way to panchromatic copper(I)-based dye-sensitized solar cells: co-sensitization with the organic dye SQ2. *J. Mater. Chem. A* **2017**, 5, 13717-13729, 10.1039/c7ta02575k; Dragonetti, C.; Magni, M.; Colombo, A.; Fagnani, F.; Roberto, D.; Melchiorre, F.; Biagini, P.; Fantacci, S. Towards efficient sustainable full-copper dye-sensitized solar cells. *Dalton Trans.* **2019**, 48, 9703-9711, 10.1039/c9dt00790c; Karpacheva, M.; Malzner, F. J.; Wobill, C.; Büttner, A.; Constable, E. C.; Housecroft, C. E. Cuprophilia:

Dye-sensitized solar cells with copper(I) dyes and copper(I)/(II) redox shuttles. *Dyes Pigm.* **2018**, *156*, 410-416, <https://doi.org/10.1016/j.dyepig.2018.04.033>.

(3) Khnayzer, R. S.; McCusker, C. E.; Olaiya, B. S.; Castellano, F. N. Robust Cuprous Phenanthroline Sensitizer for Solar Hydrogen Photocatalysis. *J. Am. Chem. Soc.* **2013**, *135*, 14068-14070, 10.1021/ja407816f; Windisch, J.; Oraziotti, M.; Hamm, P.; Alberto, R.; Probst, B. General Scheme for Oxidative Quenching of a Copper Bis-Phenanthroline Photosensitizer for Light-Driven Hydrogen Production. *ChemSusChem* **2016**, *9*, 1719-1726, doi:10.1002/cssc.201600422; Mejía, E.; Luo, S.-P.; Karnahl, M.; Friedrich, A.; Tschierlei, S.; Surkus, A.-E.; Junge, H.; Gladiali, S.; Lochbrunner, S.; Beller, M. A Noble-Metal-Free System for Photocatalytic Hydrogen Production from Water. *Chem. - Eur. J.* **2013**, *19*, 15972-15978, 10.1002/chem.201302091; Heberle, M.; Tschierlei, S.; Rockstroh, N.; Ringenberg, M.; Frey, W.; Junge, H.; Beller, M.; Lochbrunner, S.; Karnahl, M. Heteroleptic copper photosensitizers: Why an extended π -system does not automatically lead to enhanced hydrogen production. *Chem. - Eur. J.* **2017**, *23*, 312-319, 10.1002/chem.201604005; Fischer, S.; Hollmann, D.; Tschierlei, S.; Karnahl, M.; Rockstroh, N.; Barsch, E.; Schwarzbach, P.; Luo, S.-P.; Junge, H.; Beller, M.; Lochbrunner, S.; Ludwig, R.; Brueckner, A. Death and Rebirth: Photocatalytic Hydrogen Production by a Self-Organizing Copper-Iron System. *ACS Catal.* **2014**, *4*, 1845-1849, 10.1021/cs500387e; Takeda, H.; Monma, Y.; Sugiyama, H.; Uekusa, H.; Ishitani, O. Development of Visible-Light Driven Cu(I) Complex Photosensitizers for Photocatalytic CO₂ Reduction. *Frontiers in Chemistry* **2019**, *7*, 10.3389/fchem.2019.00418; Gracia, L.-L.; Luci, L.; Bruschi, C.; Sambri, L.; Weis, P.; Fuhr, O.; Bizzarri, C. New Photosensitizers Based on Heteroleptic CuI Complexes and CO₂ Photocatalytic Reduction with [NiII(cyclam)]Cl₂. *Chem. Eur. J.* **2020**, *26*, 9929-9937, <https://doi.org/10.1002/chem.202001279>; Lennox, A. J. J.; Fischer, S.; Jurrat, M.; Luo, S.-P.; Rockstroh, N.; Junge, H.; Ludwig, R.; Beller, M. Copper-Based Photosensitizers in Water Reduction: A More Efficient In Situ Formed System and Improved Mechanistic Understanding. *Chem. Eur. J.* **2016**, *22*, 1233-1238, 10.1002/chem.201503812; Luo, S.-P.; Chen, N.-Y.; Sun, Y.-Y.; Xia, L.-M.; Wu, Z.-C.; Junge, H.; Beller, M.; Wu, Q.-A. Heteroleptic copper(I) photosensitizers of dibenzo[b,j]-1,10-phenanthroline derivatives driven hydrogen generation from water reduction. *Dyes Pigm.* **2016**, *134*, 580-585, <https://doi.org/10.1016/j.dyepig.2016.07.040>; Takeda, H.; Kamiyama, H.; Okamoto, K.; Irimajiri, M.; Mizutani, T.; Koike, K.; Sekine, A.; Ishitani, O. Highly Efficient and Robust Photocatalytic Systems for CO₂ Reduction Consisting of a Cu(I) Photosensitizer and Mn(I) Catalysts. *J. Am. Chem. Soc.* **2018**, *140*, 17241-17254, 10.1021/jacs.8b10619.

(4) Baralle, A.; Fensterbank, L.; Goddard, J.-P.; Ollivier, C. Aryl Radical Formation by Copper(I) Photocatalyzed Reduction of Diaryliodonium Salts: NMR Evidence for a CuI/CuI Mechanism. *Chem. Eur. J.* **2013**, *19*, 10809-10813, 10.1002/chem.201301449; Michelet, B.; Deldaele, C.; Kajouj, S.; Moucheron, C.; Evano, G. A General Copper Catalyst for Photoredox Transformations of Organic Halides. *Org. Lett.* **2017**, *19*, 3576-3579, 10.1021/acs.orglett.7b01518; Reiser, O. Shining Light on Copper: Unique Opportunities for Visible-Light-Catalyzed Atom Transfer Radical Addition Reactions and Related Processes. *Acc. Chem. Res.* **2016**, *49*, 1990-1996, 10.1021/acs.accounts.6b00296; Engl, S.; Reiser, O. Copper Makes the Difference: Visible Light-Mediated Atom Transfer Radical Addition Reactions of Iodoform with Olefins. *ACS Catal.* **2020**, *10*, 9899-9906, 10.1021/acscatal.0c02984; Hossain, A.; Bhattacharyya, A.; Reiser, O. Copper's rapid ascent in visible-light photoredox catalysis. *Science* **2019**, *364*, eaav9713, 10.1126/science.aav9713; Sandoval-Pauker, C.; Molina-Aguirre, G.; Pinter, B. Status report on copper (I) complexes in photoredox catalysis; photophysical and electrochemical properties and future prospects. *Polyhedron* **2021**, *199*, 115105, <https://doi.org/10.1016/j.poly.2021.115105>; Hunter, C. J.; Boyd, M. J.; May, G. D.; Fimognari, R. Visible-Light-Mediated N-Desulfonylation of N-Heterocycles Using a Heteroleptic Copper(I) Complex as a Photocatalyst. *The Journal of Organic Chemistry* **2020**, *85*, 8732-8739, 10.1021/acs.joc.0c00983; Hernandez-Perez, A. C.; Collins, S. K. Heteroleptic Cu-Based Sensitizers in Photoredox Catalysis. *Acc. Chem. Res.* **2016**, *49*, 1557-1565, 10.1021/acs.accounts.6b00250.

(5) Meyer, M.; Albrecht-Gary, A.-M.; Dietrich-Buchecker, C. O.; Sauvage, J.-P. π - π Stacking-Induced Cooperativity in Copper(I) Complexes with Phenanthroline Ligands. *Inorg. Chem.* **1999**, *38*, 2279-2287, 10.1021/ic981259+.

(6) Kern, J.-M.; Sauvage, J.-P. Photoassisted C-C coupling via electron transfer to benzylic halides by a bis(di-imine) copper(I) complex. *Journal of the Chemical Society, Chemical Communications* **1987**, 546-548, 10.1039/c39870000546.


(7) McCusker, C. E.; Castellano, F. N. Design of a Long-Lifetime, Earth-Abundant, Aqueous Compatible Cu(I) Photosensitizer Using Cooperative Steric Effects. *Inorg. Chem.* **2013**, *52*, 8114-8120, 10.1021/ic401213p.

(8) Green, O.; Gandhi, B. A.; Burstyn, J. N. Photophysical Characteristics and Reactivity of Bis(2,9-di-tert-butyl-1,10-phenanthroline)copper(I). *Inorg. Chem.* **2009**, *48*, 5704-5714, 10.1021/ic802361q.

- (9) McMillin, D. R.; Buckner, M. T.; Ahn, B. T. A light-induced redox reaction of bis(2,9-dimethyl-1,10-phenanthroline)copper(I). *Inorg. Chem.* **1977**, *16*, 943-945, 10.1021/ic50170a046; Buckner, M. T.; McMillin, D. R. Photoluminescence from copper(I) complexes with low-lying metal-to-ligand charge transfer excited states. *J. Chem. Soc., Chem. Commun.* **1978**, 759,
- (10) Blaskie, M. W.; McMillin, D. R. Photostudies of copper(I) systems. 6. Room-temperature emission and quenching studies of bis(2,9-dimethyl-1,10-phenanthroline)copper(I). *Inorg. Chem.* **1980**, *19*, 3519-3522, 10.1021/ic50213a062.
- (11) Miller, M. T.; Gantzel, P. K.; Karpishin, T. B. Effects of Sterics and Electronic Delocalization on the Photophysical, Structural, and Electrochemical Properties of 2,9-Disubstituted 1,10-Phenanthroline Copper(I) Complexes. *Inorg. Chem.* **1999**, *38*, 3414-3422, 10.1021/ic9900399.
- (12) Gimeno, L.; Blart, E.; Rebilly, J.-N.; Coupeau, M.; Allain, M.; Roisnel, T.; Quarré de Verneuil, A.; Gourlaouen, C.; Daniel, C.; Pellegrin, Y. Non-Symmetrical Sterically Challenged Phenanthroline Ligands and Their Homoleptic Copper(I) Complexes with Improved Excited-State Properties. *Chem. Eur. J.* **2020**, *26*, 11887-11899, <https://doi.org/10.1002/chem.202001209>.
- (13) Garakyaraghi, S.; Crapps, P. D.; McCusker, C. E.; Castellano, F. N. Cuprous Phenanthroline MLCT Chromophore Featuring Synthetically Tailored Photophysics. *Inorg. Chem.* **2016**, *55*, 10628-10636, 10.1021/acs.inorgchem.6b01880.
- (14) Eggleston, M. K.; McMillin, D. R.; Koenig, K. S.; Pallenberg, A. J. Steric Effects in the Ground and Excited States of Cu(II) Systems. *Inorg. Chem.* **1997**, *36*, 172-176, 10.1021/ic960698a; Kohler, L.; Hayes, D.; Hong, J.; Carter, T. J.; Shelby, M. L.; Fransted, K. A.; Chen, L. X.; Mulfort, K. L. Synthesis, structure, ultrafast kinetics, and light-induced dynamics of CuHETPHEN chromophores. *Dalton Trans.* **2016**, 10.1039/c6dt00324a; Cunningham, C. T.; Cunningham, K. L. H.; Michalec, J. F.; McMillin, D. R. Cooperative Substituent Effects on the Excited States of Copper Phenanthrolines. *Inorg. Chem.* **1999**, *38*, 4388-4392, 10.1021/ic9906611; Miller, M. T.; Gantzel, P. K.; Karpishin, T. B. A Photoluminescent Copper(I) Complex with an Exceptionally High Cu(II)/Cu(I) Redox Potential: [Cu(bfp)₂]⁺ (bfp = 2,9-bis(trifluoromethyl)-1,10-phenanthroline). *Angew. Chem., Int. Ed.* **1998**, *37*, 1556-1558, 10.1002/(sici)1521-3773(19980619)37:11<1556::aid-anie1556>3.0.co;2-w; Miller, M. T.; Karpishin, T. B. Phenylethynyl Substituent Effects on the Photophysics and Electrochemistry of [Cu(dpp)₂]⁺ (dpp = 2,9-Diphenyl-1,10-phenanthroline). *Inorg. Chem.* **1999**, *38*, 5246,
- (15) Gandhi, B. A.; Green, O.; Burstyn, J. N. Facile Oxidation-Based Synthesis of Sterically Encumbered Four-Coordinate Bis(2,9-di-tert-butyl-1,10-phenanthroline)copper(I) and Related Three-Coordinate Copper(I) Complexes. *Inorg. Chem.* **2007**, *46*, 3816-3825, 10.1021/ic0615224.
- (16) Miller, M. T.; Gantzel, P. K.; Karpishin, T. B. A Highly Emissive Heteroleptic Copper(I) Bis(phenanthroline) Complex: [Cu(dbp)(dmp)]⁺ (dbp = 2,9-Di-tert-butyl-1,10-phenanthroline; dmp = 2,9-Dimethyl-1,10-phenanthroline). *J. Am. Chem. Soc.* **1999**, *121*, 4292-4293, 10.1021/ja9901415.
- (17) Brown-Xu, S.; Fumanal, M.; Gourlaouen, C.; Gimeno, L.; Quatela, A.; Thobie-Gautier, C.; Blart, E.; Planchat, A.; Riobé, F.; Monnereau, C.; Chen, L. X.; Daniel, C.; Pellegrin, Y. Intriguing Effects of Halogen Substitution on the Photophysical Properties of 2,9-(Bis)halo-Substituted Phenanthrolinecopper(I) Complexes. *Inorg. Chem.* **2019**, *58*, 7730-7745, 10.1021/acs.inorgchem.9b00042.
- (18) Yang, L.; Powell, D. R.; Houser, R. P. Structural variation in copper(I) complexes with pyridylmethylamide ligands: structural analysis with a new four-coordinate geometry index, τ_4 . *Dalton Trans.* **2007**, 955-964, 10.1039/b617136b.
- (19) Smith, C. S.; Mann, K. R. Void Space Containing Crystalline Cu(I) Phenanthroline Complexes As Molecular Oxygen Sensors. *Chem. Mater.* **2009**, *21*, 5042-5049, 10.1021/cm901109n.
- (20) Felder, D.; Nierengarten, J. F.; Barigelletti, F.; Ventura, B.; Armaroli, N. Highly luminescent Cu(I)-phenanthroline complexes in rigid matrix and temperature dependence of the photophysical properties. *J. Am. Chem. Soc.* **2001**, *123*, 6291-6299, 10.1021/ja0043439.
- (21) Siddique, Z. A.; Yamamoto, Y.; Ohno, T.; Nozaki, K. Structure-Dependent Photophysical Properties of Singlet and Triplet Metal-to-Ligand Charge Transfer States in Copper(I) Bis(diimine) Compounds. *Inorg. Chem.* **2003**, *42*, 6366-6378, 10.1021/ic034412v.
- (22) Everly, R. M.; McMillin, D. R. Reinvestigation of the absorbing and emitting charge-transfer excited states of [Cu(II)]²⁺ systems. *The Journal of Physical Chemistry* **1991**, *95*, 9071-9075, 10.1021/j100176a011.
- (23) Shaw, G. B.; Grant, C. D.; Shiota, H.; Castner, E. W.; Meyer, G. J.; Chen, L. X. Ultrafast Structural Rearrangements in the MLCT Excited State for Copper(I) bis-Phenanthrolines in Solution. *J. Am. Chem. Soc.* **2007**, *129*, 2147-2160, 10.1021/ja067271f; Iwamura, M.; Takeuchi, S.; Tahara, T. Real-Time Observation of the Photoinduced Structural Change of Bis(2,9-dimethyl-1,10-phenanthroline)copper(I) by Femtosecond Fluorescence Spectroscopy: A Realistic Potential Curve of the Jahn-Teller Distortion. *J. Am. Chem. Soc.* **2007**, *129*, 5248-5256, 10.1021/ja069300s; Capano, G.; Chergui, M.; Rothlisberger,

- U.; Tavernelli, I.; Penfold, T. J. A Quantum Dynamics Study of the Ultrafast Relaxation in a Prototypical Cu(I)–Phenanthroline. *J. Phys. Chem. A* **2014**, *118*, 9861-9869, 10.1021/jp509728m.
- (24) Gothard, N. A.; Mara, M. W.; Huang, J.; Szarko, J. M.; Rolczynski, B.; Lockard, J. V.; Chen, L. X. Strong Steric Hindrance Effect on Excited State Structural Dynamics of Cu(I) Diimine Complexes. *J. Phys. Chem. A* **2012**, *116*, 1984-1992, 10.1021/jp211646p.
- (25) Garakyaraghi, S.; Danilov, E. O.; McCusker, C. E.; Castellano, F. N. Transient Absorption Dynamics of Sterically Congested Cu(I) MLCT Excited States. *J. Phys. Chem. A* **2015**, *119*, 3181-3193, 10.1021/acs.jpca.5b00901.
- (26) Lockard, J. V.; Kabehie, S.; Zink, J. I.; Smolentsev, G.; Soldatov, A.; Chen, L. X. Influence of Ligand Substitution on Excited State Structural Dynamics in Cu(I) Bisphenanthroline Complexes. *The Journal of Physical Chemistry B* **2010**, *114*, 14521-14527, 10.1021/jp102278u.
- (27) Cunningham, K. L.; McMillin, D. R. Reductive Quenching of Photoexcited Cu(dipp)₂⁺ and Cu(tptap)₂⁺ by Ferrocenes (dipp = 2,9-Diisopropyl-1,10-phenanthroline and tptap = 2,3,6,7-Tetraaryl-1,4,5,8-tetraazaphenanthrene). *Inorg. Chem.* **1998**, *37*, 4114-4119, 10.1021/ic980213d.
- (28) Cunningham, K. L.; Hecker, C. R.; McMillin, D. R. Competitive energy-transfer and reductive quenching of the CT excited states of copper(I) phenanthrolines. *Inorganica Chim. Acta* **1996**, *242*, 143-147, [http://dx.doi.org/10.1016/0020-1693\(95\)04859-6](http://dx.doi.org/10.1016/0020-1693(95)04859-6).
- (29) Gushurst, A. K. I.; McMillin, D. R.; Dietrich-Buchecker, C. O.; Sauvage, J. P. Comparative studies of the photophysical properties of copper phenanthrolines: from Cu(dmp)₂⁺ to the copper(I) catenates. *Inorg. Chem.* **1989**, *28*, 4070-4072, 10.1021/ic00321a008.
- (30) Pallenberg, A. J.; Koenig, K. S.; Barnhart, D. M. Synthesis and Characterization of Some Copper(I) Phenanthroline Complexes. *Inorg. Chem.* **1995**, *34*, 2833-2840, 10.1021/ic00115a009.
- (31) Sheldrick, G. SHELXT - Integrated space-group and crystal-structure determination. *Acta Crystallographica Section A* **2015**, *71*, 3-8, doi:10.1107/S2053273314026370.
- (32) Sheldrick, G. Crystal structure refinement with SHELXL. *Acta Crystallographica Section C* **2015**, *71*, 3-8, doi:10.1107/S2053229614024218.
- (33) Hayes, D.; Kohler, L.; Hadt, R. G.; Zhang, X.; Liu, C.; Mulfort, Karen L.; Chen, L. X. Excited state electron and energy relays in supramolecular dinuclear complexes revealed by ultrafast optical and X-ray transient absorption spectroscopy. *Chem. Sci.* **2018**, *9*, 860-875, 10.1039/c7sc04055e.
- (34) te Velde, G.; Bickelhaupt, F. M.; Baerends, E. J.; Fonseca Guerra, C.; van Gisbergen, S. J. A.; Snijders, J. G.; Ziegler, T. Chemistry with ADF. *Journal of Computational Chemistry* **2001**, *22*, 931-967, 10.1002/jcc.1056.

Table Of Content



Caption: "A stable copper(I) complex was prepared, whose steric hindrance is halfway between benchmark $[\text{Cu}(\text{dipp})_2]^+$ and $[\text{Cu}(\text{dtbp})_2]^+$. The coordination sphere of the complex is not geometrically symmetric and this causes a geometric imbalance, which favors rotational and rocking motions of the ligands in the excited state, causing the premature quenching of the excited state."