



**HAL**  
open science

## Sulfur disproportionation is exergonic in the vicinity of marine hydrothermal vents

Karine Alain, Heidi Aronson, Maxime Allieux, Stéven Yvenou, Jan Amend

### ► To cite this version:

Karine Alain, Heidi Aronson, Maxime Allieux, Stéven Yvenou, Jan Amend. Sulfur disproportionation is exergonic in the vicinity of marine hydrothermal vents. *Environmental Microbiology*, 2022, *Microbial Ecology of Aquatic Systems*, 24 (5), pp.2210-2219. 10.1111/1462-2920.15975 . hal-03690392v2

**HAL Id: hal-03690392**

**<https://hal.science/hal-03690392v2>**

Submitted on 6 Jul 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Correspondence

## Sulfur disproportionation is exergonic in the vicinity of marine hydrothermal vents

Karine Alain <sup>1\*</sup>, Heidi S. Aronson,<sup>2†</sup> Maxime Allieux,<sup>1§</sup> Stéven Yvenou<sup>1§</sup> and Jan P. Amend <sup>2,3</sup>


<sup>1</sup>Univ Brest, CNRS, Ifremer, Unité Biologie et Ecologie des Ecosystèmes marins Profonds BEEP, UMR 6197, IRP 1211 MicrobSea, IUEM, Rue Dumont d'Urville, Plouzané, F-29280, France.

<sup>2</sup>Department of Biological Sciences, University of Southern California, Los Angeles, CA, 90089.

<sup>3</sup>Department of Earth Sciences, University of Southern California, Los Angeles, CA, 90089.

## Introduction

Sulfur compounds are ubiquitous in marine hydrothermal environments and several are central in catabolic reactions involved in primary production, recycling and terminal degradation of organic matter (Cao *et al.*, 2014). Sulfur is found in various oxidation states ( $\text{H}_2\text{S}$ ,  $\text{FeS}$ ,  $\text{FeS}_2$ ,  $\text{S}^0$ ,  $\text{SO}_3^{2-}$ ,  $\text{S}_2\text{O}_3^{2-}$ ,  $\text{S}_n\text{O}_6^{2-}$ ,  $\text{S}_n\text{O}_6^-$ ,  $\text{SO}_4^{2-}$ , etc.) in the chimney structures, vent fluids, surrounding seawater, and in pore spaces and fractures of the seafloor, and consequently used as an electron donor or terminal electron acceptor in aerobic and anaerobic reactions. Sulfide/sulfur-oxidizing, sulfur-reducing and sulfate-reducing microbial taxa of hydrothermal habitats are well known (Zeng *et al.*, 2021; Godfroy *et al.*, 2022), and models of the biogenic hydrothermal cycle of sulfur are based on these reactions. Another redox reaction involving sulfur species takes place in this unique ecosystem: microbial sulfur disproportionation (=dismutation) (MSD), most commonly represented as



In this reaction, an intermediate valence inorganic sulfur compound is simultaneously reduced and oxidized into sulfide and sulfate, respectively (Bak and Cypionka, 1987; Finster, 2008; Slobodkin and Slobodkina, 2019). In addition to  $\text{S}^0$ , other studied substrates for MSD include thiosulfate ( $\text{S}_2\text{O}_3^{2-}$ ) and sulfite ( $\text{SO}_3^{2-}$ ) (Finster, 2008; Slobodkin and Slobodkina, 2019). MSD, also called inorganic fermentation, is poorly documented in hydrothermal ecosystems and the first hydrothermal microbial taxa able to disproportionate inorganic sulfur compounds under anoxic conditions have been isolated quite recently (Slobodkin *et al.*, 2012; Slobodkin *et al.*, 2013; Slobodkina *et al.*, 2016; Slobodkina *et al.*, 2017; Frolova *et al.*, 2018).

The metabolic pathways and enzymatic mechanisms of anaerobic sulfur disproportionation are only partially resolved (Finster, 2008; Mardanov *et al.*, 2016; Slobodkin and Slobodkina, 2019; Allieux *et al.*, 2020). A few archaea and bacteria, including one hydrothermal vent bacterium *Aquifex aeolicus*, are also capable of oxygen-dependent disproportionation of  $\text{S}^0$  using the SOR (sulfur oxygenase reductase) enzymatic system (whose disproportionation sub-reaction is identical to Reaction 1 noted above) (Kletzin *et al.*, 2004; Pelletier *et al.*, 2008; Bertran *et al.*, 2020). Although it has not been comprehensively investigated in hydrothermal environments, MSD is an important contributor to the sulfur cycle in other marine ecosystems and may prove to be integral to the hydrothermal sulfur cycle as well. Microorganisms capable of disproportionating intermediate sulfur species are widespread in marine surface sediments and are partly responsible for the isotopic signatures of sulfidic minerals in young and old sediments (Bak and Pfenning, 1987; Thamdrup *et al.*, 1993; Finster, 2008; Jørgensen *et al.*, 2019). Using radioisotope tracing of thiosulfate, it has even been shown that a significant proportion of the thiosulfate present in oxidized and reduced sediments is disproportionated (Bak and Cypionka, 1987; Fossing and Jørgensen, 1990; Jørgensen, 1990). However, the precise extent and distribution of the MSD process in natural environments are not known because (i) there is no specific genetic marker for this reaction, which prevents any

Received 21 December, 2021; revised 2 March, 2022; accepted 12 March, 2022. \*For correspondence. E-mail [karine.alain@univ-brest.fr](mailto:karine.alain@univ-brest.fr); Tel. (+33)2 98 49 88 53; Fax +33(2) 98 49 87 05. †These authors share first authorship. §These authors have contributed equally to this work.

application of genomic investigation methods to identify and quantify taxa implementing this process in natural habitats, because (ii) intermediate sulfur species are transient and generally short-lived *in situ* (Zopfi *et al.*, 2004; Jørgensen *et al.*, 2019), and (iii) because it is difficult to disentangle the relative magnitude of different sulfur compound-based redox metabolisms from the net isotopic signals (Jørgensen *et al.*, 2019; Allieux *et al.*, 2020).

In this context, calculations of the Gibbs energy of reaction ( $\Delta G_r$ ) at environmentally relevant conditions can identify whether MSD is thermodynamically favourable and consequently point to niches where sulfur disproportionators could be present and thriving. A common message in the literature is that MSD yields minimal energy or is endergonic (energy consuming) (Finster, 2008). It should be noted, however, that this claim is typically (and incorrectly) based on the value of the standard state Gibbs energy ( $\Delta G_r^0$ ) for Reaction 1 at 25°C and 1 bar, which is +120.5 kJ mol<sup>-1</sup> (Amend and LaRowe, 2019). At low concentrations, and hence activities, of sulfate and sulfide, together with circumneutral to alkaline pHs, Reaction 1 can indeed be exergonic (Amend and LaRowe, 2019).

Deep-sea hydrothermal vents are seabed manifestations of hydrothermal convection of seawater through the permeable ocean floor, driven by heat from magma or subsurface rock (Humphris and Klein, 2018). These ecosystems occur in a range of tectonic settings, such as mid-ocean ridges, back-arc spreading basins, and volcanoes and seamounts at hotspots. They are characterized by a variety of fluid compositions, with acidic or alkaline pHs, and wide ranges of fluid temperatures (Orcutt *et al.*, 2011; Dick, 2019a). These singular habitats host a multitude of ecological niches with contrasting physico-chemical conditions along the mixing gradient between the venting fluid and seawater, allowing the settlement of luxuriant microbial and animal communities deriving their energy by chemosynthesis through oxidation–reduction reactions (Dick, 2019a). In addition, efficient sulfide scavenging manganese (IV) and iron (III) minerals can also increase the energy yield of MSD in these ecosystems (Slobodkin and Slobodkina, 2019).

In this article, we use a thermodynamic approach and a range of physico-chemical parameters (*i.e.* temperature, pressure, salinity, pH, sulfate, sulfide) to predict the Gibbs energy of reaction for S<sup>0</sup> disproportionation in seawater. We then calculate the free-energy values of S<sup>0</sup> disproportionation from environmental data at four hydrothermal sites located in different geological settings and characterized by contrasted fluid compositions. We show that S<sup>0</sup> disproportionation is energetically favourable in most, if not all, niches of the hydrothermal vent ecosystem, whatever the geological and geochemical context and depth.

## Results and discussion

In marine hydrothermal vents, particulate elemental sulfur is actively formed by oxidation of hydrogen sulfide during mixing of hydrothermal fluid with oxygenated seawater (McCollom, 2000), at interface of reduced free hydrogen sulfide with other oxidants (Fe<sup>3+</sup> or NO<sub>3</sub><sup>-</sup>) (Zopfi *et al.*, 2004), and by weathering of sulfide minerals (<https://www.mindat.org/>). To date, only six sulfur disproportionating bacteria have been isolated from marine hydrothermal vents, with five of them isolated in the last decade (Zillig *et al.*, 1986; Slobodkin *et al.*, 2012; Slobodkin *et al.*, 2013; Slobodkin *et al.*, 2016; Slobodkina *et al.*, 2016; Slobodkina *et al.*, 2017; Frolova *et al.*, 2018; Bertran *et al.*, 2020). At this point, we have limited knowledge about how much microbial disproportionation of sulfur contributes to the biotic sulfur cycle of hydrothermal ecosystems. In this work, we have tried to provide some clues to this question through the lens of thermodynamics, by performing two series of calculations, which are described below.


### Free-energy yield of sulfur disproportionation in seawater

In a first set of calculations, we determined the free-energy change of sulfur disproportionation (Reaction 1) across a broad range of physicochemical parameters (Table 1) commonly observed in hydrothermal systems. In Fig. 1A–E, we depict the energy yields as a function of pressure, salinity, pH, activity of sulfide and activity of sulfate, respectively, *versus* temperature; in Fig. 1F, energy yields are depicted as functions of activities of sulfide *versus* sulfate. Note that the other geochemical parameters (*i.e.* those not represented on the axes) are fixed at levels given in Table 1. It can be seen in these six panels that  $\Delta G_r < 0$  (exergonic) for Reaction 1 approaches and even exceeds energy yields of 100 kJ mol<sup>-1</sup> at certain conditions. Pressure even up to

**Table 1.** Gradients and fixed data used for Gibbs energy calculations for sulfur disproportionation in seawater.

	Gradient analyses	Fixed data
Temperature (°C)	0–130	25
Salinity (g L <sup>-1</sup> )	0–40	545 mmol kg <sup>-1</sup>
Pressure (MPa)	0–120	26 <sup>a</sup>
pH	0–12	7
Total sulfide concentration (mM)	0–10	0
Total sulfate concentration (mM)	0–30	28

<sup>a</sup>The average depth of mid-ocean ridges is 2600 m; we set the pressure at 26 MPa because it corresponds to the hydrostatic pressure at this depth.



**Fig. 1.** Values of  $\Delta G_r$  for elemental sulfur disproportionation (Reaction 1) as a function of pressure (A), salinity (B), pH (C), sulfide (D) and sulfate (E) versus temperature, and sulfide versus sulfate (F), calculated with the Equation  $\text{S}^0 + \text{H}_2\text{O} = 0.25\text{SO}_4^{2-} + 0.75\text{H}_2\text{S}(\text{aq}) + 0.5\text{H}^+$  (Eq. 1bis). Geochemical parameters, when not allowed to vary along the x- and y-axes, are fixed as follows:  $T = 25^\circ\text{C}$ , pressure = 26 MPa, salinity = 545 mmol  $\text{kg}^{-1}$ , pH = 7.0, sulfide concentration = 0 mM, sulfate concentration = 28 mM. Note that the colour scale changes for each panel in this figure.

120 MPa (Fig. 1A) has little to no effect on  $\Delta G_r$  of this reaction. Similarly, salinity (Fig. 1B), which affects the ionic strength of the system and hence the activity coefficients (see below), has only a tiny effect on the free-energy values of the reaction, with  $\Delta G_r$  slightly more exergonic at elevated salinities. In Fig. 1C, it can be seen that pH has a large effect on  $\Delta G_r$  of  $\text{S}^0$  disproportionation, with energy yields increasing from  $<20 \text{ kJ mol}^{-1}$  at low temperature and pH  $< 2$  to near  $100 \text{ kJ mol}^{-1}$  at temperatures above  $100^\circ\text{C}$  and pH in the 10–12 range. The effects of variable sulfide and sulfate activities (and concentrations) are demonstrated, respectively, in Fig. 1D

and E. In both cases, values of  $\Delta G_r$  change demonstrably, with relatively low energy yields at low temperature and high sulfide or sulfate levels, and ever higher energy yields as temperature increases and sulfide or sulfate levels drop. Lastly, the coupled effects of sulfide and sulfate activities on  $\Delta G_r$  of  $\text{S}^0$  disproportionation can be seen in Fig. 1F. Consistent with observations in panels D and E, the energy yields increase by  $\sim 20 \text{ kJ mol}^{-1}$  as the activities of both of these aqueous sulfur species decrease by several orders of magnitude.

Overall, these careful but still rough predictions made on the basis of seawater ionic composition suggest that

microorganisms can grow *via* sulfur disproportionation under almost any combination of physicochemical conditions tested here, that cover all conditions encountered in hydrothermal systems, except when sulfide concentration reaches 10 mM in a niche where the temperature is below 5°C. At very acidic pH ( $\text{pH} \leq 2$ ) and low temperature ( $T^\circ \leq 25^\circ\text{C}$ ), free-energy change values are close to the microbial Gibbs energy minimum amount for a catabolic reaction ( $\Delta G_{\text{min}}$ , cost of minimal quantum of ATP synthesis – less than one-third of ATP unit), somewhere in between  $-9$  and  $-20$  kJ mole $^{-1}$  of reaction turnover (Schink, 1997; Hoehler, 2004). Conversely, at certain conditions (e.g.  $T^\circ > 100^\circ\text{C}$  and  $\text{pH} > 10$ , or  $T^\circ > 110^\circ\text{C}$  and sulfate concentration  $< 5$  mM),  $\text{S}^0$ -disproportionation free-energy yields can be significant and reach or exceed 100 kJ mol $^{-1}$ . In summary, these predictions show that this catabolic reaction is thermodynamically more favourable at high temperature, high pH, and low sulfide and low sulfate concentrations, regardless of depth. Nevertheless, these predictions do not lead to the conclusion that the reaction will actually occur *in situ* under all these combinations of conditions, as microbial systems may face additional biological and physiological challenges to thrive in their natural extreme habitat. In the current state of knowledge, five thermophilic anaerobic  $\text{S}^0$ -disproportionating bacteria, and one hyperthermophilic microaerophilic  $\text{S}^0$ -disproportionating bacterium, all with optimal growth pH close to neutrality, have been isolated from marine hydrothermal environments (Huber and Stetter, 2001; Slobodkin *et al.*, 2012; Slobodkin *et al.*, 2013; Slobodkina *et al.*, 2016; Slobodkina *et al.*, 2017; Frolova *et al.*, 2018) and three other anaerobic mesophilic ones are being characterized (Alain, personal communication). These strains have all been isolated and are all growing under conditions predicted here to be favourable for sulfur disproportionation, and their growth is enhanced in the presence of sulfide scavengers, predicted by thermodynamics to be more favourable. These thermodynamic calculations encourage the search for hydrothermal sulfur disproportionators at more acidic or more alkaline pHs than those tested to date, and to lower sulfate and sulfide concentrations in cultures.

To further refine these thermodynamic predictions, the free-energy yields of  $\text{S}^0$ -disproportionation were calculated from real environmental data.

*Free-energy yield of  $\text{S}^0$  disproportionation along a dilution gradient from vent fluid to seawater: example of four contrasted vent sites*

In a second set of calculations, we calculated the free-energy change of elemental sulfur disproportionation (Reaction 1) at various points in the mixing zone between pure seawater and pure hydrothermal fluid for four well-


characterized hydrothermal sites. These calculations were carried out, at *in situ* pressure, from the compositions of hydrothermal fluids and seawater measured *in situ*. Given that there are 721 hydrothermal fields identified to date (confirmed or inferred, active and inactive fields) (InterRidge database Version 3.4. PANGAEA, <https://doi.org/10.1594/PANGAEA.917894>), we selected active hydrothermal sites located in diverse marine geological and tectonic settings and characterized by contrasted fluid compositions to have an overview of the different types of existing hydrothermal systems.

The first sets of calculation were done for the TAG active mound located on the low spreading Mid-Atlantic Ridge. This massive sulfide deposit is located on 100 000 year-old basaltic crust at 3620 m water depth (Thompson *et al.*, 1988). Hydrothermal fluids at this site are emitted in the form of iron-rich acidic black smokers at a temperature of 369°C and a pH of 3.4 (at 25°C) (Humphris and Klein, 2018). The main dissolved gas is  $\text{H}_2\text{S}$  and the concentrations of  $\text{CO}_2$ ,  $\text{CH}_4$  and  $\text{H}_2$  are low (Humphris and Klein, 2018). The second set of calculations focused on the Fenway felsic hydrothermal site, located at 1710 m depth in the PACMANUS hydrothermal system in the eastern part of the Manus back-arc basin (Western Pacific). Fluids of the large black smoker chimney complex at the summit of this system are characterized by a high emission temperature (358°C), a very acidic pH (2.7 at 25°C) and high concentrations of dissolved  $\text{CO}_2$  and  $\text{H}_2\text{S}$  (Reeves *et al.*, 2011; Humphris and Klein, 2018). The third round of calculations concerned the Rainbow vent field, located in ultramafic rocks, at 2300 m at the Mid-Atlantic Ridge. The Rainbow fluids are very acidic (pH 2.99 at 25°C), hot temperature (367°C), rich in dissolved  $\text{H}_2$  and with very low Fe and Mn content (Douville *et al.*, 2002; Humphris and Klein, 2018). The last calculations series focused on the Upper Cone of Brothers submarine volcano, located along the Kermadec arc, offshore New Zealand, at 1220 m depth. The end-member fluid of this white smoker is characterized by a mild temperature (122°C), an extremely low pH (1.9 at 25°C), a high concentration of Mg, presence of dissolved  $\text{CO}_2$  and no detectable  $\text{H}_2\text{S}$  in the selected dataset (de Ronde *et al.*, 2005; Humphris and Klein, 2018). The presence of native sulfur was reported for the TAG, PACMANUS and Brothers volcano sites (<https://www.mindat.org/>). Unfortunately, no complete geochemical data set of ultrabasic end-member fluid was available to perform such calculations.

As seen in Fig. 2, Reaction 1 is exergonic at all four of these sites along mixing zones between pure seawater and pure hydrothermal fluid. It should be noted that at some of these points (e.g. pure vent fluid; 75% vent fluid for all sites; 50% vent fluid at TAG, Fenway, and Rainbow) the temperatures are above the apparent limit for

life. At present, microbial growth has not been demonstrated at temperature above 122°C (at 20 MPa, in the archaeon *Methanopyrus kandleri* strain 116) (Takai *et al.*, 2008) but could exist a little beyond, up to the temperature that would allow the maintenance of the integrity of the genome and proteome, of the bonds governing the structure of macromolecules and of the precise 3D conformation of proteins with the degree of flexibility necessary to perform biological functions. On the outermost

parts of the chimney edifices, where the temperature allows life, the  $S^0$  disproportionation reaction is exergonic in the four examples studied here, providing  $\Delta G_r$  ranging from  $\sim -6$  to  $< -42$  kJ mol $^{-1}$  (Fig. 2). The lowest values, between  $-6$  kJ mol $^{-1}$  and  $-13$  kJ mol $^{-1}$ , are close to the critical free-energy amount for a catabolic reaction ( $\Delta G_{\min}$ ) reported so far for microbial metabolisms in field and culture studies (Hoehler, 2004), and well above the power minimum for microbial cells to remain viable or



**Fig. 2.** Gibbs energies of sulfur disproportionation in the mixing zone between pure fluid and seawater, in various geological settings. A. Schematic representation of the gradient points used to make the calculations from seawater and fluid compositions measured *in situ*. B. Temperature, log activities of H<sup>+</sup>, H<sub>2</sub>S, and SO<sub>4</sub><sup>2-</sup>, standard state Gibbs energies, and Gibbs energies of sulfur disproportionation (kJ mol $^{-1}$  S) along a dilution gradient from seawater to pure hydrothermal vent fluid at TAG, Fenway Basin, Rainbow Vent, and Brothers Volcano Upper Cone. The suffixes after the vent names refer to the ratio of vent fluid to seawater (*i.e.* TAG.75 is a mixture of 75% vent fluid and 25% seawater). The standard state Gibbs energy and Gibbs energy of Fenway Basin hydrothermal vent fluid are listed as NA because it is below the minimum density for applicability.

**Table 2.** Iron (III) minerals present in the hydrothermal vent fields of the four sites studied

TAG	PACMANUS (Fenway)	Rainbow	Brothers volcano (upper cone)
Goethite ( $\alpha\text{-FeO(OH)}$ )	Ferrihydrite ( $\text{Fe}^{3+}_{10}\text{O}_{14}(\text{OH})_2$ )	Magnetite ( $\text{Fe}^{2+}\text{Fe}^{3+}_2\text{O}_4$ )	Celadonite ( $\text{K}(\text{Mg}, \text{Fe}^{2+})$ ( $\text{Fe}^{3+}, \text{Al}$ ) [ $\text{Si}_4\text{O}_{10}$ ]( $\text{OH}$ ) <sub>2</sub> )
Haematite ( $\alpha\text{-Fe}_2\text{O}_3$ )	Goethite ( $\alpha\text{-FeO(OH)}$ )	Limonite ( $\text{FeO(OH)} \cdot n\text{H}_2\text{O}$ )	Goethite ( $\alpha\text{-FeO(OH)}$ )
Jarosite ( $\text{KFe}^{3+}_2(\text{SO}_4)_2(\text{OH})_6$ )	Haematite ( $\alpha\text{-Fe}_2\text{O}_3$ )		Haematite ( $\alpha\text{-Fe}_2\text{O}_3$ )
Nontronite ( $\text{Na}_{0.3}\text{Fe}_2(\text{Si}, \text{Al})_4\text{O}_{10}$ ( $\text{OH}$ ) <sub>2</sub> · $n\text{H}_2\text{O}$ )	Nontronite ( $\text{Na}_{0.3}\text{Fe}_2(\text{Si}, \text{Al})_4\text{O}_{10}$ ( $\text{OH}$ ) <sub>2</sub> · $n\text{H}_2\text{O}$ )		Jarosite ( $\text{KFe}^{3+}_3(\text{SO}_4)_2(\text{OH})_6$ )
			Nontronite ( $\text{Na}_{0.3}\text{Fe}_2(\text{Si}, \text{Al})_4\text{O}_{10}$ ( $\text{OH}$ ) <sub>2</sub> · $n\text{H}_2\text{O}$ )

Source: <https://www.mindat.org/>.

subsist at ultra-low energy fluxes (LaRowe and Amend, 2015; Bradley *et al.*, 2020). Sulfur disproportionation is thus thermodynamically favourable in the vicinity of these four hydrothermal vents located in very different geological settings and provides energy yields of up to 42 kJ mol<sup>-1</sup>, based on calculations taking into account the *in situ* geochemical composition of the seawater and hydrothermal fluid, and a dilution model of the two solutions. These energy yields are rather low but sufficient to allow microbial growth. Nevertheless, given that chimney rocks contain various minerals that can strongly chelate sulfides, such as manganese (IV) and iron (III) minerals (Thamdrup *et al.*, 1993; Ollivier *et al.*, 2018; Saji, 2021), and other less potent metal-based chelating minerals (containing Zn, Cu, Fe(II), Co, Ni, Mg, etc.) (Saji, 2021), which were not included in our calculations, the amount of energy released by the reaction could be even higher inside the chimney walls, since the energy yield increases significantly as the sulfide concentration decreases. Indeed, chimneys at all four sites were reported to contain two to five iron(III) mineral species (Table 2), and numerous less powerful metal-based chelating minerals (<https://www.mindat.org/>), that could chelate the sulfide *in situ* and make the reaction even more energy-producing locally (Thamdrup *et al.*, 1993; Ollivier *et al.*, 2018). Because the composition of the chimney walls evolves with time and mineral precipitation and is uneven (e.g. Meng *et al.*, 2019), it is difficult to make more accurate predictions without accurate mapping of the mineral composition of the chimney walls on a sub-millimetre scale at a given time. However, one can speculate that the reaction is likely much more exergonic locally in the chimney walls, in pores and fractures containing sulfide-chelating minerals, than calculated here, as the  $\Delta G_r$  of sulfur disproportionation changes demonstrably with decreasing sulfide levels (Fig. 1D).

#### Implications of an exergonic microbial sulfur disproportionation reaction

The thermodynamic approach implemented here indicates that MSD is sufficiently exergonic in most, if not all,

marine hydrothermal niches at non-lethal temperatures, to drive cellular functions. This information is crucial because in the current absence of genetic markers of this process and given the difficulty of interpreting *in situ* isotopic data of sulfur species, it was not known whether this process could take place in this ecosystem. These results based on thermodynamics indicate that the reaction is possible and favourable but do not prove that it occurs *in situ*. Strains isolated from hydrothermal ecosystems capable of disproportionating S<sup>0</sup>, among other things, could express this catabolism *in situ* to grow. In the future, it will be crucial to be able to document this process *in situ* and to determine what is the contribution of MSD to the fluxes of sulfur species in hydrothermal habitats compared with those of sulfur-oxidation and sulfate-reduction, as MSD is confused with these pathways in global budgets, since it leads to the production of sulfates and sulfides. Sulfur-disproportionating taxa do not necessarily have the same ecophysiological properties as sulfur-oxidizers and sulfate-reducers, and this could have a significant impact on our understanding of the biotic cycle of sulfur. At present, with exception of *Dissulfurimicrobium hydrothermale* Sh68<sup>T</sup>, sulfur disproportionating strains isolated from marine hydrothermal ecosystems are all capable of using other more energy-yielding catabolic pathways (sulfate reduction, or dissimilatory nitrate reduction to ammonium DNRA) to produce energy and to grow (Slobodkin and Slobodkina, 2019). It will be important to determine whether MSD is used in the same way as the other catabolic pathways and confers a metabolic versatility to the taxa allowing them to adapt to the changing physico-chemical conditions of their natural habitat, or if it is simply an accessory metabolism used by the microorganisms to wait for more favourable conditions as this process is vastly less energetically favourable than sulfate reduction or DNRA.

#### Conclusion

Hydrothermal habitats are characterized by numerous geochemical and geochemical/lithological interfaces containing reduced free hydrogen sulfide on one side and

strong oxidants on the other, where intermediate sulfide species (*i.e.* substrates for sulfur disproportionation) are thus produced. Despite the popular belief that  $S^0$  disproportionation is endergonic under standard conditions, the predictions made here from theoretical gradients and real data sets from a variety of hydrothermal geological settings indicate that this reaction is exergonic under the vast majority of physicochemical conditions tested. These calculations show that this reaction could produce enough energy to sustain growth in all hydrothermal niches of the four hydrothermal sites studied where the temperature is compatible with life. This reaction could therefore be widespread in hydrothermal habitats, provided that the cellular systems are adapted to environmental constraints. Furthermore, as the first set of calculations of free-energy changes in  $S^0$  disproportionation were performed over very broad gradients of physical parameters and chemical species, these theoretical data could also inform the occurrence of this process in other ecosystems, such as seafloor environments, for example. The sulfur cycle is known to play a critical role in the redox balance of the ocean due to the multiple oxidation states of this element, and furthermore marine hydrothermal vents are widespread on the seafloor and the seafloor ecosystem represents one of the largest biomes on Earth (Orcutt *et al.*, 2011). By affecting the sulfur cycle, microbial activities in these ecosystems, including sulfur disproportionation, could influence the oxidation state of the global ocean and, by extension, the Earth's climate (Bowles *et al.*, 2014; D'Hondt *et al.*, 2019).

## Experimental procedures

### Thermodynamic modelling

The Gibbs energy yields ( $\Delta G_r$ ) for sulfur disproportionation (Reaction 1) was calculated with the relation

$$\Delta G_r = \Delta G_r^0 + RT \ln Q_r \quad (2)$$

where  $\Delta G_r^0$  is as defined above,  $R$  indicates the universal gas constant,  $T$  stands for the temperature in Kelvin, and  $Q_r$  refers to the reaction activity quotient.  $\Delta G_r^0$  values were calculated at the temperature and pressure of interest with the revised Helgeson–Kirkham–Flowers (HKF) equations of state (Helgeson *et al.*, 1981; Tanger and Helgeson, 1988; Shock *et al.*, 1992) using the 'subcrt' command from the R software package CHNOSZ v1.4.1 (Dick, 2019b). Thermodynamic data in CHNOSZ are derived from the OrganoBioGeoTherm database (Helgeson *et al.*, 2009), and the sources of these data are provided in the CHNOSZ package documentation. Values of  $Q_r$  were calculated with the relation

$$Q_r = \prod a_i^{v_{i,r}} \quad (3)$$

where  $a_i$  represents the activity of the  $i$ th species raised to its stoichiometric reaction coefficient  $v_{i,r}$  in the  $r$ th reaction, which is positive for products and negative for reactants. Concentration data were converted to chemical activities using

$$a_i = m_i \gamma_i \quad (4)$$

where  $m_i$  and  $\gamma_i$  are the molality and activity coefficients of the  $i$ th species. Activity coefficients were calculated using the aqueous speciation package AqEquil v0.9.1 (Boyer *et al.*, 2021), which interfaces with the geochemical speciation software EQ3/6 (Wolery, 1979). The activities of pure minerals [elemental sulfur ( $S^0$ )] and water were taken to be unity ( $a_i = 1$ ).  $\Delta G_r$  represents the Gibbs energy of reaction under *in situ* conditions and is expressed in units of kilojoules per mole of sulfur.

### In situ data and gradient analyses

Gibbs energy yields for sulfur disproportionation were calculated using concentration and temperature data from seawater, Manus Basin (Fenway), Brothers Volcano Upper Cone, Mid-Atlantic Ridge Trans-Atlantic Geotraverse (TAG), and Mid-Atlantic Ridge Rainbow vents (Humphris and Klein, 2018). Seawater concentrations which were reported as below detection limit were assumed to be at the detection limit (*i.e.*  $0.001 \mu\text{mol kg}^{-1}$  was used in place of  $<0.001 \mu\text{mol kg}^{-1}$ ). Zero values were reported as  $10^{-9}$  mol. Pressures were assumed to be 36 MPa for TAG vent field, 17 MPa for Fenway, 12 MPa for Brothers volcano, and 23 MPa for Rainbow, on the basis of the water depth at which these hydrothermal fields are located.

Gibbs energy yields for sulfur disproportionation were calculated along a linear dilution gradient from pure vent fluid to pure seawater at the four sites using endmember values as well as vent fluid:seawater mixing ratios of 75:25, 50:50, and 25:75. Calculated temperatures of the mixed fluids incorporated the specific heat capacity of water at that temperature, following the method described in the study by Lu *et al.* (2021). Activity coefficients from these dilution gradients were calculated after speciating with AqEquil. Energy yields for disproportionation in seawater were also calculated as functions of various geochemical parameters while holding other variables fixed (Table 1). Activity coefficients were recalculated using AqEquil v0.9.1 based on the parameters for each gradient. The CHNOSZ v1.4.1 'mosaic' function was used to speciate  $H_2S$  and  $SO_4^{2-}$  for energy yields that were calculated as a function of pH.



The mineral composition of the hydrothermal edifices is based on the open mineral database [mindat.org](http://mindat.org) (<https://www.mindat.org/>).

### Acknowledgements

This work was supported by the Sino-French IRP 1211 MicrobSea to K.A., by the ISblue project, Interdisciplinary graduate school for the blue planet (ANR-17-EURE-0015) and co-funded by a grant from the French government under the programme 'Investissements d'Avenir' (Theme 2, project DISMUT) to K.A. The study was supported by two grants from the French Ministry of Higher Education and Research, and from the Region Bretagne, to M.A. and S.Y. The authors would like to thank Doug LaRowe for helpful discussions on fluid mixing. The authors also thank two anonymous reviewers whose insightful comments have greatly improved this contribution.

### Author Contributions

K.A., S.Y. and M.A. designed the experiment; H.S.A. performed the calculations. J.P.A. supervised the calculations methods. K.A. and J.P.A supervised the project. All authors interpreted the data; K.A. wrote the manuscript. All authors reviewed the manuscript.

### References

- Allioux, M., Yvenou, S., Slobodkina, G., Slobodkin, A., Shao, Z., Jebbar, M., and Alain, K. (2020) Genomic characterization and environmental distribution of a Thermophilic anaerobe *Dissulfurirhabdus thermomarina* SH388<sup>T</sup> involved in disproportionation of sulfur compounds in shallow sea hydrothermal vents. *Microorganisms* **8**: 1132. <https://doi.org/10.3390/microorganisms8081132>.
- Amend, J.P., and LaRowe, D.E. (2019) Minireview: demystifying microbial reaction energetics. *Environ Microbiol* **21**: 3539–3547. <https://doi.org/10.1111/1462-2920.14778>.
- Bak, F., and Cypionka, H. (1987) A novel type of energy metabolism involving fermentation of inorganic sulphur compounds. *Nature* **326**: 891–892. <https://doi.org/10.1038/326891a0>.
- Bak, F., and Pfennig, N. (1987) Chemolithotrophic growth of *Desulfovibrio sulfodismutans* sp. nov. by disproportionation of inorganic sulfur compounds. *Arch Microbiol* **147**: 184–189.
- Bertran, E., Ward, L.M., and Johnston, D.T. (2020) Draft genome sequence of *Acidianus ambivalens* DSM 3772, an aerobic Thermoacidophilic sulfur Disproportionator. *Microbiol Resour Announc* **9**: e01415–e01419. <https://doi.org/10.1128/MRA.01415-19>.
- Bowles, M.W., Mogollon, J.M., Kasten, S., Zabel, M., and Hinrichs, K.-U. (2014) Global rates of marine sulfate reduction rates and implications for seafloor metabolic activities. *Science* **344**: 889–891. <https://doi.org/10.1126/science.1249213>.
- Boyer, G., Ely, T., and Shock, E.. (2021) AqEquil v0.9.1 (v0.9.1). Zenodo. <https://doi.org/10.5281/zenodo.5534832>
- Bradley, J.A., Arndt, S., Amend, J.P., Burwicz, E., Dale, A. W., Egger, M., and LaRowe, D.E. (2020) Widespread energy limitation to life in global seafloor sediments. *Sci Adv* **6**: eaba0697. <https://doi.org/10.1126/sciadv.aba0697>.
- Cao, H., Wang, Y., Lee, O.O., Zeng, X., Shao, Z., and Qian, P.Y. (2014) Microbial sulfur cycle in two hydrothermal chimneys on the southwest Indian ridge. *MBio* **5**: e00980–e00913. <https://doi.org/10.1128/mBio.00980-13>.
- D'Hondt, S., Inagaki, F., Orcutt, B.N., and Hinrichs, K.-U. (2019) IODP advances in understanding of seafloor life. *Oceanography* **32**: 198–207. <https://doi.org/10.5670/oceanog.2019.146>.
- de Ronde, C.E.J., Hannington, M.F., Stoffers, P., Wright, I. C., Ditchburn, R.G., Reyes, A.G., et al. (2005) Evolution of a submarine magmatic-hydrothermal system: brothers volcano, southern Kermadec arc, New Zealand. *Econ Geol* **100**: 1097–1133.
- Dick, G.J. (2019a) The microbiomes of deep-sea hydrothermal vents: distributed globally, shaped locally. *Nat Rev Microbiol* **17**: 271–283. <https://doi.org/10.1038/s41579-019-0160-2>.
- Dick, J.M. (2019b) CHNOSZ: thermodynamic calculations and diagrams for geochemistry. *Front Earth Sci* **7**: 1–18. <https://doi.org/10.3389/feart.2019.00180>.
- Douville, E., Charlou, J.L., Oelkers, E.H., Bienvu, P., Colon, C.F.J., Donval, J.P., et al. (2002) The rainbow vent fluids (36°14'N, MAR): the influence of ultramafic rocks and phase separation on trace metal content in mid-Atlantic ridge hydrothermal fluids. *Chem Geol* **184**: 37–48. [https://doi.org/10.1016/S0009-2541\(01\)00351-5](https://doi.org/10.1016/S0009-2541(01)00351-5).
- Finster, K. (2008) Microbiological disproportionation of inorganic sulfur compounds. *J Sulfur Chem* **29**: 281–292. <https://doi.org/10.1080/17415990802105770>.
- Fossing, H., and Jørgensen, B.B. (1990) Oxidation and reduction of radiolabeled inorganic sulfur compounds in an estuarine sediment, Kysing Fjord, Denmark. *Geochim Cosmochim Acta* **54**: 2731–2742.
- Frolova, A.A., Slobodkina, G.B., Baslerov, R.V., Novikov, A. A., Bonch-Osmolovskaya, E.A., and Slobodkin, A.I. (2018) *Thermosulfurimonas marina* sp. nov., an autotrophic sulfur-disproportionating and nitrate-reducing bacterium isolated from a shallow-sea hydrothermal vent. *Microbiol* **87**: 502–507. <https://doi.org/10.1134/S0026261718040082>.
- Godfroy, A., François, D., Hartunians, J., Moalic, Y., and Alain, K. (2022) Physiology, metabolism and ecology of thermophiles from deep-sea vents. In *The Microbiology of Deep-Sea*, Vetriani, C., and Giovannelli, D. (eds). New York, NY: Springer International (in press).
- Helgeson, H.C., Kirkham, D.H., and Flowers, G.C. (1981) Theoretical prediction of the thermodynamic behavior of aqueous electrolytes by high pressures and temperatures; IV, calculation of activity coefficients, osmotic coefficients, and apparent molal and standard and relative partial molal properties to 600 degrees C and 5kb. *Am J Sci* **281**: 1249–1516.
- Helgeson, H.C., Richard, L., McKenzie, W.F., Norton, D.L., and Schmitt, A. (2009) A chemical and thermodynamic

- model of oil generation in hydrocarbon source rocks. *Geochim Cosmochim Acta* **73**: 594–695.
- Hoehler, T.M. (2004) Biological energy requirements as quantitative boundary conditions for life in the subsurface. *Geobiology* **2**: 205–215.
- Huber, R., and Stetter, K.O. (2001) Genus I. *Aquifex* Huber and Stetter 1992e, 656VP. In *Bergey's Manual of Systematic Bacteriology (The Archaea and the deeply branching and phototrophic Bacteria)*, 2nd ed, Vol. **1**, Boone, D.R., Castenholz, R.W., and Garrity, G.M. (eds). New York: Springer-Verlag, pp. 360–362.
- Humphris, S.E., and Klein, F. (2018) Progress in deciphering the controls on the geochemistry of fluids in seafloor hydrothermal systems. *Annu Rev Mar Sci* **10**: 315–343.
- Jørgensen, B.B. (1990) A thiosulfate shunt in the sulfur cycle of marine sediments. *Science* **249**: 152–154. <https://doi.org/10.1126/science.249.4965.152>.
- Jørgensen, B.B., Findlay, A.F., and Pellerin, A. (2019) The biogeochemical sulfur cycle of marine sediments. *Front Microbiol* **10**: 849. <https://doi.org/10.3389/fmicb.2019.00849>.
- Kletzin, A., Urich, T., Müller, F., Bandejas, T.M., and Gomes, C.M. (2004) Dissimilatory oxidation and reduction of elemental sulfur in thermophilic archaea. *J Bioenerg Biomembr* **36**: 77–91. <https://doi.org/10.1023/B:JOB.0000019600.36757.8c>.
- LaRowe, D.E., and Amend, J.P. (2015) Power limits for microbial life. *Front Microbiol* **6**: 718. <https://doi.org/10.3389/fmicb.2015.00718>.
- Lu, G.S., LaRowe, D.E., and Amend, J.P. (2021) Bioenergetic potentials in terrestrial, shallow-sea and deep-sea hydrothermal systems. *Chem Geol* **583**: 120449. <https://doi.org/10.1016/j.chemgeo.2021.120449>.
- Mardanov, A.V., Beletsky, A.V., Kadnikov, V.V., Slobodkin, A.I., and Ravin, N.V. (2016) Genome analysis of *Thermosulfurimonas dismutans*, the first thermophilic sulfur-disproportionating bacterium of the phylum *Thermodesulfobacteria*. *Front Microbiol* **7**: 950. <https://doi.org/10.3389/fmicb.2016.00950>.
- McCollom, T.M. (2000) Geochemical constraints on primary productivity in submarine hydrothermal vent plumes. *Deep-Sea Res Part I* **47**: 85–101. [https://doi.org/10.1016/S0967-0637\(99\)00048-5](https://doi.org/10.1016/S0967-0637(99)00048-5).
- Meng, X., Li, X., Chu, F., Fu, B., Lei, J., Li, Z., et al. (2019) Multi-stage growth and fluid evolution of a hydrothermal sulphide chimney in the East Pacific ridge 1–2° S hydrothermal field: constraints from in situ sulphur isotopes. *Geol Mag* **156**: 989–1002. <https://doi.org/10.1017/S0016756818000316>.
- Ollivier, B., Zeyen, N., Gales, G., Hickman-Lewis, K., Gaboyer, F., Benzerara, K., and Westall, F. (2018) Importance of prokaryotes in the functioning and evolution of the present and past geosphere and biosphere. In *Prokaryotes and Evolution*, Bertrand, J.C., Normand, P., Ollivier, B., and Sime-Ngando, T. (eds). Switzerland: Springer. [https://doi.org/10.1007/978-3-319-99784-1\\_3](https://doi.org/10.1007/978-3-319-99784-1_3).
- Orcutt, B.N., Sylvan, J.B., Knab, N.J., and Edwards, K.J. (2011) Microbial ecology of the dark ocean above, at, and below the seafloor. *Microbiol Mol Biol Rev* **75**: 361–422. <https://doi.org/10.1128/MMBR.00039-10>.
- Pelletier, N., Leroy, G., Guiral, M., Giudici-Ortoni, M.T., and Aubert, C. (2008) First characterisation of the active oligomer form of sulfur oxygenase reductase from the bacterium *Aquifex aeolicus*. *Extremophiles* **12**: 205–215. <https://doi.org/10.1007/s00792-007-0119-5>.
- Reeves, E.P., Seewald, J.S., Saccocia, P., Bach, W., Craddock, P.R., Shanks, W.C., et al. (2011) Geochemistry of hydrothermal fluids from the PACMANUS, northeast Pual and Vienna woods hydrothermal fields, Manus Basin, Papua New Guinea. *Geochim Cosmochim Acta* **75**: 1088–1123. <https://doi.org/10.1016/j.gca.2010.11.008>.
- Saji, V.S. (2021) Research advancements in sulfide scavengers for oil and gas sectors. *Rev Chem Eng* **37**: 663–686. <https://doi.org/10.1515/revce-2019-0049>.
- Schink, B. (1997) Energetics of syntrophic cooperation in methanogenic degradation. *Microbiol Mol Biol Rev* **61**: 262–280. <https://doi.org/10.1128/mmb.61.2.262-280.1997>.
- Shock, E.L., Oelkers, E.H., Johnson, J.W., Sverjensky, D.A., and Helgeson, H.C. (1992) Calculation of the thermodynamic properties of aqueous species at high pressures and temperatures. Effective electrostatic radii, dissociation constants and standard partial molal properties to 1000 °C and 5 kbar. *J Chem Soc Faraday Trans* **88**: 803–826.
- Slobodkin, A.I., Reysenbach, A.-L., Slobodkina, G.B., Baslerov, R.V., Kostrikin, N.A., Wagner, I.D., and Bonch-Osmolovskaya, E.A. (2012) *Thermosulfurimonas dismutans* gen. nov., sp. nov. a novel extremely thermophilic sulfurdisproportionating bacterium from a deep-sea hydrothermal vent. *Int J Syst Evol Microbiol* **62**: 2565–2571. <https://doi.org/10.1099/ijs.0.034397-0>.
- Slobodkin, A.I., Reysenbach, A.L., Slobodkina, G.B., Kolganova, T.V., Kostrikin, N.A., and Bonch-Osmolovskaya, E.A. (2013) *Dissulfuribacter thermophilus* gen. nov., sp. nov., a thermophilic, autotrophic, sulfur-disproportionating, deeply branching deltaproteobacterium from a deep-sea hydrothermal vent. *Int J Syst Evol Microbiol* **63**: 1967–1971. <https://doi.org/10.1099/ijs.0.046938-0>.
- Slobodkin, A.I., and Slobodkina, G.B. (2019) Diversity of sulfur-disproportionating microorganisms. *Microbiol* **88**: 509–522. <https://doi.org/10.1134/S0026261719050138>.
- Slobodkin, A.I., Slobodkina, G.B., Panteleeva, A.N., Chernykh, N.A., Novikov, A.A., and Bonch-Osmolovskaya, E.A. (2016) *Dissulfurimicrobium hydrothermale* gen. nov., sp. nov., a thermophilic, autotrophic, sulfur-disproportionating deltaproteobacterium isolated from a hydrothermal pond of Uzon caldera, Kamchatka. *Int J Syst Evol Microbiol* **66**: 1022–1026. <https://doi.org/10.1099/ijsem.0.000828>.
- Slobodkina, G.B., Kolganova, T.V., Kopitsyn, D.S., Viryasov, M.B., Bonch-Osmolovskaya, E.A., and Slobodkin, A.I. (2016) *Dissulfurirhabdus thermomarina* gen. nov., sp. nov. a thermophilic, autotrophic, sulfite-reducing and disproportionating deltaproteobacterium isolated from a shallow sea hydrothermal vent. *Int J Syst Evol Microbiol* **66**: 2515–2519. <https://doi.org/10.1099/ijsem.0.001083>.
- Slobodkina, G.B., Reysenbach, A.-L., Kolganova, T.V., Novikov, A.A., Bonch-Osmolovskaya, E.A., and Slobodkin, A.I. (2017) *Thermosulfuriphilus ammonigenes* gen. nov., sp. nov., a thermophilic, chemolithoautotrophic bacterium capable of respiratory ammonification of nitrate with elemental sulfur. *Int J Syst Evol Microbiol* **67**: 3474–3479. <https://doi.org/10.1099/ijsem.0.002142>.
- Takai, K., Nakamura, K., Toki, T., Tsunogai, U., Miyazaki, M., Miyazaki, J., et al. (2008) Cell proliferation at 122°C

- and isotopically heavy CH<sub>4</sub> production by a hyperthermophilic methanogen under high-pressure cultivation. *Proc Natl Acad Sci U S A* **105**: 10949–10954. <https://doi.org/10.1073/pnas.0712334105>.
- Tanger, J.C., and Helgeson, H.C. (1988) Calculation of the thermodynamic and transport properties of aqueous species at high pressures and temperatures; revised equations of state for the standard partial molal properties of ions and electrolytes. *Am J Sci* **288**: 19–98.
- Thamdrup, B., Finster, K., Hansen, J.W., and Bak, F. (1993) Bacterial disproportionation of elemental sulfur coupled to chemical reduction of iron or manganese. *Appl Environ Microbiol* **59**: 101–108. <https://doi.org/10.1128/aem.59.1.101-108.1993>.
- Thompson, G., Humphris, S.E., Schroeder, B., Sulanowska, M., and Rona, P.A. (1988) Active vents and massive sulfides at 26 degrees N (TAG) and 23 degrees N (Snakepit) on the mid-Atlantic ridge. *Canad Mineral* **26**: 697–711.
- Wolery, T.J. (1979) *Calculation of Chemical Equilibrium Between Aqueous Solution and Minerals: The EQ3/6 Software Package*. Livermore, CA: California Univ.
- Zeng, X., Alain, K., and Shao, Z. (2021) Microorganisms from deep-sea hydrothermal vents. *Mar Life Sci Technol* **3**: 204–230. <https://doi.org/10.1007/s42995-020-00086-4>.
- Zillig, W., Yeats, S., Holz, I., Böck, A., Rettenberger, M., Gropp, F., and Simon, G. (1986) *Desulfurolobus ambivalens*, gen. Nov., sp., nov., an autotrophic archaeobacterium facultatively oxidizing or reducing sulfur. *Syst Appl Microbiol* **8**: 197–203.
- Zopfi, J., Ferdelman, T.G., and Fossing, H. (2004) Distribution and fate of sulfur intermediates - sulfide, tetrathionate, thiosulfate, and elemental sulfur – in marine sediments. In *Sulfur Biogeochemistry - Past and Present: Geological Society of America Special Paper 379*, Amend, J.P., Edwards, K.J., and Lyons, T.W. (eds). Boulder, CO: Geological Society of America, pp. 97–116. <https://doi.org/10.1130/0-8137-2379-5.97>.