



HAL
open science

Larval habitat determines the bacterial and fungal microbiota of the mosquito vector *Aedes aegypti*

Karima Zouache, Edwige Martin, Nil Rahola, Marc Gangue, Guillaume Minard, Audrey Dubost, van Tran Van, Laura B Dickson, Diego Ayala, Louis A Lambrechts, et al.

► **To cite this version:**

Karima Zouache, Edwige Martin, Nil Rahola, Marc Gangue, Guillaume Minard, et al.. Larval habitat determines the bacterial and fungal microbiota of the mosquito vector *Aedes aegypti*. *FEMS Microbiology Ecology*, 2022, 98 (1), <10.1093/femsec/fiac016>. <hal-03690756>

HAL Id: hal-03690756

<https://hal.science/hal-03690756v1>

Submitted on 13 Jun 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

1 **Larval habitat determines the bacterial and fungal microbiota of the mosquito vector**

2 *Aedes aegypti*

3

4 Karima Zouache^{1*}, Edwige Martin^{1*}, Nil Rahola^{2,3}, Marc F. Ngangue^{2,4}, Guillaume Minard¹,

5 Audrey Dubost¹, Van Tran Van¹, Laura Dickson^{5,6}, Diego Ayala^{2,3}, Louis Lambrechts⁵ and

6 Claire Valiente Moro¹

7

8 ¹Univ Lyon, Université Claude Bernard Lyon 1, CNRS, INRAE, VetAgro Sup, UMR

9 Ecologie Microbienne, F-69622 Villeurbanne, France

10

11 ²CIRMF, Franceville, Gabon

12

13 ³UMR MIVEGEC, Univ. Montpellier, CNRS, IRD, Montpellier, France

14

15 ⁴ANPN, Libreville, Gabon

16

17 ⁵Institut Pasteur, Université de Paris, CNRS UMR2000, Insect-Virus Interactions Unit, Paris,

18 France

19

20 ⁶Department of Microbiology and Immunology, University of Texas Medical Branch,

21 Galveston, TX, USA

22

23

24 *These authors contributed equally to this study.

25

26 **Correspondence:** Dr. Claire Valiente Moro, Université Claude Bernard Lyon 1, Ecologie
27 Microbienne, Bât. André Lwoff, 10 rue Raphaël Dubois, 69100 Villeurbanne Cedex, France.
28 claire.valiente-moro@univ-lyon1.fr
29

30 **Abstract**

31 Mosquito larvae are naturally exposed to microbial communities present in a variety of larval
32 development sites. Several earlier studies have highlighted that the larval habitat influences
33 the composition of the larval bacterial microbiota. However, little information is available on
34 their fungal microbiota, *i.e.* the mycobiota. In this study, we provide the first simultaneous
35 characterization of the bacterial and fungal microbiota in field-collected *Aedes aegypti* larvae
36 and their respective aquatic habitats. We evaluated whether the microbial communities
37 associated with the breeding site may affect the composition of both the bacterial and fungal
38 communities in *Ae. aegypti* larvae. Our results show a higher similarity in microbial
39 community structure for both bacteria and fungi between larvae and the water in which larvae
40 develop than between larvae from different breeding sites. This supports the hypothesis that
41 larval habitat is a major factor driving microbial composition in mosquito larvae. Since the
42 microbiota plays an important role in mosquito biology, unravelling the network of
43 interactions that operate between bacteria and fungi is essential to better understand the
44 functioning of the mosquito holobiont.

45

46 Key words: mosquito, microbiota, larva, breeding site, bacteria, mycobiota

47

48 **Introduction**

49 Like most living organisms, the mosquito microbiota includes resident (*i.e.*
50 microorganisms that constantly live in or on the insect body) and transient (*i.e.*
51 microorganisms that are only temporarily found in the insect body) microorganisms.
52 Environmentally acquired microbes are thus part of the transient/flexible microbiota and are
53 also considered as members of the mosquito microbiota. The recent and extensive scientific
54 research on mosquito microbiota has led to the recognition that mosquito-microbe
55 relationships are important for the host biology and pathogen transmission (Minard,
56 Mavingui, Moro 2013; Guégan et al. 2018b; Dada et al. 2021). Regarding pathogen
57 transmission, many examples demonstrated the ability of bacteria to affect vector competence
58 for pathogens that infect humans (Dennison et al. 2014; van Tol and Dimopoulos, 2016).
59 Some bacteria can negatively affect pathogen development and/or replication by activating
60 immune response (Gabrieli et al. 2021) or prevent pathogen infection through competition for
61 resources or the secretion of molecules against pathogens (Jupatanakul et al. 2014). In
62 contrast, other bacteria as well as fungi can enhance mosquito host susceptibility to pathogen
63 infection (Apte-Deshpande et al. 2014; Angleró-Rodríguez et al. 2017; Wu et al. 2019).
64 Besides their influence in vector competence, bacteria were also shown to play important
65 roles in mosquito development, reproduction and nutrition (Dong, Manfredini, Dimopoulos
66 2009; Coon et al. 2014; Coon, Brown, Strand 2016a and 2016b; Dickson et al. 2017; Correa
67 et al. 2018; Chen et al. 2020; Guégan et al. 2020; Scolari et al. 2021; Wang et al. 2021).

68 Bacterial microbiota of mosquitoes is a dynamic community that shifts across the insect
69 biological cycle, as previously evidenced across several mosquito species (Gimonneau et al.,
70 2014; Coon et al. 2014; Duguma et al. 2015; Alfano et al. 2019). In most observations, the
71 microbiota diversity decreases during the development of mosquitoes from larvae to adults. A
72 selective sweep occurs during the pupal stage when a renewal of the gut epithelial layers

73 induces a massive loss of bacterial communities (Alfano et al. 2019). Interestingly, a thorough
74 study conducted in artificially reared *Ae. albopictus* mosquitoes under laboratory conditions
75 highlighted that multiple factors including development stage, adult age and female feeding
76 regimes determine the structure of bacterial microbiota from egg to adult stages (Chen et al.
77 2020). Overall, multiple intrinsic and extrinsic factors may affect the acquisition and
78 settlement of microbes during early stages. However, the breeding sites and habitats of
79 mosquitoes were previously suggested as the main drivers of adult-associated bacterial
80 communities (Guégan et al. 2018b; Dickson et al. 2017). As a matter of fact, an important part
81 of the microbiota is environmentally acquired in mosquitoes (Coon et al. 2014). It was shown
82 that larvae or newly emerged adults harbour microorganisms derived from their aquatic
83 habitats (Guégan et al. 2018b; Mallasigné, Valiente Moro, Luis 2020). This means that
84 environmental determinants strongly influence important traits in the biology of mosquitoes
85 (Ranasinghe and Amarasinghe 2020; Dickson et al. 2017).

86 The yellow fever mosquito *Aedes aegypti* is a main vector for several medically
87 important arboviruses, including dengue, Zika, yellow fever, and chikungunya viruses
88 (Powell and Tabachnick 2013). Infections of humans with dengue and yellow fever viruses
89 remain an important public health concern as they infect 390 million and 200,000 people each
90 year, respectively (Barrett and Higgs 2007; Bhatt et al. 2013). To date, only few publications
91 have explored the overlap between the microbiota of *Ae. aegypti* larvae and the microbial
92 communities of their aquatic habitat (Dada et al. 2014; Ranasinghe and Amarasinghe 2020;
93 Hery et al. 2021). Moreover, in comparison to bacteria, fungal communities (also referred to
94 as mycobiota) were largely understudied across mosquito species including *Ae. aegypti*
95 (Mallasigné, Valiente Moro, Luis 2020). A better understanding of the dynamics involved in
96 microbial acquisition in this mosquito species is therefore needed. In particular, it remains
97 unclear whether the microorganisms acquired from the water of breeding sites constitute a

98 major or negligible proportion of the larval microbiota and whether similar factors drive the
99 acquisition of bacterial and fungal communities. Additional field studies comparing the
100 microbial diversity including both the simultaneous analysis of bacteria and fungi of larvae
101 and their aquatic habitats would provide important knowledge on the role of the environment
102 in modulating the microbiota as a whole. In a previous study in central Gabon, we showed
103 that the bacterial microbiota colonizing the gut of adult mosquitoes is highly correlated with
104 variations in the microbiota of water they emerged from but markedly differed from water in
105 terms of composition (Dickson et al. 2017). Indeed, bacteria found in adult guts are
106 underrepresented in water. This result is consistent with the well-known microbiota
107 reorganization that occurs between immature and adult stages (Alfano et al. 2019) and
108 suggests that water may constitute a main source of microbes but that mosquitoes filter most
109 of them out during their development. Considering this observation, we suggest that immature
110 mosquito larvae that did not yet proceed to microbiota filtering during pupation may even be
111 more influenced by the microbial composition of the water they are living in.

112 The objective of the current study was to investigate to which extent variations of the
113 microbial communities in aquatic habitats influence *Ae. aegypti* larval microbiota
114 composition and the paired structures of bacterial and fungal communities. For this, we
115 collected larvae and water from various artificial containers within the Lopé village in Gabon.
116 We hypothesized that similar environmental factors shape the composition of these two
117 communities in the immature stages of mosquitoes. Such knowledge provides new insights
118 into the environmental drivers of mosquito-associated bacterial and fungal communities.

119

120 **Material and Methods**

121

122 **Study site and sample collection**

123 Field sampling was performed in Gabon during the wet season in November - December 2017
124 and April - May 2018 in Lopé village (latitude, -0.099221; longitude, 11.600330).
125 Third- and fourth-instar larvae as well as water were sampled from artificial breeding sites
126 such as small tanks and containers (Table S1). Samples were collected with a sterile plastic
127 pipette into a sterile 50-ml falcon tube with filter-top lid. Water samples were agitated before
128 sampling to mix detritus and the upper surface to a depth of 10cm was collected representing
129 ~ 50 ml per breeding site. In the Station d'Etude des Gorilles et Chimpanzés field station,
130 larvae were separated from the water for each breeding site under a laminar flow cabinet,
131 transferred to a new sterile 50-ml falcon tube and both samples, water and larvae were then
132 snap frozen in a liquid nitrogen tank and subsequently conserved at -80°C. All larvae
133 collected from each sampled breeding site were selected for further sequencing analysis
134 (Table S1).

135

136 **DNA extraction**

137 Water samples were homogenized after vortexing for 1 min and an aliquot of 1.5 mL was
138 centrifuged for 10 min at maximum speed (13 400 rpm). The supernatant was discarded and a
139 new aliquot of 1.5 mL was added. This step was repeated from 2 to 4 times depending on the
140 volume of water available for each sample. Pooled pellets were resuspended in 500 µL of
141 extraction buffer (10 mM tris-HCl [pH 8.0], 10 mM EDTA, disodium salt [pH 8.0], 100 mM
142 sodium chloride, and 2% (v/v) SDS) and DNA was extracted as previously described
143 (Dickson et al. 2017). The extractions were performed in 6 batches and a negative control
144 (*i.e.*, extraction performed on an empty tube) was made for each batch. Prior to DNA
145 extraction, larvae were surface-sterilized by three washes in sterile water, followed by 5 min
146 soaking in 70% ethanol, and five rinses in sterile water. Each larva was crushed for 10 s using
147 a Mini-beadbeaterTP apparatus (Biospec Products) into a 2-mL Eppendorf tube containing 1-

148 mm diameter beads (Biospec, 1 mm) and 250 μ L of CTAB buffer (2% hexadecyltrimethyl
149 ammonium bromide, 1.4 M NaCl, 0.02 M EDTA, 0.1 M Tris pH8, 0.2% 2- β
150 mercaptoethanol). After incubation for 1h at 60°C on a shaker (300 rpm), 4 μ L of RNase (100
151 mg/ml) was added and incubated at 37°C for 5 min. Mixtures of phenol:chloroform:isoamyl
152 alcohol (25 :24 :1) and chloroform:isoamyl alcohol (24 :1) were added and after
153 centrifugation for 30 min at 13 200 rpm, DNA was precipitated using isopropyl alcohol. DNA
154 pellets were rinsed twice with 75% cold ethanol, air-dried under laminar flow and
155 resuspended in 22 to 27 μ L of sterile RNase free water. DNA purity and concentration were
156 measured using a NanoDrop™ spectrophotometer for both water and larvae samples.

157

158 **Mitochondrial gene amplification**

159 DNA samples from larvae were used as templates to amplify a 597-base-pair (bp) region of
160 the mitochondrial cytochrome c oxidase subunit I (COI) gene as previously described
161 (Raharimalala et al. 2012). PCR products were sent to sequencing at BIOFIDAL-DTAMB
162 (FR BioEnvironment and Health, Lyon, France). Sequences of each COI haplotype were
163 deposited on European Nucleotide Archive (<https://www.ebi.ac.uk/ena/browser/home>) under
164 the accession number PRJEB45388.

165

166 **Library preparation and sequencing**

167 For amplicon sequencing, two-step Nextera PCR libraries were created following the Illumina
168 protocol. First-step PCR were performed in triplicate on the hypervariable V5-V6 regions of
169 the 16S rRNA gene (*rrs*) of ~ 280 bp and the nuclear ribosomal internal transcribed spacer
170 (ITS2 region) for bacterial and fungal community identification, respectively. The modified
171 primers 784F (5'-AGGATTAGATACCCTGGTA-3') and 1061R (5'-
172 CRRCACGAGCTGACGAC-3') for bacteria and gITS7 (5'- GTG AAT CAT CGA RTC TTT

173 G -3') and ITS4 primers (5'- TCC TCC GCT TAT TGA TAT GC-3') for fungi were used.
174 Each primer pair was associated with the two Illumina overhanging adapter sequences. Primer
175 pair targeting the V5-V6 region was selected as these primers did not match mosquito 18S
176 and mitochondrial 16S rDNA gene sequences and allowed the better coverage of the
177 sequences belonging to the RDP Bacteria domain database (Minard et al. 2014). All PCR
178 amplifications were carried out in triplicates on a Biorad C1000 thermal cycler (Biorad, CA,
179 USA) in a 25 µl reaction. PCR reactions were performed using the 5X BioAmp® master mix
180 (Biofidal, France) containing 2 µL sample DNA, 5 µL of Mix HotStarTaq 5X, 1X of GC rich
181 Enhancer, 0.5 mg.mL⁻¹ of bovine serum albumin (New England Biolabs, Evry, France) and 1
182 µM of each primer. For bacteria, amplifications were conducted for 10 min at 96°C followed
183 by 5 cycles at 96°C for 20 s, 65°C for 30 s, 72°C for 60 s, 5 cycles at 96°C for 20 s, 61°C for
184 30 s, 72°C for 60 s, 5 cycles at 96°C for 20 s, 57°C for 30 s, 72°C for 60 s, 25 cycles at 96°C
185 for 20 s, 54°C for 30 s, 72°C for 60 s and a final extension at 72°C for 10 min. For fungi,
186 amplifications were conducted with one cycle of 3 min at 94°C, followed by 40 cycles at
187 94°C for 45 s, 55°C for 45 s, 72°C for 30 s, and a final extension step of 5 min at 72°C. The
188 three PCR product replicates from each sample (water or individual larvae) of each gene were
189 pooled, purified and quantified with the Quant-iT Picogreen dsDNA Assay Kit (Life
190 Technologies). A total of 148 amplicon libraries were constructed. A total of 21 (11 for
191 bacteria, 10 for fungi), 115 (57 for bacteria and 58 for fungi) and 12 (6 for bacteria and 6 for
192 fungi) amplicon libraries were generated for water samples, larvae and negative controls
193 respectively. Subsequently, one Illumina MiSeq platform and a v3 600 cycles kit was used to
194 sequence the PCR libraries (second-step PCR) at Biofidal (Vaulx-en-Velin, France). All
195 FastQ files were deposited in the EMBL European Nucleotide Archive
196 (<https://www.ebi.ac.uk/ena>) under the project accession number PRJEB45062.

197

198 **Bioinformatics analysis**

199 Total of 5,311,262 and 4,983,099 reads were obtained for the bacteria and fungi respectively
200 and demultiplexed. Sequence quality control and analysis of sequence data were carried out
201 using the FROGS pipeline (Escudie et al. 2016) with the previously described parameters
202 (Guégan et al. 2018a; Guégan et al., 2020). Taxonomic affiliation was performed against the
203 SILVA database 132 for bacteria (Yilmaz et al. 2014) and ITS UNITE database for fungi
204 (Kõljalg et al. 2013) with the Mothur pipeline (Schloss et al. 2009) at 80% minimum
205 bootstrap using a naïve Bayesian classifier (Wang et al. 2007). Contaminants were filtered out
206 using negative control (blank extraction and PCR). OTUs were removed if they were detected
207 in the negative control sample and their relative abundance was not at least 10 times greater
208 than that observed in the negative control (Minard et al. 2015). Sequences were grouped into
209 operational taxonomic units (OTUs) by clustering at 97% similarity. To compare samples,
210 normalization was performed at 21,629 and 18500 sequences for the bacterial and fungal
211 sequences, respectively. A total of 877 and 768 operational taxonomic units (OTUs) were
212 obtained for bacteria and fungi, respectively. Venn diagrams were generated with the Venn
213 Diagrams software from the Van de Peer Lab Bioinformatics and Evolutionary Genomics
214 (<http://bioinformatics.psb.ugent.be/webtools/Venn/>).

215

216 **Statistical analysis**

217 Data analysis and statistical tests (Mann-Whitney-Wilcoxon and Adonis-ANOVA) were
218 carried out using R software (R Core Team 2018) with phyloseq (McMurdie and Holmes
219 2013) and vegan (Oksanen 2018) for diversity analysis as well as ggplot2 (Wickham 2018)
220 for graphical representations. Bacterial and fungal within individual diversity also referred as
221 α -diversity was estimated with the Shannon index (H'). Community similarity between
222 individuals also referred as β -diversity was estimated with the Bray-Curtis index.

223 Comparisons of α -diversity between larvae and water were performed with an ANOVA.
224 Correlations between larvae and water microbial communities (either bacteria and fungi)
225 collected in the same habitats were performed with a Spearman rank-based analysis.
226 Comparisons of microbial communities' β -diversity were performed with the *adonis*-ANOVA
227 permutational analysis of variance with the package *vegan*. Spearman correlations between
228 communities associated with water and larvae were computed with the *lineup2* package
229 (Broman 2021) and heatmap graphs were represented with *complexHeatmap* (Gu 2021).

230

231 **Ethical statement**

232 Mosquito collections inside Lopé National Park were conducted under permit number
233 AE17012/PR/ANPN/SE/CS/AFKP and the national research authorisation number
234 AR0013/117/MESRSFC/CENAREST/CG/ CST/CSAR.

235

236 **RESULTS**

237 **Diversity and correlations between microbial communities of larvae and larval habitat**

238 **OTU diversity.** The microbiota of water and *Ae. aegypti* larvae samples were examined by
239 sequencing of the bacterial 16S rRNA gene and ITS2 region, for bacterial and fungal
240 communities respectively. Sequences of a total of 148 libraries (*i.e.*, 21 water samples; 115
241 individual larval samples, 6 DNA extraction negative controls for each gene) resulted in
242 5,311,262 sequence reads for bacteria, ranging from 57,562 to 116,144 (with an average of
243 73,763 and 78,945 for water and larvae, respectively) and 4,983,099 sequence reads for fungi,
244 ranging from 47,493 to 101,501 (with an average of 61,967 and 75,232 for water and larvae,
245 respectively). Larvae and water exhibited a higher number of OTUs for bacteria than for fungi
246 and the number of OTUs was higher in water than in larvae irrespective of the microbial
247 community (Table S2 and S3). The mean number of OTUs detected in larvae sharing the

248 same breeding site varied from 146 ± 35 to 332 ± 67 for bacteria and from 58 ± 13 to $268 \pm$
249 62 for fungi (Table S2 and S3). The average numbers of bacterial and fungal OTUs in water
250 were 202 ± 40 (range 138-266) and 84 ± 47 (range 38-202), respectively (Table S2 and S3).

251 **Microbial communities comparison between water samples and larvae.** Larvae belonging
252 to the same breeding site showed a relatively conserved microbiota for both bacteria (Fig 1A)
253 and fungal communities (Fig 1B). Permutational analysis of variance showed that bacterial
254 and fungal microbiota of larvae differed among individuals belonging to different breeding
255 sites (*adonis*-ANOVA, $F = 5.137$, $R^2 = 0.678$, $p = 0.001$; $F = 7.249$, $R^2 = 0.73$, $p = 0.001$ for
256 bacteria and fungi respectively) suggesting a strong impact of the larval breeding site on
257 larval microbial communities. Water samples and larvae significantly differ from each other
258 in terms of bacterial (*adonis*-ANOVA, $F = 4.871$, $R^2 = 0.033$, $p = 0.001$) and fungal
259 communities (*adonis*-ANOVA, $F = 1.386$, $R^2 = 0.02$, $p = 0.001$) (Fig 2A and Fig 2B).

260 Comparison analysis performed with the Shannon index indicated no significant difference in
261 the α -diversity of bacterial communities between larvae and water (ANOVA, $F = 0.305$, $p =$
262 0.582). However, significant differences were observed between larvae and water in the
263 mycobiota diversity, the fungal microbiota being more diverse in larvae than in water
264 (ANOVA, $F = 4.345$, $p = 0.04$). Furthermore a positive correlation was detected between the
265 Shannon index of the water and the Shannon index of larvae for fungi (Spearman correlation
266 $\rho = 0.74$, $p = 3.379 \times 10^{-7}$; Fig. 3B) but not for bacteria (Spearman correlation $\rho = -0.22$, $p =$
267 0.18 ; Fig. 3A). A total of 542 bacterial OTUs and 375 fungal OTUs (*i.e.* 61.8 % and 48.8 %
268 of the total OTUs) were shared between larvae and water (Fig. 4A and Fig. 4B). Larvae-
269 specific bacterial and fungal OTUs counted for 251 and 346, respectively while that of water-
270 specific OTUs was lower with 27 and 21 respectively for bacteria (Fig. 4A) and fungi (Fig.
271 4B).

272 **Correlation between the microbial composition of larvae and water.** A clear correlation of
273 the microbiota of larvae and water was detected for individuals belonging to the same
274 breeding site for both bacteria (Fig. S1) and fungi (Fig. S2). The same analysis performed at
275 the scale of OTUs showed that OTUs split into different groups highlighting some patterns of
276 positive correlations between larval and water microbiota composition (Fig. 5 and 6).
277 Regarding bacteria, one group consisted of 5 OTUs in larval samples (*Clostridium sensu*
278 *stricto*, *Camelimonas*, *Candidatus_Proteochlamydia*, *Bosea* and *Vogesselia*) that showed
279 correlated abundances with 4 OTUs in water samples (*Chryseobacterium*, *Legionella*,
280 *Camelimonas* and *Bosea*) (Fig. 5). The identification of *Camelimonas* and *Bosea* OTUs in the
281 two groups of both categories of samples, suggests that water of breeding sites was a major
282 determinant of bacterial microbiota composition in larvae of these two taxa. Another group
283 harbouring the OTUs *Chromobacterium*, *Arcicella* and one unclassified taxa in larvae
284 samples was positively correlated with the OTUs *Novosphingobium*, uncultured *Chlamydiales*
285 (*CvE6_ge*) and uncultured *Beijerinckiaceae* (*FukuN57*) in water samples. Finally, a third
286 group of OTUs composed of *Microbacterium*, *Phreatobacter* and 2 unclassified taxa were
287 positively correlated with the OTUs *Leifsonia*, *Rudanella* and 5 other unclassified taxa.
288 Regarding fungi, 5 groups of positive correlations were highlighted (Fig. 6). All together
289 these results indicate that water of breeding sites is a strong driver of microbial assemblages
290 in larvae, for both bacterial and fungal communities.

291

292 **Composition of microbial communities in *Aedes aegypti* larvae and their associated** 293 **aquatic habitats**

294 **Bacterial taxonomic composition.** A total of 877 OTUs belonging to 11 bacterial phyla and
295 254 genera were identified across all the samples. The percentage of unclassified bacterial
296 sequences represented less than 1% irrespective of the sample type except for one larval

297 sample, which exhibited 21% of unclassified sequences (Fig. 7). In larvae, 11 bacterial phyla
298 were identified and 3 of them (Actinobacteria, Proteobacteria and Firmicutes) were prevalent
299 and abundant (taken together range 47.07% - 99.97% of relative abundance, average 89.69%).
300 Their relative abundance was variable among larval samples between breeding sites ranging
301 from 3.34% to 88.11% for Actinobacteria, 10.25% to 56.54% for Proteobacteria and 1.19% to
302 85.26% for Firmicutes. For water samples, only Proteobacteria and Actinobacteria were
303 retrieved in all samples. Taken together, the mean abundance of the three phyla
304 (Actinobacteria, Proteobacteria and Firmicutes) per larval habitat is more than 67.68 %
305 (average 88.59%). The relative abundance of Proteobacteria ranged from 34.01% to 95.44%
306 (average 76.73%) and was higher compared to that of Actinobacteria (range 0.82% - 35.49%,
307 average 10.99%). When comparing water and larval samples, the proportion of Proteobacteria
308 was more important in water compared to larvae (76.73 ± 19.93 vs 32.32 ± 16.79) while
309 higher proportions of Actinobacteria were reported in larvae compared to water ($34.55 \pm$
310 25.78 vs 10.99 ± 13.52) (Fig. 7). A total of 62 bacterial genera out of 254 (24.4%) were
311 specific to larvae (among which the most abundant were *Ralstonia* and
312 *Candidatus_Proteochlamydia*) and 8 genera (3.1%) were specific for water (among which the
313 most abundant were unclassified Sporichthyaceae, *Pseudarcicella* and *Rudanella*). A
314 combined analysis of prevalence and abundance was also performed. Prevalence refers to the
315 overall infection rate of a given OTU overall samples (number of samples, i.e. larvae or
316 water, in which the OTU was detected / total number of samples). Abundance refers to
317 proportion occupied by an OTU within a sample among either the bacterial or fungal
318 community (number of normalized reads assigned to the OTU in a given samples / total
319 number of normalized reads). This analysis revealed the 5 OTUs *Mycobacterium*, *Clostridium*
320 *sensu stricto*, unclassified *Peptostreptococcaceae*, unclassified *Burkholderiaceae* and
321 unclassified *Microbacteriaceae* were the most prevalent (>75%) and abundant (>10²) across

322 larvae (Fig. 8A). In water, 2 OTUs identified as unclassified *Rhizobiales* were identified as
323 the most prevalent (100%) with high abundance ($>10^2$) (Fig. 8B).

324 **Fungal taxonomic composition.** A total of 668 OTUs belonging to 9 sub-phyla and 49 order
325 were identified in *Ae. aegypti* larvae. Compared to bacterial communities, the percentage of
326 unclassified fungal sequences was important and variable across samples. It was comprised
327 between 6.65% and 99.79% in larvae or between 12.58% and 99.94% in water (Fig. 9).

328 Larvae and water samples exhibited similar communities dominated by the sub-phylum
329 Pezizomycotina (from 9% to 91.59% in larvae and from 4.94% to 83.67% in water) and
330 followed by the sub-phylum Agaromycotina (from 0,14% to 15,34% in larvae and from
331 0,37% to 20,29% in water). The Zygomycota incertae sedis phylum was absent from water
332 but present in 3 out of 17 samples of larvae while conversely the phylum Chytridiomycota
333 was absent from larvae but identified in one sample of water. At the order level, similar
334 fungal compositions were observed between larvae and their associated aquatic habitats
335 within the phyla Ascomycota and Basidiomycota (Fig. S3 and S4). Within the phylum
336 Ascomycota, the majority of reads of water samples was assigned to the order Pleosporales
337 (33.73%) followed by Capnodiales (27.02%) and Hypocreales (10.04%) (Fig. S3). For larvae
338 samples (at least one larva of the breeding site), it was Pleosporales (27.64%),
339 Trichosphaeriales (13.46%), Eurotiales (13.39%) and Capnodiales (10.46%) (Fig. S3). Reads
340 assigned to Pleosporales were detected in all of the water and mosquito samples (Fig. S3).

341 Within the Basidiomycota, the majority of reads were associated with the order Tremellales
342 (46.87%) followed by Rhyzophydiales (20.19%) (Fig. S4). For larvae samples, it was
343 Tremellales (32%), Polyporales (12.78%) and Wallemiales (11.92%) (Fig. S4). A combined
344 analysis of prevalence and abundance revealed two OTUs (one unclassified *Trichosphaeriales*
345 and another OTU identified as unclassified) as being the most prevalent and abundante ($>60\%$
346 and 10^2 , respectively) across larvae followed by the OTU *Penicillium citrinum* ($>50\%$ and

347 10²) (Fig. 8C). In water, one OTU identified as unclassified Pleosporales was found to be the
348 most prevalent (>80%) and abundant (5.10²) followed by the two OTUs *Toxidocladosporium*
349 *irritans* and *Aureobasidium* (70%, 10¹) (Fig. 8D).

350

351 **Discussion**

352 In a previous study, we demonstrated that larval exposure to different bacteria drive
353 variation in *Ae. aegypti* vector competence (Dickson et al. 2017). Such a result underlines the
354 importance to better characterize the microbial composition of mosquito larvae in relation
355 with their breeding sites since larvae acquire the major part of their microbiota from water
356 habitats (Xia et al. 2021). Despite growing information on the composition and diversity of
357 bacterial communities associated with *Ae. aegypti* at the adult stage, in comparison limited
358 data are available on larvae and their water habitats as well as on other associated microbes.
359 To fill this gap, we used high-throughput sequencing to compare bacterial and fungal
360 communities between water samples and *Ae. aegypti* larvae collected from various breeding
361 sites in the Lopé village (Gabon).

362 Overall, our study showed that there was higher similarity in microbial structure, for
363 both bacterial and fungal communities, between larvae and water in which larvae develop
364 than between larvae and water from different breeding sites. This correlation in the microbial
365 composition between larvae and their surrounding environment could reflect either an
366 influence of larval microbiota on water microbial communities (through the release of
367 microbial community from larvae feces or during molting as well as larval predation of
368 microorganisms) or on the opposite, an influence of the water microbial communities on
369 larval microbiota (through acquisition by filtration, nutrition, limited microbial vertical
370 transmission). Interestingly, our results also support the fact that larval microbiota is highly
371 similar to the microbial communities found in larval habitats contrary to what was observed

372 for adult mosquitoes. Since it is known that a massive loss of bacterial communities occurs
373 during the pupal stage, this suggests that the mosquitoes select the microbiota in late stages
374 (most likely between pupation and emergence) or that some bacteria are quickly able to re-
375 colonize newly-emerged adult mosquitoes. The first hypothesis would explain why adults
376 have a microbiota that varies symmetrically to the larval breeding site they originated from
377 but have a different microbiota composed of sub-dominant species in water.

378 While no significant differences were detected for bacteria, a more diverse microbiota
379 in larvae than in water was observed for fungal communities, associated with a positive
380 correlation between the Shannon indices of the water and the larvae. This higher diversity
381 could be explained by biotic and abiotic differences between both environments (*i.e.*
382 difference in pH, oxygen, immunity ...) and / or differences in fungi acquisition mode (*i.e.*
383 vertical transmission in insects, nutrition...). The hypoxia in the gut of larvae is a factor
384 limiting the growth of particular aerobic bacteria but little is known for fungi (Coon et al.
385 2017). Muturi et al (2020) also demonstrated that mosquito larvae are highly selective of the
386 bacterial taxa from the larval environment that colonize their bodies. Interestingly, a recent
387 study also demonstrated that amount of larval diet in the breeding water that proportionally
388 increase with bacterial loads in breeding water, has a significant effect on *Ae. aegypti*
389 mosquito microbiota size and composition and that these effects last into adulthood
390 (MacLeod, Dimopoulos, Short 2021). Microbial interactions such as competition,
391 amensalism, mutualism, cross-feeding or predation occur among microbial species in the
392 mosquito microbiota and may also impact in structuring microbial communities (Gusmao et
393 al. 2010). Thus, it is possible that bacteria or fungi already present in larvae limit the
394 colonization by new microorganisms referred to as priority effects. This was observed in a
395 recent study where the development of fungi was correlated with a reduction of bacterial
396 numbers by antibiotics in field-caught mosquitoes (Hyde et al. 2019).

397 Moreover, previous studies also reported that water with or without mosquito larvae as well
398 as larval density influence the bacterial community composition in water that in turn may
399 have consequences on larvae microbiota composition (Nilsson et al. 2018; Muturi et al. 2020).
400 However, such an impact remains to be investigated for fungal communities.

401 In agreement with other studies, Proteobacteria, Firmicutes and Actinobacteria were
402 the most abundant phyla found in field-collected *Ae. aegypti* larvae representing on average
403 89% of the total number of reads (Dada et al. 2014; Hery et al. 2021). These phyla were also
404 very abundant and common in water habitats but contrary to what was previously observed
405 (Hery et al. 2021), not all bacterial taxa identified in *Ae. aegypti* larvae were found in the
406 corresponding breeding sites. This difference could potentially be due to the difficulty of
407 sampling all microorganisms from a water source given that small volumes were sequenced
408 or differences in the total biomass and the DNA extraction protocols between the two
409 samples. Distinct patterns were observed for Proteobacteria and Actinobacteria. While the
410 proportion of Proteobacteria was more important in water compared to larvae, the contrary
411 was observed for Actinobacteria in which a higher proportion in larvae was obtained.
412 However, at the genus level the bacterial composition diverged between larvae and water
413 samples, with heterogeneity in the taxonomic profiles. The genus *Mycobacterium* was found
414 to be the most prevalent and abundant in larvae while it was an OTU belonging to an
415 unclassified *Rhizobiales* in water samples. Interestingly, the genus *Mycobacterium* was
416 recently shown to dominate in *Ae. japonicus* larvae and this genus with an unclassified
417 *Rhizobiales* occurred in disproportionally higher abundance in *Ae. japonicus* samples from
418 used tires (Juma et al. 2021). Moreover, the distribution of the other most abundant and
419 prevalent genera varied considerably among water and larvae except for an OTU belonging to
420 an uncultured *Burkholderiaceae* which was highly abundant and prevalent in both samples.

421 Interestingly, *Wolbachia* was identified in 4 individual larvae belonging to different
422 breeding sites with low number of sequences per larvae. Analysis of the V5-V6 sequences
423 showed 99% identity GenBank accession number MN046789) with *Wolbachia* previously
424 identified in field-collected *Ae. aegypti* mosquitoes in Philippines (Carvajal et al. 2019). This
425 result confirms recent studies highlighting the presence of this bacterium in *Ae. aegypti*
426 (Thongsripong et al. 2018; Balaji, Jayachandran and Prabakaran 2019; Carvajal et al. 2019;
427 Kulkarni et al. 2019, Inacio da Silva et al. 2021) while the absence of natural infection of
428 *Wolbachia* in *Ae. aegypti* mosquitoes had long been put forward (Gloria-Soria, Chiodo,
429 Powell 2018). We can not exclude the possibility that the detection of the bacterium could be
430 explained either by the infection of larvae by nematodes or parasitoids that carry *Wolbachia*
431 themselves or the ingestion by larvae of arthropods infected with *Wolbachia*. However,
432 further studies are needed to confirm the presence of naturally occurring *Wolbachia* in *A.*
433 *aegypti* as the scientific consensus is that sequencing data alone is not enough to establish this
434 (Chrostek et al. 2019; Ross et al. 2020). Screening on a larger number of individuals by using
435 additional markers such as the *Wolbachia* outer surface protein gene *wsp* and the GroE operon
436 would be performed in combination with a phylogeny of *Wolbachia* strains originating from
437 other arthropod hosts and mosquitoes.

438 In fungal taxonomy and community ecology, the most studied DNA barcode is the
439 nuclear ribosomal RNA intergenic spacer locus (Schoch et al. 2012). Due to a lack of
440 references in public databases, many of those sequences remained unassigned. Despite this
441 issue, we found that the sub-phyla Pezizomycotina and Agaromycotina were the most
442 abundant phyla found in *Ae. aegypti* larvae and their associated aquatic habitats. Within the
443 phylum Ascomycota, OTUs in the fungal orders Pleosporales were shared across all water
444 and larvae samples as previously observed in *Ae. albopictus* (Tawidian et al. 2021). In
445 addition, some specific patterns were observed between larvae and water as the majority of

446 reads was assigned to the orders Capnodiales and Hypocreales for water samples and
447 Trichosphaeriales, Eurotiales and Capnodiales for larvae samples. Within the Basidiomycota,
448 the majority of reads were associated with the order Tremellales that contrasts with previous
449 observations that showed a dominance and prevalence of the order Agaricales in all of the
450 water and *Ae. albopictus* mosquito samples (Tawidian et al. 2021). Interestingly, we showed
451 that *P. citrinum* was one of the most abundant and prevalent species across larvae. Russel et al
452 (2001) reported *P. citrinum* parasitizing eggs of *Ae. aegypti* in Australia by the production of
453 mycotoxins that inhibit the complete development of the eggs during the dry season, thereby
454 diminishing the incidence of mosquitoes during the rainy season. The fact that this species
455 was highly prevalent and abundant in larvae raises the question of its impact on this mosquito
456 species. Two OTUs belonging to the class Microbotryomycetes were found to be conserved
457 fungal OTUs between adult female mosquito species *Coquillettidia perturbans* and
458 *Ochlerotatus canadensis* collected at disparate sites (Hyde et al. 2019). OTUs belonging to
459 this class were also detected in *Ae. aegypti* larvae and aquatic habitats but they were not
460 among the most abundant and prevalent suggesting specific associations according to the
461 mosquito species or the stage.

462 The strong homogeneity between the breeding sites samples did not allow to group
463 them by type and to assess the role of differences in aquatic habitats on larval
464 microbiota/mycobiota composition and structure. To date, there have been no studies on the
465 influence of the type of habitat on both these communities among water and larvae. While
466 different habitats may result in location-driven variability in microbial composition, further
467 studies with replicates of habitats corresponding to each category would help to evaluate
468 whether the type of habitat determine the presence of specific microbial communities in water
469 and larvae. It means that the variation of some bacterial or fungal communities are linked
470 between the water and the larvae. More globally, the nature of breeding sites as well as

471 physicochemical factors are important factors that need to take into account to explain the
472 dynamics of microbial communities. Despite investigating the impact of spatial and temporal
473 variability of abiotic properties of *Ae. aegypti* breeding sites on immature abundance and
474 adult body size (David et al. 2021), no studies have yet disentangled their direct impact on the
475 microbial communities and their indirect consequences on larvae development and
476 physiological characteristics of the adult mosquito.

477 While many previous studies have examined the bacterial composition between larvae
478 and the water in which they are developing, this study is the first to examine the bacterial and
479 fungal composition of *Ae. aegypti* larval habitats in relation to that of their associated larvae.
480 To our knowledge, this is the first characterization of the mycobiota from field-collected
481 individual *Ae. aegypti* larvae and their associated aquatic habitats. We showed that larvae and
482 water share microbial communities whose diversity and richness are driven by the breeding
483 site. Further studies are needed to evaluate whether some fungi or bacteria detected in larvae
484 are transstadially passed to the adult and how the natural variability of breeding sites shapes
485 adult trait variation and vector competence.

486

487 **Supplementary data**

488 Supplementary data are available at FEMSEC online.

489

490 **Acknowledgements**

491 We thank all members of the DMTV group of the Microbial Ecology laboratory (UMR5557)
492 for their insights. We thank the IRD, CIRMF and the ANPN-SEGC in Gabon for the
493 assistance and expertise during the fieldwork. We are grateful to Christophe Paupy, Davy
494 Jiolle, Emilie Giraud and Amine Ghozlane for their input during development of the study.
495 We also thank the IBIO platform (UMR5557) for providing the computational resources. We

496 also thank Agnès Nguyen (NGS sequencing department of Biofidal) for helpful discussions.

497

498 **Funding.** This work was primarily funded by the Agence Nationale de la Recherche (grant
499 ANR-16-CE35-0004-01 to LL and CVM). It was also supported by the French Government's
500 Investissement d'Avenir program Laboratoire d'Excellence Integrative Biology of Emerging
501 Infectious Diseases (grant ANR-10-LABX-62- IBEID to LL).

502

503 ***Conflicts of interest.*** None declared.

504

505

506 **References**

507

508 Alfano N, Tagliapietra V, Rosso F et al. Changes in microbiota across developmental stages
509 of *Aedes koreicus*, an invasive mosquito vector in Europe: Indications for microbiota-Based
510 Control Strategies. *Front Microbiol* 2019; 10:2832.

511

512 Angleró-Rodríguez YI, Talyuli OA, Blumberg BJ et al. An *Aedes aegypti*-associated fungus
513 increases susceptibility to dengue virus by modulating gut trypsin activity. *Elife* 2017;
514 6:e28844.

515

516 Apte-Deshpande AD, Paingankar MS, Gokhale MD, Deobagkar DN. *Serratia odorifera*
517 mediated enhancement in susceptibility of *Aedes aegypti* for chikungunya virus. *IJMR* 2014;
518 139(5):762-768.

519

520 Balaji S, Jayachandran S, Prabakaran SR. Evidence for the natural occurrence of *Wolbachia*
521 in *Aedes aegypti* mosquitoes. *FEMS Microbiol Lett* 2019; 366(6):fnz055

522

523 Barrett AD and Higgs S. Yellow fever: a disease that has yet to be conquered. *Annu Rev*
524 *Entomol* 2007; 52:209-29.

525

526 Bhatt S, Gething PW, Brady OJ et al. The global distribution and burden of dengue. *Nature*
527 2013; 496: 504-7.

528

529 Broman KW. 2021. Lineup2: Lining up two sets of measurements.

530

531 Carvajal TM, Hashimoto K, Harnandika RK et al. Detection of *Wolbachia* in field-collected
532 *Aedes aegypti* mosquitoes in metropolitan Manila, Philippines. *Parasit Vectors* 2019;
533 *12(1)*:361.
534
535 Chen S, Zhang D, Augustinos A et al. Multiple factors determine the structure of bacterial
536 communities associated with *Aedes albopictus* under artificial rearing conditions. *Front.*
537 *Microbio* 2020; *11*:605.
538
539 Coon KL, Vogel KJ, Brown MR et al. Mosquitoes rely on their gut microbiota for
540 development. *Mol. Ecol* 2014; *23*:2727–2739.
541
542 Coon KL, Brown MR, Strand MR. Gut bacteria differentially affect egg production in the
543 anautogenous mosquito *Aedes aegypti* and facultatively autogenous mosquito *Aedes*
544 *atropalpus* (Diptera: Culicidae). *Parasit. Vectors* 2016a; *9*:1–12.
545
546 Coon KL, Brown MR, Strand MR. Mosquitoes host communities of bacteria that are essential
547 for development but vary greatly between local habitats. *Mol Ecol* 2016b; *25*:5806–26.
548
549 Coon KL, Valzania L, McKinney DA et al. Bacteria-mediated development of mosquitoes.
550 *PNAS* 2017 ; *114 (27)*:E5362-E5369.
551
552 Correa MA, Matusovsky B, Brackney DE et al. Generation of axenic *Aedes aegypti*
553 demonstrate live bacteria are not required for mosquito development. *Nat Commun* 2018;
554 *9*:4464.
555

556 Dada N, Jumas-Bilak E, Manguin S et al. Comparative assessment of the bacterial
557 communities associated with *Aedes aegypti* larvae and water from domestic water storage
558 containers. *Parasit Vectors* 2014; 7:391.
559

560 Dada, N, Jupatanakul, N, Minard, G et al. Considerations for mosquito microbiome research
561 from the Mosquito Microbiome Consortium. *Microbiome* 2021; 9:36.
562

563 David MR, Dantas ES, Maciel-de-Freitas R et al. Influence of larval habitat environmental
564 characteristics on Culicidae immature abundance and body size of adult *Aedes aegypti*. *Front*
565 *Ecol Evol* 2021; 9:626757.
566

567 Dennison NJ, Jupatanakul N, Dimopoulos G. The mosquito microbiota influences vector
568 competence for human pathogens. *Curr Opin Insect Sci* 2014 ; 3: 6–13.
569

570 Dickson LB, Jiolle D, Minard G et al. Carryover effects of larval exposure to different
571 environmental bacteria drive adult trait variation in a mosquito vector. *Sci Adv.* 2017;
572 3(8):e1700585.
573

574 Dong Y, Manfredini F, Dimopoulos G. Implication of the mosquito midgut microbiota in the
575 defense against malaria parasites. *PLoS Pathog* 2009; 5:e1000423.
576

577 Duguma D, Hall MW, Rugman-Jones P et al. Developmental succession of the microbiome of
578 *Culex* mosquitoes. *BMC Microbiol* 2015; 15:140.
579

580 Escudie, F, Auer L, Bernard M et al. FROGS: find rapidly OTU with galaxy solution.
581 F1000Research 2016; 5:1287–1294.
582
583 Gabrieli P, Caccia S, Varotto-Boccazzi I, Arnoldi I et al. Mosquito trilogy: microbiota,
584 immunity and pathogens, and their implications for the control of disease Transmission. *Front*
585 *Microbiol* 2021; 12:630438.
586
587 Gloria-Soria A, Chiodo TG, Powell JR. Lack of evidence for natural *Wolbachia* Infections in
588 *Aedes aegypti* (Diptera: Culicidae). *J Med Entomol* 2018; 55:1354-1356.
589 Gu, Z. 2021. ComplexHeatmap: Make Complex Heatmaps; Bioconductor version: Release
590 (3.12).
591
592 Guégan, M, Minard G, Tran FH et al. Short-term impacts of anthropogenic stressors on *Aedes*
593 *albopictus* mosquito vector microbiota. *FEMS Microbiol Ecol* 2018a; 94:fiy188.
594
595 Guégan M, Zouache K, Démichel C et al. The mosquito holobiont: fresh insight into
596 mosquito-microbiota interactions. *Microbiome* 2018b. 6:49.
597
598 Guégan M, Tran Van V, Martin E et al. Who is eating fructose within the *Aedes albopictus*
599 gut microbiota? *Environ Microbiol* 2020; 22(4):1193-1206.
600
601 Gusmão DS, Santos AV, Marini DC et al. Culture-dependent and culture-independent
602 characterization of microorganisms associated with *Aedes aegypti* (Diptera: Culicidae) (L.)
603 and dynamics of bacterial colonization in the midgut. *Acta Trop* 2010; 115(3):275–281.
604

605 Hery L, Guidez A, Durand AA et al. Natural variation in physicochemical profiles and
606 bacterial communities associated with *Aedes aegypti* breeding sites and larvae on Guadeloupe
607 and French Guiana. *Microb Ecol* 2021 81:93–109.
608

609 Hyde J, Gorham C, Brackney DE and Blaire S. Antibiotic resistant bacteria and commensal
610 fungi are common and conserved in the mosquito microbiome. *Plos one* 2019 14(8):
611 E0218907.
612

613 Inácio da Silva LM, Dezordi FZ, Paiva MHS et al. Systematic review of Wolbachia symbiont
614 detection in mosquitoes: An Entangled Topic about Methodological Power and True
615 Symbiosis. *Pathogens* 2021; 10:39.
616

617 Jupatanakul N, Sim S, Dimopoulos G. The insect microbiome modulates vector competence
618 for arboviruses. *Viruses* 2014; 6(11): 4294–4313.
619

620 Gimonneau G, Tchioffo MT, Abate L et al. Composition of *Anopheles coluzzii* and *Anopheles*
621 *gambiae* microbiota from larval to adult stages. *Infect Genet Evol* 2014 ; 28:715-24.
622

623 Juma EO, Allan BF, Kim CH et al. The larval environment strongly influences the bacterial
624 communities of *Aedes triseriatus* and *Aedes japonicus* (Diptera: Culicidae). *Sci Rep* 2021; 11:
625 7910.
626

627 Kõljalg U, Nilsson RH, Abarenkov K *et al.* Towards a unified paradigm for sequence-based
628 identification of fungi. *Mol Ecol* 2013; 22:5271-5277.
629

630 Kulkarni A, Yu W, Jiang J et al. *Wolbachia pipientis* occurs in *Aedes aegypti* populations in
631 New Mexico and Florida, USA. *Ecol Evol* 2019; 9:6148–6156.
632
633 MacLeod HJ, Dimopoulos G, Short SM. Larval diet abundance influences size and
634 composition of the midgut microbiota of *Aedes aegypti* mosquitoes. *Front Microbiol*
635 2021;12:645362.
636
637 Malassigné S, Valiente Moro C, Luis P. Mosquito mycobiota: an overview of non-
638 entomopathogenic fungal interactions. *Pathogens* 2020; 9(7):564.
639
640 McMurdie PJ and Holmes S. Phyloseq: An R package for reproducible interactive analysis
641 and graphics of microbiome census data. *PLoS One* 2013; 8:e61217.
642
643 Minard G, Mavingui P, Moro CV. Diversity and function of bacterial microbiota in the
644 mosquito holobiont. *Parasit Vectors* 2013; 6:146.
645
646 Minard G, Tran F-H, Dubost A, et al. Pyrosequencing 16S rRNA genes of bacteria associated
647 with wild tiger mosquito *Aedes albopictus*: a pilot study. *Front Cell Infect Microbiol* 2014;
648 4:59.
649
650 Minard G, Tran FH, Van VT et al. French invasive Asian tiger mosquito populations harbor
651 reduced bacterial microbiota and genetic diversity compared to Vietnamese autochthonous
652 relatives. *Front Microbiol* 2015; 6:970.
653

654 Minard G, Tran FH, Tran Van V et al. Shared larval rearing environment, sex, female size
655 and genetic diversity shape *Ae. albopictus* bacterial microbiota. *PLoS One* 2018;13:e0194521.
656

657 Muturi EJ, Dunlap C and Caceres CE. Microbial communities of container aquatic habitats
658 shift in response to *Culex restuans* larvae. *FEMS Microbiol. Ecol* 2020; 96:fiaa112.
659

660 Nilsson LKJ, Sharma A, Bhatnagar RK et al. Presence of *Aedes* and *Anopheles* mosquito
661 larvae is correlated to bacteria found in domestic water-storage containers. *FEMS Microbiol*
662 *Ecol* 2018; 94:fiy058.
663

664 Oksanen J, Blanchet GF, Friendly M et al. Vegan: community ecology package. R package
665 version 2.4-6. 2018; <https://CRAN.R-project.org/package=vegan>.
666

667 Powell JR and Tabachnick WJ. History of domestication and spread of *Aedes aegypti* - A
668 Review. *Mem Inst Oswaldo Cruz* 2013; 108:11-17.
669

670 R Core Team. R: A Language and Environment for Statistical Computing. Vienna, Austria: R
671 Foundation for Statistical Computing. 2018; <https://www.R-project.org/>.
672

673 Raharimalala FN, Ravaomanarivo LH, Ravelonandro P et al. Biogeography of the two major
674 arbovirus mosquito vectors, *Aedes aegypti* and *Aedes albopictus* (Diptera, Culicidae), in
675 Madagascar. *Parasit Vectors*. 2012; 5:56.
676

677 Ranasinghe AK and Amarasinghe LD. Naturally occurring microbiota associated with
678 mosquito breeding habitats and their effects on mosquito larvae. *Biomed Res Int*. 2020;

679 4065315.
680
681 Russell BM, Kay BH, Shipton W. Survival of *Aedes aegypti* (Diptera: Culicidae) Eggs in
682 Surface and Subterranean Breeding Sites During the Northern Queensland Dry Season. *J Med*
683 *Entomol* 2001; 38(3): 441–445.
684
685 Schoch CL, Seifert KA, Huhndorf S et al. Nuclear ribosomal internal transcribed spacer (ITS)
686 region as a universal DNA barcode marker for Fungi. *Proc Natl Acad Sci.* 2012; 109: 6241–
687 6246.
688
689 Schloss PD *et al.* 2009. Introducing mothur: Open-source, platform-independent, community-
690 supported software for describing and comparing microbial communities. *Appl Environ*
691 *Microbiol* 2009; 75(23):7537-41
692
693 Scolari F, Sandionigi A, Carlassara M et al. Exploring changes in the microbiota of *Aedes*
694 *albopictus*: comparison among breeding site water, larvae, and adults. *Front. Microbiol.* 2021;
695 12:624170.
696
697 Tawidian P, Coon KL, Jumpponen A et al. Host-environment interplay shapes fungal
698 diversity in mosquitoes. *MSpehere* 2021; 6:5
699
700 Thongsripong P, Chandler JA, Green AB et al. Mosquito vector-associated microbiota:
701 Metabarcoding bacteria and eukaryotic symbionts across habitat types in Thailand endemic
702 for dengue and other arthropod-borne diseases. *Ecol Evol* 2018; 8:1352–1368.
703

704 van Tol and Dimopoulos G. Influences of the mosquito microbiota on vector competence.
705 *Adv Insect Physiol* 2016; 51:243–291.
706

707 Wang Q, Garrity GM, Tiedje JM et al. Naive Bayesian classifier for rapid assignment of
708 rRNA sequences into the new bacterial taxonomy. *Appl Environ Microbiol* 2007; 73: 5261-
709 5267.
710

711 Wang X, Liu T, Wu Y et al. Bacterial microbiota assemblage in *Aedes albopictus* mosquitoes
712 and its impacts on larval development. *Mol Ecol* 2018; 27:2972-2985.
713

714 Wang Y, Hoon Eum J, Harrison RE et al. Riboflavin instability is a key factor underlying the
715 requirement of a gut microbiota for mosquito development. *Proc Natl Acad Sci* 2021;
716 118(15):e2101080118.
717


718 Wu P, Sun P, Nie K, et al. A gut commensal bacterium promotes mosquito permissiveness to
719 arboviruses. *Cell Host Microbe* 2019; 25(1):101-112.e5.
720

721 Wickham H, Francois R, Henry L et al. Dplyr: A grammar of data manipulation. URL
722 <https://CRAN.Rproject.org/package=dplyr>.
723

724 Xia S, Dweck HKM, Lutomiah J et al. Larval sites of the mosquito *Aedes aegypti formosus* in
725 forest and domestic habitats in Africa and the potential association with oviposition evolution.
726 *Ecol Evol* 2021; 11:16327-16343.
727

728 Yilmaz P, Parfrey LW, Yarza P et al. The SILVA and “All-species Living Tree Project
729 (LTP)” taxonomic frameworks. *Nucleic Acids Res* 2014; 42: D643–D648.
730

731 **Figures**



732

733 **Figure 1.** Non-metric multidimensional scaling (NMDS) ordination displaying bacterial (A)

734 and fungal (B) Operational Taxonomic Units (OTUs) composition across larvae and water

735 habitats originated from different breeding sites. NMDS ordination of samples was based on a


736 Bray-Curtis dissimilarity matrix of the square root-transformed abundance data obtained from

737 read counts. To better distinguish the larvae (indicated by a circle) collected from the same

738 water habitat (indicated by a triangle), the shade of the corresponding colour is slightly

739 different from a breeding site to another.

740



741

742 **Figure 2.** Non-metric multidimensional scaling (NMDS) ordination displaying bacterial (A)


743 and fungal (B) Operational Taxonomic Units (OTUs) composition between larvae (L) and

744 aquatic habitats (W) indicated in red and black colour, respectively. The ovals refer to the

745 standard error of the centroid and the vectors refer to the distance of each sample from the

746 centroid of its group.


747



748

749 **Figure 3.** Correlation between Shannon indices of water and larval samples. The Shannon
 750 indices for water and larval samples are positively correlated for fungi ($\rho = 0.74$, $p = 3.379 \times$
 751 10^{-7}) (B) nor for bacteria ($\rho = -0.22$, $p = 0.18$) (A).

752



753


754 **Figure 4.** Venn diagrams showing the numbers of shared or unique bacterial (A) and fungal

755 (B) OTUs among larvae and water of breeding sites. A total of 542 and 375 bacterial and

756 fungal OTUs respectively (based on the sequencing of the V5-V6 and ITS-2 regions) were

757 detected among the two samples.


758



759

760 **Figure 5.** Correlation analysis between bacterial communities associated with the breeding
 761 site and larval bacterial microbiota. OTU pairs for which the residual correlation was
 762 estimated to be positive (respectively, negative) with 95% credibility level are shown in red
 763 (respectively, green) colour.


764



765

766 **Figure 6.** Correlation analysis between fungal communities associated with the breeding site
 767 and larval mycobiota. OTU pairs for which the residual correlation was estimated to be
 768 positive (respectively, negative) with 95% credibility level are shown in red (respectively,
 769 green) colour. OTU_19_Nigrospora_oryzae (92) is indicated in white colour as it is correlated
 770 to no other OTU and it is present in water in relatively large abundance but only in a single
 771 sample.

772



773


774 **Figure 7.** Relative abundance of the bacterial taxa at the phylum level. *Aedes aegypti* larvae

775 collected within the same breeding site are indicated by L_BD and next to the water sample

776 they come from indicated by BD. L_W indicates larvae belonging to the same breeding site

777 but for which the water sample from which they originated is absent.

778



779

780

781

782

783

784

785

786

787

788

789


790

791

792

Figure 8. Prevalence and relative abundance of operational taxonomic units in the different modalities. (A) bacteria in larvae, (B) bacteria in water, (C) fungi in larvae and (D) fungi in water. OTUs with the highest prevalence and abundance across the samples are highlighted in red and labeled with their OTU ID (Panel A: Mycobacterium (OTU_11), Microbacterium (OTU_118), Unclassified Microbacteriaceae (OTU_133), Actinomycetospora (OTU_102), Uncultured Burkholderiaceae (OTU_59), Clostridium sensu stricto (OTU_47), Curtobacterium (OTU_185), Actinomycetospora (OTU_123), Williamsia (OTU_76), Uncultured Peptostreptococcaceae (OTU_10). Panel B: Uncultured Rhizobiales (OTU_181), Uncultured Rhizobiales incertae sedis (OTU_27), Uncultured Beijerinckiaceae (OTU_611), Methylobacterium (OTU_252), Fodinicola (OTU_161), Uncultured Rhizobiales (OTU_120), Uncultured Burkholderiaceae (OTU_59), Uncultured Rhizobiales incertae sedis (OTU_272), Pelomonas (OTU_234), Methylobacterium (OTU_131), Uncultured Rhizobiaceae (OTU_79). Panel C: Unclassified OTU (OTU_2), Trichosphaeriales incertae sedis (OTU_9), Penicillium

793 citrinum (OTU_31). Panel D: Uncultured Pleosporales (OTU_58), Toxicocladosporium
794 irritans (OTU_459), uncultured Aureobasidium (OTU_105). Prevalence is defined as the
795 overall infection rate of a given OTU overall samples. It was calculated for water and larvae
796 samples as follows: number of samples in which the OTU was detected / total number of
797 samples. Abundance refers to proportion occupied by an OTU within a sample among either
798 the bacterial or fungal community. It was calculated as follows: number of normalized reads
799 assigned to the OTU in a given samples / total number of normalized reads.
800



801

802 **Figure 9.** Relative abundance of the fungal taxa at the phylum and subphylum levels. *Aedes*


803 *aegypti* larvae collected within the same breeding site are indicated by L_BD and next to the

804 water sample they come from indicated by BD. L_W indicates larvae belonging to the same

805 breeding site but for which the water sample from which they originated is absent.


806

Microbiota of larvae ranked by breeding sites




808 **Figure S1.** Correlation analysis between bacterial communities associated with the aquatic
809 habitat and larval bacterial microbiota, ranked by breeding sites. A white line separates each
810 breeding site. Individuals for which the residual correlation was estimated to be positive
811 (respectively, negative) with 95% credibility level are shown in red (respectively, blue)
812 colour.

813



815 **Figure S2.** Correlation analysis between fungal communities associated with the aquatic
 816 habitat and larval mycobiota, ranked by breeding sites. Each breeding site is separated by a
 817 white line. Individuals for which the residual correlation was estimated to be positive
 818 (respectively, negative) with 95% credibility level are shown in red (respectively, blue)
 819 colour.
 820

ASCOMYCOTA



821

822 **Figure S3.** Relative abundance of the Ascomycota at the order level. *Aedes aegypti* larvae


823 collected within the same breeding site are indicated by L_BD and next to the water sample

824 they come from indicated by BD. L_W indicates larvae belonging to the same breeding site

825 but for which the water sample from which they originated is absent.

826

BASIDIOMYCOTA



827

828 **Figure S4.** Relative abundance of the Basidiomycota at the order level. *Aedes aegypti* larvae
 829 collected within the same breeding site are indicated by L_BD and next to the water sample
 830 they come from indicated by BD. L_W indicates larvae belonging to the same breeding site
 831 but for which the water sample from which they originated is absent.

832 **Table S1.** List of samples analysed with related information. Third- and fourth-instar larvae
833 as well as water were sampled in artificial breeding sites such as small flasks or containers.
834 Samples were collected with a sterile plastic pipette into a sterile 50-ml falcon tube with
835 filter-top lid. Water samples were agitated before sampling to mix detritus and the upper
836 surface to a depth of 10cm was collected representing \approx 50 ml per breeding site. In grey are
837 indicated samples that were successfully sequenced and in white those for which sequencing
838 failed.

839

840 **Table S2.** Number of bacterial OTUs and Shannon diversity index calculated for each sample
841 (one sample = one larva or one water sample).

842

843 **Table S3.** Number of fungal OTUs and Shannon diversity index calculated for each sample
844 (one sample = one larva or one water sample).

Table S1. List of samples analysed with related information. Third- and fourth-instar larvae as well as water were sampled in artificial breeding sites such as small flasks or containers. Samples were collected with a sterile plastic pipette into a sterile 50-ml falcon tube with filter-top lid. Water samples were agitated before sampling to mix detritus and the upper surface to a depth of 10cm was collected representing \approx 50 ml per breeding site. In grey are indicated samples that were successfully sequenced and in white those for which sequencing failed.

Breeding dite (BD)		Date of sampling	Larvae		Water sequencing		Larva sequencing	
Type	Sample name		Number collected / BD	Sample name	16S	ITS	16S	ITS
Plastic flask	BD1	02/12/17	2	L1-BD1 L2-BD1				
Plastic flask	BD2	02/12/17	3	L1-BD2 L2-BD2 L3-BD2 L4-BD2				
Plastic bucket	BD3	03/12/17	3	L1-BD3 L2-BD3 L3-BD3				
Metal container	BD4	03/12/17	4	L1-BD4 L2-BD4 L3-BD4 L4-BD4				
Plastic flask	BD5	04/12/17	1	L1-BD5				
Stainless steel container	BD6	20/04/18	3	L1-BD6 L2-BD6 L3-BD6				
Plastic bucket	BD7	25/04/18	3	L1-BD7 L2-BD7 L3-BD7				

Plastic flask	BD8	29/04/18	8	L1-BD8 L2-BD8 L3-BD8 L4-BD8 L5-BD8 L6-BD8 L7-BD8 L8-BD8	
Plastic flask	BD9	25/04/18	3	L1-BD9 L2-BD9 L3-BD9	
Plastic flask	BD10	29/04/18	4	L1-BD10 L2-BD10 L3-BD10 L4-BD10 L5-BD10	
Plastic flask	BD11	02/05/18	5	L1-BD11 L2-BD11 L3-BD11 L4-BD11 L5-BD11	
Plastic flask	BD12	25/04/18	3	L1-BD12 L2-BD12 L3-BD12	
Metal container	BD13	28/04/18	2	L1-BD13 L2-BD13 L3-BD13 L4-BD13	
Metal container	BD14	03/05/18	3	L1-BD14	

				L2-BD14	
				L3-BD14	
Plastic cup	BD15	02/05/18	5	L1-BD15	
				L2-BD15	
				L3-BD15	
				L4-BD15	
				L5-BD15	
Plastic flask	BD16	21/04/18	2	L1-BD16	
				L2-BD16	
Plastic flask	BD17	01/05/18	2	L1-BD17	
				L2-BD17	

Table S2. Number of bacterial OTUs and Shannon diversity index calculated for each sample (one sample = one larva or one water sample).

Sample type	Sample name	No. Of OTUs	Mean number of OTUs per larva	Shannon index
Larva	L1-BD1	147	160±18	1.17
Larva	L2-BD1	173		1.18
Water	BD1	138		2.69
Larva	L1-BD2	146	169±21	2.54
Larva	L2-BD2	174		2.93
Larva	L3-BD2	187		3.32
Water	BD2	213		2.39
Larva	L1-BD3	273	275±18	3.50
Larva	L2-BD3	258		3.49
Larva	L3-BD3	293		3.84
Water	BD3	200		2.18
Larva	L1-BD4	252	218±64	3.50
Larva	L2-BD4	257		3.53
Larva	L3-BD4	122		1.05
Larva	L4-BD4	242		3.24
Water	BD4	266		2.56
Larva	L1-BD5	215	–	2.47
Water	BD5	205		3.04
Larva	L1-BD6	112	146±35	0.75
Larva	L2-BD6	181		1.09
Larva	L3-BD6	144		0.65
Water	BD6	160		3.22
Larva	L1-BD12	172	162±22	1.45
Larva	L2-BD12	178		2.68
Larva	L3-BD12	137		1.75
Larva	L1-BD7	192	240±43	3.42

Larva	L2-BD7	272		3.34
Larva	L3-BD7	207		3.15
Water	BD7	190		2.41
Larva	L1-BD9	278		2.93
Larva	L2-BD9	229	215±71	1.67
Larva	L3-BD9	138		1.72
Water	BD9	167		1.62
Larva	L1-BD10	174		3.07
Larva	L2-BD10	219	190±24	3.16
Larva	L3-BD10	168		3.06
Larva	L4-BD10	199		3.31
Water	BD10	223		3.20
Larva	L3-BD12	291	—	
Larva	L1-BD13	270	273±4	3.35
Larva	L2-BD13	276		2.83
Larva	L1-BD14	275		2.98
Larva	L2-BD14	280	281±7	1.17
Larva	L3-BD14	289		1.20
Larva	L1-BD15	263		1.62
Larva	L2-BD15	208		1.12
Larva	L3-BD15	296	226±85	0.92
Larva	L4-BD15	276		2.54
Larva	L5-BD15	86		2.94
Larva	L6-BD8	353		2.88
Larva	L7-BD8	272		2.63
Larva	L8-BD8	338	240±72	2.85
Larva	L1-BD8	159		1.36
Larva	L2-BD8	206		2.74
Larva	L3-BD8	199		3.16

Larva	L4-BD8	195		3.11
Larva	L5-BD8	197		3.05
Water	BD8	198		3.43
Larva	L1-BD16	294	293±2	3.55
Larva	L2-BD16	291		2.95
Larva	L1-BD11	189		2.49
Larva	L2-BD11	204		3.10
Larva	L3-BD11	230	239±46	3.25
Larva	L4-BD11	296		3.56
Larva	L5-BD11	276		2.96
Water	BD11	263		3.26
Larva	L1-BD17	284	332±67	2.48
Larva	L2-BD17	379		3.51

Table S3. Number of fungal OTUs and Shannon diversity index calculated for each sample (one sample = one larva or one water sample).

Sample type	Sample name	No. Of OTUs	Mean number of OTUs per larva	Shannon index
Larva	L1-BD1	312	268±62	4.07
Larva	L2-BD1	224		3.91
Water	BD1	202		3.26
larva	L1-BD2	56	91±24	2.81
Larva	L2-BD2	100		3.28
larva	L4-BD2	97		3.19
Larva	L3-BD2	109		3.09
Water	BD2	99		1.60
larva	L1-BD3	105	83±20	1.87
Larva	L2-BD3	77		1.53
Larva	L3-BD3	66		1.59
Water	BD3	47		1.42
Larva	L1-BD4	84	78±7	1.64
Larva	L2-BD4	79		1.52
Larva	L3-BD4	69		1.36
Larva	L4-BD4	81		1.32
Water	BD4	93		2.33
Larva	L1-BD5	44	–	0.15
Water	BD5	38		0.04
Larva	L1-BD6	141	141±24	3.03
Larva	L2-BD6	186		3.13
Larva	L3-BD6	151		3.25
Water	BD6	82		2.95
Larva	L1-BD12	67	58±13	1.88
Larva	L2-BD12	43		1.74
Larva	L3-BD12	65		2.08

larva	L1-BD7	54		1.74
Larva	L2-BD7	99	76±23	2.07
Larva	L3-BD7	75		1.66
Water	BD7	81		1.40
Larva	L1-BD9	70		1.01
Larva	L2-BD9	59	62±7	1.02
Larva	L3-BD9	56		1.15
Water	BD9	39		0.23
Larva	L1-BD10	72		1.34
Larva	L2-BD10	77		1.58
Larva	L3-BD10	81	76±6	1.47
Larva	L5-BD10	68		1.73
Larva	L4-BD10	81		1.45
Water	BD10	85		1.42
Larva	L3-BD13	97		2.98
Larva	L4-BD13	104	107±9	3.29
Larva	L1-BD13	111		3.34
Larva	L2-BD13	117		3.16
Larva	L1-BD14	106		3.20
Larva	L2-BD14	115	110±5	3.19
Larva	L3-BD14	110		3.37
Larva	L1-BD15	62		2.09
Larva	L2-BD15	71		2.19
Larva	L3-BD15	62	72±18	2.26
Larva	L4-BD15	60		2.36
Larva	L5-BD15	103		0.95
Larva	L6-BD8	137		3.31
Larva	L7-BD8	85	125±34	2.67
Larva	L8-BD8	146		3.11

Larva	L1-BD8	131		3.53
Larva	L2-BD8	76		3.19
Larva	L3-BD8	179		4.11
Larva	L4-BD8	138		3.72
Larva	L5-BD8	105		3.63
Water	BD8	72		2.90
Larva	L1-BD16	135	138±4	2.51
Larva	L2-BD16	140		2.35
Larva	L1-BD11	75		2.10
Larva	L2-BD11	72		2.11
Larva	L3-BD11	84	124±66	1.47
Larva	L4-BD11	211		4.18
Larva	L5-BD11	180		4.14
Larva	L1-BD17	105	139±48	3.30
Larva	L2-BD17	173		3.24