



Fibropapillomatosis Prevalence and Distribution in Immature Green Turtles (*Chelonia mydas*) in Martinique Island (Lesser Antilles)

Thibaut Roost, Jo-Ann Schies, Marc Girondot, Jean-Patrice Robin, Pierre Lelong, Jordan Martin, Flora Siegwalt, Lorène Jeantet, Mathieu Giraudeau, Guillaume Le Loc'h, et al.

► To cite this version:

Thibaut Roost, Jo-Ann Schies, Marc Girondot, Jean-Patrice Robin, Pierre Lelong, et al.. Fibropapillomatosis Prevalence and Distribution in Immature Green Turtles (*Chelonia mydas*) in Martinique Island (Lesser Antilles). *EcoHealth*, 2022, 19 (12), pp.190-202. 10.1007/s10393-022-01601-y . hal-03692770

HAL Id: hal-03692770

<https://hal.science/hal-03692770>

Submitted on 17 Jun 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License



Fibropapillomatosis prevalence and distribution in immature green turtles in Martinique

| | |
|------------------|--|
| Journal: | <i>EcoHealth</i> |
| Manuscript ID | ECH-21-0147.R1 |
| Manuscript Type: | Original Contribution |
| Keywords: | marine turtles, infectious disease, epizootiology, environmental quality |
| | |

SCHOLARONE™
Manuscripts

Authors:

Roost, Thibaut Schies, Jo-Ann Girondot, Marc
 Robin, Jean-Patrice Lelong, Pierre Martin, Jordan
 Siegwalt, Flora Jeantet, Lorène Giraudeau, Mathieu
 Le Loch, Guillaume Bejarano, Manola Bonola, Marc
 Benhalilou, Abdelwahab Murgale, Céline Andreani, Lucas
 Jacaria, François Campistron, Guilhem Lathière, Anthony
 Martial, François Hielard, Gaëlle Arqué, Alexandre
 Régis, Sidney Lecerf, Nicolas Frouin, Cédric
 Lefebvre, Fabien Aubert, Nathalie Flora, Frédéric
 Pimentel, Esteban Lafolle, Rachelle Thobor, Florence
 Arthus, Mosiah Etienne, Denis Lecerf, Nathaël
 Allenou, Jean-Pierre Desigaux, Florian Larcher, Eugène
 Larcher, Christian Curto, Alberto Lo Befort, Joanne
 Maceno-Panevel, Myriane Lepori, Muriel Chevallier, Pascale
 Chevallier, Tao Meslier, Stéphane Landreau, Anthony
 Hibold, Caroline Le Maho, Yvon Chevallier, Damien

Fibropapillomatosis prevalence and distribution in immature green turtles (*Chelonia mydas*) in Martinique Island (Lesser Antilles)

Abstract

Fibropapillomatosis (FP) threatens the survival of green turtle (*Chelonia mydas*) populations at a global scale and human activities are regularly pointed as causes of high FP prevalence. However, the association of ecological factors with the disease's severity in complex coastal systems has not been well established and requires further studies. Based on a set of 405 individuals caught over ten years, this preliminary study provides the first insight of FP in Martinique Island, which is a critical development area for immature green turtles. Our main results are: i) 12.8% of the individuals were affected by FP, ii) FP has different prevalence and temporal evolution between very close sites, iii) green turtles are more frequently affected on the upper body part such as eyes (41.4%), fore flippers (21.9%) and the neck (9.4%), and iv) high densities of individuals are observed on restricted areas. We hypothesise that turtle's aggregation enhances horizontal transmission of the disease. FP could represent a risk for immature green turtles' survival in the French West Indies, a critical development area, which replenishes the entire Atlantic population. Continuing scientific monitoring is required to identify which factors are implicated in this panzootic disease and ensure the conservation of the green turtle at an international scale.

Keywords: marine turtles, infectious disease, epizootiology, environmental quality

1
2
3 24 **1. Introduction**
4

5
6 25 Green turtle *Chelonia mydas* (Linnaeus, 1758) populations have to face a wide range
7
8 26 of anthropogenic threats such as bycatch, boat strike, seagrass meadows destruction, dredging
9
10 27 operations, marine pollution, poaching, and tourism development (Domiciano et al., 2017;
11
12 28 Herbst & Klein, 1995; Jones et al., 2015; Rossi et al., 2019) leading to decreasing trend for the
13
14 29 global population. This resulted in the classification of the species as “endangered” on the
15
16 30 IUCN Red List (Seminoff, 2004). In addition to these threats, the epizootic disease
17
18 31 fibropapillomatosis (FP) is an emerging global threat for green turtles (Bjorndal, 1999; Herbst
19
20 32 & Klein, 1995).

21
22
23
24 33 FP is a neoplastic disease characterised by the growth of tumours mostly on soft tissues
25
26 34 and the shell (Herbst, 1994). The tumours can have a diameter of up to 30 cm and alter vision,
27
28 35 swimming, foraging, orientation abilities or even breathing (Jones et al., 2015; Williams et al.,
29
30 36 1994). In the disease’s end stage, internal tumours can develop on the lungs, kidneys, heart or
31
32 37 digestive tract, which can lead to death (Jones et al., 2015). FP has been observed on the seven
33
34 38 existing species of marine turtles but has only reached a panzootic level in the green turtle
35
36 39 (Jones et al., 2015). This disease spread to numerous regions worldwide in the 1980’s,
37
38 40 especially in the Atlantic, the Caribbean (i.e. Cayman Islands, Puerto Rico, Virgin Islands,
39
40 41 Barbados, Venezuela, Colombia, Nicaragua, Costa Rica, Panama, and Belize), and in the Indo-
41
42 42 Pacific region with prevalence varying from 1.4% to 92% (Adnyana et al., 1997; Herbst, 1994).
43
44 43 FP’s precise aetiology is not yet fully known. However, the Chelonid HerpesVirus 5 (ChHV5)
45
46 44 has been regularly associated with this disease, and it is now a consensus that this virus could
47
48 45 be the most likely cause of FP (Chaves et al., 2017; Domiciano et al., 2017; Jones et al., 2015).
49
50 46 Moreover, the hypothesis of infection through horizontal transmission (i.e. from one individual
51
52 47 to another) during turtle’s settlement into coastal habitats has been widely recognized. Indeed,
53
54 48 there has been no observation of FP clinical signs on recently recruited turtles (Jones et al.,
55
56
57
58
59
60

2020; Patricio et al., 2016; Shaver et al., 2019). Furthermore, it has been proved that horizontal transmission can be promoted by parasite marine leech, which act as mechanical vectors of ChHV5 (Rittenburg et al., 2021).

Environmental conditions have been proposed to play an important role in the emergence of FP (Herbst & Klein, 1995; Herbst et al., 2004) because of the multiple reports of high variations of prevalence between very close geographic regions (Herbst, 1994; Jones et al., 2015). Herbst & Klein (1995) suggested that higher sea temperature could induce faster tumour growth that in turn would result in more severe FP in green turtles. FP is also frequently associated with poor water quality (e.g. pollution, eutrophication) in coastal areas near human activities and/or with low hydrodynamics (Hargrove et al., 2016; Torezani et al., 2010). Metal contaminants (da Silva et al., 2016), persistent organic pollutants (Foley et al., 2005), and eutrophication coupled with a change in diet quality (Van Houtan et al., 2014) have also been suspected to enhance green turtles' susceptibility to FP.

Located in the Lesser Antilles of the West Indies in the eastern Caribbean, Martinique Island hosts an important population of immature green turtles in which clinical signs of FP infection (i.e. tumours) have been observed (Bonola et al., 2019). Moreover, Martinique has many sheltered bays that support the settlement of multi-species seagrass meadows on large shallow areas, particularly favourable to green turtles (Siegwalt et al., 2020). Indeed, the island is an important developmental area for these immatures, who show high fidelity to their foraging grounds for several years (Siegwalt et al., 2020) before performing their developmental migration in the Caribbean and the entire Atlantic (Chambault et al., 2018). In areas where green turtles tend to congregate for feeding, we hypothesize that positive interactions between individuals would facilitate horizontal transmission of FP. Moreover, an important urban development and poor sanitation of wastewater of this island lead to discharge

releases of polluted water in the marine environment (Hily et al., 2010) that could drive enhanced environmental pollution and/or trigger an eutrophication phenomenon.

This study is the first one to focus on FP prevalence in Martinique. From 2010 to 2019, capture-mark-recapture (CMR) data as well as the presence of tumours have been collected on immature green turtles along the western coast of Martinique. During 2018 and 2019, data on animal aggregation were gathered whereas, in 2021, seawater samples were collected in order to describe environmental quality. These datasets offer a unique opportunity to analyse the emergence of FP in green turtles (based on symptomatic individuals) in the critical developmental area of Martinique. The aims of this preliminary study were (i) to describe FP evolution through time between different high-fidelity grounds, (ii) look for possible environmental variables affecting the dynamics of this disease, (iii) describe turtles' distribution and compare it to FP prevalence and dynamics.

2. Materials and methods

2.1. Study area

This study was conducted along 60 km of the western coast of Martinique (14°30'9.64'N, 61°5'11.85'W, France). Green turtles were mainly present in six bays of the Caribbean coast of the island that are, from north to south, Anse Noire, Anse Dufour, Grande Anse, Anse du Bourg, Anse Chaudière and Petite Anse (Fig. 1). These sites were identified as critical for immature green turtle foraging (Siegwalt et al., 2020).

2.2. Capture-mark-recapture program (CMR)

The CMR program used for this study is based on the tagging of animals by the injection of passive integrated transponders (PIT). Since 2010, with the exception of 2014, immature green turtles have been captured by freedivers at depths up to 25 m (Fig. 1). Capture

methodology and procedures are described in Nivière et al. (2018) and in Bonola et al. (2019). Date and geographical coordinates were recorded for each turtle capture. The presence of a PIT was checked using a universal reader (GR251, TROVAN). In case of absence of a transponder, a PIT (ID-100, TROVAN) was injected into the right triceps. The search for FP was conducted by carefully examining each turtle for the presence of external tumours looking like single or multiple raised masses. Tumours were located on the turtles' bodies according to the following body parts: eyes, head, nape, neck, shoulders, fore flippers, carapace, plastron, back flippers, and tail base.

The study met the French legal and ethical requirements. The protocol was approved by the Conseil National de la Protection de la Nature and the French Ministry for Ecology (permit numbers: 2013154-0037 and 201710-0005) and followed the recommendations of the Police Prefecture of Martinique.

2.3. Environmental conditions

Sea surface temperature (SST) data were obtained from the NOAA/OAR/ESRL PSL public database (Colorado, USA, <https://psl.noaa.gov/> (accessed 5-10-2020)). Daily means of the 'Optimal Interpolation Sea Surface Temperature V2' were extracted for the closest coordinates to Martinique Island, which were 14.55° latitude and -61.25° longitude (Fig. S1). The NOAA 1/4° daily 'Optimum Interpolation Sea Surface Temperature' is an analysis constructed by combining observations from different platforms (satellites, ships, buoys, and Argo floats) on a regular global grid (0.25° latitude x 0.25° longitude grid). These values were averaged per trimester from 2010 to 2019 and applied to every site since we had only a single sample point close to Martinique Island (Fig. S1).

Net primary production (NPP) data were collected from the Ocean Productivity public database (<https://sites.science.oregonstate.edu/ocean.productivity/index.php>). Monthly means

of NPP, based on the Vertically Generalized Production Model (Behrenfeld & Falkowski, 1997) from MODIS satellite measures, were extracted for six points alongside Martinique's West coast (Fig. S1). Each turtle's capture site was assigned the NPP means of the nearest sampling point.

For seawater quality, two (for Anse Noire, Anse Dufour, Anse Chaudière) to three (for Grande Anse, Anse du Bourg, Petite Anse) sampling points per foraging site were chosen prior to field work (Siegwalt et al., 2020). On each sampling location, seawater was sampled multiple times with a plunger sampler ($V = 1\text{ L}$) at 5 m depth in order to fill a 2 L vial for total chlorophyll *a* analysis, a 250 mL vial for bacterial analysis (*E. coli* and enterococci) and four 150 mL vials for chemical analysis (ammonium, nitrites, nitrates and phosphates). Chlorophyll *a* vials were protected from any light by wrapping them in aluminium paper. Vials were stored in coolers away from sunlight. All samples were collected on the same day in 2021 and sent to the Laboratoire Territorial d'Analyses de Martinique. Chemical and chlorophyll *a* analyses were done according to Aminot & Chaussepied (1983) while *Escherichia coli* (*E. coli*) and enterococci were measured following European standards NF EN ISO 9308-3 and NF EN ISO 7899-1, respectively. We looked at chemical and biological parameters in order to seek for potential eutrophication and at European standards bacterial parameters to describe seawater sanitary safety.

141

142 **2.4. Density surveys and mapping**

Population density surveys of green turtles were carried out in 2018 and 2019 on several bays: Anse Noire, Anse Dufour, Grande Anse, Anse du Bourg, Anse Chaudière and Petite Anse. In order to count a maximum number of turtles under the water surface, one diver (observer) was connected by a 10 m rope to a boat. Transects specific to each bay, parallel to the beach and arranged from coast to sea, were previously created and integrated into a GPS

(GarminTrex) in order to follow a predefined course. The chosen distance between transects (10 to 30 m depending on the visibility) was intended to cover the greatest area and sample the largest number of individuals and to minimize the risk of missing an individual. Standardised hand signals have been established so that the observer could communicate with the operators aboard the boat to report the number of observed turtles. Each observation has been associated with its GPS coordinates.

Density maps were created to visualise the spatial distribution and density of green turtles using the software QGIS 2.18 (2016) and the legal CRS RGAF09 (EPSG:5490) in use at that time for French West Indies. Turtle sighting points and predefined transects were placed on Mapbox Satellite v9 satellite image in order to set up grids of hexagonal 1 ha-cells using the QMarxan extension, whose extents correspond to the surveyed areas of each bay. Then, turtle densities were determined in each cell.

2.5. Data analysis

FP data were obtained through CMR surveys and visual detection of FP-indicating tumours. Given the movements of individuals between Anse Noire and Anse Dufour (< 500 m apart; Siegwalt et al., 2020), the two sites were considered as a single entity called Anse Noire/Dufour. With the same considerations, Anse du Bourg and Anse Chaudière became Anse du Bourg/Chaudière.

Relationships between FP prevalence and environmental conditions were assessed with generalised linear models (GLM). Thus, in order to describe geographic disparities between close bays, interannual evolution, look for SST's influence on the disease's severity, and NPP's influence according its relationship with immature green turtle's body mass (Bonola et al., 2019), fixed factors included *capture site*, *year*, *mean SST*, and *mean NPP*, respectively. *Year* was considered as a discrete factor instead of a numeric variable. The distribution of the

173 explained variable was binomial (presence or absence of FP) and a logit link was therefore
174 used. A model selection was performed with the use of the Akaike information criterion (AIC;
175 Akaike, 1973) and relative Akaike weight. Models with a difference lower than 2 in their
176 respective AIC were considered similar (Burnham & Anderson, 2002).

177 To highlight which individual body parts were the most affected by FP, we calculated
178 infection proportions per body part by dividing the number of turtles with the presence of
179 tumour(s) on the specific body part by the total number of turtles affected by FP. Individuals
180 captured several times were counted as one turtle having FP.

181 Generalised Linear Mixed Models (GLMMs) were used to explore potential differences
182 in the seven seawater variables measured (i.e. ammonium, nitrites, nitrates, phosphates, total
183 chlorophyll *a*, *E. coli*, and enterococci) between field sites. As for FP analysis, Anse
184 Noire/Dufour and Anse du Bourg/Chaudière were considered unique sites. Sampling
185 replication was included as a random effect for every GLMM to account for non-independence
186 of data. Models were fitted using the *lme4* package (Bates et al., 2012). Likelihood Ratio Tests
187 (LRTs) were performed using the *lmtest* package (Hothorn et al., 2015) to select the best-fit
188 model between the nul GLMM and the one with capture site fixed effect. Post-hoc estimations
189 and comparisons were performed when the *capture site* effect was kept using Estimated
190 Marginal Means (EMMs) with the *emmeans* package (Lenth et al., 2021). Degrees of freedom
191 were calculated using the Kenward-Roger method and p-values were adjusted for multiple
192 comparisons with Tukey adjustment. Significance thresholds were fixed at 0.05. All data
193 analysis was performed using R version 4.0.2 (R Core Team, 2020).

194

195

196

197

3. Results

3.1. *Fibropapillomatosis prevalence and body distribution*

From 2010 to 2019, 539 immature green turtle catches were performed on the Caribbean coast of Martinique, corresponding to 405 distinct individuals. The vast majority of the individuals were captured at Grande Anse ($n = 302$), Anse du Bourg/Chaudière ($n = 181$) and Anse Noire/Dufour ($n = 56$). The change in occurrence of FP could not be properly evaluated at Le Prêcheur, Saint-Pierre, Le Carbet, Cap enragé, and Petite Anse (Fig. S2). First, immature green turtles' abundance seemed to be weaker at Le Prêcheur, Saint-Pierre, Le Carbet, and Cap enragé. Second, at Petite Anse, captures were less successful since animals escaped more often, perhaps because of important human activities in this bay (pers. obs.).

A significant increase of immature green turtles' FP prevalence from 0.000 (IC95% [0.000; 0.278]) in 2011 to 0.168 (IC95% [0.108; 0.253]) in 2019 was observed when considering the three major sites together (Odd-ratio : 1.72, p -value < 0.001 ; Fig. 2). Post-hoc Tukey between capture sites highlighted significant differences of prevalence between Grande Anse and Anse du Bourg/Chaudière ($p < 0.001$). Indeed, FP prevalence remained close to zero from 2011 to 2019 in Grande Anse and also from 2015 to 2019 in Anse Noire/Dufour. In fact, the major increase in FP prevalence on the Martinique coast seemed to be restricted to Anse du Bourg/Chaudière, where no individual with FP being observed in 2011 and 2012 and with FP prevalence increasing from 2013 (0.11) to 2019 (0.50).

A total of 128 observations on 52 individuals were used to calculate the percentage distribution of tumours on individual body parts (Fig. 3) as tumour's body location was not reported during the first years of the CMR program. For the majority of turtles, lesions were observed on several parts of the body. Eyes were the most frequently affected body parts (21% of individuals for the left and 20% for the right). The front fins and neck's ventral part were

also frequently affected, with 9% of the individuals being affected on the left fin, 13% on the right fin, and 9% on the neck.

3.2. *Fibropapillomatosis and environmental cofactors*

AICs, Δ AICs and relative Akaike weights for the 11 tested GLMs are presented in Table 1. The null model is the model without effect and FP prevalence is therefore supposed constant. The lowest AIC (79.74) has been observed for the model with *capture site* and *year* effects. This model had a relative Akaike weight of 0.45. When adding *NPP* to the two previous fixed effects (Fig. S3), the model had a Δ AIC < 2 (1.70) and a relative Akaike weight of 0.19. *Temperature* effect (Fig. S4) associated with *capture site* and *year* resulted with a Δ AIC of exactly 2 compared to the model with the lowest AIC. All other models had a Δ AIC > 2 and lower relative Akaike weight values.

To validate the pertinence of the two selected models (*capture site + year* & *capture site + year + NPP*) we used them to predict FP prevalence. These predictions were compared with the observed prevalence (Fig. S5ab). Indeed, despite large CI, the vast majority of predicted values for both models were very close to those observed. However, some points diverged, such as an observed value at 1.00 but predicted at 0.55 or those observed at 0 and predicted at 0.25. Then, predicted values of both models were compared graphically (Fig. S5c). The relationship between predicted FP prevalence of the model *capture site + year* and those from *capture site + year + NPP* seemed to be nearly linear with a slope close to 1. Focusing on the *capture site + year + NPP* model, the regression between NPP and FP prevalence (Fig. S5d) was positive and greater values of NPP were associated with higher FP prevalence. However, regression's CI happened to be important.

Ammonium and nitrites levels were significantly different between sites (Fig. 4; Table 2). Ammonium levels were significantly lower at Grande Anse compared to Petite Anse with

respective means of $0.17 \mu\text{mol.L}^{-1}$ and $1.49 \mu\text{mol.L}^{-1}$ (Table 3). For nitrites, significant higher levels were observed at both Grande Anse and Anse du Bourg/Chaudière ($0.07 \mu\text{mol.L}^{-1}$ and $0.11 \mu\text{mol.L}^{-1}$, respectively) than at Anse Noire/Dufour and Petite Anse (Table 3). Nitrates levels were quite similar and there has been no significant differences between capture sites, with values ranging from $0.50 \mu\text{mol.L}^{-1}$ for Grande Anse and Anse du Bourg/Chaudière to $0.29 \mu\text{mol.L}^{-1}$ for Petite Anse (Fig. 4). No significant differences between sites were measured for phosphate levels even if it was significantly predicted by the capture sites (Table 3). Chlorophyll *a* means were not significantly different and close to $0.30 \mu\text{g.L}^{-1}$ for Anse Noire/Dufour, Anse du Bourg/Chaudière, and Petite Anse. Grande Anse had a slightly higher level ($0.43 \mu\text{g.L}^{-1}$; Fig. 4). There was no significant difference between locations for bacterial parameters (Table 2). However, a strong value for *E. coli* (mean = $1548 \text{ MPN.100mL}^{-1}$) was observed at Grande Anse. Anse du Bourg/Chaudière had an important presence of *E. coli* too (mean = $1836 \text{ MPN.100mL}^{-1}$) with three values out of five exceeding $1000 \text{ MPN.100mL}^{-1}$ and it was the only site where the presence of enterococci was measured (mean = 5 MPN.100mL^{-1}).

3.3. *Turtles density distribution*

The following number of turtles were observed in each bay in 2018 and 2019, respectively : 10 and 24 in Anse Noire, 8 and 3 in Anse Dufour, 93 and 100 in Grande Anse, 95 and 76 in Anse du Bourg, 14 and 7 in Anse Chaudière, 95 and 114 in Petite Anse. Densities of turtles per 1-ha cell were high in some restricted areas of the bays, with some variations between 2018 and 2019 (Fig. 5). Higher density cells were found in Anse du Bourg in 2018 ($10\text{--}12 \text{ individuals.ha}^{-1}$) and Petite Anse in 2019 ($13\text{--}15 \text{ individuals.ha}^{-1}$). More precisely, medium to high-density patches were located in the northern and central parts of Anse du Bourg, in the south of Petite Anse, and in the central and northern parts of Grande Anse, where

272 individuals seemed more dispersed in 2019 than in 2018. Anse Noire, Anse Dufour and Anse
 273 Chaudière had lower turtle concentrations for both years compared to the other sites (0-6
 274 individuals.ha⁻¹). Anse du Bourg (2018) and Petite Anse (2019) seemed to be the site with the
 275 highest heterogeneity considering the distribution of immature green turtles, with 1-ha cells
 276 reaching 10-15 individuals.ha⁻¹ right next to cells with no individuals observed.

277

278 4. Discussion

279 This study provides the first long-term study on FP prevalence over time in immature
 280 green turtles in the Lesser Antilles. Looking at the spatio-temporal distribution of FP, we noted
 281 an increase of global FP prevalence between 2011 and 2019 for Anse du Bourg/Chaudière,
 282 Anse Noire/Dufour and Grande Anse. Our data suggested differences between FP evolution
 283 patterns through time of these geographically close sites. While NPP was slightly positively
 284 associated with FP prevalence, *mean SST* had no effect on FP prevalence. In parallel, we
 285 demonstrated clear differences in seawater quality between the different bays. The
 286 heterogeneous distribution of tumours on the body of individuals was highlighted. Indeed, the
 287 majority of tumours was observed on the eyes, fore fins and the neck. Finally, the analysis of
 288 immature green turtles density distributions underscored their tendency to cluster in relatively
 289 small areas in each bay. Indeed, individuals were strongly concentrated in a restricted zone in
 290 Anse du Bourg. This could be one of the reasons explaining the higher FP prevalence in Anse
 291 du Bourg/Chaudière than in other capture sites, because of the proximity between individuals
 292 which could induce an increase of FP horizontal transmission. However, this assumption needs
 293 to be verified by further studies. Our study demonstrated a significant increase of the global FP
 294 prevalence on three sites with high densities of turtle : Anse Noire/Dufour, Grande Anse, and
 295 Anse du Bourg/Chaudière. The temporal patterns of FP prevalence seemed to be different
 296 between these three sites, which are very close geographically (i.e. from 0.36 to 7.2 km, for

distance's details in Siegwalt et al., 2020, Table S1). It is possible that FP prevalences were underestimated as we based our diagnosis on external lesions while it has been demonstrated that asymptomatic turtles can present high loads of ChHV5 (Chaves et al., 2017; Page-Karjian et al., 2015). In comparison, FP clinical signs were observed on half of the green turtles located in the Indian River lagoon, but none seemed to be affected at Sabellariid Worm Reef located one kilometer away in Florida (Herbst, 1994). A significant increase of FP prevalence over time was also highlighted for the same species in Texas with a prevalence under 5% before 2015 and rising to 35.2% in only three years (Shaver et al., 2019). At Pala'au, Molokai Island, the prevalence increase was quite similar to the one we observed at Anse du Bourg/Chaudière, rising from 1% to 61% in eight years (Jones et al., 2015). Thus, the literature suggests an important influence of local conditions on the disease's development and global trends toward an increase of FP prevalence.

Higher NPP values, but not temperature, were associated with higher FP prevalence. However, the positive link between NPP and FP prevalence was slight and associated with high uncertainty. Herbst (1994) proposed that tumour growth was more important in spring and summer because of higher water temperature. Murakawa et al. (2000) and Chaloupka et al. (2008), though, found no intra-annual variation of tumour size in stranded green turtles in Hawaii. Furthermore, Torezani et al. (2010) did not find an effect of temperature on FP prevalence with a range of temperatures from 27.5°C to 33.5°C. We had only a single SST sample point for the entire Martinique Island and mean SST values had a maximum variation of 2.7°C. This range was probably not sufficient to have a notable effect on FP-associated tumour growth. Thus, the influence of temperature on FP dynamics has not been evidenced at this time. However, Anse Noire/Dufour, Grande Anse, Anse du Bourg/Chaudière and Petite Anse are shallow bays where water temperature could greatly vary at a local scale compared

321 to our SST sample point located further at sea. Finer scale temperature data are necessary to
322 verify whether SST influences FP prevalence in Martinique.

323 Herbst (1994) suggested that human activities such as agriculture, industry and urban
324 development should impact the development of tumours through different pathways and
325 mechanisms. Santos et al. (2010) found that the high FP prevalence in a green turtle
326 developmental area was associated with its poor water quality (EEI = 2, ecological evaluation
327 index based on benthic macrophyte) in Espírito Santo Bay off Brazil. Concomitantly, on Oahu,
328 Maui and Hawaii islands, high prevalence areas corresponded to those with a greater nitrogen
329 footprint (Van Houtan et al., 2010). We therefore analysed seawater chemical, biological and
330 bacterial parameters in the different bays where we captured turtles in order to characterise
331 their environmental quality. The high level of NO₂ in Anse du Bourg/Chaudière might be
332 associated with the high prevalence observed at this site. Despite the lack of significant
333 differences between capture sites regarding bacterial parameters, we could also notice that
334 Anse du Bourg/Chaudière was the only site where the presence of enterococci was observed,
335 which reflect faecal contamination and poor water quality. This capture site also had several
336 samples exceeding the European sanitary threshold of 1000 MPN.100mL⁻¹ for *E. coli*.

337 The present results highlighted a significant difference in seawater quality between the
338 capture sites where immature green turtles are most present. The presence of a damaged outfall
339 releasing wastewater from a sewage treatment plant in Anse du Bourg (Impact Mer, 2016), as
340 well as an important pressure of pleasure boats could be responsible for the releasing of faecal
341 matter and nitrogen-enriched content (pers. obs.). Therefore, further environmental quality
342 studies are necessary to seek a possible link with the FP outbreak in Martinique. Indeed, higher
343 levels of arginine, an amino-acid known to enhance the emergence of tumours in some cases,
344 have been found in marine algae in watersheds with a higher nitrogen footprint due to human
345 land use (Hargrove et al., 2016). Considering the above-mentioned environmental context and

the high fidelity of immature green turtles to their feeding zone in the south-western bays of Martinique (Siegwalt et al., 2020), Anse du Bourg/Chaudière could provide an optimal environment for the contraction, persistence and transmission of this disease in green turtles.

By studying the density of individuals within each capture site, we have highlighted the presence of areas with high densities of individuals while others were left vacant. Differences in FP prevalence between locations could therefore be explained by the fact that some sites have higher turtle densities than others and that individuals are not uniformly dispersed within the bays. This could be especially the case in Anse du Bourg/Chaudière, where immature green turtles were highly concentrated in the North part of this small shallow bay while CMR data showed the highest FP prevalence for this location. Head rubbing or higher concentration of the ChHV5 in seawater in Anse du Bourg/Chaudière due to highly clustered individuals are possible explanations for the FP situation in this bay. On the other hand, at Anse Noire/Dufour where turtle's densities were the lowest, no FP outbreak has been recorded by the CMR monitoring.

The previous hypothesis of horizontal transmission is reinforced by our results on the relative distribution of tumours over the different body parts. Similar to Rossi et al. (2019), our results show that the neck, fore fins, and eyes, were the most frequently affected body parts. This result highlighted the possibility that turtle aggregations in restricted areas influence the prevalence of FP (Patricio et al., 2016) through positive interactions between individuals. Indeed, videos from cameras fixed on green turtles' shells have shown that green turtles sometimes rub their heads against each other and their upper body parts get consequently in contact during interactions occurring on feeding areas (pers. obs.). Moreover, the DNA of ChHV5 has also been detected in saliva and ocular secretion of green turtles affected by FP, thus representing another possibility of viral excretion and FP transmission, even if direct

transmission from one turtle to another is not fully understood for the moment (Domiciano et al., 2017; Patricio et al., 2016; Rossi et al., 2019).

5. Conclusion and perspectives

Fibropapillomatosis is a potentially debilitating condition, depending on the severity of the lesions, that affects green turtle populations worldwide. We observed an evolution of the FP prevalence in Martinique with an increase of 50% in seven years in one particular bay: Anse du Bourg/Chaudière. We supposed that the high density of immature green turtles in restricted areas enhances positive interactions between individuals and can therefore promote the transmission of FP from one turtle to another. We hypothesised that the addition of factors promoting FP such as lower water quality (i.e. possible eutrophication due to high nutrient loads, high bacterial parameters) or the presence of a wastewater discharge is responsible for the disease's outbreak at Anse du Bourg/Chaudière. Further studies regarding seawater quality (e.g. eutrophication, pollutant presence), the presence and quantity of ChHV5 in the environment, and fine scale currentology, SST, and NPP in the different capture sites are necessary to find clear evidence on how these parameters influence FP dynamics.

The presence of this infectious disease on Martinique Island is of great concern since it is a key developmental area for green turtles. The knowledge of the population's health status is critical to establish conservation programs for this species. FP emergence is recent compared to the CMR program. Thus, the prosecution of the capture program for several years and with more captures in every location will allow us to describe whether Anse du Bourg/Chaudière is the only site affected by FP in Martinique. Moreover, an important number of recaptures will permit comparison of survival rates between healthy and sick turtles to determine whether FP has an impact on immature green turtle population dynamics in Martinique. However, the presence of ChHV5 DNA in green turtles without external tumours (Chaves et al., 2017; Page-

Karjian et al., 2015) suggests that FP prevalence based on external tumours is underestimated compared to real prevalence. Monitoring the health status of immature green turtles in Martinique using serological methods (Work et al., 2020; Sposato et al., 2021) and quantitative PCR (Page-Karjian et al., 2015) will be a more efficient way to take into account asymptomatic individuals in FP prevalence measurement and would allow a better understanding of FP disease.

References

- Adnyana W., Ladds P.W., Blair D., 1997 . Observations of fibropapillomatosis in green turtles (*Chelonia mydas*) in Indonesia. Australian Veterinary Journal 75:737-742; DOI: 10.1111/j.1751-0813.1997.tb12258.x.
- Akaike H. 1973. Information theory and an extension of the maximum likelihood principle. In: Proceedings of the Second International Symposium on Information Theory (eds.: Petrov B.N. & Caski F.). Akademiai Kiado, Budapest, pp 267-281; DOI: 10.1007/978-1-4612-1694-0_15.
- Aminot A. & Chaussepied M. 1983. Manuel des Analyses Chimiques en Milieu Marin. CNEXO Editions Jouve: Paris, 395 PP.
- Bates D., Maechler M., Bolker B., Walker S., Christensen R.H.B., Singmann H., Dai B., Scheipl F., Grothendieck G., Green P., Fox J., Bauer A. & Krivitsky P.N. 2012. Package ‘lme4’. CRAN. R Foundation for Statistical Computing. Retrieved from: <https://github.com/lme4/lme4/>.
- Behrenfeld M.J. & Falkowski P.G. 1997. Photosynthetic rates derived from satellite-based chlorophyll concentration. Limnology and Oceanography 42:1-20; DOI: 10.4319/lo.1997.42.1.0001.
- Bjorndal, K.A., 1999. Priorities for research in foraging habitats. In: Research and Management Techniques for the Conservation of Sea Turtles (eds.: Eckert K.L., Bjorndal K.A., Abreu-

- 421 Grobois F.A. & Donnelly M). IUCN/SSC Marine Turtle Specialist Group Publication 4:
422 12–14.
- 423 Bonola M., Girondot M., Robin J-P., Martin J., Siegwalt F., Jeantet L., Lelong P., Grand C.,
424 Chambault P., Etienne D., Gresser J., Hielard G., Arqué A., Régis S., Lecerf N., Frouin
425 C., Lefebvre F., Sutter E., Védie F., Barnerias C., Thieulle L., Bordes R., Guimera C.,
426 Aubert N., Bouaziz M., Pinson A., Flora F., Duru M., Benhalilou A., Murgale C., Maillet
427 T., Andreani L., Campistron G., Sikora M., Rateau F., George F., Eggenspieler J.,
428 Woignier T., Allenou J-P., Louis-Jean L., Chanteur B., Béranger C., Crillon J., Brador A.,
429 Habold C., Le Maho Y. & Chevallier D. 2019. Fine scale geographic residence and annual
430 primary production drive body condition of wild immature green turtles (*Chelonia mydas*)
431 in Martinique Island (Lesser Antilles). *Biology Open* 8:1-10; DOI: 10.1242/bio.048058.
- 432 Burnham K.P. & Anderson D.R. 2002. Model selection and multimodel inference, 2nd ed.
433 Springer- Verlag, New York, 485 PP.
- 434 Chaloupka M., Work T.M., Balazs G.H., Murakawa S.K.K. & Morris R. 2008. Cause-specific
435 temporal and spatial trends in green sea turtle strandings in the Hawaiian Archipelago
436 (1982-2003). *Marine Biology* 154:887-898; DOI: 10.1007/S00227-008-0981-4.
- 437 Chambault P., De Thoisy B., Huguin M., Martin J., Bonola M., Etienne D., Gresser J., Hiélard
438 G., Mailles J., Védie F., Barnerias C., Sutter E., Guillemot B., Dumont-Dayot E., Régis S.,
439 Lecerf N., Lefebvre F., Frouin C., Aubert N., Guimera C., Bordes R., Thieulle L., Duru
440 M., Bouaziz M., Pinson A., Flora F., Queneherve P., Woignier T., Allenou J-P., Cimiterra
441 N., Benhalilou A., Murgale C., Maillet T., Rangon L., Chanteux N. Chanteur B., Béranger
442 C., Le Maho Y., Petit O. & Chevallier D. 2018. Connecting paths between juvenile and
443 adult habitats in the Atlantic green turtle using genetics and satellite tracking. *Ecology and*
444 *Evolution* 8:12790-12802; DOI: 10.1002/ece3.4708.
- 445 Chaves A., Aguirre A.A., Blanco-Peña K., Moreira-Soto A., Monge O., Torres A.M., Soto-
446 Rivas J.L., Lu Y., Chacon D., Fonseca L., Jiménez M., Gutiérrez-Espeleta G. & Lierz M.
447 2017. Examining the Role of Transmission of Chelonid Alphaherpesvirus 5. *EcoHealth*
448 14:530-541; DOI: 10.1007/s10393- 017-1248-7.

- 450 da Silva C.C., Klein R.D., Barcarolli I.F. & Bianchini A. 2016. Metal contamination as a
451 possible etiology of fibropapillomatosis in juvenile female green sea turtles *Chelonia*
452 *mydas* from the southern Atlantic Ocean. *Aquatic Toxicology* 170:42-51; DOI:
453 10.1016/j.aquatox.2015.11.007.
- 454 Domiciano I.G., Domit C., Bracarense L.R.F.P.A. 2017. The green turtle *Chelonia mydas* as a
455 marine and coastal environmental sentinel: anthropogenic activities and diseases. *Semina:*
456 *Ciências Agrárias, Londrina* 38:3417-3434; DOI: 10.5433/1679-0359.2017v38n5p3417.
- 457 Foley A.M., Schroeder B.A., Redlow A.E., Fick-Child K.J. & Teas W.G. 2005.
458 Fibropapillomatosis in stranded green turtles (*Chelonia mydas*) from the Eastern United
459 States (1980-98): Trends and associations with environmental factors. *Journal of Wildlife*
460 *Diseases* 41: 29-41; DOI: 10.7589/0090-3558-41.1.29.
- 461 Hargove S., Work T., Brunson S., Foley A. M., Balazs G. 2016. Proceedings of the 2015
462 International Summit on Fibropapillomatosis: Global Status, Trends, and Population
463 Impacts. U.S. Dep. Commer., NOAA Tech. Memo., NOAA-TM-NMFS-PIFSC, 87 PP.
464 DOI:10.7289/V5/TM-PIFSC-54.
- 465 Herbst L.H. 1994. Fibropapillomatosis of marine turtles. *Annual Review of Fish Diseases* 4:
466 389-425; DOI: 10.1016/0959-8030(94)90037-X.
- 467 Herbst L.H. & Klein P.A. 1995. Green Turtle Fibropapillomatosis: Challenges to Assessing
468 the Role of Environmental Cofactors. *Environmental Health Perspectives* 103:27-30; DOI:
469 10.1289/ehp.95103s427.
- 470 Herbst L.H., Ene A., Su M., Desalle R. & Lenz J. 2004. Tumor outbreaks in marine turtles are
471 not due to recent herpesvirus mutations. *Current Biology* 14:697-699; DOI:
472 10.1016/j.cub.2004.08.040.
- 473 Hily C., Duchêne J., Bouchon C., Bouchon-Navaro Y., Gigou A., Payri C. & Védie F. 2010.
474 Les herbiers de phanérogames marines de l'outre-mer français, écosystèmes associés aux
475 récifs coralliens. IFRECOR, Conservatoire du littoral, 140 PP.
- 476 Hothorn T., Zeileis A., Farebrother R.W., Cummins C., Millo G., Mitchell D. & Zeileis M.A.
477 2015. Package 'lmtree'. *CRAN. R Foundation for Statistical Computing*. Retrieved from:
478 <https://github.com/cran/lmtree>.

- Impact Mer, 2016. Suivi du milieu récepteur de l'émissaire de la STEU des Anses d'Arlet - Rapport pour : SICSM Martinique, 22 PP.
- Jones K., Ariel E., Burgess G., Read M. 2015. A review of fibropapillomatosis in Green Turtles (*Chelonia mydas*). *The Veterinary Journal* 212:48-57; DOI: 10.1016/j.tvjl.2015.10.041.
- Jones K., Burgess G., Budd A.M., Huerlimann R., Mashkour N. & Ariel E. 2020. Molecular evidence for horizontal transmission of chelonid alphaherpesvirus 5 at green turtle (*Chelonia mydas*) foraging grounds in Queensland, Australia. *PLoS ONE* 5:e0227268; DOI: 10.1371/journal.pone.0227268.
- Lenth R.V., Buerkner P., Herve M., Love J., Riebl H. & Singmann H. 2021. Estimated marginal means, aka least-squares means. *CRAN. R foundation for Statistical Computing*. Retrieved from: <https://github.com/rvlenth/emmeans>.
- Nivière M., Chambault P., Pérez T., Etienne D., Bonola M., Martin J., Barnerias C., Védie F., Mailles J., Dumont-Dayot E., Gresser J., Hiélaud G., Régis S., Lecerf N., Thieulle L., Duru M., Lefebvre F., Milet G., Guillemot B., Bildan B., Montgolfier B., Benhalilou A., Murgale C., Maillet T., Queneherve P., Woignier T., Safi M., Le Maho Y., Petit O. & Chevallier D. 2018. Identification of marine key areas across the Caribbean to ensure the conservation of the critically endangered hawksbill turtle. *Biological Conservation* 223: 170-180; DOI: 10.1016/j.biocon.2018.05.002.
- Murakawa S.K.K., Balazs G.H., Ellis D.M., Hau S. & Eames S.M. 2000. Trends in fibropapillomatosis among green turtles stranded in the Hawaiian Islands, 1982-98. In: *Proceedings of the 19th annual symposium on sea turtle biology and conservation*. South Padre Island, Texas. NOAA Technical Memorandum NMFS-SEFSC-443, 239–241.
- Page-Karjian A., Norton T.M., Ritchie B., Brown C., Mancina C., Jackwood M., Gottdenker N.L. 2015. Quantifying chelonid herpesvirus 5 in symptomatic and asymptomatic rehabilitating green sea turtles. *Endangered Species Research* 28:135-146.; DOI: 10.3354/esr00687.
- Patricio A.R., Diez C.E., Van Dam R.P. & Godley B.J. 2016. Novel insights into the dynamics of green turtle fibropapillomatosis. *Marine Ecology Progress Series* 547:247-255; DOI: 10.3354/meps11644.

- R Core Team. 2020. R: A language and environment for statistical computing. *R Foundation for Statistical Computing*. Retrieved from <http://www.R-project.org/>.
- Rittenburg L.T., Kelley J.R., Mansfield K.L. & Savage A.E., 2021. Marine leech parasitism of sea turtles varies across host species, seasons, and the tumor disease fibropapillomatosis. *Diseases of Aquatic Organisms* 143:1-12; DOI: 10.3354/dao03549.
- Rossi S., Sánchez-Sarmiento A.M., Santos R.G., Zamana R.R., Prioste F.E.S., Gattamorta M.A., Ochoa P.F.C., Grisi-Filho J.H.H. & Matushima E.R. 2019. Monitoring green sea turtles in Brazilian feeding areas: relating body condition index to fibropapillomatosis prevalence. *Journal of the Marine Biological Association of the United Kingdom* 99: 1879-1887; DOI: 10.1017/S0025315419000730.
- Santos R.G., Martins A.S., Torezani E., Baptistotte C., Da Nobrega Farias J., Horta P.A., Work T.M. & Balazs G.H. 2010. Relationship between fibropapillomatosis and environmental quality: a case study with *Chelonia mydas* off Brazil. *Disease of Aquatic Organisms* 89: 87-95; DOI: 10.3354/dao02178.
- Seminoff, J.A. (Southwest Fisheries Science Center, U.S.). 2004. *Chelonia mydas*. *The IUCN Red List of Threatened Species* 2004:e.T4615A11037468. Available: <https://dx.doi.org/10.2305/IUCN.UK.2004.RLTS.T4615A11037468.en>. [Accessed October 23, 2021].
- Shaver D.J., Walker J.S. & Backof T.F. 2019. Fibropapillomatosis prevalence and distribution in green turtles *Chelonia mydas* in Texas (USA). *Diseases of Aquatic Organisms* 136: 175-182; DOI: 10.3354/dao03403.
- Siegwalt F., Benhamou S., Girondot M., Jeantet L., Martin J., Bonola M., Lelong P., Grand C., Chambault P., Benhalilou A., Murgale C., Maillet T., Andreani L., Campistron G., Jacaria F., Hielard G., Arqué A., Etienne D., Gresser J., Régis S., Lecerf N., Frouin C., Lefebvre F., Aubert N., Védie F., Barnerias C., Thieulle L., Guimera C., Bouaziz M., Pinson A., Flora F., George F., Eggenspieler J., Woignier T., Allenou J-P., Louis-Jean L., Chanteur B., Béranger C., Crillon J., Brador A., Habolde C., Le Maho Y., Robin J-P. & Chevallier D. 2020. High fidelity of immature green turtle (*Chelonia mydas*) to their foraging grounds revealed by satellite tracking and capture-mark-recapture and consequences for

- 537 key marine conservation areas. *Biological Conservation* 250:108742; DOI:
538 10.1016/j.biocon.2020.108742.
- 539 Sposato, P., Keating, P., Lutz, P. L. & Milton, S. L. 2021. Evaluation of immune function
540 in two populations of green sea turtles (*Chelonia mydas*) in a degraded versus a
541 nondegraded habitat. *Journal of Wildlife Diseases* 57:761-772; DOI: 10.7589/JWD-D-20-
542 00204.
- 543 Torezani E., Baptistotte C., Mendes S. L., Barata P. C. R. 2010. Juvenile green turtles
544 (*Chelonia mydas*) in the effluent discharge channel of a steel plant, Espirito Santo, Brazil,
545 2000-2006. *Journal of the Marine Biological Association of United Kingdom*, 90:233-246;
546 DOI: 10.1017/S0025315409990579.
- 547 Van Houtan K.S., Hargrove S.K. & Balazs G.H. 2010. Land use, macroalgae, and a tumor-
548 forming disease in marine turtles. *PLoS ONE* 5:e12900; DOI:
549 10.1371/journal.pone.0012900.
- 550 Van Houtan K.S., Smith C.M., Dailer M.L. & Kawachi M. 2014. Eutrophication and the dietary
551 promotion of sea turtle tumors. *PeerJ* 2:e602; DOI: 10.7717/peerj.602.
- 552 Williams E.H., Bunkley-Williams Jr. & Bunkley-Williams L. 1994. An Epizootic of Cutaneous
553 Fibropapillomas in Green Turtles *Chelonia mydas* of the Caribbean: Part of a Panzootic?
554 *Journal of Aquatic Animal Health* 6:70-78; DOI: 10.1577/1548-
555 8667(1994)006<0070:AEOCFI>2.3.CO;2.
- 556 Work T. M., Dagenais J., Willimann A., Balazs G., Mansfield K. & Ackermann M. 2020.
557 Differences in Antibody Responses against Chelonid Alphaherpesvirus 5 (ChHV5)
558 Suggest Differences in Virus Biology in ChHV5-Seropositive Green Turtles from Hawaii
559 and ChHV5-Seropositive Green Turtles from Florida. *Journal of Virology* 94:e01658-19;
560 DOI: 10.1128/JVI.01658-19.

Figures and tables

Figure 1: Martinique map showing capture sites of immature green turtles *Chelonia mydas* (black dots).

Figure 2: Annual fibropapillomatosis prevalence from 2010 to 2019 (except 2014) of immature green turtles on the Caribbean coast of Martinique: Anse Noire/Dufour (n = 56), Grande Anse (n = 302), and Anse du Bourg/Chaudière (n = 181). Vertical bars represent 95% confidence interval.

Figure 3: Anatomical distribution of fibropapillomatosis tumours on the dorsal (left) and ventral (right) sides based on CMR data from 2010 to 2019. The percentages refer to the relative number of individuals showing tumours on the different body areas (n = 128 affected body parts on 52 distinct green turtles).

Figure 4: Water quality parameters in 2021 for several capture sites: Anse Noire/Dufour (n = 4), Grande Anse (n = 3), Anse du Bourg/Chaudière (n = 5), and Petite Anse (n = 3). Dots represent individual measures, columns the means, and vertical bars 95% confidence intervals. Ammonium, nitrites, nitrates, total nitrogen and phosphates are expressed in $\mu\text{mol.L}^{-1}$, total chlorophyll *a* in $\mu\text{g.L}^{-1}$, *Escherichia coli* and enterococci in MPN.100mL⁻¹.

Figure 5: Green turtle density maps of Anse Noire, Anse Dufour, Grand Anse, Anse du Bourg, Anse Chaudière and Petite Anse in 2018 (left) and 2019 (right). Each hexagon represents one hectare and darker hexagonal colour shades indicate a higher turtle concentration (expressed as the number of individuals per hectare).

Table 1: Synthetic results of generalised linear models applied to observed fibropapillomatosis prevalence. Models without results had too many parameters compared to the sample size. Best fit models are in bold. Δ AIC represents the difference between the AIC of a model and the one from the model with the highest relative Akaike weight.

| Model | AIC | Δ AIC | Relative Akaike weight |
|---|--------------|--------------|------------------------|
| Null | 167.94 | 88.19 | < 0.0001 |
| Capture site | 101.51 | 21.77 | < 0.0001 |
| Year | 144.13 | 64.39 | < 0.0001 |
| Temperature | 145.65 | 65.91 | < 0.0001 |
| NPP | 167.37 | 87.62 | < 0.0001 |
| Capture site + year | 79.74 | 0.00 | 0.45 |
| Capture site + temperature | 88.79 | 9.05 | 0.0049 |
| Capture site + NPP | 102.84 | 23.10 | < 0.0001 |
| Capture site + year + temperature | 81.74 | 2.00 | 0.16 |
| Capture site + year + NPP | 81.44 | 1.70 | 0.19 |
| Capture site + year + temperature + NPP | 82.42 | 2.68 | 0.12 |

Table 2: Results of likelihood ratio tests performed on generalised linear mixed models of seven different seawater parameters. Sampling replication (Replicate) was included as random effect.

| Model | df | LogLik | χ^2 | p-value |
|---|----|---------|----------|---------|
| Ammonium ~ (1 Replicate) | 3 | -15.35 | | |
| Ammonium ~ Capture site + (1 Replicate) | 6 | -9.12 | 12.47 | 0.006 |
| Nitrites ~ (1 Replicate) | 3 | 22.91 | | |
| Nitrites ~ Capture site + (1 Replicate) | 6 | 37.01 | 28.20 | < 0.001 |
| Nitrates ~ (1 Replicate) | 3 | 7.12 | | |
| Nitrates ~ Capture site + (1 Replicate) | 6 | 9.77 | 5.31 | 0.150 |
| Phosphates ~ (1 Replicate) | 3 | 26.95 | | |
| Phosphates ~ Capture site + (1 Replicate) | 6 | 31.84 | 9.78 | 0.021 |
| Chlorophyll <i>a</i> ~ (1 Replicate) | 3 | 12.35 | | |
| Chlorophyll <i>a</i> ~ Capture site + (1 Replicate) | 6 | 14.85 | 5.00 | 0.172 |
| <i>E. coli</i> ~ (1 Replicate) | 3 | -130.18 | | |
| <i>E. coli</i> ~ Capture site + (1 Replicate) | 6 | -126.89 | 6.57 | 0.087 |
| Enterococci ~ (1 Replicate) | 3 | -41.08 | | |
| Enterococci ~ Capture site + (1 Replicate) | 6 | -39.92 | 2.31 | 0.510 |

Table 3: Post-hoc comparisons between capture sites according estimated marginal means for ammonium, nitrites and phosphates. AB/C is Anse du Bourg/Chaudière, AN/D Anse Noire/Dufour, GA Grande Anse, and PA Petite Anse. Degrees of freedom were calculated according Kenward-Roger method and p-values adjusted using Tukey's method for multiple comparison. Significant differences are in bold.

| Comparison | Estimate | SE | df | t-ratio | p-value |
|--------------------|---------------|--------------|--------------|--------------|-------------------|
| <i>Ammonium</i> | | | | | |
| AB/C - AN/D | -0.733 | 0.356 | 9.64 | -2.06 | 0.233 |
| AB/C - GA | 0.293 | 0.382 | 9.35 | 0.77 | 0.867 |
| AB/C - PA | -1.033 | 0.382 | 9.35 | -2.70 | 0.091 |
| AN/D - GA | 1.026 | 0.415 | 10.12 | 2.47 | 0.125 |
| AN/D - PA | -0.301 | 0.415 | 10.12 | -0.73 | 0.884 |
| GA - PA | -1.327 | 0.424 | 9.17 | -3.13 | 0.049 |
| <i>Nitrites</i> | | | | | |
| AB/C - AN/D | 0.107 | 0.016 | 9.64 | 6.54 | < 0.001 |
| AB/C - GA | 0.038 | 0.018 | 9.35 | 2.16 | 0.205 |
| AB/C - PA | 0.107 | 0.018 | 9.35 | 6.09 | < 0.001 |
| AN/D - GA | -0.069 | 0.019 | 10.12 | -3.62 | 0.020 |
| AN/D - PA | 0.000 | 0.019 | 10.12 | 0.00 | 1.000 |
| GA - PA | 0.069 | 0.020 | 9.17 | 3.54 | 0.026 |
| <i>Phosphates</i> | | | | | |
| AB/C - AN/D | 0.027 | 0.023 | 9.64 | 1.14 | 0.674 |
| AB/C - GA | 0.074 | 0.025 | 9.35 | 2.97 | 0.061 |
| AB/C - PA | 0.004 | 0.025 | 9.35 | 0.16 | 0.998 |
| AN/D - GA | 0.048 | 0.027 | 10.12 | 1.76 | 0.346 |
| AN/D - PA | -0.023 | 0.027 | 10.12 | -0.83 | 0.838 |
| GA - PA | -0.070 | 0.028 | 9.17 | -2.53 | 0.119 |

Fibropapillomatosis prevalence and distribution in immature green turtles (*Chelonia mydas*) in Martinique Island (Lesser Antilles)

Figures




Figure 1: Martinique map showing capture sites of immature green turtles *Chelonia mydas* (black dots).




Figure 2: Annual fibropapillomatosis prevalence from 2010 to 2019 (except 2014) of immature green turtles on the Caribbean coast of Martinique: Anse Noire/Dufour (n = 56), Grande Anse (n = 302), and Anse du Bourg/Chaudière (n = 181). Vertical bars represent 95% confidence interval.




Figure 3: Anatomical distribution of fibropapillomatosis tumours on the dorsal (left) and ventral (right) sides based on CMR data from 2010 to 2019. The percentages refer to the relative number of individuals showing tumours on the different body areas (n = 128 affected body parts on 52 distinct green turtles).




Figure 4: Water quality parameters in 2021 for several capture sites: Anse Noire/Dufour (n = 4), Grande Anse (n = 3), Anse du Bourg/Chaudière (n = 5), and Petite Anse (n = 3). Dots represent individual measures, columns the means, and vertical bars 95% confidence intervals. Ammonium, nitrites, nitrates, total nitrogen and phosphates are expressed in $\mu\text{mol.L}^{-1}$, total chlorophyll *a* in $\mu\text{g.L}^{-1}$, *Escherichia coli* and enterococci in MPN.100mL $^{-1}$.




Figure 5: Green turtle density maps of Anse Noire, Anse Dufour, Grand Anse, Anse du Bourg, Anse Chaudière and Petite Anse in 2018 (left) and 2019 (right). Each hexagon represents one hectare and darker hexagonal colour shades indicate a higher turtle concentration (expressed as the number of individuals per hectare).

Supplementary Material




Figure 1: Sampling points of sea surface temperature (black circle) and net primary production (numbered triangles) on the west coast of Martinique Island.




Figure 2: Annual fibropapillomatosis prevalence from 2010 to 2019 (except 2014) of immature green turtles on the Caribbean coast of Martinique where captures were weak: Prêcheur ($n = 17$), Saint-Pierre ($n = 3$), Carbet ($n = 5$), Cap enragé ($n = 2$), and Petite Anse ($n = 4$). Vertical bars represent 95% confidence interval.




Figure 3: Trimestrial mean net primary production (NPP) on the West coast of Martinique Island for six sampling points from 2010 to 2019 (2160 x 4320 grid).




Figure 4: Trimestrial mean sea surface temperature (SST) on the West coast of Martinique Island from 2010 to 2019 on the 14.55° latitude and -61.25° longitude point (0.25° x 0.25° grid).




Figure 5: Predicted versus observed fibropapillomatosis prevalence using the *capture site + year* model (a) and the *capture site + year + NPP* model (b), comparison between predicted values of these two models (c) and regression between mean net primary production and the observed prevalence of the *capture site + year + NPP* model (d). Vertical and horizontal bars represent 66% confidence intervals. The regression's shaded area represents the 95% confidence interval.

Fibropapillomatosis prevalence and distribution in immature green turtles
(*Chelonia mydas*) in Martinique Island (Lesser Antilles)

Supplementary Material




Figure 1: Sampling points of sea surface temperature (black circle) and net primary production (numbered triangles) on the west coast of Martinique Island.




Figure 2: Annual fibropapillomatosis prevalence from 2010 to 2019 (except 2014) of immature green turtles on the Caribbean coast of Martinique where captures were weak: Prêcheur ($n = 17$), Saint-Pierre ($n = 3$), Carbet ($n = 5$), Cap enragé ($n = 2$), and Petite Anse ($n = 4$). Vertical bars represent 95% confidence interval.




Figure 3: Trimestrial mean net primary production (NPP) on the West coast of Martinique Island for six sampling points from 2010 to 2019 (2160 x 4320 grid).




Figure 4: Trimestrial mean sea surface temperature (SST) on the West coast of Martinique Island from 2010 to 2019 on the 14.55° latitude and -61.25° longitude point (0.25° x 0.25° grid).




Figure 5: Predicted versus observed fibropapillomatosis prevalence using the *capture site + year* model (a) and the *capture site + year + NPP* model (b), comparison between predicted values of these two models (c) and regression between mean net primary production and the observed prevalence of the *capture site + year + NPP* model (d). Vertical and horizontal bars represent 66% confidence intervals. The regression's shaded area represents the 95% confidence interval.